

THE TOTAL CHORD LENGTH OF MAXIMAL OUTERPLANAR GRAPHS

HALEY BROADUS AND ELENA PAVELESCU

ABSTRACT. We consider embeddings of maximal outerplanar graphs whose vertices all lie on a cycle \mathcal{C} bounding a face. Each edge of the graph that is not in \mathcal{C} , a chord, is assigned a length equal to the length of the shortest path in \mathcal{C} between its endpoints. We define the total chord length of a graph as the sum of lengths of all its chords. For each order $n \geq 5$, we find outerplanar graphs whose total chord length is minimal among all graphs of the same order, and graphs whose total chord length is maximal among all graphs of the same order. We give a complete characterization of those graphs whose total chord length is maximal. We show that every integer value in the interval determined by the minimum and maximum values is the total chord length of a maximal outerplanar graph of the same order.

1. INTRODUCTION

An outerplanar graph is a graph that has a planar embedding such that all the vertices belong to the same face of the embedding. A maximal outerplanar graph is an outerplanar graph that fails to remain outerplanar after the addition of any one edge. Outerplanarity is a hereditary property and it is characterized by the forbidden minors K_4 and $K_{3,2}$ [1]. A maximal outerplanar graph with n vertices can be drawn as an n -cycle \mathcal{C} together with a triangulation of the bounded plane region determined by the cycle. In this article, a maximal outerplanar graph means a drawing which includes \mathcal{C} and a triangulation of the bounded region. The edges of the graph that do not belong to the cycle \mathcal{C} are called *chords*. A maximal outerplanar graph with n vertices has $n - 3$ chords and each chord is assigned a length equal to the shortest path between its endpoints included in the cycle \mathcal{C} . A chord can have length as small as 2 and as large as $\lfloor \frac{n}{2} \rfloor$. A chord of length $\frac{n}{2}$ is called a *diameter* of the graph. The *total chord length* (TCL) of a maximal outerplanar graph equals the sum of the lengths of all its chords.




FIGURE 1. The two maximal outerplanar graphs on six vertices. The graph on the left has three chords of length 2 and $TCL = 6$, and the graph on the right has two chords of length 2, one chord of length 3, and $TCL = 7$.

In this article, we derive explicit formulas for the maximum and minimum total chord lengths over all maximal outerplanar graphs of a given order. These values are given in Theorems 1 and 2. We prove Theorems 1 and 2 in Sections 3 and 4, respectively, and in each case we give specific examples of graphs for which these extreme values are realized.

Theorem 1. *Let $k \geq 0$. For $3 \cdot 2^k \leq n \leq 3 \cdot 2^{k+1}$, the minimal TCL among all maximal outerplanar graphs of order n is:*

$$\min TCL(n) = n(k + 2) - 3 \cdot 2^{k+1}.$$

Theorem 2. *Let $n \geq 5$. The maximal TCL among all maximal outerplanar graphs of order n is*

$$\max TCL(n) = \begin{cases} \frac{n^2-9}{4}, & \text{if } n \text{ odd} \\ \frac{n^2-8}{4}, & \text{if } n \text{ even.} \end{cases}$$

In Section 5, we show that for every order $n \geq 5$ and every $\min TCL(n) \leq l \leq \max TCL(n)$, there exists a maximal outerplanar graph of order n whose total chord length equals l .

2. BACKGROUND AND NOTATION

A chord of length 2 in a maximal outerplanar graph is called an *ear*. Each ear corresponds to a vertex of degree 2 of the graph. The graphs in Figure 1 have three and two ears, respectively. The following theorem frequently attributed to G.H. Meisters [2] is called the “two ears theorem” and can be proved by induction on the number of vertices.

Theorem 3. *A maximal outerplanar graph with $n \geq 5$ vertices has at least two ears.*

Consider a graph G of order n and \mathcal{C} its n -cycle. For an edge $ab \in E(\mathcal{C})$, we define $n_G(ab)$ as the number of chords that “layer” ab . More precisely,

- If G does not contain a diameter: $n_G(ab) :=$ number of chords in G whose length is computed along a path in \mathcal{C} that includes ab .
- If G does contain a diameter: $n_G(ab) := \frac{1}{2} +$ number of chords in G whose length is computed along a path in \mathcal{C} that includes ab , except the diameter.

In Figure 2, $n_G(ab) = 2$ for the graph on the left, and $n_G(ab) = 2\frac{1}{2}$ for the graph on the right. We note that

$$TCL(G) = \sum_{ab \in E(\mathcal{C})} n_G(ab).$$

We also note that for a graph G of order n and an edge $ab \in E(\mathcal{C})$,

$$n_G(ab) \leq \frac{n-3}{2}.$$

An *edge contraction* means that the endpoints of the edge are identified to one vertex that inherits all the neighbors of the two endpoints. Loops and double edges created by the identification of the two endpoints are deleted. The contraction of a path within




FIGURE 2. Two maximal outerplanar graphs with twelve vertices.

a graph means contracting each edge within that path. *Subdividing the edge ab* means deleting the edge ab and adding a new vertex c together with edges ac and bc .

3. MINIMAL TOTAL CHORD LENGTH

This section starts with an algorithm to construct a maximal outerplanar graph for each order $n \geq 4$. Then the TCL for the graph obtained through this algorithm is computed and shown to be the smallest possible for all graphs of the same order.

For $n \geq 4$, consider an n -cycle \mathcal{C} whose vertices are labeled v_1, v_2, \dots, v_n , in clockwise order, as they appear in \mathcal{C} . Construct a maximal outerplanar graph as follows, where edges are added only within the bounded region determined by \mathcal{C} : (1) Include the shortest chord possible with starting endpoint at v_1 and final endpoint situated after v_1 in the clockwise direction around \mathcal{C} . (2) Make the final endpoint of the previously added chord the starting endpoint of a shortest chord whose final endpoint is positioned in the clockwise direction from its starting endpoint. (3) Continue to add chords of minimal length this way until $n - 3$ chords have been added. Two graphs constructed following this algorithm are shown in Figure 3. The chords are labeled according to the order in which they were added.

We will show that for each order, this algorithm gives a maximal outerplanar graph that minimizes the total chord length. We call graphs constructed in this manner “greedy graphs.”




FIGURE 3. Maximal outerplanar graphs of order eleven (left) and twelve (right).

Proposition 4. *Let $k \geq 0$. For $3 \cdot 2^k \leq n \leq 3 \cdot 2^{k+1}$, the total chord length of the maximal outerplanar graph G_n of order n obtained with the algorithm above is*

$$TCL(G_n) = n(k+2) - 3 \cdot 2^{k+1}.$$

Proof. Let $k \geq 0$. For $n = 3 \cdot 2^k$, the graph G_n contains a triangle whose three edges are chords of length 2^k . For each $ab \in E(\mathcal{C})$, $n_G(ab) = k$, and

$$TCL(G_n) = nk = n(k+2) - 2n = n(k+2) - 3 \cdot 2^{k+1}.$$

For each $3 \cdot 2^k \leq n \leq 3 \cdot 2^{k+1}$, a greedy graph of order n is obtained from the greedy graph on $3 \cdot 2^k$ vertices by subdividing edges of \mathcal{C} in counterclockwise order as they appear in \mathcal{C} . Figure 6 contains greedy graphs of orders 12 through 16 ($k = 2$). Graphs up to order 24 can be obtained by subdividing one by one each of the remaining eight edges of the original 12-cycle. The label v_1 marks the vertex at the beginning of the algorithm that gives the greedy graph of that order. If G' is obtained from G by one such edge subdivision, then

$$TCL(G') = TCL(G) + (k+2).$$




FIGURE 4. Maximal outerplanar graphs of orders twelve through sixteen.

This is because a chord of length 2 is added, and the length of each of k edges which layered the subdivided edge is increased by 1. In turn, for $3 \cdot 2^k \leq n \leq 3 \cdot 2^{k+1}$,

$$TCL(G_n) = 3 \cdot 2^k \cdot k + (n - 3 \cdot 2^k)(k+2) = n(k+2) - 3 \cdot 2^{k+1}.$$

□

Next, we show that for each order $n \geq 4$, the algorithm gives a maximal outerplanar graph that minimizes the total chord length.

Lemma 5. *Assume G' is a graph obtained from G by subdividing an edge $ab \in E(\mathcal{C})$ and adding the chord ab .*

Then

$$TCL(G') = \begin{cases} TCL(G) + 2 + n_G(ab), & \text{if } G \text{ has no diameter,} \\ TCL(G) + \frac{3}{2} + n_G(ab), & \text{if } G \text{ has a diameter.} \end{cases}$$

Note that for each graph of order $n \geq 5$, there are at least two ears. Consequently, for $n \geq 5$, each graph of order $n+1$ is obtained from a graph with n vertices through a subdivision, as described in the Lemma 5.

Theorem 6. For a fixed $k \geq 1$, the largest order of a maximal outerplanar graph such that $n_G(ab) \leq k$ for all $ab \in E(\mathcal{C})$ is

$$\theta(k) = 3 \cdot 2^k.$$

Proof. We may assume that for a graph of largest order with $n_G(ab) \leq k$ for all $ab \in E(\mathcal{C})$, $n_G(ab) = k$ for all $ab \in E(\mathcal{C})$. If, on the contrary, there exists an edge $ab \in E(\mathcal{C})$ with $n_G(ab) < k$, construct a new graph by subdividing the edge ab and adding the chord ab . The resulting graph has $n_G(e) \leq k$ for all edges in its cycle and has a larger order.

Maximal outerplanar graphs may possess one, two, or three chords of the greatest length. Considering that at most one of these chords contributes to $n_G(ab)$ for any edge $ab \in E(\mathcal{C})$, a graph of the largest order satisfying $n_G(ab) = k$ for all $ab \in E(\mathcal{C})$ will have three chords of the greatest length.

We proceed by induction on k . The base case, $k = 1$ is true, since the largest order n for which $n_G(ab) = 1$ is $n = 3 \cdot 2$. Assume the statement holds for k and let G be a graph of order $n = 3 \cdot 2^k$ with $n_G(ab) \leq k$ for each $ab \in E(\mathcal{C})$. Subdivide each edge ab in $E(\mathcal{C})$ and add a chord from a to b , creating a graph G' of order $3 \cdot 2^{k+1}$. The construction ensures $n_{G'}(ac) \leq k + 1$ for each $ac \in E(\mathcal{C}')$, implying $\theta(k + 1) \geq 3 \cdot 2^{k+1}$.




FIGURE 5. Maximal outerplanar graphs of orders twelve through sixteen.

Assume $\theta(k + 1) > 3 \cdot 2^{k+1}$. The graph of greatest order for which $n_G(ab) \leq k + 1$ for all $ab \in E(\mathcal{C})$ has three chords of largest length. Call the three endpoints of these three chords a, b , and c and let T be the triangle they determine. The chords ab, ac , and bc each belongs to a triangle different from T . That is, there exist vertices $x, y, z \in V(G)$ such that abx, acy , and bcz are also triangles of the graph. The vertices a, b, c, x, y , and z determine six arcs of \mathcal{C} . We construct a graph H by deleting the triangle T and contracting to one vertex each the three shortest arcs among ax, bx, by, cy, az , and cz . The graph H is of order greater than $3 \cdot 2^{k+1} \cdot \frac{1}{2} = 3 \cdot 2^k$ and $n_H(ab) \leq k$. This contradicts the induction hypothesis. This means $\theta(k) = 3 \cdot 2^{k+1}$. \square

Corollary 7. If G has order $3 \cdot 2^k \leq n < 3 \cdot 2^{k+1}$ and G' is obtained from G as in Lemma 5, then

$$TCL(G') > \begin{cases} TCL(G) + 2 + k & \text{if } G \text{ has no diameter,} \\ TCL(G) + \frac{3}{2} + k & \text{if } G \text{ has a diameter.} \end{cases}$$

This implies that $TCL(G') \geq TCL(G) + 2 + k$.

Corollary 8. *The greedy graph has minimal total chord length among all maximal outerplanar graphs of order n .*

Proof. The graph G_6 has minimal chord length among all maximal planar graphs with six vertices. Assume by induction that the greedy graph G_n has minimal total chord length among all graphs of order n . For $n \geq 6$, and $3 \cdot 2^k \leq n < 3 \cdot 2^{k+1}$,

$$TCL(G_{n+1}) = TCL(G_n) + k + 2.$$

By Theorem 5.2, for any other graph G' of order $n + 1$, where $3 \cdot 2^k \leq n < 3 \cdot 2^{k+1}$, there exists a graph G of order n such that

$$TCL(G') \geq TCL(G) + k + 2.$$

This implies

$$TCL(G') \geq TCL(G) + k + 2 \geq TCL(G_n) + k + 2 = TCL(G_{n+1}).$$

We conclude that for each $n \geq 6$ the greedy graph G_n has the minimal TCL among all graphs of order n . \square

Remark 9. In the algorithm outlined in this section, a minimal-length chord is added at each step according to a specified method. However, if chords of minimal length are added without constraining their endpoints, the resulting graph may not necessarily have minimal total chord length. It might be valuable to find other algorithms yielding graphs with minimal total chord length.

4. MAXIMAL TOTAL CHORD LENGTH

In this section we describe a particular maximal outerplanar graph and then we compute its total chord length. We show that this TCL is the largest possible for all graphs of the same order.

For $n \geq 5$, consider the graph S_n with n vertices obtained from the cycle of length n whose vertices are labeled v_1, v_2, \dots, v_n in clockwise direction by adding the chords $v_1v_3, v_1v_4, \dots, v_1v_{n-2}$. The resulting graph resembles a scallop shell, and will be referred to as the “shell graph.”




FIGURE 6. Shell graphs of orders eleven(left) and twelve(right).

Proposition 10. *Let $n \geq 5$. The shell graph S_n with n vertices has total chord length*

$$TCL(S_n) = \begin{cases} \frac{n^2-9}{4}, & n \text{ odd} \\ \frac{n^2-8}{4}, & n \text{ even} \end{cases}$$

Proof. For n odd, the graph has two chords of length i for each $i = 2, 3, \dots, \frac{n-1}{2}$. In this case

$$TCL(S_n) = 2 \cdot \sum_{i=2}^{(n-1)/2} i = \frac{n^2-9}{4}.$$

For n even, the graph has a diameter and two chords of length i for each $i = 2, 3, \dots, \frac{n-2}{2}$. In this case

$$TCL(S_n) = \frac{n}{2} + 2 \cdot \sum_{i=2}^{(n-2)/2} i = \frac{n^2-8}{4}.$$

□

Notice that there is inherent symmetry within the shell graphs: for every chord of length l on the left-hand side, there is a corresponding chord of length l on the right-hand side. Additionally, observe that the chord length grows in increments of 1, starting at length 2.

Theorem 11. *The shell graph S_n has maximal total chord length among all maximal outerplanar graphs with n vertices.*

Proof. Each graph of order $n+1$ is obtained from a graph with n vertices through a subdivision, as described in Lemma 5. Assume the graph G' is obtained from the graph G of order n by subdividing an edge $ab \in E(C)$ and adding the chord ab .

Then

$$TCL(G') = \begin{cases} TCL(G) + 2 + n_G(ab), & \text{if } G \text{ has no diameter,} \\ TCL(G) + \frac{3}{2} + n_G(ab), & \text{if } G \text{ has a diameter.} \end{cases}$$

Since $n_G(ab) \leq \frac{n-3}{2}$, we have that

$$TCL(G') \leq \begin{cases} TCL(G) + \frac{n+1}{2}, & \text{if } G \text{ has no diameter,} \\ TCL(G) + \frac{n}{2}, & \text{if } G \text{ has a diameter.} \end{cases}$$

We also have

$$TCL(S_{n+1}) = \begin{cases} TCL(S_n) + \frac{n+1}{2}, & \text{if } n \text{ odd,} \\ TCL(S_n) + \frac{n}{2}, & \text{if } n \text{ even.} \end{cases}$$

The shell graph S_5 has maximal total chord length, as the only graph of order 5. Assume by induction that S_n has maximal total chord length among graphs of order n . For any graph G' of order $n+1$ there exists a graph G of order n such that $TCL(G') - TLC(G) \leq TCL(S_{n+1}) - TCL(S_n)$. Since $TLC(G) \leq TCL(S_n)$, it follows that $TCL(G') \leq TCL(S_{n+1})$, and the proof is complete by induction. □

Proposition 12. *A maximal outerplanar graph has maximal total chord length if and only if it has exactly two ears.*

Proof. To prove the direct implication assume G has order n and at least three ears. The graph G has one, two, or three chords of maximal length. In each case, G has a non-chordal edge $ab \in E(\mathcal{C})$ such that $n_G(ab)$ doesn't have a maximal possible value. See Figure 7. This means that G is obtained from a graph \bar{G} of order $n - 1$ through a subdivision of a non-chordal edge, and $TCL(G) - TCL(\bar{G}) < TLC(S_n) - TLC(S_{n-1})$. This implies G doesn't have maximal total chord length.




FIGURE 7. Maximal outerplanar graphs with one, two, and three chords of maximal length and non-chordal edge ab with less than maximal n_G .

To prove the reverse, consider a graph G that has two exactly to ears. A chord of maximal length layers at least one ear. Since the graph has exactly two ears, then G has at most two chords of maximal length. Assume G has one chord d of maximal length. This means n is even. The chord d , together with each half of the cycle \mathcal{C} that the endpoints of d determine are boundary cycles for two maximal outerplanar graphs H_1 and H'_1 of order $\frac{n}{2} + 1$. These graphs are highlighted in red (H_1) and in blue (H'_1) in Figure 8. As maximal outerplanar graphs, each of these graphs have two ears. Since only one of these two ears is also an ear of G , the second ear of H_1 layers d and has length $\frac{n}{2} - 1$ in G . Call this chord d_1 . See Figure 8(left). The chord d_1 , together with the path along \mathcal{C} which gives its length in G form the boundary cycle for an outerplanar graph H_2 of order $\frac{n}{2}$. The graph H_2 has two ears, one of which layers d_1 and has length $\frac{n}{2} - 2$ in G . Call this chord d_2 . Continue until a chord of length 3 is constructed. This reasoning can be replicated for the graph H'_1 . The graph G has one chord of length $\frac{n}{2}$, and two chords each of lengths $2, 3, \dots, \frac{n}{2} - 1$. Its total chord length is equal to that of the shell graph.

If the graph has two chords of maximal length d and d' (n odd) the same reasoning applies. See Figure 8(right). □

5. ALL POSSIBLE VALUES FOR THE TOTAL CHORD LENGTH

Let G_n represent the greedy graph on n vertices and S_n represent the shell graph on n vertices as described in Sections 3 and 4.

Theorem 13. *For every $n \geq 5$ and $TCL(G_n) \leq l \leq TCL(S_n)$, there exists a maximal outerplanar graph of order n with total chord length equal to l .*




FIGURE 8. Maximal outerplanar graphs of even order (left) and odd order (right) with maximal total chord length.

Proof. We show an algorithm to transition from a shell graph S_n to a greedy graph of the same order. This is achieved by repeatedly creating the chords of the greedy graph G_n clockwise around the outer cycle \mathcal{C} and “re-anchoring” the remaining edges at the endpoint of a newly created chord.

The algorithm for order 12 is presented in Figures 9 and 10. The bold edges in each graph represent edges that are part of the greedy graph. Once these “permanent” edges are formed, they persist through to the final graph. The algorithm can be summarized as follows: Start with the graph S_n whose chordal edges are all incident to v_1 . Mark as permanent the chord v_1v_3 . One by one, replace the other chords incident to v_1 by chords incident to v_3 , as in the first three rows of Figure 9. In this process, v_3 serves as a new “anchor,” and the chord v_3v_5 is a permanent chord. Next, replace the chords incident to v_3 (except the two permanent chords) one by one with chords incident to v_5 . The permanent chord v_5v_7 is added, and v_5 becomes the new anchor. Since a permanent chord is added each time the anchoring vertex is changed, with each change in anchor one vertex is eliminated from the set of possible future anchors and each subsequent anchor necessitates one less chord. This process will end, and by way of construction, the final graph is the greedy graph.

It remains to prove that for each value $TCL(G_n) \leq l \leq TCL(S_n)$, there is a graph of order n whose total chord length is l . Not all values l are achieved by the graphs given by the algorithm. We will show how these missing values can be obtained by modifying the graphs produced by the algorithm. For $r \geq 1$, we say the r th iteration of the algorithm is completed when all chords of length $2^{r+1} - 1$ or smaller have been added as permanent chords.

Let $n = 2^k \cdot m$, $m \geq 1$ odd and $r \leq k$. For $k \geq 3$, a chord replacement for the iterations $r = 1, 2$, and 3 of the algorithm is as on the left, center, and right of Figure 11. In each figure, the chord ac is replaced by the chord bd . Note that on the r -th iteration, permanent chords of length 2^r are created. On the r -th iteration, if s represents the length of the path included in \mathcal{C} between b and c and t represents the length of the path included in \mathcal{C} between d and a , as in Figure 11, then a chord of length $\min\{s + 2^r, t + 2^{r-1}\}$ is replaced by a chord of length $\min\{s + 2^{r-1}, t + 2^r\}$. Such chord replacement decreases the total chord length by up to 2^{r-1} , creating a gap in the sequence of total chord lengths of at most 2^{r-1} . This gap can be remediated. We describe how this is done. A chord of length 2^r in the greedy graph implies the




FIGURE 9. Re-anchoring algorithm applied to the shell graph on twelve vertices.

existence of a chord of length 2^{r-1} . Under a chord of length 2^r in the greedy graph, chord changes can be executed to increase the total chord length with any value between 1 and $2^{r-1} - 1$. In Figure 12, under the chord ad , the chord bd of length 2^{r-1} (left) is replaced by the chord ac of length $2^{r-1} + 2^{r-2}$. This represents a 2^{r-2} gain in the total chord length. A similar chord replacement can be made under the chord ab and further, under chords incident to a of lengths 2^k , $k = r - 2, r - 3, \dots, 2$, creating gains




FIGURE 10. Re-anchoring algorithm applied to the shell graph on twelve vertices (continued)

in the total chord length of $2^{r-4}, \dots, 2, 1$, respectively. These chord replacements can be made independently of each other. Since every number between 1 and $2^{r-1} - 1$ can be written as a sum of numbers in the set $\{1, 2, \dots, 2^{r-2}\}$, maximal outerplanar




FIGURE 11. Chord-replacement.

graphs can be generated in such a way that their total chord lengths fill the gaps in the sequence of total chord lengths for the graphs generated by the algorithm.

For $n = 2^k \cdot m$, m odd, and $r = k + 1$, the r th iterations of the algorithm will give permanent chords of length 2^r but also a permanent chord of length $2^r + 2^{r-1}$. This is because the region of the disk that is being triangulated is bounded by an odd cycle. Depending on the value of n , odd cycles may be created on subsequent iterations and these longer chords will be layered over each other. It is possible that on the r th iteration chords of length of up to $2^{r+1} - 1$ are created. However, a chord replacement still decreases the total chord length by up to 2^{r-1} , and gaps in the sequence of total chord lengths can still be filled as described above.

We conclude that between $\text{TCL}(G_n)$ and $\text{TCL}(S_n)$, any total chord length is possible for a graph of order n .




FIGURE 12. Chord change under a chord of length 2^r . The chord bd (left) is replaced by the chord ac (right), resulting in a total chord length gain of 2^{r-2} .

□

REFERENCES

1. G. Chartrand and F. Harary. *Planar Permutation Graphs*, Annales de l'institut Henri Poincaré. Section B, Calcul des probabilités 3.4 (1967), 433–438
2. G.H.Meisters. *Polygons Have Ears*, The American Mathematical Monthly 82 (1975), 648–652.