FrackyFrac: A Standalone UniFrac Calculator

Amit Lavon^{1,2}, Smadar Shilo^{1,2,3,4}, Ayya Keshet^{1,2} and Eran Segal^{1,2*}

¹Department of Computer Science and Applied Mathematics, Weizmann Institute of Science, Rehovot, Israel
²Department of Molecular Cell Biology, Weizmann Institute of Science, Rehovot, Israel
³The Jesse Z and Sara Lea Shafer Institute for Endocrinology and Diabetes, National Center for Childhood Diabetes, Schneider Children's Medical Center of Israel, Petah Tikva, Israel
⁴Faculty of Medicine, Tel Aviv University, Tel-Aviv, Israel

Abstract

UniFrac is a family of distance metrics over microbial abundances, that take taxonomic relatedness into account. Current tools and libraries for calculating UniFrac have specific requirements regarding the user's technical expertise, operating system, and pre-installed software, which might exclude potential users. FrackyFrac is a native command-line tool that can run on any platform and has no requirements. It can also generate the phylogenetic trees required for the calculation. We show that FrackyFrac's performance is on par with currently existing implementations. FrackyFrac can make UniFrac accessible to researchers who may otherwise skip it due to the effort involved, and it can simplify analysis pipelines for those who already use it.

Introduction

The study of microbial communities involves analyzing their biodiversity. The two axes of diversity typically examined are alpha diversity and beta diversity. Alpha diversity measures the biodiversity within a single community, while beta diversity measures the diversity between different communities. Measures of beta diversity allow the comparison of microbial communities and facilitate downstream analysis such as clustering and visualization.

The Bray-Curtis dissimilarity index [1] is commonly used to measure beta diversity by comparing microbial abundances. This index does not use external information about the compared operational taxonomic units (OTUs). The formula is simple and easy to calculate, and thus it is included in many studies.

UniFrac [2] is a distance metric over microbial abundance vectors that takes into account the phylogenetic relatedness between OTUs. It uses a user-provided phylogenetic tree and returns the Jaccard similarity (or dissimilarity) of the samples over the branches of the tree. This way, two different OTUs are in agreement over the part of the tree that they share, starting from the root and until they diverge to separate branches, and in disagreement over the parts of the tree past their divergence. The closer the OTUs are on the tree, the higher their similarity. Two flavors of UniFrac that are commonly used are weighted and unweighted. In weighted UniFrac, every species contributes to the total similarity relative to its abundance, while in unweighted UniFrac every species has the same weight.

The advantage of UniFrac is in the differential distances between the OTUs, according to their taxonomic relation. Similar environments in different geographies may harbor similar makeups of microbial families, orders, classes, etc. even if the individual species are different. These similarities are lost when using Bray-Curtis, but are kept when using UniFrac, allowing an informative comparison of environments. Over time, improvements and variations to the UniFrac metric were developed. Generalized UniFrac [3], Variance-Adjusted Weighted UniFrac [4], Information UniFrac and Ratio UniFrac [5] were created to broaden the kinds of variance that can be measured. Performance improvements such as Fast UniFrac [6] were introduced to allow UniFrac to run on larger datasets. Recent advancements have introduced the Striped UniFrac [7] algorithm and GPU support [8], which increased the speed of computation to make it applicable to datasets with over 100K samples.

These modern implementations come with specific challenges. R and Python-based implementations require having these languages installed, installing third-party libraries, and having some level of proficiency with these languages. Some of those third-party libraries, such as OpenMP and HDF5, require setting up the environment in which the program runs in a specific way. Bioconda's implementation works with the BIOM

 $^{{\}rm *Corresponding\ author:\ eran.segal@weizmann.ac.il}$

and QZA formats, which are specific to that framework. Some of the implementations do not support Windows. Each such constraint adds friction on the way to getting UniFrac in the analysis pipeline. Scientists may give up on using UniFrac in their research if it requires more energy or expertise than what they have.

To demonstrate the growth potential for UniFrac, we searched Google Scholar for papers that mention Bray-Curtis without mentioning UniFrac, published in 2022 with source "nature" (which included journals from the Nature Publishing Group) and got 562 results. Searching for papers that mention UniFrac, published in 2022 with source "nature" yielded 282 results. Under the assumption that UniFrac is relevant to most of the studies mentioning Bray-Curtis, this suggests a potential to double the use of UniFrac.

Reviewing the latest 20 results from the latter search (papers that mention UniFrac), we found that UniFrac was used in studies with sample sizes ranging from 5 to 3211 samples. In 19 out of the 20 papers, sample sizes ranged from 5 to 247 samples, with an average of 69±66. Interestingly, 19 out of 20 of these papers used only the original variations of UniFrac - weighted and unweighted (supplementary table S1).

We infer that an inclusive implementation of UniFrac, one that is easily accessible, can substantially expand the usage of the UniFrac metric. From the surveyed studies we learn that supporting the weighted and unweighted variations, with reasonable performance on up to a few thousands of samples, can satisfy most currently published studies.

Here, we set out to create a more accessible implementation of the UniFrac metric. We created a toolkit that is optimized for ease of access and ease of use, and can be used on any platform without installation.

Results

A standalone implementation

We evaluated our implementation against criteria that may affect a method's usability (table 1). We refer to installation processes as "builtin" and "third party", where a builtin installation is one that uses a language's builtin package installation pipeline. Third-party installation mechanisms introduce more moving parts to the process, and are typically tested and maintained by smaller communities. This can lead to more fragile installation processes which take more debugging and therefore might exclude users.

Implementation	Supported	Required programming	Required pre-existing	Installation
	operating systems	language	software	
FrackyFrac	All	None	None	None
Phyloseq [9]	All	R	R libraries	Third party, with
				BioConductor
QIIME2 [10]	Linux, macOS	Anaconda Python	>400 Python libraries	Builtin, with a conda
				configuration file
Scikit-bio	Linux, macOS	Python	Python libraries	Builtin, with pip
Bioconda-	Linux, macOS	Anaconda Python	Python libraries	Builtin, with conda
binaries (SSU)				
[7]				

Table 1: Comparison of UniFrac implementations.

Performance

We ran FrackyFrac on the OTU tables of samples from the Earth Microbiome Project [11] (the "silva" reference was used). To compare run times on different input sizes, subsamples of the data were taken arbitrarily. We compared FrackyFrac with SSU which is the implementation used in QIIME2, and with Phyloseq. Scikit-bio was left out because QIIME2 is a faster Python library. FrackyFrac's computation times were higher than SSU's by a factor of about two on a single core, and by a factor of 1.4 on eight cores. Figure 1 shows the performance of the compared implementations.

¹ Query: "bray curtis" -unifrac source:nature

² Query: unifrac source:nature




Figure 1: *UniFrac computation times on Earth Microbiome Project data.* (A) Computation times on 10, 20, 50, 100, 200, 500, 1000 and 2000 samples using a single core. (B) Computation times on 1000, 2000, 5000, 10000 and 23000 (all) samples using 8 cores.

Validation

FrackyFrac's result distances from the previous section were compared with SSU's and were found to be equal. The published repository contains a directory named 'testdata' with runnable tests and comparison scripts. Figure 2 shows the separation of samples from several hosts from the Earth Microbiome Project, using FrackyFrac's weighted and unweighted UniFrac.




Figure 2: *PCoA of samples from different host species.* Distances were calculated FrackyFrac's unweighted (A) and weighted (B) UniFrac, on Earth Microbiome Project samples.

Creating phylogenetic trees

FrackyFrac features a builtin tree maker that uses the method suggested by [12], Mash distance with UPGMA clustering, as a simple way to create phylogenetic trees. Since UniFrac requires a phylogenetic tree for its calculation, we provide this method in case no such tree is available.

Conclusions

In this work we introduce a standalone implementation of weighted and unweighted UniFrac. This evolved from a need to use UniFrac without having to go through software installation, format conversions, or having to switch programming languages or operating systems. FrackyFrac meets that need. Our implementation addresses mainly scientists who are not already versed in one of the existing frameworks for running UniFrac, but also scientists who have been using UniFrac and would like more freedom as to how and where they can

use it. Faster alternatives exist, but for the sample sizes common in current studies, calculation speed has less impact than the human factor.

Some aspects are not covered by this work. In terms of scalability, it does not run on GPUs, making it less suitable for very large datasets. In terms of variation, it supports only weighted and unweighted UniFrac. As shown in the introduction, papers citing UniFrac tend to use the two original variations, so supporting only them covers the vast majority of use cases. Future improvements may include supporting additional flavors of UniFrac, and adding optional integration with third-party libraries for better scalability.

FrackyFrac has the potential to increase the use of UniFrac by reducing the amount of effort and expertise required to use it. It can expand the range of potential users to those with more diverse backgrounds and to more platforms.

Implementation

The FrackyFrac suite contains two executables: frcfrc which takes as input a TSV of abundances and a phylogenetic tree, and outputs pairwise UniFrac distances; and trtr which creates phylogenetic trees given input fasta sequences.

To support different platforms, FrackyFrac is cross-compiled into standalone binaries which contain all the required logic, with a native binary for every operating system. This removes the need for an installation in favor of downloading ready-to-use executables. There is no need to get FrackyFrac's code in order to use it, so no particular programming language is required on the user's side.

To support diverse pipelines, FrackyFrac uses the tab-separated-values (TSV) format for input and output, for which there exist encoders and decoders in most modern programming languages.

The algorithm for calculating UniFrac distances is similar to Fast UniFrac's algorithm, where samples are represented as lists of tree-nodes and abundances. FrackyFrac uses the biostuff package suite [13] for processing biological data and the gostuff package suite [14] for algorithms and data structures.

Availability and requirements

Software: https://github.com/fluhus/frackyfrac. Operating systems: Linux, Windows, MacOS. Software requirements: none. License: MIT. Non-academic use is unrestricted.

Declarations

Availability of data and materials

The Earth Microbiome Project data that was used for demonstration can be downloaded from: https://earthmicrobiome.org/data-and-code/

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

AL: conceptualization, methodology, software, writing - original draft. SS, AK: validation, data curation, writing - review & editing. ES: supervision, writing - review & editing.

References

[1] J Roger Bray and John T Curtis. "An ordination of the upland forest communities of southern Wisconsin". In: *Ecological monographs* 27.4 (1957). Publisher: JSTOR, pp. 326–349.

- [2] Catherine Lozupone and Rob Knight. "UniFrac: a new phylogenetic method for comparing microbial communities". In: *Applied and environmental microbiology* 71.12 (2005). Publisher: Am Soc Microbiol, pp. 8228–8235.
- [3] Jun Chen et al. "Associating microbiome composition with environmental covariates using generalized UniFrac distances". In: *Bioinformatics* 28.16 (2012). Publisher: Oxford University Press, pp. 2106–2113.
- [4] Qin Chang, Yihui Luan, and Fengzhu Sun. "Variance adjusted weighted UniFrac: a powerful beta diversity measure for comparing communities based on phylogeny". In: *BMC bioinformatics* 12.1 (2011). Publisher: BioMed Central, pp. 1–14.
- [5] Ruth G Wong, Jia R Wu, and Gregory B Gloor. "Expanding the UniFrac toolbox". In: *PloS one* 11.9 (2016). Publisher: Public Library of Science San Francisco, CA USA, e0161196.
- [6] Micah Hamady, Catherine Lozupone, and Rob Knight. "Fast UniFrac: facilitating high-throughput phylogenetic analyses of microbial communities including analysis of pyrosequencing and PhyloChip data". In: *The ISME journal* 4.1 (2010). Publisher: Nature Publishing Group, pp. 17–27.
- [7] Daniel McDonald et al. "Striped UniFrac: enabling microbiome analysis at unprecedented scale". In: *Nature methods* 15.11 (2018). Publisher: Nature Publishing Group US New York, pp. 847–848.
- [8] Igor Sfiligoi et al. "Optimizing UniFrac with OpenACC Yields Greater Than One Thousand Times Speed Increase". In: Msystems 7.3 (2022). Publisher: American Society for Microbiology 1752 N St., NW, Washington, DC, e00028–22.
- [9] PAUL J McMURDIE and Susan Holmes. "Phyloseq: a bioconductor package for handling and analysis of high-throughput phylogenetic sequence data". In: *Biocomputing* 2012. World Scientific, 2012, pp. 235–246.
- [10] Evan Bolyen et al. "Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2". In: *Nature biotechnology* 37.8 (2019). Publisher: Nature Publishing Group, pp. 852–857.
- [11] Jack A Gilbert, Janet K Jansson, and Rob Knight. "The Earth Microbiome project: successes and aspirations". In: *BMC biology* 12 (2014). Publisher: BioMed Central, pp. 1–4.
- [12] Brian D Ondov et al. "Mash: fast genome and metagenome distance estimation using MinHash". In: *Genome biology* 17.1 (2016). Publisher: BioMed Central, pp. 1–14.
- [13] Amit Lavon. fluhus/biostuff: v0.1.19. Aug. 2022. DOI: 10.5281/zenodo.7033579. URL: https://doi.org/10.5281/zenodo.7033579.
- [14] Amit Lavon. fluhus/gostuff: v0.1.14. Aug. 2022. DOI: 10.5281/zenodo.7033569. URL: https://doi.org/ 10.5281/zenodo.7033569.
- [15] Thapanan Jatuyosporn et al. "White spot syndrome virus impact on the expression of immune genes and gut microbiome of black tiger shrimp Penaeus monodon". In: *Scientific Reports* 13.1 (2023). Publisher: Nature Publishing Group UK London, p. 996.
- [16] Qianjun Li et al. "Obstructive sleep apnea is related to alterations in fecal microbiome and impaired intestinal barrier function". In: *Scientific Reports* 13.1 (2023). Publisher: Nature Publishing Group UK London, p. 778.
- [17] Abigail E Asangba et al. "Diagnostic and prognostic potential of the microbiome in ovarian cancer treatment response". In: *Scientific reports* 13.1 (2023). Publisher: Nature Publishing Group UK London, p. 730.
- [18] Ping Miao et al. "Exacerbation of allergic rhinitis by the commensal bacterium Streptococcus salivarius". In: *Nature Microbiology* 8.2 (2023). Publisher: Nature Publishing Group UK London, pp. 218–230.
- [19] Dongeun Yong et al. "Altered gut microbiota in individuals with episodic and chronic migraine". In: *Scientific reports* 13.1 (2023). Publisher: Nature Publishing Group UK London, p. 626.
- [20] Shiva T Radhakrishnan et al. "Rectal swabs as a viable alternative to faecal sampling for the analysis of gut microbiota functionality and composition". In: *Scientific Reports* 13.1 (2023). Publisher: Nature Publishing Group UK London, p. 493.
- [21] Linnea Hases et al. "High-fat diet and estrogen modulate the gut microbiota in a sex-dependent manner in mice". In: *Communications Biology* 6.1 (2023). Publisher: Nature Publishing Group UK London, p. 20.

- [22] Beatriz Peñalver Bernabé et al. "Interactions between perceived stress and microbial-host immune components: two demographically and geographically distinct pregnancy cohorts". In: *Translational Psychiatry* 13.1 (2023). Publisher: Nature Publishing Group UK London, p. 3.
- [23] J Philip Karl et al. "Severe, short-term sleep restriction reduces gut microbiota community richness but does not alter intestinal permeability in healthy young men". In: *Scientific Reports* 13.1 (2023). Publisher: Nature Publishing Group UK London, p. 213.
- [24] Bin Wang et al. "Stable colonization of Akkermansia muciniphila educates host intestinal microecology and immunity to battle against inflammatory intestinal diseases". In: *Experimental & Molecular Medicine* 55.1 (2023). Publisher: Nature Publishing Group UK London, pp. 55–68.
- [25] Jae Hyun Kim et al. "Long-term taxonomic and functional stability of the gut microbiome from human fecal samples". In: *Scientific Reports* 13.1 (2023). Publisher: Nature Publishing Group UK London, p. 114.
- [26] You Zhou et al. "The rhizospheric bacterial diversity of Fritillaria taipaiensis under single planting pattern over five years". In: *Scientific Reports* 12.1 (2022). Publisher: Nature Publishing Group UK London, p. 22544.
- [27] Sally L Bornbusch et al. "Microbial rewilding in the gut microbiomes of captive ring-tailed lemurs (Lemur catta) in Madagascar". In: *Scientific Reports* 12.1 (2022). Publisher: Nature Publishing Group UK London, p. 22388.
- [28] Kristine Stø et al. "Fecal level of butyric acid, a microbiome-derived metabolite, is increased in patients with severe carotid atherosclerosis". In: *Scientific Reports* 12.1 (2022). Publisher: Nature Publishing Group UK London, p. 22378.
- [29] Lei Wu et al. "Gut microbiota as an antioxidant system in centenarians associated with high antioxidant activities of gut-resident Lactobacillus". In: *npj Biofilms and Microbiomes* 8.1 (2022). Publisher: Nature Publishing Group UK London, p. 102.
- [30] Eun Hee Ahn et al. "Helicobacter hepaticus augmentation triggers Dopaminergic degeneration and motor disorders in mice with Parkinson's disease". In: *Molecular Psychiatry* 28.3 (2023). Publisher: Nature Publishing Group UK London, pp. 1337–1350.
- [31] Antonio P Camargo et al. "Plant microbiomes harbor potential to promote nutrient turnover in impoverished substrates of a Brazilian biodiversity hotspot". In: *The ISME journal* 17.3 (2023). Publisher: Nature Publishing Group UK London, pp. 354–370.
- [32] Yusuke Nakade et al. "Increased levels of oral Streptococcus-derived d-alanine in patients with chronic kidney disease and diabetes mellitus". In: *Scientific reports* 12.1 (2022). Publisher: Nature Publishing Group UK London, p. 21773.
- [33] Misaki Eto et al. "Dynamics of rumen microbiome in sika deer (Cervus nippon yakushimae) from unique subtropical ecosystem in Yakushima Island, Japan". In: *Scientific reports* 12.1 (2022). Publisher: Nature Publishing Group UK London, p. 21623.
- [34] Jos A Bosch et al. "The gut microbiota and depressive symptoms across ethnic groups". In: *Nature Communications* 13.1 (2022). Publisher: Nature Publishing Group UK London, p. 7129.

Supplementary material

Sample size	Type of UniFrac	Paper title	
12	used W	White spot syndrome virus impact on the expression of immune genes and gut microbiome of black tiger shrimp Penaeus monodon [15]	
48	U	Obstructive sleep apnea is related to alterations in fecal microbiome and impaired intestinal barrier function [16]	
64	U,W,Generalized	Diagnostic and prognostic potential of the microbiome in ovarian cancer treatment response [17]	
160	W	Exacerbation of allergic rhinitis by the commensal bacterium Streptococcus salivarius [18]	
130	U,W	Altered gut microbiota in individuals with episodic and chronic migraine [19]	
20	U	Rectal swabs as a viable alternative to faecal sampling for the analysis of gut microbiota functionality and composition [20]	
64	U,W	High-fat diet and estrogen modulate the gut microbiota in a sex-dependent manner in mice [21]	
84	Not specified	Interactions between perceived stress and microbial-host immune components: two demographically and geographically distinct pregnancy cohorts [22]	
19	U,W	Severe, short-term sleep restriction reduces gut microbiota community richness but does not alter intestinal permeability in healthy young men [23]	
21	W	Stable colonization of Akkermansia muciniphila educates host intestinal microecology and immunity to battle against inflammatory intestinal diseases [24]	
10	U,W	Long-term taxonomic and functional stability of the gut microbiome from human fecal samples [25]	
5	Not specified	The rhizospheric bacterial diversity of Fritillaria taipaiensis under single planting pattern over five years [26]	
172	U,W	Microbial rewilding in the gut microbiomes of captive ring-tailed lemurs (Lemur catta) in Madagascar [27]	
81	U	Fecal level of butyric acid, a microbiome-derived metabolite, is increased in patients with severe carotid atherosclerosis [28]	
247	U	Gut microbiota as an antioxidant system in centenarians associated with high antioxidant activities of gut-resident Lactobacillus [29]	
20	U	Helicobacter hepaticus augmentation triggers Dopaminergic degeneration and motor disorders in mice with Parkinson's disease [30]	
84	W	Plant microbiomes harbor potential to promote nutrient turnover in impoverished substrates of a Brazilian biodiversity hotspot [31]	
17	U,W	Increased levels of oral Streptococcus-derived d-alanine in patients with chronic kidney disease and diabetes mellitus [32]	
49	W	Dynamics of rumen microbiome in sika deer (Cervus nippon yakushimae) from unique subtropical ecosystem in Yakushima Island, Japan [33]	
3211	W	The gut microbiota and depressive symptoms across ethnic groups [34]	

Table S1: Latest 20 papers mentioning UniFrac with source "nature" on Google Scholar (Jan 2023). UniFrac types are weighted (W) and unweighted (U).