# HETEROCLINIC SOLUTIONS FOR SOME CLASSES OF PRESCRIBED MEAN CURVATURE EQUATIONS IN WHOLE $\mathbb{R}^2$

CLAUDIANOR O. ALVES AND RENAN J. S. ISNERI

ABSTRACT. The purpose of this paper consists in using variational methods to establish the existence of heteroclinic solutions for some classes of prescribed mean curvature equations of the type

$$-div\left(\frac{\nabla u}{\sqrt{1+|\nabla u|^2}}\right) + A(\epsilon x, y)V'(u) = 0 \quad \text{in} \quad \mathbb{R}^2,$$

where  $\epsilon > 0$  and V is a double-well potential with minima at  $t = \alpha$  and  $t = \beta$  with  $\alpha < \beta$ . Here, we consider some class of functions A(x,y) that are oscillatory in the variable y and satisfy different geometric conditions such as periodicity in all variables or asymptotically periodic at infinity.

#### 1. INTRODUCTION

The main goal of this paper is to use variational methods to show the existence of heteroclinic solutions for prescribed mean curvature equation of the type

$$-div\left(\frac{\nabla u}{\sqrt{1+|\nabla u|^2}}\right) + A(\epsilon x, y)V'(u) = 0 \quad \text{in} \quad \mathbb{R}^2 \tag{E}$$

taking into account different geometric conditions on function  $A: \mathbb{R}^2 \to \mathbb{R}$  with  $\epsilon > 0$ . Hereafter, we mean by heteroclinic solution a function u that satisfies (E) and has the following asymptotic property at infinity

 $u(x,y) \to \alpha$  as  $x \to -\infty$  and  $u(x,y) \to \beta$  as  $x \to +\infty$ , uniformly in  $y \in \mathbb{R}$ ,

where  $\alpha, \beta \in \mathbb{R}$  are global minima of the potential  $V : \mathbb{R} \to \mathbb{R}$ .

Hereafter,  $\mathcal{V} = \{V_{\alpha,\beta}\}$  denotes a set of potentials that satisfy the following properties:

- $(V_1) \ V_{\alpha,\beta} \in C^1(\mathbb{R},\mathbb{R}).$
- $\begin{array}{l} (V_2) \ \alpha < \beta \ \text{and} \ V_{\alpha,\beta}(\alpha) = V_{\alpha,\beta}(\beta) = 0. \\ (V_3) \ V_{\alpha,\beta}(t) \ge 0 \ \text{for any} \ t \in \mathbb{R} \ \text{and} \ V_{\alpha,\beta}(t) > 0 \ \text{for all} \ t \in (\alpha,\beta). \end{array}$

Moreover, we assume that the family  $\mathcal{V}$  also satisfies the following local uniform estimate involving the numbers  $\alpha$  and  $\beta$ :

 $(V_4)$  Given  $\lambda > 0$ , there exists  $M = M(\lambda) > 0$  independent of  $\alpha, \beta \in (-\lambda, \lambda)$  such that

$$\sup_{t \in (\alpha,\beta)} |V'_{\alpha,\beta}(t)| \le M, \quad \forall V_{\alpha,\beta} \in \mathcal{V} \quad \text{with} \quad \alpha,\beta \in (-\lambda,\lambda).$$

The reader is invited to see that we could consider the family  $\mathcal{V}$  where the potentials  $V_{\alpha,\beta}$  are the form

(1.1) 
$$V_{\alpha,\beta}(t) = (t-\alpha)^2 (t-\beta)^2,$$

<sup>2020</sup> Mathematics Subject Classification. Primary: 35J62, 35A15, 35J93, 34C37; Secondary: 46E30.

Key words and phrases. Heteroclinic solutions, Quasilinear elliptic equations, Variational Methods, Mean curvature operator, Orlicz space.

which was inspired by the classical double well Ginzburg-Landau potential. These family is very important, because those potentials arise in the study of heteroclinic solutions for stationary Allen Cahn type equations

$$-\Delta u + A(x)V'_{\alpha,\beta}(u) = 0 \quad \text{in} \quad \mathbb{R}^n$$

Another important family  $\mathcal{V}$  is related to the Sine-Gordon potential that is formed by potentials of the form

(1.2) 
$$V_{-\beta,\beta}(t) = \beta + \beta \cos\left(\frac{t\pi}{\beta}\right),$$

where  $\alpha = -\beta$ . For a discussion of physical applications appearing these types of potentials, we refer the interested reader to [2, 10].

In what follows, associated with function A we assume

- $(A_1)$  A is continuous and there is  $A_0 > 0$  such that  $A(x, y) \ge A_0$  for all  $(x, y) \in \mathbb{R}^2$ .
- $(A_2)$  A(x,y) = A(x,-y) for all  $(x,y) \in \mathbb{R}^2$ .
- $(A_3)$  A(x,y) = A(x,y+1) for any  $(x,y) \in \mathbb{R}^2$ .

Now let us mention the classes of A that we will considered in our work.

**Class 1:** A satisfies  $(A_1)$ - $(A_3)$  and is 1-periodic on the variable x.

**Class 2:** A satisfies  $(A_1)$ - $(A_3)$  and there exists a continuous function  $A_p : \mathbb{R}^2 \to \mathbb{R}$ , which is 1-periodic on x, satisfying  $A(x, y) < A_p(x, y)$  for all  $(x, y) \in \mathbb{R}^2$  and

$$|A(x,y) - A_p(x,y)| \to 0$$
 as  $|(x,y)| \to +\infty$ .

**Class 3:** A satisfies  $(A_1)$ - $(A_3)$  and

$$\sup_{y \in [0,1]} A(0,y) < \liminf_{|(x,y)| \to +\infty} A(x,y) = A_{\infty} < +\infty.$$

**Class 4:** A satisfies  $(A_2)$ - $(A_3)$ , is a continuous non-negative function, even in  $x, A \in L^{\infty}(\mathbb{R}^2)$ and there exists K > 0 such that

$$\inf_{|x| \ge K, y \in [0,1]} A(x,y) > 0.$$

We would like to highlight that some of these conditions are well known in the context of the Laplacian operator. For example, a condition like Class 1 was studied by Rabinowitz [23] to show the existence of heteroclinic solution for a class of second order partial differential equations in which he includes the equation of the form

(1.3) 
$$-\Delta u + A(x,y)V'(u) = 0 \text{ in } \Omega,$$

where the set  $\Omega$  is a cylindrical domain in  $\mathbb{R}^N$  given by  $\Omega = \mathbb{R} \times D$  with D being a bounded open set in  $\mathbb{R}^{N-1}$  such that  $\partial D \in C^1$ . In the literature we also find interesting works that study the equation (1.3) in the case that A(x, y) is periodic in all variables when  $\Omega = \mathbb{R}^2$ , see for example Rabinowitz and Stredulinsky [24] and Alessio, Gui and Montecchiari [1]. Related to the Classes 2 and 3 we cite a paper by Alves [3], where the author established the existence of classical solutions of (1.3) on a cylindrical domain that are heteroclinic in the variable x. Finally, the Class 4 was introduced in [5].

The prescribed mean curvature operator

(1.4) 
$$div\left(\frac{\nabla u}{\sqrt{1+|\nabla u|^2}}\right)$$

has been extensively studied in the recent years, due to the close connection with capillarity theory [9]. After the pioneering works of Young [27], Laplace [16], and Gauss [12] in the early

18th century about the mean curvature of a capillary surface, much has already been produced in the literature and it is difficult and exhaustive to measure here the vastness of physical applications involving the (1.4) operator, however for the interested reader in this subject, we could cite here some problems that appear in optimal transport [8] and in minimal surfaces [13]. Moreover, (1.4) also appears in some problems in reaction-diffusion processes which occur frequently in a wide variety of physical and biological settings. For example, in [15], Kurganov and Rosenau observed that when the saturation of the diffusion is incorporated into these processes, it may cause a deep impact on the the morphology of the transitions connecting the equilibrium states, as now not only do discontinuous equilibria become permissible, but traveling waves can arise in their place. A specific class of such processes is modeled by the following equation

(1.5) 
$$u_t = div \left(\frac{\nabla u}{\sqrt{1 + |\nabla u|^2}}\right) - aV'(u),$$

where the reaction function V is the classical double well Ginzburg-Landau potential and a is a constant. The impact of saturated diffusion on reaction-diffusion processes was investigated by them in the straight line and in the plane.

As indicated in the previous paragraph, [15] provided a significant physical motivation for the study of equation (1.5) having as main objective the existence of transition-type solutions, that is, entire solutions of (1.5) which are asymptotic in different directions to the equilibrium states of the systems. In this sense, Bonheure, Obersnel and Omari in [7] investigated the existence of a heteroclinic solution of the one-dimensional equation

(1.6) 
$$-\left(\frac{q'}{\sqrt{1+(q')^2}}\right)' + a(t)V'(q) = 0 \text{ in } \mathbb{R},$$

looking for minima of an action functional on a convex subset of  $BV_{\text{loc}}(\mathbb{R})$  made of all functions satisfying an asymptotic condition at infinity, where the authors considered as usual V a doublewell potential with minima at  $t = \pm 1$  and a asymptotic to a positive periodic function, that is,  $a \in L^{\infty}(\mathbb{R})$  with  $0 < \text{ess} \inf_{t \in \mathbb{R}} a(t)$  and there is  $a^* \in L^{\infty}(\mathbb{R})$   $\tau$ -periodic, for some  $\tau > 0$ , such that  $a(t) \leq a^*(t)$  almost everywhere on  $\mathbb{R}$  satisfying

$$\operatorname{ess\,lim}_{|t| \to +\infty} (a^*(t) - a(t)) = 0.$$

Recently in [5], Alves and Isneri also studied the existence of heteroclinic solution for (1.6). In that paper, the authors truncated the mean curvature operator to build up a variational framework inspired to the one introduced in [19] on Orlicz-Sobolev space  $W_{\text{loc}}^{1,\Phi}(\mathbb{R})$  in order to establish the existence of a heteroclinic solution for (1.6) in the case where the function  $a \in L^{\infty}(\mathbb{R})$  is an even non-negative with  $0 < \inf_{t \ge M} a(t)$  for some M > 0. Moreover, in the case where the function a is constant, for each initial conditions  $q(0) = r_1$  and  $q'(0) = r_2$ , the uniqueness of the minimal heteroclinic type solutions for (1.6) has been proved.

Motivated by [5], [7] and [19], in the present paper we intend to analyze the existence of heteroclinic solutions for (E) and their qualitative properties, as well as regularity. A part of our arguments was inspired by papers due to Rabinowitz [23] and Alves [3].

The main results of the paper can be now stated in the following form.

**Theorem 1.1.** Assume  $\epsilon = 1$  and that A belongs to Class 1 or 2. Given L > 0 there exists  $\delta > 0$  such that for each  $V_{\alpha,\beta} \in \mathcal{V}$  with  $\max\{|\alpha|, |\beta|\} \in (0, \delta)$ , the equation (E) with  $V = V_{\alpha,\beta}$  possesses a heteroclinic solution  $u_{\alpha,\beta}$  from  $\alpha$  to  $\beta$  in  $C^{1,\gamma}_{loc}(\mathbb{R}^2)$ , for some  $\gamma \in (0,1)$ , satisfying

- (a)  $u_{\alpha,\beta}$  is 1-periodic on y.
- (b)  $\alpha \leq u_{\alpha,\beta}(x,y) \leq \beta$  for any  $(x,y) \in \mathbb{R}^2$ .
- (c)  $\|\nabla u_{\alpha,\beta}\|_{L^{\infty}(\mathbb{R}^2)} \leq \sqrt{L}.$

Moreover, if  $V_{\alpha,\beta} \in C^2(\mathbb{R},\mathbb{R})$  then the inequalities in (b) are strict.

**Theorem 1.2.** Assume that A belongs to Class 3. There is  $\epsilon_0 > 0$  such that for each  $\epsilon \in (0, \epsilon_0)$ and L > 0 given, there exists  $\delta > 0$  such that for each  $V_{\alpha,\beta} \in \mathcal{V}$  with  $\max\{|\alpha|, |\beta|\} \in (0, \delta)$ , the equation (E) with  $V = V_{\alpha,\beta}$  possesses a heteroclinic solution  $u_{\alpha,\beta}$  from  $\alpha$  to  $\beta$  in  $C^{1,\gamma}_{loc}(\mathbb{R}^2)$ , for some  $\gamma \in (0,1)$ , verifying

- (a)  $u_{\alpha,\beta}$  is 1-periodic on y.
- (b)  $\alpha \leq u_{\alpha,\beta}(x,y) \leq \beta$  for any  $(x,y) \in \mathbb{R}^2$ .
- (c)  $\|\nabla u_{\alpha,\beta}\|_{L^{\infty}(\mathbb{R}^2)} \leq \sqrt{L}$ .

Moreover, if  $V_{\alpha,\beta} \in C^2(\mathbb{R},\mathbb{R})$  occurs then the inequalities in (b) are strict.

Demanding a little more of the class  $\mathcal{V}$ , we can relax the conditions on the function A to ensure the existence of a heteroclinic solution for (E), as the following result says.

**Theorem 1.3.** Assume  $\alpha = -\beta$ ,  $\epsilon = 1$  and that A belonging to Class 4. Then, for each L > 0there exists  $\delta > 0$  such that for each  $V_{-\beta,\beta} \in \mathcal{V} \cap C^2(\mathbb{R},\mathbb{R})$  with  $\beta \in (0,\delta)$  and satisfying the conditions below

- $(V_5) V''_{-\beta,\beta}(-\beta), V''_{-\beta,\beta}(\beta) > 0,$  $(V_6) V_{-\beta,\beta}(t) = V_{-\beta,\beta}(-t) \text{ for all } t \in \mathbb{R},$

the equation (E) with  $V = V_{-\beta,\beta}$  possesses a heteroclinic solution  $u_{\beta}$  from  $-\beta$  to  $\beta$  in  $C^{1,\gamma}_{loc}(\mathbb{R}^2)$ , for some  $\gamma \in (0,1)$ , verifying the following properties:

- (a)  $u_{\beta}(x,y) = -u_{\beta}(-x,y)$  for any  $(x,y) \in \mathbb{R}^2$ .
- (b)  $u_{\beta}(x,y) = u_{\beta}(x,y+1)$  for all  $(x,y) \in \mathbb{R}^2$ .
- (c)  $0 < u_{\beta}(x,y) < \beta$  for x > 0.
- (d)  $\|\nabla u_{\beta}\|_{L^{\infty}(\mathbb{R}^2)} \leq \sqrt{L}.$

The reader is invited to see that the above theorems are true for the Ginzburg-Landau (1.1)and Sine-Gordon (1.2) potentials when roots  $\alpha$  and  $\beta$  have a small distance between them.

Motivated by the ideas in [5], in the proof of the theorems above, we truncate the differential operator involved in (E) in such a way that the new operator can be seen as a quasilinear operator in divergence form. For this reason, as a first step in the present work, we study quasilinear equations of the form

$$-\Delta_{\Phi} u + A(\epsilon x, y) V'(u) = 0 \quad \text{in} \quad \mathbb{R}^2, \tag{PDE}$$

where  $\Delta_{\Phi} u = \operatorname{div}(\phi(|\nabla u|) \nabla u)$  and  $\Phi: \mathbb{R} \to [0, +\infty)$  is an N-function of the type

(1.7) 
$$\Phi(t) = \int_0^{|t|} s\phi(s)ds$$

where  $\phi: (0, +\infty) \to (0, +\infty)$  is a  $C^1$  function verifying the following conditions:

- $(\phi_1) \ \phi(t) > 0$  and  $(\phi(t)t)' > 0$  for any t > 0.
- $(\phi_2)$  There are  $l, m \in \mathbb{R}$  with  $1 < l \le m$  such that

$$l-1 \le \frac{(\phi(t)t)'}{\phi(t)} \le m-1, \quad \forall t > 0.$$

 $(\phi_3)$  There exist constants  $c_1, c_2, \delta > 0$  and q > 1 satisfying

$$c_1 t^{q-1} \le \phi(t) t \le c_2 t^{q-1}, \ t \in (0, \delta].$$

The solutions of (PDE) are found as minima of the action functional

$$I(w) = \sum_{j \in \mathbb{Z}} \left( \int_0^1 \int_j^{j+1} \left( \Phi(|\nabla w|) + A(\epsilon x, y) V(w) \right) dx dy \right)$$

on the class of admissible functions

$$\Gamma_{\Phi}(\alpha,\beta) = \left\{ w \in W^{1,\Phi}_{\text{loc}}(\mathbb{R} \times (0,1)) : \tau_k w \to \alpha \ (\beta) \text{ in } L^{\Phi}((0,1) \times (0,1)) \text{ as } k \to -\infty \ (+\infty) \right\},$$

where  $W_{\text{loc}}^{1,\Phi}(\mathbb{R}\times(0,1))$  denotes the usual Orlicz-Sobolev space. We would like to point out that in the study of quasilinear elliptic problems driven by the  $\Phi$ -Laplacian operator the conditions  $(\phi_1)$ - $(\phi_2)$  are well-known and guarantee that  $\Phi$  is an N-function that checks the so called  $\Delta_2$ condition (see for instance Appendix A). Those conditions ensure that the Orlicz-Sobolev space is reflexive and separable. Moreover, in this work, the conditions  $(\phi_1)$ - $(\phi_2)$  are enough to show the existence of heteroclinic solution, while assumption  $(\phi_3)$  is used to get more information about the behavior of the solution, because it permits to apply a Harnack type inequality found in Trudinger [26].

Our results involving the quasilinear problem (PDE) are stated below

**Theorem 1.4.** Assume  $(\phi_1)$ - $(\phi_2)$ ,  $(V_1)$ - $(V_3)$ ,  $\epsilon = 1$  and that A belongs to Class 1 or 2. Then equation (PDE) has a heteroclinic solution from  $\alpha$  to  $\beta$  in  $C^{1,\gamma}_{loc}(\mathbb{R}^2)$  for some  $\gamma \in (0,1)$  such that

- (a) u(x,y) = u(x,y+1) for any  $(x,y) \in \mathbb{R}^2$ .
- (b)  $\alpha \leq u(x,y) \leq \beta$  for all  $(x,y) \in \mathbb{R}^2$ .

Moreover, taking into account the assumptions  $(\phi_3)$  and

 $(V_7)$  There are  $d_1, d_2, d_3, d_4 > 0$  and  $\lambda > 0$  such that

$$V'(t) \leq d_1 \phi(d_2|t-\beta|)|t-\beta|$$
 for all  $t \in [\beta - \lambda, \beta + \lambda]$ 

and

$$|V'(t)| \le d_3\phi(d_4|t-\alpha|)|t-\alpha| \text{ for all } t \in [\alpha-\lambda,\alpha+\lambda],$$

then the inequalities in (b) are strict.

**Theorem 1.5.** Assume  $(\phi_1)$ - $(\phi_2)$ ,  $(V_1)$ - $(V_3)$  and that A belongs to Class 3. Then there is a constant  $\epsilon_0 > 0$  such that for each  $\epsilon \in (0, \epsilon_0)$  equation (PDE) has a heteroclinic solution from  $\alpha$  to  $\beta$  in  $C_{loc}^{1,\gamma}(\mathbb{R}^2)$  for some  $\gamma \in (0, 1)$  such that

(a) u(x,y) = u(x,y+1) for any  $(x,y) \in \mathbb{R}^2$ . (b)  $\alpha \le u(x,y) \le \beta$  for all  $(x,y) \in \mathbb{R}^2$ .

Moreover, assuming  $(\phi_3)$  and  $(V_7)$  we have that the inequalities in (b) are strict.

**Theorem 1.6.** Assume  $(\phi_1)$ - $(\phi_2)$ ,  $(V_1)$ - $(V_3)$  and  $(V_6)$  with  $\alpha = -\beta$ ,  $\epsilon = 1$  and that A belongs to Class 4. Also consider the following assumption

 $(V_8)$  There are  $\mu > 0$  and  $\theta \in (0, \beta)$  such that

$$\mu \Phi(|t - \beta|) \le V(t), \quad \forall t \in (\beta - \theta, \beta + \theta).$$

Then equation (PDE) possesses a heteroclinic solution u from  $-\beta$  to  $\beta$  in  $C^{1,\gamma}_{loc}(\mathbb{R}^2)$  for some  $\gamma \in (0,1)$  such that

- (a) u(x,y) = -u(-x,y) for any  $(x,y) \in \mathbb{R}^2$ . (b) u(x,y) = u(x,y+1) for all  $(x,y) \in \mathbb{R}^2$ .
- (c)  $0 \le u(x,y) \le \beta$  for any x > 0 and  $y \in \mathbb{R}$ .

Moreover, if  $(\phi_3)$  and  $(V_7)$  occur then the inequalities in (c) are strict.

Here it is worth mentioning that an example of potential V that satisfies the conditions  $(V_1)$ - $(V_3)$  and  $(V_6)$ - $(V_8)$  is given by

(1.8) 
$$V(t) = \Phi(|(t - \alpha)(t - \beta)|) \text{ for all } t \in \mathbb{R},$$

where  $\Phi$  is an N-function of the type (1.7) verifying  $(\phi_1)$ - $(\phi_2)$ , which was used in [6]. The reader is invited to see that the condition  $(V_7)$  is not necessary to prove the existence of heteroclinic solution from  $\alpha$  to  $\beta$  for (PDE), however it together with  $(\phi_3)$  are used to obtain more information about the behavior of the heteroclinic solution. Moreover, the classical case  $\Phi(t) = \frac{t^2}{2}$  corresponds to the Laplacian operator, and in this case, as we are considering a new class of functions A, we can rewrite Theorem 1.6 as follows

**Theorem 1.7.** Assume  $V \in C^2(\mathbb{R},\mathbb{R})$ ,  $(V_2)$ - $(V_3)$  and  $(V_5)$ - $(V_6)$  with  $\alpha = -\beta$ , and that A belongs to Class 4. Then equation (1.3) with  $\Omega = \mathbb{R}^2$  possesses a heteroclinic (classical) solution u from  $-\beta$  to  $\beta$  such that

(a) u(x, y) = -u(-x, y) for any  $(x, y) \in \mathbb{R}^2$ . (b) u(x, y) = u(x, y + 1) for all  $(x, y) \in \mathbb{R}^2$ . (c)  $0 < u(x, y) < \beta$  for any x > 0 and  $y \in \mathbb{R}$ .

Before ending this section, we do some comments about our results: First, Theorems 1.1, 1.2 and 1.3 complement the study carried out in [5] and [7], because in those articles the authors considered the one-dimensional equation (1.6), while in the present article we treat (E) in the whole plane and investigated the existence of a heteroclinic solution for (E) for other classes of functions A. Moreover, we believe that this is the first article where the existence of a heteroclinic solution for a prescribed mean curvature equation is addressed in whole  $\mathbb{R}^2$ . Secondly, Theorems 1.4 - 1.7 hold for all pair of real numbers ( $\alpha, \beta$ ) with  $\alpha < \beta$ . Furthermore, in Theorems 1.4 and 1.5 we can consider a variety of potentials as the prototypes (1.1) and (1.2), while in Theorem 1.6 the potential V must have a strong interaction with the N-function  $\Phi$ , see for example (1.8). Finally, we would also like to point out that Theorems 1.4 and 1.5 complement the results obtained in [3], because in that paper the author considered the Laplacian operator while here we considered a large class of quasilinear operators. Moreover, Theorem 1.7 brings new contributions to the classic equation (1.3), since a new class of functions A is considered.

We organize the rest of this work as follows. Motivated by [3, 4, 23], in Section 2 we present the variational framework of the quasilinear problems mentioned above, which will be useful for the next section, and moreover, the proofs of Theorems 1.4, 1.5 and 1.6 are given. Section 3 is dedicated to the proofs of Theorems 1.1, 1.2 and 1.3 that were supported by the study developed in [5, Section 3]. Finally, we write an Appendix A about some results involving Orlicz and Orlicz-Sobolev spaces for unfamiliar readers with the topic.

#### 2. EXISTENCE OF HETEROCLINIC SOLUTION FOR QUASILINEAR EQUATIONS

The goal of this section is to establish the existence of a heteroclinic solution from  $\alpha$  to  $\beta$  for (PDE) taking into account the case where A satisfies different geometric conditions. The proof of the existence of solution is given by a minimization argument. To formulate the minimization problem of this section, let us first consider the infinite strip  $\Omega = \mathbb{R} \times (0, 1)$  of  $\mathbb{R}^2$  and for each  $j \in \mathbb{Z}$  we define the functional  $a_j : W_{\text{loc}}^{1,\Phi}(\Omega) \to \mathbb{R} \cup \{+\infty\}$  by

$$a_j(w) = \iint_{\Omega_j} \mathcal{L}(w) dx dy, \quad w \in W^{1,\Phi}_{\text{loc}}(\Omega),$$

where  $\Omega_j = (j, j+1) \times (0, 1)$  and

$$\mathcal{L}(w) = \Phi(|\nabla w|) + A(\epsilon x, y)V(w)$$

Under this notation, we also define the energy functional  $I: W^{1,\Phi}_{\text{loc}}(\Omega) \to \mathbb{R} \cup \{+\infty\}$  by

$$I(w) = \sum_{j \in \mathbb{Z}} a_j(w), \quad w \in W^{1,\Phi}_{\text{loc}}(\Omega).$$

In what follows, for each  $k \in \mathbb{Z}$  and  $w \in W^{1,\Phi}_{\text{loc}}(\Omega)$  we denote

$$\tau_k w(x,y) = w(x+k,y)$$
 for all  $(x,y) \in \Omega$ .

Clearly,  $\tau_0 w \equiv w$  on  $\Omega$ . Hereafter, let us identify  $\tau_k w|_{\Omega_0}$  with  $\tau_k w$  itself. Now, for the purposes of this paper, we will designates by  $\Gamma_{\Phi}(\alpha, \beta)$  the class of admissible functions given by (2.1)

$$\Gamma_{\Phi}(\alpha,\beta) = \left\{ w \in W^{1,\Phi}_{\text{loc}}(\Omega) : \tau_k w \to \alpha \text{ in } L^{\Phi}(\Omega_0) \text{ as } k \to -\infty \text{ and } \tau_k w \to \beta \text{ in } L^{\Phi}(\Omega_0) \text{ as } k \to +\infty \right\}.$$

We would like to point out that, as  $\Phi$  satisfies  $\Delta_2$ -condition,  $\tau_k w$  goes to  $\alpha$  in  $L^{\Phi}(\Omega_0)$  as k goes to  $-\infty$  if, and only if,

$$\iint_{\Omega_k} \Phi(|w-\alpha|) dx dy \to 0 \text{ as } k \to -\infty.$$

Analogously,  $\tau_k w$  goes to  $\beta$  in  $L^{\Phi}(\Omega_0)$  as k goes to  $+\infty$  if, and only if,

$$\iint_{\Omega_k} \Phi(|w-\beta|) dx dy \to 0 \text{ as } k \to +\infty.$$

On the other hand, it is easy to check that the class  $\Gamma_{\Phi}(\alpha,\beta)$  is not empty, because the function  $\varphi_{\alpha,\beta}: \Omega \to \mathbb{R}$  defined by

(2.2) 
$$\varphi_{\alpha,\beta}(x,y) = \begin{cases} \beta, & \text{if } \beta \leq x & \text{and } y \in (0,1), \\ x, & \text{if } \alpha \leq x \leq \beta & \text{and } y \in (0,1), \\ \alpha, & \text{if } x \leq \alpha & \text{and } y \in (0,1) \end{cases}$$

belongs to  $\Gamma_{\Phi}(\alpha, \beta)$ . By the properties of  $\Phi$ , A and V,

$$a_j(w) \ge 0$$
 for all  $j \in \mathbb{Z}$  and  $w \in \Gamma_{\Phi}(\alpha, \beta)$ ,

and hence, I is bounded from below on  $\Gamma_{\Phi}(\alpha, \beta)$ . Furthermore, it is easy to see that the function given in (2.2) has finite energy, that is,  $I(\varphi_{\alpha,\beta}) < +\infty$ , and so, the real number

$$c_{\Phi}(\alpha,\beta) = \inf_{w \in \Gamma_{\Phi}(\alpha,\beta)} I(w)$$

is well defined. Here it is worth mentioning that we will see throughout this section that critical points of the functional I on the class  $\Gamma_{\Phi}(\alpha, \beta)$  are heteroclinic solution from  $\alpha$  to  $\beta$  for the equation (*PDE*).

2.1. The case periodic. In this subsection, we intend to investigate the existence of a heteroclinic solution from  $\alpha$  to  $\beta$  for (PDE) with  $\epsilon = 1$  by assuming that A belongs to Class 1 and, unless indicated, the potential V satisfies the assumptions  $(V_1)$ - $(V_3)$ . With the preliminaries contained at the beginning of this section we may state and prove our first result that will be useful in the next lemma.

**Lemma 2.1.** If  $w \in \Gamma_{\Phi}(\alpha, \beta)$ , then for all  $k \in \mathbb{Z}$  we have that  $\tau_k w \in \Gamma_{\Phi}(\alpha, \beta)$  and  $I(\tau_k w) = I(w)$ .

*Proof.* Initially, it is easy to see that  $\tau_k w \in \Gamma_{\Phi}(\alpha, \beta)$  for any  $k \in \mathbb{Z}$  and  $w \in \Gamma_{\Phi}(\alpha, \beta)$ . On the other hand, for each  $j \in \mathbb{Z}$ , a simple change variable combined with the periodicity of A in the variable x leads to

$$\begin{aligned} a_j(\tau_k w) &= \iint_{\Omega_j} \left( \Phi(|\nabla \tau_k w|) + A(x, y) V(\tau_k w) \right) dx dy \\ &= \iint_{\Omega_j} \left( \Phi(|\nabla w(x+k, y)|) + A(x+k, y) V(w(x+k, y)) \right) dx dy \\ &= \iint_{\Omega_{j+k}} \left( \Phi(|\nabla w|) + A(x, y) V(w) \right) dx dy = a_{j+k}(w), \end{aligned}$$

from which it follows that

$$I(\tau_k w) = \sum_{j \in \mathbb{Z}} a_j(\tau_k w) = \sum_{j \in \mathbb{Z}} a_{j+k}(w) = \sum_{j \in \mathbb{Z}} a_j(w) = I(w),$$

and the proof is completed.

Now we employ the Lemma 2.1 to prove that the energy functional I reaches the minimum energy in some function of  $\Gamma_{\Phi}(\alpha, \beta)$ .

**Proposition 2.2.** There exists  $u \in \Gamma_{\Phi}(\alpha, \beta)$  such that  $I(u) = c_{\Phi}(\alpha, \beta)$  and  $\alpha \leq u(x, y) \leq \beta$  almost everywhere in  $\Omega$ .

*Proof.* Let  $(u_n)$  be a minimizing sequence for I on  $\Gamma_{\Phi}(\alpha, \beta)$ , that is,  $I(u_n) \to c_{\Phi}(\alpha, \beta)$  as  $n \to +\infty$ . Thus, there is a constant M > 0 verifying

(2.3) 
$$I(u_n) \le M \text{ for all } n \in \mathbb{N}.$$

We claim that we may assume without loss of generality that the sequence  $u_n$  satisfies

 $\alpha \leq u_n(x,y) \leq \beta$  for all  $(x,y) \in \Omega$  and  $n \in \mathbb{N}$ .

Indeed, just consider

 $\tilde{u}_n(x,y) = \max\{\alpha, \min\{u_n(x,y), \beta\}\}, \ (x,y) \in \Omega,$ 

instead of  $u_n$ . Moreover, we can also assume that for each  $n \in \mathbb{N}$ ,

(2.4) 
$$\iint_{\Omega_0} \Phi(|u_n - \alpha|) dx dy > \delta \text{ and } \iint_{\Omega_{k-1}} \Phi(|u_n - \alpha|) dx dy \le \delta \text{ for } k \le 0,$$

for some  $\delta > 0$  such that

$$\delta < \Phi(\beta - \alpha).$$

To establish this statement, it suffices to observe that

$$\liminf_{k \to +\infty} \iint_{\Omega_0} \Phi(|\tau_k u_n - \alpha|) dx dy > 0 = \lim_{k \to -\infty} \iint_{\Omega_0} \Phi(|\tau_k u_n - \alpha|) dx dy.$$

Indeed, note first that for each  $n \in \mathbb{N}$  fixed,

$$\tau_k u_n \to \beta$$
 in  $L^{\Phi}(\Omega_0)$  as  $k \to +\infty$ ,

and by  $\Phi \in \Delta_2$  (see for a moment (A.1)),

$$\iint_{\Omega_0} \Phi(|\tau_k u_n - \beta|) dx dy \to 0 \text{ as } k \to +\infty.$$

Thus, since  $\alpha \neq \beta$ , there are  $\delta > 0$  and a subsequence of  $(\tau_k u_n)_{k\geq 0}$ , still denoted  $(\tau_k u_n)$ , such that

$$\iint_{\Omega_0} \Phi(|\tau_k u_n - \alpha|) dx dy > \delta, \quad \forall k > 0.$$

Note that, without loss of generality, we may assume that  $\delta$  satisfies (2.5). On the other hand, using again the fact that  $\Phi \in \Delta_2$ ,  $\tau_k u_n$  goes to  $\alpha$  in  $L^{\Phi}(\Omega_0)$  as k goes to  $-\infty$  implies

$$\iint_{\Omega_0} \Phi(|\tau_k u_n - \alpha|) dx dy \to 0 \text{ as } k \to -\infty,$$

and so, there exists an integer  $\overline{k}_n < 0$  such that

$$\iint_{\Omega_0} \Phi(|\tau_k u_n - \alpha|) dx dy \le \delta, \quad \forall k \le \overline{k}_n.$$

From this, it is possible to find the bigger integer  $k_n \in \mathbb{Z}$  such that

$$\iint_{\Omega_{k-1}} \Phi(|u_n - \alpha|) dx dy \le \delta \text{ for all } k \le k_n \text{ and } \iint_{\Omega_{k_n}} \Phi(|u_n - \alpha|) dx dy > \delta,$$

that is,

$$\iint_{\Omega_{j-1}} \Phi(|\tau_{k_n} u_n - \alpha|) dx dy \le \delta \text{ for all } j \le 0 \text{ and } \iint_{\Omega_0} \Phi(|\tau_{k_n} u_n - \alpha|) dx dy > \delta$$

Now, we can apply Lemma 2.1 to consider  $\tau_{k_n} u_n$  in the place of  $u_n$ .

Now, since  $\alpha \leq u_n \leq \beta$  in  $\Omega$ , it is straightforward to check that  $(u_n)$  is bounded in  $W_{\text{loc}}^{1,\Phi}(\Omega)$ . Thereby, in view of Lemma A.4,  $W^{1,\Phi}(K)$  is reflexive Banach spaces whenever K is relatively compact in  $\Omega$ , and so, by a classical diagonal argument, there are a subsequence of  $(u_n)$ , still denoted by  $(u_n)$ , and  $u \in W_{\text{loc}}^{1,\Phi}(\Omega)$  satisfying

(2.6) 
$$u_n \rightharpoonup u \text{ in } W^{1,\Phi}_{\text{loc}}(\Omega) \text{ as } n \to +\infty,$$

(2.7) 
$$u_n \to u \text{ in } L^{\Phi}_{\text{loc}}(\Omega) \text{ as } n \to +\infty$$

and

(2.8) 
$$u_n(x,y) \to u(x,y)$$
 a.e. in  $\Omega$  as  $n \to +\infty$ .

As a consequence of (2.8),

(2.9) 
$$\alpha \le u(x,y) \le \beta$$
 almost everywhere in  $\Omega$ .

Moreover, from (2.3), we have the inequality below

$$\int_0^1 \int_{-j}^j \mathcal{L}(u_n) dx dy \le M, \quad \forall n, j \in \mathbb{N},$$

which combines with weak lower semicontinuity of I to give

$$\int_0^1 \int_{-j}^j \mathcal{L}(u) dx dy \le M, \quad \forall j \in \mathbb{N}.$$

Therefore, since  $j \in \mathbb{N}$  is arbitrary, we conclude that  $I(u) \leq M$ . With the aid of the previous preliminaries, our goal is to ensure that u belongs to  $\Gamma_{\Phi}(\alpha, \beta)$ . Towards that end, we will show that

(2.10) 
$$\tau_k u \to \alpha \text{ in } L^{\Phi}(\Omega_0) \text{ as } k \to -\infty.$$

To show (2.10), let us consider the sequence  $(\tau_k u)_{k\leq 0}$  with  $k \in \mathbb{Z}$ . Due to Lemma 2.1 and the estimate (2.9), it is simple to prove that  $(\tau_k u)_{k\leq 0}$  is bounded in  $W^{1,\Phi}(\Omega_0)$ . Consequently, for some subsequence, there exists  $u^* \in W^{1,\Phi}(\Omega_0)$  such that

(2.11) 
$$\tau_k u \rightharpoonup u^* \text{ in } W^{1,\Phi}(\Omega_0) \text{ as } k \to -\infty,$$

(2.12) 
$$\tau_k u \to u^* \text{ in } L^{\Phi}(\Omega_0) \text{ as } k \to -\infty$$

and

(2.13) 
$$\alpha \leq u^*(x,y) \leq \beta$$
 almost everywhere on  $\Omega_0$ .

Now, since  $I(u) \leq M$ , the definition of I ensures that

$$a_k(u) \to 0$$
 as  $|k| \to +\infty$ .

This together with the periodicity of A yields that

(2.14) 
$$a_0(\tau_k u) \to 0 \text{ as } |k| \to +\infty.$$

Now, the fact that  $a_0$  is weakly lower semicontinuous on  $W^{1,\Phi}(\Omega_0)$  and  $a_0 \ge 0$  together (2.11) and (2.14) guarantee that  $a_0(u^*) = 0$ . Thereby, (2.13) together with the assumptions on functions A and V ensures that  $u^* = \alpha$  or  $u^* = \beta$  a.e. in  $\Omega_0$ . On the other hand, it follows from (2.4) and (2.7) that

$$\iint_{\Omega_0} \Phi(|u-\alpha|) dx dy \ge \delta \text{ and } \iint_{\Omega_0} \Phi(|\tau_k u - \alpha|) dx dy \le \delta \text{ for } k < 0.$$

Consequently, taking the limit as  $k \to -\infty$  in the inequality above and employing (2.12), we arrive at

$$\iint_{\Omega_0} \Phi(|u^* - \alpha|) dx dy \le \delta$$

From (2.5), one has  $u^* = \alpha$  a.e. in  $\Omega_0$ , showing that the limit (2.10) is valid. Now we claim that

(2.15) 
$$\tau_k u \to \beta \text{ in } L^{\Phi}(\Omega_0) \text{ as } k \to +\infty.$$

Indeed, considering the sequence  $(\tau_k u)_{k>0}$  with  $k \in \mathbb{N}$ , there exist  $u^{**} \in W^{1,\Phi}(\Omega_0)$  and a subsequence of  $(\tau_k u)$ , still denoted  $(\tau_k u)$ , such that

(2.16) 
$$\tau_k u \rightharpoonup u^{**} \text{ in } W^{1,\Phi}(\Omega_0) \text{ as } k \to +\infty,$$

(2.17) 
$$\tau_k u \to u^{**} \text{ in } L^{\Phi}(\Omega_0) \text{ as } k \to +\infty,$$

(2.18) 
$$\tau_k u \to u^{**} \text{ in } L^1(\Omega_0) \text{ as } k \to +\infty$$

and

(2.19) 
$$\tau_k u(x,y) \to u^{**}(x,y)$$
 a.e in  $\Omega_0$  as  $k \to +\infty$ .

Arguing as above, we will get that  $u^{**} = \alpha$  or  $u^{**} = \beta$  a.e in  $\Omega_0$ . The claim (2.15) follows if we prove that  $u^{**} = \beta$  a.e in  $\Omega_0$ , and to do that, we will split the proof into two steps. So, seeking for a contradiction we assume that  $u^{**} = \alpha$  a.e. in  $\Omega_0$ .

**Step 1:** There are  $\epsilon_0 > 0$  and  $n_1 \in \mathbb{N}$  such that

(2.20) 
$$a_{-1}(u_n) + a_0(u_n) = \int_0^1 \int_{-1}^1 \left( \Phi(|\nabla u_n|) + A(x,y)V(u_n) \right) dxdy \ge \epsilon_0, \quad \forall n \ge n_1.$$

### HETEROCLINIC SOLUTIONS FOR SOME CLASSES OF PRESCRIBED MEAN CURVATURE EQUATIONS IN WHOLE R

Indeed, if this does not hold, then there is a subsequence  $(u_{n_i})$  of  $(u_n)$  such that

$$\int_0^1 \int_{-1}^1 \left( \Phi(|\nabla u_{n_i}|) + A(x, y) V(u_{n_i}) \right) dx dy \to 0.$$

Consequently, there is  $v \in W^{1,\Phi}((-1,1) \times (0,1))$  such that

$$u_{n_i} \rightharpoonup v$$
 in  $W^{1,\Phi}((-1,1) \times (0,1)), \quad u_{n_i} \to v$  in  $L^{\Phi}((-1,1) \times (0,1))$ 

and

(2.21) 
$$v = \alpha$$
 or  $v = \beta$  a.e. in  $(-1, 1) \times (0, 1)$ 

Making a simple analysis of the estimates contained in (2.4), we infer that (2.21) is impossible, which ends this step.

To proceed to the next step, let us fix  $\tilde{\epsilon} \in (0, \epsilon_0/2)$  and  $n_0 \ge n_1$  such that

(2.22) 
$$I(u_n) \le c_{\Phi}(\alpha, \beta) + \frac{\tilde{\epsilon}}{2}, \quad \forall n \ge n_0,$$

where  $\epsilon_0$  and  $n_1$  were given in Step 1.

**Step 2:** There are  $k \in \mathbb{N}$  and  $n \ge n_0$  large enough satisfying

(2.23) 
$$a_0 \left( x(\tau_k u_n - \alpha) + \alpha \right) \le \frac{\tilde{\epsilon}}{2}$$

In order to show estimate (2.23), we will separately analyze the terms of the functional  $a_0$  that will be divided into four parts as follows:

**Part 1:** There exists  $k_0 \in \mathbb{N}$  such that for each  $k \ge k_0$  there is  $n(k) \ge n_0$  verifying

(2.24) 
$$\iint_{\Omega_0} A(x,y) V(\tau_k u_n) dx dy \le \frac{\tilde{\epsilon}}{24 \cdot 4^m}, \quad \forall n \ge n(k)$$

(2.25) 
$$\iint_{\Omega_0} A(x,y) V\left(x(\tau_k u_n - \alpha) + \alpha\right) dx dy \le \frac{\tilde{\epsilon}}{4}, \quad \forall n \ge n(k)$$

and

(2.26) 
$$\iint_{\Omega_0} \Phi(|\tau_k u_n - \alpha|) dx dy \le \frac{\tilde{\epsilon}}{12 \cdot 4^m}, \quad \forall n \ge n(k),$$

where m was given in  $(\phi_2)$ .

In fact, let us initially note that, since  $V \in C^1$  and  $\tau_k u(x, y) \in [\alpha, \beta]$  for any  $(x, y) \in \Omega_0$ , the Mean Value Theorem together with  $(V_2)$  gives us

$$V(\tau_k u), \ V(x(\tau_k u - \alpha) + \alpha) \le R|\tau_k u - \alpha|, \ \forall (x, y) \in \Omega_0,$$

for some R > 0. Consequently,

$$\iint_{\Omega_0} A(x,y) V(x(\tau_k u - \alpha) + \alpha) dx dy \le R \sup_{\Omega_0} A(x,y) \iint_{\Omega_0} |\tau_k u - \alpha| dx dy$$

and

$$\iint_{\Omega_0} A(x,y) V(\tau_k u) dx dy \le R \sup_{\Omega_0} A(x,y) \iint_{\Omega_0} |\tau_k u - \alpha| dx dy$$

Now, as we are assuming that  $u^{**} = \alpha$  a.e in  $\Omega_0$ , it follows from (2.18) that there is  $k_0 \in \mathbb{N}$  such that

(2.27) 
$$\iint_{\Omega_0} A(x,y) V(x(\tau_k u - \alpha) + \alpha) dx dy \le \frac{\tilde{\epsilon}}{8}, \quad \forall k \ge k_0$$

and

(2.28) 
$$\iint_{\Omega_0} A(x,y) V(\tau_k u) dx dy \le \frac{\tilde{\epsilon}}{48 \cdot 4^m}, \quad \forall k \ge k_0.$$

Furthermore, from (2.17), increasing  $k_0$  if necessary, one gets

(2.29) 
$$\iint_{\Omega_0} \Phi(|\tau_k u - \alpha|) dx dy \le \frac{\tilde{\epsilon}}{24 \cdot 8^m}, \quad \forall k \ge k_0.$$

On the other hand, for each  $k \in \mathbb{N}$  fixed, Lebesgue Dominated Convergence Theorem yields

$$\left| \iint_{\Omega_0} A(x,y) \left( V(x(\tau_k u_n - \alpha) + \alpha) - V(x(\tau_k u - \alpha) + \alpha) \right) dx dy \right| \to 0 \text{ as } n \to +\infty$$

and

$$\left| \iint_{\Omega_0} A(x,y) \left( V(\tau_k u_n) - V(\tau_k u) \right) dx dy \right| \to 0 \text{ as } n \to +\infty.$$

Moreover, as  $\Phi \in \Delta_2$ , for each  $k \in \mathbb{N}$  we can use the limit (2.7) to find

$$\iint_{\Omega_0} \Phi(|\tau_k u_n - \tau_k u|) dx dy \to 0 \text{ as } n \to +\infty.$$

With everything, for every  $k \ge k_0$  there exists  $n(k) \ge n_0$  satisfying (2.30)

$$\iint_{\Omega_0} A(x,y) V(x(\tau_k u_n - \alpha) + \alpha) dx dy \le \iint_{\Omega_0} A(x,y) V(x(\tau_k u - \alpha) + \alpha) dx dy + \frac{\tilde{\epsilon}}{8}, \quad \forall n \ge n(k),$$

(2.31) 
$$\iint_{\Omega_0} A(x,y)V(\tau_k u_n)dxdy \le \iint_{\Omega_0} A(x,y)V(\tau_k u)dxdy + \frac{\tilde{\epsilon}}{48 \cdot 4^m}, \quad \forall n \ge n(k),$$

and

(2.32) 
$$\iint_{\Omega_0} \Phi(|\tau_k u_n - \tau_k u|) dx dy \le \frac{\tilde{\epsilon}}{24 \cdot 8^m}, \quad \forall n \ge n(k).$$

Finally, analyzing all the estimates, a direct use of (2.27)-(2.28) and (2.30)-(2.31) lead to (2.24) and (2.25). To see the inequality (2.26), note that from Lemma A.5-(a) one gets

$$\Phi(|\tau_k u_n - \alpha|) \le 2^m \left(\Phi(|\tau_k u_n - \tau_k u|) + \Phi(|\tau_k u - \alpha|)\right).$$

Now, (2.26) follows from (2.29) and (2.32), finishing the first part. **Part 2:** There are  $k \ge k_0$  and  $n \ge n(k)$  such that

(2.33) 
$$a_0\left(\tau_k u_n\right) \le \frac{\tilde{\epsilon}}{12 \cdot 4^m}$$

If the estimate above does not occur, for any  $k \ge k_0$  one has

$$a_0(\tau_k u_j) > \frac{\tilde{\epsilon}}{12 \cdot 4^m}, \quad \forall n \ge n(k).$$

Now let  $p \in \mathbb{N}$  be such that

$$(p+1)\frac{\tilde{\epsilon}}{12\cdot 4^m} > M,$$

where M was given in (2.3). Fixing  $i \in \mathbb{N}$  such that  $i > \max\{n(k) : k_0 \le k \le k_0 + p\}$  we have

$$I(u_i) \ge \sum_{t=k_0}^{k_0+p} a_t(u_i) = \sum_{t=k_0}^{k_0+p} a_0(\tau_t u_i) \ge (p+1)\frac{\tilde{\epsilon}}{12 \cdot 4^m} > M,$$

which contradicts (2.3), showing (2.33).

**Part 3:** For k and n as in Part 2, one has

(2.34) 
$$\iint_{\Omega_0} \Phi(|\partial_x(\tau_k u_n)|) dx dy, \quad \iint_{\Omega_0} \Phi(|\partial_y(\tau_k u_n)|) dx dy \le \frac{\tilde{\epsilon}}{12 \cdot 4^m}.$$

Indeed, just notice that the inequality at (2.33) together with the facts that  $\Phi$  is increasing on  $[0, +\infty)$  and

$$|\partial_x(\tau_k u_n)|, |\partial_y(\tau_k u_n)| \le |\nabla(\tau_k u_n)|,$$

leads to estimate (2.34).

**Part 4:** For k and n as in Part 2, one has

(2.35) 
$$\iint_{\Omega_0} \Phi(|\nabla(x(\tau_k u_n - \alpha) + \alpha)|) dx dy \le \frac{\tilde{\epsilon}}{4}.$$

To show the estimate (2.35), we first observe that

$$\partial_x \left( x(\tau_k u_n - \alpha) + \alpha \right) = \tau_k u_n - \alpha + x \partial_x (\tau_k u_n)$$

and

$$\partial_y \left( x(\tau_k u_n - \alpha) + \alpha \right) = x \partial_y \left( \tau_k u_n \right).$$

Therefore, from Lemma A.5-(a),

$$\Phi(|\nabla(x(\tau_k u_n - \alpha) + \alpha)|) \le 4^m \left(\Phi(|\partial_x(\tau_k u_n)|) + \Phi(|\tau_k u_n - \alpha|) + \Phi(|\partial_y(\tau_k u_n)|)\right) \text{ on } \Omega_0,$$

and the Part 4 follows from Parts 1 and 3.

Finally, the estimate (2.23) contained in Step 2 is immediately verified from (2.25) and (2.35). We are now ready to use Steps 1 and 2 to complete the proof of Claim (2.15). To this end, fix k and n as in Step 2 and define the following function

$$U_n(x,y) = \begin{cases} \alpha, & \text{if } x \le k & \text{and } y \in (0,1), \\ (u_n(x,y) - \alpha)(x-k) + \alpha, & \text{if } k \le x \le k+1 & \text{and } y \in (0,1), \\ u_n(x,y), & \text{if } x > k+1 & \text{and } y \in (0,1). \end{cases}$$

So, it is clear that  $U_n \in \Gamma_{\Phi}(\alpha, \beta)$  and

$$a_k(U_n) = a_0(x(\tau_k u_n - \alpha) + \alpha).$$

Hence,

$$c_{\Phi}(\alpha,\beta) \le I(U_n) = a_k(U_n) + \sum_{j=k+1}^{+\infty} a_j(u_n) \le a_k(U_n) + I(u_n) - a_0(u_n) - a_{-1}(u_n),$$

that is,

 $c_{\Phi}(\alpha,\beta) \leq a_0(x(\tau_k u_n - \alpha) + \alpha) + I(u_n) - a_0(u_n) - a_{-1}(u_n).$ Invoking estimates (2.20), (2.22) and (2.23), one gets

$$(2.20)$$
,  $(2.22)$  and  $(2.25)$ , one gets

$$c_{\Phi}(\alpha,\beta) \leq \frac{\epsilon}{2} + c_{\Phi}(\alpha,\beta) + \frac{\epsilon}{2} - \epsilon_0 = \tilde{\epsilon} + c_{\Phi}(\alpha,\beta) - \epsilon_0 < c_{\Phi}(\alpha,\beta) - \frac{\epsilon_0}{2},$$

which is absurd. Therefore, (2.15) occurs and  $u \in \Gamma_{\Phi}(\alpha, \beta)$ . To conclude the proof, it remains to show that  $I(u) = c_{\Phi}(\alpha, \beta)$ . For this purpose, given  $\epsilon > 0$  there exists  $n_0 \in \mathbb{N}$  such that

$$\sum_{-j}^{j} a_k(u_n) \le c_{\Phi}(\alpha, \beta) + \epsilon, \ \forall n \ge n_0 \text{ and } \forall j \in \mathbb{N}.$$

Letting  $n \to +\infty$  and after  $j \to +\infty$ , we find

$$I(u) \le c_{\Phi}(\alpha, \beta) + \epsilon.$$

Since  $\epsilon$  is arbitrary, we derive that  $I(u) = c_{\Phi}(\alpha, \beta)$ , and the proof is completed.

In order to find a periodic solution u(x, y) in the variable y for the equation (PDE), we will consider the following class

 $K_{\Phi}(\alpha,\beta) = \{ u \in \Gamma_{\Phi}(\alpha,\beta) : I(u) = c_{\Phi}(\alpha,\beta), u(x,0) = u(x,1) \text{ in } \mathbb{R}, \alpha \le u \le \beta \text{ a.e. on } \Omega \}.$ 

Next, we are going to show that  $K_{\Phi}(\alpha, \beta)$  is not empty.

**Lemma 2.3.** It holds that  $K_{\Phi}(\alpha, \beta) \neq \emptyset$ .

*Proof.* Initially, for each  $w \in \Gamma_{\Phi}(\alpha, \beta)$  we set

$$I_1(w) = \iint_{\mathbb{R} \times (0,\frac{1}{2})} \mathcal{L}(w) dx dy \text{ and } I_2(w) = \iint_{\mathbb{R} \times (\frac{1}{2},1)} \mathcal{L}(w) dx dy$$

Now, choosing  $u \in \Gamma_{\Phi}(\alpha, \beta)$  as in Proposition 2.2, we can write

(2.36) 
$$I(u) = I_1(u) + I_2(u) = c_{\Phi}(\alpha, \beta)$$

Suppose for a moment that  $I_1(u) \leq I_2(u)$  holds. Then considering the function

$$v(x,y) = \begin{cases} u(x,y), & \text{if } x \in \mathbb{R} \text{ and } 0 \le y \le \frac{1}{2}, \\ u(x,1-y), & \text{if } x \in \mathbb{R} \text{ and } \frac{1}{2} \le y \le 1, \end{cases}$$

it is clear that  $v \in \Gamma_{\Phi}(\alpha, \beta)$  and thanks to the assumptions  $(A_2)$ - $(A_3)$  a straightforward computation gives

(2.37) 
$$I_2(v) = I_1(v) = I_1(u).$$

According to (2.36) and (2.37),

$$c_{\Phi}(\alpha,\beta) \le I(v) = I_1(v) + I_2(v) \le I(u) = c_{\Phi}(\alpha,\beta),$$

from where it follows that  $I(v) = c_{\Phi}(\alpha, \beta)$  with v(x, 0) = v(x, 1) for every  $x \in \mathbb{R}$  and  $v(x, y) \in [\alpha, \beta]$  a.e. in  $\Omega$ . On the other hand, if  $I_2(u) \leq I_1(u)$  occurs then in this case we define the function

$$\tilde{v}(x,y) = \begin{cases} u(x,1-y), & \text{if } x \in \mathbb{R} \text{ and } 0 \le y \le \frac{1}{2} \\ u(x,y), & \text{if } x \in \mathbb{R} \text{ and } \frac{1}{2} \le y \le 1. \end{cases}$$

Consequently, using the same ideas discussed just above, we obtain that  $\tilde{v} \in \Gamma_{\Phi}(\alpha, \beta)$  with  $I_1(\tilde{v}) = I_2(\tilde{v}) = I_2(u)$ , from where it follows that  $I(\tilde{v}) = c_{\Phi}(\alpha, \beta)$ ,  $\tilde{v}(x, 0) = \tilde{v}(x, 1)$  for any  $x \in \mathbb{R}$  and  $\alpha \leq \tilde{v} \leq \beta$  a.e in  $\Omega$ , which completes the proof.

We would like to emphasize here that the functions of  $K_{\Phi}(\alpha, \beta)$  can be extended periodically in the variable y on  $\mathbb{R}^2$  with period 1. For this reason, it will be convenient to assume that the elements of  $K_{\Phi}(\alpha, \beta)$  are extended to the whole real plane. Finally, we will show our best result of this section.

**Lemma 2.4.** If  $u \in K_{\Phi}(\alpha, \beta)$ , then u is a weak solution of (PDE) with  $\epsilon = 1$ . Moreover, u is a heteroclinic solution from  $\alpha$  to  $\beta$  which belongs to  $C_{loc}^{1,\gamma}(\mathbb{R}^2)$  for some  $\gamma \in (0,1)$ .

*Proof.* Initially, let be  $u \in K_{\Phi}(\alpha, \beta)$  and  $\psi \in C_0^{\infty}(\mathbb{R}^2)$ . Then, the function  $v = u + t\psi \in \Gamma_{\Phi}(\alpha, \beta)$  and

$$\iint_{\Omega} \left( \phi(|\nabla u|) \nabla u \nabla \psi + A(x, y) V'(u) \psi \right) dx dy = 0$$

The equality above allows us to use the same arguments found in the proof [4, Theorem 1.1] to prove that

$$\iint_{\mathbb{R}^2} \left( \phi(|\nabla u|) \nabla u \nabla \psi + A(x, y) V'(u) \psi \right) dx dy = 0, \quad \forall \psi \in C_0^\infty(\mathbb{R}^2),$$

from where it follows that u is a weak solution for (PDE) with  $\epsilon = 1$ . The assumption  $(\phi_2)$  permits to apply a well known regularity result developed by Lieberman [17, Theorem 1.7] to conclude that  $u \in C^{1,\gamma}_{\text{loc}}(\mathbb{R}^2)$  for some  $\gamma \in (0,1)$ . Moreover, similar to the proof of [4, Theorem 1.1], we also have that u is a heteroclinic solution from  $\alpha$  to  $\beta$ , that is,

 $u(x,y) \to \alpha$  as  $x \to -\infty$  and  $u(x,y) \to \beta$  as  $x \to +\infty$ , uniformly in  $y \in \mathbb{R}$ ,

and the lemma follows.

Now, we will show our last result in this subsection, which ends the study of the equation (PDE) in the case where A is periodic in all variables.

# **Lemma 2.5.** Assume $(\phi_3)$ and $(V_7)$ . Then, if $u \in K_{\Phi}(\alpha, \beta)$ we have that $\alpha < u(x, y) < \beta$ for all $(x, y) \in \mathbb{R}^2$ .

Proof. Let  $u \in K_{\Phi}(\alpha, \beta)$  and observe that  $\alpha \leq u \leq \beta$  on  $\mathbb{R}^2$ . In what follows, we will show that  $u(x, y) < \beta$  for any  $(x, y) \in \mathbb{R}^2$ . Indeed, assume for the sake of contradiction that there exists  $(x_0, y_0) \in \mathbb{R}^2$  such that  $u(x_0, y_0) = \beta$ . Therefore, by the geometry of u, we can consider a compact set  $\mathcal{O}$  contained in  $\mathbb{R}^2$  such that there exists  $(x_1, y_1) \in \mathcal{O}$  with  $u(x_1, y_1) < \beta$ . Having that in mind, setting the function  $\tilde{\phi} : (0, +\infty) \to (0, +\infty)$  by

$$\tilde{\phi}(t) = \begin{cases} \phi(t), & \text{if } t \in (0, R], \\ \frac{\phi(R)t^{q-2}}{R^{q-2}}, & \text{if } t \in (R, +\infty), \end{cases}$$

where  $R > \max\{\|\nabla u\|_{L^{\infty}(\mathcal{O})}, \delta\}$  and the constants q and  $\delta$  were given in  $(\phi_3)$ , a direct computation implies that there are positive real numbers  $\gamma_1$  and  $\gamma_2$ , which dependent on  $\delta$ , q, R,  $c_1$  and  $c_2$ , such that

$$\tilde{\phi}(t)t \leq \gamma_1 t^{q-1}$$
 and  $\tilde{\phi}(t)t^2 \geq \gamma_2 t^q$  for all  $t \geq 0$ .

Using the function  $\tilde{\phi}$ , let us define the vector measurable function  $G: \mathbb{R}^2 \times \mathbb{R} \times \mathbb{R}^2 \to \mathbb{R}^2$  by

$$G(z,t,p) = \frac{\tilde{\phi}(|p|)p}{\gamma_2},$$

which satisfies

$$|G(z,t,p)| \leq \frac{\gamma_1}{\gamma_2} |p|^{q-1} \text{ and } pG(z,t,p) \geq |p|^q \text{ for all } (z,t,p) \in \mathbb{R}^2 \times \mathbb{R} \times \mathbb{R}^2.$$

Furthermore, we will also consider the scalar measurable function  $B: \mathbb{R}^2 \times \mathbb{R} \times \mathbb{R}^2 \to \mathbb{R}$  given by

$$B(z,t,p) = \frac{A(z)V'(\beta - t)}{\gamma_2}.$$

Now, combining  $(\phi_3)$  with  $(V_7)$ , it is possible to ensure that for each M > 0 there exists  $C_M > 0$  satisfying

$$|B(z,t,p)| \le C_M |t|^{q-1}$$
 for all  $(z,t,p) \in \mathbb{R}^2 \times (-M,M) \times \mathbb{R}^2$ .

All these information are necessary to guarantee that G and B fulfill the structure required in the Harnack type inequality found in Trudinger [26, Theorem 1.1]. So, setting  $v(z) = \beta - u(z)$ for  $z \in \mathbb{R}^2$ , we infer that v is a weak solution of the quasilinear equation

$$div \ G(z, v, \nabla v) + B(z, v, \nabla v) = 0 \quad \text{in } \quad \mathcal{O}.$$

Employing [26, Theorem 1.1], we deduce that v = 0 on  $\mathcal{O}$ , that is,  $u = \beta$  on  $\mathcal{O}$ , which contradicts the fact that  $(x_1, y_1) \in \mathcal{O}$  with  $u(x_1, y_1) < \beta$ . Likewise, we can apply a similar argument to show that  $u(x, y) > \alpha$  for any  $(x, y) \in \mathbb{R}^2$ , and hence the proof is completed.  $\Box$  2.2. The case asymptotic at infinity to a periodic function. In this subsection we will study the existence of a heteroclinic solution for (PDE) with  $\epsilon = 1$  and A belongs to Class 2, that is, A is asymptotic at infinity to a periodic function  $A_p$ . Moreover, unless otherwise indicated, we will consider here the conditions  $(\phi_1)$ - $(\phi_2)$  on  $\phi$  and  $(V_1)$ - $(V_3)$  on V. The fact that we are assuming that the function A is only asymptotically periodic with respect to x brings a lot of difficulties and some arguments explored in the periodic case do not work anymore.

In this section, let us consider the functional  $I_p: W^{1,\Phi}_{\text{loc}}(\Omega) \to \mathbb{R} \cup \{+\infty\}$  by

$$I_p(w) = \sum_{j \in \mathbb{Z}} a_{p,j}(w), \quad w \in W^{1,\Phi}_{\text{loc}}(\Omega).$$

where

$$a_{p,j}(w) = \iint_{\Omega_j} \left( \Phi(|\nabla w|) + A_p(x,y)V(w) \right) dxdy.$$

Moreover, we use  $c_{p,\Phi}(\alpha,\beta)$  to denote the real number given by

$$c_{p,\Phi}(\alpha,\beta) = \inf_{w \in \Gamma_{\Phi}(\alpha,\beta)} I_p(w).$$

From Subsection 2.1, we know that there is  $w_0 \in \Gamma_{\Phi}(\alpha, \beta)$  such that  $I_p(w_0) = c_{p,\Phi}(\alpha, \beta)$ , and so,

(2.38) 
$$c_{\Phi}(\alpha,\beta) \leq I(w_0) < I_p(w_0) = c_{p,\Phi}(\alpha,\beta).$$

The inequality (2.38) establishes an important relation between  $c_{\Phi}(\alpha, \beta)$  and  $c_{p,\Phi}(\alpha, \beta)$ , which will be useful to achieve the objective of this subsection. With these information, we are ready to prove the main result of this subsection.

**Proposition 2.6.** There is  $u \in \Gamma_{\Phi}(\alpha, \beta)$  such that  $I(u) = c_{\Phi}(\alpha, \beta)$  satisfying  $\alpha \leq u(x, y) \leq \beta$  almost everywhere in  $\Omega$ .

*Proof.* First of all, note that there exists a minimizing sequence  $(u_n) \subset \Gamma_{\Phi}(\alpha, \beta)$  for I satisfying

$$\alpha \leq u_n(x,y) \leq \beta, \ \forall n \in \mathbb{N} \text{ and } (x,y) \in \Omega.$$

Moreover, there are  $u \in W^{1,\Phi}_{\text{loc}}(\Omega)$  and a subsequence of  $(u_n)$ , still denoted by  $(u_n)$ , such that

(2.39) 
$$u_n \rightharpoonup u \text{ in } W^{1,\Phi}_{\text{loc}}(\Omega),$$

(2.40) 
$$u_n \to u \text{ in } L^{\Phi}_{\text{loc}}(\Omega)$$

(2.41) 
$$u_n(x,y) \to u(x,y)$$
 a.e. in  $\Omega$ .

From (2.39)-(2.41),

$$(2.42) I(u) \le c_{\Phi}(\alpha, \beta)$$

and

$$\alpha \leq u(x,y) \leq \beta$$
 a.e. in  $\Omega$ 

Now our goal is to show that  $u \in \Gamma_{\Phi}(\alpha, \beta)$ . To achieve this goal, similar to the proof of Proposition 2.2, we have that  $(\tau_k u)_{k>0}$  is a bounded sequence in  $W^{1,\Phi}(\Omega_0)$ . Thereby, for some subsequence of  $(\tau_k u)$ , still denoted by itself, there is  $u^* \in W^{1,\Phi}(\Omega_0)$  such that

$$\tau_k u \to u^* \text{ in } W^{1,\Phi}(\Omega_0) \text{ as } k \to +\infty,$$
  
 $\tau_k u \to u^* \text{ in } L^{\Phi}(\Omega_0) \text{ as } k \to +\infty$ 

and

$$\tau_k u(x,y) \to u^*(x,y)$$
 a.e on  $\Omega_0$  as  $k \to +\infty$ 

We claim that  $u^* = \alpha$  or  $u^* = \beta$  a.e. in  $\Omega_0$ . Indeed, since  $I(u) \leq c_{\Phi}(\alpha, \beta)$ , we infer that  $a_k(u)$  goes to 0 as k goes to  $+\infty$ , and so, by change of variable,

$$\iint_{\Omega_0} \left( \Phi(|\nabla \tau_k u|) + A(x+k,y)V(\tau_k u) \right) dxdy \to 0 \text{ as } k \to +\infty$$

Consequently, by  $(A_1)$ ,

$$\iint_{\Omega_0} \left( \Phi(|\nabla \tau_k u|) + A_0 V(\tau_k u) \right) dx dy \to 0 \text{ as } k \to +\infty,$$

from where it follows that

$$\iint_{\Omega_0} \left( \Phi(|\nabla u^*|) + A_0 V(u^*) \right) dx dy = 0.$$

By the assumptions on  $\Phi$  and V, we derive that  $u^* = \alpha$  or  $u^* = \beta$  a.e. in  $\Omega_0$ . Next, we claim that

(2.43) 
$$u^* = \beta \text{ a.e in } \Omega_0.$$

To establish the claim above, let us assume by contradiction that  $u^* = \alpha$  a.e. in  $\Omega_0$ . So, as a consequence, we will prove that given  $\delta \in (0, \Phi(\beta - \alpha))$  there are  $(u_{n_i}) \subset (u_n), (k_i) \subset \mathbb{N}$  and  $i_* \in \mathbb{N}$  such that

(2.44) 
$$i_* < k_i \text{ for all } i \in \mathbb{N}, k_i \to +\infty \text{ and } n_i \to +\infty \text{ as } i \to +\infty,$$

(2.45) 
$$\iint_{\Omega_0} \Phi(|\tau_j u_{n_i} - \alpha|) dx dy < \delta \text{ and } \iint_{\Omega_0} \Phi(|\tau_{k_i} u_{n_i} - \alpha|) dx dy \ge \delta \ \forall j \in [i_*, k_i - 1] \cap \mathbb{N}.$$

Indeed, since  $\tau_k u$  goes to  $\alpha$  in  $L^{\Phi}(\Omega_0)$  as k goes to  $+\infty$ , given  $\delta \in (0, \Phi(\beta - \alpha))$ , there is  $i_* = i_*(\delta) \in \mathbb{N}$  satisfying

(2.46) 
$$\iint_{\Omega_0} \Phi(|\tau_k u - \alpha|) dx dy < \frac{\delta}{2^{m+1}} \quad \forall k \ge i_*.$$

In particular,

(2.47) 
$$\iint_{\Omega_0} \Phi(|\tau_{i_*}u - \alpha|) dx dy < \frac{\delta}{2^{m+1}}.$$

Gathering (2.40) and (2.47) with the fact that  $\Phi \in \Delta_2$ , we find  $n_1 \in \mathbb{N}$  satisfying

$$\iint_{\Omega_0} \Phi(|\tau_{i_*} u_{n_1} - \alpha|) dx dy < \delta.$$

Thereby, since  $u_{n_1} \in \Gamma_{\Phi}(\alpha, \beta)$ , we may fix  $k_1 \ge i_* + 1$  as the first natural number such that

$$\iint_{\Omega_0} \Phi(|\tau_j u_{n_1} - \alpha|) dx dy < \delta \text{ and } \iint_{\Omega_0} \Phi(|\tau_{k_1} u_{n_1} - \alpha|) dx dy \ge \delta \quad \forall j \in [i_*, k_1 - 1] \cap \mathbb{N}.$$

On the other hand, according to (2.46),

$$\iint_{\Omega_0} \Phi(|\tau_{i_*}u - \alpha|) dx dy, \quad \iint_{\Omega_0} \Phi(|\tau_{i_*+1}u - \alpha|) dx dy < \frac{\delta}{2^{m+1}}$$

Hence, in the same manner we can see that there is  $n_2 \in \mathbb{N}$  such that  $n_2 > n_1$  and

$$\iint_{\Omega_0} \Phi(|\tau_{i_*}u_{n_2} - \alpha|) dx dy, \quad \iint_{\Omega_0} \Phi(|\tau_{i_*+1}u_{n_2} - \alpha|) dx dy < \delta.$$

Using the fact that  $u_{n_2} \in \Gamma_{\Phi}(\alpha, \beta)$ , we can find  $k_2 \ge i_* + 2$  as the first natural number satisfying

$$\iint_{\Omega_0} \Phi(|\tau_j u_{n_2} - \alpha|) dx dy < \delta \text{ and } \iint_{\Omega_0} \Phi(|\tau_{k_2} u_{n_2} - \alpha|) dx dy \ge \delta, \quad \forall j \in [i_*, k_2 - 1] \cap \mathbb{N}.$$

Repeating the above argument, there are sequences  $(u_{n_i}) \subset (u_n)$  and  $(k_i) \subset \mathbb{N}$  such that  $k_i \geq i_* + i$  satisfying (2.44) and (2.45). So, for some subsequence, there is  $w \in W^{1,\Phi}_{\text{loc}}(\Omega)$  such that

(2.48) 
$$\tau_{k_i} u_{n_i} \rightharpoonup w \text{ in } W^{1,\Phi}_{\text{loc}}(\Omega) \text{ as } i \to +\infty,$$

(2.49) 
$$\tau_{k_i} u_{n_i} \to w \text{ in } L^{\Phi}_{\text{loc}}(\Omega) \text{ as } i \to +\infty,$$

(2.50) 
$$\tau_{k_i} u_{n_i}(x, y) \to w(x, y) \text{ a.e. in } \Omega \text{ as } i \to +\infty$$

and

(2.51) 
$$\alpha \le w(x,y) \le \beta$$
 a.e. in  $\Omega$ .

Now, setting the functional

$$a_j^i(v) = \iint_{\Omega_j} \left( \Phi(|\nabla v|) + A(x+k_i, y)V(v) \right) dxdy, \quad v \in W_{\text{loc}}^{1,\Phi}(\Omega), \ i \in \mathbb{N}, \ j \in \mathbb{Z},$$

a simple change of variables gives us

$$a_j^i(\tau_{k_i}u_{n_i}) = a_{j+k_i}(u_{n_i}),$$

and so,

(2.52) 
$$\sum_{j\in\mathbb{Z}}a_j^i(\tau_{k_i}u_{n_i}) = \sum_{j\in\mathbb{Z}}a_j(u_{n_i}) = I(u_{n_i}), \quad \forall i\in\mathbb{N}.$$

From (2.52), one has

(2.53) 
$$a_{p,0}(\tau_k w) \to 0 \text{ as } |k| \to +\infty.$$

To see this, it suffices to show that  $I_p(w) \leq c_{\Phi}(\alpha, \beta)$ . Indeed, combining the fact that  $A(x+k_i, y)$  goes to  $A_p(x, y)$  as *i* goes to  $+\infty$  with (2.50), one gets

$$A(x+k_i,y)V(\tau_{k_i}u_{n_i}(x,y)) \to A_p(x,y)V(w(x,y))$$
 a.e. in  $\Omega$ 

Therefore, the Fatou's Lemma and (2.48) provide

$$\sum_{-j}^{j} a_{p,j}(w) \le \liminf_{i \to +\infty} \sum_{-j}^{j} a_{j}^{i}(\tau_{k_{i}} u_{n_{i}}), \quad \forall j \in \mathbb{N}.$$

As j is arbitrary, (2.52) guarantees that

(2.54) 
$$I_p(w) \le \liminf_{i \to +\infty} I(u_{n_i}) = c_{\Phi}(\alpha, \beta),$$

and (2.53) is proved. Thereby, passing to a subsequence if necessary, a direct computation shows that

 $\tau_k w \to \alpha \text{ or } \beta \text{ in } L^{\Phi}(\Omega_0) \text{ as } k \to -\infty$ 

and

$$\tau_k w \to \alpha \text{ or } \beta \text{ in } L^{\Phi}(\Omega_0) \text{ as } k \to +\infty$$

Our goal now is to ensure that  $w \in \Gamma_{\Phi}(\alpha, \beta)$ . Claim 1:  $\tau_k w \to \alpha$  in  $L^{\Phi}(\Omega_0)$  as  $k \to -\infty$ . HETEROCLINIC SOLUTIONS FOR SOME CLASSES OF PRESCRIBED MEAN CURVATURE EQUATIONS IN WHOLE R3

Indeed, note first that for each  $j \in \mathbb{N}$ , there is  $i_0 = i_0(j) \in \mathbb{N}$  such that  $k_i - 1 \ge k_i - j \ge i_*$  for all  $i \ge i_0$ , where  $i_* \in \mathbb{N}$  was given in (2.44). According to (2.45),

$$\iint_{\Omega_0} \Phi(|\tau_{k_i-j}u_{n_i}-\alpha|) dx dy < \delta, \quad \forall j \in \mathbb{N},$$

that is,

$$\iint_{\Omega_{-j}} \Phi(|\tau_{k_i} u_{n_i} - \alpha|) dx dy < \delta, \quad \forall j \in \mathbb{N}.$$

Invoking (2.49), we can increase *i* if necessary to obtain

$$\iint_{\Omega_0} \Phi(|\tau_{-j}w - \alpha|) dx dy \le \delta, \ \forall j \in \mathbb{N}.$$

Our claim is proved by noting that  $\delta \in (0, \Phi(\beta - \alpha))$ .

Claim 2:  $\tau_k w \to \beta$  in  $L^{\Phi}(\Omega_0)$  as  $k \to +\infty$ .

Assume by contradiction that  $\tau_k w \to \alpha$  in  $L^{\Phi}(\Omega_0)$  as  $k \to +\infty$ . Let us break down the proof of Claim 2 into two steps.

**Step 1:** There are  $\epsilon_0 > 0$  and  $i_0 \in \mathbb{N}$  such that

(2.55) 
$$\int_{k_i-1}^{k_i+1} \int_0^1 \left( \Phi(|\nabla u_{n_i}|) + V(u_{n_i}) \right) dx dy \ge \frac{\epsilon_0}{\tilde{A}}, \quad \forall i \ge i_0,$$

where  $\hat{A} = \min\{1, A_0\}.$ 

Indeed, if this does not occur, then there is a subsequence  $(\tau_{k_i} u_{n_i})$  of  $(\tau_{k_i} u_{n_i})$  such that

$$\int_{-1}^{1} \int_{0}^{1} \left( \Phi(|\nabla \tau_{k_{i_j}} u_{n_{i_j}}|) + V(\tau_{k_{i_j}} u_{n_i}) \right) dx dy \to 0 \text{ as } j \to +\infty.$$

Recalling that  $(u_n)$  is bounded in  $W_{\text{loc}}^{1,\Phi}(\Omega)$ , then going to a subsequence if necessary, there exists  $v \in W^{1,\Phi}((-1,1) \times (0,1))$  such that

(2.56) 
$$\tau_{k_{i_j}} u_{n_{i_j}} \rightharpoonup v \text{ in } W^{1,\Phi}((-1,1) \times (0,1)) \text{ and } \tau_{k_{i_j}} u_{n_{i_j}} \rightarrow v \text{ in } L^{\Phi}((-1,1) \times (0,1)).$$

From the assumptions on  $\Phi$  and V, we have  $v = \alpha$  or  $v = \beta$  a.e. in  $(-1, 1) \times (0, 1)$ . On the other hand, from (2.45),

(2.57) 
$$\iint_{\Omega_{-1}} \Phi(|\tau_{k_i} u_{n_i} - \alpha|) dx dy < \delta \text{ and } \iint_{\Omega_0} \Phi(|\tau_{k_i} u_{n_i} - \alpha|) dx dy \ge \delta, \quad \forall i \in \mathbb{N}.$$

Finally, taking the limit of  $k_i \to +\infty$  in (2.57) and using the limit (2.56) we find a contradiction, finishing the proof of Step 1.

In what follows, fixing  $\epsilon \in (0, \frac{\epsilon_0}{2})$  and increasing  $i_0$  if necessary, we obtain

(2.58) 
$$I(u_{n_i}) \le c_{\Phi}(\alpha, \beta) + \frac{\epsilon}{4}, \quad \forall i \ge i_0.$$

**Step 2:** There exist  $j \in \mathbb{N}$  and  $i \geq i_0$  large enough satisfying

(2.59) 
$$a_{p,j}\left((\tau_{k_i}u_{n_i}-\alpha)(x-j)+\alpha\right) \le \frac{\epsilon}{2}$$

The proof of Step 2 follows as in the proof of Proposition 2.2, and so, it will be omitted. In the sequel, let us consider  $j \in \mathbb{N}$  and  $i \geq i_0$  as in Step 2. Setting the function

$$U_{j,i}(x,y) = \begin{cases} \alpha, & \text{if } x \leq j & \text{and } y \in (0,1), \\ (\tau_{k_i} u_{n_i}(x,y) - \alpha)(x-j) + \alpha, & \text{if } j \leq x \leq j+1 & \text{and } y \in (0,1), \\ \tau_{k_i} u_{n_i}(x,y), & \text{if } x > j+1 & \text{and } y \in (0,1), \end{cases}$$

it is simple to check that  $U_{j,i} \in \Gamma_{\Phi}(\alpha, \beta)$  and

$$(2.60) \quad c_{p,\Phi}(\alpha,\beta) \le I_p(U_{j,i}) = a_{p,j}(U_{j,i}) + \sum_{t=j+1}^{+\infty} a_{p,t}(\tau_{k_i}u_{n_i}) = a_{p,j}(U_{j,i}) + \sum_{t=j+1+k_i}^{+\infty} a_{p,t}(u_{n_i}).$$

We claim that increasing  $i_0$  if necessary, one gets

(2.61) 
$$\sum_{t=j+1+k_i}^{+\infty} a_{p,t}(u_{n_i}) \le \sum_{t=j+1+k_i}^{+\infty} a_t(u_{n_i}) + \frac{\epsilon}{4}$$

Indeed, since the function A belongs to Class 2, we infer that there is R > 0 such that

$$A_p(x,y) - A(x,y) \le \frac{A_0\epsilon}{4C}, \quad \forall |x| \ge R \text{ and } \forall y \in (0,1),$$

where C > 0 is a constant such that  $I(u_n) \leq C$  for all  $n \in \mathbb{N}$ . Consequently,

$$\int_{R}^{+\infty} \int_{0}^{1} \left( A_p(x,y) - A(x,y) \right) V(u_{n_i}) dx dy \le \frac{A_0 \epsilon}{4C} \int_{R}^{+\infty} \int_{0}^{1} V(u_{n_i}) dx dy \le \frac{\epsilon}{4},$$

and therefore, increasing i if necessary the last inequality is sufficient to justify (2.61). In view of (2.60) and (2.61), one has

(2.62) 
$$c_{p,\Phi}(\alpha,\beta) \le a_{p,j}(U_{j,i}) + \sum_{t=j+1+k_i}^{+\infty} a_t(u_{n_i}) + \frac{\epsilon}{4}.$$

On the other hand, according to Step 1,

$$\sum_{t=j+1+k_i}^{+\infty} a_t(u_{n_i}) + \epsilon_0 \le \sum_{t=j+1+k_i}^{+\infty} a_t(u_{n_i}) + \tilde{A} \int_{k_i-1}^{k_i+1} \int_0^1 \left(\Phi(|\nabla u_{n_i}|) + V(u_{n_i})\right) dxdy \le I(u_{n_i}),$$

which together with (2.62) yields that

$$c_{p,\Phi}(\alpha,\beta) \le a_{p,j}(U_{j,i}) + I(u_{n_i}) - \epsilon_0 + \frac{\epsilon}{4}.$$

This together with (2.58) leads to

$$c_{p,\Phi}(\alpha,\beta) \le a_{p,j}(U_{j,i}) + c_{\Phi}(\alpha,\beta) + \frac{\epsilon}{2} - \epsilon_0.$$

Recalling that  $\epsilon \in (0, \frac{\epsilon_0}{2})$  and using (2.59), we arrive at

$$c_{p,\Phi}(\alpha,\beta) \le c_{\Phi}(\alpha,\beta) + \epsilon - \epsilon_0 < c_{\Phi}(\alpha,\beta) - \frac{\epsilon_0}{2},$$

contradicting (2.38). This proves the Claim 2.

Finally, by virtue of Claims 1 and 2, we infer that  $w \in \Gamma_{\Phi}(\alpha, \beta)$ . Furthermore, from (2.54) we also have

$$c_{p,\Phi}(\alpha,\beta) \le I_p(w) \le c_{\Phi}(\alpha,\beta)$$

obtaining a new contradiction, and our claim (2.43) is proved. As a byproduct,

(2.63) 
$$\tau_k u \to \beta \text{ in } L^{\Phi}(\Omega_0) \text{ as } k \to +\infty$$

A similar argument works to prove that

(2.64) 
$$\tau_k u \to \alpha \text{ in } L^{\Phi}(\Omega_0) \text{ as } k \to -\infty.$$

Combining (2.63) and (2.64) with (2.42) we get precisely the assertion of the proposition.  $\Box$ 

Considering here  $K_{\Phi}(\alpha, \beta)$  as described in Subsection 2.1, the same argument explored in that subsection guarantees that  $K_{\Phi}(\alpha, \beta)$  is a non-empty set and allows us to write the following result.

**Lemma 2.7.** There exists a weak solution u of (PDE) with  $\epsilon = 1$ , such that  $u \in K_{\Phi}(\alpha, \beta) \cap C^{1,\gamma}_{loc}(\mathbb{R}^2)$  for some  $\gamma \in (0,1)$  and it is heteroclinic solution from  $\alpha$  to  $\beta$  and 1-periodic in y. Moreover, under conditions  $(\phi_3)$  and  $(V_7)$  we have that

$$\alpha < u(x,y) < \beta$$
 for all  $(x,y) \in \mathbb{R}^2$ .

2.3. The proof of Theorem 1.4. This theorem is an immediate consequence of the Lemmas 2.4, 2.5 and 2.7.

2.4. The proof of Theorem 1.5. The goal of this section is to establish the proof of Theorem 1.5. In this specific case, we are considering that the function A belongs to Class 3. In [3], Alves called this class of *Rabinowitz's condition*, because an assumption like that has been introduced by Rabinowitz [22, Theorem 4.33] to build up a variational framework to study the existence of solution for a partial differential equation of the type

$$-\epsilon^2 \Delta u + A(x)u = f(u)$$
 in  $\mathbb{R}^N$ 

where  $\epsilon > 0, f : \mathbb{R} \to \mathbb{R}$  is a continuous function with subcritical growth and  $A : \mathbb{R}^N \to \mathbb{R}$  is a continuous function satisfying

$$0 < \inf_{x \in \mathbb{R}^N} A(x) < \liminf_{|x| \to +\infty} A(x).$$

Now we will mainly focus on some preliminary results that are crucial in our approach. As a beginning, let us denote by  $I_{\epsilon}, I_{\infty} : W^{1,\Phi}_{\text{loc}}(\Omega) \to \mathbb{R} \cup \{+\infty\}$  the following functionals

$$I_{\epsilon}(v) = \sum_{j \in \mathbb{Z}} a_{\epsilon,j}(v) \text{ and } I_{\infty}(v) = \sum_{j \in \mathbb{Z}} a_{\infty,j}(v),$$

where

$$a_{\epsilon,j}(v) = \iint_{\Omega_j} \left( \Phi(|\nabla v|) + A(\epsilon x, y) V(v) \right) dx dy$$

and

$$a_{\infty,j}(v) = \iint_{\Omega_j} \left( \Phi(|\nabla v|) + A_{\infty}V(v) \right) dxdy.$$

Moreover, we indicate by  $c_{\epsilon,\Phi}(\alpha,\beta)$  and  $c_{\infty,\Phi}(\alpha,\beta)$  the real numbers

$$c_{\epsilon,\Phi}(\alpha,\beta) = \inf_{v\in\Gamma_{\Phi}(\alpha,\beta)} I_{\epsilon}(v) \text{ and } c_{\infty,\Phi}(\alpha,\beta) = \inf_{v\in\Gamma_{\Phi}(\alpha,\beta)} I_{\infty}(v).$$

Here we would like to emphasize that throughout this subsection the potential V satisfies the conditions  $(V_1)$ - $(V_3)$ . The next lemma establishes an important relation between the real numbers  $c_{\epsilon,\Phi}(\alpha,\beta)$  and  $c_{\infty,\Phi}(\alpha,\beta)$ , which will play an essential rule in our approach.

Lemma 2.8. According to the notation above,

$$\limsup_{\epsilon \to 0^+} c_{\epsilon,\Phi}(\alpha,\beta) < c_{\infty,\Phi}(\alpha,\beta).$$

*Proof.* The proof is similar to that discussed in the proof of [3, Lemma 4.1] and its proof is omitted.  $\Box$ 

We are now ready to prove the following result.

**Proposition 2.9.** There exists  $\epsilon_0 > 0$  such that for each  $\epsilon \in (0, \epsilon_0)$  there is  $u_{\epsilon} \in \Gamma_{\Phi}(\alpha, \beta)$  satisfying  $I_{\epsilon}(u_{\epsilon}) = c_{\epsilon,\Phi}(\alpha, \beta)$  and  $\alpha \leq u_{\epsilon}(x, y) \leq \beta$  almost everywhere  $(x, y) \in \Omega$ .

*Proof.* The idea here is to use a variant of the proof of Proposition 2.6 to establish the proposition. First of all, thanks to Lemma 2.8 we may fix  $\epsilon_0 > 0$  small enough verifying

(2.65) 
$$c_{\epsilon,\Phi}(\alpha,\beta) < c_{\infty,\Phi}(\alpha,\beta), \quad \forall \epsilon \in (0,\epsilon_0)$$

Now, arguing as in Subsection 2.1, for each  $\epsilon \in (0, \epsilon_0)$  there exist a minimizing sequence  $(u_n) \subset \Gamma_{\Phi}(\alpha, \beta)$  for  $I_{\epsilon}$  and  $u_{\epsilon} \in W^{1,\Phi}_{\text{loc}}(\Omega)$  such that

$$\begin{aligned} \alpha &\leq u_n(x,y) \leq \beta, \ \forall (x,y) \in \Omega \text{ and } \forall n \in \mathbb{N}, \\ u_n &\rightharpoonup u_\epsilon \text{ in } W^{1,\Phi}_{\text{loc}}(\Omega), \\ u_n &\to u_\epsilon \text{ in } L^{\Phi}_{\text{loc}}(\Omega), \\ u_n(x,y) &\to u_\epsilon(x,y) \text{ a.e in } \Omega, \\ \alpha &\leq u_\epsilon(x,y) \leq \beta \text{ a.e in } \Omega \end{aligned}$$

and

(2.66) 
$$I_{\epsilon}(u_{\epsilon}) \le c_{\epsilon,\Phi}(\alpha,\beta)$$

By a similar argument to the one used in the proof of Proposition 2.6, there are  $u_{\epsilon}^* \in W^{1,\Phi}(\Omega_0)$ and a subsequence of  $(\tau_k u_{\epsilon})$ , still denoted by itself, such that

$$\tau_k u_\epsilon \rightharpoonup u_\epsilon^* \text{ in } W^{1,\Phi}(\Omega_0) \text{ as } k \to +\infty$$
  
 $\tau_k u_\epsilon \to u_\epsilon^* \text{ in } L^{\Phi}(\Omega_0) \text{ as } k \to +\infty$ 

and

$$au_k u_\epsilon(x,y) \to u_\epsilon^*(x,y)$$
 a.e in  $\Omega_0$  as  $k \to +\infty$ .

where  $u_{\epsilon}^* = \alpha$  or  $u_{\epsilon}^* = \beta$  a.e. in  $\Omega_0$ . As in the proofs of Propositions 2.2 and 2.6, we want to show that  $u_{\epsilon} \in \Gamma_{\Phi}(\alpha, \beta)$ . Toward that end, we show that  $u_{\epsilon}^* = \beta$  a.e. in  $\Omega_0$ . The argument is similar to that developed in Proposition 2.6, but we present the proof in detail for the reader's convenience. Indeed, arguing by contradiction, assume that  $u_{\epsilon}^* = \alpha$  a.e. in  $\Omega_0$ . Thus, given  $\delta \in (0, \Phi(\beta - \alpha))$  there exist  $i_* \in \mathbb{N}$ , a sequence  $(k_i) \subset \mathbb{N}$  and a subsequence  $(u_{n_i})$  of  $(u_n)$  such that  $i_* < k_i$  for all  $i \in \mathbb{N}$ ,  $k_i \to +\infty$  and  $n_i \to +\infty$  as  $i \to +\infty$  and

$$\iint_{\Omega_0} \Phi(|\tau_j u_{n_i} - \alpha|) dx dy < \delta \text{ and } \iint_{\Omega_0} \Phi(|\tau_{k_i} u_{n_i} - \alpha|) dx dy \ge \delta \ \forall j \in [i_*, k_i - 1] \cap \mathbb{N}.$$

Consequently, considering the sequence  $(\tau_{k_i}u_{n_i})$ , for some subsequence, there exists  $w_{\epsilon} \in W^{1,\Phi}_{\text{loc}}(\Omega)$  satisfying

$$\tau_{k_i} u_{n_i} \rightharpoonup w_{\epsilon} \text{ in } W^{1,\Phi}_{\text{loc}}(\Omega) \text{ as } i \to +\infty,$$
  
$$\tau_{k_i} u_{n_i} \to w_{\epsilon} \text{ in } L^{\Phi}_{\text{loc}}(\Omega) \text{ as } i \to +\infty$$

and

$$\alpha \leq w_{\epsilon}(x,y) \leq \beta$$
 a.e. in  $\Omega$ .

Setting the functional

$$a_{\epsilon,j}^{i}(v) = \iint_{\Omega_{j}} \left( \Phi(|\nabla v|) + A(\epsilon x + \epsilon k_{i}, y)V(v) \right) dxdy, \quad v \in W_{\text{loc}}^{1,\Phi}(\Omega), \ i \in \mathbb{N} \text{ and } j \in \mathbb{Z},$$

it is easy to check that

$$\sum_{j \in \mathbb{Z}} a_{\epsilon,j}^i(\tau_{k_i} u_{n_i}) = \sum_{j \in \mathbb{Z}} a_{\epsilon,j}(u_{n_i}) = I_{\epsilon}(u_{n_i}), \quad \forall i \in \mathbb{N}.$$

This fact together with the limit below

$$\liminf_{i \to +\infty} A(\epsilon x + \epsilon k_i, y) = A_{\infty}$$

implies that

(2.67) 
$$I_{\infty}(w_{\epsilon}) \le c_{\epsilon,\Phi}(\alpha,\beta).$$

and so,  $a_{\infty,j}(w_{\epsilon})$  goes to 0 as j goes to  $\pm\infty$ . So, by passing to a subsequence if necessary, it is easy to see that

$$\tau_k w_\epsilon \to \alpha \text{ or } \beta \text{ in } L^{\Phi}(\Omega_0) \text{ as } k \to \pm \infty.$$

The same ideas explored in the proof of Claim 1 of Proposition 2.6 ensures that

$$\tau_k w_\epsilon \to \alpha \text{ in } L^{\Phi}(\Omega_0) \text{ as } k \to -\infty.$$

Next, we are going to prove that

(2.68) 
$$\tau_k w_\epsilon \to \beta \text{ in } L^{\Phi}(\Omega_0) \text{ as } k \to +\infty.$$

Assume for contradiction that (2.68) is not true. Arguing as in the proof of Proposition 2.6, it follows that there are  $\tilde{\epsilon}_0 > 0$ ,  $j \in \mathbb{N}$  and  $i \in \mathbb{N}$  large enough such that for some fixed  $\tilde{\epsilon} \in (0, \frac{\tilde{\epsilon}_0}{2})$  one has

(2.69) 
$$\int_{k_i-1}^{k_i+1} \int_0^1 \left( \Phi(|\nabla u_{n_i}|) + V(u_{n_i}) \right) dx dy \ge \frac{\tilde{\epsilon}_0}{\tilde{A}}$$

(2.70) 
$$I_{\epsilon}(u_{n_i}) \le c_{\epsilon,\Phi}(\alpha,\beta) + \frac{\epsilon}{4}$$

and

(2.71) 
$$a_{\infty,j}\left((\tau_{k_i}u_{n_i}-\alpha)(x-j)+\alpha\right) \leq \frac{\epsilon}{2}.$$

Using the function  $U_{j,i} \in \Gamma_{\Phi}(\alpha, \beta)$  given by

$$U_{j,i}(x,y) = \begin{cases} \alpha, & \text{if } x \leq j & \text{and } y \in (0,1), \\ (\tau_{k_i} u_{n_i}(x,y) - \alpha)(x-j) + \alpha, & \text{if } j \leq x \leq j+1 & \text{and } y \in (0,1), \\ \tau_{k_i} u_{n_i}(x,y), & \text{if } x > j+1 & \text{and } y \in (0,1), \end{cases}$$

we derive that

(2.72) 
$$c_{\infty,\Phi}(\alpha,\beta) \le I_{\infty}(U_{j,i}) = a_{\infty,j}(U_{j,i}) + \sum_{t=j+1+k_i}^{+\infty} a_{\infty,t}(u_{n_i}).$$

Now, since the function A belongs to Class 3, increasing i if necessary, an easy computation shows that

(2.73) 
$$\sum_{t=j+1+k_i}^{+\infty} a_{\infty,t}(u_{n_i}) \le \sum_{t=j+1+k_i}^{+\infty} a_{\epsilon,t}(u_{n_i}) + \frac{\tilde{\epsilon}}{4}.$$

Thus, from (2.69)-(2.73),

$$c_{\infty,\Phi}(\alpha,\beta) \le a_{\infty,j}(U_{j,i}) + I_{\epsilon}(u_{n_i}) - \epsilon_0 + \frac{\tilde{\epsilon}}{4} \le c_{\epsilon,\Phi}(\alpha,\beta) - \frac{\tilde{\epsilon}_0}{2},$$

contrary to (2.65). Therefore,  $w_{\epsilon} \in \Gamma_{\Phi}(\alpha, \beta)$  and (2.67) leads to

$$c_{\infty,\Phi}(\alpha,\beta) \leq I_{\infty}(w_{\epsilon}) \leq c_{\epsilon,\Phi}(\alpha,\beta),$$

which again contradicts (2.65). Consequently, we conclude from the study carried out here that  $\tau_k u_{\epsilon} \to \beta$  in  $L^{\Phi}(\Omega_0)$  as  $k \to +\infty$ . By a similar argument, we can conclude that  $u_{\epsilon} \in \Gamma_{\Phi}(\alpha, \beta)$  for  $\epsilon \in (0, \epsilon_0)$ . Moreover, by (2.66), we must have  $I_{\epsilon}(w_{\epsilon}) = c_{\epsilon,\Phi}(\alpha, \beta)$ , finishing the proof.  $\Box$ 

Finally, we can now prove our main result of this subsection.

Proof of Theorem 1.5.

Initially, we will consider the following set

(2.74)

 $K_{\epsilon,\Phi}(\alpha,\beta) = \left\{ u \in \Gamma_{\Phi}(\alpha,\beta) : I_{\epsilon}(u) = c_{\epsilon,\Phi}(\alpha,\beta), \ u(x,0) = u(x,1) \text{ in } \mathbb{R}, \ \alpha \leq u \leq \beta \text{ a.e. on } \Omega \right\},$ 

which consists of points of minima of  $I_{\epsilon}$  on  $\Gamma_{\Phi}(\alpha, \beta)$  that are seen as functions defined on  $\mathbb{R}^2$  being 1-periodic on the variable y. Next, from Proposition 2.9 we can proceed analogously to the proof of Lemma 2.3 for show that  $K_{\epsilon,\Phi}(\alpha,\beta)$  is non empty whenever  $\epsilon \in (0, \epsilon_0)$ . Finally, we point out that the Theorem 1.5 follows following the same steps of Subsection 2.1 and the details are left to the reader.

2.5. The proof of Theorem 1.6. We exhibit in this subsection the proof of Theorem 1.6. To build a framework for this theorem and avoid some bothersome technicalities, we will always assume here that the potential V satisfies the conditions  $(V_1)$ - $(V_3)$ ,  $(V_6)$  and  $(V_8)$ . Furthermore, we will consider assumptions  $(\phi_1)$ - $(\phi_2)$  on  $\phi$ ,  $\epsilon = 1$  and that A belongs to Class 4. Next we consider the following class of admissible functions

 $\Gamma_{\Phi}^{o}(\beta) = \{ v \in \Gamma_{\Phi}(-\beta,\beta) : v(x,y) = -v(-x,y) \text{ a.e. in } \Omega \text{ and } 0 \le v(x,y) \le \beta \text{ for a.e } x \ge 0 \}$ 

and the real number

$$c_{\Phi}^{o}(\beta) = \inf_{v \in \Gamma_{\Phi}^{o}(\beta)} I(v),$$

where  $\Gamma_{\Phi}(-\beta,\beta)$  is given as in (2.1). Now it is important to point out that  $\Gamma_{\Phi}^{o}(\beta)$  is not empty, because the function  $\varphi_{-\beta,\beta}$  defined as in (2.2) belongs to  $\Gamma_{\Phi}^{o}(\beta)$  with  $I(\varphi_{-\beta,\beta}) < +\infty$ . Having said that, we shall now explore the conditions  $(V_6)$  and  $(V_8)$  to show that the following class

(2.75) 
$$K^o_{\Phi}(\beta) = \{ v \in \Gamma^o_{\Phi}(\beta) : I(v) = c^o_{\Phi}(\beta) \text{ and } v(x,0) = v(x,1) \text{ in } \mathbb{R} \}$$

is not empty. Hereafter, we will assume that the functions of  $K^o_{\Phi}(\beta)$  are periodically extended in  $\mathbb{R}^2$  on the variable y. Therefore,  $K^o_{\Phi}(\beta)$  is constituted by (minimal) heteroclinic type solutions of (PDE) with  $\epsilon = 1$  that are 1-periodic in y and odd in x.

**Lemma 2.10.** It holds that  $K^o_{\Phi}(\beta) \neq \emptyset$ .

*Proof.* By some standard computations, one easily verifies that there exists a minimizing sequence  $(u_n) \subset \Gamma^o_{\Phi}(\beta)$  for I such that

$$0 \le u_n(x, y) \le \beta, \ \forall n \in \mathbb{N} \text{ and } x \ge 0.$$

Besides that, there exist  $u \in W^{1,\Phi}_{\text{loc}}(\Omega)$  and a subsequence of  $(u_n)$ , still denoted by  $(u_n)$ , satisfying

(2.76) 
$$u_n \rightharpoonup u \text{ in } W^{1,\Phi}_{\text{loc}}(\Omega)$$

(2.77) 
$$u_n \to u \text{ in } L^{\Phi}_{\text{loc}}(\Omega)$$

and

(2.78) 
$$u_n(x,y) \to u(x,y)$$
 a.e. on  $\Omega$ .

We conclude from (2.78) that u(x,y) = -u(-x,y) almost everywhere  $(x,y) \in \Omega$  and  $0 \leq u(x,y) \leq \beta$  for almost every  $x \geq 0$ , and finally by (2.76)-(2.77) it is easy to check that

$$(2.79) I(u) \le c_{\Phi}^{o}(\beta)$$

Now we claim that  $u \in \Gamma_{\Phi}^{o}(\beta)$ . To establish our claim, we assume for the sake of contradiction that  $\tau_{k}u$  not goes to  $\beta$  as k goes to  $+\infty$  in  $L^{\Phi}(\Omega_{0})$ . Thereby, since  $\Phi \in \Delta_{2}$  there are  $\epsilon > 0$  and a subsequence  $(k_{i})$  of natural numbers with  $k_{i} \to +\infty$  such that

(2.80) 
$$\iint_{\Omega_0} \Phi\left(|\tau_{k_i}u - \beta|\right) dx dy \ge \epsilon, \quad \forall i \in \mathbb{N}.$$

On the other hand,  $(V_1)$ - $(V_3)$  and  $(V_8)$  yield

$$\tilde{\mu}\Phi(|t-\beta|) \le V(t), \quad \forall t \in [0,\beta],$$

for some  $\tilde{\mu} > 0$ . Consequently,

$$I(u) \ge \sum_{i \in \mathbb{N}} \left( \iint_{\Omega_{k_i}} A(x, y) V(u) dx dy \right) \ge \tilde{\mu} \sum_{i \in \mathbb{N}} \left( \iint_{\Omega_{k_i}} A(x, y) \Phi\left( |u - \beta| \right) dx dy \right),$$

that is,

$$I(u) \ge \tilde{\mu} \sum_{i \in \mathbb{N}} \left( \iint_{\Omega_0} A(x+k_i, y) \Phi\left( |\tau_{k_i} u - \beta| \right) dx dy \right).$$

Now, fixing  $i_0 \in \mathbb{N}$  such that  $|x + k_i| \ge K$  for any  $x \in [0, 1]$  and  $i \ge i_0$ , the fact that A belongs to Class 4 leads to

$$I(u) \ge \tilde{\mu}a_0 \sum_{i\ge i_0} \left( \iint_{\Omega_0} \Phi\left( |\tau_{k_i}u - \beta| \right) dx dy \right),$$

where

$$a_0 = \inf_{|x| \ge K, y \in [0,1]} A(x,y) > 0.$$

Hence, by (2.80),

$$I(u) \ge a_0 \tilde{\mu} \sum_{i \ge i_0} \epsilon = +\infty,$$

contrary to (2.79). For this reason,

$$\tau_k u \to \beta$$
 in  $L^{\Phi}(\Omega_0)$  as  $k \to +\infty$ ,

and therefore, since u is odd in the variable x we conclude that  $u \in \Gamma_{\Phi}^{o}(\beta)$ . This fact combined with (2.79) produces that  $I(u) = c_{\Phi}^{o}(\beta)$ . Now assumptions  $(A_2)$ - $(A_3)$  allow us to proceed as in the proof of Lemma 2.3 to find a function  $v \in \Gamma_{\Phi}^{o}(\beta)$  dependent on u such that  $v \in K_{\Phi}^{o}(\beta)$ , and the proof is over.

We now finish this subsection by proving Theorem 1.6 as follows.

#### Proof of Theorem 1.6.

Our proof follows the method developed in [4] and we will do it in detail for the reader's convenience. To begin with, thanks to Lemma 2.10 we can take  $u \in K^o_{\Phi}(\beta)$ . Here we will first show that

(2.81) 
$$\iint_{\Omega_0} \left( \phi\left(|\nabla u|\right) \nabla u \nabla \psi + A(x,y) V'(u) \psi \right) dx dy \ge 0, \text{ for all } \psi \in C_0^\infty(\mathbb{R}^2),$$

which will guarantee that

$$\iint_{\Omega_0} \left( \phi\left( |\nabla u| \right) \nabla u \nabla \psi + A(x, y) V'(u) \psi \right) dx dy = 0, \text{ for all } \psi \in C_0^\infty(\mathbb{R}^2),$$

implying that u is a weak solution of (PDE).

In what follows, for each  $\psi \in C_0^{\infty}(\mathbb{R}^2)$ , we will use the fact that  $\psi(x, y) = \psi_o(x, y) + \psi_e(x, y)$ , where

$$\psi_e(x,y) = \frac{\psi(x,y) + \psi(-x,y)}{2}$$
 and  $\psi_o(x,y) = \frac{\psi(x,y) - \psi(-x,y)}{2}$ .

In addition to these functions, let us consider for t > 0 the function

$$\tilde{\varphi}_t(x,y) = \max\left\{-\beta, \min\{\beta, \varphi_t(x,y)\}\right\}, \quad (x,y) \in \Omega,$$

where

$$\varphi_t(x,y) = \begin{cases} u(x,y) + t\psi_o(x,y), & \text{if } x \ge 0 \text{ and } u(x,y) + t\psi_o(x,y) \ge 0, \\ -u(x,y) - t\psi_o(x,y), & \text{if } x \ge 0 \text{ and } u(x,y) + t\psi_o(x,y) \le 0, \\ -\varphi_t(-x,y), & \text{if } x < 0. \end{cases}$$

Now, a direct computation shows that  $\tilde{\varphi}_t \in \Gamma^o_{\Phi}(\beta)$ . Then  $(V_6)$  together with  $\tilde{\varphi}_t$  yields that

(2.82) 
$$I(u + t\psi_o) = I(\varphi_t) \ge I(\tilde{\varphi}_t) \ge c_{\Phi}^o(\beta) = I(u)$$

On the other hand, by Lemma A.5-(b),

$$\begin{split} I(u+t\psi) - I(u+t\psi_o) &\geq t \sum_{j \in \mathbb{Z}} \iint_{\Omega_j} \phi(|\nabla(u+t\psi_o)|) \nabla u \nabla \psi_e dx dy \\ &+ t^2 \sum_{j \in \mathbb{Z}} \iint_{\Omega_j} \phi(|\nabla(u+t\psi_o)|) \nabla \psi_o \nabla \psi_e dx dy \\ &+ \sum_{j \in \mathbb{Z}} \iint_{\Omega_j} A(x,y) \left( V(u+t\psi) - V(u+t\psi_o) \right) dx dy. \end{split}$$

Since the functions  $\phi(|\nabla(u+t\psi_o)|)\nabla u\nabla\psi_e$  and  $\phi(|\nabla(u+t\psi_o)|)\nabla\psi_o\nabla\psi_e$  are odd in the variable x, then it is easily seen that

$$(2.84) \quad \sum_{j \in \mathbb{Z}} \iint_{\Omega_j} \phi(|\nabla(u+t\psi_o)|) \nabla u \nabla \psi_e dx dy = \sum_{j \in \mathbb{Z}} \iint_{\Omega_j} \phi(|\nabla(u+t\psi_o)|) \nabla \psi_o \nabla \psi_e dx dy = 0,$$

and so, from (2.82)-(2.84),

$$I(u+t\psi) - I(u) \ge \sum_{j \in \mathbb{Z}} \iint_{\Omega_j} A(x,y) \left( V(u+t\psi) - V(u+t\psi_o) \right) dxdy.$$

Consequently, as  $A(x, y)V'(u)\psi_e$  is odd in the variable x, one gets

$$\begin{aligned} (2.85) \\ \iint_{\Omega} (\phi(|\nabla u|) \nabla u \nabla \psi + A(x,y) V'(u) \psi) dx dy &= \lim_{t \to 0^+} \frac{I(u+t\psi) - I(u)}{t} \\ &\geq \lim_{t \to 0^+} \sum_{j \in \mathbb{Z}} \iint_{\Omega_j} A(x,y) \frac{V(u+t\psi) - V(u+t\psi_o)}{t} dx dy \\ &\geq \sum_{j \in \mathbb{Z}} \iint_{\Omega_j} A(x,y) V'(u) (\psi - \psi_o) dx dy = \sum_{j \in \mathbb{Z}} \iint_{\Omega_j} A(x,y) V'(u) \psi_e dx dy = 0, \end{aligned}$$

showing that the inequality (2.81) occurs for every  $\psi \in C_0^{\infty}(\mathbb{R}^2)$ . Finally, slightly varying the same ideas discussed in the Subsection 2.1, we can conclude the proof of Theorem 1.6.

26

#### 3. Heteroclinic solution of the prescribed curvature equation

Throughout this section, we adapt for our problem the approach explored in [5] to find solutions that are periodic in the variable y and heteroclinic in x from  $\alpha$  to  $\beta$  to the prescribed mean curvature equation (E). In this section, we will return to indexing  $\alpha$  and  $\beta$  in V, denoted as  $V_{\alpha,\beta}$ , because here our focus is on choosing suitable values of  $\alpha$  and  $\beta$ . Since the ideas are so close to those of [5], the presentation will be brief. In the following, we consider for each L > 0 the quasilinear equation

$$-\Delta_{\Phi_L} u + A(\epsilon x, y) V'_{\alpha,\beta}(u) = 0 \quad \text{in} \quad \mathbb{R}^2, \tag{E}_L$$

with  $V_{\alpha,\beta} \in \mathcal{V}$ , where  $\Phi_L : \mathbb{R} \to [0, +\infty)$  is an N-function of the form

$$\Phi_L(t) = \int_0^{|t|} \phi_L(s) s ds,$$

where  $\phi_L(t) = \varphi_L(t^2)$  and  $\varphi_L$  is defined by

$$\varphi_L(t) = \begin{cases} \frac{1}{\sqrt{1+t}}, & \text{if } t \in [0,L], \\ x_L(t-L-1)^2 + y_L, & \text{if } t \in [L,L+1], \\ y_L, & \text{if } t \in [L+1,+\infty) \end{cases}$$

with

$$x_L = \frac{\sqrt{1+L}}{4(1+L)^2}$$
 and  $y_L = (4L+3)x_L$ .

We point out that the main purpose of this section is to use the arguments of Sect. 2 to investigate the existence of a heteroclinic solution  $u_{\alpha,\beta}$  from  $\alpha$  to  $\beta$  for  $(E)_L$  that satisfies

(3.1) 
$$\|\nabla u_{\alpha,\beta}\|_{L^{\infty}(\mathbb{R}^2)} \leq \sqrt{L},$$

because this inequality implies that  $u_{\alpha,\beta}$  is a heteroclinic solution from  $\alpha$  to  $\beta$  for (E). Here, we will prove that the inequality (3.1) holds when max{ $|\alpha|, |\beta|$ } is small enough. In order to do that, a control involving the roots  $\alpha$  and  $\beta$  of  $V_{\alpha,\beta}$  is necessary, and at this point the condition  $(V_4)$  applies an important rule in our argument.

The next result is about functions  $\phi_L$  and  $\Phi_L$ , which makes it clear that  $\Phi_L$  is an N-function.

**Lemma 3.1.** For each L > 0, the functions  $\phi_L$  and  $\Phi_L$  have the following properties:

(a)  $\phi_L$  is  $C^1$ . (b)  $y_L \leq \phi_L(t) \leq 1$  for all  $t \geq 0$ . (c)  $\frac{y_L}{2}t^2 \leq \Phi_L(t) \leq \frac{1}{2}t^2$  for any  $t \in \mathbb{R}$ . (d)  $\Phi_L$  is a convex function. (e)  $(\phi_L(t)t)' > 0$  for all t > 0.

*Proof.* The argument follow the same lines as the proof of [5, Lemma 3.1].

We would like to point out that our focus now is on examining if the N-function  $\Phi_L$  is in the settings of Sect. 2, that is, if  $\phi_L$  satisfies conditions  $(\phi_1)$ - $(\phi_3)$ . Indeed, it is clear that by Lemma 3.1-(e)  $\phi_L$  checks  $(\phi_1)$ , and by Lemma 3.1-(b), it checks  $(\phi_3)$  with q = 2. Moreover, with direct computations one can get that there are real numbers  $m_L, l_L > 1$  such that  $l_L \leq m_L$ and

$$l_L - 1 \le \frac{(\phi_L(t)t)'}{\phi_L(t)} \le m_L - 1 \quad \text{for any} \quad t \ge 0,$$

from which it follows that  $\phi_L$  verifies  $(\phi_2)$ . As a direct consequence the N-functions  $\Phi_L$  and  $\tilde{\Phi}_L$  satisfy  $\Delta_2$ -condition, where  $\tilde{\Phi}_L$  is the complementary function associated with  $\Phi_L$ , which

ensures that the space  $L^{\Phi_L}$  is reflexive (see for instance Appendix A). Actually, the study made in [5, Lemma 3.2] shows that the space  $L^{\Phi_L}$  is exactly  $L^2$  space and the norm of  $L^{\Phi_L}$  is equivalent to the norm of  $L^2$ .

Now, assuming for a moment that function A belongs to Class 1 or 2,  $\epsilon = 1$ , and that the potential  $V_{\alpha,\beta} \in \mathcal{V}$ , the same arguments from Subsections 2.1 and 2.2 guarantee that there exist a periodic function  $u_{\alpha,\beta} : \mathbb{R}^2 \to \mathbb{R}$  on the variable y such that  $u_{\alpha,\beta} \in K_{\Phi_L}(\alpha,\beta)$ , where

$$K_{\Phi_L}(\alpha,\beta) = \{ w \in \Gamma_{\Phi_L}(\alpha,\beta) : I(w) = c_{\Phi_L}(\alpha,\beta), \ w(x,0) = w(x,1) \text{ in } \mathbb{R}, \ \alpha \le w \le \beta \text{ a.e. on } \Omega \}.$$

Moreover,  $u_{\alpha,\beta}$  is a weak solution of equation  $(E)_L$  with  $\epsilon = 1$  in  $C_{\text{loc}}^{1,\gamma}(\mathbb{R}^2)$  for some  $\gamma \in (0,1)$  that is heteroclinic in x from  $\alpha$  to  $\beta$ . The next lemma is crucial to guarantee the existence of a heteroclinic solution for (E) and its proof is inspired by the proof of [5, Lemma 3.3].

**Lemma 3.2.** There exists  $\delta > 0$  such that for each  $V_{\alpha,\beta} \in \mathcal{V}$  with  $\max\{|\alpha|, |\beta|\} < \delta$ , the heteroclinic solution  $u_{\alpha,\beta} \in K_{\Phi_L}(\alpha,\beta)$  obtained in the previous section satisfies the estimate below

$$\|u_{\alpha,\beta}\|_{C^1(B_1(z))} < \sqrt{L}, \quad \forall z \in \mathbb{R}^2,$$

where  $B_1(z)$  denotes the ball in  $\mathbb{R}^2$  of center z and radius 1.

*Proof.* If the lemma does not hold, then for each  $n \in \mathbb{N}$  there is a potential  $V_{\alpha_n,\beta_n} \in \mathcal{V}$  with  $\max\{|\alpha_n|, |\beta_n|\} < n^{-1}$  and  $z_n = (r_n, s_n) \subset \mathbb{R}^2$  such that the heteroclinic solution  $u_{\alpha_n,\beta_n} \in K_{\Phi_L}(\alpha_n,\beta_n)$  satisfies

(3.2) 
$$\|u_{\alpha_n,\beta_n}\|_{C^1(B_1(r_n,s_n))} \ge \sqrt{L}, \quad \forall n \in \mathbb{N}.$$

Now, we note that for each  $n \in \mathbb{N}$  the function  $\tilde{u}_n$  defined by

$$\tilde{u}_n(x,y) = u_{\alpha_n,\beta_n}(x+r_n,y+s_n) \text{ for } (x,y) \in \mathbb{R}^2$$

is a weak solution of the quasilinear equation

$$-\Delta_{\Phi_L} u + B_n(x, y) = 0 \text{ in } \mathbb{R}^2,$$

where

$$B_n(x,y) = A(x+r_n, y+s_n)V'_{\alpha_n,\beta_n}(u_{\alpha_n,\beta_n}(x,y)).$$

Furthermore, the condition  $(V_4)$  is crucial to show the following inequality

(3.3) 
$$|B_n(x,y)| \le M ||A||_{L^{\infty}(\mathbb{R}^2)} \quad \forall (x,y) \in \mathbb{R}^2 \text{ and } \forall n \ge n_0,$$

where  $n_0 \in \mathbb{N}$  and M is a positive constant independent of n. Indeed, let us first observe that the condition max  $\{|\alpha_n|, |\beta_n|\} \to 0$  as  $n \to +\infty$  implies that there exists  $n_0 \in \mathbb{N}$  such that  $\alpha_n, \beta_n \in (-1, 1)$  for all  $n \ge n_0$ . Since

$$\alpha_n \le u_{\alpha_n,\beta_n}(x,y) \le \beta_n, \ \forall (x,y) \in \mathbb{R}^2 \text{ and } \forall n \in \mathbb{N},$$

by condition  $(V_4)$ , it follows that there exists M = M(1) > 0 such that

$$\left|V_{\alpha_n,\beta_n}'\left(u_{\alpha_n,\beta_n}(x,y)\right)\right| \le M, \quad \forall (x,y) \in \mathbb{R}^2 \text{ and } \forall n \ge n_0,$$

which is sufficient to guarantee estimate (3.3). Therefore, the elliptic regularity theory found in [17, Theorem 1.7] implies that  $\tilde{u}_n \in C^{1,\gamma_0}_{\text{loc}}(\mathbb{R}^2)$ , for some  $\gamma_0 \in (0,1)$ , and that there is a positive constant R independent of n verifying

$$\|\tilde{u}_n\|_{C^{1,\gamma_0}_{\text{loc}}(\mathbb{R}^2)} \le R \quad \forall \ n \in \mathbb{N}$$

The above estimate allows us to use Arzelà–Ascoli Theorem to find  $u \in C^1(B_1(0))$  and a subsequence of  $(\tilde{u}_n)$ , still denoted by  $(\tilde{u}_n)$ , such that

$$\tilde{u}_n \to u$$
 in  $C^1(B_1(0))$ .

Now since  $\|u_{\alpha_n,\beta_n}\|_{L^{\infty}(\mathbb{R}^2)}$  tends to zero as  $n \to +\infty$ , we obtain that u = 0 on  $B_1(0)$ , and so,

$$\|\tilde{u}_n\|_{C^1(B_1(0))} < \sqrt{L}, \quad \forall \ n \ge n_0$$

for some  $n_0 \in \mathbb{N}$ . Therefore,

$$||u_{\alpha_n,\beta_n}||_{C^1(B_1(r_n,s_n))} < \sqrt{L}, \quad \forall \ n \ge n_0,$$

which contradicts (3.2), and the proof is completed.

We are finally ready to prove the Theorem 1.1.

## Proof of Theorem 1.1.

To begin with, we claim that given L > 0 there exists  $\delta > 0$  such that if  $\max\{|\alpha|, |\beta|\} \in (0, \delta)$  we have

(3.4) 
$$\|\nabla u_{\alpha,\beta}\|_{L^{\infty}(\mathbb{R}^2)} \leq \sqrt{L} \text{ for all } u_{\alpha,\beta} \in K_{\Phi_L}(\alpha,\beta).$$

Indeed, for each  $(x, y) \in \mathbb{R}^2$  we can choose  $z \in \mathbb{R}^2$  verifying  $(x, y) \in B_1(z)$ . Thanks to Lemma 3.2, there is  $\delta > 0$  such that for each pair  $(\alpha, \beta)$  of real numbers with  $\alpha < \beta$  and  $\max\{|\alpha|, |\beta|\} \in (0, \delta)$  one has

$$||u_{\alpha,\beta}||_{C^1(B_1(z))} < \sqrt{L}$$

whenever  $u_{\alpha,\beta} \in K_{\Phi_L}(\alpha,\beta)$ . Now, from the arbitrariness of (x,y) it is easy to see that our claim is established. Therefore, the estimate (3.4) ensures  $u_{\alpha,\beta}$  is a heteroclinic solution of (E). To complete the proof, the fact that  $V_{\alpha,\beta} \in C^2(\mathbb{R},\mathbb{R})$  combined with the assumptions  $(V_2)$ - $(V_3)$ yields that there are  $\lambda, d_1, d_2 > 0$  such that

$$|V_{\alpha,\beta}'(t)| \le d_1 |t - \alpha|, \quad \forall t \in [\alpha - \lambda, \alpha + \lambda]$$

and

$$|V'_{\alpha,\beta}(t)| \le d_2|t-\beta|, \quad \forall t \in [\beta-\lambda,\beta+\lambda].$$

Thus, by Lemma 3.1-(b),

$$|V_{\alpha,\beta}'(t)| \le \frac{d_1}{y_L} \phi_L(|t-\alpha|)|t-\alpha|, \quad \forall t \in [\alpha - \lambda, \alpha + \lambda]$$

and

$$|V'_{\alpha,\beta}(t)| \le \frac{d_2}{y_L} \phi_L(|t-\beta|)|t-\beta|, \quad \forall t \in [\beta-\lambda,\beta+\lambda].$$

Consequently,  $V_{\alpha,\beta}$  satisfies  $(V_7)$  with  $\phi_L$ , and so, proceeding as in the proof of Lemma 2.5, we see that  $u_{\alpha,\beta}$  verifies

$$\alpha < u_{\alpha,\beta}(x,y) < \beta$$
 for all  $(x,y) \in \mathbb{R}^2$ ,

which is the desired conclusion.

Now, let us assume that A belongs to Class 3 and that  $V_{\alpha,\beta} \in \mathcal{V}$ . Considering the set  $K_{\epsilon,\Phi_L}(\alpha,\beta)$  as in (2.74), then we can argue similarly to the proof of Theorem 1.5 to obtain that  $K_{\epsilon,\Phi_L}(\alpha,\beta) \neq \emptyset$  whenever  $\epsilon > 0$  is small. With everything, proceeding analogously as in the proof of Lemma 3.2 we get the following result.

**Lemma 3.3.** There exist  $\delta > 0$  independent of  $\epsilon > 0$  and  $\epsilon_0 > 0$  such that for each  $\epsilon \in (0, \epsilon_0)$ and  $V_{\alpha,\beta} \in \mathcal{V}$  with  $\max\{|\alpha|, |\beta|\} \in (0, \delta)$ , the heteroclinic solution  $u_{\epsilon,\alpha,\beta} \in K_{\epsilon,\Phi_L}(\alpha,\beta)$  satisfies the following estimate

$$\|u_{\epsilon,\alpha,\beta}\|_{C^1(B_1(z))} < \sqrt{L}, \quad \forall z \in \mathbb{R}^2.$$

We now present the proof of Theorem 1.2.

# Proof of Theorem 1.2.

The proof can be done via a comparison argument like that of the proof Theorem 1.1 and we omit it here. Details are left to the reader.  $\Box$ 

For the final exhibition of these ideas, let's assume that the function A belongs to Class 4,  $\alpha = -\beta$ ,  $\epsilon = 1$  and that each  $V_{-\beta,\beta} \in \mathcal{V} \cap C^2(\mathbb{R},\mathbb{R})$  also satisfies the conditions  $(V_5)$  and  $(V_6)$ . We want to point out that condition  $(V_5)$  implies that the potential  $V_{-\beta,\beta}$  satisfies  $(V_8)$  with  $\Phi_L$ . In fact, note that by  $(V_5)$  there are  $\rho > 0$  and  $\theta \in (0, \frac{\beta}{2})$  such that

(3.5) 
$$\rho |t - \beta|^2 \le V_{-\beta,\beta}(t), \ \forall t \in (\beta - \theta, \beta + \theta)$$

from which it follows by Lemma 3.1-(c) and (3.5),

$$2\rho\Phi_L(|t-\beta|) \le V_{-\beta,\beta}(t), \ \forall t \in (\beta-\theta,\beta+\theta).$$

Consequently, the argument of Subsection 2.5 shows that for each L > 0 the set  $K_{\Phi_L}^o(\beta)$  is not empty, where  $K_{\Phi_L}^o(\beta)$  is given as in (2.75). We would like to remind here that each element of  $K_{\Phi}^o(\beta)$  can be seen as a function on  $\mathbb{R}^2$  being periodic in the variable y. Moreover, if  $u_{\beta} \in K_{\Phi_L}^o(\beta)$ , then  $u_{\beta}$  is a weak solution for  $(E)_L$  with  $\epsilon = 1$  in  $C_{\text{loc}}^{1,\gamma}(\mathbb{R}^2)$  for some  $\gamma \in (0,1)$ , odd and heteroclinic from  $-\beta$  to  $\beta$  in x satisfying

$$0 \le u_{\beta}(x, y) \le \beta$$
 for all  $(x, y) \in \mathbb{R}_+ \times \mathbb{R}$ .

Now, the following result is a similar version of Lemma 3.2 and is proved in an essentially identical fashion, which will play an analogous role to that developed in Theorem 1.1 in the present setting.

**Lemma 3.4.** There exists  $\delta > 0$  such that for each  $V_{-\beta,\beta} \in \mathcal{V} \cap C^2(\mathbb{R},\mathbb{R})$  satisfying  $(V_5)$ - $(V_6)$  with  $\beta \in (0, \delta)$ , the heteroclinic solution  $u_\beta \in K^o_{\Phi_T}(\beta)$  satisfies

$$\|u_{\beta}\|_{C^1(B_1(z))} < \sqrt{L}, \quad \forall z \in \mathbb{R}^2.$$

Finally, to conclude this section, we prove Theorem 1.3 using the framework discussed above. **Proof of Theorem 1.3**.

The proof is established using Lemma 3.4 and arguing as in the proof of Theorem 1.1. Detailed verification is left to the reader.  $\hfill \Box$ 

# APPENDIX A. ORLICZ AND ORLICZ-SOBOLEV SPACES

In the years 1932 and 1936 Orlicz considered and investigated in [20, 21] the following class of functions

$$L^{\Phi}(\mathcal{O}) = \left\{ u \in L^{1}_{\text{loc}}(\mathcal{O}) : \int_{\mathcal{O}} \Phi\left(\frac{|u|}{\lambda}\right) dx < +\infty \text{ for some } \lambda > 0 \right\},$$

where  $\mathcal{O}$  is an open set of  $\mathbb{R}^N$  with  $N \geq 1$  and the function  $\Phi : \mathbb{R} \to [0, +\infty)$  has the following properties:

- (a)  $\Phi$  is continuous, convex and even,
- (b)  $\Phi(t) = 0$  if and only if t = 0,
- (c)  $\lim_{t \to 0} \frac{\Phi(t)}{t} = 0$  and  $\lim_{t \to +\infty} \frac{\Phi(t)}{t} = +\infty$ .

In these configurations, we say that  $\Phi$  is an *N*-function. In Orlicz's paper [20], he also introduced an additional condition on the function  $\Phi$  the so called  $\Delta_2$ -condition ( $\Phi \in \Delta_2$  for short) which says that there exist constants C > 0 and  $t_0 \ge 0$  such that

$$\Phi(2t) \le C\Phi(t), \quad \forall t \ge t_0. \tag{\Delta}_2$$

Below are some examples of N-functions that satisfy  $(\Delta_2)$  with  $t_0 = 0$ :

1. 
$$\Phi_1(t) = \frac{|t|^p}{p}$$
 with  $p \in (1, +\infty)$ ,  
2.  $\Phi_2(t) = \frac{|t|^p}{p} + \frac{|t|^q}{q}$  for  $p < q$  and  $p, q \in (1, +\infty)$ ,  
3.  $\Phi_3(t) = |t|^p \ln(1 + |t|)$ , where  $p \in (1, +\infty)$ ,  
4.  $\Phi_4(t) = (1 + t^2)^{\gamma} - 1$  with  $\gamma > 1$ ,  
5.  $\Phi_5(t) = (\sqrt{1 + t^2} - 1)^{\gamma}$  for  $\gamma > 1$ ,  
6.  $\Phi_6(t) = \int_0^t s^{1-\gamma} (\sinh^{-1} s)^{\beta} ds$  with  $0 \le \gamma < 1$  and  $\beta > 0$ .

The following Minkowski functional

$$\|u\|_{L^{\Phi}(\mathcal{O})} = \inf\left\{\lambda > 0 : \int_{\mathcal{O}} \Phi\left(\frac{|u|}{\lambda}\right) dx \le 1\right\}, \quad u \in L^{\Phi}(\mathcal{O}),$$

was introduced by Luxemburg in his thesis [18] and the reader can verify that  $||u||_{L^{\Phi}(\mathcal{O})}$  defines a norm on  $L^{\Phi}(\mathcal{O})$ , which is called the **Luxemburg norm** over  $\mathcal{O}$ . A direct computation ensures that  $L^{\Phi}(\mathcal{O})$  endowed with the Luxemburg norm is a Banach space. To illustrate this phenomenon in particular cases, consider  $1 and <math>\Phi_p(t) = \frac{|t|^p}{p}$  to see that

$$L^{\Phi_p}(\mathcal{O}) = L^p(\mathcal{O}) \text{ and } \|u\|_{L^{\Phi_p}(\mathcal{O})} = p^{-\frac{1}{p}} \|u\|_{L^p(\mathcal{O})},$$

where  $\|\cdot\|_{L^p(\mathcal{O})}$  is the usual norm of space  $L^p(\mathcal{O})$ . This fact makes it clear that the spaces  $L^{\Phi}(\mathcal{O})$  are more general than Lebesgue's  $L^p$  spaces. However, these spaces may possess peculiar properties that do not occur in ordinary Lebesgue's spaces. Now let us note as a point of interest that if  $|\mathcal{O}| < +\infty$ , then a direct computation shows that the embedding

$$L^{\Phi}(\mathcal{O}) \hookrightarrow L^{1}(\mathcal{O})$$

is continuous. In other words, this embedding is equivalent to the simple containment  $L^{\Phi}(\mathcal{O}) \subset L^{1}(\mathcal{O})$  in which some topological properties are preserved. The  $L^{\Phi}(\mathcal{O})$  space became known in the literature as **Orlicz space** over  $\mathcal{O}$ . In [20], Orlicz restricted himself to the case  $\Phi \in \Delta_2$  and in this case we may write the Orlicz spaces as follows

$$L^{\Phi}(\mathcal{O}) = \left\{ u \in L^{1}_{\mathrm{loc}}(\mathcal{O}) : \int_{\mathcal{O}} \Phi(|u|) dx < +\infty \right\}.$$

Furthermore, still under  $\Delta_2$ -condition we have the following fact

(A.1) 
$$u_n \to u \text{ in } L^{\Phi}(\mathcal{O}) \Leftrightarrow \int_{\mathcal{O}} \Phi(|u_n - u|) dx \to 0$$

It is important to mention here that in general these facts do not occur when  $\Phi$  does not satisfy ( $\Delta_2$ ). Orlicz was the first to investigate this case, that is, without the  $\Delta_2$ -condition, in his paper [21] in the year 1936.

A natural generalization of Sobolev spaces  $W^{1,p}(\mathcal{O})$  is the following Banach space associated with an N-function  $\Phi$ 

$$W^{1,\Phi}(\mathcal{O}) = \left\{ u \in L^{\Phi}(\mathcal{O}) : \frac{\partial u}{\partial x_i} = u_{x_i} \in L^{\Phi}(\mathcal{O}), \ i = 1, ..., N \right\},\$$

equipped with the norm

$$\|u\|_{W^{1,\Phi}(\mathcal{O})} = \|\nabla u\|_{L^{\Phi}(\mathcal{O})} + \|u\|_{L^{\Phi}(\mathcal{O})}$$

where  $\nabla u = (u_{x_1}, ..., u_{x_N})$ . Space  $W^{1,\Phi}(\mathcal{O})$  is usually called **Orlicz-Sobolev space** associated with  $\Phi$  over  $\mathcal{O}$ . It is impossible for the authors to give an exhaustive account here of the vast

literature devoted to Orlicz-type spaces. For a quite comprehensive account of this topic, the interested reader might start by referring to [14, 25] and the bibliography therein.

We are going to now consider some results that will be widely used in the development of this work. Initially we observe that given an N-function  $\Phi$ , the complementary function  $\Phi$ associated with  $\Phi$  is defined by Legendre's transformation

$$\tilde{\Phi}(s) = \max_{t \ge 0} \left\{ st - \Phi(t) \right\} \text{ for } s \ge 0.$$

The function  $\tilde{\Phi}$  is also an N-function, and moreover, the functions  $\Phi$  and  $\tilde{\Phi}$  are complementary each other. An interesting example of such complementary functions are

$$\Phi(t) = \frac{|t|^p}{p}$$
 and  $\tilde{\Phi}(t) = \frac{|t|^q}{q}$  with  $1 and  $\frac{1}{p} + \frac{1}{q} = 1$ .$ 

An important result of the theory of Orlicz spaces is the following.

**Lemma A.1.** The space  $L^{\Phi}(\mathcal{O})$  is reflexive if, and only if,  $\Phi$  and  $\tilde{\Phi}$  satisfy the  $\Delta_2$ -condition. *Proof.* See for instance [25, Ch. IV, Theorem 10]. 

Before ending this section, we would like to point out that a function  $\Phi$  of the form (1.7) satisfying  $(\phi_1)$  and  $(\phi_2)$  is an N-function. Moreover, the reader can verify that examples 1 to 6 listed above are models of N-functions of the type (1.7) checking  $(\phi_1)$ - $(\phi_2)$ . Next, we list some lemmas about N-functions of the form (1.7) that will be used frequently in this work.

**Lemma A.2.** Let  $\Phi$  be an N-function of the form (1.7) satisfying  $(\phi_1)$ - $(\phi_2)$ . Set

$$\xi_0(t) = \min\{t^l, t^m\}$$
 and  $\xi_1(t) = \max\{t^l, t^m\}, \quad \forall t \ge 0.$ 

Then  $\Phi$  satisfies

$$\xi_0(t)\Phi(s) \le \Phi(st) \le \xi_1(t)\Phi(s), \quad \forall s, t \ge 0.$$

In particular,  $\Phi \in \Delta_2$ .

*Proof.* Let's first show that condition  $(\phi_2)$  leads to

(A.2) 
$$l \le \frac{\phi(t)t^2}{\Phi(t)} \le m, \quad \forall t > 0.$$

Indeed, by  $(\phi_2)$  we can write

 $l\phi(t) \le (\phi(t)t)' + \phi(t) \le m\phi(t), \quad \forall t > 0,$ 

from which it follows that

$$l\phi(t)t \le (\phi(t)t^2)' \le m\phi(t)t, \quad \forall t > 0.$$

Therefore, integrating the last inequality we get (A.2), as we wanted. Finally, due to estimation (A.2) the proof becomes similar to that given in [11, Lemma 2.1]. The details are left to the reader. 

**Lemma A.3.** Let  $\Phi$  be an N-function of the form (1.7) satisfying  $(\phi_1)$ - $(\phi_2)$ . Then,  $\tilde{\Phi}$  satisfies the  $\Delta_2$ -condition.

*Proof.* See [11, Lemma 2.7].

**Lemma A.4.** If  $\Phi$  is an N-function of the form (1.7) satisfying  $(\phi_1)$ - $(\phi_2)$ , then the spaces  $L^{\Phi}(\mathcal{O})$  and  $W^{1,\Phi}(\mathcal{O})$  are reflexive.

*Proof.* The proof follows directly from Lemmas A.1, A.2 and A.3.

For our study, it is useful to consider the following lemma whose proof is left for the reader to verify.

**Lemma A.5.** Let  $\Phi$  be an N-function of the type (1.7) satisfying  $(\phi_1)$ - $(\phi_2)$ . Then the following inequalities hold

- (a)  $\Phi(|a+b|) \le 2^m (\Phi(|a|) + \Phi(|b|))$  for all  $a, b \in \mathbb{R}$ .
- (b)  $\phi(|z|)z.(w-z) \leq \Phi(|w|) \Phi(|z|)$  for all  $w, z \in \mathbb{R}^N$  with  $z \neq 0$ , where "." denotes the usual inner product in  $\mathbb{R}^N$ .

Acknowledgments: The authors would like to thank to the anonymous referee for their invaluable suggestions, which significantly contributed to the enhancement of this paper.

## References

- F. Alessio, C. Gui and P. Montecchiari, Saddle solutions to Allen-Cahn equations in doubly periodic media, Indiana Univ. Math. J. 65, 2016, 1, 199-221.
- [2] S. Allen and J. Cahn, A microscopic theory for the antiphase boundary motion and its application to antiphase domain coarsening, Acta Metallurgica, 27, 1979, 1085-1095. 2
- [3] C. O. Alves, Existence of a heteroclinic solution for a double well potential equation in an infinite cylinder of ℝ<sup>N</sup>, Adv. Nonlinear Stud. 19, 2019, 1, 133-147. 2, 3, 6, 21
- [4] C. O. Alves, R. Isneri and P. Montecchiari, Existence of Heteroclinic and Saddle type solutions for a class of quasilinear problems in whole ℝ<sup>2</sup>, Communications in Contemporary Mathematics, 26, 2024, 02, 2250061. 6, 14, 15, 25
- [5] C. O. Alves and R. Isneri, Existence of heteroclinic solutions for the prescribed curvature equation, J. Differential Equations, 362, 2023, 484-513. 2, 3, 4, 6, 27, 28
- [6] C. O. Alves, R. Isneri and P. Montecchiari, Existence of Saddle-type solutions for a class of quasilinear problems in ℝ<sup>2</sup>, Topol. Methods Nonlinear Anal., 61, 2023, 2, 825-868. 6
- [7] D. Bonheure, F. Obersnel and P. Omari, Heteroclinic solutions of the prescribed curvature equation with a double-well potential, Differential Integral Equations, 26, 2013, 11-12, 1411-1428. 3, 6
- [8] Y. Brenier, Optimal transportation and applications, Extended Monge-Kantorovich theory, 1813, 2003, 91-121. 3
- [9] R. Finn, Equilibrium Capillary Surfaces, Springer Science & Business Media, 284, 2012. 2
- [10] J. Frenkel and T. Kontorova, On the theory of plastic deformation and twinning, Acad. Sci. U.S.S.R. J. Phys., 1, 1939, 137-149. 2
- [11] N. Fukagai, M. Ito and K. Narukawa, Positive Solutions of Quasilinear Elliptic Equations with Critical Orlicz-Sobolev Nonlinearity on R<sup>N</sup>, Funkcial. Ekvac., 49, 2006, 2, 235-267. 32
- [12] C. F. Gauss, Principia generalia theoriae figurae fluidorum in statu aequilibrii, Comment. Soc. Regiae Scient. Gottingensis Rec. 7, 1830, 1-53. 2
- [13] E. Giusti and G. H. Williams, Minimal Surfaces and Functions of Bounded Variation, Boston: Birkhäuser, vol. 80, 1984. 3
- [14] M. A. Krasnosels'kii and J. Rutic'kii, Convex functions and Orlicz spaces, Noordhoff, Groningen, 1961. 32
- [15] A. Kurganov and P. Rosenau, On reaction processes with saturating diffusion, Nonlinearity 19, 2006, 1, 171-193. 3
- [16] P. S. Laplace, Traité de mécanique céleste: suppléments au Livre X, Gauthier-Villars, Paris (1806) 2
- [17] G. M. Lieberman, The natural generalization of the natural conditions of Ladyzhenskaya and Ural'tseva for elliptic equations, Comm. Partial Differential Equations 16(2-3) (1991), 311-361. 15, 28
- [18] W. A. J. Luxemburg. Banach function spaces, thesis Delft, Inst. of Techn, 1955. 31
- [19] F. Obersnel and P. Omari, Positive solutions of the Dirichlet problem for the prescribed mean curvature equation, J. Differential Equations. 249, 2010, 7, 1674-1725. 3
- [20] W. Orlicz. Über eine gewisse Klasse von Räumen vom Typus B, Bull. Int. Acad. Polon. Sci. A, 1932, 8/9, 207-220. 30, 31
- [21] W. Orlicz. Ueber Raume (L<sup>M</sup>), Bull. Int. Acad. Polon. Sci. A, 1936, 93-107. 30, 31
- [22] P. H. Rabinowitz, On a class of nonlinear Schrödinger equations, Z. Angew. Math. Phys, 43, 1992, 2, 270-291. 21
- [23] P. H. Rabinowitz, Solutions of heteroclinic type for some classes of semilinear elliptic partial differential equations, J. Math. Sci. Univ. Tokyo, 1, 1994, 3, 525-550. 2, 3, 6

- [24] P. H. Rabinowitz and E. Stredulinsky, Mixed states for an Allen-Cahn type equation, Comm. Pure Appl. Math. 56, 2003, 8, 1078-1134. 2
- [25] M. N. Rao and Z. D. Ren, Theory of Orlicz Spaces, Marcel Dekker, New york, 1985. 32
- [26] N. S. Trudinger, On Harnack type inequalities and their application to quasilinear elliptic equations, Comm. Pure Appl. Math, 20, 1967, 721-747. 5, 15
- [27] T. Young, As essay on the cohesion of fluids, Philos. Trans. Roy. Soc. Lond. 95, 1805, 65-87. 2

(Claudianor O. Alves) UNIDADE ACADÊMICA DE MATEMÁTICA UNIVERSIDADE FEDERAL DE CAMPINA GRANDE, 58429-970, CAMPINA GRANDE - PB - BRAZIL Email address: coalves@mat.ufcg.edu.br

(Renan J. S. Isneri) UNIDADE ACADÊMICA DE MATEMÁTICA UNIVERSIDADE FEDERAL DE CAMPINA GRANDE, 58429-970, CAMPINA GRANDE - PB - BRAZIL Email address: renan.isneri@academico.ufpb.br