RESONANCE, SYZYGIES, AND RANK-3 ULRICH BUNDLES ON THE DEL PEZZO THREEFOLD V_5

MARIAN APRODU AND YEONGRAK KIM

Dedicated to the memory of Gianfranco Casnati

ABSTRACT. We investigate a geometric criterion for a smooth curve C of genus 14 and degree 18 to be described as the zero locus of a section in an Ulrich bundle of rank 3 on a del Pezzo threefold $V_5 \subset \mathbb{P}^6$. The main challenge is to read off the Pfaffian quadrics defining V_5 from geometric structures of C. We find that this problem is related to the existence of a special rank-two vector bundle on C with trivial resonance. This answers a question posed by Ciliberto-Flamini-Knutsen, in the case of degree 5 del Pezzo threefolds. From an explicit calculation of the Betti table of such a curve, we also deduce the uniqueness of the del Pezzo threefold.

1. INTRODUCTION

Let $X \subset \mathbb{P}^N$ be a projective variety of dimension n. A coherent sheaf \mathcal{E} on X is called an *Ulrich sheaf* if there is a finite linear projection $\pi : \mathbb{P}^N \dashrightarrow \mathbb{P}^n$ which is defined over X such that $\pi_*\mathcal{E} \simeq \mathcal{O}_{\mathbb{P}^n}^{\oplus k}$ for some positive integer k [ES03]. In commutative algebra, an analogous object often called an *Ulrich module* appears in the form of maximal Cohen-Macaulay modules having the maximal number of generators, and has been prominently featured in modern mathematics due to its various spectrum of applications, see for instance [Ulr84, BGS87, IMW22, Ma23]. Eisenbud and Schreyer transferred this concept in the geometric setup and discovered a remarkable connection with Cayley–Chow forms [ES03]; see also [Bea18, CMP21] for recent developments from the geometric perspective and further applications.

The most significant open problem in the Ulrich sheaf theory is the *Eisenbud–Schreyer conjecture* which predicts that any smooth projective variety must carry an Ulrich bundle. If true, this conjecture would

²⁰²⁰ Mathematics Subject Classification. Primary 14J60, Secondary 13D02, 14M15, 14H10.

Key words and phrases. Ulrich bundle, quintic del Pezzo threefold, Resonance variety.

have dramatic consequences on the expected shapes of cohomology tables [ES03, ES11]. The next step, assuming the existence was established, is to determine the possible ranks of Ulrich bundles and classify them. While all these problems are widely open, they have been resolved in a number of relevant cases, in particular, for varieties of minimal degree [AHMP19, ES03]. Hence, it is natural to raise the same questions in the next case, namely, for del Pezzo varieties. These problems are rather straightforward for smooth del Pezzo varieties of dimension n = 1, 2 [ES03, Corollary 4.5, 6.5; Bea18, Proposition 5]. More recently, the answer was also given for del Pezzo threefolds of degree das in various case studies: [Bea00, CH12, LMS15] for d = 3; [CKL21] for d = 4; [Fae05, LP21] for d = 5; [CFM16] and [CFM18] for two cases of d = 6; [CFM17] for d = 7; and [ES03, Corollary 5.3, Proposition 5.11] for d = 8. In particular, X carries Ulrich bundles of every rank $r \ge 2$ unless $X = \nu_2(\mathbb{P}^3) \subset \mathbb{P}^9$ is the 2-uple embedding of \mathbb{P}^3 where it only carries Ulrich bundles of even ranks. Very recently, Ciliberto, Flamini, and Knutsen found a single deformation argument for constructing Ulrich bundles of rank 2 and 3 over every del Pezzo threefold of degree $3 \le d \le 7$ [CFK23].

There were several attempts to construct Ulrich bundles directly using the Hartshorne-Serre correspondence, for example, [AC00] over a del Pezzo threefold of Picard number 1 and [Bea18, Proposition 8] over every del Pezzo threefold for rank 2 Ulrich bundles. The problem translates into the existence of an elliptic normal curve of degree d+2contained in a del Pezzo threefold $V_d \subset \mathbb{P}^{d+1}$ of degree d. For rank-3 Ulrich bundles, we refer to [CH12] for d = 3, and to an unpublished additional note to [CKL21] for d = 4. In this case, we need to look for an arithmetically Cohen-Macaulay curve $C \subset V_d \subset \mathbb{P}^{d+1}$ of genus 2d + 4 and degree 3d + 3 such that the twisted canonical line bundle $\omega_C(-1)$ has two global sections which generate the module of twisted global sections $H^0_*(\omega_C) = \bigoplus_{j \in \mathbb{Z}} H^0(C, \omega_C(j))$. In loc. cit., the authors showed that a general curve C of genus 2d + 4 with d = 3 or d = 4admits an embedding $C \hookrightarrow V_d \subset \mathbb{P}^{d+1}$ satisfying these conditions, and the incidence set $\{(C, V_d) \mid C \subset V_d \subset \mathbb{P}^{d+1}\}$ dominates the family of del Pezzo threefolds $\{V_d\}$ of degree d. In particular, a general del Pezzo threefold of degree d has an Ulrich bundle of rank 3. We remark that the results of [CH12, CKL21] are deeply related to the rationality problem for the Hurwitz spaces $\mathcal{H}(k, 2g + 2k - 2)$ and the moduli of smooth curves \mathcal{M}_q as in computational approaches developed by Geiß and Schrever [Gei12; CH12, Appendix].

The main purpose of this paper is to answer the analogous question for Ulrich bundles of rank 3 on a del Pezzo threefold $V_5 \subset \mathbb{P}^6$ of degree 5. As pointed out in [CFK23, Section 5], a general curve of genus 14 in \mathbb{P}^6 cannot be associated to an Ulrich bundle of rank 3 on V_5 , which represents a significant difference from the cases d = 3 and d = 4. On the other hand, any model of $V_5 \subset \mathbb{P}^6$ has an Ulrich bundle \mathcal{E} of rank 3, and two generic global sections s_1, s_2 will give rise an arithmetically Cohen-Macaulay (ACM for short) curve $C \subset V_5 \subset \mathbb{P}^6$ of genus 14, degree 18 as the zero locus of $s_1 \wedge s_2$. Hence, it is natural to inquire about the geometry of the sublocus of \mathcal{M}_{14} which parametrizes curves that induce Ulrich bundles of rank 3 over some $V_5 \subset \mathbb{P}^6$. Following the notation in [CFK23, Section 5], this question is the same as studying the image of $\varphi_{3,5} : \mathbb{G} \longrightarrow \mathcal{M}_{14}$.

The outline of the paper is as follows. In Section 2 we recall the connection between rank-3 Ulrich bundles on V_d and certain curves contained in V_d . For d = 5, which is the focus of the present work, we compute the Betti table of these curves, Proposition 3 and we compute the dimension of the locus of these curves in the moduli space, Proposition 4. Section 3 is devoted to the main result of the paper, Theorem 9. It shows that the source of an embedding in V_5 of a curve C of genus 14 and degree 18 in \mathbb{P}^6 is the existence of a rank-2 vector bundle on C with 5-dimensional space of global sections, determinant equal to $\mathcal{O}_C(1)$ and trivial resonance. In fact, the theorem proves that a rank-2 bundle with determinant $\mathcal{O}_{C}(1)$ and exactly five independent global sections gives an embedding of the curve in a linear section of the Grassmannian, and this section is smooth if and only if the resonance of the given bundle is trivial. For the proof of Theorem 9 we need an algebraic characterization of the triviality of the resonance in terms of generalized Pfaffians, which is given in Theorem 7. We conclude the paper with some remarks on the main theorem.

Throughout this paper, we work over \mathbb{C} for simplicity.

Acknowledgments. M. A. was partly supported by the PNRR grant CF 44/14.11.2022 *Cohomological Hall algebras of smooth surfaces and applications*. Y. K. is supported by Basic Science Research Program of the NRF of Korea (NRF-2022R1C1C1010052). Y. K. thanks Mateusz Michalek for his hospitality and helpful discussion.

2. Rank 3 Ulrich bundles on del Pezzo threefolds of Picard number 1

In this section, we recall the construction of rank-3 Ulrich bundles on del Pezzo threefolds of Picard number 1. We start with the general definition of Ulrich bundles on a (smooth) projective variety $X \subset \mathbb{P}^N$. Note that it depends on an embedding i.e. on the choice of a polarization $(X, \mathcal{O}_X(1))$ where $\mathcal{O}_X(1) = \mathcal{O}_{\mathbb{P}^N}(1)|_X$ is a very ample line bundle on X.

Definition 1. Let $X \subseteq \mathbb{P}^N$ be a projective variety of dimension n, and let \mathcal{E} be a nonzero coherent sheaf on X.

(i) \mathcal{E} is called ACM if \mathcal{E} is locally Cohen-Macaulay, and has no intermediate cohomology, that is,

$$H^i(X, \mathcal{E}(-j)) = 0$$

for every 0 < i < n and $j \in \mathbb{Z}$.

(ii) \mathcal{E} is called *Ulrich* if \mathcal{E} has the same cohomological behavior as the structure sheaf of \mathbb{P}^n , that is,

$$H^i(X, \mathcal{E}(-j)) = 0$$

for every $i \in \mathbb{Z}$ and $1 \leq j \leq n$.

This definition admits several variants, see for instance [ES03, Proposition 2.1]. We refer to [CH12, Section 2] for basic properties of Ulrich sheaves. Note that any Ulrich sheaf over a smooth projective variety is locally free, and hence in the smooth case it makes sense to speak about Ulrich bundles.

An *n*-dimensional projective variety $X \subseteq \mathbb{P}^N$ is called a *del Pezzo* variety if $\omega_X \simeq \mathcal{O}_X(1 - \dim X)$. It is known that there are exactly four types of smooth del Pezzo threefolds of Picard number 1, namely,

- $V_3 \subset \mathbb{P}^4$ a cubic threefold;
- $V_4 \subset \mathbb{P}^5$ a complete intersection of two quadrics;
- $V_5 \subset \mathbb{P}^6$ a 3-dimensional linear section of the Grassmannian $\operatorname{Gr}(2,5) \subset \mathbb{P}^9;$
- $v_2(\mathbb{P}^3) \subset \mathbb{P}^9$ a 2-uple embedding of \mathbb{P}^3 .

The Ulrich geography for the Veronese threefold in \mathbb{P}^9 is very simple, it has a unique Ulrich bundle of rank 2 and no Ulrich bundle of odd ranks [ES03, Proposition 5.11, Corollary 5.3].

The remaining cases $V_d \subseteq \mathbb{P}^{d+1}$ are del Pezzo threefolds of degree d = 3, 4, 5 embedded by a very ample line bundle $\mathcal{O}_{V_d}(1)$ which is a generator of the Picard group $\operatorname{Pic}(V_d)$ in each of the cases. Arrondo and Costa classified possible Chern classes of ACM bundles of small ranks over V_d [AC00, Theorem 3.4, 4.9], and constructed some of them in their list. For instance, a general global section of an Ulrich bundle \mathcal{E} of rank 2 over V_d degenerates along an elliptic normal curve in \mathbb{P}^{d+1} , which is of degree d + 2.

By the Hartshorne-Serre correspondence (see [Arr07] for more details), the existence of rank-2 Ulrich bundles over V_d reduces to the question whether or not there exist elliptic normal curves contained in V_d . We refer to [AC00, Example 3.3; Bea18, Proposition 8] for the construction of rank-2 Ulrich bundles on V_d in this manner.

Similarly, the existence problem for rank-3 Ulrich bundles over V_d is translated in terms of the existence of certain curves lying on V_d . Let \mathcal{E} be an Ulrich bundle of rank 3 on a del Pezzo threefold $V_d \subset \mathbb{P}^{d+1}$ of degree d, and let $V \subset H^0(V_d, \mathcal{E})$ be a general subspace of dimension 2. The inclusion $V \otimes \mathcal{O}_{V_d} \to \mathcal{E}$ degenerates along a smooth ACM curve C_d of genus 2d + 4 and degree 3d + 3 (cf. [CFK23, Theorem 3.5]) such that $\omega_{C_d}(-1)$ is generated by a 2-dimensional space W and the multiplication map $\nu : W \otimes H^0(\mathbb{P}^{d+1}, \mathcal{O}_{\mathbb{P}^{d+1}}(j)) \to H^0(\omega_{C_d}(j-1))$ has full rank for every $j \in \mathbb{Z}$. Thanks to the base-point-free pencil trick, the last condition is satisfied when the multiplication map $W \otimes$ $H^0(C_d, \mathcal{O}_{C_d}(1)) \to H^0(C_d, \omega_{C_d})$ is surjective.

Conversely, if there is such a curve C_d embedded in V_d , the Hartshorne-Serre correspondence implies the existence of a rank 3 vector bundle \mathcal{E} which fits into the short exact sequence

$$0 \to \mathcal{O}_{V_d}^{\oplus 2} \to \mathcal{E} \to \mathcal{I}_{C_d/V_d}(3) \to 0.$$

A direct computation shows that \mathcal{E} is an Ulrich bundle on V_d . We remark that the generating condition on $\omega_{C_d}(-1)$ is essential, see [CH12, Remark 5.5] for a discussion on this issue.

Constructions of rank-3 Ulrich bundles on V_3 and V_4 using syzygies of algebraic curves and the Hartshorne-Serre correspondence are well understood and are contained in the paper by Casanellas-Hartshorne-Geiß-Schreyer [CH12, Proposition 5.4, Appendix A], and in an unpublished appendix to the paper by Cho, the second named author, and Lee [CKL21]. The key idea behind these constructions was first made clear by Geiß in his Ph. D. dissertation [Gei13, Section 3.4.2], in relation with the rationality problem for the moduli space of curves \mathcal{M}_g and for the Hurwitz space $\mathcal{H}(k, 2g + 2k - 2)$, for small values of g, k.

We briefly describe these constructions below.

- (i) Construct a random curve C_d of genus g = 2d + 4 together with a base-point-free pencil \mathfrak{g}_{d+3}^1 using a bigraded projective model in $\mathbb{P}^1 \times \mathbb{P}^2$. Note that such a pencil of degree d + 3 always exists thanks to the Brill-Noether theory.
- (ii) The linear system $|\omega_{C_d} \mathfrak{g}_{d+3}^1|$ is a very ample $\mathfrak{g}_{3d+3}^{d+1}$ on C_d . Verify that $C_d \subset \mathbb{P}^{d+1}$ is contained in a smooth del Pezzo threefold V_d .

- (iii) Verify that C_d induces an Ulrich bundle of rank 3 via the Hartshorne-Serre correspondence.
- (iv) Verify that the incidence family $\{(C_d, V_d) \mid C_d \subset V_d\}$ from this construction dominates the space of del Pezzo threefolds $\{V_d\}$.

As a result, we see that a general V_d carries rank-3 Ulrich bundles for d = 3 and for d = 4.

The case of the del Pezzo threefold $V_5 \subset \mathbb{P}^6$ of degree 5, the central object in this paper, is treated separately. First note that $V_5 =$ $\operatorname{Gr}(2,5) \cap H_1 \cap H_2 \cap H_3 \subset \mathbb{P}^6$ is a smooth \mathbb{P}^6 -section of the Grassmannian $\operatorname{Gr}(2,5) \subset \mathbb{P}^9$ of lines in \mathbb{P}^4 . From the Euler sequence over the Grassmannian $\operatorname{Gr}(2,5)$, we have the following short exact sequence of vector bundles on V_5

$$0 \to \mathcal{U} \to V \otimes \mathcal{O}_{V_5} \to \mathcal{Q} \to 0$$

so that \mathcal{U} is the restriction of the universal subbundle on $\operatorname{Gr}(2,5)$ and \mathcal{Q} is the restriction of the quotient bundle on $\operatorname{Gr}(2,5)$. Thanks to the Bott formula, one readily verifies that the bundles \mathcal{U} and \mathcal{Q}^{\vee} are ACM bundles on V_5 . Furthermore, Orlov showed that there is a full exceptional collection in the derived category of coherent sheaves on V_5 [Orl91], namely,

$$D^b(V_5) = \langle \mathcal{U}, \mathcal{Q}^{\vee}, \mathcal{O}_{V_5}, \mathcal{O}_{V_5}(1) \rangle.$$

We refer to [Fae05, Section 3; LP21, Section 2.3] for more details on the notion and computations for cohomology groups of these bundles.

It is well-known that V_5 supports an Ulrich bundle \mathcal{E} of rank 3, namely, $\operatorname{Sym}^2(\mathcal{U}^*)$ and its deformations. Note also that there is a 10-dimensional family of stable Ulrich bundles of rank 3 from the computation $h^1(\mathcal{E} \otimes \mathcal{E}^{\vee}) = 10$ [AC00, Example 4.4; Fae05, Proposition 6.8; LP21, Theorem 1.1]. In particular, Ulrich bundles on V_5 have a very nice presentation using universal bundles as follows.

Proposition 2. [LP21, Proposition 3.4] For any $r \ge 2$, an Ulrich bundle \mathcal{E} of rank r on V_5 corresponds to a quiver representation and fits into the short exact sequence

$$0 \to \mathcal{U}^{\oplus r} \to \mathcal{Q}^{\vee \oplus r} \to \mathcal{E}(-1) \to 0.$$

We now consider an analogous construction using the Hartshorne-Serre correspondence. Note that two generic global sections of an Ulrich bundle \mathcal{E} of rank 3 degenerates along an ACM curve $C \subset V_5$ which fits into the short exact sequence

$$0 \to \mathcal{O}_{V_5}^{\oplus 2} \to \mathcal{E} \to \mathcal{I}_{C/V_5}(3) \to 0.$$

 $\mathbf{6}$

From this short exact sequence, we may read off algebraic information about C as follows.

Proposition 3. Notation as above. C is an ACM curve of genus 2d+4 = 14 and degree 3d+3 = 18 in \mathbb{P}^6 . The Betti table of C is given by

$j \setminus i$	0	1	2	3	4	5
0	1	_	_	_	_	_
1	—	5	5	_	_	_
2	—	13	45	56	25	—
3	—	—	—	—	—	2

In particular, C can only lie on one single del Pezzo threefold in \mathbb{P}^6 .

Proof. The minimal free resolutions of $2\mathcal{O}_{V_5}$ and of \mathcal{E} are

$$0 \to 2\mathcal{O}_{\mathbb{P}^6}(-5) \to 10\mathcal{O}_{\mathbb{P}^6}(-3) \to 10\mathcal{O}_{\mathbb{P}^6}(-2) \to 2\mathcal{O}_{\mathbb{P}^6} \to 2\mathcal{O}_{V_5} \to 0,$$

and

$$0 \to 15\mathcal{O}_{\mathbb{P}^6}(-3) \to 45\mathcal{O}_{\mathbb{P}^6}(-2) \to 45\mathcal{O}_{\mathbb{P}^6}(-1) \to 15\mathcal{O}_{\mathbb{P}^6} \to \mathcal{E} \to 0,$$

respectively. The mapping cone gives a free resolution of $\mathcal{I}_{C/V_5}(3)$ as

$$0 \to 2\mathcal{O}_{\mathbb{P}^6}(-5) \to 25\mathcal{O}_{\mathbb{P}^6}(-3) \to 55\mathcal{O}_{\mathbb{P}^6}(-2)$$
$$\to 45\mathcal{O}_{\mathbb{P}^6}(-1) \to 13\mathcal{O}_{\mathbb{P}^6} \to \mathcal{I}_{C/V_5}(3) \to 0.$$

We recover a free resolution of C in \mathbb{P}^6 from the short exact sequence $0 \to \mathcal{I}_{V_5/\mathbb{P}^6} \to \mathcal{I}_{C/\mathbb{P}^6} \to \mathcal{I}_{C/V_5} \to 0$, and note that its Betti table coincides with the Betti table in the conclusion. Since $C \subset V_5$, the ideal of C and V_5 share the same quadrics, and thus there are 5 linear syzygies among these quadrics, hence, the free resolution of C we obtained above is indeed minimal.

The last part of the conclusion follows immediately again by the fact that the ideal of V_5 in \mathbb{P}^6 is generated by five quadrics.

Note that if we have such a curve $C \subset V_5$, then we can reverse the construction and obtain an Ulrich bundle of rank 3 on V_5 . To apply the Hartshorne-Serre correspondence, we only need to show that $\omega_C(-1)$ has two global sections which generate the graded module $H^0_*(\omega_C)$. Since C is an ACM curve, it suffices to establish the surjectivity of the multiplication maps

$$H^0(C, \omega_C(-1)) \otimes H^0(C, \mathcal{O}_C(j)) \to H^0(C, \omega_C(j-1))$$

for all j > 0. Thanks to the base–point–free pencil trick, it reduces to the surjectivity of the multiplication map

$$\mu: H^0(C, \omega_C(-1)) \otimes H^0(C, \mathcal{O}_C(1)) \to H^0(C, \omega_C),$$

which, by dimension reasons, is equivalent to the injectivity of μ in this case. By considering the dual resolution of C obtained by applying Hom_{P⁶} $(-, \omega_{\mathbb{P}^6})$ which gives a minimal free resolution for ω_C , we notice that the line bundle $\omega_C(-1)$ is globally generated by two sections. Furthermore, the kernel of the multiplication map μ equals to the Koszul cohomology group $K_{1,-1}(C, \omega_C, \mathcal{O}_C(1))$. By Green's duality [Gre84, Theorem 2.c.6], [Gre89, p. 188], we have an isomorphism

$$K_{1,-1}(C, \omega_C, \mathcal{O}_C(1)) \simeq K_{4,3}(C, \mathcal{O}_C(1))^{\vee} = 0.$$

Hence, such a curve C provides a rank-3 bundle \mathcal{E} on V_5 as an extension $0 \to \mathcal{O}_{V_5}^{\oplus 2} \to \mathcal{E} \to \mathcal{I}_{C/V_5}(3) \to 0$. It is straightforward that this \mathcal{E} is Ulrich.

From the construction, one can prove that the family of ACM curves associated to rank 3 Ulrich bundles on V_5 cannot dominate \mathcal{M}_{14} . This fact distinguishes the del Pezzo threefold V_5 from V_3 and V_4 .

Proposition 4. The locus of curves C in \mathcal{M}_{14} which can be obtained as the degeneracy locus of two global sections of an Ulrich bundle of rank 3 on $V_5 \subset \mathbb{P}^6$ has dimension 33.

Proof. Note first that the moduli of smooth curves of genus 14 has dimension dim $\mathcal{M}_{14} = 39$, and the Brill-Noether number is $\rho(14, 6, 18) = 0$. Hence, the family of smooth projective curves $\{C \subset \mathbb{P}^6\}$ of genus 14 and degree 18 curves in \mathbb{P}^6 has dimension

$$\dim(\mathcal{M}_{14}) + \dim(\mathrm{PGL}(7)) = 39 + 48 = 87.$$

Since any del Pezzo variety $V_5 \subset \mathbb{P}^6$ is a smooth linear section of the Grassmannian $\operatorname{Gr}(2,5) \subset \mathbb{P}^5$, and is projectively equivalent to each other [Isk80, Theorem II-1.1], there are $45 = \dim(\operatorname{PGL}(7)) - \dim(\operatorname{Aut}(V_5))$ dimensional family of del Pezzo threefolds of degree 5 in \mathbb{P}^6 . Let us choose and fix a model, and call it $V_5 \subset \mathbb{P}^6$ again. There is a 10-dimensional family of rank 3 Ulrich bundles \mathcal{E} on V_5 [Fae05, Proposition 6.8-(v); LP21, Theorem 1.1], and for each choice of \mathcal{E} , a general choice of 2-dimensional subspace $V \subset H^0(\mathcal{E})$ defines an ACM curve C as the degeneracy locus of the evaluation map $0 \to V \otimes \mathcal{O}_{V_5} \to \mathcal{E}$. Hence, from Proposition 3, the family of ACM curves

 $\{C \subset \mathbb{P}^6 \mid C \text{ is associated to a rank-3 Ulrich bundle } \mathcal{E} \text{ on some } V_5\}$ has dimension

$$45 + 10 + \dim(\operatorname{Gr}(2, H^0(\mathcal{E}))) = 45 + 10 + 26 = 81$$

Choose a dominating component \mathcal{X} of this family, and consider the curves C parametrized by this component. Since such a C is stable, the fiber of $\mathcal{X} \to \mathcal{M}_{14}$ containing C has dimension $48 = \dim(\mathrm{PGL}(7))$.

We conclude that the image of these curves in \mathcal{M}_{14} has dimension 81 - 48 = 33.

Proposition 4 implies that a general curve $C \subset \mathbb{P}^6$ of genus 14, degree 18 does not appear as a subvariety of a smooth del Pezzo threefold $V_5 \subset \mathbb{P}^6$. This phenomenon was also observed by Ciliberto, Flamini, and Knutsen [CFK23, Section 5] using Grassmannian bundles over the moduli of smooth curves \mathcal{M}_g . Alternatively, one can compare the Betti table of a curve on V_5 as found in Proposition 3 to the Betti table of a general curve (2).

Hence, it is natural to ask the following question whose answer is the main objective of this paper.

Question 5. Describe geometric conditions on a curve C of genus 14 so that C can be embedded into $C \hookrightarrow V_5 \subset \mathbb{P}^6$ and C is the degeneracy locus of two sections of some Ulrich bundles of rank 3 on V_5 .

3. Vanishing resonance and embeddings of curves into V_5

We show next that the answer to Question 5 is related to the existence of a certain rank-2 vector bundle with trivial resonance. The first part of the section is algebraic in nature and is applicable to various other situations. We shall utilize these algebraic facts in the proof of the main result of the paper, Theorem 9 that represents the content of the second part of this section.

3.1. Resonance of a skew-symmetric matrix. We recast here the definition of resonance from [PS15]. Instead of working with subspaces, for practical reasons, we prefer working with maps and matrices. We consider V an n-dimensional complex vector space with $n \geq 4$, W another complex vector space, and $\partial : \wedge^2 V^{\vee} \to W^{\vee}$ a linear map. Denote the kernel of ∂ by K^{\perp} . We define the *resonance variety of* ∂ to be the affine variety

 $\mathcal{R}(\partial) := \{ a \in V^{\vee} \mid \exists \ b \in V^{\vee} \text{ such that } a \land b \neq 0, \text{ and } \partial(a \land b) = 0 \} \cup \{0\}.$

It coincides with the usual resonance variety (see [PS15]) $\mathcal{R}(V, K)$ of the pair (V, K) where K^{\vee} is the image of the map ∂ . It was shown in [AFRS23] that the resonance carries a natural scheme structure, however, we will not be concerned with this finer structure here, and we simply consider the reduced structure.

Example 6. Let X be a complex algebraic variety, and let \mathcal{F} be a vector bundle on X. Consider the second determinant map

$$d_2: \wedge^2 H^0(X, \mathcal{F}) \to H^0(X, \wedge^2 \mathcal{F}).$$

The resonance variety $\mathcal{R}(X, \mathcal{F})$ of \mathcal{F} is by definition $\mathcal{R}(d_2)$. In the rank-two case, the map d_2 coincides with the determinant map

$$\det : \wedge^2 H^0(X, \mathcal{F}) \to H^0(X, \det(\mathcal{F})).$$

We refer to [AFRW24] and [AFRS23] for an extensive discussion on the resonance of vector bundles and several applications.

We fix a basis $\{x_1, \ldots, x_n\}$ in V, denote by $\{e_1, \ldots, e_n\}$ its dual basis in V^{\vee} , and put $z_{ij} = e_i \wedge e_j$. Denote also by $A = (\ell_{ij})_{i,j=\overline{1,n}}$ the skewsymmetric matrix of linear forms on W associated to the map ∂ in the chosen bases, i.e. $\ell_{ij} := \partial(e_i \wedge e_j) \in W^{\vee}$.

Recall from [KS89] that A is said to have a generalized zero if there exists an invertible matrix U such that the skew-symmetric matrix UAU^t has a zero off the diagonal. It means, up to a base-change in V, the matrix associated to ∂ has a zero off the diagonal. A 4 × 4 Pfaffian of such a matrix UAU^t is called a generalized Pfaffian of A.

The generalized Pfaffians are directly connected to the resonance, as shown below.

Theorem 7. Notation as above. The following are equivalent:

- (i) The resonance $\mathcal{R}(\partial)$ is nontrivial;
- (ii) The matrix A has a generalized zero;
- (iii) There exists a 4×4 generalized Pfaffian of A of rank at most four.

Proof. (i) \Rightarrow (ii). Let e_1 and e_2 be two non-collinear vectors in V^{\vee} such that $\partial(e_1 \wedge e_2) = 0$. We complete $\{e_1, e_2\}$ to a basis $\{e_1, \ldots, e_n\}$ of V^{\vee} . In this new basis, the matrix associated to ∂ has a zero element as its (1, 2)-th entry, and hence A has a generalized zero.

(ii) \Rightarrow (iii). Without loss of generality, we may assume $\ell_{12} = 0$. Consider the submatrix

(1)
$$A' = \begin{pmatrix} 0 & \ell_{12} & \ell_{13} & \ell_{14} \\ -\ell_{12} & 0 & \ell_{23} & \ell_{24} \\ -\ell_{13} & -\ell_{23} & 0 & \ell_{34} \\ -\ell_{14} & -\ell_{24} & -\ell_{34} & 0 \end{pmatrix}$$

of A. Since $\ell_{12} = 0$ it follows that its Pfaffian equals $\ell_{14}\ell_{23} - \ell_{13}\ell_{24}$ which is a quadric of rank ≤ 4 .

(iii) \Rightarrow (i). Recall that the definition of the resonance is independent on the choice of a basis in V, and hence it only depends on the equivalence class of A. In particular, we may assume that A has a 4×4 Pfaffian of rank ≤ 4 and we want to prove that $\mathbb{P}K^{\perp} \cap \operatorname{Gr}(2, V^{\vee}) \neq \emptyset$ in $\mathbb{P}(\bigwedge^2 V^{\vee})$. Without loss of generality, we may assume n = 4 and A looks like in (1), i.e. A = A'. In this case, the Grassmanian $\operatorname{Gr}(2, V)$ is the quadric in $\mathbb{P}(\bigwedge^2 V)$ given by the equation Q = 0 where $Q = z_{12}z_{34} - z_{13}z_{24} + z_{14}z_{23}$. Its dual quadric is the Grassmannian $\operatorname{Gr}(2, V^{\vee})$.

We note that the dimension of K cannot be six, for in this case, the rank of the Pfaffian would be six, too. If the dimension of Kis ≤ 4 , then the dimension of K^{\perp} would be at least two, and hence $\mathbb{P}K^{\perp} \cap \operatorname{Gr}(2, V^{\vee}) \neq \emptyset$.

We may assume then that K is five-dimensional, and hence $\mathbb{P}K^{\perp}$ is a point in $\mathbb{P}(\bigwedge^2 V^{\vee})$ which corresponds to a hyperplane H of $\mathbb{P}(\bigwedge^2 V)$. We prove that this point belongs to $\operatorname{Gr}(2, V^{\vee})$, equivalently H is tangent to the dual quadric $\operatorname{Gr}(2, V)$.

Via the identification of K with the hyperplane H via the dual of ∂ , the rank of the Pfaffian of A equals the rank of $Q|_H$ and the hypothesis implies that the hyperplane section $H \cap \text{Gr}(2, V)$ is a singular quadric. This is only possible when H is tangent to the quadric Gr(2, V), which we wanted to prove.

Remark 8. In the previous result, the notion of generalized Pfaffians when dim V > 4 is used in an essential way. We notice that the ranks of 4×4 –Pfaffians might change in the equivalence class of A. Let

$$A = \begin{pmatrix} 0 & x_0 & x_1 & x_2 & x_3 \\ -x_0 & 0 & x_2 & x_3 & x_4 \\ -x_1 & -x_2 & 0 & x_4 & x_5 \\ -x_2 & -x_3 & -x_4 & 0 & x_0 + x_5 \\ -x_3 & -x_4 & -x_5 & -x_0 - x_5 & 0 \end{pmatrix}$$

be a 5×5 skew-symmetric matrix of linear forms in \mathbb{P}^5 . One can immediately check that the five 4×4 -Pfaffians of A are quadrics of rank ≥ 5 (indeed, two are of rank 6 and the other three are of rank 5).

On the other hand, these Pfaffian quadrics define a smooth del Pezzo surface $S_5 \subset \mathbb{P}^5$ of degree 5. Since every smooth del Pezzo surface of degree 5 are projectively equivalent to each other, and the ideal defining S_5 can be generated by 5 Pfaffian quadrics of rank ≤ 4 after a suitable base-change [MP23, Theorem 6.1; KMP23, Proposition 6.3-(2)].

3.2. Curves on V_5 . We focus next on finding geometric conditions on C so that C can be embedded into a del Pezzo threefold $V_5 \subset \mathbb{P}^6$. Let us recall the generic behavior of a curve of genus 14 and its projective model in \mathbb{P}^6 embedded by a complete linear system $|\mathcal{O}_C(1)|$ of degree 18. Since $\mathcal{O}_C(j)$ is nonspecial for $j \geq 2$, we may read off its Hilbert

function so that the Hilbert series of C is

$$HS(\mathcal{O}_C) = \frac{1 - 5t^2 - 8t^3 + 45t^4 - 56t^5 + 25t^6 - 2t^8}{(1 - t)^7}$$

Hence, we expect that such a generic curve $C \subset \mathbb{P}^6$ has the Betti table

	$j \setminus i$	0	1	2	3	4	5
	0	1	—	—	—	—	_
(2)	1	_	5	_	_	_	_
	2	_	8	45	56	25	_
	3	—	_	_	_	_	2

These 5 quadrics define a reducible curve of degree 32 containing C which is a union of C and a residual curve P of genus 8 and degree 14. Indeed, a general curve $C \subset \mathbb{P}^6$ of genus 14 and degree 18 has the same Betti table as above, and is obtained as the residual curve of a paracanonical curve $P \subset \mathbb{P}^6$ of genus 8 and degree 14 [Ver05, Theorem 4.5, 5.14].

The key issue is to develop a criterion to distinguish whether or not a given curve $C \subset \mathbb{P}^6$ can be embedded into a del Pezzo threefold $V_5 \subset \mathbb{P}^6$. If this is the case, the 5 quadrics defining C will also define V_5 , which is a linear section of the Grassmannian $\operatorname{Gr}(2,5)$. In particular, these quadrics can be obtained as Pfaffians of a 5×5 skew–symmetric matrix by the Buchsbaum-Eisenbud structure theorem [BE77, Theorem 2.1]. Hence, the core of the question is knowing how to extract information about Pfaffian quadrics that can generate (linear sections of) the Grassmannian $\operatorname{Gr}(2,5)$ from a given curve $C \subset \mathbb{P}^6$. We see that the resonance of a certain vector bundle on C nicely captures the required information as in the next few paragraphs.

Theorem 9. Let $C \subset \mathbb{P}^6$ be an ACM curve of genus 14, degree 18, and $K_{4,3}(C, \mathcal{O}_C(1)) = 0$. Then the following are equivalent:

- (i) $C \subset V_5$ for some del Pezzo threefold $V_5 \subset \mathbb{P}^6$;
- (ii) $C \subset V_5$ for some del Pezzo threefold V_5 , and is the zero locus of 2 global sections of an Ulrich bundle \mathcal{E} of rank 3 on V_5 ;
- (iii) there is a rank 2 vector bundle \mathcal{F} on C such that $\wedge^2 \mathcal{F} = \mathcal{O}_C(1)$, $h^0(C, \mathcal{F}) = 5$, and $\mathcal{R}(C, \mathcal{F}) = 0$.

Proof. (i) \Rightarrow (ii). Since C is ACM and 4-regular, we must have

$$\dim K_{5,3}(C, \mathcal{O}_C(1)) = 2$$

from the Hilbert function of $C \subset \mathbb{P}^6$. This implies that $\omega_C(-1)$ is globally generated and $h^0(C, \omega_C(-1)) = 2$ by considering the dual resolution as we discussed above. Moreover, we have $K_{1,-1}(C, \omega_C, \mathcal{O}_C(1)) \simeq$

 $K_{4,3}(C, \mathcal{O}_C(1))^{\vee} = 0$, and thus the multiplication map

$$\mu: H^0(C, \omega_C(-1)) \otimes H^0(\mathbb{P}^6, \mathcal{O}_{\mathbb{P}^6}(1)) \to H^0(C, \omega_C)$$

is an isomorphism. Hence, C provides a vector bundle ${\mathcal E}$ of rank 3 on V_5 so that

$$0 \to \mathcal{O}_{V_5}^{\oplus 2} \to \mathcal{E} \to \mathcal{I}_{C/V_5}(3) \to 0$$

by the Hartshorne-Serre correspondence. It is clear that \mathcal{E} is Ulrich.

(ii) \Rightarrow (iii). Recall that V_5 is a linear section of the Grassmannian $\operatorname{Gr}(2,5)$, so let us denote the (restriction of) universal subbundle on V_5 by \mathcal{U} . We choose $\mathcal{F} = \mathcal{U}^{\vee}|_C$ as the restriction of the dual of the universal bundle \mathcal{U} . Since $\wedge^2 \mathcal{U}^{\vee} = \mathcal{O}_{V_5}(1)$, the restriction \mathcal{F} is a rank 2 vector bundle with $\wedge^2 \mathcal{F} = \mathcal{O}_C(1)$ as well.

We first claim that $h^0(V_5, \mathcal{U}^{\vee}) = h^0(C, \mathcal{F}) = 5$. From the exact sequence

$$0 \to \mathcal{O}_{V_5}^{\oplus 2} \to \mathcal{E} \to \mathcal{I}_{C/V_5}(3) \to 0,$$

we have a 4-term exact sequence

$$0 \to \mathcal{U}^{\vee}(-3)^{\oplus 2} \to \mathcal{E} \otimes \mathcal{U}^{\vee}(-3) \to \mathcal{U}^{\vee} \to \mathcal{F} \to 0$$

by tensoring $\mathcal{U}^{\vee}(-3)$. Since $H^1(V_5, \mathcal{U}^{\vee}) = 0$, it is enough to show that $H^1(V_5, \mathcal{E} \otimes \mathcal{U}^{\vee}(-3)) = H^2(V_5, \mathcal{U}^{\vee}(-3)) = 0$ to verify the claim $h^0(C, \mathcal{F}) = h^0(V_5, \mathcal{U}^{\vee}) = 5$. It is well-known that the universal bundle \mathcal{U} is an ACM bundle on V_5 , and thus $\mathcal{U}^{\vee} \simeq \mathcal{U}(1)$ has no intermediate cohomology, in particular, $H^2(V_5, \mathcal{U}^{\vee}(-3)) = 0$. For the vanishing of $H^1(V_5, \mathcal{E} \otimes \mathcal{U}^{\vee}(-3))$, we use the correspondence between Ulrich bundles on V_5 and quiver representations in [LP21, Proposition 3.4]. In particular, we have the short exact sequence $0 \to \mathcal{U}^{\oplus 3} \to \mathcal{Q}^{\vee \oplus 3} \to \mathcal{E}(-1) \to 0$, so the desired cohomology vanishing is provided when

$$H^1(V_5, \mathcal{Q}^{\vee} \otimes \mathcal{U}^{\vee}(-2)) = H^2(V_5, \mathcal{U} \otimes \mathcal{U}^{\vee}(-2)) = 0.$$

By the Serre duality, it is equivalent to the vanishing $H^2(V_5, \mathcal{Q} \otimes \mathcal{U}) = H^1(V_5, \mathcal{U} \otimes \mathcal{U}^{\vee}) = 0$. For the first vanishing, we have $H^2(V_5, \mathcal{Q} \otimes \mathcal{U}) = \text{Ext}_{V_5}^2(\mathcal{Q}^{\vee}, \mathcal{U}) = 0$ which is used to prove that \mathcal{U} is left-orthogonal to \mathcal{Q}^{\vee} . The second vanishing is also the key for the fact that \mathcal{U} is an exceptional bundle in $D^b(V_5)$, and thus $\text{Ext}_{V_5}^i(\mathcal{U}, \mathcal{U}) = 0$ for $i \neq 0$.

Next, we compute the resonance variety $\mathcal{R}(C, \mathcal{F})$. Suppose that it is nonzero. Let det : $\wedge^2 H^0(C, \mathcal{F}) \to H^0(C, \mathcal{O}_C(1)) \simeq H^0(\mathbb{P}^6, \mathcal{O}_{\mathbb{P}^6}(1))$ be the determinant map. Note that \mathcal{F} defines a nonzero Koszul cycle in $K_{2,1}(C, \mathcal{O}_C(1))$ as in [Voi05, Section 1; AN10, Section 3.4.1]. In particular, the 5 quadrics in the ideal I(C) of C can be obtained as 4-Pfaffians of the map det. By Theorem 7, there is a 4-Pfaffian of det of rank ≤ 4 among them. However, the quadrics in the ideal I(C)and $I(V_5)$ are the same, and there is no quadric of rank ≤ 4 in $I(V_5)$ [KMP23, Proposition 6.3-(3)] which contradicts to the assumption. We conclude that the resonance $\mathcal{R}(C, \mathcal{F}) = 0$ is zero.

Note that such a vector bundle \mathcal{F} does not exist over a general curve $C \subset \mathbb{P}^6$ of genus 14 and degree 18. Suppose that $C \subset \mathbb{P}^6$ has such a rank 2 vector bundle \mathcal{F} . Then \mathcal{F} defines a nonzero Koszul cycle in $K_{2,1}(C, \mathcal{O}_C(1))$, in particular, the 5 quadrics in the ideal I(C) of C must allow a linear syzygy. On the other hand, the quadrics in the ideal of a general model $\widetilde{C} \subset \mathbb{P}^6$ of genus 14 and degree 18 have no linear syzygies among them.

(iii) \Rightarrow (i). Let det : $\wedge^2 H^0(C, \mathcal{F}) \to H^0(C, \mathcal{O}_C(1)) \simeq H^0(\mathbb{P}^6, \mathcal{O}_{\mathbb{P}^6}(1))$ be the determinant map, which is represented by a 5×5 skew-symmetric matrix composed of linear forms in \mathbb{P}^6 . We follow the construction of Grassmannian syzygies introduced by Koh-Stillman [KS89, Lemma 1.3]. The five principal 4-Pfaffians of det are contained in I(C), we consider the variety X defined by these Pfaffian quadrics which contains C as its subvariety. It is clear that X is a \mathbb{P}^6 -linear section of the Grassmannian Gr(2,5), and hence it remains to show that X is isomorphic to a smooth del Pezzo threefold $V_5 \subset \mathbb{P}^6$.

Since the resonance $\mathcal{R}(C, \mathcal{F})$ is trivial, we see that the map det is nondegenerate and surjective by Theorem 7. Via this surjection, we have an inclusion $H^0(\mathbb{P}^6, \mathcal{O}_{\mathbb{P}^6}(1))^{\vee} \subset \wedge^2 H^0(C, \mathcal{F})^{\vee}$. Moreover, $\mathcal{R}(C, \mathcal{F}) = 0$ also implies

$$Y = \mathbb{P}(H^0(\mathcal{O}_{\mathbb{P}^6}(1))^{\vee \perp}) \cap \operatorname{Gr}(2, H^0(\mathcal{F}))$$
$$= \mathbb{P}(\ker(\det)) \cap \operatorname{Gr}(2, H^0(\mathcal{F})) = \emptyset$$

in $\mathbb{P}(\wedge^2 H^0(C,\mathcal{F})).$

By the projective duality, the linear section of the Grassmannian $X = \operatorname{Gr}(2, H^0(C, \mathcal{F})^{\vee}) \cap \mathbb{P}(H^0(\mathbb{P}^6, \mathcal{O}_{\mathbb{P}^6}(1))^{\vee})$ is smooth and (dimensionally) transverse if and only if Y is smooth and transverse [DK18, Proposition 2.24]. In our case, $H^0(\mathbb{P}^6, \mathcal{O}_{\mathbb{P}^6}(1))^{\vee} \subset \wedge^2 H^0(C, \mathcal{F})^{\vee}$ is a linear subspace of codimension 3, being Y smooth and transverse is equivalent to $Y = \emptyset$ as similar as in [Kuz18, Lemma 3.3]. We conclude that X is a smooth 3-dimensional linear section of the Grassmannian $\operatorname{Gr}(2,5)$, and thus projectively equivalent to a smooth del Pezzo three-fold $V_5 \subset \mathbb{P}^6$ [Isk80, Theorem II-1.1].

Remark 10. The condition $K_{4,3}(C, \mathcal{O}_C(1)) = 0$ is essential. Indeed, by following the instructions in [AC00, Example 4.4], one can construct an ACM curve C of genus 14 and degree 18 contained in a smooth del Pezzo threefold V_5 but $K_{4,3}(C, \mathcal{O}_C(1)) \neq 0$. In this example, two global sections of $\omega_C(-1)$ have a linear syzygy which implies dim $K_{4,3}(C, \mathcal{O}_C(1)) = 1$, and such C does not satisfy the generating condition to apply the Hartshorne-Serre correspondence. We refer to [CH12, Remark 5.5; CKL21, p. 17284] for details since the reasons are exactly the same.

Remark 11. By using similar techniques as in the proof above, one can also show that $h^0(V_5, \mathcal{U} \otimes \mathcal{U}^{\vee}) = h^0(C, \mathcal{F} \otimes \mathcal{F}^{\vee}) = 1$ where \mathcal{F} is a rank 2 vector bundle on C that appeared in statement of the theorem. In particular, \mathcal{F} is a simple vector bundle on C.

References

- [AHMP19] M. Aprodu, S. Huh, F. Malaspina, and J. Pons-Llopis, Ulrich bundles on smooth projective varieties of minimal degree, Proc. Amer. Math. Soc. 147 (2019), 5117–5129.
- [AFRW24] M. Aprodu, G. Farkas, C. Raicu, and J. Weyman, Koszul modules with vanishing resonance in algebraic geometry, Selecta Math. 30 (2024), Paper No. 24, 33pp
- [AFRS23] M. Aprodu, G. Farkas, C. Raicu, and A. Suciu, *Reduced resonance* schemes and Chen ranks, preprint available at arXiv:2303.07855.
- [AN10] M. Aprodu and J. Nagel, Koszul cohomology and algebraic geometry, Univ. Lecture Ser. 52, American Mathematical Society, Providence, RI (2010).
- [AC00] E. Arrondo and L. Costa, Vector bundles on Fano 3-folds without intermediate cohomology, Comm. Algebra 28 (2000), 3899–3911.
- [Arr07] E. Arrondo, A home-made Hartshorne-Serre correspondence, Rev. Mat. Compult. **20** (2007), 423–443.
- [Bea00] A. Beauville, determinantal hypersurfaces, Michigan Math. J. 48 (2000), 39–64.
- [Bea18] A. Beauville, An introduction to Ulrich bundles, Eur. J. Math. 4 (2018), 26–36.
- [BE77] D. A. Buchsbaum and D. Eisenbud, Algebra structure for finite free resolutions, and some structure theorems for ideals of codimension 3, Amer. J. Math. 99 (1977), 447–485.
- [BGS87] R.-O. Buchweitz, G.-M. Greuel, and F.-O. Schreyer, Cohen-Macaulay modules on hypersurface singularities II, Invent. Math. 88 (1987), 165– 182.
- [CH12] M. Casanellas and R. Hartshorne, Stable Ulrich bundles, Internat. J. Math. 23 (2012), 1250083.
- [CFM16] G. Casnati, D. Faenzi, and F. Malaspina, Moduli spaces of rank two aCM bundles on the Segre product of three projective lines, J. Pure Appl. Algebra 220 (2016), 1554–1575.
- [CFM18] G. Casnati, D. Faenzi, and F. Malaspina, Rank two aCM bundles on the del Pezzo fourfold of degree 6 and its general hyperplane section, J. Pure Appl. Algebra 222 (2018), 585–609.
- [CFM17] G. Casnati, M. Filip, and F. Malaspina, Rank two aCM bundles on the del Pezzo threefold of degree 7, Rev. Math Compultense 37 (2017), 129–165.

- [CFK23] C. Ciliberto, F. Flamini, and A. L. Knutsen, Ulrich bundles on Del Pezzo threefolds, J. Algebra 634 (2023), 209–236.
- [CKL21] Y. Cho, Y. Kim, and K.-S. Lee, Ulrich bundles on intersections of two 4dimensional quadrics, Int. Math. Res. Not. IMRN (2021), 17277–17303.
- [CMP21] L. Costa, R. M. Miró-Roig, and J. F. Pons-Llopis, Ulrich bundles: From commutative algebra to algebraic geometry, De gruyter Studies in Mathematics 77, De Gruyter, Berlin (2021).
- [DK18] O. Debarre and A. Kuznetsov, Gushel-Mukai varieties: classification and birationalities, Algebr. Geom. 5 (2018), 15–76.
- [ES03] D. Eisenbud and F.-O. Schreyer, Resultants and Chow forms via exterior syzygies, J. Amer. Math. Soc. 16 (2003), 537–579.
- [ES11] D. Eisenbud and F.-O. Schreyer, *Boij-Söderberg theory*, in: Combinatorial aspects of commutative algebra and algebraic geometry, Abel Symp. 6, Springer, Berlin (2011), 35–48.
- [Fae05] D. Faenzi, Bundles over the Fano threefold V_5 , Comm. Algebra **33** (2005), 3061–3080.
- [Gei12] F. Geiß, The unirationality of Hurwitz spaces of 6-gonal curves of small genus, Doc. Math. 17 (2012), 627–640.
- [Gei13] F. Geiß, The unirationality of Hurwitz spaces of hexagonal curves of small genus, Ph. D. Dissertation in Universität des Saarlandes (2013).
- [Gre84] M. L. Green, Koszul cohomology and the geometry of projective varieties, J. Differential Geom. 19 (1984), 125–171.
- [Gre89] M. L. Green, Koszul cohomology and the geometry, Lectures on Riemann Surfaces, Proceedings of the College on Riemann Surfaces, ICTP, Italy, 9 Nov-18 Dec 1987, 177-200 (1989).
- [Isk80] V. A. Iskovskih, Anticanonical models of three-dimensional algebraic varieties, J. Soviet Math. 13 (1980), 745–814.
- [IMW22] S. B. Iyengar, L. Ma, and M. E. Walker, Multiplicities and Betti numbers in local algebra via lim Ulrich points, Algebra Number Theory 16 (2022), 1213–1257.
- [KMP23] Y. Kim, H. Moon, and E. Park, Some remarks on the $\mathcal{K}_{p,1}$ -theorem, preprint avilable at arXiv:2404.03293.
- [KS89] J. Koh and M. Stillman, Linear syzygies and line bundles on an algebraic curve, J. Algebra 125 (1989), 120–132.
- [Kuz18] A. Kuznetsov, On linear sections of the spinor tenfold. I., Izv. Math. 82 (2018), 694–751.
- [LMS15] M. Lahoz, E. Macrì, and P. Stellari, Arithmetically Cohen-Macaulay bundles on cubic threefolds, Algebr. Geom. 2 (2015), 231–269.
- [LP21] K.-S. Lee and K.-D. Park, Moduli spaces of Ulrich bundles on the Fano 3-fold V₅, J. Algebra 574 (2021), 262–277.
- [Ma23] L. Ma, Lim Ulrich sequences and Lech's conjecture, Invent. Math. 231 (2023), 407–429.
- [MP23] H. Moon and E. Park, On the rank index of some quadratic varieties, Mediterr. J. Math. 20 (2023), Paper No. 260, 19 pp.
- [Orl91] D. O. Orlov, Exceptional set of vector bundles on the variety V_5 , Moscow Univ. Math. Bull. **46** (1991), 48–50.
- [PS15] S. Papadima and A. Suciu, Vanishing resonance and representations of Lie algebras, J. Reine Angew. Math. 706 (2015), 83–101.

- [Ulr84] B. Ulrich, Gorenstein rings and modules with high numbers of generators, Math. Z. 188 (1984), 23–32.
- [Ver05] A. Verra, The unirationality of the moduli spaces of curves of genus 14 or lower, Compos. Math. 141 (2005), 1425–1444.
- [Voi05] C. Voisin, Green's canonical syzygy conjecture for generic curves of odd genus, Compos. Math. 141 (2005), 1163–1190.

UNIVERSITY OF BUCHAREST, DEPARTMENT OF MATHEMATICS & INSTITUTE OF MATHEMATICS "SIMION STOILOW" OF THE ROMANIAN ACADEMY

Email address: marian.aprodu@fmi.unibuc.ro, marian.aprodu@imar.ro

DEPARTMENT OF MATHEMATICS & INSTITUTE OF MATHEMATICAL SCIENCE, PUSAN NATIONAL UNIVERSITY, 2 BUSANDAEHAK-RO 63BEON-GIL, GEUMJEONG-GU, 46241 BUSAN, KOREA

Email address: yeongrak.kim@pusan.ac.kr