# Low-rank alternating direction doubling algorithm for solving large-scale continuous time algebraic Riccati equations

Juan ${\rm Zhang}^{1*}$  and Wenlu  ${\rm Xun}^2$ 

<sup>1</sup>Key Laboratory of Intelligent Computing and Information Processing of Ministry of Education, Hunan Key Laboratory for Computation and Simulation in Science and Engineering, School of Mathematics and Computational Science, Xiangtan University, Xiangtan, Hunan, China. <sup>2</sup>School of Mathematics and Computational Science, Xiangtan University, Xiangtan, Hunan, China.

\*Corresponding author(s). E-mail(s): zhangjuan@xtu.edu.cn;

#### Abstract

This paper proposes an effective low-rank alternating direction doubling algorithm (R-ADDA) for computing numerical low-rank solutions to large-scale sparse continuous-time algebraic Riccati matrix equations. The method is based on the alternating direction doubling algorithm (ADDA), utilizing the low-rank property of matrices and employing Cholesky factorization for solving. The advantage of the new algorithm lies in computing only the  $2^{k}$ -th approximation during the iterative process, instead of every approximation. Its efficient low-rank formula saves storage space and is highly effective from a computational perspective. Finally, the effectiveness of the new algorithm is demonstrated through theoretical analysis and numerical experiments.

**Keywords:** low-rank alternating direction doubling algorithm, Riccati matrix equation, ADDA, Cholesky factorization

## 1 Introduction

In this paper, we consider the large-scale continuous-time algebraic Riccati equation (CARE):

$$A^T X + XA + Q - XGX = 0, (1)$$

and the complementary equation of (1) is

$$AY + YA^T - YQY + G = 0, (2)$$

where  $A, G, Q \in \mathbb{R}^{n \times n}, B \in \mathbb{R}^{n \times m}, C \in \mathbb{R}^{p \times n}, Q = C^T C$ , and  $G = BB^T$ , with  $m, p \ll n$ .

The CARE (1) mainly arises in the context of the quadratic optimal control problem for the following continuous-time linear time-invariant control system:

$$\begin{cases} \dot{x}(t) = Ax(t) + Bu(t), & x(0) = x_0, \\ y(t) = Cx(t). \end{cases}$$

Here,  $x(t) \in \mathbb{R}^n$  is the state vector,  $u(t) \in \mathbb{R}^m$  is the control vector, and  $y(t) \in \mathbb{R}^p$  is the output vector. The objective of quadratic optimal control is to find a control u(t) that minimizes the following function:

$$J(x_0, u) = \frac{1}{2} \int_0^{+\infty} (y(t)^T y(t) + u(t)^T u(t)) dt$$

Assuming that (A, B) is stabilizable and (C, A) is detectable, there exists a unique optimal solution  $\bar{u}$  that minimizes the functional  $J(x_0, u)[1]$ . Furthermore, this optimal solution can be determined using the feedback operator PP, such that  $\bar{u}(t) = Px(t)$ , where  $P = B^T$  and  $X \in \mathbb{R}^{n \times n}$  is the unique symmetric positive semi-definite stable solution of the CARE (1).

We consider the  $2n \times 2n$  Hamiltonian matrix H related to the CARE (1):

$$H = \begin{pmatrix} A & -G \\ -Q & -A^T \end{pmatrix},\tag{3}$$

which satisfies the relation

$$HJ = -JH^T, \quad J = \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix},$$

where  $I_n$  denotes the identity matrix of order n.

Laub[2] proposed a numerically backward stable algorithm that applies a reordered QR algorithm[3–5] to the eigenvalue problem  $Hx = \lambda x$  for computing X. Unfortunately, the QR algorithm does not preserve the structure of the Hamiltonian matrix H and the splitting of its eigenvalues. Ammar and Mehrmann[6] introduced a structured-preserving algorithm that utilizes orthogonal symplectic transformations to compute a basis for the stable invariant subspace of H. Byers[14] presented a stable symplectic

orthogonal method, but it is only applicable to systems with a single input or output. Over the past few decades, numerous iterative methods have been proposed for solving algebraic Riccati equations. The Newton method has been widely applied in the literature[8–11]. Mehrmann and Tan[12] also proposed a refinement method to correct approximate solution deficiencies. These methods require a good initial approximate solution and can be viewed as iterative refinement methods that can be combined with other direct methods. Gardiner and Laub[13, 14] extended the Structured Preserved Matrix Sign Function Method (MSGM)[21, 22].

Additionally, a class of methods known as doubling algorithms (DA)[15] garnered widespread interest in the 1970s and 1980s. These methods stem from fixed-point iterations derived from discrete-time algebraic Riccati equations (DARE):

$$X_{k+1} = \widehat{A}^T X_k (I + \widehat{G} X_k)^{-1} \widehat{A} + \widehat{Q}.$$

Doubling algorithms generate sequences  $X_{2^k}$  instead of sequences  $X_k$ . It is necessary to transform CAREs into DAREs for post-processing. However, the convergence of this algorithm has only been proven when  $\hat{A}$  is non-singular[15],  $(\hat{A}, \hat{G}, \hat{Q})$  is stable and detectable[16]. The structurally preserved doubling algorithm (SDA), proposed by Guo, Lin, and Xu [17], leverages the Sherman–Morrison–Woodbury formula and various iterations of sparse and low-rank representations. The resulting large-scale doubling algorithm exhibits O(n) computational complexity and memory requirements per iteration and converges essentially quadratically. Furthermore, Li, Kuo, and Lin [18] introduced a structurally preserved doubling algorithm (SDA-ls- $\varepsilon$ ) for solving large-scale nonsymmetric algebraic Riccati equations. They conducted a detailed error analysis of the iterative truncation effect on the approximate solution obtained by SDA.

In this paper, based on the alternating direction doubling algorithm (ADDA) proposed by Wang[19], we introduce a new low-rank doubling algorithm, called R-ADDA. By transforming the Hamiltonian matrix into skew-Hermitian matrix pairs using the corresponding Cayley transformation, the low-rank alternating direction doubling algorithm inherits good convergence properties. Numerical results demonstrate that the algorithm exhibits competitiveness and effectiveness.

The structure of the remaining sections of this paper is as follows. In section 2, we present the iterative framework for the low-rank alternating direction doubling algorithm for solving CARE (1) and introduce the R-ADDA algorithm. Section 3 provides theoretical proofs of the structural preservation properties and convergence results of the new algorithm. In section 4, we demonstrate the efficiency of the R-ADDA algorithm through numerical experiments. Finally, in section 5, we present some conclusions and remarks to conclude this paper.

In this paper, we introduce some necessary symbols and terminology. Let  $\mathbb{R}^{n \times m}$  denote the set of all  $n \times m$  real matrices. For any matrices  $A = [a_{ij}], B = [b_{ij}] \in \mathbb{R}^{n \times m}$ , if  $a_{ij} \geq b_{ij}$   $(a_{ij} > b_{ij})$  for all i, j, we denote  $A \geq B$  (A > B) and define  $|A| \doteq [|a_{ij}|]$ . The symbols  $A^T$  and  $A^{-1}$  represent the transpose and inverse of matrix A, respectively. Additionally,  $\rho(A) = \max_{1 \leq i \leq n} \{|\lambda_i(A)|\}$  denotes the spectral radius of  $A, ||A||_2$  represents the 2-norm of matrix A, and  $A \otimes I$  denotes the Kronecker product of A and I.

## 2 Low-rank alternating direction doubling algorithm

**Definition 1.** For  $M, L \in \mathbb{R}^{2n \times 2n}$ , let  $M - \lambda L$  be a symplectic matrix pencil, i.e.,

$$MJM^T = LJL^T, \quad J = \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix},$$

 $and \ define$ 

$$\mathcal{N}(M,L) = \left\{ [M_*, L_*] : M_*, L_* \in \mathbb{R}^{2n \times 2n}, \ rank[M_*, L_*] = 2n, \ [M_*, L_*] \begin{bmatrix} L \\ -M \end{bmatrix} = 0 \right\} \neq \varnothing.$$

**Definition 2.** For any given  $[M_*, L_*] \in \mathcal{N}(M, L)$ , define

$$\widehat{M} = M_*M, \quad \widehat{L} = L_*L.$$

The transformation

$$M - \lambda L \to \widehat{M} - \lambda \widehat{L}$$

is called a doubling transformation.

An important characteristic of this transformation is that it preserves the structure, eigenvalue spaces, and squares of eigenvalues.

Assuming  $X \ge 0$  is a non-negative solution to the CARE (1), then the CARE (1) can be rewritten as

$$H\begin{pmatrix}I\\X\end{pmatrix} = \begin{pmatrix}I\\X\end{pmatrix}R.$$
(4)

Similarly, (2) yields

$$H\begin{pmatrix} -Y\\I \end{pmatrix} = \begin{pmatrix} -Y\\I \end{pmatrix} (-S),\tag{5}$$

where

$$R = A - GX, \quad S = A^T - QY,$$

and the matrix H is the Hamiltonian matrix defined in (3).

By choosing appropriate parameters  $\alpha > 0$  to optimize conditions for inverting certain matrices, and using Cayley transformation, we can transform (4) and (5) into the following forms

$$(H + \alpha I) \begin{pmatrix} I \\ X \end{pmatrix} (R - \alpha I) = (H - \alpha I) \begin{pmatrix} I \\ X \end{pmatrix} (R + \alpha I),$$
$$(H + \alpha I) \begin{pmatrix} -Y \\ I \end{pmatrix} (-S - \alpha I) = (H - \alpha I) \begin{pmatrix} -Y \\ I \end{pmatrix} (-S + \alpha I).$$

If  $R - \alpha I$  and  $S - \alpha I$  are nonsingular, then we have

$$(H + \alpha I) \begin{pmatrix} I \\ X \end{pmatrix} = (H - \alpha I) \begin{pmatrix} I \\ X \end{pmatrix} \mathscr{C}(R; \alpha),$$
  

$$(H + \alpha I) \begin{pmatrix} -Y \\ I \end{pmatrix} \mathscr{C}(S; \alpha) = (H - \alpha I) \begin{pmatrix} -Y \\ I \end{pmatrix},$$
(6)

where

$$\mathscr{C}(R;\alpha) = (R+\alpha I)(R-\alpha I)^{-1} = (R-\alpha I)^{-1}(R+\alpha I),$$
  
$$\mathscr{C}(S;\alpha) = (S+\alpha I)(S-\alpha I)^{-1} = (S-\alpha I)^{-1}(S+\alpha I).$$

Assuming  $A - \alpha I$  is nonsingular, and let

6

$$A_{\alpha} = A - \alpha I, \quad \widetilde{A}_{\alpha} = A + \alpha I, \quad U_{\alpha} = A_{\alpha}^{T} + Q A_{\alpha}^{-1} G, \quad V_{\alpha} = A_{\alpha} + G A_{\alpha}^{-T} Q,$$
$$Z_{1} = \begin{pmatrix} A_{\alpha}^{-1} & 0 \\ Q A_{\alpha}^{-1} & I \end{pmatrix}, \quad Z_{2} = \begin{pmatrix} I & 0 \\ 0 & -U_{\alpha}^{-1} \end{pmatrix}, \quad Z_{3} = \begin{pmatrix} I & A_{\alpha}^{-1} G \\ 0 & I \end{pmatrix},$$
we can prove that

then w р

$$M_{0} = Z_{3}Z_{2}Z_{1}(H + \alpha I) = \begin{pmatrix} \widehat{A}_{0} & 0 \\ -X_{0} & I \end{pmatrix},$$

$$L_{0} = Z_{3}Z_{2}Z_{1}(H - \alpha I) = \begin{pmatrix} I & Y_{0} \\ 0 & \widehat{A}_{0}^{T} \end{pmatrix},$$

$$\widehat{A}_{0} = I + 2\alpha V_{\alpha}^{-1},$$

$$X_{0} = 2\alpha U_{\alpha}^{-1}QA_{\alpha}^{-1},$$

$$Y_{0} = 2\alpha A_{\alpha}^{-1}GU_{\alpha}^{-1}.$$
(7)

(7)

where

Next, utilizing the Sherman–Morrison–Woodbury (SMW) formula

$$(A + UV^{T})^{-1} = A^{-1} - A^{-1}U(I + V^{T}A^{-1}U)^{-1}V^{T}A^{-1},$$

we can efficiently compute  $U_{\alpha}^{-1}$  and  $V_{\alpha}^{-1},$  then

$$U_{\alpha}^{-1} = A_{\alpha}^{-T} - A_{\alpha}^{-T} C^{T} (I_{p} + CA_{\alpha}^{-1} GA_{\alpha}^{-T} C^{T})^{-1} CA_{\alpha}^{-1} GA_{\alpha}^{-T},$$
(8)

$$V_{\alpha}^{-1} = A_{\alpha}^{-1} - A_{\alpha}^{-1} B (I_m + B^T A_{\alpha}^{-T} Q A_{\alpha}^{-1} B)^{-1} B^T A_{\alpha}^{-T} Q A_{\alpha}^{-1}.$$
 (9)

We take

$$D_{0} = A_{\alpha}^{-T}C^{T}, \quad P_{0} = A_{\alpha}^{-1}B,$$
  

$$\Sigma_{0} = 2\alpha[I_{p} - (I_{p} + D_{0}^{T}GD_{0})^{-1}D_{0}^{T}GD_{0}],$$
  

$$\Gamma_{0} = 2\alpha[I_{m} - P_{0}^{T}QP_{0}(I_{m} + P_{0}^{T}QP_{0})^{-1}],$$
(10)

then we can get

$$X_0 = D_0 \Sigma_0 D_0^T, \quad Y_0 = P_0 \Gamma_0 P_0^T, \tag{11}$$

where  $Q_0$  and  $P_0$  are full-column rank matrices. Multiplying both sides of (6) by  $Z_3Z_2Z_1$ , we get:

$$M_0 \begin{pmatrix} I \\ X \end{pmatrix} = L_0 \begin{pmatrix} I \\ X \end{pmatrix} \mathscr{C}(R; \alpha),$$
  

$$M_0 \begin{pmatrix} -Y \\ I \end{pmatrix} \mathscr{C}(S; \alpha) = L_0 \begin{pmatrix} -Y \\ I \end{pmatrix}.$$
(12)

This iterative method can construct a pair of sequences  $\{M_k, L_k\}$ , for  $k = 0, 1, 2, \cdots$ , such that:

$$M_{k}\begin{pmatrix} I\\ X \end{pmatrix} = L_{k}\begin{pmatrix} I\\ X \end{pmatrix} [\mathscr{C}(R;\alpha)]^{2^{k}},$$

$$M_{k}\begin{pmatrix} -Y\\ I \end{pmatrix} [\mathscr{C}(S;\alpha)]^{2^{k}} = L_{k}\begin{pmatrix} -Y\\ I \end{pmatrix}.$$
(13)

Here,  $M_k$  and  $L_k$  have the same form as  $M_0$  and  $L_0$ , such as:

$$M_k = \begin{pmatrix} \widehat{A}_k & 0 \\ -X_k & I \end{pmatrix}, \quad L_k = \begin{pmatrix} I & Y_k \\ 0 & \widehat{A}_k^T \end{pmatrix}.$$

Next, we need construct  $\{M_{k+1}, L_{k+1}\}$  to find suitable  $\widetilde{M}, \widetilde{L} \in \mathbb{R}^{2n \times 2n}$  such that:

$$\operatorname{rank}((\widetilde{M}, \ \widetilde{L})) = 2n, \ \ (\widetilde{M}, \ \widetilde{L}) \begin{pmatrix} L_k \\ -M_k \end{pmatrix} = 0.$$

Let  $M_{k+1} = \widetilde{M}M_k$  and  $L_{k+1} = \widetilde{L}L_k$ , through calculations we can obtain:  $\{\widetilde{M}, \widetilde{L}\},\$ 

$$\widetilde{M} = \begin{pmatrix} \widehat{A}_k (I_n + Y_k X_k)^{-1} & 0\\ -\widehat{A}_k^T (I_n + X_k Y_k)^{-1} X_k & I_n \end{pmatrix}, \quad \widetilde{L} = \begin{pmatrix} I_n & \widehat{A}_k Y_k (I_n + X_k Y_k)^{-1}\\ 0 & \widehat{A}_k^T (I_n + X_k Y_k)^{-1} \end{pmatrix}.$$

Then, we can get the iterative format of the Alternating Direction Doubling Algorithm (ADDA) as follows:

$$\widehat{A}_{k+1} = \widehat{A}_k (I_n + Y_k X_k)^{-1} \widehat{A}_k, 
X_{k+1} = X_k + \widehat{A}_k^T (I_n + X_k Y_k)^{-1} X_k \widehat{A}_k, 
Y_{k+1} = Y_k + \widehat{A}_k Y_k (I_n + X_k Y_k)^{-1} \widehat{A}_k^T.$$
(14)

Obviously, from the iterative format (14), we can see that this algorithm has a computational complexity of  $O(n^3)$ . By leveraging the low-rank properties of  $Q = C^T C$ and  $G = BB^T$ , we can derive the iterative framework of the Low-Rank alternating direction doubling algorithm (R-ADDA), such that for  $k = 1, 2, \cdots$ , the R-ADDA iteration follows a recursive form:

$$\widehat{A}_{k} = \widehat{A}_{k-1}^{2} + \widehat{A}_{1k} \widehat{A}_{2k}^{T}, 
X_{k} = D_{k} \Sigma_{k} D_{k}^{T}, 
Y_{k} = P_{k} \Gamma_{k} P_{k}^{T},$$
(15)

where  $\widehat{A}_{ik} \in \mathbb{R}^{n \times p_{k-1}}$   $(i = 1, 2), D_k \in \mathbb{R}^{n \times p_k}, P_k \in \mathbb{R}^{n \times m_k}$ , and  $\Sigma_k \in \mathbb{R}^{p_k \times p_k}, \Gamma_k \in \mathbb{R}^{m_k \times m_k}$ .

In the iteration, for all previous k, we need to store  $D_k, \Sigma_k, P_k, \Gamma_k, \widehat{A}_{ik}$ , and apply the Sherman-Morrison-Woodbury (SMW) formula again to obtain:

$$(I_n + Y_k X_k)^{-1} = I_n - Y_k D_k \Sigma_k (I_{p_k} + D_k^T Y_k D_k \Sigma_k)^{-1} D_k^T$$

$$= I_n - P_k (I_{m_k} + \Gamma_k P_k^T X_k P_k)^{-1} \Gamma_k P_k^T X_k,$$
  
$$(I_n + X_k Y_k)^{-1} = I_n - D_k (I_{p_k} + \Sigma_k D_k^T Y_k D_k)^{-1} \Sigma_k D_k^T Y_k$$
  
$$= I_n - X_k P_k \Gamma_k (I_{m_k} + P_k^T X_k P_k \Gamma_k)^{-1} P_k^T,$$

then we can obtain that

$$\hat{A}_{k+1} = \hat{A}_k [I_n - Y_k D_k \Sigma_k (I_{p_k} + D_k^T Y_k D_k \Sigma_k)^{-1} D_k^T] \hat{A}_k$$
$$= \hat{A}_k [I_n - P_k (I_{m_k} + \Gamma_k P_k^T X_k P_k)^{-1} \Gamma_k P_k^T X_k] \hat{A}_k,$$

$$\begin{aligned} X_{k+1} &= X_k + \widehat{A}_k^T (I_n + X_k Y_k)^{-1} X_k \widehat{A}_k \\ &= X_k + \widehat{A}_k^T X_k \widehat{A}_k - \widehat{A}_k^T D_k (I_{p_k} + \Sigma_k D_k^T Y_k D_k)^{-1} \Sigma_k D_k^T Y_k X_k \widehat{A}_k \\ &= X_k + \widehat{A}_k^T X_k \widehat{A}_k - \widehat{A}_k^T X_k P_k \Gamma_k (I_{m_k} + P_k^T X_k P_k \Gamma_k)^{-1} P_k^T X_k \widehat{A}_k, \end{aligned}$$

$$Y_{k+1} = Y_k + \widehat{A}_k Y_k (I_n + X_k Y_k)^{-1} \widehat{A}_k^T$$
  
=  $Y_k + \widehat{A}_k Y_k \widehat{A}_k^T - \widehat{A}_k Y_k D_k (I_{p_k} + \Sigma_k D_k^T Y_k D_k)^{-1} \Sigma_k D_k^T Y_k \widehat{A}_k^T$   
=  $Y_k + \widehat{A}_k Y_k \widehat{A}_k^T - \widehat{A}_k Y_k X_k P_k \Gamma_k (I_{m_k} + P_k^T X_k P_k \Gamma_k)^{-1} P_k^T \widehat{A}_k^T$ 

Here,  $\oplus$  denotes the direct sum of matrices. Based on the iterative scheme above and reference [18] , we can choose the recursive matrices as follows:

$$\begin{aligned} \widehat{A}_{1,k+1} &= \widehat{A}_k Y_k D_k \Sigma_k (I_{p_k} + D_k^T Y_k D_k \Sigma_k)^{-1} \\ &= \widehat{A}_k P_k (I_{m_k} + \Gamma_k P_k^T X_k P_k)^{-1} \Gamma_k P_k^T D_k \Sigma_k, \\ \widehat{A}_{2,k+1} &= \widehat{A}_k^T D_k, \\ D_{k+1} &= [D_k, \widehat{A}_k^T D_k], \\ P_{k+1} &= [P_k, \widehat{A}_k P_k], \end{aligned}$$
$$\begin{aligned} \Sigma_{k+1} &= \Sigma_k \oplus [\Sigma_k - (I_{p_k} + \Sigma_k D_k^T Y_k D_k)^{-1} \Sigma_k D_k^T Y_k D_k \Sigma_k] \\ &= \Sigma_k \oplus [\Sigma_k - \Sigma_k D_k^T P_{1k} \Gamma_k (I_{m_k} + P_{2k}^T X_k P_k \Gamma_k)^{-1} P_k^T D_k \Sigma_k] \\ &= \Sigma_k \oplus \widetilde{\Sigma}_k, \end{aligned}$$

$$\Gamma_{k+1} = \Gamma_k \oplus [\Gamma_k - \Gamma_k P_k^T D_k (I_{p_k} + \Sigma_k D_k^T Y_k D_k)^{-1} \Sigma_k D_k^T P_k \Gamma_k]$$
  
=  $\Gamma_k \oplus [\Gamma_k - \Gamma_k P_k^T X_k P_k \Gamma_k (I_{m_k} + P_k^T X_k P_k \Gamma_k)^{-1}]$   
=  $\Gamma_k \oplus \widetilde{\Gamma}_k.$ 

Therefore, we can see that in the process of solving large-scale continuous-time algebraic Riccati equations using the R-ADDA iterative method, the dimensions of  $D_k$  and  $P_k$  grow exponentially. Moreover, the following conclusion holds:

$$\operatorname{rank}(X_k) \le \operatorname{rank}(D_k) \le 2^k p, \quad \operatorname{rank}(Y_k) \le \operatorname{rank}(P_k) \le 2^k m,$$

where the number of columns of  $Q_{ik}$  and  $P_{ik}$  are  $2^k p$  and  $2^k m$ , respectively. The advantage of the R-ADDA algorithm depends on the accuracy of its approximations as well as the CPU time and memory requirements.

The following is the R-ADDA algorithm for solving large-scale continuous-time algebraic Riccati equations (1):

Algorithm I It-MDDA algorithm for solving OARD (	g CARE (1)	solving	for a	algorithm	DDA	R-A	1	lgorithm	Al
--------------------------------------------------	------------	---------	-------	-----------	-----	-----	---	----------	----

**Input:** Matrices  $A \in \mathbb{R}^{n \times n}$ ,  $B \in \mathbb{R}^{n \times m}$ ,  $C \in \mathbb{R}^{p \times n}$ , where  $G = C^T C$ ,  $Q = BB^T$ , and parameters  $\alpha$  and residual limit  $\varepsilon$ ;

**Output:**  $X_k = D_k \Sigma_k D_k^T$ , such that  $X_k \approx X$ , where X is the solution of CARE (1). 1: Compute

$$A_{\alpha} = A - \alpha I, \ U_{\alpha} = A_{\alpha}^{T} + Q A_{\alpha}^{-1} G, \ V_{\alpha} = A_{\alpha} + G A_{\alpha}^{-T} Q,$$

and  $A_{\alpha}^{-1}$ ,  $U_{\alpha}^{-1}$ ,  $V_{\alpha}^{-1}$ ; 2: Set k = 0;

3: Compute

$$D_0 = A_{\alpha}^{-T} C^T, \quad P_0 = A_{\alpha}^{-1} B,$$
  

$$\Sigma_0 = 2\alpha [I_p - (I_p + D_0^T G D_0)^{-1} D_0^T G D_0],$$
  

$$\Gamma_0 = 2\alpha [I_m - P_0^T Q P_0 (I_m + P_0^T Q P_0)^{-1}];$$

4: for k = 1 until convergence do 5: Compute  $D_k$ ,  $P_k$ ,  $\hat{A}_{ik}$  (i = 1, 2), and  $\Sigma_k$ ,  $\Gamma_k$ ; 6: Update  $k \leftarrow k + 1$ ,  $X_{k+1} = D_{k+1}\Sigma_{k+1}D_{k+1}^T$ ; 7: Compute  $\varepsilon_{k+1} = \frac{\|A^T X_{k+1} - X_{k+1}A + Q - X_{k+1}G X_{k+1}\|_2}{\|Q\|_2}$ ; 8: if  $\varepsilon_{k+1} < \varepsilon$  then 9: stop (interrupt); 10: end if 11: end for

Next, we consider the computational complexity of Algorithm 1, the operations involved include: computing  $A_{\alpha} = A - \alpha I$  with a complexity of n; calculating  $D_0 = A_{\alpha}^{-T}C^T$ ,  $P_0 = A_{\alpha}^{-1}B$  with a complexity of 4(p+m)n; determining the complexity of  $\Sigma_0$  and  $\Gamma_0$  as  $4(p^2 + m^2)n$ . During the iteration, the computation complexity for  $\widehat{A}_{1,k+1}$  and  $\widehat{A}_{2,k+1}$  are  $O((p_k^3 + m_k^3))$ , while for  $\Sigma_{k+1}$ ,  $\Gamma_{k+1}$ , it is  $4p_k m_k n$ . Subsequently,

computing  $D_{k+1}$ ,  $P_{k+1}$  involves a complexity of  $4(p_k^2 + m_k^2)n$ . Therefore, the overall computational complexity sums up to  $[4(p_k^2 + m_k^2 + p_km_k + p^2 + m^2 + p + m) + 1]n$ .

## **3** Convergence analysis

Since the R-ADDA iteration (15) is a low-rank version of the ADDA iteration (14), we only need to present the convergence theory of the ADDA iteration (14). **Theorem 1.** [20] Assuming that the matrix bundle  $\widehat{M} - \lambda \widehat{L}$  is the doubling transformation of the symplectic matrix bundle  $M - \lambda L$ , we have the following conclusions:

(a) The matrix bundle  $\widehat{M} - \lambda \widehat{L}$  is also a symplectic matrix bundle; (b) If  $M\begin{bmatrix}U\\V\end{bmatrix} = L\begin{bmatrix}U\\V\end{bmatrix}S$ , where  $U, V \in \mathbb{R}^{n \times m}$  and  $S \in \mathbb{R}^{n \times m}$ , then

$$\widehat{M} \begin{bmatrix} U \\ V \end{bmatrix} = \widehat{L} \begin{bmatrix} U \\ V \end{bmatrix} S^2;$$

(c) If  $M - \lambda L$  has a Kronecker product canonical form

$$WMZ = \begin{pmatrix} J_r & 0\\ 0 & I_{2n-r} \end{pmatrix}, \quad WLZ = \begin{pmatrix} I_r & 0\\ 0 & N_{2n-r} \end{pmatrix},$$

where W and Z are non-singular,  $J_r$  is a Jordan matrix, and  $N_{2n-r}$  is a nilpotent matrix, then there exists a non-singular matrix  $\widehat{W}$  such that

$$\widehat{W}\widehat{M}Z = \begin{pmatrix} J_r^2 & 0\\ 0 & I_{2n-r} \end{pmatrix}, \quad \widehat{W}\widehat{L}Z = \begin{pmatrix} I_r & 0\\ 0 & N_{2n-r}^2 \end{pmatrix}.$$

**Theorem 2.** Assuming  $X, Y \ge 0$  are symmetric positive semi-definite solutions of equations (1) and (2) respectively, the sequences  $\{\widehat{A}_k\}, \{X_k\}, \{Y_k\}$  generated by the ADDA iteration (14) satisfy: (a)  $\widehat{A}_k = (I_n + Y_k X)[\mathscr{C}(R;\alpha)]^{2^k}$ ; (b)  $0 \le X_k \le X_{k+1} \le X$  and

$$0 \le X - X_k = (I_n + X_k Y) [\mathscr{C}(S;\alpha)]^{2^k} X [\mathscr{C}(R;\alpha)]^{2^k} \le (I + XY) [\mathscr{C}(S;\alpha)]^{2^k} X [\mathscr{C}(R;\alpha)]^{2^k};$$
  
(c)  $0 \le Y_k \le Y_{k+1} \le Y$  and

$$0 \le Y - Y_k = (I_n + Y_k X) [\mathscr{C}(R;\alpha)]^{2^k} Y [\mathscr{C}(S;\alpha)]^{2^k} \le (I + YX) [\mathscr{C}(R;\alpha)]^{2^k} Y [\mathscr{C}(S;\alpha)]^{2^k}.$$

**Proof.** We will prove this by induction. Firstly, we observe that  $U, V \ge 0$  implies that I+UV is nonsingular, and  $V(I+UV)^{-1}, (I+UV)^{-1}U \ge 0$ . Given the definitions

$$X_0 = 2\alpha U_{\alpha}^{-1} Q A_{\alpha}^{-1}, \quad Y_0 = 2\alpha A_{\alpha}^{-1} G U_{\alpha}^{-1} \ge 0,$$

we have

$$X_1 = X_0 + \hat{A}^T 0 (In + X_0 Y_0)^{-1} X_k \hat{A}_0 \ge X_0,$$

 $Y_1 = Y_0 + \hat{A}0Y0(I_n + X_0Y_0)^{-1}\hat{A}^T 0 \ge Y0.$ 

Since  $M_1 - \lambda L_1$  is a doubling transformation of  $M_0 - \lambda L_0$ , by equation (14), we have:

$$\begin{pmatrix} \widehat{A}_1 & 0 \\ -X_1 & I \end{pmatrix} \begin{pmatrix} I \\ X \end{pmatrix} = \begin{pmatrix} I & Y_1 \\ 0 & \widehat{A}_1^T \end{pmatrix} \begin{pmatrix} I \\ X \end{pmatrix} [\mathscr{C}(R;\alpha)]^2,$$
$$\begin{pmatrix} \widehat{A}_1 & 0 \\ -X_1 & I \end{pmatrix} \begin{pmatrix} -Y \\ I \end{pmatrix} [\mathscr{C}(S;\alpha)]^2 = \begin{pmatrix} I & Y_1 \\ 0 & \widehat{A}_1^T \end{pmatrix} \begin{pmatrix} -Y \\ I \end{pmatrix},$$

then we can get

$$\begin{split} \widehat{A}_1 &= (I+Y_1X)[\mathscr{C}(R;\alpha)]^2, \quad X - X_1 = \widehat{A}_1^T X[\mathscr{C}(R;\alpha)]^2, \\ \widehat{A}_1^T &= (I+X_1Y)[\mathscr{C}(S;\alpha)]^2, \quad Y - Y_1 = \widehat{A}_1Y[\mathscr{C}(S;\alpha)]^2, \end{split}$$

which implies that  $X \ge X_1$  and  $Y \ge Y_1$ .

Similarly, we also have

$$X - X_1 = (I + X_1 Y) [\mathscr{C}(S;\alpha)]^2 X [\mathscr{C}(R;\alpha)]^2 \le (I + XY) [\mathscr{C}(S;\alpha)]^2 X [\mathscr{C}(R;\alpha)]^2,$$

$$Y - Y_1 = (I + Y_1 X) [\mathscr{C}(R;\alpha)]^2 Y [\mathscr{C}(S;\alpha)]^2 \le (I + Y X) [\mathscr{C}(R;\alpha)]^2 Y [\mathscr{C}(S;\alpha)]^2,$$
  
thus, we have proved the conclusion for  $k = 1$ .

Next, assuming the conclusion holds for all positive integers less than or equal to k, we now consider the case for k + 1. By the definition of  $\hat{A}_{k+1}$ ,  $X_{k+1}$ ,  $Y_{k+1}$  in the ADDA iteration as per equation (14), we can similarly deduce that

$$0 \le X_k \le X_{k+1}, \quad 0 \le Y_k \le Y_{k+1}.$$

Since  $M_{j+1} - \lambda L_{j+1}$  is a doubling transformation of  $M_j - \lambda L_j$ , (j = 1, 1, k), then by equation (14), we have

$$\hat{A}_{k+1} = (I + Y_{k+1}X)[\mathscr{C}(R;\alpha)]^{2^{k+1}}, \quad X - X_{k+1} = \hat{A}_{k+1}^T X[\mathscr{C}(R;\alpha)]^{2^{k+1}},$$
$$\hat{A}_{k+1}^T = (I + X_{k+1}Y)[\mathscr{C}(S;\alpha)]^{2^{k+1}}, \quad Y - Y_{k+1} = \hat{A}_{k+1}Y[\mathscr{C}(S;\alpha)]^{2^{k+1}}.$$

Similarly, we can obtain

$$0 \le X - X_{k+1} = (I + X_{k+1}Y)[\mathscr{C}(S;\alpha)]^{2^{k+1}}X[\mathscr{C}(R;\alpha)]^{2^{k+1}} \le (I + XY)[\mathscr{C}(S;\alpha)]^{2^{k+1}}X[\mathscr{C}(R;\alpha)]^{2^{k+1}},$$

$$0 \le Y - Y_{k+1} = (I + Y_{k+1}X)[\mathscr{C}(R;\alpha)]^{2^{k+1}}Y[\mathscr{C}(S;\alpha)]^{2^{k+1}} \le (I + YX)[\mathscr{C}(R;\alpha)]^{2^{k+1}}Y[\mathscr{C}(S;\alpha)]^{2^{k+1}},$$

which proves  $X \ge X_{k+1}$  and  $Y \ge Y_{k+1}$ . Therefore, the conclusion holds for the k + 1 case, and by the principle of mathematical induction, the theorem is proven.

$$W = \begin{bmatrix} L \begin{bmatrix} I \\ X \end{bmatrix}, \ M \begin{bmatrix} -Y \\ I \end{bmatrix} \end{bmatrix}, \quad Z = \begin{bmatrix} I & -Y \\ X & I \end{bmatrix},$$

where  $M_0 = M$ ,  $L_0 = L$ ,  $X, Y \ge 0$ . From equation (12), it follows that W and Z are nonsingular and satisfy

$$W^{-1}MZ = \begin{pmatrix} \mathscr{C}(R;\alpha) & 0\\ 0 & I \end{pmatrix}, \quad W^{-1}LZ = \begin{pmatrix} I & 0\\ 0 & \mathscr{C}(S;\alpha) \end{pmatrix}$$

Therefore, according to the spectral properties of symplectic matrix pencils,  $if\rho(\mathscr{C}(R;\alpha)) < 1$ , then  $\rho(\mathscr{C}(S;\alpha)) = \rho(\mathscr{C}(R;\alpha)) < 1$ . Additionally, if  $0 \le U \le V$ , then  $||U||_2 \le ||V||_2$ . Thus, from Theorem 2, we can obtain the convergence results of the ADDA algorithm as follows.

**Corollary 1.** Under the assumptions of Theorem 2, if  $\rho(\mathscr{C}(R; \alpha)) < 1$ , then we have the following conclusions:

 $\begin{aligned} &(a) \|\widehat{A}_{k}\|_{2} \leq (1+\|Y\|_{2}\|X\|_{2})\|[\mathscr{C}(R;\alpha)]^{2^{k}}\|_{2} \to 0 \quad (k\to\infty);\\ &(b) \|X-X_{k}\| \leq (\|X\|_{2}+\|X\|_{2}^{2}\|Y\|_{2})\|[\mathscr{C}(S;\alpha)]^{2^{k}}\|_{2}\|[\mathscr{C}(R;\alpha)]^{2^{k}}\|_{2} \to 0 \quad (k\to\infty);\\ &(c) \|Y-Y_{k}\| \leq (\|Y\|_{2}+\|Y\|_{2}^{2}\|X\|_{2})\|[\mathscr{C}(R;\alpha)]^{2^{k}}\|_{2}\|[\mathscr{C}(S;\alpha)]^{2^{k}}\|_{2} \to 0 \quad (k\to\infty). \end{aligned}$ 

## 4 Numerical experiments

**Example 1.** We take the coefficient matrix of the CARE (1) to be

$$A = \begin{pmatrix} -12 & -3 & 0 & \cdots & 0 \\ 2 & -12 & -3 & \cdots & 0 \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & \cdots & 2 & -12 & -3 \\ 0 & \cdots & 0 & 2 & -12 \end{pmatrix}_{n \times n}, \quad B = \begin{pmatrix} 0.02 \\ 0.02 \\ \vdots \\ 0.02 \\ 0.02 \end{pmatrix}_{n \times 1}$$

$$C = (0.01, 0.01, \cdots, 0.01, 0.01)_{1 \times n}.$$

Here, we choose the relative Residual (Res) as the iteration stopping criterion, where  $\_$ 

$$\operatorname{Res}(X_k) = \frac{\|A^T X_k + X_k A - X_k G X_k + Q\|_2}{\|Q\|_2}.$$

We use the low-rank Kleinman-Newton GADI (K-N-R-GADI) method and the R-ADDA method to solve this example, and the numerical results are shown in Table 1. From the table data, it is apparent that the R-ADDA iterative method is more effective in solving this example as the matrix dimension increases multiplicatively. Moreover, Figure 1 clearly illustrates the iteration steps and the variation of relative residuals for both methods when the matrix dimension is n = 1024, demonstrating that the convergence speed of the R-ADDA method is faster. Figure 2 depicts the time consumption of these two iterative methods as the matrix dimension increases, highlighting the efficiency of the R-ADDA method.

11

Let

n	algorithm	Res	IT	CPU
128	K-N-R-GADI	3.7511e-15	8	0.22s
128	R-ADDA	6.3853e-15	4	0.02s
256	K-N-R-GADI	4.3294e-14	7	0.61s
256	R-ADDA	6.6167e-15	4	0.07s
512	K-N-R-GADI	2.5746e-15	8	4.65s
512	R-ADDA	9.1141e-15	4	0.55s
1024	K-N-R-GADI	1.8153e-14	7	43.56s
1024	R-ADDA	2.9441e-14	4	5.95s
2048	K-N-R-GADI	1.6358e-14	7	567.73s
2048	R-ADDA	1.9252e-13	4	80.83s
4096	K-N-R-GADI	3.4587e-14	7	3246.6s
4096	R-ADDA	1.5886e-12	4	580.02s

**Table 1**: Numerical results for Example 1.



Fig. 1: The residual curve of Example Fig. 2: The time curve of Example 1. 1. n = 1024



$$A = \begin{pmatrix} -10 & -3 & -2 & 0 & 0 & \cdots & 0 \\ 2 & -10 & -3 & -2 & 0 & \cdots & 0 \\ 1 & 2 & -10 & -3 & -2 & \cdots & 0 \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \vdots \\ 0 & \cdots & 1 & 2 & -10 & -3 & -2 \\ 0 & \cdots & 0 & 1 & 2 & -10 & -3 \\ 0 & \cdots & 0 & 0 & 1 & 2 & -10 \end{pmatrix}_{n \times n}, \quad B = \begin{pmatrix} 0.005 \\ 0.005 \\ \vdots \\ 0.005 \\ 0.005 \end{pmatrix}_{n \times 1},$$

$$C = (0.001, 0.001, \cdots, 0.001, 0.001)_{1 \times n}$$

We use the same method as the previous example to solve this problem, and the numerical results are shown in Table 2. Similarly, we can conclude that compared with the K-N-R-GADI method, the R-ADDA method has a faster convergence speed and requires less iteration time, making it more effective.

n	algorithm	Res	IT	CPU
128	K-N-R-GADI	4.2138e-14	10	0.34s
128	R-ADDA	6.9657e-14	5	0.03s
256	K-N-R-GADI	2.252e-13	9	0.85s
256	R-ADDA	2.5169e-13	5	0.09s
512	K-N-R-GADI	1.1262e-13	9	5.31s
512	R-ADDA	9.5031e-13	5	0.62s
1024	K-N-R-GADI	$6.8534e{-}12$	7	37.96s
1024	R-ADDA	3.6833e-12	4	5.16s
2048	K-N-R-GADI	3.9627e-11	6	389.17s
2048	R-ADDA	1.4499e-11	4	61.53s
4096	K-N-R-GADI	1.9814e-11	6	5082.9s
4096	R-ADDA	5.7516e-11	4	635.19s

**Table 2**: Numerical results for Example 2.

## 5 Conclusions

In this paper, we propose a new low-rank alternating direction doubling algorithm utilizing the low-rank properties of matrices to compute low-rank approximate solutions for large-scale algebraic Riccati equations. By combining with the ADDA algorithm, we further introduce the low-rank ADDA algorithm. Moreover, we discover that theoretically, the low-rank ADDA algorithm and the ADDA algorithm exhibit the same convergence properties. Finally, we provide numerical examples to compare the effectiveness of the low-rank Kleinman-Newton GADI algorithm and the low-rank ADDA algorithm. The results demonstrate that the low-rank ADDA algorithm is more efficient. However, like other solvers, the performance of this algorithm heavily depends on the choice of parameters, which remains a challenging issue.

## References

- W.M.W., On a matrix Riccati equation of stochastic control, SIAM J. Control Optim. 6 (1968), 681–697.
- [2] A.J. Laub, A Schur method for solving algebraic Riccati equations, IEEE Trans. Automat. Control 24 (1979), 913–921.
- [3] Z. Bai, J. Demmel, On swapping diagonal blocks in real Schur form, Linear Algebra Appl. 186 (1993), 73–95.
- [4] J.H. Brandts, Matlab code for sorting real Schur forms, Numer. Linear Algebra Appl. 9 (2002), 249–261.
- [5] G.W. Stewart, HQR3 and EXCHNG: Fortran subroutines for calculating and ordering the eigenvalues of a real upper Hessenberg matrix, ACM Trans. Math. Software 2 (1976), 275–280.
- [6] G. Ammar, V. Mehrmann, On Hamiltonian and symplectic Hessenberg forms, Linear Algebra Appl. 149 (1991), 55–72.

- [7] R. Byers, A Hamiltonian QR-algorithm, SIAM J. Sci. Statist. Comput. 7 (1986), 212–229.
- [8] L. Dieci, Some numerical considerations and Newton's method revisited for solving algebraic Riccati equations, IEEE Trans. Automat. Control 36 (1991), 608–616.
- [9] S. Hammarling, Newton's method for solving the algebraic Riccati equation, NPL Rep. DITC 12/82, Nat. Phys. Lab., Teddington, Middlesex TW11 OLW, UK, 1982.
- [10] D. Kleinman, On an iterative technique for Riccati equation computations, IEEE Trans. Automat. Control 13 (1968), 114–115.
- [11] N. Sandell, On Newton's method for Riccati equation solution, IEEE Trans. Automat. Control 19 (1974), 254–255.
- [12] V. Mehrmann, E. Tan, Defect correction methods for the solution of algebraic Riccati equations, IEEE Trans. Automat. Control 33 (1988), 695–698.
- [13] J. Gardiner, A.J. Laub, A generalization of the matrix-sign-function solution to the algebraic Riccati equations, Internat. J. Control 44 (1986), 823–832.
- [14] R. Byers, Solving the algebraic Riccati equation with the matrix sign function, Linear Algebra Appl. 85 (1987), 267–279.
- [15] B.D.O. Anderson, Second-order convergent algorithms for the steady-state Riccati equation, Internat. J. Control 28 (1978) 295–306.
- [16] M. Kimura, Convergence of the doubling algorithm for the discrete-time algebraic Riccati equation, Internat. J. Systems Sci. 19 (1988) 701–711.
- [17] X.X. Guo, W.W. Lin, and S.F. Xu, A structure-preserving doubling algorithm for nonsymmetric algebraic Riccati equations, Numer. Math. 103 (2006), 393–412.
- [18] T.X. Li, Y.C. Kuo, and W.W.,Lin, Solving large-scale nonsymmetric algebraic Riccati equations by doubling, SIAM J. Matrix Ana. Appl. 34 (2013) 1129-1147.
- [19] W.G. Wang, R. C. Li, Alternating-directional doubling algorithm for M-matrix algebraic Riccati equation, SIAM J. Matrix Ana. Appl. 33 (2012), 170–194.
- [20] W.W. Lin, S. F. Xu, Convergence analysis of structure-preserving doubling algorithms for Riccati-type matrix equations, SIAM J. Matrix Anal. Appl. 28 (2006), 26–39.
- [21] J. Roberts, Linear model reduction and solution of the algebraic Riccati equation by the use of the sign function, Internat. J. Control 32 (1980), 667–687.
- [22] J.L. Howland, The sign matrix and the separation of matrix eigenvalues, Linear Algebra Appl. 49 (1983), 221–332.