Reduced density matrix formulation of quantum linear response

Theo Juncker von Buchwald,*^{,†} Karl Michael Ziems,^{*,†} Erik Rosendahl

Kjellgren,*^{,‡} Stephan P. A. Sauer,[¶] Jacob Kongsted,[‡] and Sonia Coriani[†]

Department of Chemistry, Technical University of Denmark, Kemitorvet Building 207, DK-2800 Kongens Lyngby, Denmark.

Department of Physics, Chemistry and Pharmacy, University of Southern Denmark, Campusvej 55, 5230 Odense, Denmark.

¶Department of Chemistry, University of Copenhagen, DK-2100 Copenhagen Ø.

E-mail: tjvbu@kemi.dtu.dk; kmizi@kemi.dtu.dk; kjellgren@sdu.dk

Abstract

The prediction of spectral properties via linear response (LR) theory is an important tool in quantum chemistry for understanding photo-induced processes in molecular systems. With the advances of quantum computing, we recently adapted this method for near-term quantum hardware using a truncated active space approximation with orbital rotation, named quantum linear response (qLR). In an effort to reduce the classic cost of this hybrid approach, we here derive and implement a reduced density matrix (RDM) driven approach of qLR. This allows for the calculation of spectral properties of moderately sized molecules with much larger basis sets than so far possible. We report qLR results for benzene and *R*-methyloxirane with a ccpVTZ basis set and study the effect of shot noise on the valence and oxygen K-edge absorption spectra of H_2O in the cc-pVTZ basis.

1 Introduction

The ability to predict molecular properties such as excitation energies, oscillator and rotational strengths, is important for the interpretation of spectroscopic data, and a cornerstone in many scientific contexts, e.g., photochemistry, photophysics, photocatalysis, and optogenetics. A well-established framework to calculate such properties (on conventional hardware) is molecular response theory $^{1-4}$. Within such theory, the response of the molecular system to (frequency-dependent) perturbing fields is rationalized in terms of linear and non-linear response functions $^{1-4}$. Through the calculation of the linear response function, for example, one can obtain the electric dipole polarizability and the optical rotation tensor. From the poles of the frequency-dependent response functions one can get the excited state energies, whereas the residues yield various transition properties (e.g. for multi-photon absorption processes) and properties of excited states. Response theory has been implemented for most electronic structure methods, including Hartree-Fock self-consistent field (SCF)^{1,5}, multi-configurational SCF (MCSCF)^{1,6,7}, coupled cluster (CC)^{2,8}, Møller-Plesset perturbation theory⁹ and time-dependent density functional theory^{10,11}.

With the ongoing promising developments in quantum computing and its expected impact on quantum chemistry, it comes as no surprise that linear response theory has recently been expanded to quantum hardware, and named quantum linear response (qLR)^{12,13}. In the approach by Kumar et al.¹², the authors perform wave function optimization and linear response in the complete orbital space to the level of singles and doubles. Therein, they introduce two full-space qLR approaches, namely self-consistent qLR and projected qLR. However, since current quantum hardware (referred to as noisy intermediate-scale quantum, NISQ) is extremely prone to errors due to noise and decoherence, chemical calculations are restricted to only a few qubits and necessitate hybrid approaches^{14,15}. The latter are defined as splitting the algorithm workload into quantum and classic parts, outsourcing less correlated calculations to classical machines.

The qLR approach by Ziems et al.¹³ builds upon this premise with a NISQ-era formulation of qLR. Here, the authors combine an active space approximation with truncated excitation ranks and orbital optimization. The hybrid approach comes into play by only mapping the active space

contributions onto the quantum computer. By doing so, they can dramatically reduce the number of parameters needed to describe the linear response properties and, importantly, can rely on less converged wave functions, which translates into shallower circuits and allows for the treatment of larger molecules and basis sets on near-term hardware. In their work, Ziems et al. introduce various parametrization schemes for the orbital rotation and active space excitation operators. This yields eight different qLR methods, three of which are classified as near-term. These are the naive, proj, and all-proj qLR ansätze with active spaces. Note that this qLR approach is similar to the complete active space self-consistent field (CASSCF) method^{16–20} commonly used on classical hardware with the difference of truncating the active space excitation rank and the use of various novel parameterization schemes. Closely related to the qLR approaches are also the qEOM approaches^{21,22}.

Our goal here is to reformulate the qLR approach with active spaces by Ziems et al.¹³ using reduced density matrices (RDMs). RDMs have been used in the past to efficiently formulate approaches on classical hardware resulting in reduced computational scaling^{23–26}. This is especially important for the classical part of the hybrid approach behind qLR. With increasing basis set size, the orbital rotation parameters can become the bottleneck over the quantum part. Using RDMs will allow performing qLR with even larger basis sets and reduced classical costs.

When truncating the linear response active space excitation rank above singles (qLRS), the naive qLRSD is the only ansatz that requires no more than 4-electron RDMs. The other near-term qLR approaches, proj qLRSD and all-proj qLRSD, both require 5- and 6-electron RDMs, which becomes quickly unfeasible due to the exponential memory scaling of the RDMs. On this basis, in this work, the naive qLRSD ansatz with RDMs is derived and subsequently compared against classic CASSCF LR for the absorption spectrum of benzene, using an active space of 6 electrons in 6 active orbitals [in short (6,6)], and the electronic circular dichroism (ECD) spectrum of *R*-methyloxirane, also with a (6,6) active space. Using the RDM ansatz of naive qLRSD allows us to simulate these systems up to and including the cc-pVTZ basis set. Additionally, a shot noise analysis investigation of the RDM qLR formulation is performed on H₂O (4,4)/cc-pVTZ.

This paper is organized as follows. In Section 2, we shortly review unitary coupled cluster theory as well as the active space approximation and linear response. We then introduce reduced density matrices and briefly discuss the scaling of our RDM-driven naive qLR approach. In Section 3, we provide computational details, followed by Section 4 where we present and discuss the results for the absorption spectrum of benzene (Section 4.1) and the ECD spectrum of *R*-methyloxirane (Section 4.2) using the RDM formalism of linear response and compare these against CASSCF results. In Section 4.3, we investigate the effects of shot noise when using the RDM-based naive qLR approach. Finally, some concluding remarks are given in Section 5.

2 Theory

In this section we formulate and provide working equations for a reduced density matrix-based approach to linear response theory with a truncated active space. We start by introducing the orbital-optimized unitary coupled cluster (oo-UCC) method in Sec. 2.1, followed by an introduction to the active space decomposition in Sec. 2.2, and the linear response theory in an active space framework in Sec. 2.3. After this, a brief overview of reduced density matrices (RDMs) is given in Sec. 2.4 and a short discussion of scaling in Sec. 2.5. It is important to note that the linear response equations derived here may be utilized with any variational reference state and not just oo-UCC. Throughout the manuscript, unless otherwise stated, this index notation will be used: p, q, r, s, t, u, m, and n are general indices; a, b, c, and d are virtual indices; i, j, k, and l are inactive indices; and v_a and v_i are active space orbitals that are, respectively, virtual and inactive in the Hartree-Fock reference state.

2.1 Unitary coupled cluster

In unitary coupled cluster (UCC) theory²⁷, the UCC wave function is given by an exponential ansatz of a reference state

$$|\text{UCC}\rangle = e^{\hat{T}(\theta) - \hat{T}^{\dagger}(\theta)} |0\rangle \tag{1}$$

where $\hat{T}(\theta)$ is the cluster operator and $\hat{T}^{\dagger}(\theta)$ is its Hermitian conjugate, both depending on the quantum circuit parameters θ . The cluster operator can be expanded by excitation ranks

$$\hat{T}(\boldsymbol{\theta}) = \hat{T}_1(\boldsymbol{\theta}) + \hat{T}_2(\boldsymbol{\theta}) + \dots$$
(2)

where

$$\hat{T}_{1}(\theta) = \sum_{ai} \theta_{ai} \hat{E}_{ai}$$

$$\hat{T}_{2}(\theta) = \frac{1}{2} \sum_{aibj} \theta_{aibj} \hat{E}_{ai} \hat{E}_{bj}$$

$$\vdots$$

$$(3)$$

and \hat{E}_{ai} is the singlet one-electron excitation operator. The UCC energy expression is

$$E(\theta) = \langle \text{UCC}(\theta) | \hat{H} | \text{UCC}(\theta) \rangle \tag{4}$$

with the Hamiltonian defined as

$$\hat{H} = \sum_{pq} h_{pq} \hat{E}_{pq} + \frac{1}{2} \sum_{pqrs} g_{pqrs} \hat{e}_{pqrs}$$
(5)

where \hat{e}_{pqrs} is the two-electron excitation operator, defined as

$$\hat{e}_{pqrs} = \hat{E}_{pq}\hat{E}_{rs} - \delta_{rq}\hat{E}_{ps} ; \qquad (6)$$

 h_{pq} and g_{pqrs} are the one- and two-electron integrals in the molecular orbital (MO) basis. The θ circuit parameters are found by minimizing the energy with respect to the circuit parameter values. A continuation of the UCC formalism is the orbital-optimized UCC (oo-UCC)^{28,29}, where the UCC wavefunction is exponentially parameterized by the orbital rotation operator

$$\hat{\boldsymbol{\kappa}}(\boldsymbol{\kappa}) = \sum_{p>q} \boldsymbol{\kappa}_{pq} \hat{E}_{pq}^{-} , \qquad (7)$$

with

$$\hat{E}_{pq}^{-} = \hat{E}_{pq} - \hat{E}_{qp} , \qquad (8)$$

as

$$|\text{oo-UCC}(\kappa, \theta)\rangle = e^{-\hat{\kappa}(\kappa)} |\text{UCC}(\theta)\rangle$$
 (9)

When combined with UCC in an active space, the only non-redundant rotations are *inactive* to active (AI), inactive to virtual (VI), and active to virtual (VA), $pq \in \{AI, VI, VA\}$. Here, I, A, and V refer to the set of orbitals contained in, respectively, the inactive, active, and virtual space. Instead of including the orbital rotation in the wave function, it is convenient to transform the Hamiltonian (integrals) using the orbital rotation parameters

$$\hat{H}(\kappa) = \sum_{pq} h_{pq}(\kappa) \hat{E}_{pq} + \frac{1}{2} \sum_{pqrs} g_{pqrs}(\kappa) \hat{e}_{pqrs}$$
(10)

using

$$h_{pq}(\kappa) = \sum_{p'q'} \left[e^{\kappa} \right]_{q'q} h_{p'q'} \left[e^{-\kappa} \right]_{p'p}$$
(11)

$$g_{pqrs}(\kappa) = \sum_{p'q'r's'} [e^{\kappa}]_{s's} [e^{\kappa}]_{q'q} g_{p'q'r's'} [e^{-\kappa}]_{p'p} [e^{-\kappa}]_{r'r}.$$
 (12)

The oo-UCC energy expression is then

$$E(\theta) = \langle \text{UCC}(\theta) | \hat{H}(\kappa) | \text{UCC}(\theta) \rangle.$$
(13)

Performing a minimization of the energy with respect to the circuit parameters, which are optimized on the quantum hardware, and the orbital rotation parameters, which are split into classic and quantum workload, is known as the orbital optimized variational quantum eigensolver (oo-VOE)^{28,29}.

2.2 Active space

A well-known active space method in 'classic' quantum chemistry is CASSCF^{18–20}. In CASSCF, the full space is split into subspaces consisting of an inactive space where all orbitals are doubly occupied, an active space that is treated using a complete configuration interaction (CI) expansion, and a virtual space where all orbitals are empty. In the quantum approach we use here, the CI expansion in the active space is truncated to singles and doubles excitations¹³. Applying an active space approximation to the wave function yields

$$|0(\theta)\rangle = |I\rangle \otimes |A(\theta)\rangle \otimes |V\rangle , \qquad (14)$$

where $|I\rangle$ is the inactive part of the wave function, $|A(\theta)\rangle$ is the active part, and $|V\rangle$ is the virtual part. The active space is parameterized by θ by using a unitary transformation

$$|A(\theta)\rangle = U(\theta)|A\rangle \tag{15}$$

Operators can likewise be decomposed as

$$\hat{O} = \hat{O}_I \otimes \hat{O}_A \otimes \hat{O}_V. \tag{16}$$

By applying the active space approximation, expectation values of any operator now have the form

$$\langle 0(\theta) | \hat{O} | 0(\theta) \rangle = \langle I | \hat{O}_I | I \rangle \langle A(\theta) | \hat{O}_A | A(\theta) \rangle \langle V | \hat{O}_V | V \rangle.$$
(17)

Using the active space approximation on the oo-UCC energy expression [Eq. (13)], only the active space part is calculated on the quantum hardware.

2.3 Linear response

Here, we adopt the qLR approach with active spaces introduced by Ziems et al.¹³ that uses, as a starting point, the MCSCF wave function,

$$|\tilde{0}(t)\rangle = e^{\hat{\kappa}(t)} e^{S(t)} |0\rangle$$
(18)

$$\hat{\kappa}(t) = \sum_{\mu} \left(\kappa_{\mu}(t)\hat{q}_{\mu} + \kappa_{\mu}^{*}(t)\hat{q}_{\mu}^{\dagger} \right)$$
(19)

$$\hat{S}(t) = \sum_{n} \left(S_n(t)\hat{G}_n + S_n^*(t)\hat{G}_n^\dagger \right)$$
(20)

where the wave function $|0\rangle$ is parameterized through the time-dependent orbital operator $\hat{\kappa}(t)$ and active space rotation operator $\hat{S}(t)$, which contain, respectively, the orbital rotation operators \hat{q}_{μ} and the active space excitation operators \hat{G}_n . Ziems et al. then introduce various ansätze for linear response on near-term quantum computers using a variety of choices for \hat{q}_{μ} and \hat{G}_n^{13} . Here, we specifically focus on the naive operators

$$\hat{q}_{pq} = \frac{1}{\sqrt{2}} \hat{E}_{pq} \tag{21}$$

$$\hat{G} \in \left\{ \frac{1}{\sqrt{2}} \hat{E}_{\nu_{a}\nu_{i}}, \frac{1}{2\sqrt{(1+\delta_{\nu_{a}\nu_{b}})(1+\delta_{\nu_{i}\nu_{j}})}} \left(\hat{E}_{\nu_{a}\nu_{i}} \hat{E}_{\nu_{b}\nu_{j}} + \hat{E}_{\nu_{a}\nu_{j}} \hat{E}_{\nu_{b}\nu_{i}} \right), \qquad (22)$$

$$\frac{1}{2\sqrt{3}} \left(\hat{E}_{\nu_{a}\nu_{i}} \hat{E}_{\nu_{b}\nu_{j}} - \hat{E}_{\nu_{a}\nu_{j}} \hat{E}_{\nu_{b}\nu_{i}} \right) \right\}.$$

We will refer to the method as naive qLRSD, since only singles and doubles excitations are included in the linear response excitation space.

The key quantity in linear response theory is the frequency-dependent linear response function $\langle \langle \hat{A}; \hat{B} \rangle \rangle_{\omega_B}$, for two generic operators \hat{A} and \hat{B} . The linear response function can be computed as³

$$\langle\langle \hat{A}; \hat{B} \rangle\rangle_{\omega_B} = -\mathbf{V}_A^{[1]\dagger} \boldsymbol{\beta}_B.$$
⁽²³⁾

This requires solving sets of linear response equations ^{1,3,30}

$$\left(\mathbf{E}^{[2]} - \boldsymbol{\omega}_{B} \mathbf{S}^{[2]}\right) \boldsymbol{\beta}_{B} = \mathbf{V}_{B}^{[1]}$$
(24)

where $\mathbf{E}^{[2]}$ is the electronic Hessian, $\mathbf{S}^{[2]}$ is the Hermitian metric, β_B is the linear response vector, and $\mathbf{V}_B^{[1]}$ is the property gradient. The Hessian and Hermitian metric can be expressed as

$$\mathbf{E}^{[2]} = \begin{pmatrix} A & B \\ B^* & A^* \end{pmatrix}, \quad \mathbf{S}^{[2]} = \begin{pmatrix} \Sigma & \Delta \\ -\Delta^* & -\Sigma^* \end{pmatrix}$$
(25)

with the submatrices

$$A = \begin{pmatrix} \left\langle 0 \left| \left[\hat{q}_{\mu}^{\dagger}, \left[\hat{H}, \hat{q}_{\nu} \right] \right] \right| 0 \right\rangle & \left\langle 0 \left| \left[\hat{G}_{n}, \left[\hat{H}, \hat{q}_{\nu}^{\dagger} \right] \right] \right| 0 \right\rangle \\ \left\langle 0 \left| \left[\hat{G}_{n}^{\dagger}, \left[\hat{H}, \hat{q}_{\nu} \right] \right] \right| 0 \right\rangle & \left\langle 0 \left| \left[\hat{G}_{n}^{\dagger}, \left[\hat{H}, \hat{G}_{m} \right] \right] \right| 0 \right\rangle \end{pmatrix} \end{cases}$$
(26)

$$B = \begin{pmatrix} \left\langle 0 \left| \left[\hat{q}_{\mu}^{\dagger}, \left[\hat{H}, \hat{q}_{\nu}^{\dagger} \right] \right] \right| 0 \right\rangle & \left\langle 0 \left| \left[\hat{G}_{n}^{\dagger}, \left[\hat{H}, \hat{q}_{\nu}^{\dagger} \right] \right] \right| 0 \right\rangle \\ \left\langle 0 \left| \left[\hat{G}_{n}^{\dagger}, \left[\hat{H}, \hat{q}_{\nu}^{\dagger} \right] \right] \right| 0 \right\rangle & \left\langle 0 \left| \left[\hat{G}_{n}^{\dagger}, \left[\hat{H}, \hat{G}_{m}^{\dagger} \right] \right] \right| 0 \right\rangle \end{pmatrix}$$

$$(27)$$

$$\Sigma = \begin{pmatrix} \left\langle 0 \left| \left[\hat{q}_{\mu}^{\dagger}, \hat{q}_{\nu} \right] \right| 0 \right\rangle & \left\langle 0 \left| \left[\hat{q}_{\mu}^{\dagger}, \hat{G}_{m} \right] \right| 0 \right\rangle \\ \left\langle 0 \left| \left[\hat{G}_{n}^{\dagger}, \hat{q}_{\nu} \right] \right| 0 \right\rangle & \left\langle 0 \left| \left[\hat{G}_{n}^{\dagger}, \hat{G}_{m} \right] \right| 0 \right\rangle \end{pmatrix} \end{cases}$$
(28)

$$\Delta = \begin{pmatrix} \left\langle 0 \left| \left[\hat{q}_{\mu}^{\dagger}, \hat{q}_{\nu}^{\dagger} \right] \right| 0 \right\rangle & \left\langle 0 \left| \left[\hat{q}_{\mu}^{\dagger}, \hat{G}_{m}^{\dagger} \right] \right| 0 \right\rangle \\ \left\langle 0 \left| \left[\hat{G}_{n}^{\dagger}, \hat{q}_{\nu}^{\dagger} \right] \right| 0 \right\rangle & \left\langle 0 \left| \left[\hat{G}_{n}^{\dagger}, \hat{G}_{m}^{\dagger} \right] \right| 0 \right\rangle \end{pmatrix}.$$

$$(29)$$

The property gradient has the form

$$\mathbf{V}_{B}^{[1]} = \begin{pmatrix} \langle 0 | [\hat{q}_{\mu}, \hat{B}] | 0 \rangle \\ \langle 0 | [\hat{G}_{n}, \hat{B}] | 0 \rangle \\ \langle 0 | [\hat{q}_{\mu}^{\dagger}, \hat{B}] | 0 \rangle \\ \langle 0 | [\hat{G}_{n}^{\dagger}, \hat{B}] | 0 \rangle \end{pmatrix}.$$
(30)

The linear response vector is given as

$$\beta_B = \begin{pmatrix} Z_B \\ Y_B^* \end{pmatrix} = \begin{pmatrix} \kappa_B \\ S_B \\ \kappa_{-B}^* \\ S_{-B}^* \end{pmatrix}$$
(31)

where κ_B and S_B are the first order expansions of the orbital parameters in Eqs. (19) and (20). The spectral representation of the linear response function can be written as

$$\langle\langle \hat{A}; \hat{B} \rangle\rangle_{\omega_{B}} = \sum_{k>0} \frac{\langle 0|[\hat{A}, \hat{\tilde{O}}_{k}]|0\rangle \langle 0|[\hat{\tilde{O}}_{k}^{\dagger}, \hat{B}]|0\rangle}{\omega - \omega_{k}} - \sum_{k>0} \frac{\langle 0|[\hat{B}, \hat{\tilde{O}}_{k}]|0\rangle \langle 0|[\hat{\tilde{O}}_{k}^{\dagger}, \hat{A}]|0\rangle}{\omega + \omega_{k}} , \qquad (32)$$

where $\hat{\tilde{O}}_k$ is a normalized (excitation) operator

$$\hat{\hat{O}}_{k} = \frac{\hat{O}_{k}}{\sqrt{\langle k|k\rangle}} = \frac{\hat{O}_{k}}{\sqrt{\langle 0|[\hat{O}_{k},\hat{O}_{k}^{\dagger}]|0\rangle}}$$
(33)

$$\hat{O}_k = \sum_{l \in \mu, n} \left(Z_{k,l} \hat{X}_l^{\dagger} + Y_{k,l} \hat{X}_l \right).$$
(34)

In the equation above, \hat{X}_l is the set of orbital rotation and active space excitation operators $\{\hat{q}_{\mu}, \hat{G}_n\}$, and $Z_{k,l}$ and $Y_{k,l}$ are their weights. The (one-electron) operators \hat{A} and \hat{B} are in their second quantization form, e.g.

$$\hat{A} = \sum_{pq} \mathcal{A}_{pq} \hat{E}_{pq} , \qquad (35)$$

where A_{pq} are the \hat{A} operator's integrals over the MO basis.

From the poles and residues of the linear response function, transition properties like excitation energies and transition strengths are obtained. The excitation energies and excitation vectors are found by solving the eigenvalue equation

$$\mathbf{E}^{[2]}\boldsymbol{\beta}_k = \boldsymbol{\omega}_k \mathbf{S}^{[2]}\boldsymbol{\beta}_k. \tag{36}$$

which is the homogeneous version of the linear response equations in Eq. (23), and where β_k is the excitation vector containing the $Z_{k,l}$ and $Y_{k,l}$ parameters in Eq. (34). The transition strengths are the residues of the linear response function and are found according to

$$\lim_{\omega \to \omega_k} (\omega - \omega_k) \langle \langle \hat{A}; \hat{B} \rangle \rangle_{\omega} = \langle 0 | [\hat{A}, \hat{\tilde{O}}_k] | 0 \rangle \langle 0 | [\hat{\tilde{O}}_k^{\dagger}, \hat{B}] | 0 \rangle = \left(\mathbf{V}_A^{[1]\dagger} \beta_k \right) \left(\beta_k^{\dagger} \mathbf{V}_B^{[1]} \right)$$
(37)

which, as seen, requires the property gradient and the excitation vector.

A prototypical example of a linear response function is the dipole polarizability

$$\alpha_{\gamma\delta}(\omega) = -\langle \langle \hat{\mu}_{\gamma}; \hat{\mu}_{\delta} \rangle \rangle_{\omega} \tag{38}$$

where $\hat{\mu}_{\gamma}$ is a Cartesian component of the one-electron electric dipole moment operator ($\gamma \in \{x, y, z\}$), in second quantization given by

$$\hat{\mu}_{\gamma} = -\sum_{pq} (\vec{r}_{\gamma})_{pq} \hat{E}_{pq}$$
(39)

$$(\vec{r}_{\gamma})_{pq} = \left\langle p \left| \vec{r}_{\gamma} \right| q \right\rangle.$$
(40)

In this paper, we focus on the oscillator strengths f_k of one-photon absorption, which are related

to the residues of the electric dipole polarizability,

$$f_k = \frac{2}{3}\omega_k \sum_{\gamma} \left| \langle 0|[\hat{\mu}_{\gamma}, \hat{\tilde{O}}_k]|0\rangle \right|^2 , \qquad (41)$$

and on the rotational strengths R_k of ECD, connected to the residues of the optical rotation tensor,

$$R_{k} = \operatorname{Im}(\langle 0|[\hat{\mu}, \hat{\tilde{O}}_{k}]|0\rangle \cdot \langle 0|[\hat{\tilde{O}}_{k}^{\dagger}, \hat{m}]|0\rangle)$$

$$(42)$$

where \hat{m} is the magnetic dipole moment operator

$$\hat{\boldsymbol{m}} = -\frac{i}{2} \sum_{pq} (\vec{r} \times \vec{\nabla})_{pq} \hat{E}_{pq}$$
(43)

$$(\vec{r} \times \vec{\nabla})_{pq} = \left\langle p \left| \vec{r} \times \vec{\nabla} \right| q \right\rangle.$$
(44)

2.4 Reduced density matrices

As anticipated, this work formulates naive qLRSD in terms of reduced density matrices. This entails calculating the Hessian, the metric and the property gradients of the linear response equation [Eq. (24)] using RDMs. To this end, the largest terms we can obtain from Eq. (24) within a singles and doubles parameterization are

$$\left\langle 0 \left| \left[\hat{E}_{\boldsymbol{v}_k \boldsymbol{v}_c} \hat{E}_{\boldsymbol{v}_l \boldsymbol{v}_d}, \left[\sum_{pqrs} g_{pqrs} \hat{e}_{pqrs}, \hat{E}_{\boldsymbol{v}_a \boldsymbol{v}_i} \hat{E}_{\boldsymbol{v}_b \boldsymbol{v}_j} \right] \right] \right| 0 \right\rangle$$
(45)

$$\left\langle 0 \left| \left[\hat{E}_{\boldsymbol{\nu}_{k}\boldsymbol{\nu}_{c}} \hat{E}_{\boldsymbol{\nu}_{l}\boldsymbol{\nu}_{d}}, \left[\sum_{pqrs} g_{pqrs} \hat{e}_{pqrs}, \hat{E}_{\boldsymbol{\nu}_{i}\boldsymbol{\nu}_{a}} \hat{E}_{\boldsymbol{\nu}_{j}\boldsymbol{\nu}_{b}} \right] \right] \right| 0 \right\rangle.$$
(46)

They are both matrix elements of the Hessian, obtained by inserting Eq. (22) in their respective matrix elements. The first, Eq. (45), is obtained by inserting Eq. (22) in $\langle 0 | [\hat{G}_n^{\dagger} [\hat{H}, \hat{G}_m]] | 0 \rangle$ of Eq. (26) and the second, Eq. (46), is obtained by inserting Eq. (22) in $\langle 0 | [\hat{G}_n^{\dagger} [\hat{H}, \hat{G}_m^{\dagger}]] | 0 \rangle$ of Eq. (27). Due to rank reduction³¹, these matrix elements will contain at most the 4-electron reduced density matrix (4-RDM).

As the terms in Eqs. (45) and (46) are the largest elements in the naive qLRSD approach, we need to consider all RDMs up to four particles (4-RDM) to be able to formulate naive qLRSD in terms of reduced density matrices. These are given in second quantization as

$$\Gamma_{pq}^{[1]} = \sum_{\tau \in \{\alpha,\beta\}} \left\langle 0 \left| \hat{a}_{p\tau}^{\dagger} \hat{a}_{q\tau} \right| 0 \right\rangle = \left\langle 0 \left| \hat{E}_{pq} \right| 0 \right\rangle \tag{47}$$

$$\Gamma_{pqrs}^{[2]} = \sum_{\tau\gamma \in \{\alpha,\beta\}} \left\langle 0 \left| \hat{a}_{p\tau}^{\dagger} \hat{a}_{r\gamma}^{\dagger} \hat{a}_{s\gamma} \hat{a}_{q\tau} \right| 0 \right\rangle = \left\langle 0 \left| \hat{e}_{pqrs} \right| 0 \right\rangle$$
(48)

$$\Gamma_{pqrstu}^{[3]} = \sum_{\tau\gamma\delta\in\{\alpha,\beta\}} \left\langle 0 \left| \hat{a}_{p\tau}^{\dagger} \hat{a}_{r\gamma}^{\dagger} \hat{a}_{t\delta}^{\dagger} \hat{a}_{u\delta} \hat{a}_{s\gamma} \hat{a}_{q\tau} \right| 0 \right\rangle$$

$$\tag{49}$$

$$= \left\langle 0 \left| \hat{E}_{pq} \hat{E}_{rs} \hat{E}_{tu} \right| 0 \right\rangle - \delta_{ts} \Gamma_{pqru}^{[2]} - \delta_{rq} \Gamma_{pstu}^{[2]} - \delta_{tq} \Gamma_{purs}^{[2]} - \delta_{ts} \delta_{rq} \Gamma_{pu}^{[1]} \right.$$

$$\Gamma_{pqrstumn}^{[4]} = \sum_{\tau\gamma\delta\sigma\in\{\alpha,\beta\}} \left\langle 0 \left| \hat{a}_{p\tau}^{\dagger} \hat{a}_{r\gamma}^{\dagger} \hat{a}_{t\delta}^{\dagger} \hat{a}_{m\sigma}^{\dagger} \hat{a}_{n\sigma} \hat{a}_{u\delta} \hat{a}_{s\gamma} \hat{a}_{q\tau} \right| 0 \right\rangle$$

$$= \left\langle 0 \left| \hat{E}_{pq} \hat{E}_{rs} \hat{E}_{tu} \hat{E}_{mn} \right| 0 \right\rangle - \delta_{rq} \Gamma_{pstumn}^{[3]} - \delta_{tq} \Gamma_{pursmn}^{[3]} - \delta_{mq} \Gamma_{pnrstu}^{[3]} - \delta_{mu} \Gamma_{pqrstn}^{[3]} - \delta_{ts} \Gamma_{pqrmn}^{[3]} - \delta_{ms} \Gamma_{pqrmtu}^{[2]} - \delta_{mu} \delta_{rq} \Gamma_{pstn}^{[2]} - \delta_{mu} \delta_{tq} \Gamma_{pqrs}^{[2]} - \delta_{ts} \delta_{mu} \Gamma_{pqrn}^{[2]} - \delta_{ts} \delta_{mq} \Gamma_{pnru}^{[2]} - \delta_{ms} \delta_{rq} \Gamma_{pnru}^{[2]} - \delta_{ms} \delta_{tq} \Gamma_{purn}^{[2]} - \delta_{mu} \delta_{ts} \delta_{rq} \Gamma_{pn}^{[1]} \right.$$

where a^{\dagger} and a are the creation and annihilation operators.

Using the separation of the wave function into inactive, active, and virtual components in Eq. (17), the RDMs may be expressed by lower order RDMs when not all indices are in the active space and equals zero when any virtual indices are present. Crucially, inactive indices trivially reduce to Kronecker deltas and thus we only need to calculate the RDMs with active space indices on quantum hardware. Moreover, since the RDMs fulfill the symmetries shown in Eqs. (S1) to (S4) in the SI (Section S1), it is only necessary to consider the non-redundant combinations of inactive and active indices as shown in Eqs. (51) to (54) reducing the number of elements that need to be measured on the quantum hardware by a factor equal to the number of symmetries. For

the 4-RDM there are 48 symmetries, and so only $\frac{1}{48}$ of the elements need to be measured.

$$\Gamma_{pq}^{[1]} = \begin{cases} 2\delta_{pq} & pq \in II \\ \langle 0 | \hat{E}_{pq} | 0 \rangle & pq \in AA \\ 0 & \text{otherwise} \end{cases}$$
(51)

$$\Gamma_{pqrs}^{[2]} = \begin{cases} 4\delta_{pq}\delta_{rs} - 2\delta_{rq}\delta_{ps} & pqrs \in IIII \\ 2\delta_{rs}\Gamma_{pq}^{[1]} & pqrs \in AAII \\ -\delta_{ps}\Gamma_{rq}^{[1]} & pqrs \in IAAI \\ \langle 0 | \hat{e}_{pqrs} | 0 \rangle & pqrs \in AAAA \\ 0 & \text{otherwise} \end{cases}$$
(52)

$$\Gamma_{pqrstu}^{[3]} = \begin{cases} 2\delta_{pq}\Gamma_{rstu}^{[2]} - \delta_{rq}\Gamma_{pstu}^{[2]} - \delta_{tq}\Gamma_{purs}^{[2]} & pqrstu \in IIIIII \\ 4\delta_{rs}\delta_{tu}\Gamma_{pq}^{[1]} - 2\delta_{ts}\delta_{ru}\Gamma_{pq}^{[1]} & pqrstu \in AAIIII \\ \delta_{pu}\delta_{ts}\Gamma_{rq}^{[1]} - 2\delta_{ps}\delta_{tu}\Gamma_{rq}^{[1]} & pqrstu \in IAAIII \\ 2\delta_{tu}\Gamma_{pqrs}^{[2]} & pqrstu \in AAAAII \\ -\delta_{ru}\Gamma_{pqts}^{[2]} & pqrstu \in AAIAAI \\ \langle 0 | \hat{E}_{qsu}^{prt} | 0 \rangle & pqrstu \in AAAAAA \\ 0 & \text{otherwise} \end{cases}$$
(53)

$$\Gamma_{pqrstumn}^{[4]} = \begin{cases} 2\delta_{pq}\Gamma_{rstumn}^{[3]} - \delta_{rq}\Gamma_{pstumn}^{[3]} - \delta_{tq}\Gamma_{pursmn}^{[3]} - \delta_{mq}\Gamma_{pnrstu}^{[3]} & pqrstumn \in IIIIIIII \\ \Gamma_{pq}^{[1]}\Gamma_{rstumn}^{[3]} & pqrstumn \in AAIIIIII \\ -\frac{1}{2}\Gamma_{rq}^{[1]}\Gamma_{pstumn}^{[3]} & pqrstumn \in IAAIIIII \\ (4\delta_{tu}\delta_{mn} - 2\delta_{mu}\delta_{tn})\Gamma_{pqrs}^{[2]} & pqrstumn \in AAAAIIII \\ (\delta_{rn}\delta_{mu} - 2\delta_{ru}\delta_{mn})\Gamma_{pqts}^{[2]} & pqrstumn \in AAIAAIII \\ \delta_{ps}\delta_{tn}\Gamma_{rqmu}^{[2]} + \delta_{pn}\delta_{ts}\Gamma_{rumq}^{[2]} & pqrstumn \in IAAIIAAI \\ 2\delta_{mn}\Gamma_{pqrstu}^{[3]} & pqrstumn \in AAAAAAII \\ -\delta_{tn}\Gamma_{pqrsmu}^{[3]} & pqrstumn \in AAAAAIAI \\ (0 | \hat{E}_{qsun}^{prtm} | 0 \rangle & pqrstumn \in AAAAAAAA \\ 0 & otherwise \end{cases}$$

$$(54)$$

We have also only included one permutation of the unique combinations of inactive-active pairs. As an example, let us take the term $\Gamma_{pqrs}^{[2]} = -\delta_{ps}\Gamma_{rq}^{[1]}$ with $pqrs \in IAAI$. For this combination of inactive and active orbitals, there are two possible permutations

$$\Gamma_{pqrs}^{[2]} = -\delta_{ps}\Gamma_{rq}^{[1]} \quad pqrs \in IAAI \tag{55}$$

$$\Gamma_{pqrs}^{[2]} = -\delta_{rq}\Gamma_{ps}^{[1]} \quad pqrs \in AIIA , \qquad (56)$$

where the permutations on the right hand side of the equations have been moved according to the permutation in the inactive and active indices ($pq \leftrightarrow rs$). Working equations for all matrix elements needed to solve the linear response equation can be found in Section S2 in the SI.

2.5 Scaling

In terms of the number of elements, the three terms with the largest scaling are

$$\left\langle 0 \left| \left[\hat{G}, \left[\hat{H}, \hat{G} \right] \right] \right| 0 \right\rangle \to N_{A_{\text{occ}}}^4 N_{A_{\text{unocc}}}^4 N_{I+A}^2 \tag{57}$$

$$\langle 0 \left| \left[\hat{q}, \left[\hat{H}, \hat{G} \right] \right] \right| 0 \rangle \to N_{A_{\text{occ}}}^2 N_{A_{\text{unocc}}}^2 \max \{ N_{I+A}^2 N_I N_A, N_{\text{orb}} N_{I+A} N_I N_V, N_{\text{orb}} N_{I+A} N_A N_V \}$$
(58)

$$\left\langle 0 \left| \left[\hat{q}, \left[\hat{H}, \hat{q} \right] \right] \right| 0 \right\rangle \to \max\left\{ N_{I+A}^2 N_I^2 N_A^2, N_{\text{orb}}^2 N_I^2 N_V^2, N_{\text{orb}}^2 N_A^2 N_V^2 \right\}.$$

$$\tag{59}$$

In the equations above, various indices have been used to refer to the number of orbitals in different spaces. These are the number of orbitals in the inactive space (N_I) , active space (N_A) , virtual space (N_V) , active space that is inactive in the Hartree-Fock reference state $(N_{A_{occ}})$, active space that is virtual in the Hartree-Fock reference state $(N_{A_{unocc}})$, inactive and active space (N_{I+A}) , and in total (N_{orb}) . For a minimal basis set Eq. (57) will be the dominating term. However, with an increasing basis set size the virtual indices will dominate the scaling eventually causing Eq. (59) to become the dominant term. Regarding memory usage, the two-electron integrals and the 4-RDM are of interest as they scale as N_{orb}^4 and N_A^8 , respectively. Since we construct the entire Hessian, it is also important to consider its size for larger systems, as it scales

$$\mathbf{E}^{[2]} \to (2(N_{\hat{q}} + N_{\hat{G}}))^2 = 4(N_{\hat{q}} + N_{\hat{G}})^2 \tag{60}$$

where $N_{\hat{q}}$ and $N_{\hat{G}}$ are the number of orbital rotation and active space excitation operators. $N_{\hat{G}}$ scale with the active space size whereas $N_{\hat{q}}$ scale with the size of the inactive, active, and virtual spaces. To reduce the impact of the scaling of the Hessian it is possible to use a subspace approach instead of constructing the full Hessian explicitly³². This is a topic for future research.

3 Computational Details

As the oo-UCC wave function is decomposed into an inactive, active, and virtual space we here adopt the CAS-like notation oo-UCC(n, σ), where n is the number of electrons in the active space and σ is the number of orbitals in the active space. The excitation ranks included in the wave function's active space is truncated at the singles and doubles level. Meaning, oo-UCCSD(6,6) uses a (6,6) active space with cluster operators to the level of singles and doubles excitations in the active space.

The oo-UCC calculations were performed using our in-house quantum chemistry software SlowQuant³³ which is interfaced with PySCF^{34,35} for one and two-electron integrals as well as the initial MP2 optimization for the MP2 natural orbitals, and with Qiskit-aer³⁶ as the quantum simulation backend. All calculations used the MP2 natural orbitals as their initial guess. After the oo-UCC optimization, the RDMs in the active space were calculated using SlowQuant and then fed into our standalone DensityMatrixDrivenModule (DMDM) for the calculation of the complete space RDMs according to Eqs. (51) to (54) and, subsequently, of the linear response properties, specifically excitation energies and spectral strengths. The classic CASSCF reference calculations were performed using Dalton³⁷ and MP2 natural orbitals as the initial guess.

To show the performance capabilities of using RDMs for naive qLRSD, excitation energies and oscillator strengths were calculated for benzene and the excitation energies, oscillator strengths, and rotational strengths of *R*-methyloxirane were calculated using active space RDMs from an oo-UCCSD(6,6)/ cc-pVTZ³⁸ calculation in SlowQuant. The absorption spectrum of benzene and the ECD spectrum of *R*-methyloxirane were then compared to CASSCF(6,6)/cc-pVTZ results obtained from Dalton.

Finally, a shot noise analysis was performed. For this, 10 samples of the absorption spectrum of water were calculated on top of a noise-free oo-UCCSD(4,4)/cc-pVTZ wave function using 10k, 50k, and 100k shots. The active space RDMs were simulated through SlowQuant using Qiskit-aer as the quantum simulator backend. The wave function on the quantum simulator used a UCCSD ansatz with Parity mapping. The reference noiseless absorption spectrum was

obtained using SlowQuant.

All geometries used are available in Section S3 of the SI, and examples of code calculating excitation energies, oscillator strengths, and rotational strengths using DMDM are available as well in Section S4 of the SI. All spectra use a Lorentzian convolution with FWHM broadening of 0.2 eV and compare the first 399 excitation energies as this is the maximum that Dalton allows by default. Note that this exceedingly large number of states is used with the sole purpose of comparing CASSCF with oo-UCCSD on a broad spectral range; not all states are expected to be physically significant.

4 **Results and Discussion**

Having derived working equations for an RDM formulation of naive qLRSD, we will in the following section compare its results against CASSCF(6,6) LR for the calculation of excitation energies, oscillator strengths, and rotational strengths. For this, we provide calculated absorption spectra of benzene in Sec. 4.1 and ECD spectra of *R*-methyloxirane in Sec. 4.2. Finally, we present an analysis of our formulation's resilience to shot noise in Sec. 4.3.

4.1 Benzene

In Fig. 1, we show the absorption spectra of benzene using CASSCF(6,6) LR (blue), and qLRSD (dashed orange). Note that the CASSCF(6,6) LR calculation includes the complete active excitation space whereas qLRSD only includes singles and doubles excitations in the active excitation space, leading to a reduction in active space parameters from 174 for CASSCF(6,6) LR to 54 for qLRSD. The reduced excitation space does not result in a significant change in the absorption spectrum, with most peaks being identical in both excitation energy and oscillator strength. The minor deviations between the two spectra are caused by the truncated active excitation space in the linear response. To gain a more accurate description for these excitations, a higher active excitation space is needed¹³.



Figure 1: Benzene. Absorption spectrum of the lowest 399 excitation energies using CASSCF(6,6) LR (blue), and qLRSD (dashed orange) based on the oo-UCCSD(6,6) wave function using the cc-pVTZ basis set.

4.2 *R*-methyloxirane

The ECD spectra of *R*-methyloxirane using CASSCF(6,6) LR and qLRSD are shown in Fig. 2. In the SI, the ECD spectrum containing all 399 calculated excitation energies (Fig. S4) and the absorption spectrum (S3) can be found (Section S5).

The truncated active excitation space of qLRSD leads to the same parameter reduction as for benzene, whilst no discernible difference between the CASSCF(6,6) and the qLRSD spectra is visible. Therefore, the truncated excitation space is sufficient in describing the excitation energies and rotational strengths of *R*-methyloxirane. As can be appreciated from Fig. S3, this also applies (not surprisingly) to the oscillator strengths of *R*-methyloxirane where truncating the excitation space to singles and doubles proves sufficient for oscillator strengths of comparable quality to CASSCF(6,6).



Figure 2: *R*-methyloxirane. ECD spectrum of the lowest 97 excitation energies using CASSCF(6,6) LR (blue), and qLRSD (dashed orange) based on the oo-UCCSD(6,6) wave function using the cc-pVTZ basis set.

4.3 Shot noise analysis

In Fig. 3, the absorption spectra of $H_2O(4,4)/cc$ -pVTZ with shot noise from a quantum simulator can be found. We show, respectively, part of the valence excitation energy region (8 to 16 eV) and around the oxygen K-edge (starting at 540 eV) using qLRSD. Exact values for variance and standard deviations in the two energy regions can be found in the SI in table S1 (Section S5). It is evident that a higher amount of shots leads to more reliable results. Moreover, the peaks show differences in their convergence to the ideal spectrum with increasing number of shots. Looking at individual peaks, the first three peaks in the low energy region and the peaks between 560 and 570 eV of Fig. 3 have low errors when comparing the simulated spectra with shot noise to the ideal spectrum. The remaining peaks converge toward the ideal result but still show some deviations like the fourth peak in the low energy region and the first peak in the high energy region of Fig. 3. Significant variance in the region >580 eV of Fig. 3 also appears. A more in-depth analysis of the excitation energies and oscillator strengths to identify why some peaks have larger variance and standard deviation is a topic for future research.



Figure 3: Water. Absorption spectrum in the low energy region of 8 to 16 eV (valence excitations, left) and in the high energy region of 540 to 600 eV (oxygen K-edge, right) on simulated hard-ware to model shot noise. Top panels: 10 000 shots; middle panels: 50 000 shots; bottom panels: 100 000 shots. Each panel contains ten spectra including shot noise and one 'ideal' spectrum based on a state-vector calculation.

5 Summary

In this paper, we have formulated and implemented a density-matrix-driven approach to our recent naive qLRSD approach, that drastically reduces the classical computational costs. This will allow for more demanding calculations employing larger basis sets and on larger molecules. To illustrate our approach, we reported the absorption spectrum of benzene and the ECD spectrum of *R*-methyloxirane calculated in a (6,6) active space using the cc-pVTZ basis set. These constitute the largest basis sets employed so far for quantum approaches to molecular properties on moderate sized molecules and is an important step to achieving meaningful results on NISQ-era quantum devices. The truncation of the excitation rank in the active space leads to a significant reduction in the number of active space parameters with results almost indistinguishable from classic CASSCF simulations with a complete active space.

Using the active space approximation allows us to separate the RDMs in the complete space into a sum of RDMs over different indices in the inactive and active spaces, where only the latter necessitates evaluation on quantum device. Crucially, when not all indices are in the active space, the order of the RDMs can be significantly reduced. Moreover, the symmetries of the RDMs leads to a reduction of the number of elements in the RDMs that need to be measured. Depending on the number of symmetries in the different order of RDMs, this can lead to a reduction by a factor of up to 48 for the 4-RDM.

Additionally, we provide proof-of-principle quantum simulations with shot noise from a simulated quantum backend. In the analysis of shot noise, we observe the trivial trend that more shots converge to the ideal spectrum. However, some peaks converge quickly, while others have larger variances. The reason behind this observation is a point for further investigation. We are currently investigating the impact of using a cumulant-based density matrix approach where higher-order RDMs, such as the 3- and 4-electron RDMs, are approximated from lower order RDMs^{39–41}. This would lead to a further reduction in classic computational costs and in quantum measurements.

Acknowledgments

We acknowledge support from the Novo Nordisk Foundation (NNF) for the focused research project *Hybrid Quantum Chemistry on Hybrid Quantum Computers* (HQC)² (grant number: NN-FSA220080996).

References

- Olsen, J.; Jørgensen, P. Linear and nonlinear response functions for an exact state and for an MCSCF state. J. Chem. Phys. 1985, 82, 3235–3264.
- (2) Christiansen, O.; Jørgensen, P.; Hättig, C. Response functions from Fourier component variational perturbation theory applied to a time-averaged quasienergy. *Int. J. Quantum Chem.* 1998, 68, 1–52.
- (3) Helgaker, T.; Coriani, S.; Jørgensen, P.; Kristensen, K.; Olsen, J.; Ruud, K. Recent advances in wave function-based methods of molecular-property calculations. *Chem. Rev.* 2012, *112*, 543–631.
- (4) Pawłowski, F.; Olsen, J.; Jørgensen, P. Molecular response properties from a Hermitian eigenvalue equation for a time-periodic Hamiltonian. *J. Chem. Phys.* **2015**, *142*, 114109.
- (5) Norman, P.; Jonsson, D.; Vahtras, O.; Ågren, H. Cubic response functions in the random phase approximation. *Chem. Phys. Lett.* **1995**, *242*, 7–16.
- (6) Hettema, H.; Jensen, H. J. A.; Jørgensen, P.; Olsen, J. Quadratic response functions for a multiconfigurational self-consistent field wave function. J. Chem. Phys. 1992, 97, 1174–1190.
- (7) Jonsson, D.; Norman, P.; Ågren, H. Cubic response functions in the multiconfiguration selfconsistent field approximation. *J. Chem. Phys.* **1996**, *105*, 6401–6419.
- (8) Koch, H.; Jørgensen, P. Coupled cluster response functions. J. Chem. Phys. 1990, 93, 3333–3344.
- (9) Olsen, J.; Jørgensen, P.; Helgaker, T.; Oddershede, J. Quadratic response functions in a second-order polarization propagator framework. *J. Phys. Chem. A* **2005**, *109*, 11618–11628.
- (10) Sałek, P.; Vahtras, O.; Helgaker, T.; Ågren, H. Density-functional theory of linear and nonlinear time-dependent molecular properties. J. Chem. Phys. 2002, 117, 9630 – 9645.

- (11) Parker, S. M.; Rappoport, D.; Furche, F. Quadratic Response Properties from TDDFT: Trials and Tribulations. J. Chem. Theory Comput. 2018, 14, 807–819.
- (12) Kumar, A.; Asthana, A.; Abraham, V.; Crawford, T. D.; Mayhall, N. J.; Zhang, Y.; Cincio, L.; Tretiak, S.; Dub, P. A. Quantum Simulation of Molecular Response Properties in the NISQ Era. J. Chem. Theory and Comput. 2023, 19, 9136–9150.
- (13) Ziems, K. M.; Kjellgren, E. R.; Reinholdt, P.; Jensen, P. W. K.; Sauer, S. P. A.; Kongsted, J.; Coriani, S. Which options exist for NISQ-friendly linear response formulations? *J. Chem. Theory Comput.* 2024,
- (14) Chen, S.; Cotler, J.; Huang, H.-Y.; Li, J. The complexity of NISQ. *Nat. Commun.* 2023, 14, 6001.
- (15) McClean, J. R.; Romero, J.; Babbush, R.; Aspuru-Guzik, A. The theory of variational hybrid quantum-classical algorithms. *New J. Phys.* **2016**, *18*, 023023.
- (16) Yeager, D. L.; Jørgensen, P. A multiconfigurational time-dependent Hartree-Fock approach. *Chem. Phys. Lett.* **1979**, *65*, 77–80.
- (17) Dalgaard, E. Time-dependent multiconfigurational Hartree–Fock theory. J. Chem. Phys. 1980, 72, 816–823.
- (18) Roos, B. O.; Taylor, P. R.; Sigbahn, P. E. A complete active space SCF method (CASSCF) using a density matrix formulated super-CI approach. *Chem. Phys.* **1980**, *48*, 157–173.
- (19) Siegbahn, P.; Heiberg, A.; Roos, B.; Levy, B. A Comparison of the Super-CI and the Newton-Raphson Scheme in the Complete Active Space SCF Method. *Phys. Scr.* **1980**, *21*, 323–327.
- (20) Siegbahn, P. E. M.; Almlöf, J.; Heiberg, A.; Roos, B. O. The complete active space SCF (CASSCF) method in a Newton–Raphson formulation with application to the HNO molecule. *J. Chem. Phys.* 1981, 74, 2384–2396.

- (21) Ollitrault, P. J.; Kandala, A.; Chen, C.-F.; Barkoutsos, P. K.; Mezzacapo, A.; Pistoia, M.; Sheldon, S.; Woerner, S.; Gambetta, J. M.; Tavernelli, I. Quantum equation of motion for computing molecular excitation energies on a noisy quantum processor. *Phys. Rev. Res.* 2020, 2, 043140.
- (22) Jensen, P. W. K.; Kjellgren, E. R.; Reinholdt, P.; Ziems, K. M.; Coriani, S.; Kongsted, J.; Sauer, S. P. A. Quantum Equation of Motion with Orbital Optimization for Computing Molecular Properties in Near-Term Quantum Computing. *J. Chem. Theory Comput.* 2024, in press.
- (23) Fosso-Tande, J.; Nguyen, T.-S.; Gidofalvi, G.; DePrince III, A. E. Large-scale variational two-electron reduced-density-matrix-driven complete active space self-consistent field methods. J. Chem. Theory Comput. 2016, 12, 2260–2271.
- (24) Mazziotti, D. A. Parametrization of the two-electron reduced density matrix for its direct calculation without the many-electron wave function. *Phys. Rev. Lett.* **2008**, *101*, 253002.
- (25) McWeeny, R. The density matrix in self-consistent field theory I. Iterative construction of the density matrix. *Proc. R. Soc. Lond. A Math. Phys. Sci.* **1956**, *235*, 496–509.
- (26) Rice, J.; Amos, R. On the efficient evaluation of analytic energy gradients. *Chem. Phys. Lett.* 1985, *122*, 585–590.
- (27) Bartlett, R. J.; Kucharski, S. A.; Noga, J. Alternative coupled-cluster ansätze II. The unitary coupled-cluster method. *Chem. Phys. Lett.* **1989**, *155*, 133–140.
- (28) Mizukami, W.; Mitarai, K.; Nakagawa, Y. O.; Yamamoto, T.; Yan, T.; Ohnishi, Y. Orbital optimized unitary coupled cluster theory for quantum computer. *Phys. Rev. Res.* **2020**, *2*, 033421.
- (29) Sokolov, I. O.; Barkoutsos, P. K.; Ollitrault, P. J.; Greenberg, D.; Rice, J.; Pistoia, M.; Tavernelli, I. Quantum orbital-optimized unitary coupled cluster methods in the strongly corre-

lated regime: Can quantum algorithms outperform their classical equivalents? *J. Chem. Phys.* **2020**, *152*.

- (30) Jørgensen, P.; Jensen, H. J. A.; Olsen, J. Linear response calculations for large scale multiconfiguration self-consistent field wave functions. *J. Chem. Phys.* **1988**, *89*, 3654–3661.
- (31) Helgaker, T.; Jørgensen, P.; Olsen, J. Molecular electronic-structure theory; John Wiley & Sons: Nashville, TN, 2013.
- (32) Reinholdt, P.; Kjellgren, E. R.; Fuglsbjerg, J. H.; Ziems, K. M.; Coriani, S.; Sauer, S.; Kongsted, J. Subspace methods for the simulation of molecular response properties on a quantum computer. *arXiv preprint arXiv:2402.12186* **2024**,
- (33) Kjellgren, E.; Ziems, K. M. SlowQuant.
- (34) Sun, Q. Libcint: An efficient general integral library for Gaussian basis functions. J. Comput. Chem. 2015, 36, 1664–1671.
- (35) Sun, Q.; Zhang, X.; Banerjee, S.; Bao, P.; Barbry, M.; Blunt, N. S.; Bogdanov, N. A.; Booth, G. H.; Chen, J.; Cui, Z.-H.; Eriksen, J. J.; Gao, Y.; Guo, S.; Hermann, J.; Hermes, M. R.; Koh, K.; Koval, P.; Lehtola, S.; Li, Z.; Liu, J.; Mardirossian, N.; McClain, J. D.; Motta, M.; Mussard, B.; Pham, H. Q.; Pulkin, A.; Purwanto, W.; Robinson, P. J.; Ronca, E.; Sayfutyarova, E. R.; Scheurer, M.; Schurkus, H. F.; Smith, J. E. T.; Sun, C.; Sun, S.-N.; Upadhyay, S.; Wagner, L. K.; Wang, X.; White, A.; Whitfield, J. D.; Williamson, M. J.; Wouters, S.; Yang, J.; Yu, J. M.; Zhu, T.; Berkelbach, T. C.; Sharma, S.; Sokolov, A. Y.; Chan, G. K.-L. Recent developments in the PySCF program package. *J. Chem. Phys.* 2020, *153*.
- (36) Qiskit contributors Qiskit: An Open-source Framework for Quantum Computing. 2023.
- (37) Aidas, K.; Angeli, C.; Bak, K. L.; Bakken, V.; Bast, R.; Boman, L.; Christiansen, O.; Cimiraglia, R.; Coriani, S.; Dahle, P.; Dalskov, E. K.; Ekström, U.; Enevoldsen, T.; Eriksen, J. J.;

Ettenhuber, P.; Fernández, B.; Ferrighi, L.; Fliegl, H.; Frediani, L.; Hald, K.; Halkier, A.; Hättig, C.; Heiberg, H.; Helgaker, T.; Hennum, A. C.; Hettema, H.; Hjertenaes, E.; Høst, S.; Høyvik, I.-M.; Iozzi, M. F.; Jansík, B.; Jensen, H. J. A.; Jonsson, D.; Jørgensen, P.; Kauczor, J.; Kirpekar, S.; Kjaergaard, T.; Klopper, W.; Knecht, S.; Kobayashi, R.; Koch, H.; Kongsted, J.; Krapp, A.; Kristensen, K.; Ligabue, A.; Lutnaes, O. B.; Melo, J. I.; Mikkelsen, K. V.; Myhre, R. H.; Neiss, C.; Nielsen, C. B.; Norman, P.; Olsen, J.; Olsen, J. M. H.; Osted, A.; Packer, M. J.; Pawlowski, F.; Pedersen, T. B.; Provasi, P. F.; Reine, S.; Rinkevicius, Z.; Ruden, T. A.; Ruud, K.; Rybkin, V. V.; Sałek, P.; Samson, C. C. M.; de Merás, A. S.; Saue, T.; Sauer, S. P. A.; Schimmelpfennig, B.; Sneskov, K.; Steindal, A. H.; Sylvester-Hvid, K. O.; Taylor, P. R.; Teale, A. M.; Tellgren, E. I.; Tew, D. P.; Thorvaldsen, A. J.; Thøgersen, L.; Vahtras, O.; Watson, M. A.; Wilson, D. J. D.; Ziolkowski, M.; Ågren, H. The Dalton quantum chemistry program system. *WIREs Comput Mol Sci* **2013**, *4*, 269–284.

- (38) Dunning, T. H. Gaussian basis sets for use in correlated molecular calculations. I. The atoms boron through neon and hydrogen. J. Chem. Phys. 1989, 90, 1007–1023.
- (39) Harris, F. E. Cumulant-based approximations to reduced density matrices. *Int. J. Quantum Chem.* **2002**, *90*, 105–113.
- (40) Zgid, D.; Ghosh, D.; Neuscamman, E.; Chan, G. K.-L. A study of cumulant approximations to n-electron valence multireference perturbation theory. J. Chem. Phys. 2009, 130.
- (41) Saitow, M.; Kurashige, Y.; Yanai, T. Multireference configuration interaction theory using cumulant reconstruction with internal contraction of density matrix renormalization group wave function. J. Chem. Phys. 2013, 139.

Supplementary Information for Reduced density matrix formulation of quantum linear response

Theo Juncker von Buchwald,*,† Karl Michael Ziems,*,† Erik Rosendahl

Kjellgren,*^{,‡} Stephan P. A. Sauer,[¶] Jacob Kongsted,[‡] and Sonia Coriani[†]

†Department of Chemistry, Technical University of Denmark, Kemitorvet Building 207, DK-2800 Kongens Lyngby, Denmark.

[‡]Department of Physics, Chemistry and Pharmacy, University of Southern Denmark, Campusvej 55, 5230 Odense, Denmark.

¶Department of Chemistry, University of Copenhagen, DK-2100 Copenhagen \emptyset .

E-mail: tjvbu@kemi.dtu.dk; kmizi@kemi.dtu.dk; kjellgren@sdu.dk

April 25, 2024

S1Symmetries

The symmetries of the RDMs are given below

$$\Gamma_{pq}^{[1]} = \Gamma_{qp}^{[1]} \tag{S1}$$

$$\Gamma_{pqrs}^{[2]} = \Gamma_{rspq}^{[2]} = \Gamma_{qpsr}^{[2]} = \Gamma_{srqp}^{[2]}$$
(S2)

[3]

[2]

$$\Gamma_{pqrstu}^{[3]} = \Gamma_{pqturs}^{[3]} = \Gamma_{rspqtu}^{[3]} = \Gamma_{rstupq}^{[3]} = \Gamma_{tupqrs}^{[3]} = \Gamma_{turspq}^{[3]}$$
(S3)

$$=\Gamma_{qpsrut}^{[3]} = \Gamma_{qputsr}^{[3]} = \Gamma_{srqput}^{[3]} = \Gamma_{srutqp}^{[3]} = \Gamma_{utqpsr}^{[3]} = \Gamma_{utsrqp}^{[3]}$$

$$\Gamma_{pqrstumn}^{[4]} = \Gamma_{pqtursmn}^{[4]} = \Gamma_{pqtursmn}^{[4]} = \Gamma_{pqtursmrs}^{[4]} = \Gamma_{pqmnrstu}^{[4]} = \Gamma_{pqmnturs}^{[4]}$$

$$=\Gamma_{rspqtumn}^{[4]} = \Gamma_{rspqmntu}^{[4]} = \Gamma_{rstupqmn}^{[4]} = \Gamma_{rstumpq}^{[4]} = \Gamma_{rsmnpqtu}^{[4]} = \Gamma_{rsmntupq}^{[4]}$$

$$=\Gamma_{tupqrsmn}^{[4]} = \Gamma_{tupqmnrs}^{[4]} = \Gamma_{turspqmn}^{[4]} = \Gamma_{tursmnpq}^{[4]} = \Gamma_{tumnpqrs}^{[4]} = \Gamma_{tumnrspq}^{[4]}$$

$$=\Gamma_{mnpqrstu}^{[4]} = \Gamma_{mnpqturs}^{[4]} = \Gamma_{mrspqtu}^{[4]} = \Gamma_{mrstupq}^{[4]} = \Gamma_{mntupqrs}^{[4]} = \Gamma_{mnturspq}^{[4]}$$

$$=\Gamma_{qpsrutnm}^{[4]} = \Gamma_{qpsrnmut}^{[4]} = \Gamma_{qputsrnm}^{[4]} = \Gamma_{qputnmsr}^{[4]} = \Gamma_{qpnmsrut}^{[4]} = \Gamma_{srnmutqp}^{[4]}$$

$$=\Gamma_{srqputnm}^{[4]} = \Gamma_{srqpnmut}^{[4]} = \Gamma_{srutqpmm}^{[4]} = \Gamma_{srutmqp}^{[4]} = \Gamma_{srnmqput}^{[4]} = \Gamma_{srnmutqp}^{[4]}$$

$$=\Gamma_{utqpsrnm}^{[4]} = \Gamma_{utqpnmsr}^{[4]} = \Gamma_{utsrqpnm}^{[4]} = \Gamma_{srutmqp}^{[4]} = \Gamma_{srnmqput}^{[4]} = \Gamma_{utnmsrqp}^{[4]}$$

$$=\Gamma_{utqpsrnm}^{[4]} = \Gamma_{utqpnmsr}^{[4]} = \Gamma_{utsrqpnm}^{[4]} = \Gamma_{utsrmqp}^{[4]} = \Gamma_{utnmqpsr}^{[4]} = \Gamma_{utnmsrqp}^{[4]}$$

S2Working equations

The key equations of linear reponse are the linear response equation, Eq. (24), and its homogeneous eigenvalue equation, Eq. (36). To solve them, the Hessian and the Hermitian metric must be calculated first. The Hessian and the Hermitian metric contain the submatrices in Eqs. (26) to (29). With the definition of the orbital rotation operators \hat{q}_{μ} and \hat{G}_n in Eqs. (21) and (22), these may be expressed by the matrix elements in Eqs. (S5) to (S28) as RDMs. All indices used in this section should be read as general indices.

$$\left\langle 0 \left| \left[\hat{E}_{mn}, \sum_{pq} h_{pq} \hat{E}_{pq} \right] \right| 0 \right\rangle$$

$$= \sum_{p} h_{np} \Gamma_{mp}^{[1]} - \sum_{p} h_{pm} \Gamma_{pn}^{[1]}$$
(S5)

$$\left\langle 0 \left| \left[\hat{E}_{mn}, \sum_{pqrs} g_{pqrs} \hat{e}_{pqrs} \right] \right| 0 \right\rangle$$

$$= 2 \sum_{pqr} g_{npqr} \Gamma^{[2]}_{mpqr} - 2 \sum_{pqr} g_{pmqr} \Gamma^{[2]}_{pnqr}$$
(S6)

$$\left\langle 0 \left| \left[\hat{E}_{tu} \hat{E}_{mn}, \sum_{pq} h_{pq} \hat{E}_{pq} \right] \right| 0 \right\rangle$$

$$= \sum_{q} h_{uq} \Gamma_{tqmn}^{(2)} - \sum_{p} h_{pt} \left(\Gamma_{pumn}^{(2)} - \delta_{mu} \Gamma_{pn}^{(1)} \right)$$

$$+ \sum_{q} h_{nq} \left(\Gamma_{tumq}^{(2)} + \delta_{mu} \Gamma_{tq}^{(1)} \right) - \sum_{p} h_{pm} \Gamma_{tupn}^{(2)}$$
(S7)

$$\left\langle 0 \left| \left[\hat{E}_{tu} \hat{E}_{mn}, \sum_{pqrs} g_{pqrs} \hat{e}_{pqrs} \right] \right| 0 \right\rangle$$

$$= 2 \left(\sum_{pqr} g_{npqr} \Gamma_{tumpqr}^{[3]} + \sum_{pqr} g_{upqr} \Gamma_{mntpqr}^{[3]} - \sum_{pqr} g_{pmqr} \Gamma_{tupnqr}^{[3]} - \sum_{pqr} g_{ptqr} \Gamma_{mnpuqr}^{[3]} \right)$$

$$+ \sum_{pqr} \delta_{mu} g_{pqnr} \Gamma_{trpq}^{[2]} - \sum_{pqr} \delta_{mu} g_{ptqr} \Gamma_{pnqr}^{[2]} + \sum_{pq} g_{upnq} \Gamma_{tpmq}^{[2]} - \sum_{pq} g_{pmqt} \Gamma_{pnqu}^{[2]} \right)$$

$$(S8)$$

$$\left\langle 0 \left| \left[\hat{E}_{tu}, \left[\sum_{pq} h_{pq} \hat{E}_{pq}, \hat{E}_{mn} \right] \right] \right| 0 \right\rangle$$

$$= -\sum_{p} h_{np} \delta_{mu} \Gamma_{tp}^{[1]} + h_{nt} \Gamma_{mu}^{[1]} + h_{um} \Gamma_{tn}^{[1]} - \sum_{p} h_{pm} \delta_{tn} \Gamma_{pu}^{[1]}$$
(S10)

$$\left\langle 0 \left| \left[\hat{E}_{tu}, \left[\sum_{pqrs} g_{pqrs} \hat{e}_{pqrs}, \hat{E}_{mn} \right] \right] \right| 0 \right\rangle$$

$$= -2 \sum_{pqr} g_{npqr} \delta_{mu} \Gamma_{tpqr}^{[2]} + 2 \sum_{pq} g_{ntpq} \Gamma_{mupq}^{[2]} - 2 \sum_{pq} g_{npuq} \Gamma_{mptq}^{[2]} + 2 \sum_{pq} g_{npqt} \Gamma_{mpqu}^{[2]}$$

$$+ 2 \sum_{pq} g_{umpq} \Gamma_{tnpq}^{[2]} - 2 \sum_{pqr} g_{pmqr} \delta_{tn} \Gamma_{puqr}^{[2]} + 2 \sum_{pq} g_{pmuq} \Gamma_{pntq}^{[2]} - 2 \sum_{pq} g_{pmqt} \Gamma_{pnqu}^{[2]}$$

$$(S11)$$

$$\left\langle 0 \left| \left[\hat{E}_{ij} \hat{E}_{kl}, \left[\sum_{pq} h_{pq} \hat{E}_{pq}, \hat{E}_{tu} \right] \right] \right| 0 \right\rangle$$

$$= h_{ui} \Gamma_{tjkl}^{[2]} + h_{uk} \Gamma_{ijtl}^{[2]} + h_{jk} \Gamma_{iukl}^{[2]} + h_{lt} \Gamma_{ijku}^{[2]} + \delta_{kj} h_{ui} \Gamma_{tl}^{[1]} + \delta_{kj} h_{lt} \Gamma_{iu}^{[1]}$$

$$- \sum_{p} \delta_{tj} h_{up} \Gamma_{ipkl}^{[2]} - \sum_{p} \delta_{tl} h_{up} \Gamma_{ijkp}^{[2]} - \sum_{p} \delta_{iu} h_{pt} \Gamma_{pjkl}^{[2]} - \sum_{p} \delta_{ku} h_{pt} \Gamma_{ijpl}^{[2]}$$

$$- \sum_{p} \delta_{tl} \delta_{kj} h_{up} \Gamma_{ip}^{[1]} - \sum_{p} \delta_{iu} \delta_{kj} h_{up} \Gamma_{pl}^{[1]}$$
(S12)

$$\left\langle 0 \left| \left[\hat{E}_{ij} \hat{E}_{kl}, \left[\sum_{pqrs} g_{pqrs} \hat{e}_{pqrs}, \hat{E}_{tu} \right] \right] \right| 0 \right\rangle$$

$$= 2 \left(\sum_{p} g_{jtlp} \Gamma_{iukp}^{[2]} + \sum_{p} g_{ltjp} \Gamma_{ipku}^{[2]} + \sum_{p} g_{uipk} \Gamma_{tjpl}^{[2]} + \sum_{p} g_{piuk} \Gamma_{pjtl}^{[2]} + \sum_{pq} g_{ukpq} \Gamma_{ijtpq}^{[3]} \right.$$

$$+ \sum_{pq} g_{upqk} \Gamma_{ijtpql}^{[3]} + \sum_{pq} g_{uipq} \Gamma_{tjpqkl}^{[3]} + \sum_{pq} g_{upqi} \Gamma_{tpqjkl}^{[3]} + \sum_{pq} g_{ptlq} \Gamma_{ijpukq}^{[3]} + \sum_{pq} g_{ltpq} \Gamma_{ijpkq}^{[3]} \right.$$

$$+ \sum_{pq} g_{jtpq} \Gamma_{iupqkl}^{[3]} + \sum_{pq} g_{ptjq} \Gamma_{puiqkl}^{[3]} - \sum_{pq} g_{upjq} \Gamma_{tpiqkl}^{[3]} - \sum_{pq} g_{ptjq} \Gamma_{ijpkq}^{[3]} - \sum_{pq} g_{ptqk} \Gamma_{ijpuql}^{[3]} - \sum_{pq} g_{ptqi} \Gamma_{piqkl}^{[2]} + \sum_{pq} \delta_{kj} g_{upqi} \Gamma_{tpq}^{[2]} + \sum_{pq} \delta_{kj} g_{upqi} \Gamma_{ipql}^{[2]} + \sum_{pq} \delta_{kj} g_{ptqi} \Gamma_{ipq}^{[2]} \right.$$

$$+ \sum_{pq} g_{uplq} \Gamma_{ijtpkq}^{[3]} - \sum_{pq} \delta_{iu} g_{ptqk} \Gamma_{pjql}^{[2]} + \sum_{pq} \delta_{kj} g_{upqi} \Gamma_{tpq}^{[2]} + \sum_{pq} \delta_{kj} g_{upqi} \Gamma_{tpq}^{[2]} \right.$$

$$+ \sum_{pq} \delta_{kj} g_{ltpq} \Gamma_{iupq}^{[2]} - \sum_{pq} \delta_{kj} g_{uplq} \Gamma_{iqtp}^{[2]} - \sum_{pq} \delta_{kj} g_{uplq} \Gamma_{ipq}^{[2]} - \sum_{pq} \delta_{kj} g_{uplq} \Gamma_{ipqkl}^{[2]} - \sum_{pq} \delta_{tj} g_{upq} \Gamma_{ipqkl}^{[2]} - \sum_{pq} \delta_{tj} g_{upq} \Gamma_{ipqkl}^{[2]} - \sum_{pq} \delta_{tj} \delta_{tj} \delta_{tj} g_{upq} \Gamma_$$

$$\left\langle 0 \left| \left[\hat{E}_{ij}, \left[\sum_{pq} h_{pq} \hat{E}_{pq}, \hat{E}_{tu} \hat{E}_{mn} \right] \right] \right| 0 \right\rangle$$

$$= h_{jt} \Gamma_{iumn}^{[2]} + h_{ui} \Gamma_{tjmn}^{[2]} + h_{jm} \Gamma_{tuin}^{[2]} + h_{ni} \Gamma_{tumj}^{[2]} + \delta_{mu} h_{jt} \Gamma_{in}^{[1]} + \delta_{mu} h_{ni} \Gamma_{tj}^{[1]} \right.$$

$$+ \sum_{p} \delta_{iu} h_{np} \Gamma_{tjmp}^{[2]} - \sum_{p} \delta_{iu} h_{pq} \Gamma_{pjmn}^{[2]} - \sum_{p} \delta_{iu} h_{pm} \Gamma_{tjpn}^{[2]} + \sum_{p} \delta_{mj} h_{pt} \Gamma_{puin}^{[2]} \right.$$

$$- \sum_{p} \delta_{mj} h_{up} \Gamma_{tpin}^{[2]} - \sum_{p} \delta_{mj} h_{np} \Gamma_{tuip}^{[2]} + \sum_{p} \delta_{in} h_{up} \Gamma_{tpmj}^{[2]} - \sum_{p} \delta_{in} h_{pt} \Gamma_{pumj}^{[2]} \right.$$

$$- \sum_{p} \delta_{in} h_{pm} \Gamma_{tupj}^{[2]} + \sum_{p} \delta_{tj} h_{pm} \Gamma_{iupn}^{[2]} - \sum_{p} \delta_{tj} h_{up} \Gamma_{ipmn}^{[2]} - \sum_{p} \delta_{tj} h_{np} \Gamma_{iump}^{[2]} - \sum_{p} \delta_{in} \delta_{mu} h_{pt} \Gamma_{pj}^{[1]} \right.$$

$$- \sum_{p} \delta_{in} \delta_{mu} h_{pt} \Gamma_{pj}^{[1]} - \sum_{p} \delta_{tj} \delta_{mu} h_{np} \Gamma_{ip}^{[1]} \right.$$

$$\left\langle 0 \left| \left[\hat{E}_{ij}, \left[\sum_{pqrs} g_{pqrs} \hat{e}_{pqrs}, \hat{E}_{tu} \hat{E}_{mn} \right] \right] \right| 0 \right\rangle$$

$$= 2 \left(\sum_{p} g_{niup} \Gamma_{tpmj}^{[2]} + \sum_{p} g_{npui} \Gamma_{tjmp}^{[2]} + \sum_{p} g_{jtpm} \Gamma_{iupn}^{[2]} + \sum_{p} g_{ptjm} \Gamma_{puin}^{[2]} + \sum_{pq} g_{jmpq} \Gamma_{tunpq}^{[3]} \right. \\ \left. + \sum_{pq} g_{pmjq} \Gamma_{tupniq}^{[3]} + \sum_{pq} g_{nipq} \Gamma_{tumjpq}^{[3]} + \sum_{pq} g_{npqi} \Gamma_{tupnqj}^{[3]} + \sum_{pq} g_{jtpq} \Gamma_{iupqmn}^{[3]} + \sum_{pq} g_{ptjq} \Gamma_{puin}^{[3]} + \sum_{pq} g_{ptjq} \Gamma_{puinq}^{[3]} + \sum_{pq} g_{ptjq} \Gamma_{puinqn}^{[3]} + \sum_{pq} g_{ptjq} \Gamma_{puinqn}^{[3]} + \sum_{pq} g_{npjq} \Gamma_{puinqn}^{[3]} + \sum_{pq} g_{npjq} \Gamma_{puinqn}^{[3]} + \sum_{pq} g_{mu} g_{npiq} \Gamma_{puinqn}^{[2]} + \sum_{pq} \delta_{mu} g_{ptjq} \Gamma_{puinqn}^{[3]} + \sum_{pq} \delta_{mu} g_{ptqn} \Gamma_{puinqn}^{[3]} + \sum_{pq} \delta_{mu} g_{ptqn} \Gamma_{puinqn}^{[3]} + \sum_{pq} \delta_{iu} g_{pnqn} \Gamma_{iupnqn}^{[3]} + \sum_{pq} \delta_{iu} g_{pnqn} \Gamma_{ipnqn}^{[3]} + \sum_{pq} \delta_{iu} g_{pnqn} \Gamma_{ipnqn}^{[3]} + \sum_{pq} \delta_{iu} g_{pnqn} \Gamma_{iupnqn}^{[3]} + \sum_{pq} \delta_{iu} g_{pnqn} \Gamma_{iupnqn}^{[3]} - \sum_{pqr} \delta_{iu} g_{pnqr} \Gamma_{iupnqr}^{[3]} + \sum_{pqr} \delta_{iu} g_{pnqr} \Gamma_{iupnqr}^{[3]} - \sum_{pqr} \delta_{iu} g_{pnqr} \Gamma_{iupnqr}^{[3]} + \sum_{pqr} \delta_{iu} g_{pnqr} \Gamma_{iupnqr}^{[3]} - \sum_{pqr} \delta_{iu} g_{pnqr} \Gamma_{iupnqr}^{[3]} + \sum_{pqr} \delta_{in} g_{pnqr} \Gamma_{iupnqr}^{[3]} - \sum_{pqr} \delta_{in} g_{pnqr} \Gamma_{iupnqr}^{[3]} - \sum_{pqr} \delta_{in} g_{pnqr} \Gamma_{iupnq$$

$$\left\langle 0 \left| \left[\hat{E}_{ij} \hat{E}_{kl}, \left[\sum_{pq} h_{pq} \hat{E}_{pq}, \hat{E}_{lu} \hat{E}_{mn} \right] \right] \right| 0 \right\rangle$$

$$(S16)$$

$$= h_{uk} \Gamma_{ijtmn}^{[3]} + h_{ul} \Gamma_{ijmnkl}^{[3]} + h_{lt} \Gamma_{ijkmnn}^{[3]} + h_{jl} \Gamma_{immkl}^{[3]} + h_{lk} \Gamma_{ijtmnl}^{[3]} + h_{ml} \Gamma_{lamjkl}^{[3]} + h_{lm} \Gamma_{lamjkl}^{[3]} + h_{lm} \Gamma_{lim}^{[3]} + \delta_{kj} h_{ml} \Gamma_{lim}^{[2]} + \delta_{kj} h_{mn} \Gamma_{limn}^{[2]} + \delta_{kj} h_{mn} \Gamma_{limn}^{[2]} + \sum_{p} \delta_{ki} h_{mp} \Gamma_{limnkl}^{[2]} - \sum_{p} \delta_{ki} h_{pm} \Gamma_{limnkl}^{[2]} - \sum_{p} \delta_{ki} h_{pm} \Gamma_{limnkl}^{[2]} - \sum_{p} \delta_{ki} h_{pm} \Gamma_{limpkl}^{[2]} - \sum_{p} \delta_{ki} h_{kj} h_{mn} \Gamma_{limp}^{[2]} - \sum_{p} \delta_{ki} h_{kj} h_{mn} \Gamma_{limp}^{[2]} + \sum_{p} \delta_{ki} h_{kj} h_{mn} \Gamma_{limp}^{[2]} - \sum_{p$$

$$\left\langle 0 \left| \left[\hat{E}_{ij} \hat{E}_{kl}, \left[\sum_{pqrs} g_{pqrs} \hat{e}_{pqrs}, \hat{E}_{tu} \hat{E}_{mn} \right] \right] \right| 0 \right\rangle$$

$$= 2 \left(\mathcal{A} + \mathcal{B} + \mathcal{C} + \mathcal{D} + \mathcal{E} + \mathcal{F} + \mathcal{G} + \mathcal{H} \right)$$
(S17)

$$\mathcal{A} = g_{niuk}\Gamma^{[2]}_{tlmj} + g_{nkui}\Gamma^{[2]}_{tjml} + g_{ltjm}\Gamma^{[2]}_{inku} + g_{jtlm}\Gamma^{[2]}_{iukn}$$
(S18)

$$\mathcal{B} = \sum_{p} g_{lmjp} \Gamma_{tuipkn}^{[3]} + \sum_{p} g_{jmlp} \Gamma_{tuinkp}^{[3]} + \sum_{p} g_{nkup} \Gamma_{tpijml}^{[3]} + \sum_{p} g_{npuk} \Gamma_{tlijmp}^{[3]}$$
(S19)
+
$$\sum_{p} g_{nipk} \Gamma_{tumjpl}^{[3]} + \sum_{p} g_{niup} \Gamma_{tpmjpl}^{[3]} + \sum_{p} g_{npui} \Gamma_{tjmpkl}^{[3]} + \sum_{p} g_{nkpi} \Gamma_{tumlpj}^{[3]}$$
+
$$\sum_{p} g_{ltpm} \Gamma_{ijkupn}^{[3]} + \sum_{p} g_{ltjp} \Gamma_{ipkumn}^{[3]} + \sum_{p} g_{ptlm} \Gamma_{ijpukn}^{[3]} + \sum_{p} g_{jtlp} \Gamma_{iukpmn}^{[3]}$$
+
$$\sum_{p} g_{jtpm} \Gamma_{iupnkl}^{[3]} + \sum_{p} g_{ptjm} \Gamma_{puinkl}^{[3]} + \sum_{p} g_{uipk} \Gamma_{tjplmn}^{[3]} + \sum_{p} g_{ukpi} \Gamma_{tlpjmn}^{[3]}$$
+
$$\sum_{p} \delta_{kj} g_{niup} \Gamma_{tpml}^{[2]} + \sum_{p} \delta_{kj} g_{npui} \Gamma_{tlmp}^{[2]} + \sum_{p} \delta_{kj} g_{ltpm} \Gamma_{iupn}^{[2]} + \sum_{p} \delta_{mu} g_{nkpi} \Gamma_{tpp}^{[2]} + \sum_{p} \delta_{mu} g_{nkpi} \Gamma_{ippn}^{[2]} + \sum_{p} \delta_{mu} g_{nkpi} \Gamma_{ippn}^{[2]} + \sum_{p} \delta_{mu} g_{nkpi} \Gamma_{ippn}^{[2]} + \sum_{p} \delta_{mu} g_{nkpi} \Gamma_{tpp}^{[2]} + \sum_{p} \delta_{mu} g_{nkpi} \Gamma_{tpp}^{[2]} + \sum_{p} \delta_{mu} g_{nkpi} \Gamma_{ippn}^{[2]} + \sum_{p} \delta_{mu} g_{nkpi} \Gamma_{tpp}^{[2]} + \sum_{p} \delta_{mu} g_{nkpi} \Gamma_{tpp}^{[2]} + \sum_{p} \delta_{mu} g_{nkpi} \Gamma_{tpp}^{[2]} + \sum_{p} \delta_{mu} g_{nkpi} \Gamma_{ippn}^{[2]} + \sum_{p} \delta_{mu} g_{nkpi} \Gamma_{ippn}^{[2]} + \sum_{p} \delta_{mu} g_{nkpi} \Gamma_{tppn}^{[2]} + \sum_{p} \delta_{mu} g_{nkpi} \Gamma_{tppn}^{[2]} + \sum_{p} \delta_{mu} g_{nkpi} \Gamma_{ippn}^{[2]} + \sum_{p} \delta_{mu} g_{nkpi} \Gamma_{tppn}^{[2]} + \sum_{p} \delta$$

$$\mathcal{C} = \sum_{pq} g_{lmpq} \Gamma_{tuijknpq}^{[4]} + \sum_{pq} g_{pmlq} \Gamma_{tuijpnkq}^{[4]} + \sum_{pq} g_{nkpq} \Gamma_{tuijmlpq}^{[4]} + \sum_{pq} g_{npqk} \Gamma_{tuijmpql}^{[4]}$$
(S20)
+
$$\sum_{pq} g_{jmpq} \Gamma_{tuinpqkl}^{[4]} + \sum_{pq} g_{pmjq} \Gamma_{tupniqkl}^{[4]} + \sum_{pq} g_{nipq} \Gamma_{tumjpqkl}^{[4]} + \sum_{pq} g_{npqi} \Gamma_{tumpqjkl}^{[4]}$$
+
$$\sum_{pq} g_{ltpq} \Gamma_{ijkupqmn}^{[4]} + \sum_{pq} g_{ptlq} \Gamma_{ijpukqmn}^{[4]} + \sum_{pq} g_{ukpq} \Gamma_{ijtpqmn}^{[4]} + \sum_{pq} g_{upqk} \Gamma_{ijtpqmn}^{[4]}$$
+
$$\sum_{pq} g_{jtpq} \Gamma_{iupqklmn}^{[4]} + \sum_{pq} g_{ptjq} \Gamma_{puiqklmn}^{[4]} + \sum_{pq} g_{ukpq} \Gamma_{ijpqklmn}^{[4]} + \sum_{pq} g_{upqi} \Gamma_{ijpqklmn}^{[4]}$$
+
$$\sum_{pq} g_{ptqq} \Gamma_{iupqklmn}^{[4]} + \sum_{pq} g_{ptjq} \Gamma_{puiqklmn}^{[4]} + \sum_{pq} g_{upqq} \Gamma_{ijpqklmn}^{[4]} + \sum_{pq} g_{upqi} \Gamma_{tupnqjkl}^{[4]}$$
-
$$\sum_{pq} g_{nplq} \Gamma_{tuijmpkq}^{[4]} - \sum_{pq} g_{pmqk} \Gamma_{ijpkqmn}^{[4]} - \sum_{pq} g_{ptqi} \Gamma_{tupnqjkl}^{[4]} - \sum_{pq} g_{upqq} \Gamma_{ipqklmn}^{[4]}$$
-
$$\sum_{pq} g_{ptqk} \Gamma_{ijpuqlmn}^{[4]} - \sum_{pq} g_{uplq} \Gamma_{ijpkqmn}^{[4]} - \sum_{pq} g_{ptqi} \Gamma_{puiqklmn}^{[4]} - \sum_{pq} g_{upjq} \Gamma_{tupnqjkl}^{[4]}$$

$$\begin{aligned} \mathcal{D} &= \sum_{pq} \delta_{in} g_{upqk} \Gamma_{ipqlmj}^{[3]} + \sum_{pq} \delta_{in} g_{ukpq} \Gamma_{ilpqmj}^{[3]} + \sum_{pq} \delta_{in} g_{nkpq} \Gamma_{ijmpql}^{[3]} + \sum_{pq} \delta_{in} g_{npqk} \Gamma_{ijmpql}^{[3]} + \sum_{pq} \delta_{in} g_{npqk} \Gamma_{impql}^{[3]} + \sum_{pq} \delta_{kj} g_{npqq} \Gamma_{iumpql}^{[3]} + \sum_{pq} \delta_{kj} g_{npqq} \Gamma_{ipqmmn}^{[3]} + \sum_{pq} \delta_{kn} g_{upqn} \Gamma_{ipqmn}^{[3]} + \sum_{pq} \delta_{kn} g_{upqn} \Gamma_{ipqmn}^{[3]} + \sum_{pq} \delta_{kn} g_{upqn} \Gamma_{ipqmn}^{[3]} + \sum_{pq} \delta_{kn} g_{upqn} \Gamma_{ipqmq}^{[3]} + \sum_{pq} \delta_{kn} g_{upqn} \Gamma_{ipqmq}^{[3]} + \sum_{pq} \delta_{mn} g_{npqn} \Gamma_{ippqmq}^{[3]} + \sum_{pq} \delta_{in} g_{npqn} \Gamma_{ippqmq}^{[3]} + \sum_{pq} \delta_{in} g_{npqnq} \Gamma_{inpqknq}^{[3]} + \sum_{pq} \delta_{in} g_{npqnq} \Gamma_{ippqmq}^{[3]} + \sum_{pq} \delta_{in} g_{npqnq} \Gamma_{inpqknq}^{[3]} + \sum_{pq} \delta_{in} g_{npqqnq} \Gamma_{inpqknq}^{[3]} + \sum_{pq} \delta_{i$$

$$\mathcal{E} = \sum_{pq} \delta_{kj} \delta_{mu} g_{nipq} \Gamma_{tlpq}^{[2]} + \sum_{pq} \delta_{kj} \delta_{mu} g_{npqi} \Gamma_{tpql}^{[2]} + \sum_{pq} \delta_{kj} \delta_{mu} g_{ltpq} \Gamma_{inpq}^{[2]}$$
(S22)
+
$$\sum_{pq} \delta_{kj} \delta_{mu} g_{ptlq} \Gamma_{iqpn}^{[2]} + \sum_{pq} \delta_{kj} \delta_{tl} g_{pmuq} \Gamma_{iqpn}^{[2]} - \sum_{pq} \delta_{in} \delta_{kj} g_{ptqm} \Gamma_{puql}^{[2]}$$

-
$$\sum_{pq} \delta_{in} \delta_{ku} g_{ptqm} \Gamma_{plqj}^{[2]} - \sum_{pq} \delta_{in} \delta_{mu} g_{ptqk} \Gamma_{pjql}^{[2]} - \sum_{pq} \delta_{iu} \delta_{kj} g_{ptqm} \Gamma_{plqn}^{[2]}$$

-
$$\sum_{pq} \delta_{iu} \delta_{kn} g_{ptqm} \Gamma_{pjql}^{[2]} - \sum_{pq} \delta_{kj} \delta_{mu} g_{nplq} \Gamma_{tpiq}^{[2]} - \sum_{pq} \delta_{kj} \delta_{mu} g_{ptqi} \Gamma_{pnql}^{[2]}$$

-
$$\sum_{pq} \delta_{kj} \delta_{ml} g_{npuq} \Gamma_{tqip}^{[2]} - \sum_{pq} \delta_{kj} \delta_{tl} g_{upqm} \Gamma_{ipqn}^{[2]} - \sum_{pq} \delta_{kj} \delta_{tl} g_{npuq} \Gamma_{ipqn}^{[2]}$$

-
$$\sum_{pq} \delta_{kn} \delta_{mu} g_{ptqi} \Gamma_{plqj}^{[2]} - \sum_{pq} \delta_{mj} \delta_{tl} g_{npuq} \Gamma_{ipkq}^{[2]} - \sum_{pq} \delta_{mu} \delta_{tj} g_{nplq} \Gamma_{ipkq}^{[2]}$$

-
$$\sum_{pq} \delta_{mu} \delta_{tl} g_{npjq} \Gamma_{iqkp}^{[2]} - \sum_{pq} \delta_{ml} \delta_{tj} g_{npuq} \Gamma_{ipkq}^{[2]}$$

-
$$\sum_{pq} \delta_{mu} \delta_{tl} g_{npjq} \Gamma_{iqkp}^{[2]} - \sum_{pq} \delta_{ml} \delta_{tj} g_{npuq} \Gamma_{iqkp}^{[2]}$$

$$\mathcal{F} = \sum_{pqr} \delta_{in} g_{upqr} \Gamma_{tpqrmjkl}^{[4]} + \sum_{pqr} \delta_{iu} g_{npqr} \Gamma_{tjklmpqr}^{[4]} + \sum_{pqr} \delta_{kn} g_{upqr} \Gamma_{tpqrijml}^{[4]}$$
(S23)

$$+ \sum_{pqr} \delta_{ku} g_{npqr} \Gamma_{ijtlmpqr}^{[4]} + \sum_{pqr} \delta_{mj} g_{ptqr} \Gamma_{puqrinkl}^{[4]} + \sum_{pqr} \delta_{ml} g_{ptqr} \Gamma_{puqrijkn}^{[4]}$$
(S23)

$$+ \sum_{pqr} \delta_{tj} g_{pmqr} \Gamma_{iuklpnqr}^{[4]} + \sum_{pqr} \delta_{tl} g_{pmqr} \Gamma_{ijkupnqr}^{[4]} - \sum_{pqr} \delta_{in} g_{ptqr} \Gamma_{puqrmjkl}^{[4]}$$
(S23)

$$- \sum_{pqr} \delta_{in} g_{pmqr} \Gamma_{tupjqrkl}^{[4]} - \sum_{pqr} \delta_{il} g_{pmqr} \Gamma_{ijkupnqr}^{[4]} - \sum_{pqr} \delta_{iu} g_{ptqr} \Gamma_{pjqrklmn}^{[4]}$$
(S23)

$$- \sum_{pqr} \delta_{kn} g_{ptqr} \Gamma_{puqrijml}^{[4]} - \sum_{pqr} \delta_{iu} g_{pmqr} \Gamma_{ijklpnqr}^{[4]} - \sum_{pqr} \delta_{iu} g_{ptqr} \Gamma_{jjtlpnqr}^{[4]}$$
(S23)

$$- \sum_{pqr} \delta_{kn} g_{ptqr} \Gamma_{ijplqrmn}^{[4]} - \sum_{pqr} \delta_{kn} g_{pmqr} \Gamma_{tuipplqr}^{[4]} - \sum_{pqr} \delta_{ku} g_{pmqr} \Gamma_{ijtlpnqr}^{[4]}$$
(S23)

$$\mathcal{G} = \sum_{pqr} \delta_{in} \delta_{kj} g_{upqr} \Gamma_{tpqrml}^{[3]} + \sum_{pqr} \delta_{iu} \delta_{kj} g_{npqr} \Gamma_{tlmpqr}^{[3]} + \sum_{pqr} \delta_{ml} \delta_{kj} g_{ptqr} \Gamma_{puqrnn}^{[3]}$$
(S24)
+
$$\sum_{pqr} \delta_{tl} \delta_{kj} g_{pmqr} \Gamma_{iupnqr}^{[3]} - \sum_{pqr} \delta_{in} \delta_{kj} g_{ptqr} \Gamma_{puqrml}^{[3]} - \sum_{pqr} \delta_{in} \delta_{kj} g_{pmqr} \Gamma_{tuplqr}^{[3]}$$

-
$$\sum_{pqr} \delta_{in} \delta_{ku} g_{ptqr} \Gamma_{plqrmj}^{[3]} - \sum_{pqr} \delta_{in} \delta_{ku} g_{pmqr} \Gamma_{tlpjqr}^{[3]} - \sum_{pqr} \delta_{in} \delta_{mu} g_{ptqr} \Gamma_{plqrmn}^{[3]}$$

-
$$\sum_{pqr} \delta_{iu} \delta_{kn} g_{ptqr} \Gamma_{plqrmj}^{[3]} - \sum_{pqr} \delta_{iu} \delta_{kn} g_{pmqr} \Gamma_{tjplqr}^{[3]} - \sum_{pqr} \delta_{iu} \delta_{kj} g_{ptqr} \Gamma_{plqrmn}^{[3]}$$

-
$$\sum_{pqr} \delta_{iu} \delta_{kj} g_{pmqr} \Gamma_{tlpmqr}^{[3]} - \sum_{pqr} \delta_{kn} \delta_{mu} g_{ptqr} \Gamma_{plqrij}^{[3]} - \sum_{pqr} \delta_{iu} \delta_{kj} g_{ptqr} \Gamma_{plqrmn}^{[3]}$$

-
$$\sum_{pqr} \delta_{iu} \delta_{kj} g_{pmqr} \Gamma_{tlpmqr}^{[3]} - \sum_{pqr} \delta_{kn} \delta_{mu} g_{ptqr} \Gamma_{plqrij}^{[3]} - \sum_{pqr} \delta_{mj} \delta_{tl} g_{npqr} \Gamma_{ipkuqr}^{[3]}$$

-
$$\sum_{pqr} \delta_{ml} \delta_{tj} g_{npqr} \Gamma_{inkpqr}^{[3]} - \sum_{pqr} \delta_{ml} \delta_{kj} g_{upqr} \Gamma_{tpqrin}^{[3]} - \sum_{pqr} \delta_{ml} \delta_{kj} g_{npqr} \Gamma_{tuinqr}^{[3]}$$

-
$$\sum_{pqr} \delta_{ml} \delta_{tj} g_{npqr} \Gamma_{iukpqr}^{[3]} - \sum_{pqr} \delta_{ml} \delta_{tj} g_{upqr} \Gamma_{ipqrkn}^{[3]} - \sum_{pqr} \delta_{tj} \delta_{mu} g_{npqr} \Gamma_{ipklqr}^{[3]}$$

-
$$\sum_{pqr} \delta_{tl} \delta_{kj} g_{npqr} \Gamma_{iukpqr}^{[3]} - \sum_{pqr} \delta_{tl} \delta_{kj} g_{upqr} \Gamma_{ipqrkn}^{[3]} - \sum_{pqr} \delta_{tj} \delta_{mu} g_{npqr} \Gamma_{ipklqr}^{[3]}$$

$$\mathcal{H} = -\sum_{pqr} \delta_{in} \delta_{kj} \delta_{mu} g_{ptqr} \Gamma^{[2]}_{plqr} - \sum_{pqr} \delta_{tl} \delta_{kj} \delta_{mu} g_{npqr} \Gamma^{[2]}_{ipqr}$$
(S25)

$$\left[\hat{E}_{mn}, \hat{E}_{pq}\right] = \delta_{pn} \Gamma_{mq}^{[1]} - \delta_{mq} \Gamma_{pn}^{[1]}$$
(S26)

$$\left\langle 0 \left| \left[\hat{E}_{mn}, \hat{E}_{pq} \hat{E}_{rs} \right] \right| 0 \right\rangle$$

$$= \delta_{pn} \Gamma_{mqrs}^{[2]} + \delta_{pn} \delta_{qr} \Gamma_{ms}^{[1]} - \delta_{mq} \Gamma_{pnrs}^{[2]}$$

$$+ \delta_{rn} \Gamma_{pqms}^{[2]} - \delta_{ms} \Gamma_{pqrn}^{[2]} - \delta_{ms} \delta_{qr} \Gamma_{pn}^{[1]}$$

$$(S27)$$

$$\left\langle 0 \left| \left[\hat{E}_{tu} \hat{E}_{mn}, \hat{E}_{pq} \hat{E}_{rs} \right] \right| 0 \right\rangle$$

$$= -^{-} P_{qn}^{mp} \delta_{mq} \left(\Gamma_{tupnrs}^{[3]} + \delta_{nr} \left(\Gamma_{tups}^{[2]} + \delta_{up} \Gamma_{ts}^{[1]} \right) + \delta_{up} \Gamma_{tnrs}^{[2]} + \delta_{ur} \Gamma_{tspn}^{[2]} \right)$$

$$-^{-} P_{sn}^{mr} \delta_{ms} \left(\Gamma_{tupqrn}^{[3]} + \delta_{qr} \left(\Gamma_{tupn}^{[2]} + \delta_{up} \Gamma_{tn}^{[1]} \right) + \delta_{up} \Gamma_{tqrn}^{[2]} + \delta_{ur} \Gamma_{tnpq}^{[2]} \right)$$

$$-^{-} P_{qu}^{tp} \delta_{tq} \left(\Gamma_{pursmn}^{[3]} + \delta_{sm} \left(\Gamma_{purn}^{[2]} + \delta_{ur} \Gamma_{pn}^{[1]} \right) + \delta_{ur} \Gamma_{psmn}^{[2]} + \delta_{um} \Gamma_{pnrs}^{[2]} \right)$$

$$-^{-} P_{su}^{tr} \delta_{ts} \left(\Gamma_{pqrnn}^{[3]} + \delta_{um} \left(\Gamma_{pqrn}^{[2]} + \delta_{qr} \Gamma_{pn}^{[1]} \right) + \delta_{qr} \Gamma_{pumn}^{[2]} + \delta_{qm} \Gamma_{pnru}^{[2]} \right)$$

$$(S28)$$

S3 Geometries

Benzene:

С	-1.210	0.698	0.004
С	-1.210	-0.698	0.000
С	0.000	-1.397	-0.003
С	1.210	-0.698	-0.003
С	1.210	0.698	0.000
С	0.000	1.397	-0.003
Η	-2.164	1.249	0.007
Η	-2.164	-1.249	0.007
Η	0.000	-2.500	-0.005
H	2.164	-1.249	-0.006
Η	2.164	1.249	0.000
H	0.000	2.500	0.000

R-methyloxirane:

0	0.8171	-0.7825	-0.2447
С	-1.5018	0.1019	-0.1473
Η	-1.3989	0.3323	-1.2066
Η	-2.0652	-0.8262	-0.0524
Η	-2.0715	0.8983	0.3329
С	-0.1488	-0.0393	0.4879
Η	-0.1505	-0.2633	1.5506
С	1.0387	0.6105	-0.0580
Η	0.9518	1.2157	-0.9531
Н	1.8684	0.8649	0.5908

- Water:
- 0 0.0 0.0 0.1035174918
- Н 0.0 0.7955612117 -0.4640237459
- Н 0.0 -0.7955612117 -0.4640237459

S4 Code examples

```
from dmdm.interface import DMDM
dmdm = DMDM(
    int1e, int2e,
    num_inact_orb, num_act_orb, num_virt_orb, num_elec,
    rdm1, rdm2, rdm3, rdm4
)
excitation_energies = dmdm.get_excitation_energies()
oscillator_strengths = dmdm.get_oscillator_strength(elec_dip_int)
```

Figure S1: Excitation energies and oscillator strengths calculation. intle are the oneelectron integrals from the Hamiltonian in MO. int2e are the two-electron integrals from the Hamiltonian in MO; num_inact_orb is the number of inactive spatial orbitals; num_act_orb is the number of active spatial orbitals; num_virt_orb is the number of virtual spatial orbitals;num_elec is the number of electrons. rdm1, rdm2, rdm3 and rdm4 are the n-RDMs in only the active space. elec_dip_int are the electric dipole integrals in MO (x, y, zcomponents).

```
from dmdm.interface import DMDM
dmdm = DMDM(
    int1e, int2e,
    num_inact_orb, num_act_orb, num_virt_orb, num_elec,
    rdm1, rdm2, rdm3, rdm4
)
excitation_energies = dmdm.get_excitation_energies()
rotational_strengths = dmdm.get_rotational_strength(elec_dip_int,
    mag_dip_int)
```

Figure S2: Excitation energies and rotational strengths calculation. intle are the oneelectron integrals from the Hamiltonian in MO. int2e are the two-electron integrals from the Hamiltonian in MO. num_inact_orb is the number of inactive spatial orbitals. num_act_orb is the number of active spatial orbitals. num_virt_orb is the number of virtual spatial orbitals. num_elec is the number of electrons. rdm1, rdm2, rdm3, and rdm4 are the n-RDMs in only the active space. elec_dip_int are the electric dipole integrals in MO (x, y, z); mag_dip_int are the magnetic dipole integrals in MO (x, y, z)

S5 Tables and figures



Figure S3: *R*-methyloxirane. Absorption spectra of the first 399 excitation energies using CASSCF(6,6) LR (blue), and qLRSD (dashed orange) based on the oo-UCCSD(6,6) wave function using the cc-pVTZ basis set.



Figure S4: R-methyloxirane. ECD spectra of the first 399 excitation energies using CASSCF(6,6) LR (blue), and qLRSD (dashed orange) based on the oo-UCCSD(6,6) wave function using the cc-pVTZ basis set.

			1				1	1																	
	shots	Osc.	0.0003	0.0000	0.0003	0.0012	0.0018	0.0016	0.0016	0.0006	0.0001	0.0000	0.0002	0.0001	0.0000	0.0000	0.0000	0.0000	0.0000	0.0032	0.0084	0.0063	0.0005	0.0000	0.0000
г	100k	Exc.	0.0287	0.0290	0.0134	0.0147	0.0430	0.0994	0.0787	0.0362	0.0248	0.0171	0.0226	0.0200	0.0276	0.0217	0.0211	0.0191	0.0228	0.5420	0.1610	0.5621	1.7710	0.0185	0.0185
deviation	hots	Osc.	0.0008	0.0002	0.0005	0.0017	0.0020	0.0027	0.0034	0.0008	0.0002	0.0000	0.0002	0.0004	0.0000	0.0000	0.0000	0.0000	0.0040	0.0038	0.0081	0.0079	0.0013	0.0000	0.0000
tandard	50k s	Exc.	0.0915	0.0529	0.0248	0.0307	0.0708	0.1813	0.2287	0.0658	0.0800	0.0336	0.0444	0.0435	0.0560	0.0425	0.0415	0.0375	0.1488	1.8971	0.3002	1.0487	1.8779	0.0365	0.0364
\mathbf{v}	$_{\rm shots}$	Osc.	0.0034	0.0018	0.0008	0.0036	0.1076	0.0106	0.0124	0.0017	0.0005	0.0001	0.0005	0.0009	0.0001	0.0000	0.0000	0.0003	0.0068	0.0063	0.0106	0.0106	0.0032	0.0000	0.0019
	10k s	Exc.	0.4007	0.1086	0.0657	0.0734	0.2072	0.4237	0.3549	0.0970	0.1678	0.0632	0.0819	0.0833	0.0891	0.0813	0.0789	0.0721	1.6102	2.5327	0.7460	0.7943	4.1260	0.5629	0.6480
	100k shots	Osc.	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0000	0.0000	0.0000	0.0000
		Exc.	0.0008	0.0008	0.0002	0.0002	0.0019	0.0099	0.0062	0.0013	0.0006	0.0003	0.0005	0.0004	0.0008	0.0005	0.0004	0.0004	0.0005	0.2938	0.0259	0.3160	3.1363	0.0003	0.0003
nce	hots	Osc.	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0001	0.0000	0.0000	0.0000
Varia	50k s	Exc.	0.0084	0.0028	0.0006	0.0009	0.0050	0.0329	0.0523	0.0043	0.0064	0.0011	0.0020	0.0019	0.0031	0.0018	0.0017	0.0014	0.0222	3.5990	0.0901	1.0999	3.5265	0.0013	0.0013
	lots	Osc.	0.0000	0.0000	0.0000	0.0000	0.0116	0.0001	0.0002	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0001	0.0001	0.0000	0.0000	0.0000
	10k s	Exc.	0.1605	0.0118	0.0043	0.0054	0.0429	0.1795	0.1260	0.0094	0.0282	0.0040	0.0067	0.0069	0.0079	0.0066	0.0062	0.0052	2.5928	6.4146	0.5565	0.6310	17.0240	0.3169	0.4200
		Osc.	0.0319	0.0000	0.1133	0.0976	0.2517	0.0427	0.0735	0.0303	0.0014	0.0779	0.0094	0.0236	0.0001	0.0000	0.0000	0.0021	0.0003	0.0135	0.0267	0.0007	0.0061	0.0000	0.0000
	Idea	Exc.	8.5433	10.4364	11.1083	12.8695	14.8841	546.6519	547.0419	560.3159	561.5096	561.7025	563.5124	567.0508	569.4280	570.4970	573.9153	574.3891	580.8528	586.0916	586.3814	587.7218	588.2594	596.3517	597.7355
	Variance Standard deviation	IdealVarianceVarianceStandard deviationIdeal $10k \text{ shots}$ $50k \text{ shots}$ $100k \text{ shots}$ $50k \text{ shots}$ $100k \text{ shots}$	$\begin{tabular}{c c c c c c c c c c c c c c c c c c c $	$ \begin{array}{c c c c c c c c c c c c c c c c c c c $					$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	$ \begin{array}{l lllllllllllllllllllllllllllllllllll$	$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	$ \begin{array}{c c c c c c c c c c c c c c c c c c c $	$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	$ \begin{array}{l lllllllllllllllllllllllllllllllllll$	$ \begin{array}{l lllllllllllllllllllllllllllllllllll$	$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	Ideal Variance Variance Standard deviation Ideal 10k shots 50k shots 100 0000 0.000 0.000 0.000 0.000 0.000 0.000 0.000 0.000 0.000 0.000 0.000 0.000 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.0011 0.001 0.001		Variance Standard deviation Ideal In Nationa Jack shots Jok shots Standard deviation Jok Jok shots Jok shots Jok shots Jok shots Exc. Osc. Exc. Osc. Jok Jok shots Jok shots S-433 0.0000 Jok shots Jok Jok Jok S-433 Johds Johds Johds Johds Johds S-433 Johds Johds Johds Johds Johds Johds Johds Johds Johds Johds Johds Johds Johds Johds Johds Johds Johds

Table S1: Ideal excitation energies and oscillator energies shown in Fig. 3 with the variance and standard deviation of the 10