# On the global dynamics of a forest model with monotone positive feedback and memory

Franco Herrera<sup>a</sup>, Sergei Trofimchuk<sup>a,\*</sup>

<sup>a</sup>Instituto de Matemáticas, Universidad de Talca, Casilla 747, Talca, Chile

# Abstract

We continue to study (see [6]) a renewal equation  $\phi(t) = \mathfrak{F}\phi_t$  proposed in [1, 2] to model trees growth. This time we are considering the case when the per capita reproduction rate  $\beta(x)$  is a non-monotone (unimodal) function of tree's height x. Note that the height of some species of trees can impact negatively seed viability, cf. [3, p. 524], in a kind of autogamy depression. Similarly to previous works, it is also assumed that the growth rate g(x) of an individual of height x is a strictly decreasing function. As in [5, 6], we analyse the connection between dynamics of the associated one-dimensional map  $F(b) = \mathfrak{F}b, \ b \in \mathbb{R}_+$ , and the delayed (hence infinite-dimensional) model  $\phi(t) = \mathfrak{F}\phi_t$ . Our key observation is that this model is of monotone positive feedback type since F is strictly increasing on  $\mathbb{R}_+$  independently on the monotonicity properties of  $\beta$ .

Keywords: size-structured model, semiflow, global attractor, stability, Allee effect

Preprint submitted to MATRIX Annals

*e-mtails addresses*: franco.herrera@utalca.cl (F. Herrera); trofimch@inst-mat.utalca.cl (S. Trofimchuk, corresponding author)

#### 1. Introduction

In this note, we consider the renewal equation

$$b(t) = \mathfrak{F}b_t := \int_0^\infty \beta \left( x_m + \int_0^a g \left( e^{-\mu(\tau-a)} \int_a^\infty e^{-\mu s} b(t-s) \, ds \right) d\tau \right) e^{-\mu a} b(t-a) da, \quad (1)$$

where  $b_t(s) := b(t+s)$ ,  $s \leq 0$ , and functional  $\mathfrak{F}$  is given by

$$\mathfrak{F}\phi := \int_0^\infty \beta\left(x_m + \int_0^a g\left(e^{-\mu(\tau-a)}\int_a^\infty e^{-\mu s}\phi(-s)\,ds\right)d\tau\right)e^{-\mu a}\phi(-a)da,$$

This equation was proposed recently in [1, 2] to model trees growth and presents one of possible approaches to analyse the forest population dynamics, several other models are mentioned in [1, 2, 6]. In equation (1), function b(t) corresponds to the tree population birth rate at time t; continuous strictly decreasing function  $g(x), g: [0, +\infty) \to (0, +\infty), g(+\infty) = 0$ , is the growth rate of an individual of height  $x; \mu > 0$  represents the per capita death rate; globally Lipschitzian function  $\beta(x), \beta: [x_m, +\infty) \to [0, +\infty)$ , is the per capita reproduction rate (depending only on height x). This model postulates the minimal height  $x_m \ge 0$  for all newborn trees. By [1, 2], b(t) gives the influx at  $x_m$  originating from reproduction by the extant population. To ensure good persistence properties of solutions, it suffices to assume that  $\beta(x) > 0$  a.e. on  $[x_0, +\infty)$  for some  $x_0 \ge x_m$  [6].

Clearly, (1) can be considered as an autonomous functional equation with infinite delay once it is provided with non-negative initial conditions  $b(s) = \phi(s) \ge 0$ ,  $s \le 0$ , where  $\phi$  belongs to the cone  $\mathfrak{K}$  of non-negative measurable functions of the Banach space

$$L^1_\rho(\mathbb{R}_-) = \{\phi: |\phi|_\rho < \infty\}, \quad (\text{here we use the notation} \quad |\phi|_q := \int_{-\infty}^0 |\phi(s)| e^{qs} ds),$$

with  $\rho < \mu$ . It was proved in [1, 2, 6] that the renewal equation (1) generates a continuous semiflow  $\mathfrak{S} : \mathfrak{K} \times \mathbb{R}_+ \to \mathfrak{K}$  by  $(\mathfrak{S}^t \phi)(s) = b(t+s), s \leq 0$ . It has a zero steady state  $b(t) \equiv 0$ while the positive equilibria of  $\mathfrak{S}^t$  are defined from the equation

$$1 = R(b) := \int_0^\infty \beta\left(x_m + \int_0^a g\left(\frac{e^{-\mu\tau}}{\mu}b\right)d\tau\right) e^{-\mu a}da.$$
 (2)

Note that equation (2) has at least one positive solution  $b_*$  if R(0) > 1 and  $R(+\infty) = \beta(x_m)/\mu < 1$ . If additionally  $\beta$  increases on  $\mathbb{R}_+$ , then R(b) is strictly decreasing so that solution  $b_*$  is unique, cf. [1, Theorem 4.1]. Moreover, this unique positive steady state is locally stable and globally attracting solution of (1), see [1, 2, 6]. The key argument in proofs here is that semiflow  $\mathfrak{F}^t \phi$  is monotone [12] once  $\beta$  is increasing on  $\mathbb{R}_+$ .

In some cases, however, the assumption of monotonicity of  $\beta$  is not realistic. For example, the height of some species of trees can impact negatively seed viability, cf. [3, p. 524], in a kind of autogamy depression. Thus it would be interesting to study the dynamics in (1) in the case when the per capita reproduction rate  $\beta(x)$  is a nonmonotone (for instance, unimodal) function of tree's height x. Now, a general description of this dynamics is known from [6], where it was proven that under all assumed (except monotonicity of  $\beta$ ) restrictions on  $\beta$ , g and  $0 \leq \beta(x_m) < \mu$ , R(0) > 1, the semiflow  $\mathfrak{F}^t \phi$  has a compact global attractor  $\mathcal{A}$  attracting each non-zero solution. Moreover, each element  $\psi$  of  $\mathcal{A}$  satisfies  $0 < \theta_1 \leq \psi(s) \leq \theta_2$ ,  $|\psi'(s)| \leq \theta_3$ ,  $s \leq 0$ , for some universal positive numbers  $\theta_1 \leq \theta_2, \theta_3$ .

In the sequel we present several analytical and numerical arguments shedding some light on possible dynamical structures of the global attractor  $\mathcal{A}$  without assumption of the monotonicity for  $\beta$ . As in [5, 6], we will analyse the connection between dynamics of the associated one-dimensional map  $F(b) = \mathfrak{F}b, \ b \in \mathbb{R}_+$ , and the delayed (hence infinite-dimensional) model  $\phi(t) = \mathfrak{F}\phi_t$ .

## 2. Monotonicity of the function $F(b) = bR(b), b \in \mathbb{R}_+$ .

**Theorem 1.** Assume all conditions (except monotonicity of  $\beta$ ) imposed on  $\beta$  and g in Section 1. Then function F(b) = bR(b) is strictly increasing on  $\mathbb{R}_+$ .

PROOF. Assuming that  $\phi(s) > 0$  a.e. on  $\mathbb{R}_{-}$  and setting  $h(s) = \phi(s)e^{\mu s}$  and  $\theta = H(u) = \int_{-\infty}^{u} h(s)ds$ , we find that

$$\mathfrak{F}\phi = \int_{-\infty}^{0} \beta \left( x_m + \int_{u}^{0} g(e^{-\mu s} H(u)) ds \right) h(u) du = \int_{0}^{H(0)} \beta \left( x_m + \int_{H^{-1}(\theta)}^{0} g(e^{-\mu s} \theta) ds \right) d\theta$$

Clearly, H(u) is strictly increasing and positive on  $\mathbb{R}$ , with  $H(-\infty) = 0$ ,

$$\frac{d}{d\theta}H^{-1}(\theta) = \frac{1}{h(H^{-1}(\theta))} \quad \text{a.e. on } (0, H(0)).$$

If  $\phi_2(s) > \phi_1(s) > 0$ , almost everywhere on  $\mathbb{R}_-$ , then

$$H_2(u) = \int_{-\infty}^{u} h_2(s) ds > H_1(u) = \int_{-\infty}^{u} h_1(s) ds, \quad u \le 0,$$

so that

$$H_2^{-1}(\theta) < H_1^{-1}(\theta) \le 0, \quad 0 < \theta \le H_1(0)$$

Since g is strictly decreasing, we find that the function

$$\gamma(\theta,\phi) := x_m + \int_{H^{-1}(\theta)}^0 g(e^{-\mu s}\theta) ds = x_m + \int_{H^{-1}(\theta)+\mu^{-1}\ln(1/\theta)}^{\mu^{-1}\ln(1/\theta)} g(e^{-\mu s}) ds,$$

is continuous and strictly decreasing in  $\theta$  for each fixed  $\phi > 0$ ,

$$\partial \gamma(\theta,\phi)/\partial \theta = -\frac{1}{\mu} \left( \frac{g(\theta)}{\theta} - g(\theta e^{-\mu H^{-1}(\theta)}) \left[ \frac{1}{\theta} - \frac{\mu}{h(H^{-1}(\theta))} \right] \right) < 0.$$

Also, for a fixed  $\theta$  and  $\phi_2(s) > \phi_1(s) > 0$ , it holds that

$$\gamma(\theta, \phi_2) > \gamma(\theta, \phi_1).$$

Thus the equation  $u = \gamma(\theta, \phi)$  can be solved with respect to  $\theta$ , we denote this solution by  $\theta = \sigma(u, \phi)$ . Clearly,  $\sigma$  is strictly decreasing in the first argument and is strictly increasing in  $\phi$ . Therefore  $g(\sigma(u, \phi))/\sigma(u, \phi)$  is strictly decreasing in  $\phi$  In this way,

$$\mathfrak{F}\phi = \mu \int_{x_m}^{+\infty} \beta(u) \left(\frac{g(\theta)}{\theta} - g(\theta e^{-\mu H^{-1}(\theta)}) \left[\frac{1}{\theta} - \frac{\mu}{h(H^{-1}(\theta))}\right]\right)^{-1} du, \text{ where } \theta = \sigma(u,\phi).$$

Particularly,

$$F(b) = \mathfrak{F}\phi = \mu \int_{x_m}^{+\infty} \beta(u) \frac{\theta}{g(\theta)} du, \text{ where } \theta = \sigma(u, b).$$

so that F(b) is a strictly increasing function independently on the monotonicity properties of  $\beta$ .

Theorem 1 suggests to explore a possible connection between dynamical structures for the delayed equation (1) and seemingly simpler equation

$$b(t) = \mathfrak{G}b_t := \int_0^\infty f(b(t-a-h)) e^{-\mu a} da,$$
 (3)

where  $F = \mu^{-1} f : \mathbb{R} \to \mathbb{R}$  is a strictly increasing smooth function, f'(x) > 0 for all  $x \in \mathbb{R}$ , having three equilibria  $e_1 < e_2 := 0 < e_3$ ,  $F(e_j) = e_j$ . Equation (3) describes bounded solutions from the global attractor for the semiflow generated by the delayed equation

$$b'(t) = -\mu b(t) + f(b(t-h)), \quad \mu > 0.$$

Specific applications of such a model can be found in neural network theory [7, 8]. Obviously, each equilibrium for (3) is determined from the equation

$$b = \mathfrak{G}b = \mu^{-1}f(b),$$

where the right hand side is given by a strictly increasing function. The global attractor  $\mathcal{A}$  for (3) has relatively simple structure described in [8] (if  $e_2$  is "sufficiently unstable", then  $\mathcal{A}$  is a smooth solid 3-dimensional spindle separated by the invariant disk into the basins of attraction toward the tips  $e_1$  and  $e_3$ ). See also [9] for further results and references on this topic.

#### 3. Characteristic equations and principle of linearized stability

Formal linearization of the operator  $\mathfrak{F}$  at an equilibrium  $b(t) \equiv b$  yields the following action

$$\mathfrak{F}'(b)\psi = \int_0^\infty \beta\left(x_m + \int_0^a g\left(\frac{e^{-\mu\tau}b}{\mu}\right)d\tau\right) e^{-\mu a}\psi(-a)da + \\ \int_0^\infty \beta'\left(x_m + \int_0^a g\left(\frac{e^{-\mu\tau}b}{\mu}\right)d\tau\right) \left[g\left(\frac{b}{\mu}\right) - g\left(\frac{e^{-\mu a}b}{\mu}\right)\right] \int_a^\infty e^{-\mu s}\psi(-s)\,dsda = \\ \beta(x_m)\int_0^\infty e^{-\mu s}\psi(-s)\,ds + g\left(\frac{b}{\mu}\right)\int_0^\infty \beta'\left(x_m + \int_0^a g\left(\frac{e^{-\mu\tau}b}{\mu}\right)d\tau\right) \int_a^\infty e^{-\mu s}\psi(-s)\,dsda = \\ \mathbf{F}(x_m)\int_0^\infty e^{-\mu s}\psi(-s)\,ds + g\left(\frac{b}{\mu}\right)\int_0^\infty \beta'\left(x_m + \int_0^a g\left(\frac{e^{-\mu\tau}b}{\mu}\right)d\tau\right)\int_a^\infty e^{-\mu s}\psi(-s)\,dsda = \\ \mathbf{F}(x_m)\int_0^\infty e^{-\mu s}\psi(-s)\,ds + g\left(\frac{b}{\mu}\right)\int_0^\infty \beta'\left(x_m + \int_0^a g\left(\frac{e^{-\mu\tau}b}{\mu}\right)d\tau\right)\int_a^\infty e^{-\mu s}\psi(-s)\,dsda = \\ \mathbf{F}(x_m)\int_0^\infty e^{-\mu s}\psi(-s)\,ds + g\left(\frac{b}{\mu}\right)\int_0^\infty \beta'\left(x_m + \int_0^a g\left(\frac{e^{-\mu\tau}b}{\mu}\right)d\tau\right)\int_a^\infty e^{-\mu s}\psi(-s)\,dsda = \\ \mathbf{F}(x_m)\int_0^\infty e^{-\mu s}\psi(-s)\,ds + g\left(\frac{b}{\mu}\right)\int_0^\infty \beta'\left(x_m + \int_0^a g\left(\frac{e^{-\mu\tau}b}{\mu}\right)d\tau\right)\int_a^\infty e^{-\mu s}\psi(-s)\,dsda = \\ \mathbf{F}(x_m)\int_0^\infty e^{-\mu s}\psi(-s)\,ds + g\left(\frac{b}{\mu}\right)\int_0^\infty \beta'\left(x_m + \int_0^a g\left(\frac{e^{-\mu\tau}b}{\mu}\right)d\tau\right)\int_a^\infty e^{-\mu s}\psi(-s)\,dsda = \\ \mathbf{F}(x_m)\int_0^\infty e^{-\mu s}\psi(-s)\,ds + g\left(\frac{b}{\mu}\right)\int_0^\infty \beta'\left(x_m + \int_0^a g\left(\frac{e^{-\mu\tau}b}{\mu}\right)d\tau\right)\int_a^\infty e^{-\mu s}\psi(-s)\,dsda = \\ \mathbf{F}(x_m)\int_0^\infty e^{-\mu s}\psi(-s)\,ds + g\left(\frac{b}{\mu}\right)\int_0^\infty \beta'\left(x_m + \int_0^a g\left(\frac{e^{-\mu\tau}b}{\mu}\right)d\tau\right)\int_a^\infty e^{-\mu s}\psi(-s)\,dsda = \\ \mathbf{F}(x_m)\int_0^\infty e^{-\mu s}\psi(-s)\,ds + g\left(\frac{b}{\mu}\right)\int_0^\infty \beta'\left(x_m + \int_0^a g\left(\frac{e^{-\mu\tau}b}{\mu}\right)d\tau\right)\int_a^\infty e^{-\mu s}\psi(-s)\,dsda = \\ \mathbf{F}(x_m)\int_0^\infty e^{-\mu s}\psi(-s)\,ds + g\left(\frac{b}{\mu}\right)\int_0^\infty \beta'\left(x_m + \int_0^a g\left(\frac{e^{-\mu\tau}b}{\mu}\right)d\tau\right)\int_a^\infty e^{-\mu s}\psi(-s)\,dsda = \\ \mathbf{F}(x_m)\int_0^\infty e^{-\mu s}\psi(-s)\,ds + g\left(\frac{b}{\mu}\right)\int_0^\infty e^{-\mu s}\psi($$

Looking for the exponential eigenfunctions  $\psi(s) = \exp(\lambda s)$  of the linearized equation  $\psi(t) = \mathfrak{F}'(b)\psi_t$ , we obtain the characteristic equation for  $\lambda$ , cf. [1, 2]:

$$\lambda + (\mu - \beta(x_m)) = g\left(\frac{b}{\mu}\right) \int_0^\infty \beta' \left(x_m + \int_0^a g\left(\frac{e^{-\mu\tau}b}{\mu}\right) d\tau\right) e^{-(\mu+\lambda)a} da =: \chi_b(\lambda).$$

First, we consider the simpler case when  $\beta$  is a non-decreasing function and  $\mu > \beta(x_m)$  (so that  $F'(b) \leq 1$  at the unique positive equilibrium):

**Lemma 2.** Assume that  $\beta'(x_m + s) \ge 0$ , g(s) > 0, for  $s \ge 0$  and  $\mu - \beta(x_m) > 0$ . Then the characteristic equation has exactly one real solution  $\lambda = \lambda_0$ . Moreover,  $\lambda_0 \le 0$  and  $\Re \lambda_j < \lambda_0$  for each other (complex) solution  $\lambda_j$ . If g is strictly decreasing on  $\mathbb{R}_+$ , then  $\lambda_0 < 0$ .

PROOF. The proof is obvious if  $\beta'(s) = 0$  a.e. so that we consider the case when  $\beta' \neq 0$ . Clearly each eigenvalues satisfies the inequality  $\lambda_j + (\mu - \beta(x_m)) \geq 0$ . Taking into account opposite types of monotonicity of  $\lambda + (\mu - \beta(x_m))$  and  $\chi_b(\lambda)$  and the fact that  $\chi_b(\lambda) > 0$ and  $\chi_b(0) = \mu F'(b) - \beta(x_m) \leq \mu - \beta(x_m)$ , we deduce the existence of a unique real (non-positive) solution  $\lambda_0$ .

Assuming that there is another eigenvalue  $\lambda_j$  with  $\Re \lambda_j \geq \lambda_0$ , we get a contradiction:

$$\lambda_{0} + (\mu - \beta(x_{m})) = |\lambda_{0} + (\mu - \beta(x_{m}))| < |\lambda_{j} + (\mu - \beta(x_{m}))| \le$$

$$g\left(\frac{b}{\mu}\right) \int_{0}^{\infty} \beta'\left(x_{m} + \int_{0}^{a} g\left(\frac{e^{-\mu\tau}b}{\mu}\right) d\tau\right) e^{-(\mu + \Re\lambda_{j})a} da \le$$

$$g\left(\frac{b}{\mu}\right) \int_{0}^{\infty} \beta'\left(x_{m} + \int_{0}^{a} g\left(\frac{e^{-\mu\tau}b}{\mu}\right) d\tau\right) e^{-(\mu + \lambda_{0}a)} da = \lambda_{0} + (\mu - \beta(x_{m}))$$

Finally, suppose that g is a strictly decreasing and  $\lambda_0 = 0$ . Then we get the following contradiction:

$$\mu - \beta(x_m) = \int_0^\infty \beta' \left( x_m + \int_0^a g\left(\frac{e^{-\mu\tau}b}{\mu}\right) d\tau \right) g\left(\frac{b}{\mu}\right) e^{-\mu a} da < \int_0^\infty \beta' \left( x_m + \int_0^a g\left(\frac{e^{-\mu\tau}b}{\mu}\right) d\tau \right) g\left(\frac{e^{-\mu a}b}{\mu}\right) e^{-\mu a} da = -\beta(x_m) + \mu \int_0^\infty \beta \left( x_m + \int_0^a g\left(\frac{e^{-\mu\tau}b}{\mu}\right) d\tau \right) e^{-\mu a} da = -\beta(x_m) + \mu. \quad \Box$$

**Remark 3.** Lemma 2 simplifies and improves arguments in [1, Section 5] (apparently, the uniqueness statement of this lemma contradicts to the existence of two negative eigenvalues in the particular case presented in [1, Subsection 5.1]; however, one of these eigenvalues is false and should be excluded to allow the convergence of integrals in  $\chi_b(\lambda)$ .

Next, we analyse the situation when  $\beta$  is not necessarily monotone function and  $\mu > \beta(x_m)$  (we assume that the model possesses at least one positive equilibrium b):

**Lemma 4.** Assume that g > 0 is strictly decreasing Lipschitz continuous function on  $\mathbb{R}_+$ ,  $\mu > \beta(x_m)$ . Then in the half-plane  $\{\Re z \ge -\mu\}$  the characteristic equation has exactly one real solution  $\lambda = \lambda_0$  and  $\Re \lambda_j < \lambda_0$  for each other (complex) solution  $\lambda_j$ . Moreover, if F'(b) < 1 then  $\lambda_0 < 0$ ,  $\lambda_0 = 0$  when F'(b) = 1 and  $\lambda_0 > 0$  if F'(b) > 1.

**PROOF.** Note that

$$\begin{split} \lambda + (\mu - \beta(x_m)) &= g\left(\frac{b}{\mu}\right) \int_0^\infty \beta' \left(x_m + \int_0^a g\left(\frac{e^{-\mu\tau}b}{\mu}\right) d\tau\right) e^{-(\mu+\lambda)a} da = \\ g\left(\frac{b}{\mu}\right) \int_0^\infty \frac{e^{-(\mu+\lambda)a}}{g\left(\frac{e^{-\mu ab}}{\mu}\right)} d\beta \left(x_m + \int_0^a g\left(\frac{e^{-\mu\tau}b}{\mu}\right) d\tau\right) = \\ -\beta(x_m) - g\left(\frac{b}{\mu}\right) \int_0^\infty \beta \left(x_m + \int_0^a g\left(\frac{e^{-\mu\tau}b}{\mu}\right) d\tau\right) \left[\frac{e^{-(\mu+\lambda)a}}{g\left(\frac{e^{-\mu ab}}{\mu}\right)}\right]' da. \end{split}$$

Since

$$\left[\frac{e^{-(\mu+\lambda)a}}{g\left(\frac{e^{-\mu a}b}{\mu}\right)}\right]' = e^{-(\mu+\lambda)a}\frac{-(\mu+\lambda)g\left(\frac{e^{-\mu a}b}{\mu}\right) + e^{-\mu a}bg'\left(\frac{e^{-\mu a}b}{\mu}\right)}{\left(g\left(\frac{e^{-\mu a}b}{\mu}\right)\right)^2} < 0,$$

we obtain the following equivalent form of the characteristic equation:

$$1 = g\left(\frac{b}{\mu}\right) \int_0^\infty \beta\left(x_m + \int_0^a g\left(\frac{e^{-\mu\tau}b}{\mu}\right) d\tau\right) \frac{e^{-(\mu+\lambda)a}}{g\left(\frac{e^{-\mu a}b}{\mu}\right)} \left[1 - \frac{e^{-\mu a}bg'\left(\frac{e^{-\mu a}b}{\mu}\right)}{(\mu+\lambda)g\left(\frac{e^{-\mu a}b}{\mu}\right)}\right] da.$$

The right side of this equation (we will denote it by  $\xi_b(\lambda)$ , clearly  $\xi_b(\lambda) = (\chi_b(\lambda) + \beta(x_m))/(\mu + \lambda)$  and  $\xi_b(0) = (\chi_b(0) + \beta(x_m))/\mu = F'(b))$  is a strictly decreasing function of  $\lambda$  on the interval  $(-\mu, +\infty)$ ,  $\xi_b(-\mu^+) = +\infty$ ,  $\xi_b(+\infty) = 0$ , so that the equation has exactly one real root  $\lambda_0$  on this interval. Furthermore, we see that  $\lambda_0$  is a dominating eigenvalue. Indeed if  $\lambda_j$  is another eigenvalue with  $\Re \lambda_j \geq \lambda_0$  then  $\lambda_0 + \mu < |\lambda_j + \mu|$  and we get a contradiction:

$$1 \leq g\left(\frac{b}{\mu}\right) \int_0^\infty \beta\left(x_m + \int_0^a g\left(\frac{e^{-\mu\tau}b}{\mu}\right) d\tau\right) \frac{e^{-(\mu+\Re\lambda_j)a}}{g\left(\frac{e^{-\mu a}b}{\mu}\right)} \left[1 + \frac{e^{-\mu a}b|g'\left(\frac{e^{-\mu a}b}{\mu}\right)|}{|\mu + \lambda_j|g\left(\frac{e^{-\mu a}b}{\mu}\right)|}\right] da < g\left(\frac{b}{\mu}\right) \int_0^\infty \beta\left(x_m + \int_0^a g\left(\frac{e^{-\mu\tau}b}{\mu}\right) d\tau\right) \frac{e^{-(\mu+\lambda_0)a}}{g\left(\frac{e^{-\mu a}b}{\mu}\right)} \left[1 + \frac{e^{-\mu a}b|g'\left(\frac{e^{-\mu a}b}{\mu}\right)|}{|\mu + \lambda_0|g\left(\frac{e^{-\mu a}b}{\mu}\right)|}\right] da = 1$$

Finally, since  $\xi_b(0) = F'(b)$  we conclude that  $\lambda_0 < 0$  when F'(b) < 1,  $\lambda_0 = 0$  when F'(b) = 1 and  $\lambda_0 > 0$  when F'(b) > 1.

Lemma 4 implies that, except for the critical case F'(b) = 1, the local stability/instability of an equilibrium b in (1) is equivalent to the local stability/instability of the same equilibrium for one-dimensional system  $b \to F(b)$ :

**Corollary 5.** Assume that functions  $g : \mathbb{R} \to \mathbb{R}$ ,  $\beta : \mathbb{R}_+ \to \mathbb{R}$  have a bounded and globally Lipschitzian first derivative and  $\rho < \mu/5$ . Then an equilibrium b of continuous semiflow  $\mathfrak{S} : \mathfrak{K} \times \mathbb{R}_+ \to \mathfrak{K}$  is locally asymptotically stable if F'(b) < 1 and is unstable when F'(b) > 1.

PROOF. As it was established in [1, Theorem 3.1], under additional assumptions of this corollary, the functional  $\mathfrak{F} : L^1_{\rho}(\mathbb{R}_-) \to \mathbb{R}$  is continuously differentiable with bounded derivative. Then an application of [1, Theorem 3.3] (based on Theorem 3.15 in [4]) completes the proof.

**Remark 6.** Note that the inequalities F'(b) < 1 and F'(b) > 1 are equivalent, respectively, to the inequalities R'(b) < 0 and R'(b) > 0, where

$$R'(b) = -\frac{1}{b} - \frac{1}{\mu b} g\left(\frac{b}{\mu}\right) \int_0^\infty \beta\left(x_m + \int_0^a g\left(\frac{e^{-\mu\tau}b}{\mu}\right) d\tau\right) \left[\frac{e^{-\mu a}}{g\left(\frac{e^{-\mu a}b}{\mu}\right)}\right] da.$$

## 4. Numerical simulations and open problems

Stability analysis and discussion presented in the previous sections suggest the existence of a 'large' (we believe, open dense) set  $\mathcal{O}$  in the phase space  $\mathfrak{K}$  such that solutions with initial data from  $\mathcal{O}$  converge to one of stable equilibria. Thus we can expect that a typical numerical solution of equation (1) is asymptotically constant independently on the monotonicity properties of  $\beta$ . To check this conclusion and to find out possible configurations for the set of equilibria for (1) with the unimodal fertility function  $\beta$ , we took  $\beta(x) = \alpha x e^{-x}$  (which is used in the Nicholson blowflies equation and other models, cf. [10]);  $x_m = 0$ ,  $\mu = 1$ , and  $g(x) = p e^{-x}$ , for some specific parameters  $\alpha, p > 0$ . With these choices, our computations exhibit certain diversity in dynamical patterns for (1).



Figure 1: Graphics of y = F(b) and y = b for different pairs of parameters  $\alpha$  and p

First, in full accordance with Theorem 1, Figure 1 shows that the function F is strictly increasing even in lack of monotonicity for  $\beta$ . Nevertheless, the function R is not necessarily monotone and we can observe that for different values of p, R can be a decreasing function as well as a unimodal one, see Figure 2. It is important to mention that these shapes for R do not change essentially even if instead of  $g(x) = pe^{-x}$  we take g with other orders of decay (e.g. polynomial) at  $+\infty$ .



Figure 2: Graphics of R for different pairs of parameters  $\alpha$  and p



Figure 3: Simulations of b(t) using constant initial data  $\phi$ 



Figure 4: Simulations of b(t) using periodic initial data of the form  $\phi(a) = b_* + \varepsilon \sin(\omega a)$ 

Interestingly, regardless the fact R(0) < 1, equation (1) with  $\alpha = 6$  and p = 5 has three equilibria,  $b_1 = 0 < b_2 \approx 0.47 < b_3 \approx 3.2$ . Since F'(0) = R(0) < 1,  $F'(b_2) = 1 + b_2 R'(b_2) > 1$ ,  $F'(b_3) = 1 + b_3 R'(b_3) < 1$ , we deduce that equilibria  $b_1 = 0$  and  $b_3$ are locally asymptotically stable and  $b_2$  is unstable (see Corollary 5). Our computations show that the linearization of (1) at  $b_2$  has only one (real) eigenvalue with non-negative real part for  $\alpha = 6$ , p = 5, therefore we cannot invoke the Hopf bifurcation approach to prove the existence of a periodic orbit for (1). Simulating solutions b(t) of (1) for different initial data  $\phi$ , we took constant and periodic initial functions close to the intermediate equilibrium  $b_2$ . Figures 3 and 4 show that these solutions are asymptotically constant. Clearly, the nonlinearity F(b) is bistable here and we observe a kind of the Allee effect for the forest dynamics.

#### Acknowledgments

This research was supported in part by the projects FONDECYT 1231169 and AM-SUD220002 (ANID, Chile). The first author was also supported by ANID-Subdirección de Capital Humano/Doctorado Nacional/2024–21240616. The second author gratefully acknowledges the hospitality and the support of the MATRIX Institute allowing him to participate in the Research Program "Delay Differential Equations and Their Applications" (Australia, Creswick, 12–20 Dec 2023).

## References

- C. Barril, À. Calsina, O. Diekmann, J. Z. Farkas, On competition through growth reduction, e-print arXiv:2303.02981, https://doi.org/10.48550/arXiv.2303.02981
- [2] C. Barril, A. Calsina, O. Diekmann, J. Z. Farkas, On hierarchical competition through reduction of individual growth, Journal of Mathematical Biology, (2024) 88:66 https://doi.org/10.1007/s00285-024-02084-x
- [3] M. A. Caraballo-Ortiz, E. Santiago-Valentín, T. A. Carlo, Flower number and distance to neighbours affect the fecundity of Goetzea elegans (Solanaceae) Journal of Tropical Ecology, 27 (2011) 521–528.
- [4] O. Diekmann, M. Gyllenberg, Equations with infinite delay: blending the abstract and the concrete, J. Diff. Equations, 252 (2012), 819–851.
- [5] F. Herrera, S. Trofimchuk, Dynamics of one-dimensional maps and Gurtin-MacCamy's population model. Part I: asymptotically constant solutions, Ukrainian Math. J. 75 (2023), 1635–1651 https://doi.org/10.3842/umzh.v75i12.7678
- [6] F. Herrera, S. Trofimchuk, Global dynamics of a size-structured forest model, e-print arXiv:2401.08618, https://doi.org/10.48550/arXiv.2401.08618
- [7] T. Krisztin, H.-O. Walther and J. Wu, Shape, smoothness and invariant stratification of an attracting set for delayed monotone positive feedback, Fields Institute Monograph Series, Vol. 11, AMS, Providence, RI, 1999.
- [8] T. Krisztin, H.-O. Walther, Unique periodic orbits for delayed positive feedback and the global attractor. J. Dyn. Differ. Equ. 13, 1–57 (2001).
- [9] T. Krisztin, G. Vas, The unstable set of a periodic orbit for delayed positive feedback, J. Dyn. Differ. Equ. 28, (2016) 805–855.
- [10] Z. Ma, P. Magal, Global asymptotic stability for Gurtin-MacCamy's population dynamics model, Proceedings of the AMS,152 (2024), 765–780.
- [11] J.A.J Metz, O. Diekmann, The dynamics of physiologically structured populations, Lecture Notes in Biomathematics, 68, 1986.
- [12] H.L. Smith, Monotone dynamical systems, Amer. Math. Soc., Providence, 1995.