ON SMOOTH INFINITE DIMENSIONAL GRASSMANNIANS, SPLITTINGS AND NON-COMMUTATIVE GENERALIZED CROSS-RATIO MAPPINGS

JEAN-PIERRE MAGNOT

ABSTRACT. We describe basic diffeological structures related to splittings and Grassmannians for infinite dimensional vector spaces. We analyze and expand the notion of non-commutative cross-ratio and prove its smoothness. Then we illustrate this theory by examples, with some of them extracted from the existing literature related to infinite dimensional (Banach) Grassmannians, and others where the diffeological setting is a key primary step for rigorous definitions.

Keywords: Diffeological spaces, infinite dimensional Grassmanians, infinite dimensional groups, non-commutative cross-ratio, W-algebras.

MSC(2020): 53C30, 58A40, 57P99, 14M15, 53C15, 37K20.

1. INTRODUCTION

The notion of cross-ratio plays an important role in modern geometry, and its basic application to characterization of cocyclic points in the plane has been for a while at the curriculum of high school students in some EU states. A clear exposition of its importance can be found in [32]. Some various generalizations in symplectic and non-commutative geometry arise in recent algebraic and geometric works, see e.g. [14, 17, 21, 33, 53, 54, 64]. We analyze here one non-commutative analog to the cross-ratio present, algebraically, in a remark of [54] following the pioneering work [64]. We show that there are non necessary hypothesis in its definition, which gathers previously described compatible approaches. Therefore, we refresh the algebraic approach already highlighted in a *smooth* or *geometric* way. Our enhanced definition of non commutative cross-ratio between two admissible splittings is shown to produce a vector space isomorphism which depends smoothly on the underlying admissible couple of splittings. Then we build up remarkable involutions on the space of admissible couples splittings, and derive from these involutions and from our initial non-commutative cross-ratio mapping that we note by DV, two other mappings, that we note by DV and Ξ , from admissible couples of splittings to GL(V). For this, a safe notion of smoothness on Grassmanians, splittings, and on the group GL(V) is necessary.

The geometry of finite dimensional Grassmanians in a finite dimensional vector space is an underlying object for many methods in differential geometry. Its generalizations to infinite dimensional settings are, in the actual state of knowledge, subject to technical restrictions, and the same holds for the geometric structures that can be developed on Grassmanian "manifolds" in view of applications. In most examples, the Grassmanian space of an infinite dimensional vector space V is constituted by finite dimensional and finite codimensional vector subspaces of V, or it

JEAN-PIERRE MAGNOT

is defined through a fixed splitting $V = V_+ \oplus V_-$ by two closed supplementary infinite dimensional vector spaces of V. In each of these cases, this is obviously only one part of the full space of closed vector subspaces (with topological supplement) of Vthat is under consideration. In particular, the important example of the restricted Grassmanian defined in the works of Pressley, Segal and Wilson (see e.g. [47, 52] for two expositary textbooks on this subject) is defined through the eigenspaces of the sign of the Dirac operator acting on $L^2(S^1, \mathbb{C})$, partly inspired by Sato's geometric approaches on integrable systems. These initial works had ramifications until very recent works, in related but different subjects. A selected bibliography on these developments is [1, 2, 3, 6, 10, 15, 21, 26, 37, 48, 54, 59, 60].

In all these works, one main technical limit to the investigations is the necessary underlying Hilbert or Banach space topology on the vector space V, that enables, using smart restrictions, to define an atlas on the space of so-called restricted Grassmanians. We propose here to describe a differential geometry in the generalized setting of diffeologies. We study spaces of splittings of an infinite dimensional vector space that can be in the category of Hilbert, Banach, Fréchet, locally convex, or even more generally, Frölicher or diffeological vector spaces. Splittings are understood as eigenspaces of linear symmetries of V. Within diffeological considerations, technical difficulties vanish to define a well-suited smoothness on spaces of splittings, and then on the space of Grassmanians constituted by vector subspaces (closed, with supplement).

The same way, the same diffeologies enable to talk safely of smooth linear operators acting on an infinite dimensional vector space V which is not in the Hilbert or Banach setting, and to define smoothness on L(V) and GL(V). Therefore, this is natural for us to start this paper by necessary statements on diffeologies, known and new (section 2). In particular, a short exposition on diffeological vector spaces, diffeological algebras and their groups of the units, and on the particular case of the linear algebra L(V) and the linear group GL(V) of a diffeological vector space V, completes the existing statements in the literature. Therefore, we have all the necessary technical tools to define, given a diffeological algebra A, the space of splittings of V from the space S(A) of involutive elements (or symmetries, section 3.1) of A. Then the non-commutative cross-ratio DV already announced can be defined on *admissible* couples of splittings Adm(V) in section 3.2 and is easily proven to be smooth, with values in a linear group. From these basic descriptions, we highlight involutions on $\mathbf{S}(A)$ and on Adm(V) that are intrinsically linked with the construction of the non-commutative cross-ratio. Then we extend DV to two GL(V)-valued maps in section 3.4 and in section 3.5. The first one, DV, is a direct extension to the cross-ratio map. The second one, noted by Ξ , defines a smooth diffeomorphism on the space of topological supplement to a vector subspace of V. We call this diffeological space the Grassmanian relative to V. After these abstract constructions, this is highly needed to develop examples in section 4. The Pressley-Segal-Wilson restricted Grassmanian gives us a first framework in section 4.1 that enables to highlight basic geometric properties of admissible splittings as they are described in our approach. But this setting is not adapted for a simple calculation of a non-trivial non-commutative cross-ratio in the actual state of knowledge and in the context of an illustration of this work. In order to produce a non trivial crossratio in an accessible and straightforward computation, we produce in section 4.2.2 a second example of admissible splittings in the algebra of formal classical pseudodifferential operators which can be understood as a locally convex non complete vector space. The admissible couple of splittings that we consider is derived, from one side, from Kontsevich and Vishik's works on spectral analysis [29], see e.g. [57], and also from the Pressley-Segal-Wilson polarization. In this example, the non-commutative cross-ratio DV is proven to be non-trivial. Finally, we consider splittings on spaces of mappings defined by signed measures. We prove that two different signed measures do not produce any admissible couple of splittings in the context that we consider.

These three settings intend to produce examples and to give illustrations to the complex constructions that are developed. The main novelty of this paper is, to our own feeling, on one side the smooth structures on Grassmanians and splittings that are no longer technically restrained, and on the other side the maps DV, DV and Ξ , that extend the (commutative) cross-ratio in non-commutative setting, are now safely defined on admissible couples of splittings. Moreover, two among the three of our testing examples (formal pseudo-differential operators, signed measures) could not be reached in a Banach manifold category. Our approach has now to be compared to the other existing ones.

2. A presentation of diffeologies and Frölicher spaces

2.1. **Diffeological and Frölicher spaces.** In this section we follow the expositions appearing in [41, 43, 44], and complete the classical exposition [27] along the lines of the recent review [25].

Definition 1. Let X be a set.

• A **p-parametrization** of dimension p on X is a map from an open subset O of \mathbb{R}^p to X.

• A diffeology on X is a set \mathcal{P} of parametrizations on X such that:

- For each $p \in \mathbb{N}$, any constant map $\mathbb{R}^p \to X$ is in \mathcal{P} ;

- For any set of indexes I and any family $\{f_i: O_i \to X\}_{i \in I}$ of compatible maps that extend to a map $f: \bigcup_{i \in I} O_i \to X$, if $\{f_i: O_i \to X\}_{i \in I} \subset \mathcal{P}$, then $f \in \mathcal{P}$.

- $\forall f \in \mathcal{P}, f: O \subset \mathbb{R}^p \to X, g \forall g \in C^{\infty}(O', O), in which O' is an open subset of an Euclidean space, we have <math>f \circ g \in \mathcal{P}$.

 (X, \mathcal{P}) is called if called a **diffeological space** \mathcal{P} is a diffeology on X and, if (X, \mathcal{P}) and (X', \mathcal{P}') are two diffeological spaces, a map $f : X \to X'$ is **smooth** if and only if $f \circ \mathcal{P} \subset \mathcal{P}'$.

For an historical introduction to diffeologies, see [58], inspired by the remarks from Chen in [12]. The textbook [27] gives a comprehensive exposition of basic concepts.

Definition 2. Let (X, \mathcal{P}) be a diffeological space. Let \mathcal{P}' be a diffeology on X. Then \mathcal{P}' is a **subdiffeology** of \mathcal{P} if and only if $\mathcal{P}' \subset \mathcal{P}$ or, equivalently, $Id_X : (X, \mathcal{P}') \to (X, \mathcal{P})$ is smooth.

Example 3. Let X be a nonempty set. There is a minimal diffeology for inclusion. This diffeology is made of constant parametrizations and is called the **discrete diffeology**.

The category of diffeological spaces is very large, and carries many different pathological examples. Therefore, a restricted category can be useful, the category of Frölicher spaces.

Definition 4. A Frölicher space is a triple $(X, \mathcal{F}, \mathcal{C})$ such that

- \mathcal{C} is a set of paths $\mathbb{R} \to X$, called set of contours.

- \mathcal{F} is the set of functions from X to \mathbb{R} , such that a function $f: X \to \mathbb{R}$ is in \mathcal{F} if and only if for any $c \in C$, $f \circ c \in C^{\infty}(\mathbb{R}, \mathbb{R})$;

- Let $c : \mathbb{R} \to X$. $c \in \mathcal{C} \Leftrightarrow \forall f \in \mathcal{F}, f \circ c \in C^{\infty}(\mathbb{R}, \mathbb{R})$.

If $(X, \mathcal{F}, \mathcal{C})$ and $(X', \mathcal{F}', \mathcal{C}')$ are two Frölicher spaces, a map $f : X \to X'$ is smooth if and only if $\mathcal{F}' \circ f \circ \mathcal{C} \subset C^{\infty}(\mathbb{R}, \mathbb{R})$.

This definition first appeared in [20] but the actual terminology was fixed to our knowledge in [31]. The comparison of these two frameworks was published in [39] in its first steps, the reader can also see [4, 41, 42, 43, 61] for extended expositions. In particular, it is explained in [43] that Diffeological, Frölicher and Gâteaux smoothness are the same notion if we restrict ourselves to a Fréchet context. For this, let us analyze how we generate a Frölicher space, that is, how we implement a Frölicher structures on a given set X. Any family of maps \mathcal{F}_q from X to \mathbb{R} generates a Frölicher structure $(X, \mathcal{F}, \mathcal{C})$ by setting, after [31]:

- $\begin{aligned} \mathcal{C} &= \{ c : \mathbb{R} \to X \text{ such that } \mathcal{F}_g \circ c \subset C^{\infty}(\mathbb{R}, \mathbb{R}) \} \\ \mathcal{F} &= \{ f : X \to \mathbb{R} \text{ such that } f \circ \mathcal{C} \subset C^{\infty}(\mathbb{R}, \mathbb{R}) \}. \end{aligned}$

We call \mathcal{F}_q a generating set of functions for the Frölicher structure $(X, \mathcal{F}, \mathcal{C})$. A Frölicher space $(X, \mathcal{F}, \mathcal{C})$ carries a natural topology, the pull-back topology of \mathbb{R} via \mathcal{F} . Alternatively, one can start alternatively from a generating set of contours \mathcal{C}_{q} , and set:

- $\mathcal{F} = \{ f : X \to \mathbb{R} \text{ such that } f \circ \mathcal{C}_q \subset C^\infty(\mathbb{R}, \mathbb{R}) \}$

 $-\mathcal{C} = \{c : \mathbb{R} \to X \text{ such that } \mathcal{F} \circ c \subset C^{\infty}(\mathbb{R}, \mathbb{R}) \}.$

In the case of a finite dimensional differentiable manifold X we can take \mathcal{F} as the set of all smooth maps from X to \mathbb{R} , and \mathcal{C} the set of all smooth paths from \mathbb{R} to X. Then, the underlying topology of the Frölicher structure is the same as the manifold topology [31]. We also remark that if $(X, \mathcal{F}, \mathcal{C})$ is a Frölicher space, we can define a natural diffeology on X by using the following family of maps fdefined on open domains D(f) of Euclidean spaces, see [39]:

$$\mathcal{P}_{\infty}(\mathcal{F}) = \prod_{p \in \mathbb{N}^*} \{ f : D(f) \to X \, | \, D(f) \text{ is open in } \mathbb{R}^p \text{ and } \mathcal{F} \circ f \in C^{\infty}(D(f), \mathbb{R}) \}.$$

If X is a finite-dimensional differentiable manifold, setting $\mathcal{F} = C^{\infty}(X, \mathbb{R})$, this diffeology is called the *nebulae diffeology*, see e.g. [27]. We extend this diffeology to any Frölicher space. Now, we can easily show the following:

Proposition 5. [39] $Let(X, \mathcal{F}, \mathcal{C})$ and $(X', \mathcal{F}', \mathcal{C}')$ be two Frölicher spaces. A map $f: X \to X'$ is smooth in the sense of Frölicher if and only if it is smooth for the underlying diffeologies $\mathcal{P}_{\infty}(\mathcal{F})$ and $\mathcal{P}_{\infty}(\mathcal{F}')$.

Thus, Proposition 5 and the foregoing remarks imply that the following implications hold:

smooth manifold \Rightarrow Frölicher space \Rightarrow diffeological space

2.2. Diffeologies on spaces of mappings, products, quotients, subsets and algebraic structures.

Proposition 6. [27, 58] Let (X, \mathcal{P}) and (X', \mathcal{P}') be two diffeological spaces. There exists a diffeology $\mathcal{P} \times \mathcal{P}'$ on $X \times X'$ made of plots $g : O \to X \times X'$ that decompose as $g = f \times f'$, where $f : O \to X \in \mathcal{P}$ and $f' : O \to X' \in \mathcal{P}'$. We call it the **product diffeology**, and this construction extends to an infinite (maybe not countable) product.

The same holds on contours of Frölicher spaces in a cartesian product (compare with [31]). We can also state the above result for infinite products; we simply take Cartesian products of the plots, or of the contours.

After [58] and [27, p. 27], let (X, \mathcal{P}) be a diffeological space, and let X' be a set. Let $f: X \to X'$ be a map. We define the **push-forward diffeology**, noted by $f_*(\mathcal{P})$, as the coarsest (i.e. minimal for inclusion) among the diffeologies on X', which contains $f \circ \mathcal{P}$, that is, for which f is smooth. Conversely, if (X', \mathcal{P}') is a diffeological space and if X is a set, if $f: X \to X'$, the **pull-back diffeology** noted by $f^*(\mathcal{P}')$ is the diffeology on X, maximal for inclusion, for which f is smooth.

Proposition 7. [27, 58] Let (X, \mathcal{P}) b a diffeological space and \mathcal{R} an equivalence relation on X. There is a natural diffeology on X/\mathcal{R} , noted by \mathcal{P}/\mathcal{R} , defined as the push-forward diffeology on X/\mathcal{R} by the quotient projection $X \to X/\mathcal{R}$. This diffeology is made of plots that read locally as maps $\pi \circ p$, where $p \in \mathcal{P}$.

Given a subset $X_0 \subset X$, where X is a Frölicher space or a diffeological space, we equip X_0 with structures induced by X as follows:

(1) If X is equipped with a diffeology \mathcal{P} , we define a diffeology \mathcal{P}_0 on X_0 called the **subset or trace diffeology**, see [58, 27], by setting

 $\mathcal{P}_0 = \{ p \in \mathcal{P} \text{ such that the image of } p \text{ is a subset of } X_0 \}.$

(2) If $(X, \mathcal{F}, \mathcal{C})$ is a Frölicher space, we take as a generating set of maps \mathcal{F}_g on X_0 the restrictions of the maps $f \in \mathcal{F}$. In this case, the contours (resp. the induced diffeology) on X_0 are the contours (resp. the plots) on X whose images are a subset of X_0 .

Example 8. Let us consider the infinite product diffeology on Y^X . This is the largest diffeology for which the evaluation maps ev_x are smooth for each $x \in X$. Therefore, any other diffeology on any subset of Y^X for which all the evaluation maps are smooth is a subdiffeology of the subset diffeology inherited from this product diffeology.

Following [27], let (X, \mathcal{P}) and (X', \mathcal{P}') be diffeological spaces. Let $M \subset C^{\infty}(X, X')$ be a set of smooth maps. The **functional diffeology** on S is the diffeology \mathcal{P}_S made of plots

$$\rho: D(\rho) \subset \mathbb{R}^k \to S$$

such that, for each $p \in \mathcal{P}$, the maps $\Phi_{\rho,p} : (x, y) \in D(p) \times D(\rho) \mapsto \rho(y)(x) \in X'$ are plots of \mathcal{P}' . We have, see [27, Paragraph 1.60]:

Proposition 9. Let X, Y, Z be diffeological spaces. Then,

$$C^{\infty}(X \times Y, Z) = C^{\infty}(X, C^{\infty}(Y, Z)) = C^{\infty}(Y, C^{\infty}(X, Z))$$

as diffeological spaces equipped with functional diffeologies.

2.3. Groups, vector spaces, algebra and fiber bundles in diffeologies. Given an algebraic structure, we can define a corresponding compatible diffeological (resp. Frölicher) structure, see for instance [35]. For example, see [27, pp. 66-68], if \mathbb{R} is equipped with its canonical diffeology (resp. Frölicher structure), we say that an \mathbb{R} -vector space equipped with a diffeology (resp. Frölicher structure) is a diffeological (resp. Frölicher) vector space if addition and scalar multiplication are smooth. We state the same way, following [35, 41]:

Definition 10. Let G be a group equipped with a diffeology (resp. Frölicher structure). We call it a **diffeological (resp. Frölicher) group** if both multiplication and inversion are smooth.

2.3.1. The group of diffeomorphisms of a diffeological space. Let (X, \mathcal{P}) be a diffeological space. The space $C^{\infty}(X, X)$ is endowed with its functional diffeology and a diffeomorphism on X is a map $g \in C^{\infty}(X, X)$ such that there exists an inverse map $g^{-1} \in C^{\infty}(X, X)$ such that $g \circ g^{-1} = g^{-1} \circ g = Id_X$. The elementary algebraic properties of the inverse assert that the inverse is unique. The set of diffeomorphisms on X is noted by Diff(X). Unlike for the Fréchet Lie group of diffeomorphisms of a finite dimensional compact manifold, the group Diff(X) is not a priori a diffeological group because the inverse map $g \mapsto g^{-1}$ is not a priori smooth for the (trace of) the functional diffeology $\mathcal{P}_{Diff(X)}$ on Diff(X). In fact, only the composition of diffeomorphisms is smooth. Our unability to state smoothness of the inversion is due to a lack of inverse functions theorem. Therefore, we have to define the following diffeology.

Proposition 11. Let $\mathcal{P}_{Diff(X)}^{(-1)}$ be the diffeology on Diff(X) defined by pull-back

$$\mathcal{P}_{Diff(X)}^{(-1)} = \Phi^* \left(\mathcal{P}_{Diff(X)} \times \mathcal{P}_{Diff(X)} \right)$$

where

$$\Phi: g \in Diff(X) \mapsto (g, g^{-1}) \in Diff(X) \times Diff(X).$$

Then $\left(Diff(X), \mathcal{P}_{Diff(X)}^{(-1)}\right)$ is a diffeological group.

The proof is straightforward. Therefore, considering now a diffeological ring R, the set of invertible elements R^* of R has itself a specific diffeology $\mathcal{P}_{R^*}^{(-1)}$ which differ from the subset diffeology \mathcal{P}_{R^*} inherited from the inclusion $R^* \subset R$. We only have in general

$$\mathcal{P}_{R^*}^{(-1)} \subset \mathcal{P}_{R^*}.$$

The same construction holds for a diffeological field, a diffeological module, a diffeological algebra: the group of the units (i.e. the group of the invertible elements) can be endowed with a diffeology "of $\mathcal{P}^{(-1)}$ -type".

Remark 12. In the case of a \mathbb{R} -algebra or a \mathbb{C} -algebras A, the group A^* may carry this pathology and there is an identified class of examples of infinite dimensional algebras for which A^* is a Lie group for the subset diffeology, in the setting of locally convex, Mackey-complete algebras [23, 24]. Particular examples of this class are Banach algebras. In that case, $\mathcal{P}_{A^*}^{(-1)} = \mathcal{P}_{A^*}$.

Example 13. When considering fields, the groups \mathbb{R}^* and \mathbb{C}^* have inversion which is smooth for the subset diffeology, in other words, $\mathcal{P}_{\mathbb{R}^*}^{(-1)} = \mathcal{P}_{\mathbb{R}^*}$ and $\mathcal{P}_{\mathbb{C}^*}^{(-1)} = \mathcal{P}_{\mathbb{C}^*}$.

2.3.2. Linear group of a diffeological vector space. Let V, V' be two diffeological vector spaces. We note by L(V, V') the space of smooth linear maps from V to V' equipped with the functional diffeology (understood as a subset diffeology inherited from the inclusion $L(V; V') \subset C^{\infty}(V, V')$). We note L(V) for short, instead of L(V, V).

We recall that, with no local inverse functions theorem at hand, it is impossible to prove that the group GL(V) of invertible maps has a smooth inversion, and following [25], we equip the (algebraic) group GL(V) with the diffeology of Diff(V)(understood as the subset diffeology inherited by the inclusion $GL(V) \subset Diff(V)$). Since composition and inversion is smooth on Diff(V), so it is on GL(V). This construction provides a concrete class of examples of diffeological algebras and their diffeological group of the units that complete the classical examples recalled in Remark 12 and where the diffeology $\mathcal{P}_{A^*}^{(-1)}$ may differ from the diffeology \mathcal{P}_{A^*} . These examples carry a wide range of linear algebras of vector spaces with adequate properties to define smoothness, including vector spaces with pathological properties, see e.g. [51, 63] for actual concerns on "exotic" diffeological vector spaces and their technical properties.

3. Projections, symmetries, Grassmannians, splittings and Non-commutative cross-ratio

3.1. Symmetries and projections. Let A be a diffeological unital algebra with group of the units A^* . This algebra is assumed to act smoothly on a diffeological vector space V. We note by

$$\mathbf{P}(A) = \left\{ p \in A \,|\, p^2 = p \right\}$$

the set of projectors in A and by

$$\mathbf{S}(A) = \{ s \in A \, | \, \exists p \in \mathbf{P}(A), s = 2p - 1_A \} \,.$$

Remark 14. The classical condition $s^2 = 1_A$ is equivalent, that is,

$$\forall x \in V, x \mapsto \frac{1}{2}(x + s(x))$$

defines the projection $p \in \mathbf{P}(A)$ such that $s = 2p - 1_A$.

When A = L(V), this picture specialize to the following. We note by

$$\mathbf{P}(V) = \{ p \in L(V) \, | \, p^2 = p \}$$

the set of (diffeological) projectors in V and by

$$\mathbf{S}(V) = \left\{ s \in L(V) \, | \, s^2 = Id_V \right\}$$

the set of (diffeological) symmetries on V.

Lemma 15. Let i_A and i_{A^*} be the inclusion maps from $\mathbf{S}(A)$ to A and to A^* respectively, then $i_A^*(\mathcal{P}_A) = i_{A^*}^*(\mathcal{P}_{A^*}) = i_{A^*}^*(\mathcal{P}_{A^*}^{(-1)})$.

Proof. The map

$$\Phi: u \in A^* \mapsto (u, u^{-1}) \in A^2$$

restricts to the canonical (diagonal) inclusion $u \mapsto (u, u)$ from $\mathbf{S}(A)$ to A^2 .

We also note by \mathcal{P}_A the subset diffeology of $\mathbf{P}(A) \subset A$ and as well as the subset diffeology of $\mathbf{S}(A) \subset A$. We can state:

Proposition 16. There is a diffeomorphism

$$p \in \mathbf{P}(A) \mapsto 2p - 1_A \in \mathbf{S}(A)$$

for the diffeology \mathcal{P}_A on $\mathbf{P}(A)$.

Proof. It follows from the smoothness of addition and multiplication on A.

The previous results apply to $\mathbf{P}(V)$ and to $\mathbf{S}(V)$ according to the description of L(V) as a diffeological algebra given in section 2.3.2. Let us recall, following the classical algebraic decompositions, that given $s \in \mathbf{S}(A)$ and $u \in A$,

(1)
$$(s+u) \in \mathbf{S}(A) \quad \Leftrightarrow \quad su+us+u^2 = 0$$

Let

$$AS(s): u \in A \mapsto us + su \in A.$$

Let $s \in \mathbf{S}(A)$. We define

$$\pi_s^T : a \in A \mapsto \frac{1}{2} \left(a - sas \right)$$

and

$$\pi^N_s: a \in A \mapsto \frac{1}{2} \left(a + sas \right)$$

Let us now give an easy but useful lemma:

Lemma 17. $\forall s \in \mathbf{S}(A)$,

- $\pi_s^T \in \mathbf{P}(L(A))$ and $\pi_s^N \in \mathbf{P}(L(A))$ $Im(\pi_s^T) = Ker(\pi_s^N)$ and $Im(\pi_s^N) = Ker(\pi_s^T)$ $Im(\pi_s^T) = KerAS(s)$
- A is isomorphic, as a diffeological vector space, with

$$Im(\pi_s^T) \oplus Im(\pi_s^N)$$

where each component is equipped with the subset diffeology. In other words, each $a \in A$ splits uniquely into $a = a_s^T + a_s^N$, with $a_s^T \in Im(\pi_s^T)$, $a_s^N \in Im(\pi_s^N)$, and the mapping $a \mapsto (a_s^T, a_s^N) = (\pi_s^T(a), \pi_s^N(a))$ is a diffeomorphism.

Proof. In the results that have to be proven, all the algebraic parts are elementary and well-established.

For the diffeological parts, it follows from the smoothness of scalar multiplication, addition and multiplication in A. \square

Let us now analyze the interaction between the splitting $a = a_s^T + a_s^N$ with the multiplication. We calculate, for $s \in \mathbf{S}(A)$ and for $(a, b) \in A^2$,

$$\begin{split} a_s^T b_s^T &= \frac{1}{4} \left(a - sas \right) \left(b - sbs \right) \\ &= \frac{1}{2} \pi_s^N (ab) - \frac{1}{4} \left(sasb + asbs \right), \\ a_s^N b_s^N &= \frac{1}{4} \left(a + sas \right) \left(b + sbs \right) \\ &= \frac{1}{2} \pi_s^N (ab) + \frac{1}{4} \left(sasb + asbs \right), \end{split}$$

$$\begin{aligned} a_s^T b_s^N &= \frac{1}{4} (a - sas) (b + sbs) \\ &= \frac{1}{2} \pi_s^T (ab) - \frac{1}{4} (sasb - asbs), \\ a_s^N b_s^T &= \frac{1}{4} (a + sas) (b - sbs) \\ &= \frac{1}{2} \pi_s^T (ab) + \frac{1}{4} (sasb - asbs). \end{aligned}$$

These are all intermediate results which have as a consequence that $ab = (ab)_s^T + (ab)_s^N$ but we can now precise a "polarization like" property which is a straightforward consequence of our computations:

Lemma 18. Let $s \in \mathbf{S}(A)$ and let $(a, b) \in A^2$.

$$(ab)_s^T = a_s^T b_s^N + a_s^N b_s^T$$

and

$$(ab)_s^N = a_s^T b_s^T + a_s^N b_s^N.$$

Next theorem follows straightway from our preliminary constructions.

Theorem 19. for each $s \in \mathbf{S}(A)$, we define a \mathbb{Z}_2 -grading on $A = Im(\pi_s^T) \oplus Im(\pi_s^N)$ by assigning $deg_s(a_s^N) = 1$ and $deg_s(a_s^T) = -1$, and the algebra A reads as a diffeological \mathbb{Z}_2 -graded algebra with respect to the decomposition $A = Im(\pi_s^N) \oplus Im(\pi_s^T)$.

Moreover, the map

$$(a,s) \in A \times \mathbf{S}(A) \mapsto (a_s^N, a_s^T) \in A^2$$

is smooth.

Remark 20. $\forall s \in \mathbf{S}(A), s \in Im(\pi_s^N), Id_A \in Im(\pi_s^N) \text{ and } p = \frac{1}{2}(s + Id_A) \in Im(\pi_s^N).$

Let us now investigate the following problem: let $a \in A$. For which $s \in \mathbf{S}(A)$ do we have $a \in Im(\pi_s^N)$? or $a \in Im(\pi_s^N)$?

Proposition 21. Let $a \in A$. Then $a \in Im(\pi_s^T)$ if and only if

(2)
$$(a+s)^2 = a^2 + Id_A$$

Proof. $a \in Im(\pi_s^T) \leftrightarrow a \in Ker(\pi_s^N) \leftrightarrow as + sa = 0$. Then expanding

$$(a+s)^2 - a^2 - Id = as + sa$$

we get the result.

Proposition 22. Let $a \in A$. Then $a \in Im(\pi_s^N)$ if and only if

[a,s] = 0

which is equivalent to

$$(3) \qquad (a+s)(a-s) = a^2 - Id_A$$

or

(4)
$$(a+s)(a-s) = (a - Id_A)(a + Id_A)$$

Proof. $a \in Im(\pi_s^N) \leftrightarrow a \in Ker(\pi_s^T) \leftrightarrow as - sa = 0$. Then expanding

$$(a+s)(a-s) - a^2 + Id_A = as + sa$$

we get the result, since $[a, Id_A] = 0 \Leftrightarrow a^2 - Id_A = (a - Id_A)(a + Id_A).$

3.2. From splittings to Grassmannians and to projections and non-commutative cross-ratio.

Definition 23. The space of (diffeological) splittings of A is the space

$$\mathbf{Spl}(A) = \left\{ (V, W) \subset A^2 \mid \exists s \in \mathbf{S}(A), \ V = Ker(s - Id) \land W = Ker(s + Id) \right\}.$$

Proposition 24. We have equivalently

- $\mathbf{Spl}(A) = \left\{ (V, W) \subset A^2 \,|\, \exists p \in \mathbf{P}(A), \, V = Im(p) \land W = Ker(p) \right\}.$
- We equip $\mathbf{Spl}(A)$ with its push-forward diffeology $\mathcal{P}_{\mathbf{Spl}} = (Im \times Ker)_* \mathcal{P}_L$.

We define the map $J_{(-)} : a \in A \mapsto -a \in A$. Here are few easy properties:

- **Proposition 25.** (1) The map $Im \times Ker : \mathbf{P}(A) \to \mathbf{Spl}(A)$ is a diffeomorphism, and induces a diffeomorphism $Pol : \mathbf{S}(A) \to \mathbf{Spl}(A)$.
 - (2) $\forall (V, W) \in \mathbf{Spl}(A),$

 $J_{(-)}^{\mathbf{Spl}}(V,W) = Pol \circ J_{(-)} \circ Pol^{-1}(V,W) = (W,V).$

This enables to define a smooth involution on $\mathbf{Spl}(A)$ that we also note by $J_{(-)}$ instead of $J_{(-)}^{\mathbf{Spl}}$ when it carries no ambiguity.

The map *Pol* is such named because it is often called Polarization map in the literature. Therefore, let us analyze the non-commutative cross-ratio from [21, 64] which algebraic aspects are refreshed in [54]. We recall its definition along the lines of our notations, and both extend and complete the statements which are, to our opinion, a little too fastly stated in [54, 64], or at least, developed in a too reductive setting.

Definition 26. Let $((V_1, W_1), (V_2, W_2)) \in \operatorname{Spl}(A)^2$ such that $((V_1, W_2), (V_2, W_1)) \in \operatorname{Spl}(A)^2$. We call such two splittings an admissible couple of polarizations or admissible couple of splittings. We denote by Adm(V) the set of admissible couples of splittings. Then we define on Adm(V) the map DV with values in $L(V_1)$ defined by

 $DV(V_1, W_1, V_2, W_2) = (Im \times Ker)^{-1}(V_1, W_2) \circ (Im \times Ker)^{-1}(V_2, W_1)|_{V_1}.$

Remark 27. This definition corresponds to the composition diagram

$$V_1 \xrightarrow{W_1} V_2 \xrightarrow{W_2} V_1$$

present in [54, 64]. Comparing our statement to those present in [54, 64], the assumption $((V_1, W_2), (V_2, W_1)) \in \mathbf{Spl}(A)^2$ replaces the existence of a fixed isomorphism $\psi \in GL(V_1, W_1)$ which is not a priori necessary at this step. Therefore, our construction can be understood as a tentative of generalization of the setting described [54, 64] even if we are not sure that this is strictly speaking a "generalization", but maybe only an "adaptation". Let us now give more comments on the assumptions on (V_1, W_1) and (V_2, W_2) that one can find in [54, 64]. The assumption of isomorphism between V_i and W_i , for $i \in \{1, 2\}$, is not assumed here. However, we can check that the condition $((V_1, W_1), (V_2, W_2)) \in Adm(V)$, the diffeological vector spaces V_1 and V_2 are isomorphic. For this, we note by $p_{(V,W)}$ the projection on Vparallel to W, with $(V,W) \in \mathbf{Spl}(A)$. Indeed, given $x_1 \in V_1$ and $x \in \pi_{(V_1,W_1)}^{-1}(x_1)$, this is an easy exercise to check that $x_2 = p_{(V_2,W_1)}(x)$ only depends on x_1 and that $p_{(V_1,W_1)}(x_2) = x_1$. We can state a bit quickly that $(p_{(V_2,W_1)}|_{V_1})^{-1} = p_{(V_1,W_1)}|_{V_2}$. Therefore, our construction on admissible couples of splittings only implies that V_1 and V_2 are isomorphic, and that W_1 and W_2 are also isomorphic.

We have the following:

Proposition 28. The map DV is a smooth $GL(V_1)$ -valued map, and its inverse reads as

$$DV(V_1, W_1, V_2, W_2)^{-1} = DV(V_1, W_2, V_2, W_1).$$

Proof. Considering the diagram

$$V_1 \xrightarrow{W_2} V_2 \xrightarrow{W_1} V_1$$

(which is possible under our assumptions), we see that we construct the inverse of the map $DV(V_1, W_1, V_2, W_2)$. The smoothness properties follow from the smoothness of the elementary operations that we use in this construction.

Remark 29. The set of admissible couples of splittings, as it is defined here, is stable under the action of $J_{(-)}$ which is a smooth map on Adm(A).

Let us equip the space of admissible polarizations with its subset diffeology inherited from its inclusion in $\mathbf{Spl}(A)^2$.

Definition 30. The Grassmanian on A is the space

$$\mathbf{Gr}(A) = \{ V \subset A \mid \exists s \in \mathbf{S}(A), V = Im(s) \}.$$

We equip $\mathbf{Gr}(A)$ with its push-forward diffeology $\mathcal{P}_{\mathbf{Gr}} = Im_*\mathcal{P}_L$.

Denoting π_i the canonical projection on the *i*-th component of $\mathbf{Gr}(A)^2$ with $i \in \{1; 2\}$, and denoting the same way the restriction of the projection maps to $\mathbf{Spl}(A)$, we have the following commutative diagram:



where the horizontal arrow is the canonical inclusion map.

3.3. Yet another involutions on Adm(V). Let us now analyze the maps

$$J: ((V_1, W_1), (V_2, W_2)) \in Adm(V) \mapsto ((V_1, W_2), (V_2, W_1))$$

and

$$J_{sw}: ((V_1, W_1), (V_2, W_2)) \in Adm(V) \mapsto ((V_2, W_2), (V_1, W_1))$$

Proposition 31. The maps \tilde{J} and J_{sw} are smooth on Adm(V) with values in Adm(V).

Proof. By definition, we already know that $Im\tilde{J} = Adm(V)$. The same way, it is straightforward to prove that if $((V_1, W_1), (V_2, W_2)) \in Adm(V)$, then $((V_2, W_2), (V_1, W_1)) \in Adm(V)$. Moreover, since the map

$$sw: (a,b) \in \mathbf{S}(A)^2 \to (b,a) \in \mathbf{S}(A)^2$$

is smooth, we get that

$$J_{sw} = (Pol \times Pol) \circ sw \circ (Pol^{-1} \times Pol^{-1})$$

is smooth. Moreover, the diffeology on Adm(V) is the subset diffeology inherited from $\mathbf{Gr}(A)^4$. Therefore, \tilde{J} is smooth.

Remark 32. We recall that

$$\mathfrak{D}_4 = \{ Id_{\mathbb{N}_4}, (12)(34), (13)(24), (14)(23) \}$$

is a subgroup of \mathfrak{S}_4 . The map J_{sw} corresponds indexwise to the action of $(13)(24) \in \mathfrak{D}_4$. The same way, $J_{(-)}$ corresponds to $(12)(34) \in \mathfrak{D}_4$. But \tilde{J} is not in \mathfrak{D}_4 but correspond to $(24) \in \mathfrak{S}_4 - \mathfrak{D}_4$.

3.4. Non-commutative cross-ratio generates GL(V)-valued mappings.

Theorem 33. The non-commutative cross ratio DV on the space of admissible couples $((V_1, W_1), (V_2, W_2))$ of polarizations defines the map

$$\Phi: ((V_1, W_1), (V_2, W_2)) \mapsto DV(V_1, W_1, V_2, W_2) \circ (Im \times Ker)^{-1}(V_1, W_1)$$

which is a smooth map with values in $L(V, V_1) \subset L(V)$ and moreover, if $a = \phi((V_1, W_1), (V_2, W_2))$, we have

$$Ker(a) \supset W_1 \land Im(a) \subset V_1.$$

Remark 34. We have to precise that the statement of smoothness is two-fold in this theorem: for any fixed admissible couple of polarizations $((V_1, W_1), (V_2, W_2))$, the map $\Phi((V_1, W_1), (V_2, W_2)) : V \to V$ is smooth and the maps $\Phi((V_1, W_1), (V_2, W_2)) \in L(V)$ depend smoothly on $((V_1, W_1), (V_2, W_2))$, where smoothness in with respect of the diffeology on Adm(A) in this second aspect.

Proof. Smoothness follows from the smoothness of the diffeomorphism $Im \times Ker$, and the algebraic part is direct by construction.

Moreover, we can define and state:

Definition 35. Adm(V), as it is defined here, is stable under the action of $J_{(-)}$. Therefore, we can define a global non-commutative cross-ratio map DV on admissible couples of splittings by

$$DV((V_1, W_1), (V_2, W_2)) = \Phi((V_1, W_1), (V_2, W_2)) + \Phi(J_{(-)}(V_1, W_1), J_{(-)}(V_2, W_2)).$$

Remark 36. By direct computations on projections, we have:

$$\forall ((V_1, W_1)(V_2, W_2)) \in Adm(V), [(V_1 = V_2) \lor (W_1 = W_2)] \Rightarrow \tilde{DV}((V_1, W_1), (V_2, W_2)) = Id_V.$$

Let us give now an elementary example in which $DV((V_1, W_1), (V_2, W_2)) \neq Id_V$.

Example 37. Let $V = \mathbb{R}^2$,

$$V_1 = \{ (x, y) \in \mathbb{R}^2 | y = 0 \},\$$

$$V_2 = \{ (x, y) \in \mathbb{R}^2 | x = y \},\$$

$$\begin{split} W_1 &= \{(x,y) \in \mathbb{R}^2 \,|\, x = 0\}\\ and \; W_2 &= \{(x,y) \in \mathbb{R}^2 \,|\, x + y = 0\}.\\ Then \; \tilde{DV}((V_1,W_1),(V_2,W_2))(x,y) &= (2x,2y), \; therefore \end{split}$$

$$\tilde{DV}((V_1, W_1), (V_2, W_2)) = 2Id_{\mathbb{R}^2}.$$

More generally, for $\theta \in (0; \frac{\pi}{2})$ with $V = \mathbb{R}^2$,

$$V_1 = \{(x, y) \in \mathbb{R}^2 | y = 0\},\$$

$$V_2 = \{(x, y) \in \mathbb{R}^2 | y = \tan(\theta)x\},\$$

$$W_1 = \{(x, y) \in \mathbb{R}^2 | x = 0\}\$$
and $W_2 = \{(x, y) \in \mathbb{R}^2 | x + \tan(\theta)y = 0\}.\$

Then

$$\tilde{DV}((V_1, W_1), (V_2, W_2)) = \frac{1}{\cos^2(\theta)} Id_{\mathbb{R}^2}.$$

More complex examples will be given in the dedicated section.

Lemma 38. Let $((V_1, W_1)(V_2, W_2)) \in Adm(V)$.

$$\left(\tilde{DV}((V_1, W_1), (V_2, W_2))\right)^{-1} = \Phi \circ \tilde{J} + \Phi \circ \tilde{J} \circ J_{(-)},$$

therefore the map

$$((V_1, W_1)(V_2, W_2)) \in Adm(V) \mapsto \left(\tilde{DV}((V_1, W_1), (V_2, W_2))\right)^{-1}$$

is smooth in L(V).

Theorem 39. Let $((V_1, V_2), (W_1, W_2)) \in Adm(V)$.

- if $s = Pol^{-1}(V_1, W_1), \Phi((V_1, W_1), (V_2, W_2)) \in Im(\pi_s^N).$
- If $((V_1, W_1), (V_2, W_2))$ is an admissible couple of splittings,

 $Im\tilde{DV}((V_1, W_1), (V_2, W_2)) \in GL(V_1) \times GL(W_1) \subset GL(V)$

• If Adm(V) is endowed with its subset diffeology as a subset of Spl(V), then

$$DV: Adm(V) \to GL(V)$$

is smooth.

The proof is direct from previous constructions.

Theorem 40. Let $((V_1, W_1), (V_2, W_2)) \in Adm(V)$. Then the map $\Xi((V_1, W_1), (V_2, W_2))$ defined by

 $\Xi((V_1, W_1), (V_2, W_2)) = \tilde{DV}((V_1, W_1), (V_2, W_2)) \circ \tilde{DV}((V_1, W_2), (V_2, W_1)) \in GL(V)$

is in GL(V) and satisfies:

$$\Xi((V_1, W_1), (V_2, W_2))|_{V_1} = Id_{V_1}.$$

Moreover the map

$$\Xi: Adm(V) \to GL(V)$$

is smooth.

Proof. By Theorem 39, $\tilde{DV} \in C^{\infty}(Adm(V), GL(V))$, therefore we also have that $\Xi \in C^{\infty}(Adm(V), GL(V))$, writing $\Xi(.) = \tilde{DV}(.) \circ \tilde{DV}(\tilde{J}(.))$. Let us now analyze $\Xi((V_1, W_1), (V_2, W_2))|_{V_1} = \tilde{DV}((V_1, W_1), (V_2, W_2)) \circ \tilde{DV}((V_1, W_2), (V_2, W_1))|_{V_1}$ $= \tilde{DV}((V_1, W_1), (V_2, W_2))|_{V_1} \circ DV((V_1, W_2), (V_2, W_1))|_{V_1}$ $= DV((V_1, W_1), (V_2, W_2)) \circ DV((V_1, W_2), (V_2, W_1))|_{V_1}$ $= Id_{V_1}$ by Proposition 28.

3.5. From non-commutative cross-ratio to relative Grassmannians. Let $V_1 \in \mathbf{Gr}(A)$. Let us now define the following domain:

Definition 41. We define

$$Gr_{V_1}(A) = \pi_1^{-1}(V_1) = \{ W_1 \in Gr(A) \, | \, (V_1, W_1) \in Spl(A) \}$$

and we call it the Grassmannian relative to V_1 , or V_1 -Grassmannian for short.

This is the main object studied in this section. Before that, we analyze the map Ξ in the blockwise decomposition $V_1 \oplus W_1$. We have that

$$\Xi((V_1, W_1), (V_2, W_2)) = \begin{pmatrix} Id_{V_1} & * \\ 0 & * \end{pmatrix}.$$

Let us now analyze, for $w_2 \in W_2$!

$$\begin{aligned} \Xi((V_1, W_1), (V_2, W_2))|_{W_2} &= \tilde{DV}((V_1, W_1), (V_2, W_2)) \circ \tilde{DV}((V_1, W_2), (V_2, W_1))|_{W_2} \\ &= \tilde{DV}((V_1, W_1), (V_2, W_2))|_{W_2} \circ DV((W_2, V_1), (W_1, V_2)). \end{aligned}$$

Let $w'_2 = DV((W_2, V_1), (W_1, V_2))(w_2)$. In the decomposition $V = V_1 \oplus W_1$,

$$w_2' = (w_2')_{V_1} + (w_2')_{W_1},$$

and

Since $DV((W_2, V_1), (W_1, V_2)) \in GL(W_2)$ and $DV((V_1, W_1), (V_2, W_2)) \in GL(V_1)$, the first term of the decomposition $DV((V_1, W_1), (V_2, W_2))((w'_2)_{V_1})$ vanishes identically if and only if

$$\forall w_2 \in W_2, (w_2')_{V_1} = 0 \Leftrightarrow W_1 = W_2.$$

Theorem 42. Let $V_1 \in \mathbf{Gr}(A)$. Then, $\forall (W_1, V_2, W_2) \in \mathbf{Gr}(A)$ such that $((V_1, W_1), (V_2, W_2)) \in Adm(A)$, then the map $\Xi((V_1, W_1), (V_2, W_2))$ defines a diffeomorphism on $\mathbf{Gr}_{V_1}(A)$ defined by

$$W \in \mathbf{Gr}_{V_1}(A) \mapsto \Xi((V_1, W_1), (V_2, W_2))(W).$$

Moreover, setting

$$D_{V_1} = \{ (W_1, V_2, W_2) \in \mathbf{Gr}(A) \mid ((V_1, W_1), (V_2, W_2)) \in Adm(A) \}$$

The map

$$\Xi: (W_1, V_2, W_2) \in D_{V_1} \mapsto \Xi((V_1, W_1), (V_2, W_2)) \in Diff(\mathbf{Gr}_{V_1}(A))$$

is smooth.

Proof. For $((V_1, W_1), (V_2, W_2)) \in Adm(A)$,

 $\begin{cases} \Xi((V_1, W_1), (V_2, W_2))(V_1) &= V_1 \\ \Xi((V_1, W_1), (V_2, W_2))(W) &= W' \end{cases}$

and since the map $\Xi((V_1, W_1), (V_2, W_2))$ is in GL(V),

$$V = \Xi((V_1, W_1), (V_2, W_2))(V_1 \oplus W) = V_1 + W' = V_1 \oplus W'.$$

Therefore, $(V_1, W') \in \mathbf{Spl}(A)$ and $\Xi((V_1, W_1), (V_2, W_2))$ defines a map from $\mathbf{Gr}_{V_1}(A)$ to itself. Moreover,

$$\left(\Xi((V_1, W_1), (V_2, W_2))\right)^{-1} = \left(\tilde{DV}((V_1, W_2), (V_2, W_1))\right)^{-1} \circ \left(\tilde{DV}((V_1, W_1), (V_2, W_2))\right)^{-1}$$

and

$$(V_1, W') = Pol^{-1} \left(\Xi((V_1, W_1), (V_2, W_2)) \circ Pol(V_1, W) \circ (\Xi((V_1, W_1), (V_2, W_2)))^{-1} \right)$$

which proves simultaneously that the map $\Xi((V_1, W_1), (V_2, W_2))$ from $\mathbf{Gr}_{V_1}(A)$ to itself is smooth, that the map $\Xi: (W_1, V_2, W_2) \in D_{V_1} \mapsto \Xi((V_1, W_1), (V_2, W_2))$ is smooth and that

$$\Xi((V_1, W_1), (V_2, W_2))^{-1} : W' \mapsto$$

$$\pi_2 \circ Pol^{-1} \left(\Xi((V_1, W_1), (V_2, W_2)) \circ Pol(V_1, W') \circ (\Xi((V_1, W_1), (V_2, W_2)))^{-1} \right)$$

is smooth by smoothness of each operation in these formulas.

4. Selected examples

4.1. On two classical polarizations in quantum field theory. We describe here the splittings derived from the spectrum of a Dirac operator. The main reference for a classical description of these objects is [52].

4.1.1. On the polarizations of $Map(S^1, \mathbb{C})$. Lett us start with the complex Hilbert space $L^2(S^1, \mathbb{C})$, with $S^1 = \mathbb{R}/2\pi\mathbb{Z} = \{z \in \mathbb{C} \mid |z| = 1\}$. The Fourier series are based on the orthonormal base $(z^n)_{n \in \mathbb{Z}}$, (where $z = e^{ix}$) defined by the eigenvectors of the (positive) Laplacian $\Delta = -\frac{\partial^2}{\partial x^2}$. The associated Dirac operator $D = -i\frac{\partial}{\partial x}$ splits $E = L^2(S^1, \mathbb{C})$ into three spaces:

- $E_{+} = Span(z^{n} | n > 0)$ associated to the positive eigenvalues of D,
- $E_{-} = Span(z^{n} | n < 0)$ associated to the negative eigenvalues of D,
- E₀ ~ C the space of constant maps, which is the eigenspace associated to the eigenvalue 0 of D.

This decomposition is L^2 -orthogonal. In the sequel, we note by π_+, π_- and π_0 the L^2 -orthogonal projections on E_+, E_- and E_0 respectively.

4.1.2. An admissible pair with respect to E_0 and its non-commutative cross-ratio. The splitting (E_+, E_-) is a realization of the standard splitting of $L^2(S^1, \mathbb{C})/E_0$. Following e.g. [19], two realizations of $L^2(S^1, \mathbb{C})/E_0$ in $L^2(S^1, \mathbb{C})$ can be highlighted:

- (1) H_1 is the closure of $\{f \in C^{\infty}(S^1, \mathbb{C}) \mid f(0) = 0\}$
- (2) $H_2 = \{f \in L^2(S^1, \mathbb{C}) \mid \int_{S^1} f = 0\} = E_- \oplus E_+.$ The condition $\int_{S^1} f = 0$ is equivalent to $\pi_0(f) = 0.$

Next proposition is straightforward:

Proposition 43. $((E_0, H_1), (E_0, H_2))$ is an admissible pair.

Let $i \in \mathbb{N}_2$. We note by p_i the projection on H_i with respect to E_0 . Then $q_i = Id_E - p_i$ the projection on E_0 with respect to H_i . Let $s_i = p_i - q_i$. Then,

Proposition 44.

$$DV((E_0, H_1), (E_0, H_2)) = Id_E$$

Proof. First, since the two projections on H_1 and H_2 are both with respect to E_0 , we have that $DV((E_0, H_1), (E_0, H_2)) = p_1 \circ p_2|_{H_1} = Id_{H_1}$.

 $4.1.3. \ Restricted \ Grassmannians, \ admissible \ pairs \ and \ non-commutative \ cross-ratio. \\ We \ choose \ to \ set$

- $H_+ = E_+ \oplus E_0$
- $H_- = E_-$.

All what follows holds true for other choices of splittings $H = L^2(S^1, \mathbb{C}) = H_+ \oplus H_$ with $E_{\pm} \subset H_{\pm}$. We recall that, with our notations, if $\epsilon = Pol^{-1}(H_+, H_-)$, the Pressley-Segal-Wilson restricted linear group is the group

$$GL_{res} = \{g \in GL(H) \mid [\epsilon, g] \in \tau_2\}$$

where τ_2 is the ideal of L(H) made of Hilbert-Schmidt operators

$$\tau_2 = \left\{ a \in L(H) \, | \, ||a||_2 = tr(aa^*) = \sum_{n \in \mathbb{Z}} \langle a(z^n), a(z^n) \rangle_H < +\infty \right\}.$$

The group GL_{res} is the group of the units of the algebra

$$L_{res} = \{ a \in L(H) \mid [\epsilon, a] \in \tau_2 \}$$

equipped with the norm

$$||a|| = ||a||_{L(H)} + ||[\epsilon, a]||_2.$$

As a group of the units of a Banach algebra, GL_{res} is a Banach Lie group open in L_{res} which implies that

$$\mathcal{P}_{\infty}^{(-1)}(GL_{res}) = \mathcal{P}_{\infty}(GL_{res}).$$

We then work with:

$$A = L_{res}, \quad A^* = GL_{res}$$

and

$$\mathbf{S}(A) = \left\{ g \in GL_{res} \,|\, g^2 = Id_H \right\}$$

We decompose an operator $a \in L_{res}$ blockwise with respect to $(H_+, H_-) = Pol(\epsilon)$, traditionally written as

$$a = \left(\begin{array}{cc} a_{++} & a_{+-} \\ a_{-+} & a_{--} \end{array}\right).$$

Proposition 45. Let

$$a = \left(\begin{array}{cc} a_{++} & a_{+-} \\ a_{-+} & a_{--} \end{array}\right) \in GL_{res}.$$

Then

$$\pi_{\epsilon}^{T} \left(\begin{array}{cc} a_{++} & a_{+-} \\ a_{-+} & a_{--} \end{array} \right) = \left(\begin{array}{cc} 0 & a_{+-} \\ a_{-+} & 0 \end{array} \right) \in \tau_{2}(H)$$

and

$$\pi_{\epsilon}^{N} \left(\begin{array}{cc} a_{++} & a_{+-} \\ a_{-+} & a_{--} \end{array} \right) = \left(\begin{array}{cc} a_{++} & 0 \\ 0 & a_{--} \end{array} \right) \in GL(H_{+}) \times GL(H_{-}).$$

Let us now concentrate on the splitting (H_+, H_-) and the corresponding splittings $(V, W) \in \mathbf{Spl}(L_{res})$ such $((H_+, H_-), (V, W)) \in Adm(L_{res})$. This last condition requires that $W \cap H_+ = \{0\}$ and that $V \cap H_- = \{0\}$.

Lemma 46. Let $(V, W) \in$ Spl (L_{res}) , and let $s = Pol^{-1}(V, W)$. Then

$$((H_+, H_-), (V, W)) \in Adm(L_{res}) \Leftrightarrow Ker(s_{-+}) = Ker(s_{-+}) = \{0\}.$$

Proof. The condition $W \cap H_+ = \{0\}$ is equivalent to

$$\forall x_+ \in H_+, \frac{1}{2}(x_+ + s(x_+)) = x_+ \Leftrightarrow x_+ = 0.$$

We have that

$$\frac{1}{2}(x_+ + s(x_+)) = \frac{1}{2}(x_+ + s_{++}(x_+)) + \frac{1}{2}s_{-+}(x_+).$$

We remark that $\frac{1}{2}s_{-+}(x_{+}) \in H_{-}$, which implies that s_{-+} must be injective. The same condition holds for s_{+-} , and the reciprocal statement is trivial.

By Remark 27, if $((H_+, H_-), (V, W)) \in Adm(L_{res})$, the spaces H_+ and V are isomorphic, as well as the spaces H_- and W. Let $g \in GL(H)$ (in the diffeological sense) which satisfies $(g(H_+), g(H_-)) = (V, W)$, then g must satisfy the following system:

$$\begin{cases} \frac{1}{2}(Id_{H}+s) \circ g \circ Id_{++} &= g \circ Id_{++} \\ \frac{1}{2}(Id_{H}-s) \circ g \circ Id_{--} &= g \circ Id_{--} \end{cases}$$

$$\Leftrightarrow \quad (s-Id_{H}) \circ g \circ \epsilon = 0$$

$$\Leftrightarrow \quad \begin{cases} s_{++}g_{++} + s_{+-}g_{-+} + s_{-+}g_{++} + s_{--}g_{-+} &= g_{++} + g_{-+} \\ s_{++}g_{+-} + s_{+-}g_{--} + s_{-+}g_{+-} + s_{--}g_{--} &= -g_{--} - g_{+-} \end{cases}$$

$$\Leftrightarrow \quad \begin{cases} g_{++} &= s_{++}g_{++} + s_{+-}g_{-+} \\ g_{+-} &= -s_{++}g_{+-} - s_{+-}g_{--} \\ g_{-+} &= s_{-+}g_{++} + s_{--}g_{-+} \\ g_{--} &= -s_{-+}g_{+-} - s_{--}g_{--} \end{cases}$$

Such an equation is a linear equation on spaces of linear operators L(H), understood as a L_{res} -module, for which the study is not in the scope of this paper.

4.2. On spaces of pseudo-differential operators. We denote by $DO^k(S^1), k \ge 0$, the differential operators of order less or equal than k. The algebra DO(E) is filtered by the order. It is a subalgebra of the algebra of classical pseudo-differential operators $Cl(S^1, V)$ that we describe shortly hereafter, focusing on its necessary aspects. This is an algebra that contains, for example, the square root of the Laplacian

(5)
$$|D| = \Delta^{1/2} = \int_{\Gamma} \lambda^{1/2} (\Delta - \lambda I d)^{-1} d\lambda,$$

where $\Delta = -\frac{d^2}{dx^2}$ is the positive Laplacian and Γ is a contour around the spectrum of the Laplacian, see e.g. [55, 50] for an exposition on contour integrals of pseudo-differential operators. $Cl(S^1, V)$ contains also the inverse of $Id + \Delta$, and all smoothing operators on $L^2(S^1, V)$. Among smoothing operators one can find the heat operator

$$e^{-\Delta} = \int_{\Gamma} e^{-\lambda} (\Delta - \lambda I d)^{-1} d\lambda.$$

pseudo-differential operators (maybe non-scalar) are linear operators acting on $C^{\infty}(S^1, V)$ which reads locally as

$$A(f) = \int e^{ix \cdot \xi} \sigma(x,\xi) \hat{f}(\xi) d\xi$$

where $\sigma \in C^{\infty}(T^*S^1, M_n(\mathbb{C}))$ satisfying additional estimates on its partial derivatives and \hat{f} means the Fourier transform of f. Basic facts on pseudo-differential operators defined on a vector bundle $E \to S^1$ can be found e.g. in [22].

Remark 47. Since V is finite dimensional, there exists $n \in \mathbb{N}^*$ such that $V \sim \mathcal{C}^n$. Through this identification, a pseudo-differential operator $A \in Cl(S^1, V)$ can be identified with a matrix $(A_{i,j})_{(i,j)\in\mathbb{N}^2_+}$ with coefficients

$$A_{i,j} \in Cl(S^1, \mathbb{C}).$$

In other words, the identification $V \sim \mathbb{C}^n$ that we fix induces the isomorphism of algebras

$$Cl(S^1, V) \sim M_n(Cl(S^1, \mathbb{C})).$$

This identification will remain true and useful in the successive constructions below, and will be recalled if appropriate. When it will not carry any ambiguity, we will use the notation $DO(S^1)$, $Cl(S^1)$, etc. instead of $DO(S^1, \mathbb{C})$, $Cl(S^1, \mathbb{C})$, etc. for operators acting on the space of smooth functions from S^1 to \mathbb{C} .

Pseudo-differential operators can be also described by their kernel

$$K(x,y) = \int_{\mathbb{R}} e^{i(x-y)\xi} \sigma(x,\xi) d\xi$$

which is off-diagonal smooth. Pseudo-differential operators with infinitely smooth kernel (or "smoothing" operators), i.e. that are maps: $L^2 \to C^{\infty}$ form a twosided ideal that we note by $Cl^{-\infty}(S^1, V)$. Their symbols are those which are in the Schwartz space $\mathcal{S}(T^*S^1, M_n(\mathbb{C}))$. Remarkably, we have that $Cl^0 \subset L_{res}$ and, for $a \in Cl^0$, following the notations previously defined, $a_{+-} \in Cl^{-\infty}$ and $a_{-+} \in$ $Cl^{-\infty}$ following [36, 38]. The quotient $\mathcal{F}Cl(S^1, V) = Cl(S^1, V)/Cl^{-\infty}(S^1, V)$ of the algebra of pseudo-differential operators by $Cl^{-\infty}(S^1, V)$ forms the algebra of formal pseudo-differential operators. Another algebra, which is actually known as a subalgebra of $\mathcal{F}Cl(S^1, V)$ following [?], is also called algebra of formal pseudodifferential operators. This algebra is generated by formal Laurent series

$$\Psi DO(S^1, V) = C^{\infty}(S^1, V)((\partial^{-1})) = \bigcup_{d \in \mathbb{Z}} \left\{ \sum_{k \le d} a_k \partial^k \right\}$$

where each $a_k \in C^{\infty}(S^1, M_n(\mathbb{C}))$ and $\partial = \frac{d}{dx}$. Let us precise hereafter a short but complete description of basic correspondence between $\Psi DO(S^1, V)$ and $\mathcal{F}Cl(S^1, V)$.

Symbols σ project to formal symbols and there is an isomorphism between formal pseudo-differential operators and formal symbols. A detailed study can be found in [16, Tome VII]. Classical pseudo-differential operators are operators A which associated formal symbol $\sigma(A)$ reads as an asymptotic expansion

$$\sigma(A)(x,\xi) \sim \sum_{k \in \mathbb{Z}, k \le o} \sigma_k(A)(x,\xi)$$

where the partial symbol of order k

$$\sigma_k(A): (x,\xi) \in T^*S^1 \setminus S^1 \mapsto \sigma_k(A)(x,\xi) \in M_n(\mathcal{C})$$

is k-positively homogeneous in the ξ -variable, smooth on $T^*S^1 \setminus S^1 = \{(x,\xi) \in T^*S^1 | \xi \neq 0\}$ and such that $d \in \mathbb{Z}$ is the order of the operator A. The order of a smoothing operator we put equal to $-\infty$ and the formal symbol of a smoothing operator is 0.

The set $\mathcal{F}Cl(S^1, V)$ is not the same as the space of formal operators $\Psi DO(S^1, V)$ which naturally arises in the algebraic theory of PDEs, see e.g. [?] for an overview, but here the partial symbols $\sigma_k(A)$ of $A \in \Psi DO(S^1, V)$ are k-homogeneous. By the way one only has $\Psi DO(S^1, V) \subset \mathcal{F}Cl(S^1, V)$. Two approaches for a global symbolic calculus of pseudo-differential operators have been described in [8, 62].

Notations. We shall denote note by $Cl^d(S^1, V)$ the vector space of classical pseudo-differential operators of order $\leq d$. We also denote by $Cl^*(S^1, V)$ the group of invertible in $Cl(S^1, V)$ operators. We denote the sets of formal operators adding the script \mathcal{F} . The algebra of formal pseudo-differential operators, identified wih formal symbols, is noted by $\mathcal{F}Cl(S^1, V)$, and its group of invertible element is $\mathcal{F}Cl^*(S^1, V)$, while formal pseudo-differential operators of order less or equal to $d \in \mathbb{Z}$ is noted by $\mathcal{F}Cl^d(S^1, V)$.

Through identification of $\mathcal{F}Cl(S^1, V)$ with the corresponding space of formal symbols, the space $\mathcal{F}Cl(S^1, V)$ is equipped with the natural locally convex topology inherited from the space of formal symbols. A formal symbol σ_k is a smooth function in $C^{\infty}(T^*S^1 \setminus S^1, M_n(\mathbb{C}))$ which is k- positively homogeneous (i.e. homogeneous for k > 0), and hence it can be identified with an element of $C^{\infty}(S^1, M_n(\mathbb{C}))^2$ evaluating σ_k at $\xi = 1$ and $\xi = -1$. Identifyting $\mathcal{F}Cl^d(S^1, V)$ with

$$\prod_{k \le d} C^{\infty}(S^1, M_n(\mathbb{C}))^2,$$

the vector space $\mathcal{F}Cl^d(S^1, V)$ is a Fréchet space, and hence

$$\mathcal{F}Cl(S^1, V) = \bigcup_{d \in \mathbb{Z}} \mathcal{F}Cl^d(S^1, V)$$

is a locally convex topological algebra. For our needs, it will be useful to consider it as a diffeological vector space (equiped with the nebulae diffeology).

4.2.1. Cl- and PDO- Grassmanians. Let us now consider $F = C^{\infty}(S^1, \mathbb{C}$ and

$$F_+ = F \cap H_+, \quad F_- = F \cap H_-$$

Because the Fourier base is made of smooth functions, the decomposition $F = F_+ \oplus F_-$ holds true topologically, and the full space $Cl(S^1, \mathbb{C})$ of classical, maybe unbounded, pseudo-differential operators is acting smoothly on F. The same holds for $PDO(S^1, \mathbb{C})$. Therefore, the same constructions of $\mathbf{Gr}(A)$, $\mathbf{Spl}(A)$ for $A = Cl(S^1, \mathbb{C})$ and for $A = PDO(S^1, \mathbb{C})$ hold true.

4.2.2. Splittings on spaces of formal pseudo-differential operators. Now we will introduce our main classes of classical pseudo-differential operators. We define the operator s_{KV} on $\mathcal{F}Cl(M, V)$ which extends the operator $T^*M \to T^*S^1$ defined by $(x, \xi) \mapsto (x, -\xi)$ by

$$s: \sum_{n} \sigma_n(x,\xi) \mapsto \sum_{n} (-1)^n \sigma_n\left(s(x,\xi)\right).$$

This operator obviously satisfies $s_{KV}^2 = Id$. The same way, let us recall that $\sigma(\epsilon(D)) = \xi . |\xi|^{-1}$. We define $s_D : \sum_n \sigma_n(x,\xi) \mapsto \xi . |\xi|^{-1} . \sum_n \sigma_n(x,\xi)$.

Definition 48. A formal classical pseudo-differential operator A on E is called

• odd class if and only if for all $n \in \mathbb{Z}$ and all $(x, \xi) \in T^*M$ we have:

$$\sigma_n(A)(x,-\xi) = (-1)^n \sigma_n(A)(x,\xi) \Leftrightarrow s_{KV}(A) = A.$$

We note it as $A \in \mathcal{F}Cl_{odd}(S^1, V)$.

• even class if and only if for all $n \in \mathbb{Z}$ and all $(x, \xi) \in T^*M$ we have:

$$\sigma_n(A)(x,-\xi) = (-1)^{n+1} \sigma_n(A)(x,\xi) \Leftrightarrow s_{KV}(A) = -A.$$

We note it as $A \in \mathcal{F}Cl_{even}(S^1, V)$. • We define $\mathcal{F}Cl_+(S^1, V) = Ker(Id-s_D)$ and $\mathcal{F}Cl_+(S^1, V) = Ker(Id+s_D)$.

Odd class pseudo-differential operators were introduced in [29, 30]; they are called "even-even pseudo-differential operators" in the treatise [57]. In what follows, the subscript $_{odd}$ (resp. $_{even}$) is used in an obvious way. We have the splitting

$$\mathcal{F}Cl(S^1, V) = \mathcal{F}Cl_{odd}(S^1, V) \oplus \mathcal{F}Cl_{even}(S^1, V)$$
.

 $\mathcal{F}Cl(S^1, V)$ also splits into the two ideals $\mathcal{F}Cl(S^1, V) = \mathcal{F}Cl_+(S^1, V) \oplus \mathcal{F}Cl_-(S^1, V)$. This decomposition is explicit in [28, section 4.4., p. 216], and we give an explicit description here following [36, 38].

Similar to this identification, following [46], we have other injections for $\lambda \in \mathbb{R}^*$:

$$\Phi_{\lambda,\mu}: \sum_{k\in\mathbb{Z}} a_k \left(\frac{d}{dx}\right)^k \in \Psi DO(S^1, V) \mapsto \sum_{k\in\mathbb{Z}} a_k \left(\lambda^k \left(\frac{d}{dx}\right)_+^k + \mu^k \left(\frac{d}{dx}\right)_-^k\right) \in \mathcal{F}Cl(S^1, V)$$

for $(\lambda, \mu) \in \mathcal{C}^2 \setminus \{(0; 0)\}$, with unusual convention $0^k = 0 \ \forall k \in \mathbb{Z}$. Moreover, changing a little the notations of [46], we define for $\lambda \in \mathbb{C}^*$

$$\Phi_{\epsilon(D),\lambda} : \sum_{k \in \mathbb{Z}} a_k \left(\frac{d}{dx}\right)^k \in \Psi DO(S^1, V) \mapsto \sum_{k \in \mathbb{Z}} a_k \epsilon(D) \left(\lambda \frac{d}{dx}\right)^k \in \mathcal{F}Cl(S^1, V)$$

Remark 49. We have the following identifications:

- $Im\Phi_{1,1} = \mathcal{F}Cl_{odd}(S^1, V),$
- $Im\Phi_{\epsilon,1} = \mathcal{F}Cl_{even}(S^1, V),$ $Im\Phi_{1,0} = \mathcal{F}Cl_+(S^1, V),$
- $Im\Phi_{0,1} = \mathcal{F}Cl_{-}(S^1, V).$

A direct calculation shows the following:

Lemma 50. •
$$(\mathcal{F}Cl_+(S^1, V), Im\Phi_{\lambda,\mu}) \in \mathbf{Spl}(\mathcal{F}Cl(S^1, V))$$
 if and only if $(\lambda, \mu) \in (\mathbb{C}^*)^2 - \{(\lambda, 0) \mid \lambda \in \mathbb{C}^*\}.$

- $(\mathcal{F}Cl_{-}(S^{1}, V), Im\Phi_{\lambda,\mu}) \in \mathbf{Spl}(\mathcal{F}Cl(S^{1}, V))$ if and only if $(\lambda, \mu) \in (\mathbb{C}^{*})^{2} \mathbb{C}^{*}$
- $\{(0,\mu) \mid \mu \in \mathbb{C}^*\}.$ $((\mathcal{F}Cl_+(S^1,V), \mathcal{F}Cl_-(S^1,V)), (Im\Phi_{\lambda,\lambda}, Im\Phi_{\epsilon(D),\lambda})) \in Adm(\mathcal{F}Cl(S^1,V))$ for any $\lambda \in \mathbb{C}^*$.

For $\lambda = 1$, we get the admissible couple of splittings

$$\left((\mathcal{F}Cl_+(S^1, V), \mathcal{F}Cl_-(S^1, V)), (\mathcal{F}Cl_{odd}(S^1, V), \mathcal{F}Cl_{even}(S^1, V)) \right)$$

and again a direct computation finds

Theorem 51.

 $\tilde{DV}\left((\mathcal{F}Cl_{+}(S^{1},V),\mathcal{F}Cl_{-}(S^{1},V)),(\mathcal{F}Cl_{odd}(S^{1},V),\mathcal{F}Cl_{even}(S^{1},V))\right) = 2Id_{\mathcal{F}Cl(S^{1},V)}.$

4.3. On polarizations with respect to signed measures. Let X be a possibly infinite dimensional diffeological space. Let T be a topology which is a sub-topology of the D- topology on X. Let $\sigma(T)$ be the Borel algebra generated by open subsets of the topology T. Let us recall a key property, see e.g. [18]:

Proposition 52 (Jordan decomposition theorem). Any finite signed measure μ on $\sigma(T)$ splits into

$$\mu = \mu^+ - \mu^-,$$

where μ^+ and μ^- are finite positive measures.

With this result, one can heuristically say that X can split into two subsets defined by $supp(\mu^+)$ and $supp(\mu^-)$. This decomposition is only heuristic in most cases since elementary topological arguments show that $supp(\mu^+) \cap supp(\mu^-) = \emptyset$ if and only if $supp(\mu^+)$ and $supp(\mu^-)$ are topologically disconnected, that is, they are open and closed subsets. Therefore, we have to define

Definition 53. Let μ be a finite signed measure on X. Let (X^+, X^-) be a partition of X of measurable sets. We say that this is a μ -partition if

$$supp(\mu^{\pm}) \subset X^{\pm}.$$

With the pair (X^+, X^-) at hand, we can define related symmetries and projections.

Definition 54. Let V be a diffeological vector space. The μ -partition (X^+, X^-) space of (set theoretical) maps V^X into two spaces that are the kernels $Ker(p_{X^+})$ and $Ker(p_{X^-})$ of the projectors

$$p_{X^+}: f \in V^X \mapsto \mathbf{1}_{X^+} f$$

and

$$p_{X^-}: f \in V^X \mapsto \mathbf{1}_{X^-} f,$$

where $\mathbf{1}_{X^{-}}$ and $\mathbf{1}_{X^{+}}$ are the indicator maps

$$\mathbf{1}_{X^{\pm}}(x) = \begin{cases} 1 & \text{if} \quad x \in X^{\pm} \\ 0 & \text{if} \quad x \notin X^{\pm} \end{cases}$$

We define

$$s_{(X^+,X^-)} = 2p_{X^+} - Id_{V^X}.$$

Remark 55. The terminology of μ -partition is justified by $\mu(f) = |\mu| \left(s_{(X^+, X^-)}(f) \right)$ for any bounded measurable function f.

Let us first fix a μ -partition (X^+, X^-) , and we define

$$V = Im(p_{X^+}) = \left\{ f \in V^X \, | \, \forall x \in X^-, f(x) = 0 \right\}$$

and

$$W = Im(p_{X^{-}}) = \left\{ f \in V^{X} \, | \, \forall x \in X^{+}, f(x) = 0 \right\}$$

We have the following easy property:

Proposition 56. V^X , as a cartesian product indexed by V, is a diffeological vector space for the pointwise operations, $(V, W) \in \mathbf{Spl}(V^X)$ and $(V, W) = Pol(s_{(X^+, X^-)})$.

However, we have the following "no-go" properties.

JEAN-PIERRE MAGNOT

- **Theorem 57.** Unless X is equipped with the discrete diffeology or $V \neq \{0\}$, the action of Diff(X) on V^X by composition on the right is not smooth (in fact, not continuous for the underlying D-topologies.)
 - Given two different partitions (X₁⁺, X₁⁻) and (X₂⁺, X₂⁻) associated to two (possibly identical) signed measures, the corresponding couple of splittings ((V₁, W₁), (V₂, W₂)) is never admissible.
 - The set of symetries of the type $s(X^+, X^-)$, defined for any partition (X^+, X^-) of X related to any signed finite measure, is totally disconnected in $\mathbf{S}(V^X)$.

Proof. If $V \neq \{0\}$, there exists $v \in V - \{0\}$. Let $x \in X$. We define the Dirac like function δ_x^v defined by

$$\forall x' \in , \delta_x^v(x') = \begin{cases} v & \text{if } x' = x \\ 0 & \text{if } x' \neq x \end{cases}$$

If X is not discrete, Diff(X) is not discrete, and we choose a non constant path g in Diff(X) for which we can assume without restriction that $g(0) = Id_X$. and we assume that we choose x such that there exists $t \neq 0$ such that $g(t)(x) \neq x$. Then, the smooth path $ev_x \circ \delta_x^v \circ g$ from \mathbb{R} to V takes only two values: O and v. This is impossible since the image of a connected set by a smooth map is connected. This proves the first point.

Let (X_1^+, X_1^-) and (X_2^+, X_2^-) be two different partitions of X. Since the partitions are different, we have either $X_1^+ \cap X_2^- \neq \emptyset$ or $X_2^+ \cap X_1^- \neq \emptyset$. Let $x \in X_1^+ \Delta X_2^+ =$ $(X_1^+ \cap X_2^-) \cup (X_2^+ \cap X_1^-)$. Then the Dirac function $\delta_x \in (V_1 \cap W_2) \cup (V_2 \cap W_1)$. Therefore, $V_1 \cap W_2 \neq \emptyset$ or $V_2 \cap W_1 \neq \emptyset$, which shows that $((V_1, W_2), (V_2, W_1)) \notin$ $\mathbf{Spl}(V^X)$. Therefore, $((V_1, W_1), (V_2, W_2))$ is not admissible. For the third point, we first assume that $V = \mathbb{R}$. Let us now assume that s_1 and s_2 are the two symmetries constructed from (X_1^+, X_1^-) and (X_2^+, X_2^-) and that there is a smooth path c in the set of symetries of the type $s(X^+, X^-)$, defined for any partition (X^+, X^-) of X related to any signed finite measure, such that $c(0) = s_1$ and $c(1) = s_2$. Let us consider now the evaluation map ev_x . This map is smooth on V^X , and hence the map $t \mapsto ev_x \circ c(t) \circ \delta_x$ is also smooth from \mathbb{R} to \mathbb{R} . But the image set of this map is $\{-1, 1\}$ which is disconnected. Therefore the assumption is impossible. Passing from \mathbb{R}^X to V^X , we change δ_x to δ_x^v .

Remark 58. Therefore, if one has to consider smooth structures on V^X , one has to enrich or adapt the cartesian product diffeology. Let us make a review of few apparently interesting properties.

- The cartesian product diffeology is the natural diffeology for the smoothness of the evaluation maps ev_x. Indeed, it is the pull-back diffeology ∩_{x∈X} ev^{*}P_V.
- The same way, let us consider the map

$$\Phi: (f,g) \in V^X \times Diff(X) \to f \circ g \in V^X.$$

Then $\Phi(V^X, \mathcal{P}_{Diff(X)})$ generates the smallest diffeology for which the action of Diff(X) on V^X is smooth.

These two diffeologies are completely different.

These two points suggests that we need to restrict our space of functions in order to be interesting. We now consider the space of *smooth* functions $C^{\infty}(X, V)$ instead of V^X . The space $C^{\infty}(X, V)$ is equipped with its functional diffeology. We hope

to be able to expand this approach, envolving extra-structures, in a forthcoming work.

Acknowledgements

The author would like to thank Vladimir Rubtsov for informal introductions to his works on generalizations of cross-ratios to non-commutative settings, and for explanations on his algebraic motivations.

The starting point of this work is an idea that emerged during the XL workshop on methods in Goemetry and Physics in Bialowieza, july 2023, in which the author was invited for a plenary lecture. Helpful discussions principally with Daniel Beltità, but also with Gerald Goldin, Tomasź Goliński, Tudor Ratiu and Alice Barbara Tumpach, discussions on their own research interests, convinced the author about the accuracy of this topic. May they all be acknowledged, as well as the organizers of the workshop for this invitation and this very stimulating week.

References

- Andruchow, E.; Chiumiento, E.; Di Iorio y Lucero, M. E. The compatible Grassmannian. Differ. Geom. Appl. 32, 1-27 (2014).
- [2] Andruchow, E.; Chiumiento, E.; Larotonda, G.; Geometric significance of Toeplitz kernels. J. Funct. Anal. 275 no. 2, 329-355 (2018).
- [3] Andruchow, E.; Larotonda, G.; Recht, L.; Finsler geometry and actions of the p-Schatten unitary groups. Trans. Am. Math. Soc. 362 no. 1, 319-344 (2010).
- [4] Batubenge, A.; Karshon, Y.; Watts, J.; Diffeological, Frölicher, and differential spaces. Rocky Mountain J. Math. (to appear)
- [5] Batubenge, A.; Ntumba, P.; On the way to Frölicher Lie groups Quaestionnes mathematicae (2005) 28 no1, 73–93
- [6] Beltita, D.; Ratiu, T.; Tumpach, B. A.; The restricted Grassmannian, Banach Lie-Poisson spaces, and coadjoint orbits. J. Funct. Anal. 247 no 1, 138–168 (2007)
- [7] Berger, M.; A panoramic overview of Riemannian geometry Springer (2003)
- [8] Bokobza-Haggiag, J.; Opérateurs pseudo-différentiels sur une variété différentiable; Ann. Inst. Fourier, Grenoble 19,1 125-177 (1969)
- [9] Berline, N.; Getzleer, E.; Vergne, M.; Heat Kernels and Dirac Operators Springer (2004)
- [10] Cardona, A.; Ducourtioux, C.; Magnot, J-P.; Paycha, S.; Weighted traces on pseudodifferential operators and geometry on loop groups; *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* 5 no4 503-541 (2002)
- [11] Canarutto, D.; Gauge Field Theory in Natural Geometric Language: A revisitation of mathematical notions of quantum physics Oxford University Press (2020)
- [12] K.T. Chen, Iterated Path Integrals. Bulletin AMS 83 (1977), 831-879.
- [13] Christensen, D.; Sinnamon, G.; Wu, E.; The D-topology for diffeological spaces, Pacific J. Math. (2014) 272 no1, 87—110.
- [14] Conley, C. H.; Ovsienko, V.; Lagrangian configurations and symplectic cross-ratios. Math. Ann. 375 No. 3–4, 1105–1145 (2019).
- [15] Cui, M.; Ji, G; Pencils of pairs of projections. Stud. Math. 249 no. 2, 117-141 (2019).
- [16] Dieudonné, J.; Eléments d'analyse Gauthier-Villars, Paris (1978)
- [17] Dupré, M. J.; Glazebrook, J. F.; Previato, E.; Differential algebras with Banach-algebra coefficients. II: The operator cross-ratio tau-function and the Schwarzian derivative. *Complex Anal. Oper. Theory* 7 no. 6, 1713–1734 (2013)
- [18] Fischer, T.; Existence, uniqueness, and minimality of the Jordan measure decomposition arXiv:1206.5449
- [19] Freed, D.; The Geometry of loop groups J. Diff. Geome. 28 223-276 (1988)
- [20] Frölicher, A; Kriegl, A; Linear spaces and differentiation theory (1988) Wiley series in Pure and Applied Mathematics, Wiley Interscience
- [21] Gelfand, I.; Gelfand, S.; Retakh, V.; Wilson, R.; Quasideterminants. Adv. Math. 193 no 1, 56–141 (2005)

JEAN-PIERRE MAGNOT

- [22] Gilkey, P; Invariance theory, the heat equation and the Atiyah-Singer index theorem Publish or Perish (1984)
- [23] Glöckner, H.; Algebras whose groups of units are Lie groups. Stud. Math. 153 no. 2, 147-177 (2002).
- [24] Glöckner, H.; Neeb, K-H.; When unit groups of continuous inverse algebras are regular Lie groups. Stud. Math. 211 no. 2, 95-109 (2012).
- [25] Goldammer, N.; Magnot, J-P.; Welker, K.; On diffeologies from infinite dimensional geometry to PDE constrained optimization. Magnot, Jean-Pierre (ed.), Recent advances in diffeologies and their applications. AMS-EMS-SMF special session, Université de Grenoble-Alpes, Grenoble, France, July 18–20, 2022. Contemp. Math. 794 1-48 (2024).
- [26] Goliński, T.; Odzijewicz, A.; Hierarchy of Hamilton equations on Banach Lie-Poisson spaces related to restricted Grassmannian. J. Funct. Anal. 258 no. 10, 3266-3294 (2010).
- [27] Iglesias-Zemmour, P. Diffeology Mathematical Surveys and Monographs 185 AMS (2013).
- [28] Kassel, Ch.; Le résidu non commutatif (d'après M. Wodzicki) Séminaire Bourbaki, Vol. 1988/89. Astérisque 177-178, Exp. No. 708, 199-229 (1989)
- [29] Kontsevich, M.; Vishik, S.; Determinants of elliptic pseudo-differential operators Max Plank Institut fur Mathematik, Bonn, Germany, preprint n. 94-30 (1994)
- [30] Kontsevich, M.; Vishik, S.; Geometry of determinants of elliptic operators. Functional analysis on the eve of the 21st century, Vol. 1 (New Brunswick, NJ, 1993), Progr. Math. 131,173-197 (1995)
- [31] Kriegl, A.; Michor, P.W.; The convenient setting for global analysis (1997); AMS Math. Surveys and Monographs 53, AMS, Providence
- [32] Labourie, F., What is a Cross Ratio. Notices of the AMS 55 no 10, 1234-1235 (2008)
- [33] Labourie, F.; McShane, G.; Cross ratios and identities for higher Teichmüller-Thurston theory. Duke Math. J. 149 no. 2, 279-345 (2009).
- [34] Laubinger, M.; A Lie algebra for Frölicher groups Indag. Math. 21 no 3-4, 156–174 (2011)
- [35] Leslie, J.; On a Diffeological Group Realization of certain Generalized symmetrizable Kac-Moody Lie Algebras J. Lie Theory 13 (2003), 427-442.
- [36] Magnot, J-P.; The Kähler form on the loop group and the Radul cocycle on Pseudo-differential Operators; GROUP'24: Physical and Mathematical aspects of symmetries, Proceedings of the 24th International Colloquium on Group Theorical Methods in Physics, Paris, France, 15-20 July 2002; Institut of Physic conferences Publishing **173**, 671-675, IOP Bristol and Philadelphia (2003)
- [37] Magnot, J-P.; Chern forms on mapping spaces, Acta Appl. Math. 91, no. 1, 67-95 (2006).
- [38] Magnot, J-P.; Renormalized traces and cocycles on the algebra of S¹-pseudo-differential operators; Lett. Math. Phys. 75 no2, 111-127 (2006)
- [39] Magnot, J-P.; Difféologie du fibré d'Holonomie en dimension infinie, C. R. Math. Soc. Roy. Can. 28 no4 (2006) 121-127.
- [40] Magnot, J-P.; The Schwinger cocycle on algebras with unbounded operators. Bull. Sci. Math. 132, no. 2, 112-127 (2008).
- [41] Magnot, J-P.; Ambrose-Singer theorem on diffeological bundles and complete integrability of KP equations. Int. J. Geom. Meth. Mod. Phys. 10, no 9 (2013) Article ID 1350043.
- [42] Magnot, J-P.; The group of diffeomorphisms of a non-compact manifold is not regular Demonstr. Math. 51, No. 1, 8-16 (2018)
- [43] Magnot, J-P.; Reyes, E. G.; Well-posedness of the Kadomtsev-Petviashvili hierarchy, Mulase factorization, and Frölicher Lie groups Ann. Henri Poincaré 21, No. 6, 1893-1945 (2020)
- [44] Magnot, J-P. Reyes, E.G.; The Cauchy problem of the Kadomtsev-Petviashvili hierarchy and infinite-dimensional groups. in Nonlinear Systems and Their Remarkable Mathematical Structures, Volume 2; Norbert Euler and Maria Clara Nucci Editors, CRC press (2019) section B6
- [45] Magnot, J-P.; On the geometry of $Diff(S^1)$ -pseudodifferential operators based on renormalized traces. Proceedings of the International Geometry Center 14 No. 1, 19-48 (2021)
- [46] Magnot, J-P.; Rubtsov, V.; On the Kadomtsev-Petviashvili hierarchy in an extended class of formal pseudo-differential operators *Theoret. Math. Phys.* 207, No. 3, 799-826 (2021)
- [47] Mickelsson, J.; *Current algebras and groups*. Plenum monographs in Nonlinear Physics, Springer (1989)
- [48] Mickelsson, J.; Wodzicki residue and anomalies on current algebras Integrable models and strings A. Alekseev and al. eds. Lecture notes in Physics 436, Springer (1994)

- [49] Omori, H.; Infinite Dimensional Lie Groups (1997) AMS Translations of Mathematical Monographs no 158 Amer. Math. Soc., Providence, R.I.
- [50] Paycha, S; Regularised integrals, sums and traces. An analytic point of view. University Lecture Series 59, AMS (2012).
- [51] Pervova, E.; Finite-dimensional diffeological vector spaces being and not being coproducts. Magnot, Jean-Pierre (ed.), Recent advances in diffeologies and their applications. AMS-EMS-SMF special session, Université de Grenoble-Alpes, Grenoble, France, July 18–20, 2022. Contemp. Math. 794, 129-138 (2024).
- [52] Pressley, A.; Segal, G.; Loop Groups Oxford Univ. Press (1988)
- [53] Retakh, V.; Noncommutative cross-ratios. J. Geom. Phys. 82 13-17 (2014).
- [54] Retakh, V.; Rubtsov, V.; Sharygin, G.; The non-commutative cross-ratio and Schwartz derivatives. in:Donagi, Ron (ed.) et al., Integrable systems and algebraic geometry. A celebration of Emma Previato's 65th birthday. Volume 2. Lond. Math. Soc. Lect. Note Ser. 459, 499-528 (2020).
- [55] Seeley, R.T.; Complex powers of an elliptic operator AMS Proc. Symp. Pure Math. 10, 288-307 (1968)
- [56] Schwinger, J.; Field theory of commutators; Phys. Rev. Lett. 3, 296-297 (1959)
- [57] Scott, S.; Traces and determinants of pseudodifferential operators; OUP (2010)
- [58] Souriau, J-M.; un algorithme générateur de structures quantiques Astérisque (hors série) 341-399 (1985)
- [59] Shi, W.; Ji, G.; Anti-selfadjoint operators as commutators of projections. J. Math. Anal. Appl. 478 no. 2, 539-559 (2019).
- [60] Tumpach, A.B.; Banach Poisson-Lie groups and Bruhat-Poisson structure of the restricted Grassmannian. Comm. Math. Phys. 373 no. 3, 795–858 (2020)
- [61] Watts, J.; Diffeologies, differentiable spaces and symplectic geometry. University of Toronto, PhD thesis (2013). arXiv:1208.3634v1.
- [62] Widom, H.; A complete symbolic calculus for pseudo-differential operators; Bull. Sc. Math. 2e serie 104 19-63 (1980)
- [63] Wu, E.; A survey on diffeological vector spaces and applications. Magnot, Jean-Pierre (ed.), Recent advances in diffeologies and their applications. AMS-EMS-SMF special session, Université de Grenoble-Alpes, Grenoble, France, July 18–20, 2022. Contemp. Math. 794, 113–128 (2024)
- [64] Zelikin, M. I.; Geometry of operator cross-ratio. Sb. Math. 197 no 1, 37-51 (2006)
- J.-P.M.: UNIV. ANGERS, CNRS, LAREMA, SFR MATHSTIC, F-49000 ANGERS, FRANCE, AND, LYCÉE JEANNE D'ARC, AVENUE DE GRANDE BRETAGNE,, 63000 CLERMONT-FERRAND, FRANCE Email address: magnot@math.cnrs.fr