

Generic differentiability and P -minimal groups

Will Johnson

April 29, 2024

Abstract

We prove generic differentiability in P -minimal theories, strengthening an earlier result of Kuijpers and Leenknegt. Using this, we prove Onshuus and Pillay’s P -minimal analogue of Pillay’s conjectures on o-minimal groups. Specifically, let G be an n -dimensional definable group in a highly saturated model M of a P -minimal theory. Then there is an open definable subgroup $H \subseteq G$ such that H is compactly dominated by H/H^{00} , and H/H^{00} is a p -adic Lie group of the expected dimension.

1 Introduction

Let K be a p -adic field, i.e., a finite extension of \mathbb{Q}_p . An expansion T of $\text{Th}(K)$ is P -minimal in the sense of Haskell and Macpherson [12] if for every model $M \models T$, every 1-variable definable set $D \subseteq M^1$ is already definable in the reduct $M \upharpoonright \mathcal{L}_{rings}$. Haskell and Macpherson show that P -minimality has many properties similar to o-minimality, such as a good dimension theory and cell decomposition theorem. There are non-trivial examples of P -minimal theories arising from restricted analytic functions [7].

One of the main results of this paper is the following generic differentiability theorem:

Theorem 1.1. *Let M be a model of a P -minimal theory T . Let $U \subseteq M^n$ be a non-empty definable open set. Let $f : U \rightarrow M^m$ be a definable function. Then there is a definable open set $U_0 \subseteq U$ such that $\dim(U \setminus U_0) < n$ and f is differentiable on U_0 .*

In fact, we even get generic *strict* differentiability. See Definition 5.1 and Theorem 5.14. We also prove an inverse function theorem for strictly differentiable functions, using the standard proof (Theorem 5.18).

The generic differentiability theorem generalizes work of Kuijpers and Leenknegt [18, Theorem 1.8], who proved generic differentiability in the special case where T is “strictly P -minimal”, meaning that T is the theory of some P -minimal expansion of the p -adic field K . Most P -minimal theories arising in practice are strictly P -minimal. However, there are many simple examples of P -minimal theories which fail to be strictly P -minimal, so it is better to have generic differentiability without any additional assumptions.

We apply generic differentiability and the inverse function theorem to the study of definable groups in P -minimal structures. Recall that a definable group G in a monster model \mathbb{M}

is *compactly dominated* if G^{00} exists and the quotient map $f : G \rightarrow G/G^{00}$ has the property that for any definable set $D \subseteq G$, we can partition G/G^{00} into three sets D_0, D_1, D_{err} such that

$$\begin{aligned} f(x) \in D_0 &\implies x \notin D \\ f(x) \in D_1 &\implies x \in D \\ D_{\text{err}} &\text{ has Haar measure 0.} \end{aligned}$$

Onshuus and Pillay conjectured the following in [20], by analogy with Pillay's conjectures on o-minimal groups:

Conjecture 1.2 (Onshuus-Pillay). *Let G be an n -dimensional definable group in a highly saturated P -minimal structure M . Then there is an n -dimensional definable open subgroup $H \subseteq G$ with the following properties:*

1. H/H^{00} is isomorphic to an n -dimensional Lie group over K .
2. H is compactly dominated by H/H^{00} .

Here K is the finite extension of \mathbb{Q}_p whose theory T expands. We prove Conjecture 1.2, and in fact prove something a little stronger:

Theorem 1.3 (Theorem 8.4). *Let G be an n -dimensional definable group in a highly saturated P -minimal structure M . Then there is an n -dimensional definable open subgroup $H \subseteq G$ with the following properties:*

1. H is compactly dominated by H/H^{00} .
2. H/H^{00} is isomorphic to $(\mathcal{O}_K, +)^n$, where \mathcal{O}_K is the ring of integers in K . In particular, H/H^{00} is isomorphic to an n -dimensional Lie group over K .

The proof strategy is a bit idiotic: one simply takes H to be an infinitesimally small neighborhood of 0 and checks that everything works. Generic differentiability helps control the structure of H , showing that $(H, +)$ looks like a deformation of $(\mathcal{O}^n, +)$.

1.1 Variants of Theorem 1.3

While Theorem 1.3 *technically* resolves the Onshuus-Pillay conjecture, it certainly goes against the spirit of the conjecture—the intention was that H would be close to G , and the p -adic Lie group H/H^{00} should have interesting structure related to the structure of G and H . But the H in Theorem 1.3 is much smaller than G , and the group H/H^{00} has a boring structure solely determined by $\dim(G)$.

In fact, we are essentially exploiting a loophole in Onshuus and Pillay's specific formulation of their conjecture. To be more precise, I do *not* know how to prove the following variant conjecture:

Conjecture 1.4 (Modified Onshuus-Pillay conjecture). *Let G be an n -dimensional definable group in a highly saturated P -minimal structure M . Let $M_0 \prec M$ be a small model defining G . Then there is an n -dimensional M_0 -definable open subgroup $H \subseteq G$ with the following properties:*

1. H/H^{00} is isomorphic to an n -dimensional Lie group over K .
2. H is compactly dominated by H/H^{00} .

(This is the same as Conjecture 1.2, except we are now requiring H to be defined over the same parameters that define G .) The best I could prove was the following two theorems, each of which has a drawback:

Theorem 1.5 (Theorem 8.5). *In the setting of Conjecture 1.4, there is an M_0 -definable open subgroup $H \subseteq G$ such that H is compactly dominated by H/H^{00} , and H/H^{00} is isomorphic to an $n \cdot [K : \mathbb{Q}_p]$ -dimensional Lie group over \mathbb{Q}_p .*

So the case where $K = \mathbb{Q}_p$ is completely handled, but when $[K : \mathbb{Q}_p] > 1$ we get a slightly defective result, only getting a Lie group over \mathbb{Q}_p rather than a Lie group over K . The proof of Theorem 1.5 depends on the following deep results of Lazard [19]:

- (i) If G is a Lie group over \mathbb{Q}_p , then the topological group structure on G determines the Lie group structure.
- (ii) If G is an abstract topological group, then G is a Lie group over \mathbb{Q}_p if and only if G satisfies a certain abstract group-theoretic condition (see [9, Theorem 8.1] for details).

If we replace \mathbb{Q}_p with a finite extension K , then (i) certainly fails. This makes (ii) seem more unlikely, at least to a non-expert like me. Part (ii) is the specific thing we would need to generalize Theorem 1.5 to resolve Conjecture 1.4.

Using very different techniques, we also resolve Conjecture 1.4 in the special case where T is the pure theory of p -adically closed fields $\text{Th}(K)$:

Theorem 1.6 (Theorem 9.23). *Conjecture 1.4 holds if T is $\text{Th}(K)$, the theory of pure p -adically closed fields.*

The proof depends heavily on the fact that definable functions on K are piecewise analytic [22, Theorem 1.1], something which has no analogue in a general P -minimal theory. Consequently, I do not expect the proof of Theorem 1.6 to usefully generalize.

1.2 The proof of generic differentiability

In proving generic differentiability (Theorem 1.1), the key technical tool is the “definable compactness” of closed bounded sets (Proposition 2.6). Definable compactness was essentially proved by Cubides-Kovacsics and Delon [6], but we give a slightly improved proof in Section 2. The improved version isn’t strictly necessary (Remark 2.9), but makes the subsequent proofs more streamlined.

To a first approximation, definable compactness ensures that the limit

$$\lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}$$

exists, though it might take the value ∞ and might only be a “one-sided limit”—the limit might depend on which “direction” x approaches a from. To proceed further, we analyze the asymptotic behavior of definable functions $f(x)$ as x approaches 0. In Section 4, we show that any definable function behaves asymptotically like

$$f(x) \propto Cx^q \text{ as } x \rightarrow 0 \quad (*)$$

for some constant C and rational number q , though again $(*)$ may only hold “on one side” of 0. To show this, we first analyze the valuation $v(f(x))$, relying on Cluckers’ theorem that the value group is a pure model of Presburger arithmetic [3, Theorem 6]. After controlling the valuation, we use definable compactness to get the more precise statement $(*)$.

To prove generic differentiability, one reduces to the case where

$$f(a+x) - f(a) \propto C(a)x^q \text{ as } x \rightarrow 0$$

for any a in the domain of f , for some fixed q independent of a . By carefully analyzing the behavior of f , one can show that q must be 1,¹ and the asymptotic behavior of $f(a+x) - f(a)$ is independent of which direction x approaches 0 from. This gives differentiability.

1.3 Conventions

We write group operations as $x \star y$, since we will frequently need to distinguish from both addition $x + y$ and multiplication $x \cdot y$. We write inverses as x^{-1} however. (Usually it is clear from context whether we mean the group inverse or the multiplicative inverse.)

We generally work in the following setting: K is \mathbb{Q}_p or a finite extension, $\text{Th}(K)$ is the theory of K in the language of rings \mathcal{L}_{Rings} , T is a P -minimal expansion of $\text{Th}(K)$, and \mathbb{M} is a highly saturated monster model of T . If M is an elementary substructure of \mathbb{M} , or more generally of $\mathbb{M} \restriction \mathcal{L}_{Rings}$, then \mathcal{O}_M , \mathfrak{m}_M , k_M , and Γ_M denote the valuation ring, maximal ideal, residue field, and value group, respectively. When no subscript is given, assume $M = \mathbb{M}$. The valuation is written as $v(x)$ with the additive conventions, so that

$$\begin{aligned} v(xy) &= v(x) + v(y) \\ v(x+y) &\geq \min(v(x), v(y)). \end{aligned}$$

The ball of radius γ around a , written $B_\gamma(a)$ is the *closed* ball $\{x \in \mathbb{M} : v(x - a) \geq \gamma\}$.

The *radius* of $B_\gamma(a)$ is γ . We write the radius of a ball B as $\text{rad}(B)$. The *parent* of $B_\gamma(a)$ is $B_{\gamma-1}(a)$. We say that B is a *child* of B' if B' is the parent of B . Two balls are *siblings* if

¹For example, you cannot have a definable function f such that $f(a+x) - f(a) \propto x^2$ at every point a . Contrast this with $\text{ACVF}_{2,2}$, where the function $f(x) = x^2$ *does* have the asymptotic expansion $f(a+x) = f(a) + x^2$.

they have the same parent. We say that B is an *ancestor* of B' and B' is a *descendant* of B if $B \supseteq B'$.

If $\bar{x} = (x_1, \dots, x_n)$ is a vector, then $v(\bar{x})$ denotes $\min(v(x_1), v(x_2), \dots, v(x_n))$.

We review the definition of differentiability and strict differentiability in Section 5.1.

We let P_n denote the set of non-zero n th powers, and $\overline{P_n}$ denote its closure, which is $P_n \cup \{0\}$.

2 Definable compactness

If M is any structure and X is a definable topological space in M , say that X is *definably compact* if for any definable downward-directed family \mathcal{F} of closed non-empty subsets of X , the intersection $\bigcap \mathcal{F}$ is non-empty. More generally, a definable subset $D \subseteq X$ is definably compact if D is definably compact with respect to the induced subspace topology. This notion was investigated independently by Fornasiero [11] and the author [15, §3.1].

Definable compactness has the following good properties, proven in [15, §3.1], though only (4) is non-trivial.

Fact 2.1 ([15, §3.1]).

1. If X is Hausdorff and $D \subseteq X$ is definably compact, then D is closed.
2. If X is definably compact, any closed subset $X \subseteq D$ is definably compact.
3. If D is compact, then D is definably compact.
4. A direct product of two definably compact sets is definably compact, with respect to the product topology.
5. A finite set is definably compact.
6. A finite union of definably compact sets is definably compact.
7. The image of a definably compact set under a continuous function is definably compact.
8. Definable compactness is preserved in elementary extensions: if $N \succeq M$, then $D(M)$ is definably compact iff $D(N)$ is definably compact.

Return to the setting of a P -minimal monster model \mathbb{M} .

Lemma 2.2. *The valuation ring \mathcal{O} is definably compact. In other words, any definable downward-directed family \mathcal{F} of closed non-empty subsets of \mathcal{O} has $\bigcap \mathcal{F} \neq \emptyset$.*

This is *almost* the same thing as Cubides-Kovacsics and Delon's Theorem (A) [6], but slightly more general as we do not require \mathcal{F} to be nested (or indexed by Γ). Luckily, it is not hard to deduce Lemma 2.2 from their work.

Proof. Suppose for the sake of contradiction that $\bigcap \mathcal{F} = \emptyset$. Say that a ball $B \subseteq \mathcal{O}$ is *good* if B intersects every $C \in \mathcal{F}$. Otherwise, say that B is *bad*. The following properties are straightforward:

1. The ball \mathcal{O} is good.
2. Every descendant of a bad ball is bad; every ancestor of a good ball is good.
3. A ball is good if and only if one of its children is good. This follows by directedness of \mathcal{F} , and the fact that each ball has only finitely many children. In particular, every good ball has a good child.
4. If $a \in \mathcal{O}$, then some ball $B_\gamma(a)$ is bad. Otherwise, every ball around a intersects every $C \in \mathcal{F}$. As the sets in \mathcal{F} are closed, this means $a \in \bigcap \mathcal{F}$, a contradiction.

Let $\Theta \subseteq \Gamma$ be the set $\{\text{rad}(B) : B \subseteq \mathcal{O}, B \text{ is good}\}$. By (1), Θ is non-empty. By (2), Θ is downward-closed (in $\Gamma_{\geq 0}$). By (3), Θ has no greatest element. The value group Γ is a pure model of Presburger arithmetic [3, Theorem 6], so it is definably complete. By definable completeness of Γ , Θ must be cofinal in Γ , and so $\Theta = \Gamma_{\geq 0}$. In particular, there are arbitrarily small good balls (good balls of arbitrarily high radius).

Let $f : \mathcal{O} \rightarrow \Gamma$ be the function

$$f(a) = \max\{\gamma \in \Gamma : B_\gamma(a) \text{ is good}\}.$$

The set inside the maximum is non-empty by (1) and bounded above by (4), so the maximum exists by definable completeness of Γ . Spiritually, the function $f(x)$ is measuring the distance from x to $\bigcap \mathcal{F}$.

The function f is locally constant. In fact, if $a \in \mathcal{O}$ and $\gamma_0 = f(a) + 1$, then f is constant on the ball $B_{\gamma_0}(a)$. Indeed, if $b \in B_{\gamma_0}(a)$, then

$$\begin{aligned} B_{\gamma_0}(b) &= B_{\gamma_0}(a) \text{ is bad} \\ B_{\gamma_0-1}(b) &= B_{\gamma_0-1}(a) \text{ is good,} \end{aligned}$$

and so $f(b) = \gamma_0 - 1 = f(a)$.

By Fact 2.3 below, f has a maximum value γ_0 on the set \mathcal{O} . Then every ball of radius $> \gamma_0$ is bad, contradicting the fact that Θ is cofinal in Γ . \square

Fact 2.3. *Let $f : \mathcal{O} \rightarrow \Gamma$ be a definable function which is continuous, i.e., locally constant. Then f has a maximum.*

Fact 2.3 is an instance of Theorem (B) in [6]. The proof in [6] seems somewhat complicated, so we give an alternate proof, which may be of independent interest. First we make a couple remarks.

Remark 2.4. If $\Delta \subseteq \Gamma$ is definable, say that an integer $n < \omega$ is a *modulus* for Δ if $\Delta \cap (k + n\Gamma)$ is a finite union of intervals in $k + n\Gamma$ for $k = 0, 1, 2, \dots, n-1$.

1. Quantifier elimination in Presburger arithmetic implies that every definable $\Delta \subseteq \Gamma$ has a modulus. This is related to the quasi-o-minimality of Presburger arithmetic.
2. By compactness, the following is true: if $\{\Delta_i\}_{i \in I}$ is a definable family of subsets of Γ , then there is an integer n which is uniformly a modulus for every set Δ_i .
3. If Δ is infinite and n is a modulus for Δ , then $\{\gamma, \gamma + n\} \subseteq \Delta$ for some γ . That is, Δ must contain two consecutive elements of $k + n\Gamma$ for some k . Otherwise, $\Delta \cap (k + n\Gamma)$ is a finite union of points for every k , and Δ is finite.

Remark 2.5. Let D be a finite set and let $f : \mathcal{O} \rightarrow D$ be a definable function which is continuous (i.e., locally constant). Then f is uniformly continuous: there is γ such that f is constant on every ball of radius γ . To see this, note that f is definable in the reduct $M \upharpoonright \mathcal{L}_{rings}$ by P -minimality, so we can reduce to the base theory $\text{Th}(K)$. Then we can transfer the statement to the elementarily equivalent model K , where it holds by compactness of \mathcal{O}_K .

Proof (of Fact 2.3). Let $f : \mathcal{O} \rightarrow \Gamma$ be locally constant. If f is bounded above, then a maximum exists by definable completeness of Γ . Assume f is unbounded, i.e., $f(\mathcal{O})$ is cofinal in Γ . Say a ball $B \subseteq \mathcal{O}$ is *good* if f is unbounded on B , and *bad* otherwise. Then \mathcal{O} is good, and every good ball has a good child. As in the proof of Lemma 2.2, there are arbitrarily small good balls, i.e., good balls of radius $> \gamma$ for any γ .

Let n be a uniform modulus for the sets

$$f(B) = \{f(x) : x \in B\},$$

as B ranges over descendants of \mathcal{O} . Let $g(x)$ be $f(x) \bmod 2n$. Then g is a continuous function from \mathcal{O} to the finite set $\Gamma/2n\Gamma$. By Remark 2.5, g is uniformly continuous: there is γ_0 such that

$$v(a - b) > \gamma_0 \implies g(a) = g(b) \iff f(a) \equiv f(b) \pmod{2n}.$$

Take a good ball B of radius $> \gamma_0$. Then g is constant on B . Because B is good, the set $f(B)$ is cofinal in Γ , hence infinite. As n is a modulus for $f(B)$, there are $\gamma, \gamma + n$ in $f(B)$ by Remark 2.4(3). That is, there are $a, b \in B$ such that

$$f(a) + n = f(b).$$

Then $f(a) \not\equiv f(b) \pmod{2n}$, a contradiction. □

Proposition 2.6. *Let D be a definable subset of \mathbb{M}^n . Then D is definably compact if and only if D is closed and bounded.*

Proof. If D is not closed or not bounded, it is easy to see that D is not definably compact. Indeed, the proof from [17, Lemma 2.4] works.

Conversely, suppose D is closed and bounded. Then D is a closed subset of an n -dimensional closed ball $B \subseteq \mathbb{M}^n$. There is a definable homeomorphism between B and \mathcal{O}^n . By the properties of definable compactness discussed at the start of this section

$$\begin{aligned} \mathcal{O} \text{ is definably compact} &\implies \mathcal{O}^n \text{ is definably compact} \\ \implies B \text{ is definably compact} &\implies D \text{ is definably compact.} \end{aligned} \quad \square$$

2.1 Applications of definable compactness

Corollary 2.7. *Let $D \subseteq \mathbb{M}^n$ be closed and bounded. Let $f : D \rightarrow \mathbb{M}^m$ be definable and continuous. Then f is uniformly continuous.*

Proof. Otherwise, there is some $\epsilon \in \Gamma$ such that for every $\delta \in \Gamma$, there are $\bar{a}, \bar{b} \in D$ with

$$v(\bar{a} - \bar{b}) \geq \delta \text{ but } v(f(\bar{a}) - f(\bar{b})) \leq \epsilon.$$

Here, $v(\bar{a})$ is short for $\min_{1 \leq i \leq n} v(a_i)$.

Fix such an ϵ . For each δ , let

$$C_\delta = \{(\bar{a}, \bar{b}) \in D^2 : v(\bar{a} - \bar{b}) \geq \delta \text{ and } v(f(\bar{a}) - f(\bar{b})) \leq \epsilon\}.$$

Then C_δ is closed in D^2 , by continuity of f . By assumption, C_δ is non-empty. The family $\{C_\delta\}_{\delta \in \Gamma}$ is nested, so by definable compactness of D^2 , there is some $(\bar{a}, \bar{b}) \in \bigcap_\delta C_\delta$. Then

$$v(\bar{a} - \bar{b}) = +\infty \text{ and } v(f(\bar{a}) - f(\bar{b})) \leq \epsilon,$$

which is absurd. \square

Let $f : D \rightarrow C$ be a definable function, for some definable sets $D \subseteq \mathbb{M}^n$ and $C \subseteq \mathbb{M}^m$. Let x_0 be a point in the frontier $\partial D = \overline{D} \setminus D$. Say that $b \in \mathbb{M}^m$ is a *cluster point* of $f(x)$ as $x \rightarrow x_0$ if for every $\epsilon \in \Gamma$, there is $x \in B_\epsilon(x_0) \cap D$ such that $f(x) \in B_\epsilon(b)$. Equivalently, (x_0, b) is in the closure of the graph of f .

Lemma 2.8. *In the above setting, suppose C is definably compact.*

1. *There is at least one cluster point as $x \rightarrow x_0$.*
2. *$\lim_{x \rightarrow x_0} f(x)$ exists if and only if there is a unique cluster point b , in which case $\lim_{x \rightarrow x_0} f(x) = b$.*

Proof. 1. For $\epsilon \in \Gamma$, let G_ϵ be the graph of f restricted to $B_\epsilon(x_0) \cap D$. By definable compactness², there is a point $(a, b) \in \bigcap_\epsilon \overline{G_\epsilon}$. For any ϵ we have

$$\begin{aligned} G_\epsilon &\subseteq B_\epsilon(x_0) \times \mathbb{M}^m \\ \overline{G_\epsilon} &\subseteq B_\epsilon(x_0) \times \mathbb{M}^m \\ a &\in B_\epsilon(x_0). \end{aligned}$$

²of $B_\epsilon(x_0) \times C$

Thus $a = x_0$. Fix any ϵ . If G is the graph of f , then

$$(x_0, b) = (a, b) \in \overline{G_\epsilon} \subseteq \overline{G},$$

and so b is a cluster point.

2. First suppose $\lim_{x \rightarrow x_0} f(x)$ exists and equals b . Then $(x_0, b) = \lim_{x \rightarrow x_0} (x, f(x))$, and so (x_0, b) is in the closure of the graph of f and b is a cluster point. Suppose for the sake of contradiction that b' is another cluster point. Take ϵ so large that $B_\epsilon(b') \cap B_\epsilon(b) = \emptyset$. By existence of the limit, there is $\delta > \epsilon$ such that

$$x \in B_\delta(x_0) \cap D \implies f(x) \in B_\epsilon(b).$$

As b' is a cluster point, there is $x \in B_\delta(x_0) \cap D$ such that $f(x) \in B_\delta(b') \subseteq B_\epsilon(b')$, contradicting the fact that $f(x) \in B_\epsilon(b)$. This shows that b is the unique cluster point.

Conversely, suppose that b is the unique cluster point. We claim that $\lim_{x \rightarrow x_0} f(x) = b$. Otherwise, there is some ϵ such that for any δ , there is $x \in B_\delta(x_0) \cap D$ with $f(x) \notin B_\epsilon(b)$. Fix such an ϵ . For each δ , let G_δ be the non-empty definable set

$$G_\delta = \{(x, f(x)) : x \in B_\delta(x_0) \cap D, f(x) \notin B_\epsilon(b)\}.$$

By definable compactness, there is some $(a, b') \in \bigcap_\delta \overline{G_\delta}$. As in part (1), $a = x_0$ and b' is a cluster point of f . The set G_δ is contained in the closed set $\mathbb{M}^n \times (\mathbb{M}^m \setminus B_\epsilon(b))$ for each δ , so we must have $b' \notin B_\epsilon(b)$. Then $b' \neq b$, contradicting the fact that b is the unique cluster point. \square

Remark 2.9. A Γ -family is a definable family of non-empty sets of the form $\{D_\gamma\}_{\gamma \in \Gamma}$, such that $\gamma < \gamma' \implies D_\gamma \subseteq D_{\gamma'}$. Say that a definable topological space X is Γ -compact if every Γ -family of closed sets has non-empty intersection. This condition is slightly weaker than our definition of definable compactness.

In principle, we could use Γ -compactness rather than definable compactness, making Lemma 2.2 unnecessary. Theorem (A) of [6] says that closed bounded definable subsets of \mathbb{M} are Γ -compact. The applications of definable compactness, such as Corollary 2.7, only use Γ -compactness. Lastly, there is an analogue of Fact 2.1 for Γ -compactness. Unfortunately, this analogue is a bit clumsy—one needs to restrict to definable topological spaces X that are “definably first countable” in the sense that every point $a \in X$ has a Γ -family neighborhood basis.³ This isn’t a problem in our case, because the spaces \mathbb{M}^n are definably first countable.

3 More tools

3.1 Chains of nowhere dense sets

Lemma 3.1. *Let $\{D_\gamma\}_{\gamma \in \Gamma}$ be a Γ -family of subsets of \mathbb{M}^k , in the sense of Remark 2.9. Suppose every D_γ is nowhere dense. Then the union $\bigcup_{\gamma \in \Gamma} D_\gamma$ is also nowhere dense.*

³In the base theory $\text{Th}(K)$ of p -adically closed fields, every definable (or interpretable) topological space is definably first countable, and in fact Γ -compactness agrees with definable compactness [1, Theorem 8.11]. I don’t know whether this continues to hold in P -minimal expansions.

Proof. By dimension theory, the following are equivalent for a definable set $D \subseteq \mathbb{M}^k$:

1. $\dim(D) = k$.
2. D has non-empty interior.
3. $\dim(\overline{D}) = k$.
4. \overline{D} has non-empty interior.

Thus D is nowhere dense iff $\dim(D) < k$. We can rephrase the lemma as follows:

If $\{D_\gamma\}$ is a Γ -family of subsets of \mathbb{M}^k and $\bigcup_{\gamma \in \Gamma} D_\gamma$ has dimension k , then some D_γ has dimension k .

In fact, this holds in any dp-minimal theory. By Fact 3.2 below, the condition $\dim\left(\bigcup_{\gamma \in \Gamma} D_\gamma\right) = k$ means that there are infinite subsets $S_1, \dots, S_k \subseteq \mathbb{M}$ such that

$$\bigcup_{\gamma \in \Gamma} D_\gamma \supseteq \prod_{i=1}^k S_i.$$

We may assume the S_i are countable (or merely small). Then by saturation, we can take γ sufficiently large that

$$D_\gamma \supseteq \prod_{i=1}^k S_i.$$

Again, by Fact 3.2, this makes D_γ have dimension k . □

Fact 3.2. *If \mathbb{M} is a highly saturated model of a dp-minimal theory T , and $D \subseteq \mathbb{M}^n$ is definable, then the following are equivalent:*

1. $\text{dp-rk}(D) = n$.
2. *There are countable infinite sets $S_1, \dots, S_n \subseteq \mathbb{M}$ such that $S_1 \times S_2 \times \dots \times S_n \subseteq D$.*

This is implicit in work of Simon [23], though I am having trouble finding a precise reference. At any rate, it is explicit in [16, Theorem 3.25].

3.2 Strengthening limits

Lemma 3.3. *Let $U \subseteq \mathbb{M}^k$ be open, definable, and non-empty. Let $X \subseteq \mathbb{M}^k \times \mathbb{M}^\times$ be a definable set such that $U \times \{0\} \subseteq \overline{X}$. Then there is an open definable non-empty subset $U_0 \subseteq U$ such that $0 \in \overline{X_a}$ for every $a \in U_0$, where $X_a = \{b \in \mathbb{M}^\times : (a, b) \in X\}$.*

Proof. For any $\epsilon \in \Gamma$, let D_ϵ be the set of $a \in U$ such that $B_\epsilon(0) \cap X_a = \emptyset$. Then D_ϵ has empty interior. Otherwise, if U_0 is the interior of D_ϵ , we have

$$\begin{aligned}(U_0 \times B_\epsilon(0)) \cap X &= \emptyset \\ (U_0 \times B_\epsilon(0)) \cap \overline{X} &= \emptyset \\ (U_0 \times \{0\}) \cap \overline{X} &= \emptyset,\end{aligned}$$

contradicting the fact that $U_0 \times \{0\} \subseteq U \times \{0\} \subseteq \overline{X}$. Thus D_ϵ has empty interior as claimed. By Lemma 3.1, the union $\bigcup_\epsilon D_\epsilon$ has empty interior. The complement $U \setminus \bigcup_\epsilon D_\epsilon$ then has non-empty interior. Take U_0 to be the interior of $U \setminus \bigcup_\epsilon D_\epsilon$. If $a \in U_0$, then $a \notin D_\epsilon$ for any ϵ , which means $B_\epsilon(0) \cap X_a \neq \emptyset$, and $0 \in \overline{X}_a$. \square

Lemma 3.4. *Let U be a non-empty, open, definable subset of \mathbb{M}^k . Let $D \subseteq \mathbb{M}$ be a definable set with $0 \in \partial D$. Let $f : U \times D \rightarrow \mathbb{M}$ be a definable function. Suppose that*

$$\lim_{y \rightarrow 0} f(a, y) \text{ exists}$$

for every $a \in U$. Then there is a non-empty, open, definable subset $U_0 \subseteq U$ such that

$$\lim_{(x,y) \rightarrow (a,0)} f(x, y) \text{ exists and equals } \lim_{y \rightarrow 0} f(a, y)$$

for every $a \in U_0$.

Proof. Let $g(a) = \lim_{y \rightarrow 0} f(a, y)$ for $a \in U$.

Note that the projective line $\mathbb{P}^1(\mathbb{M})$ is definably compact. Regard f as a function to \mathbb{P}^1 . For each $a \in U$, let C_a be the set of cluster points of f around $(a, 0)$, i.e.,

$$C_a = \{b \in \mathbb{P}^1(\mathbb{M}) : (a, 0, b) \in \overline{G(f)}\} = \{b : (a, 0, b) \in \partial G(f)\},$$

where $G(f)$ is the graph of f , regarded as a subset of $\mathbb{M}^k \times \mathbb{P}^1(\mathbb{M})$. Then C_a is non-empty for each $a \in U$, by Lemma 2.8. In fact, clearly $g(a) \in C_a$ for each a .

The set C_a must be finite for generic $a \in U$. Otherwise, $\dim \partial G(f) \geq \dim(U) + 1 = k + 1$, contradicting the small boundaries property [5, Theorem 3.5]:

$$\dim \partial G(f) < \dim G(f) = \dim(U \times D) = \dim U + \dim D = k + 1.$$

So there is a non-empty, open definable subset $U_0 \subseteq U$ such that C_a is finite for $a \in U_0$. Shrinking U_0 further, we may assume that the cardinality of C_a is constant on U_0 .

If $|C_a| = 1$ for every $a \in U_0$, then $C_a = \{g(a)\}$ for each a , and $\lim_{(x,y) \rightarrow (a,0)} f(x, y) = g(a)$ by Lemma 2.8.

Suppose $|C_a| = N > 1$ for every $a \in U_0$. P -minimal theories have definable finite choice [5, Corollary 2.5], so there is a definable function h such that $h(a) \in C_a \setminus \{g(a)\}$ for every $a \in U_0$. P -minimal theories have generic continuity [5, Theorem 4.6], so by shrinking U_0 further, we may assume that g and h are continuous on U_0 . Fix some $a_0 \in U_0$ and let V, W

be clopen neighborhoods in $\mathbb{P}^1(\mathbb{M})$ separating $g(a_0)$ and $h(a_0)$. By continuity, we may shrink U_0 further and assume that

$$\begin{aligned} g(a) &\in V \\ h(a) &\in W \end{aligned}$$

for every $a \in U_0$. Let $X = \{(a, b) \in U \times D : f(a, b) \in W\}$. The fact that $h(a) \in C_a$ means that $(a, 0, h(a))$ is in the closure of the graph of f . Then for any ϵ , there are $(x, y) \in U \times D$ such that $x \in B_\epsilon(a)$, $y \in B_\epsilon(0)$, and $f(x, y) \in W$, i.e., $(x, y) \in X$. This shows that $(a, 0) \in \overline{X}$. By Lemma 3.3, we may shrink U_0 further and ensure that $0 \in \overline{X_a}$ for every $a \in U_0$. But the fact that $\lim_{y \rightarrow 0} f(a, y) = g(a) \in V$ implies that there is some ϵ such that if $y \in B_\epsilon(0)$, then $f(a, y) \in V$ and so $f(a, y) \notin W$, $(a, y) \notin X$, and $y \notin X_a$. Then $B_\epsilon(0)$ shows that $0 \notin \overline{X_a}$, a contradiction. \square

3.3 Rational powers

Recall that P_n denotes the set of non-zero n th powers in \mathbb{M} , and $\overline{P_n}$ denotes its closure $P_n \cup \{0\}$, the set of n th powers. The following fact is well-known, or an easy exercise:

Fact 3.5. *There is some ℓ such that P_ℓ is torsion-free.*

Definition 3.6. Let $q = a/b$ be a rational number in lowest terms. A q th power map on P_n is a multiplicative homomorphism

$$f : P_n \rightarrow \mathbb{M}^\times$$

such that f is 0-definable in the pure language of rings, and $f(x)^b = x^a$.

When such an f exists, we write it as x^q rather than $f(x)$.⁴

Example 3.7. If K contains $\sqrt{-1}$, then there is no square root map $x^{1/2}$ on P_2 . Otherwise,

$$1 = 1^{1/2} = (-1)^{1/2}(-1)^{1/2} = -1.$$

Note that if there is a q th power map on P_n , then there is a q th power map on P_m for any $n \mid m$, simply by restricting from P_n to P_m . The next lemma says that for fixed q , there is a q th power map on P_n for all sufficiently divisible n .

Lemma 3.8. *For any rational number $q = a/b$, there is some n such that there is a q th power map on n .*

Proof. Let P_ℓ be torsion-free, as in Fact 3.5. Then the b th power map $P_\ell \rightarrow P_{\ell b}$ is a 0-definable bijection. Let $g : P_{\ell b} \rightarrow P_\ell$ be its inverse. Let x^q be the map

$$\begin{aligned} P_{\ell b} &\rightarrow P_{\ell a} \\ x &\mapsto g(x)^a. \end{aligned}$$

\square

⁴Multiple q th power maps may exist on P_n , but the ambiguity won't cause us any problems. We will construct some canonical q th power maps in Lemma 3.8 below. If you like, you may assume that x^q only means the map constructed in Lemma 3.8.

4 Asymptotic behavior of functions in one variable

In this section, we look at the asymptotic behavior of definable functions

$$f : P_n \rightarrow \mathbb{M}$$

as $x \in P_n$ approaches 0. We show that after restricting to a smaller set $P_m \subseteq P_n$, the function f looks asymptotically like Cx^q for some constant $C \in \mathbb{M}$ and rational number q :

$$\lim_{x \in P_m, x \rightarrow 0} \frac{f(x)}{Cx^q} = 1.$$

The following fact is well-known, and can be deduced from [2, Lemma 4] or [8, Theorem 1.8].⁵

Fact 4.1 (1-dimensional cell decomposition). *If $D \subseteq \mathbb{M}$ is definable, then D can be partitioned into finitely many sets of the form*

$$\{x \in \mathbb{M} : \gamma_1 \square_1 v(x - c) \square_2 \gamma_2, x - c \in \lambda P_m\}$$

where $c \in \mathbb{M}$, $\gamma_1, \gamma_2 \in \Gamma$, $\lambda \in \mathbb{M}$, and each relation \square_1, \square_2 is either $<$ or \leq or no condition.

Fact 4.1 easily implies the following:

Corollary 4.2. *Let S be a finite set and $f : B_\epsilon(0) \cap P_n \rightarrow S$ be a definable function. Then there is $\delta \geq \epsilon$ and m a multiple of n such that the restriction*

$$f : B_\delta(0) \cap P_m \rightarrow S$$

is constant.

⁵I spent an hour unsuccessfully trying to find a place where Fact 4.1 is stated and proved directly, so I should give some explanation of where one can find the proof. By P -minimality, one may assume that D is definable in the pure field reduct, so the results of [2, 8] on semialgebraic sets apply. The results in [2, 8] are statements about multivariable definable sets in the standard model K rather than 1-variable sets in the monster model $\mathbb{M} \succ K$. They imply Fact 4.1 in the same way that the cell decomposition theorem in an o-minimal structure M implies the o-minimality of elementary extensions $N \succ M$. Of the two references just given, Cluckers' Lemma 4 in [2] more directly matches what we need for Fact 4.1, but unfortunately he omits the proof. With more work, Denef's Theorem 1.8 in [8] implies Fact 4.1. Specifically, Denef's Theorem 1.8 implies the statement that given any finite set of polynomials $f_1, \dots, f_n \in \mathbb{M}[t]$ and an integer n , we can partition \mathbb{M} into finitely many annuli A with centers c such that on A , each f_i has the form $u_i(t)^n \cdot h_i \cdot (t - c)^{\nu_i}$ with $u_i(t)$ a definable function taking values in \mathcal{O}^\times . For $k \mid n$, the truth value of $P_k(f_i(t))$ on A is determined by the coset $(t - c)P_n$ in \mathbb{M}^\times / P_n . So if we further split each annulus A into the cosets $A \cap \lambda P_n$, then the truth value of $P_k(f_i(t))$ is constant on each coset. Combined with Macintyre's quantifier elimination, this gives Fact 4.1. This proof is certainly well-known, implicit in many of the papers of Denef and Cluckers [8, 2, 4], such as Denef's re-proof of Macintyre's quantifier elimination in [8, Section 3].

Lemma 4.3. *Let $f : B_\epsilon(0) \cap P_n \rightarrow \mathcal{O}^\times$ be a definable function. Then there is $\delta \geq \epsilon$ and m a multiple of n such that the restriction*

$$f : B_\delta(0) \cap P_m \rightarrow \mathcal{O}^\times$$

extends to a continuous function

$$g : B_\delta(0) \cap \overline{P_m} \rightarrow \mathcal{O}^\times.$$

Equivalently, $\lim_{x \rightarrow 0} f(x)$ exists, if we restrict x to P_m .

Proof. Let G be the graph of f . Let C be the set of cluster points of f at $x = 0$:

$$\begin{aligned} C &= \{b : (0, b) \in \overline{G}\} \\ &= \{b : (0, b) \in \partial G\}. \end{aligned}$$

The set G has dimension 1, so its frontier has dimension 0 by the small boundaries property [5, Theorem 3.5]. Therefore C is finite. On the other hand, C is non-empty by Lemma 2.8, and definable compactness of \mathcal{O}^\times . Let $C = \{b_1, \dots, b_\ell\}$. Take pairwise disjoint clopen neighborhoods $V_i \ni b_i$, and let V_0 be the complement $\mathcal{O}^\times \setminus (V_1 \cup \dots \cup V_n)$. Let $h(x)$ be the unique $i \in \{0, \dots, n\}$ such that $f(x) \in V_i$. By Corollary 4.2, there are $\delta \geq \epsilon$ and $m \in n\mathbb{Z}$ such that $h(x)$ is a constant value i on $B_\delta(0) \cap P_m$. Then

$$f(x) \in V_i \text{ for } x \in B_\delta(0) \cap P_m.$$

Let f' be the restriction of f to $B_\delta(0) \cap P_m$ and let C' be the set of cluster points of f' at 0. Again, C' is non-empty by Lemma 2.8. Clearly $C' \subseteq C$. Because the graph of f' lies inside the closed set $\mathbb{M} \times V_i$, we must have $C' \subseteq V_i$. Then

$$C' \subseteq C \cap V_i = \begin{cases} \emptyset & \text{if } i = 0 \\ \{b_i\} & \text{if } 1 \leq i \leq n. \end{cases}$$

As C' is non-empty, we must have $i > 0$ and $C' = \{b_i\}$. Then Lemma 2.8 gives $\lim_{x \rightarrow 0} f'(x) = b_i$. \square

By a theorem of Raf Cluckers [3, Theorem 6], the value group Γ is a pure model of Presburger arithmetic. Therefore, Γ is stably embedded, and eliminates imaginaries.

Cluckers also proves a cell decomposition theorem for definable sets in Presburger arithmetic. For $i_1, \dots, i_n \in \{0, 1\}$, he defines a class of “ (i_1, \dots, i_n) -cells” in Γ^n , and shows that any definable set in Γ^n can be partitioned into finitely many cells. We only need the cases $n = 1, 2$. For subsets of Γ^1 , a (0)-cell is a singleton $\{a\}$, and a (1)-cell is an infinite set of the form $[a, b] \cap (k + n\Gamma)$, where $a, b \in (k + n\Gamma) \cup \{\pm\infty\}$. For subsets of Γ^2 ,

- A (0, 0)-cell is a singleton $\{(a, b)\}$.
- A (0, 1)-cell is a vertical segment $\{a\} \times C$, where $C \subseteq \Gamma$ is a (1)-cell.

- A $(1, 0)$ -cell is the graph of a linear function on a (1) -cell.
- There is also a notion of $(1, 1)$ -cell.

We don't need the precise definition of $(1, 1)$ -cell; the only important thing to know is that $(1, 1)$ -cells have acl-rank 2, or equivalently, dp-rank 2. (Algebraic closure satisfies exchange in Presburger arithmetic, so dp-rank agrees with acl-rank .) Therefore, if $D \subseteq \Gamma^2$ is a definable set with $\text{dp-rank} \leq 1$, then D is a finite union of $(0, 0)$ -cells, $(0, 1)$ -cells, and $(1, 0)$ -cells.

Lemma 4.4. *Let $f : B_\epsilon(0) \cap P_n \rightarrow \mathbb{M}^\times$ be a definable function. Then there is $\delta \geq \epsilon$, m a multiple of n , a rational number q , and an element $\gamma_0 \in \Gamma$ such that*

$$v(f(x)) = q \cdot v(x) + \gamma_0$$

for $x \in B_\delta(0) \cap P_m$.

Proof. The graph of f has dimension 1, or equivalently, dp-rank 1. Let Θ be its image under the valuation map:

$$\Theta = \{(v(x), v(f(x))) : x \in B_\epsilon(0) \cap P_n\} \subseteq \Gamma^2.$$

Then Θ has dp-rank at most 1, so it is a union

$$\Theta = C_1 \cup \dots \cup C_\ell,$$

where each C_i is a $(0, 0)$ -cell, a $(0, 1)$ -cell, or a $(1, 0)$ -cell. Let $h : B_\epsilon(0) \cap P_n \rightarrow \{1, \dots, \ell\}$ be the function such that $h(x) = i$ iff $(v(x), v(f(x))) \in C_i$. By Corollary 4.2, there are $\delta \geq \epsilon$ and $m \in n\mathbb{Z}$ such that h is constant on $B_\delta(0) \cap P_m$. Then there is a cell C such that

$$x \in B_\delta(0) \cap P_m \implies (v(x), v(f(x))) \in C.$$

The set

$$\{v(x) : x \in B_\delta(0) \cap P_m\}$$

has no upper bound, so there can be no upper bound on the first coordinate of points in C . This prevents C from being a $(0, 0)$ -cell or a $(0, 1)$ -cell. It must instead be a $(1, 0)$ -cell, i.e., the graph of a linear function $qx + \gamma_0$ on a (1) -cell $D \subseteq \Gamma$. Then

$$x \in B_\delta(0) \cap P_m \implies (v(x), v(f(x))) \in C \implies v(f(x)) = q \cdot v(x) + \gamma_0. \quad \square$$

Proposition 4.5. *Let $f : B_\epsilon(0) \cap P_n \rightarrow \mathbb{M}$ be a definable function. Then there is $\delta \geq \epsilon$ and m a multiple of n such that one of the following holds:*

1. f is identically zero on $B_\delta(0) \cap P_m$.
2. After restricting f to $B_\delta(0) \cap P_m$, there is a rational number q and constant $C \in \mathbb{M}^\times$ such that P_m has a q th power map and

$$\lim_{x \rightarrow 0} \frac{f(x)}{Cx^q} = 1.$$

Proof. Let

$$h(x) = \begin{cases} 1 & \text{if } f(x) = 0 \\ 0 & \text{if } f(x) \neq 0. \end{cases}$$

By increasing ϵ and m , we may assume that h is constant (Corollary 4.2). Then f is everywhere zero or everywhere non-zero. In the first case, we are done. Assume the second case. Lemma 4.4 gives $\gamma_0 \in \Gamma$ and $q \in \mathbb{Q}$ such that

$$v(f(x)) = q \cdot v(x) + \gamma_0$$

after possibly restricting the domain of f . Take C_0 with $v(C_0) = \gamma_0$. Increasing m , we may assume P_m has a q th power map (Lemma 3.8). Then

$$v\left(\frac{f(x)}{C_0 x^q}\right) = 0$$

for every x . By Lemma 4.3, we may shrink the domain further, and ensure that

$$\lim_{x \rightarrow 0} \frac{f(x)}{C_0 x^q} = C_1$$

for some constant $C_1 \in \mathcal{O}^\times$. Then

$$\lim_{x \rightarrow 0} \frac{f(x)}{C_0 C_1 x^q} = 1. \quad \square$$

5 Generic differentiability

5.1 Review of strict differentiability

Let K be a topological field (Hausdorff, non-discrete). The following definition should be standard, or equivalent to the standard definition.

Definition 5.1. Let $U \subseteq K^n$ be an open set and $f : U \rightarrow K^m$ be a function. Then f is *strictly differentiable* at $\bar{a} \in U$ if for every $i \leq n$, the limit

$$g_i(\bar{a}) = \lim_{\substack{\bar{x} \rightarrow \bar{a} \\ \epsilon \rightarrow 0}} \frac{f(\bar{x} + \epsilon \bar{e}_i) - f(\bar{x})}{\epsilon}$$

exists, where \bar{e}_i is the i th standard basis vector. The *strict derivative* $Df(\bar{a})$ is the $m \times n$ matrix whose i th column is $g_i(\bar{a})$.

Fact 5.2. 1. A matrix μ is the strict derivative of f at \bar{a} if and only if the following condition holds: for any neighborhood W of μ there is a neighborhood V of \bar{a} such that if $\bar{x}, \bar{y} \in V$, then

$$f(\bar{y}) - f(\bar{x}) = \mu' \cdot (\bar{y} - \bar{x})$$

for some matrix $\mu' \in W$.

2. If f is strictly differentiable on U , then the strict derivative Df is continuous on U .
3. A function $f : U \rightarrow K^m$ is strictly differentiable at $\bar{a} \in U$ if and only if each of the component functions $f_1, \dots, f_m : U \rightarrow K$ is strictly differentiable. In this case, the strict derivative of f is the matrix whose i th row is the strict derivative of f_i .
4. If f is strictly differentiable at \bar{a} , then f is continuous at \bar{a} .
5. If f is strictly differentiable at \bar{a} and $Df(\bar{a})$ has trivial kernel (i.e., $Df(\bar{a})$ has rank n), then f is injective on a neighborhood of \bar{a} .
6. If f is strictly differentiable at \bar{a} and g is strictly differentiable at $\bar{b} = f(\bar{a})$, then $g \circ f$ is strictly differentiable at \bar{a} , and $D(g \circ f)(\bar{a}) = Dg(\bar{b}) \cdot Df(\bar{a})$.
7. If $f : U \rightarrow V$ is a homeomorphism between two open sets $U, V \subseteq K^n$, and f is strictly differentiable at $\bar{a} \in U$, and $Df(\bar{a})$ is invertible, then f^{-1} is strictly differentiable at $\bar{b} = f(\bar{a})$, and $Df^{-1}(\bar{b}) = Df(\bar{a})^{-1}$.

Proof sketch. Part (3) is trivial. For part (2), given any $i \leq n$ and $\bar{a} \in U$ and closed neighborhood $N_0 \ni g_i(\bar{a})$, there is a neighborhood N of \bar{a} such that

$$\bar{x} \in N \text{ and } \epsilon \approx 0 \implies \frac{f(\bar{x} + \epsilon \bar{e}_i) - f(\bar{x})}{\epsilon} \in N_0.$$

Taking the limit $\epsilon \rightarrow 0$, it follows that

$$\bar{x} \in N \implies g_i(\bar{x}) \in N_0,$$

and we have shown that g_i is continuous at \bar{a} .

The other points can be seen most easily through non-standard analysis. In non-standard terms, a matrix μ is the strict derivative of f at \bar{a} if the following equivalent conditions hold:

- (a) If $i \leq n$ and $\bar{x} \approx \bar{a}$ and $\epsilon \approx 0$, then

$$\frac{f(\bar{x} + \epsilon \bar{e}_i) - f(\bar{x})}{\epsilon} \approx \mu \bar{e}_i$$

- (b) If $i \leq n$ and $\bar{x} \approx \bar{a}$ and $\epsilon \approx 0$, then

$$f(\bar{x} + \epsilon \bar{e}_i) = f(\bar{x}) + \mu \epsilon \bar{e}_i + o(\epsilon)$$

- (c) If $\bar{x} \approx \bar{a}$ and $\bar{\epsilon} \approx \bar{0}$, then

$$f(\bar{x} + \bar{\epsilon}) = f(\bar{x}) + \mu \bar{\epsilon} + o(\bar{\epsilon})$$

- (d) If $\bar{x} \approx \bar{a}$ and $\bar{\epsilon} \approx \bar{0}$, then there is a matrix of infinitesimals μ_ϵ such that

$$f(\bar{x} + \bar{\epsilon}) = f(\bar{x}) + \mu_\epsilon \bar{\epsilon}.$$

(e) If $\bar{x} \approx \bar{y} \approx \bar{a}$, then there is a matrix $\mu' \approx \mu$ such that

$$f(\bar{y}) = f(\bar{x}) + \mu'(\bar{y} - \bar{x}).$$

Condition (a) is a non-standard reformulation of Definition 5.1, and condition (e) is a non-standard reformulation of the condition in part (1). The implications

$$(a) \iff (b) \iff (c) \iff (d) \iff (e)$$

are straightforward, and (b) implies (c) by changing coordinates one-by-one. This verifies part (1).

For part (4), suppose μ is the strict derivative of f at \bar{a} . If $\bar{x} \approx \bar{a}$ then

$$f(\bar{x}) = f(\bar{a}) + \mu(\bar{x} - \bar{a}) + o(\bar{x} - \bar{a}) \approx f(\bar{a})$$

since $\bar{x} - \bar{a}$ is infinitesimal. Thus f is continuous at \bar{a} .

Part (5) holds because if a standard matrix μ has trivial kernel, then so does every non-standard $\mu' \approx \mu$. Then

$$\bar{y} - \bar{x} \neq \bar{0} \implies f(\bar{y}) - f(\bar{x}) = \mu'(\bar{y} - \bar{x}) \neq \bar{0}$$

where μ' is the matrix from (e).

For part (6), suppose $\mu_1 = Df(\bar{a})$ and $\mu_2 = Dg(\bar{b})$. If $\bar{x} \approx \bar{y} \approx \bar{a}$, then $f(\bar{x}) \approx f(\bar{y}) \approx f(\bar{a}) = \bar{b}$ by continuity, and so

$$g(f(\bar{y})) - g(f(\bar{x})) = \mu'_2(f(\bar{y}) - f(\bar{x})) = \mu'_2\mu'_1(\bar{y} - \bar{x})$$

for some matrices $\mu'_2 \approx \mu_2$ and $\mu'_1 \approx \mu_1$. Then $\mu'_2\mu'_1 \approx \mu_2\mu_1$, proving (6).

The proof of (7) is similar: if $\mu = Df(\bar{a})$ and $\bar{x} \approx \bar{y} \approx f(\bar{a})$, then $f^{-1}(\bar{y}) \approx f^{-1}(\bar{x}) \approx f^{-1}(\bar{a})$ because f is a homeomorphism, and so

$$\bar{y} - \bar{x} = \mu'(f^{-1}(\bar{y}) - f^{-1}(\bar{x})) \text{ for some } \mu' \approx \mu.$$

Then

$$f^{-1}(\bar{y}) - f^{-1}(\bar{x}) = (\mu')^{-1}(\bar{y} - \bar{x}) \text{ and } (\mu')^{-1} \approx \mu^{-1}. \quad \square$$

Fact 5.3. *If (K, v) is a valued field, then a matrix μ is the strict derivative $Df(\bar{a})$ if and only if the following holds: for any $\gamma \in \Gamma_K$, there is a neighborhood $N \ni \bar{a}$ such that*

$$\bar{x}, \bar{y} \in N \implies v(f(\bar{x}) - f(\bar{y}) - \mu(\bar{x} - \bar{y})) > \gamma + v(\bar{x} - \bar{y}).$$

Proof sketch. In non-standard terms, this condition says

$$\bar{x} \approx \bar{y} \approx \bar{a} \implies v(f(\bar{x}) - f(\bar{y}) - \mu(\bar{x} - \bar{y})) \gg v(\bar{x} - \bar{y}),$$

where $\gamma_1 \gg \gamma_2$ means that $\gamma_1 - \gamma_2$ is greater than every standard element of the value group. Equivalently,

$$(\bar{x} \approx \bar{a} \text{ and } \bar{\epsilon} \approx \bar{0}) \implies v(f(\bar{x} + \bar{\epsilon}) - f(\bar{x}) - \mu\bar{\epsilon}) \gg v(\bar{\epsilon}).$$

But $v(\bar{b}) \gg v(\bar{c})$ iff \bar{b} is $o(\bar{c})$, so we can rephrase this as

$$(\bar{x} \approx \bar{a} \text{ and } \bar{\epsilon} \approx \bar{0}) \implies (f(\bar{x} + \bar{\epsilon}) - f(\bar{x}) - \mu\bar{\epsilon} \text{ is } o(\bar{\epsilon})).$$

This is Condition (c) in the proof of Fact 5.2. \square

For reference, here is the definition of multi-variable differentiability:

Definition 5.4. A function $f : U \rightarrow \mathbb{M}^m$ is *differentiable* at $\bar{a} \in U$ if there is an $m \times n$ matrix μ such that for every neighborhood W of μ , there is a neighborhood V of \bar{a} such that if $\bar{x} \in V$, then

$$f(\bar{x}) - f(\bar{a}) = \mu' \cdot (\bar{x} - \bar{a})$$

for some matrix $\mu' \in W$. The matrix μ is called the *derivative* of f at \bar{a} .

In non-standard terms, this says that if $\bar{x} \approx \bar{a}$, then

$$f(\bar{x}) = f(\bar{a}) + \mu \cdot (\bar{x} - \bar{a}) + o(\bar{x} - \bar{a}).$$

The derivative is unique, when it exists. Strict differentiability implies differentiability, and the strict derivative equals the derivative when it exists.

5.2 Generic (strict) differentiability in one variable

Return to the P -minimal monster model \mathbb{M} . Let U be a definable non-empty open set in \mathbb{M}^1 and let $f : U \rightarrow \mathbb{M}$ be a definable function.

Definition 5.5. If $m \geq 1$ and $\lambda \in \mathbb{M}^\times$ and $q \in \mathbb{Q}$ and P_m has a q th power map, then f is (λ, m, q) -*differentiable* at a if

$$\lim_{\substack{x \rightarrow 0 \\ x \in \lambda P_m}} \frac{f(a+x) - f(a)}{x^q}$$

exists, in which case the limit is the (λ, m, q) -*derivative* at a . Similarly, f is *strictly* (λ, m, q) -*differentiable* at a if

$$\lim_{\substack{(w,x) \rightarrow (a,0) \\ x \in \lambda P_m}} \frac{f(w+x) - f(w)}{x^q}$$

exists, in which case the limit is the *strict* (λ, m, q) -*derivative*. In both definitions, we shorten $(\lambda, m, 1)$ to (λ, m) .

For example, the (λ, m) -derivative is like a directional derivative in the direction λP_m , and a $(1, 1)$ -derivative is an ordinary derivative. Note that (λ, m, q) -differentiability depends only on the multiplicative coset λP_m . For fixed m , there are only finitely many possibilities for λP_m because P_m has finite index.

Remark 5.6. If $f : U \rightarrow \mathbb{M}$ is (λ, m, q) -differentiable on U , then it is strictly (λ, m, q) -differentiable on a smaller open set $U_0 \subseteq U$, by Lemma 3.4 (with $D = \lambda P_m$).

Say that $f : U \rightarrow \mathbb{M}$ is *somewhere locally constant* if there is a ball $B \subseteq U$ such that $f \upharpoonright B$ is constant, and *nowhere locally constant* otherwise.

Lemma 5.7. Suppose $f : U \rightarrow \mathbb{M}$ is definable and nowhere locally constant, $\lambda \in \mathbb{M}^\times$, and $a \in U$. Then there is some m and q such that f is $(1, m, q)$ -differentiable at a and the $(1, m, q)$ -derivative is non-zero.

Proof. This comes directly from Proposition 4.5. In more detail, replacing $f(x)$ with $f(x + a) - f(a)$, we may assume that $a = f(a) = 0$. Applying Proposition 4.5, we get an $m \geq 1$ and a ball $B_\epsilon(0)$ such that one of two things happens:

- f is identically zero on $P_m \cap B_\epsilon(0)$, contradicting the assumption on f .
- There is a constant $C \in \mathbb{M}^\times$ and a rational number q such that

$$\lim_{\substack{x \rightarrow 0 \\ x \in B_\epsilon(0) \cap P_m}} \frac{f(x)}{Cx^q} = 1$$

Then C^{-1} is the $(1, m, q)$ -derivative of f at 0. \square

Lemma 5.8. *If $f : U \rightarrow \mathbb{M}$ is definable and nowhere locally constant, and $\lambda \in \mathbb{M}^\times$, then there is an integer m and a smaller non-empty open set $U_0 \subseteq U$ such that f is strictly (λ, m) -differentiable on U_0 .*

Proof. The case where f is somewhere locally constant is trivial, so assume f is nowhere locally constant. Changing coordinates ($x' = x/\lambda$) we reduce to the case where $\lambda = 1$. Let $D_{q,m}$ be the set of $a \in U$ such that f is $(1, m, q)$ -differentiable at a , and the $(1, m, q)$ -derivative is non-zero. By Lemma 5.7, every $a \in U$ belongs to some $D_{q,m}$. By saturation, U is covered by finitely many of the $D_{q,m}$. Then one of them must be infinite, and hence have non-empty interior. Shrinking U , we may assume that $U \subseteq D_{q,m}$. So, for every $a \in U$, the $(1, m, q)$ -derivative exists and is non-zero. By Remark 5.6, we can shrink U further and arrange for f to be strictly $(1, m, q)$ -differentiable on U . It remains to show that $q = 1$.

Let $g(a)$ be the strict $(1, m, q)$ -derivative of f at $a \in U$:

$$g(a) = \lim_{\substack{(w,x) \rightarrow (a,0) \\ x \in P_m}} \frac{f(w+x) - f(w)}{x^q}. \quad (*)$$

Remember that we arranged $g(a) \neq 0$ for all a . If $c \in P_m$, then

$$\lim_{\substack{(w,x) \rightarrow (a,0) \\ x \in P_m}} \frac{f(w+cx) - f(w)}{x^q} = \lim_{\substack{(w,y) \rightarrow (a,0) \\ y \in P_m}} \frac{f(w+y) - f(w)}{y^q/c^q} = c^q g(a) \quad (\dagger)$$

by the change of coordinates $y = cx$.

Take some positive integer $k > 1$ such that $k \in P_m$. For example, $k = p^n + 1$ works for $n \gg 0$. Fix some $a \in U$. By $(*)$,

$$\begin{aligned} \lim_{\substack{x \rightarrow 0 \\ x \in P_m}} \frac{f(a+kx) - f(a)}{x^q} &= \lim_{\substack{x \rightarrow 0 \\ x \in P_m}} \sum_{i=0}^{k-1} \frac{f(a+ix+x) - f(a+ix)}{x^q} \\ &= \sum_{i=0}^{k-1} \lim_{\substack{x \rightarrow 0 \\ x \in P_m}} \frac{f(a+ix+x) - f(a+ix)}{x^q} \\ &= \sum_{i=0}^{k-1} g(a) = k \cdot g(a), \end{aligned}$$

because $(a + ix, x) \rightarrow (a, 0)$ as $x \rightarrow 0$. On the other hand,

$$\lim_{\substack{x \rightarrow 0 \\ x \in P_m}} \frac{f(a + kx) - f(a)}{x^q} = k^q \cdot g(a)$$

by (\dagger) . Thus $k = k^q$. Regardless of how one chooses k^q , this is impossible unless $q = 1$.

Then $q = 1$ and f is strictly $(1, m)$ -differentiable on U . \square

Corollary 5.9. *Let $f : U \rightarrow \mathbb{M}$ be definable.*

1. *For every ball $B \subseteq U$ and $\lambda \in \mathbb{M}^\times$, there is a smaller ball $B' \subseteq B$ and an $m \geq 1$ such that f is strictly (λ, m) -differentiable on B' .*
2. *In the previous point, m can be chosen independently of λ and B . In other words, there is an $m \geq 1$ such that for every ball $B \subseteq U$ and $\lambda \in \mathbb{M}^\times$, there is a smaller ball $B' \subseteq B$ on which f is strictly (λ, m) -differentiable.*

Proof. The first point follows by applying Lemma 5.8 to $f \upharpoonright B$. The second point follows by compactness. \square

Lemma 5.10. *Let $f : U \rightarrow \mathbb{M}$ be definable. Then there is a non-empty open $U_0 \subseteq U$ such that f is differentiable on U_0 .*

Proof. Let m be as in Corollary 5.9(2). Let $\lambda_1 P_m, \dots, \lambda_k P_m$ enumerate the finitely many cosets of P_m . By Corollary 5.9(2) we can shrink U and assume that f is strictly (λ_1, m) -differentiable on U . Repeating this for $\lambda_2, \dots, \lambda_k$ we may assume that f is strictly (λ_i, m) -differentiable for every i . Then f is strictly (λ, m) -differentiable for every λ . Let $g_\lambda(a)$ denote the strict (λ, m) -derivative at $a \in U$:

$$g_\lambda(a) = \lim_{\substack{(w,x) \rightarrow (a,0) \\ x \in \lambda P_m}} \frac{f(w+x) - f(w)}{x}.$$

Note that if $a \in U$ and $c \in \lambda P_m$, then

$$\lim_{\substack{(w,x) \rightarrow (a,0) \\ x \in P_m}} \frac{f(w+cx) - f(w)}{x} = \lim_{\substack{(w,y) \rightarrow (a,0) \\ y \in cP_m}} \frac{f(w+y) - f(w)}{y/c} = cg_\lambda(a) \quad (\dagger)$$

via the change of coordinates $y = cx$.

Claim 5.11. *For any $a \in U$ and $\lambda \in \mathbb{M}^\times$, $g_\lambda(a) = g_1(a)$.*

Proof. Take $b \in \lambda P_m$ so small that $1+b \in P_m$. Then

$$\lim_{\substack{x \rightarrow 0 \\ x \in P_m}} \frac{f(a + (1+b)x) - f(a)}{x} = (1+b)g_1(a)$$

by (†). But the same value can alternatively be calculated as

$$\begin{aligned} \lim_{\substack{x \rightarrow 0 \\ x \in P_m}} \frac{f(a+x+bx) - f(a+x)}{x} + \lim_{\substack{x \rightarrow 0 \\ x \in P_m}} \frac{f(a+x) - f(a)}{x} \\ = bg_\lambda(a) + g_1(a) \end{aligned}$$

by (†). Thus

$$(1+b)g_1(a) = bg_\lambda(a) + g_1(a),$$

implying $g_\lambda(a) = g_1(a)$. □_{Claim}

It follows that

$$\lim_{\substack{x \rightarrow 0 \\ x \in \lambda P_m}} \frac{f(a+x) - f(a)}{x} = g_\lambda(a) = g_1(a)$$

for any coset λP_m . Since there are only finitely many cosets,

$$\lim_{x \rightarrow 0} \frac{f(a+x) - f(a)}{x} = g_1(a),$$

and we see that $g_1(x)$ is the derivative of f . We have shown that f is differentiable on U . □

Proposition 5.12. *If $f : U \rightarrow \mathbb{M}$ is definable, then f is strictly differentiable at all but finitely many points.*

Proof. Let X be the set of points on which f is *not* strictly differentiable. If X is infinite, it contains an open set by P -minimality. Restricting f to this open set, we may assume that f is nowhere strictly differentiable. Lemma 5.10 gives a smaller open set on which f is differentiable, and then Remark 5.6 gives a further smaller open set on which f is strictly differentiable, a contradiction. □

Remark 5.13. Proposition 5.12 shows that if $f : \mathbb{M} \rightarrow \mathbb{M}$ is definable and $a \in \mathbb{M}$ is generic, then f has the following asymptotic expansion around a :

$$f(a+\epsilon) = f(a) + f'(a)\epsilon + o(\epsilon).$$

Now consider the remainder $f(a+\epsilon) - (f(a) + f'(a)\epsilon)$. Using Proposition 4.5, the remainder must look like a power of ϵ :

$$f(a+\epsilon) - f(a) - f'(a)\epsilon \propto C_a \epsilon^q, \tag{*}$$

at least for ϵ in some P_m . Using the proof strategy from Lemmas 5.8 and 5.10, one can show that $q = 2$, that $(*)$ holds on *all* cosets of P_m , and that $C_a = f''(a)/2$. Thus

$$f(a+\epsilon) = f(a) + f'(a)\epsilon + \frac{f''(a)}{2}\epsilon^2 + o(\epsilon^2).$$

Continuing on in this way, one should get a Taylor series expansion

$$f(a+\epsilon) = f(a) + f'(a)\epsilon + \dots + \frac{f^{(n)}(a)}{n!}\epsilon^n + o(\epsilon^n)$$

for each n . There should also be a multivariable version of these statements. Unfortunately, the calculations are too complicated to carry out in this paper.

5.3 Multivariable generic differentiability

Theorem 5.14. *Let $U \subseteq \mathbb{M}^n$ be non-empty, open, and definable. Let $f : U \rightarrow \mathbb{M}^m$ be a definable function. Then f is strictly differentiable on a definable open set $U_0 \subseteq U$ with $\dim U \setminus U_0 < \dim U$.*

Proof. We will only consider the case $m = 1$, which is sufficient, by Fact 5.2(3). By the usual methods, it suffices to find a non-empty open subset $U_0 \subseteq U$ on which f is strictly differentiable.

Let U_i be the set of points in U such that the i th partial derivative exists:

$$\lim_{y \rightarrow 0} \frac{f(\bar{a} + y\bar{e}_i) - f(\bar{a})}{y} \text{ exists.}$$

The complement $U \setminus U_i$ cannot contain a ball, by Proposition 5.12. Therefore, U_i contains a ball. Shrinking U , we may assume that the i th partial derivative exists everywhere on U . By Lemma 3.4, we may further shrink U and assume that

$$\lim_{(\bar{x}, y) \rightarrow (\bar{a}, 0)} \frac{f(\bar{x} + y\bar{e}_i) - f(\bar{x})}{y} \text{ exists}$$

for any $\bar{a} \in U$. Repeating this for $i = 1, 2, \dots, n$, we can arrange this to hold for every i . Then f is strictly differentiable (Definition 5.1). \square

5.4 The inverse function theorem

We quickly check that one of the standard proofs of the inverse function theorem works in our context.

Lemma 5.15. *Let $B \subseteq \mathbb{M}^n$ be a ball around 0. Let $f : B \rightarrow B$ be a function which is contracting:*

$$v(f(x) - f(y)) > v(x - y) \text{ for distinct } x, y \in B.$$

1. *There is a unique $x \in B$ such that $f(x) = x$.*
2. *For any $c \in B$, there is a unique $x \in B$ such that $f(x) = x - c$.*

Proof. 1. Uniqueness is clear. Suppose existence fails (f has no fixed point). Note that f is continuous. Let $g : B \rightarrow \Gamma$ be the function $g(x) = v(f(x) - x)$. Then

$$g(f(x)) = v(f(f(x)) - f(x)) > v(f(x) - x) = g(x),$$

so the set $\{g(x) : x \in B\}$ has no maximum. Since Γ is definably well-ordered, $\{g(x) : x \in B\}$ has no upper bound. Then each of the closed sets

$$D_\gamma = \{x \in B : g(x) \geq \gamma\} = \{x \in B : v(f(x) - x) \geq \gamma\}$$

is non-empty. By definable compactness, there is $a \in \bigcap_{\gamma \in \Gamma} D_\gamma$. But then $v(f(a) - a) = +\infty$, so a is a fixed point.

2. Apply the previous point to the function $g : B \rightarrow B$ given by $g(x) = f(x) + c$. \square

Lemma 5.16. *Let U be a definable neighborhood of $\bar{a} \in \mathbb{M}^n$ and let $f : U \rightarrow \mathbb{M}^n$ be a definable function which is strictly differentiable on U , such that $Df(\bar{a})$ is invertible. Then the image of f contains a ball around $f(\bar{a})$.*

Proof. Changing coordinates, we may assume $\bar{a} = f(\bar{a}) = \bar{0}$, and $Df(\bar{a})$ is the $n \times n$ identity matrix. Let $h : U \rightarrow \mathbb{M}^n$ be the function $h(\bar{x}) = \bar{x} - f(\bar{x})$. Then h is strictly differentiable on U , and $Dh(\bar{0})$ vanishes. By Fact 5.3 (with $\gamma = 0$), there is a ball $B \ni \bar{0}$ with $B \subseteq U$ such that

$$\bar{x}, \bar{y} \in B \implies v(h(\bar{x}) - h(\bar{y})) > v(\bar{x} - \bar{y}). \quad (*)$$

In particular, h is contracting. Note that $h(\bar{0}) = \bar{0}$, so if $\bar{x} \in B$, then

$$v(h(\bar{x})) = v(h(\bar{x}) - h(\bar{0})) > v(\bar{x} - \bar{0}) = v(\bar{x}).$$

Thus h maps B into B . By Lemma 5.15(2), for any $\bar{c} \in B$, there is $\bar{x} \in B$ such that

$$\bar{x} - f(\bar{x}) = h(\bar{x}) = \bar{x} - \bar{c},$$

so $f(\bar{x}) = \bar{c}$. We have shown that $\text{im}(f)$ contains B . \square

Corollary 5.17. *Let $U \subseteq \mathbb{M}^n$ be a definable open set and $f : U \rightarrow \mathbb{M}^n$ be definable and strictly differentiable. If $Df(\bar{a})$ is invertible for every $\bar{a} \in U$, then f is an open map.*

Theorem 5.18 (Inverse function theorem). *Let $U \subseteq \mathbb{M}^n$ be a definable open set and $f : U \rightarrow \mathbb{M}^n$ be a definable, strictly differentiable function. Let $\bar{a} \in U$ be a point such that the strict derivative $Df(\bar{a})$ is invertible. Then there is a smaller open set $\bar{a} \in U_0 \subseteq U$ such that f restricts to a homeomorphism $U_0 \rightarrow V_0$ for some neighborhood $V_0 \ni f(\bar{a})$, and both $f : U_0 \rightarrow V_0$ and $f^{-1} : V_0 \rightarrow U_0$ are strictly differentiable.*

Proof. The derivative $Df(\bar{x})$ is continuous by Fact 5.2(2). So we can find a neighborhood $U_0 \ni \bar{a}$ such that $Df(\bar{x})$ is invertible for every $\bar{x} \in U_0$. The fact that $Df(\bar{a})$ is invertible implies that f is injective on a neighborhood of \bar{a} , by Fact 5.2(5). Shrinking U_0 further, we may assume that $f \upharpoonright U_0$ is injective. Then $f \upharpoonright U_0$ is a continuous, injective, open map, so it is a homeomorphism onto its image V_0 . The inverse function $f^{-1} : V_0 \rightarrow U_0$ is strictly differentiable by Fact 5.2(7). \square

6 Recognizing p -adic Lie groups

Recall that a profinite group G is a *pro- p group* if it is an inverse limit of finite p -groups [9, Proposition 1.12]. Using work of Lazard [19], as reported in [9], one can prove the following:

Fact 6.1. *Let G be a pro- p group. Let S_i be the image of the p^i th power map $G \rightarrow G$. Let n be an integer. Suppose the following conditions hold:*

1. G has no p -torsion.
2. S_1 and S_2 are normal open subgroups of G , and the quotients G/S_1 and G/S_2 are abelian.
3. G/S_1 has size p^n .

Then G is isomorphic, as a topological group, to an n -dimensional Lie group over \mathbb{Q}_p .

Fact 6.1 is probably obvious or well-known to experts on p -adic Lie groups, but for the rest of us, we need to chain together some references from [9]. First we recall some definitions and facts:

1. A pro- p group G is *powerful* [9, Definition 3.1] if p is odd and $G/\overline{G^p}$ is abelian, or $p = 2$ and $G/\overline{G^4}$ is abelian, where G^n denotes the subgroup generated by n th powers.
2. If G is a profinite group, the *Frattini subgroup* $\Phi(G)$ is the intersection of all maximal open proper subgroups of G [9, Definition 1.8].
3. If G is a pro- p group, then the Frattini subgroup $\Phi(G)$ is the closure of the subgroup generated by p th powers and commutators [9, Proposition 1.13]:

$$\Phi(G) = \overline{G^p[G, G]}.$$

4. A *topological generating set* of G is a subset X generating a dense subgroup of G , and G is *finitely generated* if there is a finite topological generating set [9, p. 20].
5. A pro- p group is finitely generated iff $\Phi(G)$ is open [9, Proposition 1.14].
6. If G is a finitely generated powerful pro- p group, then $\Phi(G)$ is exactly the set of p th powers.
7. If G is a topological group, then $d(G)$ is the minimum cardinality of a topological generating set for G . When G is a pro- p group, $d(G)$ equals the dimension of $G/\Phi(G)$ as a vector space over \mathbb{F}_p . (See the paragraph above [9, Theorem 3.8].)
8. A pro- p group is *uniformly powerful* if it is powerful, finitely generated, and has no p -torsion [9, Theorem 4.5].⁶
9. If G is a uniformly powerful pro- p group, then G is a Lie group over \mathbb{Q}_p [9, Theorem 8.18]. Moreover, the dimension of G as a Lie group over \mathbb{Q}_p equals $d(G)$ by Theorem 8.36 and Definition 4.7 of [9].

Combining these ingredients, we prove Fact 6.1:

⁶Theorem 4.5 in [9] says “torsion-free” rather than “ p -torsion free”, but as noted in the third sentence of the proof, these conditions are equivalent for pro- p groups.

Proof. By our assumptions on G ,

$$\begin{aligned}\overline{G^p} &= \overline{\langle S_1 \rangle} = \overline{S_1} = S_1 \text{ for any } p \\ \overline{G^4} &= \overline{\langle S_2 \rangle} = \overline{S_2} = S_2 \text{ when } p = 2,\end{aligned}$$

and both the quotients G/S_1 and G/S_2 are abelian. Thus G is a powerful pro- p group. The Frattini subgroup $\Phi(G)$ is clopen because it contains the clopen subgroup S_1 :

$$\Phi(G) = \overline{G^p[G, G]} \supseteq \overline{G^p} = S_1.$$

Therefore G is finitely generated, which implies that $\Phi(G) = \{x^p : x \in G\} = S_1$. Moreover, G is uniformly powerful because G has no p -torsion. Then G is a p -adic Lie group of dimension

$$\dim(G) = d(G) = \dim_{\mathbb{F}_p} G/\Phi(G) = \dim_{\mathbb{F}_p} G/S_1 = n,$$

since G/S_1 has cardinality p^n . □

We will apply this to the following setting:

Lemma 6.2. *Let K be a degree e extension of \mathbb{Q}_p , and let \mathcal{O}_K be the ring of integers in K . Let d be an integer. Let \star be a group operation on \mathcal{O}_K^d . Let $f : \mathcal{O}_K^d \rightarrow \mathcal{O}_K^d$ be the p th power map (with respect to \star), i.e.,*

$$f(x) = \underbrace{x \star x \star \cdots \star x}_{p \text{ times}}.$$

Suppose the following conditions hold:

1. (\mathcal{O}_K^d, \star) is a topological group with respect to the usual topology on \mathcal{O}_K .
2. The identity element is $\bar{0}$.
3. For every $n < \omega$, the set $p^n \mathcal{O}_K^d$ is a normal subgroup of (\mathcal{O}_K^d, \star) whose cosets are the additive cosets $\bar{a} + p^n \mathcal{O}_K^d$. Equivalently, the equivalence relation

$$\bar{x} \equiv_n \bar{y} \iff \bar{x} - \bar{y} \in p^n \mathcal{O}_K^d$$

is a congruence on the group (\mathcal{O}_K^d, \star) .

4. For $n \leq 2$, the group structure on $(\mathcal{O}_K^d, \star)/p^n \mathcal{O}_K^d$ agrees with the additive structure $(\mathcal{O}_K^d, +)/p^n \mathcal{O}_K^d$. In other words,

$$\bar{x} + \bar{y} \equiv \bar{x} \star \bar{y} \pmod{p^n \mathcal{O}_K^d}.$$

5. The map f scales distances by a factor of p :

$$v(f(\bar{x}) - f(\bar{y})) = v(p) + v(\bar{x} - \bar{y}).$$

6. The image of f contains $p \mathcal{O}_K^d$.

Then (\mathcal{O}_K, \star) is an n -dimensional p -adic Lie group for $n = de$.

Proof. Let $G = (\mathcal{O}_K^d, \star)$ and let G_n be the normal subgroup $(p^n \mathcal{O}_K^d, \star)$ from Assumption 3. Since we are taking the standard topology on \mathcal{O}_K^d (Assumption 1), the descending chain $G \supseteq G_1 \supseteq G_2 \supseteq \cdots$ is a neighborhood basis of the identity element $\bar{0}$ (Assumption 2). By Assumption 3, the index $|G : G_i|$ equals the additive index $|(G, +) : (G_i, +)|$. Since $(\mathcal{O}_K^d, +)$ is isomorphic to $(\mathbb{Z}_p^{de}, +)$, the index $|G : G_i|$ equals p^{dei} . In particular,

$$G \cong \varprojlim_{i \rightarrow \infty} G/G_i \text{ is a pro-} p \text{ group.}$$

Taking $\bar{y} = \bar{0}$ in Assumption 5, we see that

$$v(f(\bar{x})) = v(\bar{x}) + v(p). \quad (*)$$

In particular, if $\bar{x} \neq \bar{0}$, then $v(f(\bar{x})) = v(\bar{x}) + v(p) < \infty$, so $f(\bar{x}) \neq \bar{0}$. This shows that the group (\mathcal{O}_K^d, \star) has no p -torsion.

Equation $(*)$ also shows that f maps $p^i \mathcal{O}_K^d$ into $p^{i+1} \mathcal{O}_K^d$. In fact,

$$f(p^i \mathcal{O}_K^d) = p^{i+1} \mathcal{O}_K^d$$

because if \bar{y} is on the right hand side, then $\bar{y} \in p^{i+1} \mathcal{O}_K^d \subseteq p \mathcal{O}_K^d \subseteq \text{im}(f)$ by Assumption 6, so $\bar{y} = f(\bar{x})$ for some $\bar{x} \in \mathcal{O}_K^d$. But then Equation $(*)$ shows that $v(\bar{x}) = v(\bar{y}) - v(p) \geq i \cdot v(p)$, so $\bar{x} \in p^i \mathcal{O}_K^d$.

Consequently, the image of the p^i th power map is exactly $p^i \mathcal{O}_K^d = G_i$. In the notation of Fact 6.1, we have $S_i = G_i$. In particular, S_i is a normal open subgroup of G , and G/S_i has size p^{dei} . Then G/S_1 has size p^{de} . For $i \leq 2$, the quotient G/S_i is abelian by Assumption 4.

Now all the conditions of Fact 6.1 are satisfied, and so G is a p -adic Lie group of dimension de . \square

7 Conditions \mathfrak{A} , \mathfrak{B} , \mathfrak{C} , \mathfrak{D} , \mathfrak{E}

Work in a model M of the P -minimal theory T . Let \mathcal{O} denote the valuation ring of M .

Definition 7.1. Let $G = (\mathcal{O}^d, \star)$ be a definable group with underlying set \mathcal{O}^d , such that $\bar{0} \in \mathcal{O}^d$ is the identity element. We define the following conditions:

(\mathfrak{A}_n) For $n < \omega$, the group G satisfies condition \mathfrak{A}_n if the ball $p^n \mathcal{O}^d$ is a normal subgroup of G , and the \star -cosets of $p^n \mathcal{O}^d$ agree with the additive cosets:

$$\{a \star p^n \mathcal{O}^d : a \in \mathcal{O}^d\} = \{a + p^n \mathcal{O}^d : a \in \mathcal{O}^d\}.$$

Equivalently, the relation $v(\bar{x} - \bar{y}) \geq n \cdot v(p)$ is a congruence on G .

(\mathfrak{A}_ω) The group G satisfies \mathfrak{A}_ω if it satisfies \mathfrak{A}_n for all $n < \omega$.

(\mathfrak{A}_∞) The group G satisfies \mathfrak{A}_∞ if for any $\bar{x}, \bar{y}, \bar{z}$ in \mathcal{O}^n and $\gamma \in \Gamma$, the relation $v(\bar{x} - \bar{y}) \geq \gamma$ if a congruence on G .

(\mathfrak{B}_n) For $n < \omega$, the group G satisfies condition \mathfrak{B}_n if it satisfies \mathfrak{A}_n , and moreover the group operation on the \star -cosets of $p^n \mathcal{O}^d$ is the usual addition.

Equivalently, $G = (\mathcal{O}^d, \star)$ satisfies \mathfrak{B}_n if

$$\bar{x} \star \bar{y} \equiv \bar{x} + \bar{y} \pmod{p^n \mathcal{O}^d}$$

for every $\bar{x}, \bar{y} \in \mathcal{O}^d$.

(\mathfrak{B}_ω) The group G satisfies \mathfrak{B}_ω if it satisfies \mathfrak{B}_n for all $n < \omega$.

(\mathfrak{C}_n) For $n < \omega$, the group G satisfies condition \mathfrak{C}_n if

- It satisfies \mathfrak{A}_n
- Let $f(\bar{x}, \bar{y}) = \bar{x} \star \bar{y}$. For any multi-indices I, J with $|I| + |J| \leq n$, the Taylor series coefficient

$$\bar{c}_{I,J} = \frac{1}{I!J!} \frac{\partial f}{\partial \bar{x}^I \partial \bar{y}^J}(\bar{0}, \bar{0})$$

exists and is in $\mathcal{O}^d \cap p^{|I|+|J|-1} \mathcal{O}^d$.

- The quotient group $G/p^n \mathcal{O}^d$ has the following group structure:

$$\bar{x} \star \bar{y} \equiv \sum_{\substack{I,J \\ |I|+|J| \leq n}} \bar{c}_{I,J} \bar{x}^I \bar{y}^J \pmod{p^n \mathcal{O}^d}.$$

(\mathfrak{C}_ω) G satisfies \mathfrak{C}_ω if it satisfies \mathfrak{C}_n for all $n < \omega$.

(\mathfrak{D}) G satisfies \mathfrak{D} if the p th power map $f : G \rightarrow G$ satisfies the condition

$$v(f(\bar{x}) - f(\bar{y})) = v(\bar{x} - \bar{y}) + v(p)$$

for any distinct $\bar{x}, \bar{y} \in G$.

(\mathfrak{E}) G satisfies \mathfrak{E} if the image of the p th power map $f : G \rightarrow G$ contains the ball $p \mathcal{O}^d$.

Remark 7.2. Condition \mathfrak{B}_ω says something like $x \star y \approx x + y$, i.e., \star is close to addition. We could define \mathfrak{B}_∞ by analogy to \mathfrak{A}_∞ , but it would imply $x \star y = x + y$ for all x, y .

Remark 7.3. Conditions $\mathfrak{A}_n, \mathfrak{A}_\infty, \mathfrak{B}_n, \mathfrak{C}_n, \mathfrak{D}$, and \mathfrak{E} are definable in families, and preserved in elementary extensions. Conditions $\mathfrak{A}_\omega, \mathfrak{B}_\omega$, and \mathfrak{C}_ω are type-definable in families, and preserved in elementary extensions.

Example 7.4. In the standard model K , suppose that (\mathcal{O}^n, \star) is a definable group structure with $\bar{0}$ as the identity element, such that \star is given by a formal power series

$$\bar{x} \star \bar{y} = \sum_{I,J} \bar{c}_{I,J} \bar{x}^I \bar{y}^J.$$

Since $\bar{x} \star \bar{0} = \bar{0} \star \bar{x} = \bar{x}$, we in fact have

$$\bar{x} \star \bar{y} = \bar{x} + \bar{y} + \sum_{\substack{I,J \\ |I| \geq 1, |J| \geq 1}} \bar{c}_{I,J} \bar{x}^I \bar{y}^J.$$

Suppose $\bar{c}_{I,J} \in p^{|I|+|J|-1}\mathcal{O}^n$ for each I, J with $|I|, |J| \geq 1$. Then $\bar{c}_{I,J} \in \mathcal{O}^n \cap p^{|I|+|J|-1}\mathcal{O}^n$ for each I, J , and condition \mathfrak{C}_ω is easy to verify. The fact that \mathfrak{A}_n holds for all n also implies that \mathfrak{A}_∞ holds, since we are in the standard model.

7.1 From \mathfrak{A}_ω to compact domination

First, we recall a few well-known facts.

Remark 7.5. P -minimal theories are dp-minimal, because p -adically closed fields are dp-minimal [10, §6], and dp-minimality depends only on the collection of definable sets in one variable.

Remark 7.6. P -minimal theories are distal. This is well-known, but I don't have a reference on hand, so here is a proof:

Proof. p -adically closed fields have definable Skolem functions, so there is a definable function $f_0(x, y)$ such that

$$f_0(x + y, xy) \in \{x, y\}.$$

Letting $f(x, y) = f_0(x + y, xy)$, we get a definable function f such that

$$\begin{aligned} f(x, y) &= f(y, x) \\ f(x, y) &\in \{x, y\}. \end{aligned}$$

Using f , we see that $\text{tp}(a, b) \neq \text{tp}(b, a)$ for any $a \neq b$. Consequently, the only totally indiscernible sequences are the constant sequences. The function f also exists in P -minimal expansions, of course. Therefore, P -minimal theories also have the property that totally indiscernible sequences are constant. If p is a global type which is generically stable, then the Morley sequence of p is totally indiscernible [14, Proposition 3.2(ii)], hence constant, which makes p be a constant/realized type. Thus P -minimal theories have no non-constant generically stable types. By [24, Corollary 9.19], P -minimal theories are distal. \square

Recall that a definable group G has *finitely satisfiable generics* (fsg) if there is a global type p on G and a small set A_0 such that every translate of p is finitely satisfiable in A_0 [13, Definition 4.1]. Equivalently, G has fsg if there is a global type p , finitely satisfiable in a small set A_0 , such that p is “almost translation-invariant” in the sense that $\{g \cdot p : g \in G\}$ is small.

Remark 7.7. If G is a definable group in a distal theory, then the following are equivalent:

1. G has finitely satisfiable generics (fsg).
2. G is compactly dominated.
3. G is compactly dominated by (normalized) Haar measure on G/G^{00} .

The equivalence of (2) and (3) is [24, Lemma 8.36]. The equivalence of (1) and (2) is stated in [24, Example 8.42], but the proof is a bit hidden. For reference:

- G is compactly dominated iff G has a smooth left-invariant measure [24, Theorem 8.37].
- G is fsg iff G is definably amenable and G has a generically stable left-invariant measure [24, Proposition 8.33].
- G is definably amenable iff G has a left-invariant measure [24, Definition 8.12]. Thus G is fsg iff G has a generically stable left-invariant measure.
- Smooth measures are generically stable [24, Lemma 7.17 and §7.5]
- In distal theories, generically stable measures are smooth [24, Proposition 9.26].
- Therefore, in distal theories, generically stable measures are the same thing as smooth measures, and fsg groups are the same thing as compactly dominated groups.

Now work in a monster model \mathbb{M} of the P -minimal theory T .

Proposition 7.8. *Suppose $G = (\mathcal{O}^d, \star)$ satisfies condition \mathfrak{A}_ω .*

1. G is fsg and compactly dominated.
2. G^{00} is the subgroup $\bigcap_{n < \omega} p^n \mathcal{O}^d$.

Proof. The proof requires several steps. We first need to analyze the groups $(\mathcal{O}, +)$ and $(\mathcal{O}^d, +)$.

In the base theory $\text{Th}(K)$ of p -adically closed fields (i.e., the reduct to $\mathcal{L}_{\text{Rings}}$) the additive group $(\mathcal{O}, +)$ is fsg and the connected component $(\mathcal{O}, +)^{00}$ is $\bigcap_{n < \omega} p^n \mathcal{O}$, by [20, Corollaries 2.3–2.4].

The group $(\mathcal{O}, +)$ is also fsg in the P -minimal expansion T . To see this, take an almost translation-invariant global type q in the $\mathcal{L}_{\text{Rings}}$ -theory, finitely satisfiable in a small set A . Then q extends uniquely to a global type \hat{q} in the expansion, because the boolean algebra of definable sets is the same in the two languages (by P -minimality), so the spaces of global 1-types are the same. It is clear that \hat{q} is almost translation-invariant and finitely satisfiable in A , and so $(\mathcal{O}, +)$ is fsg in the expansion.

Similarly, $(\mathcal{O}, +)^{00}$ in the expansion is still $\bigcap_{n < \omega} p^n \mathcal{O}$. This holds because the collection of type-definable subsets of $(\mathcal{O}, +)$ is the same in both languages, by P -minimality again. So the two collections

$$\begin{aligned} & \{H : H \text{ is a type-definable, } H \triangleleft (\mathcal{O}, +), \text{ and } (\mathcal{O}, +)/H \text{ is small}\} \text{ in } \text{Th}(K) \\ & \{H : H \text{ is a type-definable, } H \triangleleft (\mathcal{O}, +), \text{ and } (\mathcal{O}, +)/H \text{ is small}\} \text{ in } T \end{aligned}$$

are identical, and have the same minimum element.

For the rest of the proof, we remain in the P -minimal expansion T , rather than the base theory $\text{Th}(K)$.

An extension of an fsg group by an fsg group is fsg [13, Proposition 4.5], so in particular a product of two fsg groups is fsg. Therefore $(\mathcal{O}^d, +)$ is fsg (in the P -minimal expansion T). Moreover, $(G \times H)^{00} = G^{00} \times H^{00}$. Therefore

$$(\mathcal{O}^d, +)^{00} = \bigcap_{n < \omega} p^n \mathcal{O}^d$$

holds (in the P -minimal expansion T). Note that $(\mathcal{O}^d, +)/(\mathcal{O}^d, +)^{00} \cong \mathcal{O}_K^d$. (For example, it's \mathbb{Z}_p^d when \mathbb{M} is elementarily equivalent to \mathbb{Q}_p .) By Remark 7.7, the group $(\mathcal{O}^d, +)^{00}$ is compactly dominated by Haar measure on \mathcal{O}_K^d .

Let μ be the unique smooth, translation invariant measure on the group $(\mathcal{O}^d, +)$. By the proof of Proposition 8.38 in [24], we know that μ is related to Haar measure on \mathcal{O}_K^d as follows. Let $D \subseteq \mathcal{O}^d$ be definable. Let $X_0, X_{1/2}, X_1 \subseteq (\mathcal{O}^d, +)/(\mathcal{O}^d, +)^{00} \cong \mathcal{O}_K^d$ be the following sets:

- X_0 is the set of cosets $a + (\mathcal{O}^d, +)^{00}$ disjoint from D .
- X_1 is the set of cosets $a + (\mathcal{O}^d, +)^{00}$ contained in D .
- $X_{1/2}$ is the remaining cosets.

Compact domination says that the Haar measure of $X_{1/2}$ is zero. By the proof of [24, Proposition 8.38], $\mu(D)$ is the Haar measure of X_1 .

Let $f : \mathcal{O}^d \rightarrow \mathcal{O}^d$ be a definable bijection such that

$$f(\bar{x}) \equiv f(\bar{y}) \pmod{p^n \mathcal{O}^d} \iff \bar{x} \equiv \bar{y} \pmod{p^n \mathcal{O}^d} \quad (*)$$

for each n . Then f preserves the relation $\bar{x} \equiv \bar{y} \pmod{(\mathcal{O}^d, +)^{00}}$, so f induces a map \tilde{f} on the quotient $(\mathcal{O}^d, +)/(\mathcal{O}^d, +)^{00} \cong \mathcal{O}_K^d$. It's easy to see that \tilde{f} is a measure-preserving homeomorphism. By compact domination, f preserves μ .

For any $a \in \mathcal{O}^d$, the left translation $f(x) = a \star x$ satisfies $(*)$, because the group (\mathcal{O}^d, \star) satisfies \mathfrak{A}_n . Therefore μ is invariant under left translations in (\mathcal{O}^d, \star) . The measure μ is also smooth. (Smoothness is unrelated to the group structure, and we chose μ to be the smooth translation invariant measure on $(\mathcal{O}^d, +)$.)

Therefore $G = (\mathcal{O}^d, \star)$ has a smooth measure invariant under left translations. By [24, Theorem 8.37], G is compactly dominated.

We know that $(\mathcal{O}^d, +)^{00} = \bigcap_{n < \omega} p^n \mathcal{O}^d$. It remains to show that $(\mathcal{O}^d, \star)^{00} \stackrel{?}{=} (\mathcal{O}^d, +)^{00}$. For each n , the set $p^n \mathcal{O}^d$ is a normal subgroup of both groups, and the cosets are the same in both groups (Condition \mathfrak{A}_n). Therefore $p^n \mathcal{O}^d$ has finite index in both groups, and so

$$(\mathcal{O}^d, \star)^{00} \subseteq (\mathcal{O}^d, \star)^0 \subseteq \bigcap_{n < \omega} p^n \mathcal{O}^d = (\mathcal{O}^d, +)^{00}.$$

Claim 7.9. *If D is a definable set containing $(\mathcal{O}^d, \star)^{00}$, then $D \div D := \{\bar{x} \star \bar{y}^{-1} : \bar{x}, \bar{y} \in D\}$ contains $(\mathcal{O}^d, +)^{00}$.*

Proof. As D contains $(\mathcal{O}^d, \star)^{00}$, boundedly many left \star -translates of D cover \mathcal{O}^d . By compactness, finitely many left \star -translates cover \mathcal{O}^d . These translates have the same measure with respect to μ , because μ is left-invariant for \star . Then D must have positive μ -measure. Because of the connection between μ and Haar measure, some additive coset $\bar{a} + (\mathcal{O}^d, +)^{00}$ must lie in D . By compactness, there is some n such that $\bar{a} + p^n \mathcal{O}^d \subseteq D$. By Condition \mathfrak{A}_n , the set $\bar{a} + p^n \mathcal{O}^d$ is a \star -coset of $p^n \mathcal{O}^d \triangleleft (\mathcal{O}^d, \star)$. Then $D \div D$ contains $p^n \mathcal{O}^d \supseteq (\mathcal{O}^d, +)^{00}$. □_{Claim}

Because $(\mathcal{O}^d, \star)^{00}$ is a type-definable subgroup of (\mathcal{O}^d, \star) , we have

$$\begin{aligned} (\mathcal{O}^d, \star)^{00} &= (\mathcal{O}^d, \star)^{00} \div (\mathcal{O}^d, \star)^{00} \\ &\stackrel{*}{=} \bigcap \{D \div D : D \text{ is definable and } D \supseteq (\mathcal{O}^d, \star)^{00}\} \stackrel{(\text{Claim})}{\supseteq} (\mathcal{O}^d, +)^{00}. \end{aligned}$$

The starred equality holds by compactness: if $\Sigma(\bar{x})$ is the partial type defining $(\mathcal{O}^d, \star)^{00}$ and \bar{c} is in \mathcal{O}^d , then the following are equivalent:

- $\bar{c} \in (\mathcal{O}^d, \star)^{00}$.
- There are $\bar{a}, \bar{b} \in (\mathcal{O}^d, \star)^{00}$ such that $\bar{c} = \bar{a} \star \bar{b}^{-1}$.
- The partial type in the variables \bar{x}, \bar{y} saying

$$\Sigma(\bar{x}) \text{ and } \Sigma(\bar{y}) \text{ and } \bar{c} = \bar{x} \star \bar{y}^{-1}$$

is finitely satisfiable.

- $\bar{c} \in \bigcap \{\phi(\mathbb{M}) \div \phi(\mathbb{M}) : \phi \in \Sigma\}$.
- $\bar{c} \in \bigcap \{D \div D : D \text{ is definable and } D \supseteq (\mathcal{O}^d, \star)^{00}\}$.

So we conclude that

$$(\mathcal{O}^d, \star)^{00} = (\mathcal{O}^d, +)^{00} = \bigcap_{n < \omega} p^n \mathcal{O}^d. \quad \square$$

7.2 Getting a p -adic Lie group

Continue to work in a monster model \mathbb{M} of a P -minimal expansion T of $\text{Th}(K)$ for some finite extension K/\mathbb{Q}_p .

Proposition 7.10. *Suppose $G = (\mathcal{O}^d, \star)$ satisfies condition \mathfrak{B}_ω . Then G is fsg and compactly dominated, and the compact topological group G/G^{00} is isomorphic to \mathcal{O}_K^d .*

Proof. By Proposition 7.8, G^{00} is the subgroup $\bigcap_{n < \omega} p^n \mathcal{O}^d$, and so G/G^{00} is the inverse limit

$$\varprojlim_{n \rightarrow \infty} (\mathcal{O}^d, \star) / p^n \mathcal{O}^d.$$

By condition \mathfrak{B}_ω , this is the same as

$$\varprojlim_{n \rightarrow \infty} (\mathcal{O}^d, +) / p^n \mathcal{O}^d,$$

which is just \mathcal{O}_K^d . □

Proposition 7.11. *Suppose $G = (\mathcal{O}^d, \star)$ satisfies condition \mathfrak{C}_ω . Then G is fsg and compactly dominated, and G/G^{00} is isomorphic to a d -dimensional Lie group over K .*

Proof. Proposition 7.8 gives the first part, and shows that $G^{00} = \bigcap_{k < \omega} p^k \mathcal{O}^n$. Then G/G^{00} is homeomorphic to \mathcal{O}_K^n via the standard part map $\text{st} : \mathcal{O} \rightarrow \mathcal{O}_K$. That is, the following two maps are equivalent:

$$\begin{aligned} G &\rightarrow G/G^{00} \\ \mathcal{O}^n &\xrightarrow{\text{st}} \mathcal{O}_K^n. \end{aligned}$$

Condition \mathfrak{C}_ω shows that the induced group structure on \mathcal{O}_K^n is given by

$$\begin{aligned} \star : \mathcal{O}_K^n \times \mathcal{O}_K^n &\rightarrow \mathcal{O}_K^n \\ \bar{x} \star \bar{y} &= \sum_{I, J} \text{st}(\bar{c}_{I, J}) \bar{x}^I \bar{y}^J. \end{aligned}$$

Therefore the induced structure is a Lie group of dimension n . □

Proposition 7.12. *Suppose $G = (\mathcal{O}^d, \star)$ satisfies the conditions \mathfrak{A}_ω , \mathfrak{B}_1 , \mathfrak{B}_2 , \mathfrak{D} , and \mathfrak{C} . Then G is fsg and compactly dominated, and G/G^{00} is isomorphic to a de -dimensional Lie group over \mathbb{Q}_p , where $e = [K : \mathbb{Q}_p]$.*

Proof. Proposition 7.8 gives the first part, and shows that $G^{00} = \bigcap_{k < \omega} p^k \mathcal{O}^n$. Then G/G^{00} is canonically isomorphic to a topological group $(\mathcal{O}_K^d, *)$. The assumptions on \star precisely imply that $*$ satisfies all the requirements in Lemma 6.2, and so G/G^{00} is a de -dimensional Lie group over \mathbb{Q}_p . □

8 The first two proofs of the Onshuus-Pillay conjecture

Lemma 8.1. *Suppose $M \models T$ and $D \subseteq M^n$ is definable, with $\dim(D) = k$. Then there is a definable injection $D \rightarrow M^k$.*

Proof. Write $X \leq Y$ if there is a definable injection from X to Y . Note that $M \leq \mathcal{O} \times \{0, 1\}$, because of the map

$$f(x) = \begin{cases} (x, 0) & \text{if } x \in \mathcal{O} \\ (1/x, 1) & \text{if } x \notin \mathcal{O}. \end{cases}$$

Thus $M \leq \mathcal{O} + \mathcal{O}$, where $+$ denotes disjoint union. On the other hand, $\mathcal{O} \leq B$ and $B \leq \mathcal{O}$ for any ball B , and we can find two disjoint balls inside \mathcal{O} , so $\mathcal{O} + \mathcal{O} \leq B + B \leq \mathcal{O}$. Then $M \leq \mathcal{O}$, so

$$M + M \leq \mathcal{O} + \mathcal{O} \leq \mathcal{O} \leq M.$$

Multiplying by M^{n-1} , we see that $M^k + M^k \leq M^k$ for any k . And of course $M^j \leq M^k$ for $j < k$. In light of this, it suffices to cover D with finitely many definable sets D_i , such that $D_i \leq M^{k_i}$ for some $k_i \leq \dim(D)$. Such a decomposition is provided by the “topological cell decomposition” of [5]. \square

If (G, \star) is a definable group and $a \in G$, define

$$\begin{aligned} \star_a : G \times G &\rightarrow G \\ x \star_a y &= xa^{-1}y. \end{aligned}$$

Then (G, \star_a) is a definable group with identity element a , definably isomorphic to (G, \star) by the following isomorphism:

$$\begin{aligned} (G, \star) &\rightarrow (G, \star_a) \\ x &\mapsto ax. \end{aligned}$$

In particular, if we prove Conjecture 1.2 for (G, \star_a) , then we prove it for (G, \star) .

Lemma 8.2. *Let $G \subseteq \mathbb{M}^n$ be an n -dimensional definable set and \star be a definable group operation on G . Then there is a definable set U in the interior of G with $\dim(G \setminus U) < n$, such that for any $a \in U$, the group operation \star_a is strictly differentiable on a neighborhood of a .*

Proof. Morally, this comes from the fact that definable functions in \mathbb{M} are generically strictly differentiable (Theorem 5.14). However, the proof is not completely trivial, and we should give details.

Conceptually, the correct proof is to consider the canonical strict C^1 -manifold structure on G . Apply generic differentiability to the map from G (with its strict C^1 -manifold structure) to G (as a subset of \mathbb{M}^n). Fix a point a where this map is strictly differentiable. Then an open neighborhood $a \in U \subseteq \mathbb{M}^n$ is also a local chart for the manifold structure. (That is,

the two C^1 -structures on G agree with each other close to the point a .) Because the group structure is C^1 , the function $(x, y) \mapsto x \star a^{-1} \star y$ is strictly differentiable.

Rather than formally developing strict C^1 -manifolds, we instead give a more direct proof. By generic strict differentiability, there is a definable open set Δ in the interior of $G \times G \subseteq \mathbb{M}^{2n}$, such that $\dim((G \times G) \setminus \Delta) < \dim(G \times G) = 2n$, and the group operations are strictly differentiable on Δ . Let ∇ be the set of pairs $(a, b) \in G \times G$ such that the following conditions hold:

$$\begin{aligned} (b^{-1}, a) &\in \Delta \\ (b^{-1} \star a, a^{-1}) &\in \Delta \\ (b, b^{-1} \star a) &\in \Delta. \end{aligned}$$

Let M_0 be a small model over which everything is defined. If the pair $(a, b) \in G^2$ is jointly generic, in the sense that $\dim(a, b/M_0) = 2 \dim(G)$, then so are the pairs (b^{-1}, a) , $(b^{-1} \star a, a^{-1})$, $(b, b^{-1} \star a)$, and so all three pairs belong to Δ , and $(a, b) \in \nabla$. Let U be the projection of ∇ onto the first coordinate, intersected with the interior of G :

$$U = \{a : (a, b) \in \nabla\} \cap \text{int}(G).$$

If $a \in G$ is generic over M_0 , we can find $b \in G$ generic over $M_0 a$, and then (a, b) is generic over M_0 , $(a, b) \in \nabla$, and $a \in U$. It follows that U contains all $a \in G$ which are generic over M_0 , and so $\dim(G \setminus U) < \dim(G)$.

For $a \in U$, we will prove that \star_a is strictly differentiable around a . Take $b \in G$ such that $(a, b) \in \nabla$. Then the composition

$$b \star \left(\left((b^{-1} \star x) \star a^{-1} \right) \star y \right) = x \star_a y$$

is strictly differentiable for $(x, y) \approx (a, a)$, because for such x and y we have

$$\begin{aligned} (b^{-1}, x) &\approx (b^{-1}, a) \in \Delta \\ \left((b^{-1} \star x), a^{-1} \right) &\approx (b^{-1} \star a, a^{-1}) \in \Delta \\ \left(\left((b^{-1} \star x) \star a^{-1} \right), y \right) &\approx (b^{-1} \star a \star a^{-1}, a) = (b^{-1}, a) \in \Delta \\ \left(b, \left(\left((b^{-1} \star x) \star a^{-1} \right) \star y \right) \right) &\approx (b, b^{-1} \star a) \in \Delta. \end{aligned} \quad \square$$

Lemma 8.3. *Let $G \subseteq \mathbb{M}^n$ be an n -dimensional definable set and \star be a definable group operation on G . Let a be the identity element of G , and suppose a is in the interior of G and \star is strictly differentiable at (a, a) . Then there is some non-zero ϵ such that $a + \epsilon \mathcal{O}^n$ is a subgroup of G and $a + \epsilon \mathcal{O}^n$ satisfies conditions \mathfrak{A}_∞ , \mathfrak{B}_ω , \mathfrak{D} and \mathfrak{E} . In fact, any sufficiently small ϵ works.*

Proof. Moving G by a translation, we may assume $a = \bar{0}$. Let $f : G \rightarrow G$ be the p th power map. A straightforward calculation shows that at $\bar{0}$, the functions $\bar{x} \star \bar{y}$, \bar{x}^{-1} , and $f(\bar{x})$ have

the following strict derivatives:

$$\begin{aligned}\frac{\partial}{\partial \bar{x}}(\bar{x} \star \bar{y}) &= I_n \\ \frac{\partial}{\partial \bar{y}}(\bar{x} \star \bar{y}) &= I_n \\ \frac{\partial}{\partial \bar{x}}\bar{x}^{-1} &= -I_n \\ \frac{\partial}{\partial \bar{x}}f(\bar{x}) &= pI_n,\end{aligned}$$

where I_n is the $n \times n$ identity matrix.⁷

Take a small model M defining (G, \star) . Let $\epsilon \in \mathbb{M}$ be a non-zero M -infinitesimal, i.e., a non-zero element satisfying the equivalent conditions:

- ϵ is contained in any M -definable neighborhood of 0.
- ϵ is contained in any M -definable ball $B_\gamma(0)$.
- $v(\epsilon) > \Gamma_M$.

Note that ϵa is an M -infinitesimal for any $a \in \mathcal{O} = \mathcal{O}_{\mathbb{M}}$, including a outside of M .

We first verify that $\epsilon\mathcal{O}^n$ is a subgroup of G . Let \bar{x}, \bar{y} be tuples in $\epsilon\mathcal{O}^n$. Then \bar{x} and \bar{y} are tuples of M -infinitesimals. By differentiability,

$$\bar{x} \star \bar{y} = \bar{x} + \bar{y} + \mu_1 \bar{x} + \mu_2 \bar{y}$$

for some $n \times n$ matrices μ_1, μ_2 of M -infinitesimals. The right hand side is in $\epsilon\mathcal{O}^n$. A similar argument shows that $\bar{x} \in \epsilon\mathcal{O}^n \implies \bar{x}^{-1} \in \epsilon\mathcal{O}^n$. Now we check each of the required conditions:

(\mathfrak{A}_∞) Let I be a principal ideal contained in $\epsilon\mathcal{O}$. We must show that the group operation \star on $\epsilon\mathcal{O}^n$ respects congruence modulo I^n . Suppose $\bar{x}, \bar{y}, \bar{z} \in \epsilon\mathcal{O}^n$ with $\bar{x} \equiv \bar{y} \pmod{I^n}$. Then $\bar{x}, \bar{y}, \bar{z}$ are tuples of M -infinitesimals. By *strict* differentiability of the group operation,

$$(\bar{x} \star \bar{z}) - (\bar{y} \star \bar{z}) = (\bar{x} - \bar{y}) + \mu(\bar{x} - \bar{y})$$

for some matrix μ of M -infinitesimals. The right hand side is in I^n , so $\bar{x} \star \bar{z} \equiv \bar{y} \star \bar{z} \pmod{I^n}$. Similar arguments show

$$\begin{aligned}\bar{y} \equiv \bar{z} \pmod{I^n} &\implies \bar{x} \star \bar{y} \equiv \bar{x} \star \bar{z} \pmod{I^n} \\ \bar{x} \equiv \bar{y} \pmod{I^n} &\implies \bar{x}^{-1} \equiv \bar{y}^{-1} \pmod{I^n}.\end{aligned}$$

Thus, congruence modulo I^n is a congruence in the group $(\epsilon\mathcal{O}^n, \star)$, which is condition (\mathfrak{A}_∞).

⁷For the group inverse map \bar{x}^{-1} , strict differentiability follows by applying the inverse function theorem, Theorem 5.18, to the map $(\bar{x}, \bar{y}) \mapsto (\bar{x}, \bar{x} \star \bar{y})$.

(\mathfrak{B}_ω) If \bar{x}, \bar{y} are tuples in $\epsilon\mathcal{O}^n$, then \bar{x}, \bar{y} are M -infinitesimals, so differentiability of \star gives

$$\bar{x} \star \bar{y} = \bar{x} + \bar{y} + \mu_1 \bar{x} + \mu_2 \bar{y}$$

for some M -infinitesimal matrices μ_1, μ_2 . The entries in μ_1, μ_2 have valuation greater than $v(p^k)$ for any k , so

$$\bar{x} \star \bar{y} - \bar{x} - \bar{y} \in p^k \epsilon\mathcal{O}^n.$$

This is condition (\mathfrak{B}_k), for arbitrary k .

(\mathfrak{D}) If \bar{x}, \bar{y} are in $\epsilon\mathcal{O}^n$, then they are M -infinitesimal, and so the strong differentiability of f shows that

$$f(\bar{x}) - f(\bar{y}) = p(\bar{x} - \bar{y}) + \mu(\bar{x} - \bar{y})$$

for some matrix μ of M -infinitesimals. The right hand side has valuation $v(p) + v(\bar{x} - \bar{y})$ because μ is infinitesimal compared to p . Thus condition \mathfrak{D} holds.

(\mathfrak{E}) The derivative of f at $\bar{0}$ is pI_n , which is invertible, so the inverse function theorem (Theorem 5.18) shows that f maps the set of M -infinitesimal vectors onto the set of M -infinitesimal vectors. If $\bar{x} \in p\epsilon\mathcal{O}^d$, then \bar{x} is an M -infinitesimal vector, so $\bar{x} = f(\bar{y})$ for some M -infinitesimal vector \bar{y} . By the differentiability of f at $\bar{0}$,

$$\bar{x} = f(\bar{y}) = p\bar{y} + \mu\bar{y}$$

for some M -infinitesimal matrix μ . Then $v(\bar{x}) = v(p) + v(\bar{y})$ so $\bar{y} \in p^{-1}p\epsilon\mathcal{O}^n = \epsilon\mathcal{O}^n$. Thus every element of $p\epsilon\mathcal{O}^n$ is a p th power of an element in $\epsilon\mathcal{O}^n$, which is condition \mathfrak{E} . \square

Theorem 8.4. *Let G be an n -dimensional definable group in a highly saturated model \mathbb{M} of a P -minimal expansion T of $\text{Th}(K)$ for some finite extension K/\mathbb{Q}_p . Then there is a definable open subgroup $H \subseteq G$ such that H is fsg, $H/H^{00} \cong \mathcal{O}_K^n$, and H is compactly dominated by H/H^{00} . In particular, H/H^{00} is an n -dimensional Lie group over K .*

Proof. By Lemma 8.1, we may assume $G \subseteq \mathbb{M}^n$. By Lemma 8.2 we may assume that the group operation is differentiable at the identity element. Lemma 8.3 gives an n -dimensional definable subgroup $H \subseteq G$ which (up to isomorphism) satisfies \mathfrak{A}_∞ , \mathfrak{B}_ω , \mathfrak{D} , and \mathfrak{E} . In particular, it satisfies \mathfrak{B}_ω . Proposition 7.10 shows that H has the desired properties. \square

Theorem 8.5. *Let M be a model of a P -minimal expansion T of $\text{Th}(K)$ for some finite extension K/\mathbb{Q}_p with $e = [K : \mathbb{Q}_p]$. In a monster model $\mathbb{M} \succeq M$, let G be an n -dimensional M -definable group. Then there is an M -definable open subgroup $H \subseteq G$ such that H is fsg, H/H^{00} is an ne -dimensional Lie group over \mathbb{Q}_p and H is compactly dominated by H/H^{00} .*

Proof. As in Theorem 8.4, we can find a n -dimensional definable subgroup $H \subseteq G$ which satisfies \mathfrak{A}_∞ , \mathfrak{B}_ω , \mathfrak{D} , and \mathfrak{E} . In particular, it satisfies the combination $\mathfrak{A}_\infty \wedge \mathfrak{B}_1 \wedge \mathfrak{B}_2 \wedge \mathfrak{D} \wedge \mathfrak{E}$. This combination of properties is definable (Remark 7.3), so we can change H to be M -definable. Then Proposition 7.12 shows that H has the desired properties. \square

9 Third proof, for pure p -adically closed fields

In this section, we restrict to the case where $T = \text{Th}(K)$, i.e., pure p -adically closed fields. Recall that K is a fixed p -adic field, i.e., a finite extension of \mathbb{Q}_p . We assume the language \mathcal{L} contains constant symbols naming a basis of K over \mathbb{Q}_p . This ensures that the theory has definable Skolem functions and that $\text{dcl}(\emptyset)$ is dense in K . The density implies that every K -definable open ball is 0-definable, a fact we will need later.

9.1 Splendid and locally splendid functions

Definition 9.1. A function $f : \mathcal{O}_K^n \rightarrow \mathcal{O}_K^m$ is *pre-splendid* if it's given by a convergent power series with coefficients in \mathcal{O}_K . That is,

$$f(\bar{x}) = \sum_I \bar{c}_I \bar{x}^I$$

where $\bar{c}_I \in \mathcal{O}^m$ and $\lim_{|I| \rightarrow \infty} v(\bar{c}_I) = +\infty$.

- Observation 9.2.**
1. If $f : \mathcal{O}_K^n \rightarrow \mathcal{O}_K^m$ is constant, then f is pre-splendid.
 2. The identity function $\mathcal{O}_K^n \rightarrow \mathcal{O}_K^n$ and coordinate projections $\pi_1, \dots, \pi_n : \mathcal{O}_K^n \rightarrow \mathcal{O}_K$ are pre-splendid. More generally, any polynomial map $f : \mathcal{O}_K^n \rightarrow \mathcal{O}_K^m$ is pre-splendid, *provided* that the coefficients are in \mathcal{O}_K .
 3. If $f : \mathcal{O}_K^n \rightarrow \mathcal{O}_K^m$ and $g : \mathcal{O}_K^m \rightarrow \mathcal{O}_K^\ell$ are pre-splendid, then the composition $g \circ f : \mathcal{O}_K^n \rightarrow \mathcal{O}_K^\ell$ is pre-splendid.
 4. If $f : \mathcal{O}_K^n \rightarrow \mathcal{O}_K^m$ is a function, then f is pre-splendid if and only if the component functions $f_1, \dots, f_m : \mathcal{O}_K^n \rightarrow \mathcal{O}_K$ are pre-splendid.
 5. The set of pre-splendid functions from \mathcal{O}_K^n to \mathcal{O}_K is an \mathcal{O}_K -algebra, i.e., it is closed under the ring operations and multiplication by \mathcal{O}_K .
 6. If $f : \mathcal{O}_K^n \rightarrow \mathcal{O}_K^m$ is pre-splendid, then f is strictly differentiable, and the strict derivative $Df : \mathcal{O}_K^n \rightarrow \mathcal{O}_K^{nm}$ is pre-splendid.

Now consider a monster model $\mathbb{M} \succ K$. Recall that \mathcal{O} denotes $\mathcal{O}_{\mathbb{M}}$.

Definition 9.3. Let $\mathcal{F} = \{f_a\}_{a \in D}$ be a 0-definable family of functions from $\mathcal{O}^n \rightarrow \mathcal{O}^m$. Then \mathcal{F} is a *splendid family* if for every $a \in D(K)$, the function $f_a(K) : \mathcal{O}_K^n \rightarrow \mathcal{O}_K^m$ is pre-splendid. A definable function $f : \mathcal{O}^n \rightarrow \mathcal{O}^m$ is *splendid* if it belongs to some splendid family.

We can transfer the properties of pre-splendid functions in Observation 9.2 to splendid functions:

Proposition 9.4. 1. If $f : \mathcal{O}^n \rightarrow \mathcal{O}^m$ is constant, then f is splendid.

2. If $f : \mathcal{O}^n \rightarrow \mathcal{O}^m$ is a polynomial map whose coefficients are in \mathcal{O} , then f is splendid. For example, the identity map $\mathcal{O}^n \rightarrow \mathcal{O}^n$ and coordinate projections $\pi_1, \dots, \pi_n : \mathcal{O}^n \rightarrow \mathcal{O}$ are splendid.
3. If $f : \mathcal{O}^n \rightarrow \mathcal{O}^m$ and $g : \mathcal{O}^m \rightarrow \mathcal{O}^\ell$ are splendid, then the composition $g \circ f : \mathcal{O}^n \rightarrow \mathcal{O}^\ell$ is splendid.
4. If $f : \mathcal{O}^n \rightarrow \mathcal{O}^m$ is a definable function, then f is splendid if and only if the component functions $f_1, \dots, f_m : \mathcal{O}^n \rightarrow \mathcal{O}$ are splendid.
5. The set of splendid functions from \mathcal{O}^n to \mathcal{O} is an \mathcal{O} -algebra, i.e., it is closed under the ring operations and multiplication by \mathcal{O} .
6. If $f : \mathcal{O}^n \rightarrow \mathcal{O}^m$ is splendid, then f is strictly differentiable, and the strict derivative $Df : \mathcal{O}^n \rightarrow \mathcal{O}^{nm}$ is splendid.

Proof. 1. The family of constant functions from $\mathcal{O}^n \rightarrow \mathcal{O}^m$ is a splendid family, by Observation 9.2(1).

2. Suppose $f : \mathcal{O}^n \rightarrow \mathcal{O}^m$ is polynomial, with all coefficients in \mathcal{O} . Let d be the degree of f , and let $\mathcal{F}_{\leq d}$ be the family of all polynomial maps $g : \mathcal{O}^n \rightarrow \mathcal{O}^m$ such that the coefficients of g are in \mathcal{O} and $\deg(g) \leq d$. Then $\mathcal{F}_{\leq d}$ is a 0-definable family (because of the bound on degree) and $\mathcal{F}_{\leq d}$ is a splendid family by Observation 9.2(2).
3. Take splendid families $\mathcal{F} \ni f$ and $\mathcal{G} \ni g$. By Observation 9.2(3), the family $\{g' \circ f' : f' \in \mathcal{F}, g' \in \mathcal{G}\}$ is a splendid family containing $g \circ f$.
4. Similar.
5. Similar.
6. Take a splendid family $\mathcal{F} \ni f$. By Observation 9.2(6), every function in $\mathcal{F}(K)$ is strictly differentiable. Because $K \preceq \mathbb{M}$, this implies the strict differentiability of functions in \mathcal{F} (including f). Let $\mathcal{F}' = \{Dg : g \in \mathcal{F}\}$, the set of strict derivatives of functions in \mathcal{F} . Then \mathcal{F}' is a splendid family by Observation 9.2(6), so Df is splendid. \square

Additionally, 0-definable pre-splendid functions are splendid:

Proposition 9.5. *If $f : \mathcal{O}^n \rightarrow \mathcal{O}^m$ is 0-definable and $f(K) : \mathcal{O}_K^n \rightarrow \mathcal{O}_K^m$ is pre-splendid, then f is splendid.*

Proof. The singleton family $\{f\}$ is a splendid family. \square

Lemma 9.6. *Let $f : \mathcal{O}^n \rightarrow \mathcal{O}^m$ be splendid, with $f(\bar{0}) = \bar{0}$. Suppose $\epsilon \in \mathcal{O}$ is non-zero. Let $g : \mathcal{O}^n \rightarrow \mathbb{M}^m$ be the function $g(\bar{x}) = f(\epsilon\bar{x})/\epsilon$. Then $\text{im}(g) \subseteq \mathcal{O}^m$, and the function $g : \mathcal{O}^n \rightarrow \mathcal{O}^m$ is splendid.*

Proof. It suffices to check the analogous statement for pre-splendid functions. Let $f : \mathcal{O}_K^n \rightarrow \mathcal{O}_K^m$ be pre-splendid with $f(\bar{0}) = \bar{0}$. Then $f(\bar{x}) = \sum_I \bar{c}_I \bar{x}^I$, where each \bar{c}_I is in \mathcal{O}_K^m and the constant term \bar{c}_0 vanishes. Then

$$g(\bar{x}) = f(\epsilon \bar{x})/\epsilon = \sum_I \epsilon^{|I|-1} \bar{c}_I \bar{x}^I.$$

All the coefficients $\epsilon^{|I|-1} \bar{c}_I$ are still in \mathcal{O} , since the constant term vanishes. Thus g is pre-splendid. \square

Definition 9.7. Let $U \subseteq \mathbb{M}^n$ be open and $f : U \rightarrow \mathbb{M}^m$ be a definable function. Then f is *locally splendid* at $a \in U$ if there are some non-zero ϵ and δ such that

$$\begin{aligned} a + \epsilon \mathcal{O}^n &\subseteq U \\ f(a + \epsilon \mathcal{O}^n) &\subseteq f(a) + \delta \mathcal{O}^m \end{aligned}$$

and the function

$$a + \epsilon \mathcal{O}^n \xrightarrow{f} f(a) + \delta \mathcal{O}^m$$

is splendid, or more precisely, the following function is splendid:

$$\begin{aligned} \mathcal{O}^n &\rightarrow \mathcal{O}^m \\ x &\mapsto (f(a + \epsilon x) - f(a))\delta^{-1}. \end{aligned}$$

Finally, we say that f is *locally splendid* if it is splendid at every $a \in U$.

We first carry out some sanity checks: splendid functions are locally splendid, and local splendidness is really a local property.

Proposition 9.8. *If $f : \mathcal{O}^n \rightarrow \mathcal{O}^m$ is splendid, then f is locally splendid.*

Proof. For any $a \in \mathcal{O}^n$, take $\epsilon = \delta = 1$. Then

$$\begin{aligned} a + \mathcal{O}^n &= \mathcal{O}^n = \text{dom}(f) \\ f(a + \mathcal{O}^n) &= f(\mathcal{O}^n) = \text{im}(f) \subseteq \mathcal{O}^m = f(a) + \mathcal{O}^m, \end{aligned}$$

and the function

$$\begin{aligned} \mathcal{O}^n &\rightarrow \mathcal{O}^m \\ x &\mapsto f(a + x) - f(a) \end{aligned}$$

is splendid by Proposition 9.4. \square

Lemma 9.9. *Fix a definable function $f : U \rightarrow \mathbb{M}^n$ and a point $a \in U$. Say that a pair (ϵ, δ) is “suitable” if it satisfies the conditions in Definition 9.7:*

$$\begin{aligned} a + \epsilon \mathcal{O}^n &\subseteq U \\ f(a + \epsilon \mathcal{O}^n) &\subseteq f(a) + \delta \mathcal{O}^m \end{aligned}$$

and the following function is splendid:

$$\begin{aligned} \mathcal{O}^n &\rightarrow \mathcal{O}^m \\ x &\mapsto (f(a + \epsilon x) - f(a))\delta^{-1}. \end{aligned}$$

Suppose that f is locally splendid at a , i.e., some pair (ϵ_0, δ_0) is suitable.

1. If (ϵ, δ) is suitable, then (ϵ', δ) is suitable for any ϵ' with $v(\epsilon') \geq v(\epsilon)$.
2. If (ϵ, δ) is suitable, then (ϵ, δ') is suitable for any δ' with $v(\delta') \leq v(\delta)$.
3. For any δ , there is some ϵ such that (ϵ, δ) is suitable.

Proof. 1. Let $g : \mathcal{O}^n \rightarrow \mathcal{O}^m$ be the map $g(x) = (f(a + \epsilon x) - f(a))\delta^{-1}$. The multiplication by ϵ'/ϵ map $\mathcal{O}^n \rightarrow \mathcal{O}^n$ is splendid, so the composition

$$g(x\epsilon'/\epsilon) = (f(a + \epsilon'x) - f(a))\delta^{-1}$$

is splendid, showing that (ϵ', δ) is suitable.

2. Similar, using the fact that the multiplication by δ/δ' map $\mathcal{O}^m \rightarrow \mathcal{O}^m$ is splendid.
3. If $v(\delta) \leq v(\delta_0)$, then (ϵ_0, δ) is splendid by the previous point. Suppose $v(\delta) \geq v(\delta_0)$. Let $\rho = \delta/\delta_0 \in \mathcal{O}$. Because the pair (ϵ_0, δ_0) is suitable, the following function is splendid:

$$g(x) = (f(a + \epsilon_0 x) - f(a))\delta_0^{-1}.$$

Note that $g(0) = 0$. By Lemma 9.6, the composition

$$g(\rho x)\rho^{-1} = (f(a + \rho\epsilon_0 x) - f(a))\delta^{-1}$$

is splendid. Thus $(\rho\epsilon_0, \delta)$ is suitable. \square

Lemma 9.9(3) shows that in the definition of “locally splendid” (Definition 9.7), we could have said “for every δ there is ϵ ” rather than “there exist δ and ϵ ”.

Proposition 9.10. *Let U, U', U_1, \dots, U_k be open sets in \mathbb{M}^n .*

1. *If $a \in U' \subseteq U$ and $f : U \rightarrow \mathbb{M}^m$ is definable, then f is locally splendid at a iff $f \upharpoonright U'$ is locally splendid at a .*
2. *If $U = U_1 \cup \dots \cup U_k$ and $f : U \rightarrow \mathbb{M}^m$ is definable, then f is locally splendid if and only if $f \upharpoonright U_i$ is locally splendid for each i .*

Proof. Clearly (1) implies (2). We prove (1). First suppose f is splendid at a . Take (ϵ, δ) which is suitable for f in the sense of Lemma 9.9. Take ϵ' so small that $a + \epsilon'\mathcal{O}^n \subseteq U'$ and $v(\epsilon') \geq v(\epsilon)$. By Lemma 9.9(1), (ϵ', δ) is suitable for f , which implies it is suitable for $f \upharpoonright U'$.

Conversely, suppose $f \upharpoonright U$ is splendid at a . If (ϵ, δ) is suitable for $f \upharpoonright U$, then (ϵ, δ) is suitable for f . \square

Lemma 9.11. *Suppose $f : U \rightarrow \mathbb{M}^m$ is locally splendid at $a \in U$. If f, U, a are M -definable for some small $M \preceq \mathbb{M}$, then we can take the (ϵ, δ) witnessing local splendidness to be M -definable.*

Proof. Take some (ϵ_0, δ_0) which is suitable for f at a . Let \mathcal{F} be a splendid family containing the splendid function

$$x \mapsto (f(a + \epsilon_0 x) - f(a))\delta_0^{-1}.$$

In particular, \mathcal{F} is 0-definable. Let D be the set of pairs (ϵ, δ) such that

$$\begin{aligned} a + \epsilon \mathcal{O}^n &\subseteq U \\ f(a + \epsilon \mathcal{O}^n) &\subseteq f(a) + \delta \mathcal{O}^m, \end{aligned}$$

and the function

$$x \mapsto (f(a + \epsilon x) - f(a))\delta^{-1}$$

is in \mathcal{F} . Then D is M -definable and contains (ϵ_0, δ_0) . By Tarski-Vaught, there is some M -definable pair (ϵ, δ) in D . Then (ϵ, δ) is suitable, i.e., (ϵ, δ) witnesses local splendidness of f at a . \square

Remark 9.12. Recall that a family \mathcal{F} is *ind-definable* if it is a small union of definable families. For example, the family of splendid functions is ind-definable by definition. The family of locally splendid functions is also ind-definable. Since we will not need this fact, we leave the proof as an exercise to the reader.

Locally splendid functions are closed under similar operations as splendid functions:

Proposition 9.13. *Let $U \subseteq \mathbb{M}^n$ be open and $f : U \rightarrow \mathbb{M}^m$ be definable.*

1. *If f is a polynomial map, then f is locally splendid.*
2. *If $f : U \rightarrow V$ is locally splendid and $g : V \rightarrow \mathbb{M}^\ell$ is locally splendid, then $g \circ f : U \rightarrow \mathbb{M}^\ell$ is locally splendid.*
3. *f is locally splendid iff the component functions $f_1, \dots, f_m : U \rightarrow \mathbb{M}$ are locally splendid.*
4. *The set of splendid functions $U \rightarrow \mathbb{M}$ is an \mathbb{M} -algebra, i.e., closed under the ring operations and multiplication by \mathbb{M} .*
5. *If $f : U \rightarrow \mathbb{M}^m$ is locally splendid, then f is strictly differentiable and the strict derivative $Df : U \rightarrow \mathbb{M}^{nm}$ is locally splendid.*

Proof. 1. Given $a \in U$, take ϵ so small that $a + \epsilon \mathcal{O}^n \subseteq U$. The map

$$x \mapsto f(a + \epsilon x) - f(a)$$

is polynomial, so if we take δ very large, then $x \mapsto (f(a + \epsilon x) - f(a))\delta^{-1}$ will be polynomial with coefficients in \mathcal{O} , and therefore splendid.

2. Take any non-zero δ_3 . By Lemma 9.9(3) there is δ_2 such that

$$\begin{aligned} f(a) + \delta_2 \mathcal{O}^m &\subseteq V \\ g(f(a) + \delta_2 \mathcal{O}^m) &\subseteq g(f(a)) + \delta_3 \mathcal{O}^\ell \end{aligned}$$

and the following map is splendid.

$$g : (f(a) + \delta_2 \mathcal{O}^m) \rightarrow (g(f(a)) + \delta_3 \mathcal{O}^\ell)$$

By another application of Lemma 9.9(3) there is δ_1 such that

$$\begin{aligned} a + \delta_1 \mathcal{O}^n &\subseteq U \\ f(a + \delta_1 \mathcal{O}^n) &\subseteq f(a) + \delta_2 \mathcal{O}^m \end{aligned}$$

and the following map is splendid

$$f : (a + \delta_1 \mathcal{O}^n) \rightarrow (f(a) + \delta_2 \mathcal{O}^m).$$

Then

$$\begin{aligned} a + \delta_1 \mathcal{O}^n &\subseteq U \\ g(f(a + \delta_1 \mathcal{O}^n)) &\subseteq g(f(a) + \delta_2 \mathcal{O}^m) \subseteq g(f(a)) + \delta_3 \mathcal{O}^\ell \end{aligned}$$

and the composition

$$g \circ f : (a + \delta_1 \mathcal{O}^n) \rightarrow (g(f(a)) + \delta_3 \mathcal{O}^\ell)$$

is splendid. Then (δ_1, δ_3) shows that $g \circ f$ is locally splendid at a .

3. If f is locally splendid, then the components of f are locally splendid by the previous two points (compose f with the coordinate projections $\mathbb{M}^m \rightarrow \mathbb{M}$). Conversely, suppose the components f_1, \dots, f_m are locally splendid. Fix $a \in U$. For each i , there is a pair (ϵ_i, δ_i) which is “suitable” for f_i at a , in the sense of Lemma 9.9. Take ϵ and δ such that $v(\epsilon) \geq \max_i v(\epsilon_i)$ and $v(\delta) \leq \min_i v(\delta_i)$. Then (ϵ, δ) is suitable for each f_i at a by Lemma 9.8(1,2). In particular,

$$\begin{aligned} a + \epsilon \mathcal{O}^n &\subseteq U \\ f_i(a + \epsilon \mathcal{O}^n) &\subseteq f_i(a) + \delta \mathcal{O} \end{aligned} \tag{*}$$

and the map $f_i : (a + \epsilon \mathcal{O}^n) \rightarrow (f_i(a) + \delta \mathcal{O})$ is splendid. Equation $(*)$ implies $f(a + \epsilon \mathcal{O}^n) \subseteq f(a) + \delta \mathcal{O}^m$. The map $f : (a + \epsilon \mathcal{O}^n) \rightarrow (f(a) + \delta \mathcal{O}^m)$ is splendid because each component is splendid.

4. This follows from the previous three points.
5. Strict differentiability is clear because, up to a change of coordinate, f looks locally like a splendid function, and splendid functions are strictly differentiable. Locally, the strict derivative is a scaled version of the strict derivative of a splendid function, so the strict derivative is locally splendid. \square

9.2 Generic local splendidness

Next, we work towards proving that definable functions are generically locally splendid. To do this, we will essentially show that the theory of p -adically closed fields has locally splendid definable Skolem functions.

Say that a function on K is *strongly analytic* if it is given by a single convergent power series, and *analytic* if it is locally strongly analytic. (Some authors say “locally analytic” rather than “analytic”.)

Lemma 9.14. *Let $f : U \rightarrow \mathbb{M}^m$ be a 0-definable function such that $U \subseteq \mathbb{M}^n$ is open, $U(K)$ is compact, and $f(K) : U(K) \rightarrow K^m$ is analytic. Then f is locally splendid.*

Proof. Because K is locally compact and $U(K)$ is compact, we can cover U with K -definable balls $U = B_1 \cup \dots \cup B_\ell$ such that $f(K)$ is strongly analytic on each $B_i(K)$. By the assumption on \mathcal{L} at the start of Section 9, the balls B_i are 0-definable. Each of the restrictions $f(K) \upharpoonright B_i(K)$ is strongly analytic, given by a convergent power series. Replacing f with $\delta \cdot f$ for a small enough non-zero $\delta \in \text{dcl}(\emptyset)$, we can arrange that all the coefficients in these power series are in \mathcal{O}_K . Then $f(K) \upharpoonright B_i(K)$ is (essentially) a 0-definable pre-splendid function for each i , and so $f \upharpoonright B_i$ is splendid (Proposition 9.5). By Proposition 9.10(2), f is locally splendid. \square

Lemma 9.15. *The field operations are locally splendid on their domains.*

Proof. Addition, multiplication, and subtraction are polynomials, so they are locally splendid by Proposition 9.13(1). It remains to show that the map $f(x) = 1/x$ is locally splendid on \mathbb{M}^\times . Lemma 9.14 shows that f is locally splendid on the subset \mathcal{O}^\times . Then $f : c\mathcal{O}^\times \rightarrow c^{-1}\mathcal{O}^\times$ is also locally splendid for any $c \in \mathbb{M}^\times$, because it's the composition

$$c\mathcal{O}^\times \xrightarrow{x \mapsto x/c} \mathcal{O}^\times \xrightarrow{f} \mathcal{O}^\times \xrightarrow{x \mapsto x/c} c^{-1}\mathcal{O}^\times$$

and the two maps on the outside are polynomial. Then we have covered $\text{dom}(f)$ with sets $c\mathcal{O}^\times$ on which f is locally splendid, so f is locally splendid. \square

Lemma 9.16. *1. Let \mathfrak{m} denote the maximal ideal. Let*

$$h_n : \mathfrak{m} \times \mathcal{O}^\times \times \mathcal{O}^{n-2} \rightarrow \mathfrak{m}$$

be the definable function mapping $(a_0, a_1, \dots, a_{n-1})$ to the unique root in \mathfrak{m} of

$$x^n + a_{n-1}x^{n-1} + \dots + a_0 = 0,$$

the root guaranteed to exist by Hensel's lemma. Then h_n is locally splendid.

2. Let P_n denote the set of non-zero n th powers. Then there is a locally splendid 0-definable map

$$\begin{aligned} P_n &\rightarrow \mathbb{M} \\ x &\mapsto \sqrt[n]{x} \end{aligned}$$

assigning an n th root to each $x \in P_n$.

In (2), we are not claiming that $\sqrt[n]{x}$ is a $\frac{1}{n}$ th power map in the sense of Definition 3.6, i.e., $\sqrt[n]{xy}$ need not equal $\sqrt[n]{x}\sqrt[n]{y}$.

Proof. 1. The function $h_n(K) : \mathfrak{m}_K \times \mathcal{O}_K^\times \times \mathcal{O}_K^{n-2} \rightarrow \mathfrak{m}_K$ is analytic, by the implicit function theorem for analytic functions. (See [21, Proposition 5.9], at least for the *inverse* function theorem for analytic functions, which easily implies the implicit function theorem.) Then Lemma 9.14 shows that h_n is locally splendid.

2. By Lemma 3.8, there is some m such that P_m has an n th power map $x^{1/n}$ in the sense of Definition 3.6. Increasing m , we may assume $n \mid m$. We first show that this map $x^{1/n}$ on P_m is locally splendid.

The map $x^{1/n}$ is analytic on $P_m(K)$ by the inverse function theorem for analytic functions.⁸ However, $P_m(K)$ is not compact, so we cannot immediately apply Lemma 9.14. Nevertheless, the subset $P_m(K) \cap \mathcal{O}_K^\times$ is compact, so we at least see that $x^{1/n}$ is locally splendid on $P_m \cap \mathcal{O}^\times$ by Lemma 9.14. Using the same method as in Lemma 9.15, this implies that $x^{1/n}$ is locally splendid on $P_m \cap c\mathcal{O}^\times$ for every $c \in P_m \cap \mathcal{O}^\times$, and so $x^{1/n}$ is locally splendid on P_m .

Finally, take a_1, \dots, a_k coset representatives of P_m in P_n , and take b_i to be any n th root of a_i . By definable Skolem functions, we can take $a_i, b_i \in \text{dcl}(\emptyset)$. Finally, define

$$\begin{aligned} \sqrt[n]{-} : P_n &\rightarrow \mathbb{M}^\times \\ \sqrt[n]{x} &= b_i(x/a_i)^{1/n} \text{ if } x \in a_i P_m. \end{aligned}$$

The map $\sqrt[n]{x}$ is defined by gluing together locally splendid functions on the cosets $a_1 P_m, a_2 P_m, \dots, a_k P_m$, so $\sqrt[n]{x}$ is locally splendid by Proposition 9.10(2). \square

Corollary 9.17. *Let $M \prec \mathbb{M}$ be a small model. Let \bar{a} be an n -tuple in \mathbb{M} . Let b be an element in $\text{dcl}(\bar{a}M)$. Then $b = f(\bar{a})$ for some M -definable locally splendid function $f : U \rightarrow \mathbb{M}$ with $\bar{a} \in U \subseteq \mathbb{M}^n$.*

Proof. Let A be the set of elements of the form $f(\bar{a})$, where f is M -definable and locally splendid. It suffices to show that $\text{dcl}(\bar{a}M) \subseteq A$. If g is M -definable and locally splendid and $\bar{b} \in A^n$, then $g(\bar{b}) \in A$ because a composition of locally splendid functions is locally splendid (Proposition 9.13). By Lemma 9.15, A is a subfield of \mathbb{M} . As constant functions and coordinate projections are locally splendid, A contains M and the tuple \bar{a} . By Lemma 9.16(1), A is a Henselian valued field. Lastly, Lemma 9.16(2) shows that if $a \in A$ and $a \in P_n = P_n(\mathbb{M})$, then $\sqrt[n]{a}$ exists in A .

Claim 9.18. *If $\gamma \in \Gamma_A$ and γ is a multiple of n in $\Gamma_{\mathbb{M}}$, then $\gamma/n \in \Gamma_A$.*

⁸Alternatively, $x^{1/n}$ is analytic at almost all points by the generic analyticity of definable functions in K [22, Theorem 1.1]. Since $x^{1/n}$ is a homomorphism, as soon as it is locally analytic at one point, it is analytic everywhere.

Proof. Take an element $b \in A$ with $v(b) = \gamma$. Because P_n has finite index in the multiplicative group, it has only finitely many cosets, and all of them are M -definable. In particular, bP_n is M -definable, so it contains some M -definable element c . Then $b/c \in P_n$. It follows that $v(b/c)$ is a multiple of n (in $\Gamma_{\mathbb{M}}$). As $v(b) = \gamma$ is a multiple of n , we also see that $v(c)$ is a multiple of n (in Γ_M or $\Gamma_{\mathbb{M}}$). Then

$$\frac{\gamma}{n} = \frac{v(b)}{n} = \frac{v(b/c) + v(c)}{n} = \frac{v(b/c)}{n} + \frac{v(c)}{n} = v\left(\sqrt[n]{b/c}\right) + \frac{v(c)}{n}.$$

Because A is a field containing M , we have $b/c \in A$. Because A is closed under n th roots (when they exist), we have $\sqrt[n]{b/c} \in A$. Then $v\left(\sqrt[n]{b/c}\right) \in \Gamma_A$. Finally, $v(c)/n \in \Gamma_M \subseteq \Gamma_A$. It follows that $\gamma/n \in \Gamma_A$, proving the Claim. \square_{Claim}

By the Claim, Γ_A is an elementary substructure of $\Gamma_{\mathbb{M}}$. By Ax-Kochen-Ershov principles, or the axiomatization of $\text{Th}(K)$ plus model completeness, we see that $M \preceq A \preceq \mathbb{M}$. Then $\text{dcl}(M\bar{a}) \subseteq A$. \square

Using this, we see that definable functions are generically locally splendid:

Proposition 9.19. *Let U be a non-empty definable open set in \mathbb{M}^n and $f : U \rightarrow \mathbb{M}^m$ be definable. Then there is a smaller definable open set $U_0 \subseteq U$ such that $f \upharpoonright U_0$ is locally splendid, and $\dim(U \setminus U_0) < \dim(U)$. Moreover, if M is a small model defining f and U , then we can take U_0 to be M -definable.*

Proof. Fix a small $M \preceq \mathbb{M}$ defining U and f , if none was given.

Claim 9.20. *If $\bar{a} \in U$, then one of the following holds:*

- $\bar{a} \in D$ for some M -definable set $D \subseteq \mathbb{M}^n$ with $\dim(D) < n$.
- $\bar{a} \in D$ for some M -definable open set $D \subseteq \mathbb{M}^n$ such that $f \upharpoonright D$ is splendid.

Proof. This follows from Corollary 9.17. If $\dim(\bar{a}/M) < n$, then the first case holds. Suppose that instead, $\dim(\bar{a}/M) = n$. Then Corollary 9.17 shows that $f(\bar{a}) = g(\bar{a})$ for some locally splendid M -definable function g . Let $D_0 = \{\bar{x} \in \text{dom}(f) \cap \text{dom}(g) : f(\bar{x}) = g(\bar{x})\}$. Then D_0 is M -definable and $\bar{a} \in D_0$, so $\dim(D_0) = n$. Let D be the interior of D_0 . Then $\dim(D_0 \setminus D) < n$, so $\bar{a} \notin D_0 \setminus D$, and instead $\bar{a} \in D$. The restriction $f \upharpoonright D$ equals the locally splendid function $g \upharpoonright D$. \square_{Claim}

By the Claim and saturation, we can cover U with finitely many M -definable sets

$$D_1 \cup D_2 \cup \cdots \cup D_k \cup U_1 \cup \cdots \cup U_\ell$$

such that $\dim(D_i) < n$, U_i is open, and $f \upharpoonright U_i$ is locally splendid. Let $U_0 = U_1 \cup \cdots \cup U_\ell$. Then

$$\dim(U \setminus U_0) \leq \dim \bigcup_i D_i < n = \dim(U),$$

and $f \upharpoonright U$ is locally splendid by Proposition 9.10(2). \square

9.3 Back to groups

Recall the operation

$$x \star_a y = xa^{-1}y$$

making G into a definable group with identity element a , definably isomorphic to the original group.

Lemma 9.21. *Let M be a small model. Let $G \subseteq \mathbb{M}^n$ be an n -dimensional M -definable set and \star be an M -definable group operation on G . Then there is an M -definable set U in the interior of G with $\dim(G \setminus U) < n$, such that for any $a \in U$, the group operation \star_a is locally splendid at (a, a) .*

Proof. Like the “direct” proof of Lemma 8.2, using Proposition 9.19 to get an M -definable set $\Delta \subseteq G \times G$ on which the group operation is locally splendid. \square

Recall the condition \mathfrak{C}_ω from Definition 7.1.

Lemma 9.22. *Let $G \subseteq \mathbb{M}^n$ be an n -dimensional definable set and let \star be a definable group operation on G with identity element $\bar{0}$. Let ϵ, δ be non-zero elements of \mathbb{M} , with $\epsilon\mathcal{O}^n$ and $\delta\mathcal{O}^n$ contained in G . Suppose that the group operation \star maps $\epsilon\mathcal{O}^n \times \epsilon\mathcal{O}^n$ into $\delta\mathcal{O}^n$, and the map*

$$\begin{aligned} f : \mathcal{O}^n \times \mathcal{O}^n &\rightarrow \mathcal{O}^n \\ f(\bar{x}, \bar{y}) &= \delta^{-1} \cdot (\epsilon\bar{x} \star \epsilon\bar{y}) \end{aligned}$$

is splendid. Let

$$\rho = p\epsilon^2/\delta.$$

Then $(\rho\mathcal{O}^n, \star)$ is a subgroup of G satisfying \mathfrak{C}_ω .

Proof. The set of counterexamples is ind-definable, because the family of splendid functions is ind-definable (by definition) and the \mathfrak{C}_ω condition is type-definable (Remark 7.3). The set of counterexamples is even ind-definable over the empty set, by automorphism invariance. If it is non-empty, it must contain a K -definable point. In other words, if there is a counterexample, then there is a K -definable counterexample.

Therefore, we may assume that $G, \star, \epsilon, \delta$ are K -definable. Then f is a K -definable splendid map, so $f(K) : \mathcal{O}_K^n \times \mathcal{O}_K^n \rightarrow \mathcal{O}_K^n$ is pre-splendid, given by a power series

$$f(\bar{x}, \bar{y}) = \sum_{I, J} \bar{c}_{I, J} \bar{x}^I \bar{y}^J$$

with $\bar{c}_{I, J} \in \mathcal{O}_K^m$. Then the group law itself is given by

$$\begin{aligned} \bar{x} \star \bar{y} &= \delta f(\epsilon^{-1}\bar{x}, \epsilon^{-1}\bar{y}) \\ &= \sum_{I, J} \bar{c}_{I, J} \delta \epsilon^{-|I|-|J|} \bar{x}^I \bar{y}^J, \end{aligned}$$

at least for $\bar{x}, \bar{y} \in \epsilon \mathcal{O}_K^n$. By the identity law for \star , the first few terms of the power series for \star must be $\bar{x} + \bar{y} + \dots$, and so the power series for f must have the form

$$f(\bar{x}, \bar{y}) = \frac{\epsilon}{\delta} \bar{x} + \frac{\epsilon}{\delta} \bar{y} + \dots$$

In particular, $\epsilon/\delta \in \mathcal{O}$. It follows that $\rho = p(\epsilon/\delta)\epsilon$ has higher valuation than ϵ , so

$$\rho \mathcal{O}^n \subseteq \epsilon \mathcal{O}^n \subseteq G.$$

If $\bar{x}, \bar{y} \in \mathcal{O}_K^n$, then

$$\begin{aligned} \rho^{-1}(\rho \bar{x} \star \rho \bar{y}) &= \sum_{I,J} \bar{c}_{I,J} \rho^{|I|+|J|-1} \delta \epsilon^{-|I|-|J|} \bar{x}^I \bar{y}^J \\ &= \bar{x} + \bar{y} + \frac{\delta}{\epsilon} \sum_{\substack{I,J \\ |I| \geq 1 \\ |J| \geq 1}} \bar{c}_{I,J} \left(\frac{\rho}{\epsilon}\right)^{|I|+|J|-1} \bar{x}^I \bar{y}^J. \end{aligned}$$

By Example 7.4, it remains to show that

$$\frac{\delta}{\epsilon} \left(\frac{\rho}{\epsilon}\right)^{|I|+|J|-1} \bar{c}_{I,J} \stackrel{?}{\in} p^{|I|+|J|-1} \mathcal{O}^n.$$

But this is clear, because

$$\frac{\delta}{\epsilon} \left(\frac{\rho}{\epsilon}\right)^{|I|+|J|-1} = \frac{\delta}{\epsilon} \left(p \frac{\epsilon}{\delta}\right)^{|I|+|J|-1} = p^{|I|+|J|-1} (\epsilon/\delta)^{|I|+|J|-2} \in p^{|I|+|J|-1} \mathcal{O}^n,$$

using the fact that $|I| + |J| \geq 2$ and $\epsilon/\delta \in \mathcal{O}$. \square

Theorem 9.23. *Let $M \preceq \mathbb{M}$ be a model (of $T = \text{Th}(K)$) and let G be an M -definable group. There is an M -definable open subgroup $H \subseteq G$ such that*

- H is isomorphic to a definable group (\mathcal{O}^n, \star) satisfying \mathfrak{C}_ω .
- H is fsg and compactly dominated by H/H^{00} , and H/H^{00} is isomorphic to an n -dimensional p -adic Lie group.

Proof. By Lemma 8.1 we may assume $G \subseteq \mathbb{M}^n$. By Lemma 9.21, there is an M -definable non-empty set U such that \star_a is locally splendid around (a, a) for any $a \in U$. Fix an $a \in U(M)$. Then (G, \star) is definably isomorphic to (G, \star_a) . Replacing (G, \star) with (G, \star_a) , we may assume that \star is locally splendid at the identity element 1_G . Translating G , we may assume the identity element is $\bar{0} \in \mathbb{M}^n$.

By definition of “locally splendid,” there are ϵ and δ such that \star maps $\epsilon \mathcal{O}^n \times \epsilon \mathcal{O}^n$ into $\delta \mathcal{O}^n$, and the map

$$\begin{aligned} f : \mathcal{O}^n \times \mathcal{O}^n &\rightarrow \mathcal{O}^n \\ f(\bar{x}, \bar{y}) &= \delta^{-1} \cdot (\epsilon \bar{x} \star \epsilon \bar{y}) \end{aligned}$$

is splendid. We can choose $\epsilon, \delta \in M$ by Lemma 9.11. Then Lemma 9.22 gives an (M -definable) subgroup $\rho \mathcal{O}^n \subseteq G$ satisfying \mathfrak{C}_ω . By Proposition 7.11, the subgroup $\rho \mathcal{O}^n$ has the desired properties. \square

Acknowledgments. The author was supported by the National Natural Science Foundation of China (Grant No. 12101131) and the Ministry of Education of China (Grant No. 22JJD110002).

References

- [1] Pablo Andújar Guerrero and Will Johnson. Around definable types in p -adically closed fields. arXiv:2208.05815v1 [math.LO], 2022.
- [2] Raf Cluckers. Classification of semi-algebraic p -adic sets up to semi-algebraic bijection. *J. Reine Angew. Math.*, 540:105–114, 2001.
- [3] Raf Cluckers. Presburger sets and P -minimal fields. *J. Symbolic Logic*, 68(1):153–162, Mar 2003.
- [4] Raf Cluckers. Analytic p -adic cell decomposition and integrals. *Trans. Amer. Math. Soc.*, 356(4):1489–1499, 2004.
- [5] Pablo Cubides-Kovacsics, Luck Darnière, and Eva Leenknegt. Topological cell decomposition and dimension theory in P -minimal fields. *Journal of Symbolic Logic*, 82(1):347–358, 2017.
- [6] Pablo Cubides Kovacsics and Françoise Delon. Definable completeness of P -minimal fields and applications. *J. Math. Logic*, 22(2):2250004, 2022.
- [7] Lou van den Dries, Deirdre Haskell, and Dugald Macpherson. One dimensional p -adic subanalytic sets. *J. London Math. Soc.*, 59(1):1–20, February 1999.
- [8] Jan Denef. p -adic semi-algebraic sets and cell decomposition. *J. Reine Angew. Math.*, 369:154–166, 1986.
- [9] J. D. Dixon, M. P. F. du Sautoy, A. Mann, and D. Segal. *Analytic Pro- p groups*, volume 61 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, 2nd edition, 1999.
- [10] Alfred Dolich, John Goodrick, and David Lippel. Dp-minimality: basic facts and examples. *Notre Dame Journal of Formal Logic*, 52(3):267–288, 2011.
- [11] Antongiulio Fornasiero. Definable compactness for topological structures. Draft, 2015.
- [12] Deirdre Haskell and Dugald Macpherson. A version of o -minimality for the p -adics. *J. Symbolic Logic*, 62(4):1075–1092, December 1997.
- [13] Ehud Hrushovski, Ya’acov Peterzil, and Anand Pillay. Groups, measures, and the NIP. *J. Amer. Math. Soc.*, 21(2):563–596, April 2008.

- [14] Ehud Hrushovski and Anand Pillay. On NIP and invariant measures. *J. Eur. Math. Soc.*, 13(4):1005–1061, 2011.
- [15] Will Johnson. Interpretable sets in dense o-minimal structures. *J. Symbolic Logic*, 83:1477–1500, 2018.
- [16] Will Johnson. Dp-finite fields $I(A)$: the infinitesimals. *Annals of Pure and Applied Logic*, 172(6):102947, 2021.
- [17] Will Johnson and Ningyuan Yao. On non-compact p -adic definable groups. *J. Symbolic Logic*, 87(1):188–213, 2022.
- [18] Tristan Kuijpers and Eva Leenknegt. Differentiation in P -minimal structures and a p -adic Local Monotonicity Theorem. *J. Symbolic Logic*, 79(4):1133–1147, 2014.
- [19] Michel Lazard. Groupes analytiques p -adiques. *Publications Mathématiques de l’IHÉS*, 26, 1965.
- [20] A. Onshuus and A. Pillay. Definable groups and compact p -adic Lie groups. *Journal of the London Mathematical Society*, 78(1):233–247, 2008.
- [21] Peter Schneider. *p -Adic Lie groups*, volume 344 of *Grundlehren der mathematischen Wissenschaften*. Springer, 2011.
- [22] Philip Scowcroft and Lou van den Dries. On the structure of semialgebraic sets over p -adic fields. *J. Symbolic Logic*, 53(4):1138–1164, December 1988.
- [23] Pierre Simon. Dp-minimality: invariant types and dp-rank. *J. Symbolic Logic*, 79(4):1025–1045, December 2014.
- [24] Pierre Simon. *A guide to NIP theories*. Lecture Notes in Logic. Cambridge University Press, July 2015.