

# ON SETS OF LENGTHS IN MONOIDS OF PLUS-MINUS WEIGHTED ZERO-SUM SEQUENCES OVER ABELIAN GROUPS

ALFRED GEROLDINGER AND FLORIAN KAINRATH

**ABSTRACT.** Let  $G$  be an additive abelian group. A sequence  $S = g_1 \cdots g_\ell$  of terms from  $G$  is a plus-minus weighted zero-sum sequence if there are  $\varepsilon_1, \dots, \varepsilon_\ell \in \{-1, 1\}$  such that  $\varepsilon_1 g_1 + \dots + \varepsilon_\ell g_\ell = 0$ . We study sets of lengths in the monoid  $\mathcal{B}_\pm(G)$  of plus-minus weighted zero-sum sequences over  $G$ . If  $G$  is finite, then sets of lengths are highly structured. If  $G$  is infinite, then every finite, nonempty subset of  $\mathbb{N}_{\geq 2}$  is the set of lengths of some sequence  $S \in \mathcal{B}_\pm(G)$ .

## 1. INTRODUCTION

Let  $G$  be an additive abelian group. By a sequence over  $G$ , we mean a finite sequence of terms from  $G$ , where the order of terms is disregarded and repetition is allowed. We consider sequences as elements of the (multiplicatively written) free abelian monoid  $\mathcal{F}(G)$  with basis  $G$  (the multiplication of sequences in  $\mathcal{F}(G)$  means the concatenation of sequences in combinatorial language). A sequence  $S = g_1 \cdots g_\ell$ , with terms  $g_1, \dots, g_\ell$  from  $G$ , is a zero-sum sequence if  $g_1 + \dots + g_\ell = 0$  and it is a plus-minus weighted zero-sum sequence if  $\varepsilon_1 g_1 + \dots + \varepsilon_\ell g_\ell = 0$  for some  $\varepsilon_1, \dots, \varepsilon_\ell \in \{-1, 1\}$ .

Weighted zero-sum sequences (from fully weighted sequences to plus-minus weighted sequences) have been studied in additive combinatorics since the last two decades. Among others, many of the classical zero-sum invariants (such as the Davenport constant, Gao's constant, and others) gave rise to weighted analogs (for a sample see [21, Chapter 16], [1, 24, 27, 23, 22, 26]).

The monoid  $\mathcal{B}(G)$  of all zero-sum sequences over  $G$  is a Krull monoid and it plays a universal role in the arithmetic study of general Krull monoids. Pushed forward by this connection, algebraic and arithmetic properties of  $\mathcal{B}(G)$  are central topics in the factorization theory of rings and monoids. The study of the monoid  $\mathcal{B}_\pm(G)$  of plus-minus weighted zero-sum sequences over  $G$ , from an algebraic viewpoint, was initiated only a couple of years ago by Schmid and his coauthors. Algebraic topics include questions when monoids of weighted zero-sum sequences are Krull, Mori, or finitely generated. Arithmetic topics deal with questions on (various types of) Davenport constants and on invariants controlling the structure of sets of lengths. The isomorphism problem asks whether, for given abelian groups  $G_1$  and  $G_2$ , the monoids  $\mathcal{B}_\pm(G_1)$  and  $\mathcal{B}_\pm(G_2)$  are isomorphic if and only if the groups  $G_1$  and  $G_2$  are isomorphic (for all these topics see [4, 15, 7, 29]).

In the present paper, we study sets of lengths in monoids of plus-minus weighted zero-sum sequences. To fix notation, let  $H$  be a commutative and cancellative monoid. For an element  $a \in H$ , we denote by  $\mathsf{L}_H(a) \subset \mathbb{N}_0$  the set of lengths of  $a$ , and by  $\mathcal{L}(H) = \{\mathsf{L}(a) : a \in H\}$  the system of sets of lengths of  $H$  (for details, see Section 2).

We formulate our main result (more information on AAMPs and on the set of minimal distances  $\Delta^*(\mathcal{B}_\pm(G))$  will be given in Section 5).

---

2020 *Mathematics Subject Classification.* 20M13, 11B30, 13A05.

*Key words and phrases.* weighted zero-sum sequences, sets of lengths.

This work was supported by the Austrian Science Fund FWF (Project P36852-N).

**Theorem 1.1.** *Let  $G$  be an abelian group.*

1. *If  $G$  is finite, then there is a bound  $M(G) \in \mathbb{N}_0$  such that, for every plus-minus weighted sequence  $S \in \mathcal{B}_\pm(G)$ , its set of lengths  $\mathsf{L}_{\mathcal{B}_\pm(G)}(S)$  is an AAMP with difference  $d \in \Delta^*(\mathcal{B}_\pm(G))$  and bound  $M(G)$ .*
2. *If  $G$  is infinite, then, for every finite, nonempty subset  $L \subset \mathbb{N}_{\geq 2}$  and every map  $f: L \rightarrow \mathbb{N}$ , there is an  $S \in \mathcal{B}(G)$  with the following properties.*
  - (a)  *$S$  is squarefree in  $\mathcal{F}(G)$ .*
  - (b)  *$\mathsf{L}_{\mathcal{B}_\pm(G)}(S) = \mathsf{L}_{\mathcal{B}(G)}(S) = L$ .*
  - (c)  *$|\mathsf{Z}_{\mathcal{B}_\pm(G),k}(S)| \geq |\mathsf{Z}_{\mathcal{B}(G),k}(S)| \geq f(k)$  for all  $k \in L$ . Moreover, both inequalities are equalities for  $k > \min L$ .*

Before we discuss the statements in detail, consider a transfer homomorphism  $\theta: H \rightarrow B$  between monoids  $H$  and  $B$  (definitions are given in Section 2). The existence of such a homomorphism implies that  $\mathcal{L}(H) = \mathcal{L}(B)$ . Krull domains, Krull monoids, and transfer Krull monoids (see the examples discussed in the survey [19]) allow transfer homomorphisms to monoids of zero-sum sequences over subsets of abelian groups. Thus, sets of lengths in such monoids and domains can be studied in monoids of zero-sum sequences. Similarly, monoids of plus-minus weighted zero-sum sequences occur as target monoids of transfer homomorphisms which start, for example, from norm monoids of orders in Galois number fields. On the other hand, (apart from a trivial exceptional case) there is no transfer homomorphism from a monoid of plus-minus weighted zero-sum sequences to any Krull monoid ([7, Corollary 3.5]). This demonstrates that we cannot get the above result via transfer homomorphisms from an associated result on Krull monoids.

The first statement of Theorem 1.1 is a simple consequence of known results and it is formulated to highlight the difference between finite and infinite groups (see Section 5 for more on  $\Delta^*(\mathcal{B}_\pm(G))$ ). The Characterization Problem (for monoids of plus-minus weighted zero-sum sequences) asks which finite abelian groups  $G_1$  have the property that the equality of systems of sets of lengths  $\mathcal{L}(\mathcal{B}_\pm(G_1)) = \mathcal{L}(\mathcal{B}_\pm(G_2))$  implies that the groups  $G_1$  and  $G_2$  are isomorphic, for any finite abelian group  $G_2$  (see [7, 29]). All work in this direction is based on the structural description given in Theorem 1.1.1.

The second statement of Theorem 1.1 implies, in particular, that every finite, nonempty subset of  $\mathbb{N}_{\geq 2}$  occurs as a set of lengths. Thus, the monoid  $H = \mathcal{B}_\pm(G)$  satisfies the property

$$(1.1) \quad \mathcal{L}(H) = \{\{0\}, \{1\}\} \cup \{L \subset \mathbb{N}_{\geq 2} : L \text{ is finite and nonempty}\},$$

and hence  $\mathcal{L}(\mathcal{B}_\pm(G_1)) = \mathcal{L}(\mathcal{B}_\pm(G_2))$  for any two infinite abelian groups  $G_1$  and  $G_2$ .

Property (1.1) was first proved by Kainrath for Krull monoids with infinite abelian class group and prime divisors in all classes ([25]). Since then Property (1.1) was shown to hold true for various classes of integer-valued polynomials ([11, 12, 10, 8]), for some classes of primary monoids ([20, Theorem 3.6]), and others ([9]). Furthermore, there are monoid algebras which satisfy the ascending chain condition on principal ideals (whence they are atomic) and which have the property that every (not necessarily finite) nonempty subset of  $\mathbb{N}_{\geq 2}$  occurs as a set of lengths ([13]).

After the first result in [25], several distinct constructions realizing sets of lengths in Krull monoids were given (for a realization result in numerical monoids see [16]).

- (i) A realization theorem for a single finite, nonempty subset  $L \subset \mathbb{N}_{\geq 2}$  in an abstract finitely generated Krull monoid  $H_L$  ([14, Proposition 4.8.3]; note if  $\mathcal{L}$  is a family of finite subsets, then all sets of  $\mathcal{L}$  are sets of lengths in the coproduct  $\coprod_{L \in \mathcal{L}} H_L$ , and this coproduct is Krull again).
- (ii) A realization theorem for a single finite, nonempty subset  $L \subset \mathbb{N}_{\geq 2}$  in a monoid of zero-sum sequences over a finite abelian group ([32]).
- (iii) A realization theorem for all finite, nonempty subsets  $L \subset \mathbb{N}_{\geq 2}$  in a monoid of zero-sum sequences over an abelian group containing an element of infinite order ([2, Theorem 3]).

Our proof of Theorem 1.1 follows the ideas given in [25]. We recapitulate the whole construction, which allows us to prove not only Property (1.1) for  $\mathcal{B}_\pm(G)$ , but the stronger Properties (a), (b), and (c) of Theorem 1.1. Moreover, this increases the readability of the present paper.

We obtain the following corollary, which should be seen against the background of the above mentioned Characterization Problem (for related results see [33, Theorem 4], [17, Theorem 3.7]).

**Corollary 1.2.** *Let  $L \subset \mathbb{N}_{\geq 2}$  be a finite, nonempty subset. Then there are only finitely many pairwise non-isomorphic finite abelian groups  $G$  such that  $L \notin \mathcal{L}(\mathcal{B}_\pm(G))$ .*

In Section 2, we put together the background on factorizations and on monoids of (weighted) zero-sum sequences. A crucial step in the proof of Theorem 1.1 will be done in Section 3 and a further reduction step will be handled in Section 4. Finally, we complete the proofs of Theorem 1.1 and of Corollary 1.2 in Section 5. Along our way, we oftentimes (as in Theorem 1.1) consider sequences  $S \in \mathcal{B}(G)$  and find out that the set of lengths or even all factorizations of  $S$  in  $\mathcal{B}(G)$  are the same as in  $\mathcal{B}_\pm(G)$ . Needless to say, that this is far from being true in general (for a striking difference see Proposition 5.2) but holds true only in well-constructed exceptional cases.

## 2. PREREQUISITES

By a *monoid*, we mean a commutative, cancellative semigroup with identity element, and we use multiplicative notation throughout. For integers  $a, b \in \mathbb{Z}$ , let  $[a, b] = \{x \in \mathbb{Z} : a \leq x \leq b\}$  be the discrete interval between  $a$  and  $b$ . For a set  $P$ , we denote by  $\mathcal{F}(P)$  the free abelian monoid with basis  $P$ . An element  $a \in \mathcal{F}(P)$  has a unique representation in the form

$$a = \prod_{p \in P} p^{\mathbf{v}_p(a)},$$

where  $\mathbf{v}_p : \mathcal{F}(P) \rightarrow \mathbb{N}_0$  is the  $p$ -adic valuation of  $a$ . Then  $|a| = \sum_{p \in P} \mathbf{v}_p(a) \in \mathbb{N}_0$  denotes the *length* of  $a$  and  $\text{supp}(a) = \{p \in P : \mathbf{v}_p(a) > 0\} \subset P$  the *support* of  $a$ .

Let  $H$  be a monoid. We denote by  $H^\times$  its group of invertible element and by  $H_{\text{red}} = \{aH^\times : a \in H\}$  the associated reduced monoid of  $H$ . An element  $a \in H$  is said to be *irreducible* (or an *atom*) if  $a \notin H^\times$  and  $a = bc$ , with  $b, c \in H$ , implies that  $b \in H^\times$  or  $c \in H^\times$ . We denote by  $\mathcal{A}(H)$  the set of atoms of  $H$ , by  $Z(H) = \mathcal{F}(\mathcal{A}(H_{\text{red}}))$  the *factorization monoid* of  $H$ , and by  $\pi : Z(H) \rightarrow H_{\text{red}}$  the *factorization homomorphism*, defined by  $\pi(u) = u$  for all  $u \in \mathcal{A}(H_{\text{red}})$ . For an element  $a \in H$  and  $k \in \mathbb{N}_0$ , let

- (i)  $Z_H(a) = Z(a) = \pi^{-1}(a) \subset Z(H)$  be the *set of factorizations* of  $a$ ,
- (ii)  $Z_{H,k}(a) = Z_k(a) = \{z \in Z(a) : |z| = k\}$  be the set of factorizations of  $a$ , which have length  $k$ , and
- (iii)  $L_H(a) = L(a) = \{|z| : z \in Z(a)\} \subset \mathbb{N}_0$  be the *set of lengths* of  $a$ .

Thus, by definition,  $L(a) = \{1\}$  if and only if  $a$  is an atom and  $L(a) = \{0\}$  if and only if  $a \in H^\times$ . We say that  $H$  is

- (i) *atomic* if  $Z(a) \neq \emptyset$  for all  $a \in H$  (equivalently, every  $a \in H \setminus H^\times$  can be written as a finite product of atoms),
- (ii) an *FF-monoid* (finite-factorization monoid) if  $Z(a)$  is finite and nonempty for all  $a \in H$ , and
- (iii) a *BF-monoid* (bounded-factorization monoid) if  $L(a)$  is finite and nonempty for all  $a \in H$ .

Let  $H$  be a BF-monoid. We denote by

$$\mathcal{L}(H) = \{L(a) : a \in H\}$$

the *system of sets of lengths* of  $H$ . A submonoid  $S \subset H$  is called *divisor-closed* if  $a \in H$  and  $b \in S$  with  $a|b$  implies that  $a \in S$ . If  $a \in S$ , then

$$Z_S(a) = Z_H(a) \quad \text{and} \quad L_S(a) = L_H(a).$$

Let  $G$  be an additive abelian group and let  $G_0 \subset G$  be a subset. For every  $m \in \mathbb{N}$ , we set  $mG = \{mg : g \in G\}$ . The elements of  $\mathcal{F}(G_0)$  are called *sequences* over  $G_0$ . Let  $S \in \mathcal{F}(G_0)$ . Then we write

$$S = g_1 \cdot \dots \cdot g_\ell = \prod_{g \in G_0} g^{\mathbf{v}_g(S)},$$

and in this notation we tacitly assume that  $\ell \in \mathbb{N}_0$  and  $g_1, \dots, g_\ell \in G_0$ . If  $\varphi: G \rightarrow G'$  is a group homomorphism, then  $\varphi(S) = \varphi(g_1) \cdot \dots \cdot \varphi(g_\ell)$ ,  $-S = (-g_1) \cdot \dots \cdot (-g_\ell)$ ,

$$|S| = \ell = \sum_{g \in G_0} \mathbf{v}_g(S) \in \mathbb{N}_0$$

is the *length* of  $S$ , and

$$\sigma(S) = g_1 + \dots + g_\ell \in G$$

is the *sum* of  $S$ . We say that  $S$  is

- (i) *squarefree* (in  $\mathcal{F}(G_0)$ ) if  $g^2 \nmid S$  for all  $g \in G_0$ ,
- (ii) a *zero-sum sequence* if  $\sigma(S) = 0$ , and
- (iii) a *plus-minus weighted zero-sum sequence* if there are  $\varepsilon_1, \dots, \varepsilon_\ell \in \{-1, 1\}$  such that  $\varepsilon_1 g_1 + \dots + \varepsilon_\ell g_\ell = 0$ .

We denote by  $\mathcal{B}(G_0)$  the set of zero-sum sequences over  $G_0$  and by  $\mathcal{B}_\pm(G_0)$  the set of plus-minus weighted zero-sum sequences over  $G_0$ . These are submonoids of  $\mathcal{F}(G_0)$  with the obvious inclusions

$$\mathcal{B}(G_0) \subset \mathcal{B}_\pm(G_0) \subset \mathcal{F}(G_0).$$

Whenever we speak of a squarefree sequence, we mean that the sequence is squarefree in  $\mathcal{F}(G)$ . If  $G_1 \subset G_0$  is a subset, then  $\mathcal{B}(G_1) \subset \mathcal{B}(G_0)$  and  $\mathcal{B}_\pm(G_1) \subset \mathcal{B}_\pm(G_0)$  are divisor-closed submonoids. If every non-zero element  $g \in G_0$  has order two, then  $\mathcal{B}(G_0) = \mathcal{B}_\pm(G_0)$ . Both monoids,  $\mathcal{B}(G_0)$  and  $\mathcal{B}_\pm(G_0)$ , are FF-monoids and hence BF-monoids. Since the inclusion  $\mathcal{B}(G_0) \hookrightarrow \mathcal{F}(G_0)$  is a divisor homomorphism,  $\mathcal{B}(G_0)$  is a Krull monoid.

A monoid homomorphism  $\theta: H \rightarrow B$  is said to be a *transfer homomorphism* if the following two conditions are satisfied.

**(T1)**  $B = \theta(H)B^\times$  and  $\theta^{-1}(B^\times) = H^\times$ .

**(T2)** If  $u \in H$ ,  $b, c \in B$  and  $\theta(u) = bc$ , then there exist  $v, w \in H$  such that  $u = vw$ ,  $\theta(v) \in bB^\times$ , and  $\theta(w) \in cB^\times$ .

A transfer homomorphism allows to pull back various arithmetic properties from  $B$  to  $H$ . In particular, we have  $\mathcal{L}(H) = \mathcal{L}(B)$ . The classic example of a transfer homomorphism stems from the theory of Krull monoids. They allow a transfer homomorphism to monoids of zero-sum sequences  $\mathcal{B}(G_0)$ , where  $G_0$  is a subset of the class group of the Krull monoid ([14]). A monoid is said to be *transfer Krull* if it allows a transfer homomorphism to a Krull monoid (equivalently, to a monoid of zero-sums sequences) (see [19, 3] for examples of transfer Krull monoids that are not Krull). If  $G$  is an abelian group, then for the monoid of plus-minus weighted zero-sum sequences, we have the following characterizations ([7, Corollary 3.5]). There are equivalent.

- (a)  $\mathcal{B}_\pm(G)$  is Krull.
- (b)  $\mathcal{B}_\pm(G)$  is transfer Krull.
- (c)  $G$  is an elementary 2-group.

On the other hand, in [15, Theorems 3.2 and 3.5], it was proved that norm monoids of Galois number fields and monoids of positive integers, which can be represented by certain binary quadratic forms, allow transfer homomorphisms to monoids of plus-minus weighted zero-sum sequences. More on this direction can be found in [24, 4, 6].

We end this section with a simple lemma, dealing with the relationship of atoms in  $\mathcal{B}(G)$  and in  $\mathcal{B}_\pm(G)$ . Clearly, we have

$$(2.1) \quad \mathcal{A}(\mathcal{B}_\pm(G)) \cap \mathcal{B}(G) \subset \mathcal{A}(\mathcal{B}(G)).$$

The next lemma guarantees the reverse implication for groups with trivial 2-torsion (for finite groups see [4, Theorem 6.1]).

**Lemma 2.1.** *Let  $G$  be an abelian group and suppose that there is no  $g \in G$  with  $\text{ord}(g) = 2$ . Then  $\mathcal{A}(\mathcal{B}(G)) = \mathcal{A}(\mathcal{B}_\pm(G)) \cap \mathcal{B}(G)$ . In particular, we have  $\mathbb{L}_{\mathcal{B}(G)}(B) \subset \mathbb{L}_{\mathcal{B}_\pm(G)}(B)$  for all  $B \in \mathcal{B}(G)$ .*

*Proof.* Clearly, it suffice to show that statement on the atoms. Then the 'in particular' on sets of lengths follows immediately. Let  $A \in \mathcal{A}(\mathcal{B}(G))$  and suppose that there are  $B_1, B_2 \in \mathcal{B}_\pm(G)$  with  $A = B_1 B_2$  and with  $B_1 \neq 1$ . We assert that  $B_1 = A$ .

Since  $B_1, B_2 \in \mathcal{B}_\pm(G)$ , there are  $B_1^+, B_1^-, B_2^+, B_2^- \in \mathcal{F}(G)$  such that

$$B_1 = B_1^+ B_1^-, B_2 = B_2^+ B_2^-, \sigma(B_1^+) = \sigma(B_1^-), \text{ and } \sigma(B_2^+) = \sigma(B_2^-).$$

By symmetry, we may suppose without restriction that  $B_1^+ \neq 1$ . Then it follows that

$$0 = \sigma(A) = \sigma(B_1^+) + \sigma(B_1^-) + \sigma(B_2^+) + \sigma(B_2^-) = 2(\sigma(B_1^+) + \sigma(B_2^+)).$$

Since, by assumption,  $G$  has no elements of order two, we infer that  $\sigma(B_1^+) + \sigma(B_2^+) = 0$ , whence  $1 \neq B_1^+ B_2^+ \in \mathcal{B}(G)$ . Since  $A \in \mathcal{A}(\mathcal{B}(G))$ , it follows that  $A = B_1^+ B_2^+$ , whence  $B_1^- = B_2^- = 1$  and so  $B_1 = A$ .

This implies that  $\mathcal{A}(\mathcal{B}(G)) \subset \mathcal{A}(\mathcal{B}_\pm(G))$ , and hence the asserted equality holds by Relation (2.1).  $\square$

### 3. ON THE CONSTRUCTION OF A GROUP HAVING A GIVEN $L$ WITH $\min L = 2$ AS A SET OF LENGTHS

The goal in this section is to prove the following proposition.

**Proposition 3.1.** *Let  $R$  be a commutative ring, let  $L \subset \mathbb{N}_{\geq 2}$  be a finite subset with  $2 \in L$ , and let  $f: L \rightarrow \mathbb{N}$  be a map with  $\sum_{k \in L} f(k) \geq 3$ . Then there exist a finitely generated free  $R$ -module  $G$  and a sequence  $B \in \mathcal{B}(G)$  with the following properties.*

- (a)  $B$  is squarefree in  $\mathcal{F}(G)$ .
- (b)  $\mathbb{L}_{\mathcal{B}_\pm(G)}(B) = \mathbb{L}_{\mathcal{B}(G)}(B) = L$ .
- (c)  $|\mathbb{Z}_{\mathcal{B}_\pm(G),k}(B)| \geq |\mathbb{Z}_{\mathcal{B}(G),k}(B)| \geq f(k)$  for all  $k \in L$ . Moreover, both inequalities are equalities for  $k > \min L$ .

We proceed in three subsections. First, we put together the setting in which  $G$  and  $B$  are defined. Then we study the atoms in  $\mathcal{B}(G)$  and in  $\mathcal{B}_\pm(G)$ . In the third subsection, we give the proof of Proposition 3.1

#### 3.1. On the construction of the $R$ -module $G$ and the wanted sequence $B \in \mathcal{B}(G)$ .

As said in the introduction, the  $R$ -module  $G$  and the sequence  $B$  are the same as in [25]. We carefully provide all definitions and gather all the required properties. The proofs of these properties can be found in [25] and in [14, Chapter 7.4].

Let  $R$  be a commutative ring, let  $L \subset \mathbb{N}_{\geq 2}$  be a finite subset with  $2 \in L$ , and let  $f: L \rightarrow \mathbb{N}$  be a map with  $s := \sum_{k \in L} f(k) \geq 3$ . We consider a tuple of finite sets  $(X_1, \dots, X_s)$  such that

$$(3.1) \quad L = \{|X_1|, \dots, |X_s|\} \quad \text{and} \quad f(k) = |\{i \in [1, s] : |X_i| = k\}| \quad \text{for every } k \in L.$$

We set

$$X = \prod_{j=1}^s X_j \quad \text{and} \quad X_J = \prod_{j \in J} X_j \quad \text{for } \emptyset \neq J \subset [1, s].$$

For  $\emptyset \neq J \subset [1, s]$ , let  $p_J: X \rightarrow X_J$  be the canonical projection and, for  $j \in [1, s]$ , let  $p_j = p_{\{j\}}: X \rightarrow X_j$ . For  $z = (z_1, \dots, z_s) \in X$ ,  $i \in [1, s]$ , and a nonempty subset  $J \subset [1, s]$ , we define

$$X_i^{(z)} = X_i \setminus \{z_i\} \quad \text{and} \quad X_J^{(z)} = \prod_{j \in J} X_j^{(z)}.$$

Furthermore, for two subsets  $M, N \subset X$ , let

$$\Delta(M, N) = (M \setminus N) \cup (N \setminus M)$$

denote the symmetric difference of  $M$  and  $N$ . For two elements  $x = (x_1, \dots, x_s), y = (y_1, \dots, y_s) \in X$ , we set  $\Delta(x, y) = \{i \in [1, s] : x_i \neq y_i\}$ .

Now let  $R^X$  be the  $R$ -algebra of all maps  $X \rightarrow R$ , with pointwise addition, pointwise multiplication, and with scalar multiplication by elements from  $R$ . For a subset  $M \subset X$ , let  $\chi_M \in R^X$  be the characteristic function of  $M$ , defined by

$$\chi_M(x) = \begin{cases} 1, & \text{if } x \in M, \\ 0, & \text{if } x \notin M. \end{cases}$$

For an element  $x \in X$ , we set  $\chi_x = \chi_{\{x\}} \in R^X$ , and  $\mathbf{1} = \chi_X$  is the constant function with value  $1 \in R$ .

Next, we define the wanted finitely generated free  $R$ -module. We set

$$(3.2) \quad V = \langle \{\chi_{p_i^{-1}(y)} : i \in [1, s], y \in X_i\} \rangle_R, \quad G = R^X/V,$$

and, for every  $z \in X$ ,

$$W_z = \langle \{\chi_{p_J^{-1}(y)} : J \subset [1, s], |J| \geq 2, y \in X_J^{(z)}\} \rangle_R.$$

We have that  $V = \langle \{\mathbf{1}\} \cup \{\chi_{p_i^{-1}(y)} : i \in [1, s], y \in X_i^{(z)}\} \rangle_R$ ,  $R^X = V \oplus W_z$ , and so  $G$  is a finitely generated free  $R$ -module. We denote by  $P_z : R^X = V \oplus W_z \rightarrow V$  the associated projection. If  $M \subset X \setminus \{z\}$ , then

$$(3.3) \quad P_z(\chi_z) = \mathbf{1} - \sum_{i=1}^s \sum_{y \in X_i^{(z)}} \chi_{p_i^{-1}(y)}, \quad P_z(\chi_x) = \begin{cases} \chi_{p_i^{-1}(x_i)}, & \text{if } \Delta(z, x) = \{i\}, \\ \mathbf{0}, & \text{if } |\Delta(z, x)| \geq 2, \end{cases}$$

and

$$(3.4) \quad P_z(\chi_M) = \sum_{i=1}^s \chi_{p_i^{-1}(Y_i)},$$

where, for each  $i \in [1, s]$ ,  $Y_i$  is a subset of  $X_i^{(z)}$ .

For a subset  $M \subset X$ , we set

$$(3.5) \quad B_M = \prod_{x \in M} (\chi_x + V) \in \mathcal{F}(G) \quad \text{and} \quad B := B_X$$

Since  $\chi_x - \chi_z \notin V$  for any two distinct elements of  $X$ , all sequences  $B_M$  are squarefree. If  $M, M', M''$  are subsets of  $X$ , then  $B_M = B_{M'} B_{M''}$  if and only if  $M = M' \uplus M''$ . Since

$$\sigma(B_M) = \sum_{x \in M} (\chi_x + V) = \chi_M + V,$$

we have  $B_M \in \mathcal{B}(G)$  if and only if  $\chi_M \in V$ . In particular,  $\mathbf{1} = \chi_X \in V$  implies  $B \in \mathcal{B}(G) \subset \mathcal{B}_\pm(G)$ . Moreover, for every divisor  $T$  of  $B$  in  $\mathcal{F}(G)$  there is a unique subset  $M \subset X$  with  $T = B_M$ .

### 3.2. On the atoms of $\mathcal{B}(G)$ and $\mathcal{B}_\pm(G)$ .

#### Lemma 3.2.

1. If  $i \in [1, s]$ ,  $Y_i \subset X_i$  and  $M = p_i^{-1}(Y_i)$ , then  $B_M \in \mathcal{B}(G)$ , and  $B_M \in \mathcal{A}(\mathcal{B}(G))$  if and only if  $|Y_i| = 1$ .
2. If  $\text{char}(R) \neq 2$ ,  $M \subset X$ , and  $B_M \in \mathcal{B}(G)$ , then  $M = p_i^{-1}(Y_i)$  for some  $i \in [1, s]$  and some  $Y_i \subset X_i$ .
3. Let  $\text{char}(R) = 2$ ,  $M \subsetneq X$  and  $B_M \in \mathcal{B}(G)$ . Then, for every  $i \in [1, s]$ , there is a set  $Y_i \subset X_i$  such that

$$M = \{(x_1, \dots, x_s) \in X : |\{i \in [1, s] : x_i \in Y_i\}| \text{ is odd}\}.$$

If  $B_M \notin \mathcal{A}(\mathcal{B}(G))$ , then  $M = p_i^{-1}(Y_i)$  for some  $i \in [1, s]$  and some subset  $Y_i \subset X_i$ . Moreover, if  $M' \subset X$  is such that  $B_{M'} \in \mathcal{B}(G)$  and  $B_{M'} \mid B_M$ , then  $M' = p_i^{-1}(Y'_i)$  for some subset  $Y'_i \subset Y_i$ .

*Proof.* 1. If  $i \in [1, s]$ ,  $Y_i \subset X_i$  and  $M = p_i^{-1}(Y_i)$ , then

$$\chi_M = \sum_{y \in Y_i} \chi_{p_i^{-1}(y)} \in V$$

and thus  $B_M \in \mathcal{B}(G)$ . Clearly,  $B_\emptyset = 1 \in \mathcal{B}(G)$ . If  $i, j \in [1, s]$ ,  $\emptyset \neq Y_i \subset X_i$  and  $\emptyset \neq Y'_j \subset X_j$ , then  $p_i^{-1}(Y_i) \subset p_j^{-1}(Y'_j)$  if and only if  $i = j$  and  $Y_i \subset Y'_j$ . Hence  $B_M \in \mathcal{A}(\mathcal{B}(G))$  if and only if  $|Y_i| = 1$ .

2. Clearly, we have  $X = p_1^{-1}(X_1)$ . Now suppose that  $M \subsetneq X$  and  $\chi_M \in V$ . If  $z \in X \setminus M$ , then by (3.4),

$$\chi_M = P_z(\chi_M) = \sum_{i=1}^s \chi_{p_i^{-1}(Y_i)},$$

where  $Y_i \subset X_i$  for each  $i \in [1, s]$ . Since

$$\chi_M = \chi_M^2 = \sum_{i=1}^s \chi_{p_i^{-1}(Y_i)} + 2 \sum_{\substack{i,j=1 \\ i < j}}^s \chi_{p_i^{-1}(Y_i)} \chi_{p_j^{-1}(Y_j)},$$

it follows that

$$0 = 2 \sum_{\substack{i,j=1 \\ i < j}}^s \chi_{p_i^{-1}(Y_i)} \chi_{p_j^{-1}(Y_j)} = 2 \sum_{\substack{i,j=1 \\ i < j}}^s \sum_{y \in Y_i \times Y_j} \chi_{p_{\{i,j\}}^{-1}}(y).$$

By (3.3), this can happen only if  $Y_i \times Y_j = \emptyset$  for all distinct  $i, j \in [1, s]$ . Hence there is at most one  $i \in [1, s]$  with  $Y_i \neq \emptyset$ .

3. Let  $z \in X \setminus M$ . If  $B_M \in \mathcal{B}(G)$ , then (3.4) implies

$$\chi_M = P_z(\chi_M) = \sum_{i=1}^s \chi_{p_i^{-1}(Y_i)},$$

where  $Y_i \subset X_i^{(z)}$  for all  $i \in [1, s]$ . For  $x = (x_1, \dots, x_s) \in X$  we have

$$\chi_M(x) = \sum_{i=1}^s \chi_{p_i^{-1}(Y_i)}(x) = |\{i \in [1, s] : x_i \in Y_i\}| 1_R,$$

and therefore  $M = \{(x_1, \dots, x_s) \in X : |\{i \in [1, s] : x_i \in Y_i\}| \text{ is odd}\}$ .

If  $B_M \notin \mathcal{A}(\mathcal{B}(G))$ , then there is a set  $\emptyset \neq M' \subsetneq M$  such that  $B_{M'} \in \mathcal{B}(G)$ . For every  $i \in [1, s]$ , let  $Y'_i \subset X_i^{(z)}$  be such that

$$M' = \{(x_1, \dots, x_s) \in X : |\{i \in [1, s] : x_i \in Y'_i\}| \text{ is odd}\}.$$

If  $j \in [1, s]$  and  $y \in Y'_j$ , then  $(z_1, \dots, z_{j-1}, y, z_{j+1}, \dots, z_s) \in M' \subset M$ , and thus  $y \in Y_j$ . Hence we obtain  $Y'_j \subset Y_j$  for all  $j \in [1, s]$ , and  $M' \subsetneq M$  implies  $Y'_i \subsetneq Y_i$  for some  $i \in [1, s]$ , say  $Y'_1 \subsetneq Y_1$ . We assert that  $Y_j = Y'_j = \emptyset$  for all  $j \in [2, s]$ . It suffices to do the proof for  $j = 2$ . Assume the contrary. If  $Y'_2 \neq \emptyset$ , let  $y_1 \in Y_1 \setminus Y'_1$  and  $y_2 \in Y'_2$ . Then  $(y_1, y_2, z_3, \dots, z_s) \in M' \setminus M$ , a contradiction. Hence  $Y'_2 = \emptyset$ . If  $Y_2 \neq \emptyset$ , let  $y_1 \in Y'_1$  and  $y_2 \in Y_2$ . Then  $(y_1, y_2, z_3, \dots, z_s) \in M' \setminus M$ , again a contradiction.  $\square$

**Lemma 3.3.** Suppose that  $\text{char}(R) \neq 2$ . For a subset  $M \subset X$ , the following conditions are equivalent.

- (a)  $B_M \in \mathcal{B}_\pm(G) \setminus \mathcal{B}(G)$  and  $B_M$  divides  $B$  in  $\mathcal{B}_\pm(G)$ .
- (b) There are  $i, j \in [1, s]$  distinct and  $\emptyset \neq Y_i \subsetneq X_i$ ,  $\emptyset \neq Y_j \subsetneq X_j$  such that  $M = \Delta(p_i^{-1}(Y_i), p_j^{-1}(Y_j))$ .

*Proof.* (b)  $\implies$  (a) We have

$$M = \Delta(p_i^{-1}(Y_i), p_j^{-1}(Y_j)) = \underbrace{(p_i^{-1}(Y_i) \setminus p_j^{-1}(Y_j))}_{M^+} \cup \underbrace{(p_j^{-1}(Y_j) \setminus p_i^{-1}(Y_i))}_{M^-}.$$



Then  $M = M^+ \cup M^-$  and  $\chi_{M^+} - \chi_{M^-} = \chi_{p_i^{-1}(Y_i)} - \chi_{p_j^{-1}(Y_j)} \in V$ , whence  $\prod_{x \in M} p(\chi_x) \in \mathcal{B}_\pm(G)$ . The complement  $X \setminus M$  has the same form, whence it also lies in  $\mathcal{B}_\pm(G)$  and so  $B_M$  is a divisor of  $B$  in  $\mathcal{B}_\pm(G)$ .

Assume to the contrary that  $\prod_{x \in M} p(\chi_x) \in \mathcal{B}(G)$ . Then  $\chi_{M^+} + \chi_{M^-} = 0 = \chi_{M^+} - \chi_{M^-}$ , whence  $2\chi_{M^+} = 2\chi_{M^-} = 0$ . Since  $\text{char}(R) \neq 2$ , it follows that  $M^+ = M^- = \emptyset$  and so  $M = \emptyset$ . On the other hand,  $\emptyset \neq Y_i \subsetneq X_i$  and  $\emptyset \neq Y_j \subsetneq X_j$  implies that  $M = \Delta(p_i^{-1}(Y_i), p_j^{-1}(Y_j)) \neq \emptyset$ , a contradiction.

(a)  $\implies$  (b) We distinguish two cases (these cases have a large overlap).

CASE 1:  $\text{char}(R) \neq 3$ .

Let  $B_M \in \mathcal{B}_\pm(G) \setminus \mathcal{B}(G)$ . Since  $\prod_{x \in M} p(\chi_x) \in \mathcal{B}_\pm(G)$ , there are  $M^+, M^- \subset M$  such that  $M^+ \uplus M^- = M$  and  $\chi_{M^+} - \chi_{M^-} \in V$ . Since  $\prod_{x \in M} p(\chi_x) \notin \mathcal{B}(G)$ , we have  $M^+ \neq \emptyset$ ,  $M^- \neq \emptyset$ , and  $M \neq X$ .

Let  $z \in X \setminus M$  be given. By (3.3), we have, for every  $x \in M$ , that

$$\chi_x \equiv \begin{cases} 0 & |\Delta(x, z)| \geq 2 \\ \chi_{p_i^{-1}(x_i)} & \Delta(x, z) = \{i\} \end{cases} \pmod{W_z}$$

Therefore, it follows that

$$\begin{aligned} \chi_{M^+} - \chi_{M^-} &= \sum_{x \in M^+} \chi_x - \sum_{x \in M^-} \chi_x \equiv \sum_{x \in M^+, |\Delta(x, z)|=1} \chi_x - \sum_{x \in M^-, |\Delta(x, z)|=1} \chi_x \\ &= \sum_{i=1}^s \sum_{x \in M^+, \Delta(x, z)=\{i\}} \chi_{p_i^{-1}(x_i)} - \sum_{i=1}^s \sum_{x \in M^-, \Delta(x, z)=\{i\}} \chi_{p_i^{-1}(x_i)} \pmod{W_z}. \end{aligned}$$

For every  $i \in [1, s]$ , we define

$$\begin{aligned} Y_i^+ &= \{y_i \in X_i : (z_1, \dots, z_{i-1}, y_i, z_{i+1}, \dots, z_n) \in M^+\} \quad \text{and} \\ Y_i^- &= \{y_i \in X_i : (z_1, \dots, z_{i-1}, y_i, z_{i+1}, \dots, z_n) \in M^-\} \end{aligned}$$

and obtain that

$$\chi_{M^+} - \chi_{M^-} \equiv \sum_{i=1}^s \chi_{p_i^{-1}(Y_i^+)} - \sum_{i=1}^s \chi_{p_i^{-1}(Y_i^-)} \pmod{W_z}.$$

Since  $\chi_{M^+} - \chi_{M^-}$ ,  $\sum_{i=1}^s \chi_{p_i^{-1}(Y_i^+)}$ , and  $\sum_{i=1}^s \chi_{p_i^{-1}(Y_i^-)}$  are in  $V$ , it follows that

$$\chi_{M^+} - \chi_{M^-} = \sum_{i=1}^s \chi_{p_i^{-1}(Y_i^+)} - \sum_{i=1}^s \chi_{p_i^{-1}(Y_i^-)}.$$

Assume to the contrary that there are distinct  $i, j \in [1, s]$  such that  $Y_i^+ \neq \emptyset$  and  $Y_j^+ \neq \emptyset$ . We pick  $u \in X$  with  $u_i \in Y_i^+$ ,  $u_j \in Y_j^+$ , and  $u_k = z_k$  for all  $k \in [1, s] \setminus \{i, j\}$ . Since  $z \notin M$ , we have  $z_k \notin Y_k^+ \cup Y_k^-$  and therefore

$$u \notin p_k^{-1}(Y_k^+) \cup p_k^{-1}(Y_k^-) \quad \text{for any } k \in [1, s] \setminus \{i, j\}.$$

Since  $M^+ \cap M^- = \emptyset$ , we obtain that  $Y_k^+ \cap Y_k^- = \emptyset$  for all  $k \in [1, s]$ . This implies that  $u \notin p_k^{-1}(Y_k^-)$  for all  $k \in [1, s]$ , but  $u \in p_i^{-1}(Y_i^+)$  and  $u \in p_j^{-1}(Y_j^+)$ . Therefore, we obtain that

$$2 = \sum_{k=1}^s \chi_{p_k^{-1}(Y_k^+)}(u) - \sum_{k=1}^s \chi_{p_k^{-1}(Y_k^-)}(u) = \chi_{M^+}(u) - \chi_{M^-}(u) \in \{-1, 0, 1\},$$

a contradiction to  $\text{char}(R) \neq 3$ .

The same arguments yield a contradiction, if we assume that there are distinct  $i, j \in [1, s]$  such that  $Y_i^- \neq \emptyset$  and  $Y_j^- \neq \emptyset$ .

Therefore, there are  $i, j \in [1, s]$  with

$$\chi_{M^+} - \chi_{M^-} = \chi_{p_i^{-1}(Y_i^+)} - \chi_{p_j^{-1}(Y_j^-)} = \chi_{p_i^{-1}(Y_i^+) \setminus p_j^{-1}(Y_j^-)} - \chi_{p_j^{-1}(Y_j^-) \setminus p_i^{-1}(Y_i^+)}.$$



This implies that

$$M^+ = p_i^{-1}(Y_i^+) \setminus p_j^{-1}(Y_j^-) \quad \text{and} \quad M^- = p_j^{-1}(Y_j^-) \setminus p_i^{-1}(Y_i^+),$$

whence

$$M = \Delta(p_i^{-1}(Y_i^+), p_j^{-1}(Y_j^-)).$$

Assume to the contrary that  $i = j$ . Then we obtain

$$M = \Delta(p_i^{-1}(Y_i^+), p_i^{-1}(Y_i^-)) = p_i^{-1}(\Delta(Y_i^+, Y_i^-)),$$

which implies that  $\chi_M \in V$  and hence  $\prod_{x \in M} p(\chi_x) \in \mathcal{B}(G)$ , a contradiction.

Assume to the contrary that  $Y_i^+ = \emptyset$  or that  $Y_i^+ = X_i$ . This implies that

$$M = \Delta(p_i^{-1}(Y_i^+), p_j^{-1}(Y_j^-)) = p_j^{-1}(Y_j^-),$$

which yields  $\prod_{x \in M} p(\chi_x) \in \mathcal{B}(G)$ , a contradiction. The same argument shows that  $\emptyset \neq Y_j^- \neq X_j$ .

CASE 2:  $\text{char}(R)$  is either zero or odd.

Since  $B_M$  divides  $B$ , there are pairwise disjoint sets  $M^+, M^-, N^+, N^- \subset X$  with

$$M^+ \uplus M^- \uplus N^+ \uplus N^- = X$$

and with  $\sigma(B_{M^+}) = \sigma(B_{M^-})$  and  $\sigma(B_{N^+}) = \sigma(B_{N^-})$ . This implies that

$$0 = \sigma(B) = \sigma(B_{M^+ \uplus M^- \uplus N^+ \uplus N^-}) = \sigma(B_{M^+}) + \sigma(B_{M^-}) + \sigma(B_{N^+}) + \sigma(B_{N^-}) = 2(\sigma(B_{M^+}) + \sigma(B_{N^+})).$$

Since  $\text{char}(R)$  is either zero or odd, it follows that  $\sigma(B_{M^+}) + \sigma(B_{N^+}) = 0$ . Thus,  $B_{M^+} B_{N^+} \in \mathcal{B}(G)$ , whence  $M^+ \cup N^+ = p_i^{-1}(Y_i)$  for some  $i \in [1, s]$  and some  $Y_i \subset X_i$ . For its complement, we get  $M^- \cup N^- = p_i^{-1}(X_i \setminus Y_i)$ . Similarly, we get, for some  $j \in [1, s]$  and some  $W_j \subset X_j$ ,

$$M^+ \cup N^- = p_j^{-1}(X_j \setminus W_j) \quad \text{and} \quad M^- \cup N^+ = p_j^{-1}(W_j).$$

Therefore,  $M^- = (M^- \cup N^-) \cap (M^- \cup N^+) = p_i^{-1}(X_i \setminus Y_i) \cap p_j^{-1}(W_j)$  and

$$M = M^+ \cup M^- = \Delta(p_i^{-1}(Y_i), p_j^{-1}(W_j)).$$

□

**Lemma 3.4.** *Let  $i, j \in [1, s]$  distinct,  $\emptyset \neq Y_i \subsetneq X_i$ ,  $\emptyset \neq Y_j \subsetneq X_j$  and  $M = \Delta(p_i^{-1}(Y_i), p_j^{-1}(Y_j))$ . Then  $B_M \in \mathcal{A}(\mathcal{B}_\pm(G))$ .*

*Proof.* Let  $N \subset M$  be a nonempty subset with  $B_N \in \mathcal{B}_\pm(G)$ . We have to show that  $N = M$  and to do so, we distinguish two cases.

CASE 1:  $B_N \in \mathcal{B}(G)$ .

By Lemma 3.2, there are  $k \in [1, s]$  and  $W_k \subset X_k$  such that  $N = p_k^{-1}(W_k)$ . Since  $\emptyset \neq N \neq X$ , we obtain that  $\emptyset \neq W_k \subsetneq X_k$ , whence

$$p_k^{-1}(W_k) \subset \Delta(p_i^{-1}(Y_i), p_j^{-1}(Y_j)).$$

CASE 1.1:  $k \in \{i, j\}$ , say  $k = i$ .

Let  $x_i \in W_i$ . We choose  $y_j \in X_j \setminus Y_j$  and  $u \in X$  with  $u_i = x_i$  and  $u_j = y_j$ . Then we have  $u \in p_k^{-1}(W_k) \subset \Delta(p_i^{-1}(Y_i), p_j^{-1}(Y_j))$ . Since  $y_j \notin Y_j$ , it follows that  $w_i = u_i \in Y_i$ , whence  $W_i \subset Y_i$ . Now we pick some  $u \in X$  with  $u_i \in W_i \subset Y_i$  and  $u_j \in Y_j$ . This implies that

$$u \in p_i^{-1}(W_i) \quad \text{but} \quad u \notin \Delta(p_i^{-1}(Y_i), p_j^{-1}(Y_j)),$$

a contradiction.

CASE 1.2:  $k \notin \{i, j\}$ .

We pick some  $u \in X$  with  $u_k \in W_k$ ,  $u_i \in Y_i$ , and  $u_j \in Y_j$ . Then

$$u \in p_k^{-1}(W_k) \quad \text{but} \quad u \notin \Delta(p_i^{-1}(Y_i), p_j^{-1}(Y_j)),$$

a contradiction.

CASE 2:  $B_N \notin \mathcal{B}(G)$ .

By Lemma 3.2, there are  $k, \ell \in [1, s]$  distinct,  $\emptyset \neq W_k \subsetneq X_k$ , and  $\emptyset \neq W_\ell \subsetneq X_\ell$  such that  $N = \Delta(p_k^{-1}(W_k), p_\ell^{-1}(W_\ell))$ , and we obtain that

$$\Delta(p_k^{-1}(W_k), p_\ell^{-1}(W_\ell)) \subset \Delta(p_i^{-1}(Y_i), p_j^{-1}(Y_j)).$$

We distinguish three cases.

CASE 2.1:  $\{k, \ell\} \cap \{i, j\} = \emptyset$ .

We pick  $u \in X$  with  $u_k \in W_k$ ,  $u_\ell \notin W_\ell$ ,  $u_i \in Y_i$ , and  $u_j \in Y_j$ . Then

$$u \in \Delta(p_k^{-1}(W_k), p_\ell^{-1}(W_\ell)) \quad \text{but} \quad u \notin \Delta(p_i^{-1}(Y_i), p_j^{-1}(Y_j)),$$

a contradiction.

CASE 2.2:  $|\{k, \ell\} \cap \{i, j\}| = 1$ , say  $k = i$ .

Let  $w_i \in W_i$ . We choose  $u \in X$  with  $u_i = w_i$ ,  $u_\ell \notin W_\ell$ , and  $u_j \notin Y_j$ . Then

$$u \in \Delta(p_i^{-1}(W_i), p_\ell^{-1}(W_\ell)) \quad \text{and} \quad u \in \Delta(p_i^{-1}(Y_i), p_j^{-1}(Y_j)).$$

Since  $u_j \notin Y_j$ , it follows that  $w_i = u_i \in Y_i$ , and hence  $W_i \subset Y_i$ .

Now we choose  $u \in X$  with  $u_i \in W_i$ ,  $u_\ell \notin W_\ell$ , and  $u_j \in Y_j$ . Then

$$u \in \Delta(p_i^{-1}(W_i), p_\ell^{-1}(W_\ell)) \quad \text{but} \quad u \notin \Delta(p_i^{-1}(Y_i), p_\ell^{-1}(W_\ell)),$$

a contradiction.

CASE 2.3:  $\{k, \ell\} = \{i, j\}$ , say  $k = i$  and  $\ell = j$ .

Then

$$\Delta(p_i^{-1}(W_i), p_j^{-1}(W_j)) \subset \Delta(p_i^{-1}(Y_i), p_j^{-1}(Y_j)).$$

Since  $W_i$  is nonempty, we have  $W_i \not\subset Y_i$  or  $W_i \not\subset X_i \setminus Y_i$ . Since

$$\Delta(p_i^{-1}(Y_i), p_j^{-1}(Y_j)) = \Delta(p_i^{-1}(X_i \setminus Y_i), p_j^{-1}(X_j \setminus Y_j)),$$

we may assume that  $W_i \not\subset X_i \setminus Y_i$ . We choose

$$x_j \in X_j \setminus W_j, x_i \in W_i \cap Y_i, \text{ and } u \in X \text{ with } u_i = x_i \text{ and } u_j = x_j.$$

Then  $u \in \Delta(p_i^{-1}(W_i), p_j^{-1}(W_j))$  and hence  $u \in \Delta(p_i^{-1}(Y_i), p_j^{-1}(Y_j))$ . Since  $u_i = x_i \in Y_i$ , we obtain that  $x_j = u_j \in X_j \setminus Y_j$ . Thus, it follows that  $X_j \setminus W_j \subset X_j \setminus Y_j$  and so  $Y_j \subset W_j$ .

Now we choose some  $w_i \in W_i$ . Let  $u \in X$  with  $u_i = w_i$  and  $u_j \in X_j \setminus W_j$ . Thus,

$$u \in \Delta(p_i^{-1}(W_i), p_j^{-1}(W_j)) \quad \text{and hence} \quad u \in \Delta(p_i^{-1}(Y_i), p_j^{-1}(Y_j)),$$

and, because  $u_j \in X_j \setminus W_j$ , we finally obtain that  $w_i = u_i \in Y_i$ . Summing up, we obtained that  $W_i \subset Y_i$ . Since  $\emptyset \neq Y_j \subset W_j$ , we have that  $W_j \not\subset X_j \setminus Y_j$ .

Interchanging  $i$  and  $j$  in the above arguments, we also obtain that  $Y_i \subset W_i$  and  $W_j \subset Y_j$ . Thus, we obtain that  $W_i = Y_i$  and  $W_j = Y_j$ , whence  $M = N$ .  $\square$

**Lemma 3.5.** *Let  $A \in \mathcal{A}(\mathcal{B}(G))$  be a divisor of  $B$  in  $\mathcal{B}(G)$ . Then  $A \in \mathcal{A}(\mathcal{B}_\pm(G))$ .*

*Proof.* By Lemma 3.2, there are  $k \in [1, s]$  and  $x_k \in X_k$  such that  $A = \prod_{x \in p_k^{-1}(x_k)} p(\chi_x)$ . Assume to the contrary that there is  $C \in \mathcal{B}_\pm(G)$  with  $1 \neq C \neq A$  such that  $C \mid A$  (in  $\mathcal{B}_\pm(G)$ ). Since  $A \in \mathcal{A}(\mathcal{B}(G))$ , it follows that  $C \notin \mathcal{B}(G)$ . Thus, Lemma 3.3 implies that there are  $i, j \in [1, s]$  distinct and  $\emptyset \neq Y_i \subsetneq X_i$ ,  $\emptyset \neq Y_j \subsetneq X_j$  such that

$$C = B_M \quad \text{with} \quad M = \Delta(p_i^{-1}(Y_i), p_j^{-1}(Y_j)).$$

This implies that

$$\Delta(p_i^{-1}(Y_i), p_j^{-1}(Y_j)) \subset p_k^{-1}(x_k).$$

If  $k \notin \{i, j\}$ , then we choose  $u \in X$  with  $u_i \in Y_i$ ,  $u_j \notin Y_j$ , and  $u_k \neq x_k$  (note that  $|X_k| \geq 2$ ). This implies that

$$u \in \Delta(p_i^{-1}(Y_i), p_j^{-1}(Y_j)) \quad \text{and} \quad u \notin p_k^{-1}(x_k),$$

a contradiction. Thus, we infer that  $k \in \{i, j\}$ , say  $k = i$ , and then

$$\Delta(p_i^{-1}(Y_i), p_j^{-1}(Y_j)) \subset p_i^{-1}(x_i).$$

Now we pick some  $y_i \in Y_i$  and some  $u \in X$  with  $u_i = y_i$  and  $u_j \notin Y_j$ . Then  $u \in p_i^{-1}(x_i)$ , whence  $y_i = u_i = x_i$  and thus  $Y_i = \{x_i\}$ . Now we choose some  $u \in X$  with  $u_i \neq x_i$  and  $u_j \in Y_j$ . Then

$$u \in \Delta(p_i^{-1}(Y_i), p_j^{-1}(Y_j)) \quad \text{but} \quad u \notin p_i^{-1}(x_i),$$

a contradiction.  $\square$

**3.3. Proof of Proposition 3.1.** Let  $R$  be a commutative ring, let  $L \subset \mathbb{N}_{\geq 2}$  be a finite subset with  $2 \in L$ , and let  $f: L \rightarrow \mathbb{N}$  be a map with  $s := \sum_{k \in L} f(k) \geq 3$ . Furthermore, let all notation be as in (3.1). In particular, we have

$$L = \{|X_1|, \dots, |X_s|\} \quad \text{and} \quad f(k) = |\{i \in [1, s]: |X_i| = k\}| \quad \text{for every } k \in L.$$

and

$$X = \prod_{j=1}^s X_j \quad \text{and} \quad X_J = \prod_{j \in J} X_j \quad \text{for } \emptyset \neq J \subset [1, s].$$

For every  $i \in [1, s]$ , we have

$$X = \biguplus_{y \in X_i} p_i^{-1}(y), \quad \text{hence} \quad B = \prod_{y \in X_i} B_{p_i^{-1}(y)}.$$

By Lemma 3.2, all  $B_{p_i^{-1}(y)}$  are atoms of  $\mathcal{B}(G)$  and by Lemma 3.5 they are atoms of  $\mathcal{B}_{\pm}(G)$ . Thus,

$$(3.6) \quad Z_i = \prod_{y \in X_i} B_{p_i^{-1}(y)} \quad \text{is a factorization of } B \text{ in } \mathcal{B}(G)$$

and

$$Z_{\pm, i} = \prod_{y \in X_i} B_{p_i^{-1}(y)} \quad \text{is a factorization of } B \text{ in } \mathcal{B}_{\pm}(G).$$

This implies that

$$Z_{\mathcal{B}(G)}(B) \supset \{Z_1, \dots, Z_s\} \quad \text{and} \quad Z_{\mathcal{B}_{\pm}(G)}(B) \supset \{Z_{\pm, 1}, \dots, Z_{\pm, s}\}$$

We continue with the following assertion.

**A.** If there exists some  $Z \in Z_{\mathcal{B}(G)}(B) \setminus \{Z_1, \dots, Z_s\}$  or some  $Z' \in Z_{\mathcal{B}_{\pm}(G)}(B) \setminus \{Z_{\pm, 1}, \dots, Z_{\pm, s}\}$  then  $2 = |Z| = |Z'|$ .

Suppose that **A** holds. Since  $2 \in L$  and since, for all  $i \in [1, s]$ ,  $|X_i| = |Z_i| = |Z_{\pm, i}|$ , it follows that

$$\mathbb{L}_{\mathcal{B}(G)}(B) = \{|X_1|, \dots, |X_s|\} = L = \mathbb{L}_{\mathcal{B}_{\pm}(G)}(B).$$

Furthermore, for all  $k \in L \setminus \{2\}$ , we have

$$|Z_{\mathcal{B}(G), k}(B)| = |Z_{\mathcal{B}_{\pm}(G), k}(B)| = |\{i \in [1, s] \mid |X_i| = k\}| = f(k).$$

If  $k = 2 = \min L$ , then Lemma 3.5 implies that  $|Z_{\mathcal{B}(G), k}(B)| \leq |Z_{\mathcal{B}_{\pm}(G), k}(B)|$  and **A** implies that  $|Z_{\mathcal{B}(G), 2}(B)| \geq |\{i \in [1, s]: |X_i| = 2\}| = f(2)$ .

*Proof of A.* We proceed in two steps.

(i) First, we deal with  $\mathcal{B}(G)$ .

Let  $Z \in Z_{\mathcal{B}(G)}(B) \setminus \{Z_1, \dots, Z_s\}$ . If  $i, j \in [1, s]$  are distinct,  $y_i \in X_i$  and  $y_j \in X_j$ , then  $p_i^{-1}(y_i) \cap p_j^{-1}(y_j) \neq \emptyset$ , and since  $B$  is squarefree, it follows that  $B_{p_i^{-1}(y_i)} B_{p_j^{-1}(y_j)} \nmid B$ . Hence there exists some

$A \in \mathcal{A}(\mathcal{B}(G)) \setminus \{B_{p_i^{-1}(y)} \mid i \in [1, s], y \in X_i\}$  with  $A \mid Z$ . By Lemma 3.2, this can only hold for  $\text{char}(R) = 2$ . We assert also that  $V = A^{-1}B \in \mathcal{A}(\mathcal{B}(G))$ . If this holds, then we get  $|Z| = 2$ , whence we are done. Indeed, if  $V \in \mathcal{B}(G) \setminus \mathcal{A}(\mathcal{B}(G))$ , then Lemma 3.2 implies that  $V = B_{p_i^{-1}(Y_i)}$  for some  $i \in [1, s]$  and  $Y_i \subset X_i$ , and consequently  $A = V^{-1}B = B_{p_i^{-1}(X_i \setminus Y_i)}$ . But, then Lemma 3.2 implies that  $|X_i \setminus Y_i| = 1$ , a contradiction to the choice of  $A$ .

(ii) Next, we deal with  $\mathcal{B}_{\pm}(G)$ , and we may assume that  $\text{char}(R) \neq 2$ .

Let  $Z' = A_1 \cdots A_m \in \mathcal{Z}_{\mathcal{B}_{\pm}(G)}(B) \setminus \{Z_{\pm,1}, \dots, Z_{\pm,s}\}$ . If  $A_1, \dots, A_m \in \mathcal{B}(G)$ , then they are atoms in  $\mathcal{B}(G)$  by Relation (2.1), whence  $Z' \in \mathcal{Z}_{\mathcal{B}(G)}(B) \setminus \{Z_1, \dots, Z_s\}$  and so  $|Z'| = 2$  by (i). Now suppose that there is  $k \in [1, m]$  with  $A_k \notin \mathcal{B}(G)$ , say  $k = 1$ . Then  $A_2 \cdots A_m \in \mathcal{B}_{\pm}(G) \setminus \mathcal{B}(G)$ . Then Lemmas 3.3 and 3.4 imply that  $A_2 \cdots A_m \in \mathcal{A}(\mathcal{B}_{\pm}(G))$ , whence  $|Z'| = m = 2$ .  $\square$

#### 4. ON THE CONSTRUCTION OF A GROUP HAVING A GIVEN $L$ AS A SET OF LENGTHS

The goal in this section is to prove the following result.

**Proposition 4.1.** *Let  $R$  be a commutative ring, let  $L \subset \mathbb{N}_{\geq 2}$  be a finite, nonempty subset, and let  $f: L \rightarrow \mathbb{N}$  be a map. Then there exist a finitely generated free  $R$ -module  $G$  and some  $S \in \mathcal{B}(G)$  with the following properties.*

- (a)  $S$  is squarefree in  $\mathcal{F}(G)$ .
- (b)  $\mathcal{L}_{\mathcal{B}_{\pm}(G)}(S) = \mathcal{L}_{\mathcal{B}(G)}(S) = L$ .
- (c)  $|\mathcal{Z}_{\mathcal{B}_{\pm}(G),k}(S)| \geq |\mathcal{Z}_{\mathcal{B}(G),k}(S)| \geq f(k)$  for all  $k \in L$ . Moreover, both inequalities are equalities for  $k > \min L$ .

*Proof.* We set  $s = \sum_{k \in L} f(k)$  and distinguish several cases. Throughout, we use Relation (2.1) and Lemma 2.1 without further mention.

CASE 1:  $s = 1$  and  $L = \{2\}$ .

We set  $G = R^4$  and choose an  $R$ -basis  $(e_1, e_2, f_1, f_2)$  of  $G$ . Then  $U_1 = e_1 f_1 (-e_1 - f_1)$  and  $U_2 = e_2 f_2 (-e_2 - f_2)$  are atoms of  $\mathcal{B}(G)$  and of  $\mathcal{B}_{\pm}(G)$ . Thus,  $S = U_1 U_2 \in \mathcal{B}(G) \subset \mathcal{B}_{\pm}(G)$  is squarefree with  $\mathcal{L}_{\mathcal{B}(G)}(S) = \mathcal{L}_{\mathcal{B}_{\pm}(G)}(S) = \{2\}$ , and with  $|\mathcal{Z}_{\mathcal{B}(G),2}(S)| = |\mathcal{Z}_{\mathcal{B}_{\pm}(G),2}(S)| = f(2) = 1$ .

CASE 2:  $s = 2$ ,  $L = \{2\}$ , and  $\text{char}(R) \neq 2$ .

We set  $G = R^3$  and choose an  $R$ -basis  $(e_1, e_2, e_3)$  of  $G$ . Then

$$U_1 = (-e_1)e_1, U_2 = e_2 e_3 (e_1 - e_3)(-e_1 - e_2), U_3 = e_1 e_2 (-e_1 - e_2), \quad \text{and} \quad U_4 = (-e_1)e_3 (e_1 - e_3)$$

are atoms of  $\mathcal{B}(G)$  and of  $\mathcal{B}_{\pm}(G)$ . Thus,  $S = U_1 U_2 = U_3 U_4 \in \mathcal{B}(G) \subset \mathcal{B}_{\pm}(G)$  is squarefree with  $\mathcal{L}_{\mathcal{B}(G)}(S) = \mathcal{L}_{\mathcal{B}_{\pm}(G)}(S) = \{2\}$ , and with  $|\mathcal{Z}_{\mathcal{B}(G),2}(S)| = |\mathcal{Z}_{\mathcal{B}_{\pm}(G),2}(S)| = f(2) = 2$ .

CASE 3:  $s = 2$ ,  $L = \{2\}$ , and  $\text{char}(R) = 2$ .

We set  $G = R^4$  and choose an  $R$ -basis  $(e_1, e_2, e_3, e_4)$  of  $G$ . We define  $e_0 = e_1 + e_2 + e_3 + e_4$  and consider

$$\begin{aligned} S &= ((e_1 + e_2)(e_3 + e_4)e_0)(e_1 e_2 e_3 (e_0 + e_4)) \\ &= (e_1 e_2 (e_1 + e_2))(e_3 (e_0 + e_4)(e_3 + e_4)e_0) \\ &= ((e_1 + e_2)e_3 (e_0 + e_4))(e_1 e_2 (e_3 + e_4)e_0). \end{aligned}$$

Then  $S \in \mathcal{B}(G) = \mathcal{B}_{\pm}(G)$  is squarefree,  $\mathcal{L}_{\mathcal{B}(G)}(S) = \mathcal{L}_{\mathcal{B}_{\pm}(G)}(S) = 2$ , and  $|\mathcal{Z}_{\mathcal{B}(G),2}(S)| = |\mathcal{Z}_{\mathcal{B}_{\pm}(G),2}(S)| = 3 > 2 = f(2)$ .

CASE 4:  $s = 2$ ,  $L = \{2, r\}$  with  $r \geq 3$ , and  $\text{char}(R) \neq 2$ .

We set  $G = R^{r-1}$  and choose an  $R$ -basis  $(e_1, \dots, e_{r-1})$  of  $G$ . Then  $U = e_0 e_1 \cdots e_{r-1}$ , with  $e_0 = -(e_1 + \dots + e_{r-1})$ , and  $U_i = (-e_i)e_i$ , with  $i \in [0, r]$ , are atoms of  $\mathcal{B}(G)$  and of  $\mathcal{B}_{\pm}(G)$ . Thus,

$$S = (-U)U = U_0 \cdots U_{r-1} \in \mathcal{B}(G) \subset \mathcal{B}_{\pm}(G)$$

is squarefree,  $\mathcal{L}_{\mathcal{B}(G)}(S) = \mathcal{L}_{\mathcal{B}_{\pm}(G)}(S) = \{2, r\}$  and  $|\mathcal{Z}_{\mathcal{B}(G),k}(S)| = |\mathcal{Z}_{\mathcal{B}_{\pm}(G),k}(S)| = f(k) = 1$  for every  $k \in L$ .

CASE 5:  $s = 2$ ,  $L = \{2, r\}$  with  $r \geq 3$ , and  $\text{char}(R) = 2$ .

We set  $G = R^{2r-1}$  and choose an  $R$ -basis  $(e_1, \dots, e_{2r-1})$  of  $G$ . With  $e_0 = e_1 + \dots + e_{2r-2}$ , the sequences

$$U_1 = e_1 \cdot \dots \cdot e_{2r-1}(e_0 + e_{2r-1}), \quad U_2 = (e_1 + e_2)(e_3 + e_4) \cdot \dots \cdot (e_{2r-3} + e_{2r-2})e_0,$$

$$V_1 = e_1 \cdot \dots \cdot e_{2r-2}e_0, \quad \text{and} \quad V_2 = (e_1 + e_2)(e_3 + e_4) \cdot \dots \cdot (e_{2r-3} + e_{2r-2})e_{2r-1}(e_{2r-1} + e_0)$$

are in  $\mathcal{B}(G) = \mathcal{B}_\pm(G)$ . We define

$$S = U_1 U_2 = V_1 V_2 = (e_1 e_2 (e_1 + e_2)) \cdot \dots \cdot (e_{2r-3} e_{2r-2} (e_{2r-3} + e_{2r-2})) (e_{2r-1} e_0 (e_{2r-1} + e_0)).$$

By construction,  $S \in \mathcal{B}(G) = \mathcal{B}_\pm(G)$  is squarefree and, obviously,  $\mathsf{L}_{\mathcal{B}(G)}(S) = \mathsf{L}_{\mathcal{B}_\pm(G)}(S) = \{2, r\}$ ,  $|\mathsf{Z}_{\mathcal{B}(G),2}(S)| = |\mathsf{Z}_{\mathcal{B}_\pm(G),2}(S)| = 2 > 1 = f(2)$ , and  $|\mathsf{Z}_{\mathcal{B}(G),r}(S)| = |\mathsf{Z}_{\mathcal{B}_\pm(G),r}(S)| = 1 = f(r)$ .

CASE 6:  $s \geq 3$  and  $2 \in L$ .

This follows from Proposition 3.1.

CASE 7:  $2 \notin L$ .

We set  $m = -2 + \min L$ ,  $L_0 = -m + L \subset \mathbb{N}_{\geq 2}$ , and we define

$$f_0: L_0 \rightarrow \mathbb{N} \quad \text{by} \quad f_0(k) = f(k+m) \quad \text{for every } k \in L_0.$$

Since  $2 \in L_0$ , the previous cases imply that there exist a finitely generated free  $R$ -module  $G_1$  and some  $S_1 \in \mathcal{B}(G_1) \subset \mathcal{B}_\pm(G_1)$  satisfying Properties (a), (b), and (c). In particular,  $\mathsf{L}_{\mathcal{B}(G)}(S_1) = \mathsf{L}_{\mathcal{B}_\pm(G)}(S_1) = L_0$ .

Now we set  $G = G_1 \oplus R^{2m}$  and we choose an  $R$ -basis  $(e_1, f_1, \dots, e_m, f_m)$  of  $R^{2m}$ . Then

$$S_2 = \prod_{j=1}^m (e_j f_j (-e_j - f_j)) \in \mathcal{B}(R^{2m}) \subset \mathcal{B}_\pm(R^{2m})$$

has unique factorization in  $\mathcal{B}(R^{2m})$  and in  $\mathcal{B}_\pm(R^{2m})$ , say  $\mathsf{Z}_{\mathcal{B}(R^{2m})}(S_2) = \mathsf{Z}_{\mathcal{B}_\pm(R^{2m})}(S_2) = \{y\}$  with  $|y| = m$ . Now we consider the sequence

$$S = S_1 S_2 \in \mathcal{B}(G_1) \times \mathcal{B}(R^{2m}) \subset \mathcal{B}(G) \subset \mathcal{B}_\pm(G).$$

Clearly,  $S$  is squarefree in  $\mathcal{F}(G)$ ,  $\mathsf{Z}_{\mathcal{B}(G)}(S) = \{yz: z \in \mathsf{Z}_{\mathcal{B}(G_1)}(S_1)\}$  and  $\mathsf{Z}_{\mathcal{B}_\pm(G)}(S) = \{yz: z \in \mathsf{Z}_{\mathcal{B}_\pm(G_1)}(S_1)\}$ , which implies that

$$\mathsf{L}_{\mathcal{B}_\pm(G)}(S) = |y| + \mathsf{L}_{\mathcal{B}_\pm(G_1)}(S_1) = m + L_0 = L = \mathsf{L}_{\mathcal{B}(G)}(S)$$

and, for every  $k \in L$ ,

$$|\mathsf{Z}_{\mathcal{B}_\pm(G),k}(S)| = |\mathsf{Z}_{\mathcal{B}_\pm(G_1),k-m}(S_1)| \quad \text{and} \quad |\mathsf{Z}_{\mathcal{B}(G),k}(S)| = |\mathsf{Z}_{\mathcal{B}(G_1),k-m}(S_1)|,$$

Thus,  $S \in \mathcal{B}(G)$  satisfies Properties (a), (b), and (c). □

## 5. PROOF OF THEOREM 1.1 AND OF COROLLARY 1.2

For an abelian group  $G$ , we denote by  $\mathsf{T}(G)$  its torsion subgroup.

**Proposition 5.1.** *Let  $G$  be a finitely generated abelian group of torsionfree rank  $r \in \mathbb{N}$  and let  $S \in \mathcal{B}_\pm(G)$ . Then there are  $N \in \mathbb{N}$  and homomorphisms*

$$\varphi_0: G \rightarrow G_0 := \mathbb{Z} \times \mathsf{T}(G), \quad \varphi_n: G \rightarrow G_n := (\mathbb{Z}/n\mathbb{Z})^r \times \mathsf{T}(G) \quad \text{for all } n \geq N,$$

such that the following properties hold for all  $n \in \{0\} \cup \mathbb{N}_{\geq N}$ .

1. If  $S$  is squarefree in  $\mathcal{F}(G)$ , then  $\varphi_n(S)$  is squarefree in  $\mathcal{F}(G_n)$ .
2. The map  $\varphi_n$  induces a bijective map  $\overline{\varphi}_n: \mathsf{Z}_{\mathcal{B}_\pm(G)}(S) \rightarrow \mathsf{Z}_{\mathcal{B}_\pm(G_n)}(\varphi_n(S))$  such that the following three properties hold.
  - (a)  $|z| = |\overline{\varphi}_n(z)|$  for all  $z \in \mathsf{Z}_{\mathcal{B}_\pm(G)}(S)$ .
  - (b)  $\mathsf{L}_{\mathcal{B}_\pm(G)}(S) = \mathsf{L}_{\mathcal{B}_\pm(G_n)}(\varphi_n(S))$ .
  - (c)  $|\{z \in \mathsf{Z}_{\mathcal{B}_\pm(G)}(S): |z| = k\}| = |\{z \in \mathsf{Z}_{\mathcal{B}_\pm(G_n)}(\varphi_n(S)): |z| = k\}|$  for every  $k \in \mathsf{L}_{\mathcal{B}_\pm(G)}(S)$ .

3. If  $S \in \mathcal{B}(G)$ , then  $\varphi_n$  induces a bijective map  $\varphi'_n: \mathcal{Z}_{\mathcal{B}(G)}(S) \rightarrow \mathcal{Z}_{\mathcal{B}(G_n)}(\varphi_n(S))$  such that the Properties (a) - (c) of Part 2 hold for  $\mathcal{B}(G)$  and  $\mathcal{B}(G_n)$ .

*Proof.* We may suppose that  $G = \mathbb{Z}^r \times \mathsf{T}(G)$  and let  $\mathbf{p}: G \rightarrow \mathbb{Z}^r$  denote the projection. We set  $S = g_1 \cdot \dots \cdot g_\ell$ , where  $\ell \in \mathbb{N}_0$  and  $g_1, \dots, g_\ell \in G$ . It suffices to show that there are  $N \in \mathbb{N}$  and homomorphisms  $\varphi_n: G \rightarrow (\mathbb{Z}/n\mathbb{Z})^r \times \mathsf{T}(G)$ , for every  $n \in \{0\} \cup \mathbb{N}_{\geq N}$ , which are injective on the set

$$\Sigma(S) = \left\{ \sum_{\nu \in I} g_\nu : \emptyset \neq I \subset [1, \ell], \text{ for all } \nu \in I \right\},$$

which settles Part 3., and on the set

$$\Sigma_\pm(S) = \left\{ \sum_{\nu \in I} \varepsilon_\nu g_\nu : \emptyset \neq I \subset [1, \ell], \varepsilon_\nu \in \{-1, 1\} \text{ for all } \nu \in I \right\},$$

which settles Part 2. (in case when  $S \in \mathcal{B}(G)$ ). Clearly,  $\Sigma(S) \subset \Sigma_\pm(S)$  and the set

$$E = \{\mathbf{m} - \mathbf{n} : \mathbf{m}, \mathbf{n} \in \mathbf{p}(\Sigma_\pm(S)), \mathbf{m} \neq \mathbf{n}\} \subset \mathbb{Z}^r \setminus \{\mathbf{0}\}$$

is finite, say  $E = \{\mathbf{n}^{(1)}, \dots, \mathbf{n}^{(d)}\}$ . We assert that there is a homomorphism  $\psi: \mathbb{Z}^r \rightarrow \mathbb{Z}$  such that  $\psi(\mathbf{n}^{(\nu)}) \neq 0$  for all  $\nu \in [1, d]$ . If this holds, then we set  $\varphi_0: G \rightarrow G_0$  by  $\varphi_0(\mathbf{u}, g) = (\psi(\mathbf{u}), g)$ , and obviously  $\varphi_0$  has the required property.

For  $j \in [1, d]$ , let  $\mathbf{n}^{(j)} = (n_1^{(j)}, \dots, n_r^{(j)})$ , and consider the non-zero polynomial

$$f = \prod_{j=1}^d \left( \sum_{i=1}^r n_i^{(j)} X_i \right) \in \mathbb{Z}[X_1, \dots, X_r].$$

If  $(a_1, \dots, a_r) \in \mathbb{Z}^r$  is such that  $f(a_1, \dots, a_r) \neq 0$ , then  $\psi: \mathbb{Z}^r \rightarrow \mathbb{Z}$ , defined by  $\psi(x_1, \dots, x_r) = a_1 x_1 + \dots + a_r x_r$ , is a non-zero homomorphism satisfying  $\psi(\mathbf{n}^{(j)}) \neq 0$  for all  $j \in [1, d]$ .

For  $n \in \mathbb{N}$ , let  $\psi_n: \mathbb{Z}^r \rightarrow (\mathbb{Z}/n\mathbb{Z})^r$  be the canonical epimorphism. Then there exists some  $N \in \mathbb{N}$  such that  $\psi_n|_{\mathbf{p}(\Sigma_\pm(S))}$  is injective for all  $n \geq N$  and the homomorphism  $\varphi_n = \psi_n \times \text{id}_{\mathsf{T}(G)}: G \rightarrow G_n$  has the required property.  $\square$

We recall the definition of almost arithmetic multiprogressions (AAMPs) and of the set of minimal distances of a monoid. To begin with AAMPs, let  $d \in \mathbb{N}$ ,  $M \in \mathbb{N}_0$  and  $\{0, d\} \subset \mathcal{D} \subset [0, d]$ . Then  $L$  is called an *almost arithmetic multiprogression* (AAMP for short) with *difference*  $d$ , *period*  $\mathcal{D}$ , and *bound*  $M$ , if

$$L = y + (L' \cup L^* \cup L'') \subset y + \mathcal{D} + d\mathbb{Z}$$

where

- $L^*$  is finite and nonempty with  $\min L^* = 0$  and  $L^* = (\mathcal{D} + d\mathbb{Z}) \cap [0, \max L^*]$ ,
- $L' \subset [-M, -1]$  and  $L'' \subset \max L^* + [1, M]$ , which are called the *initial part* and the *end part* of  $L$ , and
- $y \in \mathbb{Z}$ .

In particular, AAMPs are finite and nonempty subsets of the integers. For an overview of monoids and domains, whose sets of lengths are AAMPs with global bounds on all parameters, we refer to [14, Chapter 4]. Schmid proved that for Krull monoids with finite class group the description of sets of lengths as AAMPs is best possible ([32]).

Next we recall the concept of minimal distances. For a finite, nonempty subset  $L = \{m_0, \dots, m_k\} \subset \mathbb{Z}$  with  $m_0 < \dots < m_k$ , let  $\Delta(L) = \{m_i - m_{i-1} : i \in [1, k]\} \subset \mathbb{N}$  denote its set of distances. Now let  $H$  be a BF-monoid. Then every divisor-closed submonoid  $S \subset H$  is a BF-monoid and

$$\Delta(S) = \bigcup_{a \in S} \Delta(\mathsf{L}_S(a)) \subset \mathbb{N}$$

denotes the *set of distances* of  $S$ . Then

$$\Delta^*(H) = \{\min \Delta(S) : S \subset H \text{ is a divisor-closed submonoid with } \Delta(S) \neq \emptyset\} \subset \Delta(H)$$

is the *set of minimal distances* of  $H$ . In the next lemma, we gather some simple properties, which highlight the differences between  $\Delta^*(\mathcal{B}(G))$  and  $\Delta^*(\mathcal{B}_\pm(G))$  and which shows how different  $\Delta^*(\mathcal{B}_\pm(G))$  can be for different groups. Let

$$D(\mathcal{B}_\pm(G)) = \max \{|S| : S \in \mathcal{A}(\mathcal{B}_\pm(G))\}$$

denote the *plus-minus weighted Davenport constant* of  $G$ , which is studied, among others, in [28, 27, 24].

**Proposition 5.2.** *Let  $G$  be an abelian group.*

1. *If  $G$  is finite with  $|G| > 2$ , exponent  $n \geq 2$ , and rank  $r \geq 1$ , then  $\max \Delta^*(\mathcal{B}(G)) = \max\{r-1, n-2\}$  and  $\max \Delta^*(\mathcal{B}_\pm(G)) \leq D(\mathcal{B}_\pm(G)) - 2$ .*
2. *If  $G$  is infinite, then  $\Delta^*(\mathcal{B}(G)) = \mathbb{N}$ .*
3. *If  $G$  is a finite elementary 2-group of rank  $r$ , then  $\Delta^*(\mathcal{B}(G)) = \Delta^*(\mathcal{B}_\pm(G)) = [1, r-1]$ .*
4. *If  $G$  is an elementary 3-group (finite or infinite), then  $\Delta^*(\mathcal{B}_\pm(G)) = \{1\}$ .*

*Proof.* 1. For the first statement see [18], and for the second see [29].

2. See [5, Theorem 1.1].

3. Since  $\mathcal{B}(G) = \mathcal{B}_\pm(G)$ , we have  $\Delta^*(\mathcal{B}(G)) = \Delta^*(\mathcal{B}_\pm(G))$ . The equality  $\Delta^*(\mathcal{B}(G)) = [1, r-1]$  follows from [14, Corollary 6.8.3].

4. Let  $S \subset \mathcal{B}_\pm(G)$  be a divisor-closed submonoid with  $\Delta(S) \neq \emptyset$ . Then there is a nonempty subset  $G_0 \subset G$  with  $\mathcal{B}_\pm(G_0) = S$ . Let  $g \in G_0$  be a nonzero element. Then  $\text{ord}(g) = 3$ ,  $A = g^3$  and  $U = g^2$  are atoms in  $\mathcal{B}_\pm(G_0)$ . Since  $A^2 = U^3$ , it follows that  $1 \in \Delta(S)$ , whence  $\min \Delta(S) = 1$ .  $\square$

We refer to [18, 34, 30] for recent progress on  $\Delta^*(\mathcal{B}(G))$ . For the monoid  $\mathcal{B}_\pm(G)$ , its set of minimal distances was recently studied by Schmid et al. [29], and it turned out, as already indicated by Proposition 5.2, that the structure of  $\Delta^*(\mathcal{B}_\pm(G))$  is in general quite different from the structure of  $\Delta^*(\mathcal{B}(G))$ .

Before completing the proofs of Theorem 1.1 and of Corollary 1.2, let us compare the statements of Part 1 and of Part 2 of Theorem 1.1 for elementary 3-groups. Let  $G$  be an elementary 3-group. If  $G \cong C_3^r$  with  $r \in \mathbb{N}$ , then, apart from a globally bounded initial and end part, all sets of lengths in  $\mathcal{B}_\pm(G)$  are intervals by Theorem 1.1.1 and by Proposition 5.2.4. Nevertheless, if  $G$  is infinite, then every finite, nonempty subset of  $\mathbb{N}_{\geq 2}$  occurs as a set of lengths. This shows that the bounds  $M(C_3^r)$ , occurring in Theorem 1.1.1, tend to infinity as  $r$  tends to infinity.

An abelian group  $G$  is said to be *bounded* if there is  $m \in \mathbb{N}$  such that  $mG = \mathbf{0}$ .

*Proof of Theorem 1.1.* Let  $G$  be an abelian group.

1. By [7, Theorem 3.7],  $\mathcal{B}_\pm(G)$  is finitely generated if and only if  $G$  is finite. Thus, if  $G$  is finite, then Theorem 4.4.11 in [14] implies that there is some  $M \in \mathbb{N}_0$  such that, for every plus-minus weighted zero-sum sequence  $S$  over  $G$ , its set of lengths  $\mathsf{L}_{\mathcal{B}_\pm(G)}(S)$  is an AAMP with difference in  $\Delta^*(\mathcal{B}_\pm(G))$  and bound  $M$ .

2. Now suppose that  $G$  is infinite. Let  $L \subset \mathbb{N}_{\geq 2}$  be a finite, nonempty subset and let  $f: L \rightarrow \mathbb{N}$  be a map. We show that there is an abelian group  $G_1$ , which is isomorphic to a subgroup of  $G_2 \subset G$ , and some  $S \in \mathcal{B}(G_1)$  having the required properties. If this holds, then the assertion follows because  $\mathcal{B}_\pm(G_2)$  is a divisor-closed submonoid of  $\mathcal{B}_\pm(G)$ .

We distinguish three cases.

CASE 1:  $G$  is not a torsion group.

Then  $G$  has a subgroup isomorphic to  $\mathbb{Z}$ . By Proposition 4.1, there exist some  $r \in \mathbb{N}$  and some  $S' \in \mathcal{B}(\mathbb{Z}^r)$  such that  $S'$  satisfies the Properties (a), (b), and (c) in Proposition 4.1. By Proposition 5.1, there exists some  $S \in \mathcal{B}(\mathbb{Z})$  satisfying the same Properties (a), (b), and (c).

CASE 2:  $G$  is an unbounded torsion group.



By CASE 1, there exists some  $S' \in \mathcal{B}(\mathbb{Z})$  satisfying the Properties (a), (b), and (c) in Proposition 4.1. By Proposition 5.1, there exists some  $N \in \mathbb{N}$  such that for every  $n \geq N$  there is some  $S_n \in \mathcal{B}_\pm(\mathbb{Z}/n\mathbb{Z})$  with the required properties. Since  $G$  contains a cyclic subgroup of order  $n$  for some  $n \geq N$ , the assertion follows.

CASE 3:  $G$  is bounded.

By [31, Chapter 4],  $G$  is a direct sum of cyclic groups. Hence there is some  $n \in \mathbb{N}_{\geq 2}$  such that  $G$  contains a subgroup isomorphic to  $(\mathbb{Z}/n\mathbb{Z})^{(\mathbb{N})}$ . In particular, for every  $r \in \mathbb{N}$ ,  $G$  contains a subgroup isomorphic to  $(\mathbb{Z}/n\mathbb{Z})^r$ , whence the assertion follows by Proposition 4.1.  $\square$

*Proof of the Corollary 1.2.* Let  $L \subset \mathbb{N}_{\geq 2}$  be a finite, nonempty subset. For a prime  $p$  and  $r \in \mathbb{N}$ , let  $C_p^r$  denote an elementary  $p$ -group of rank  $r$ . We prove the following two claims.

**C1** There is  $N \in \mathbb{N}$  such that  $L \in \mathcal{L}(\mathcal{B}_\pm(\mathbb{Z}/n\mathbb{Z}))$  for all  $n \geq N$ .

**C2** For every prime  $p$  there is  $s_p \in \mathbb{N}$  such that  $L \in \mathcal{L}(\mathcal{B}_\pm(C_p^{s_p}))$  for all  $s \geq s_p$ .

Suppose that **C1** and **C2** hold and let  $G$  be a finite abelian group with  $L \notin \mathcal{L}(\mathcal{B}_\pm(G))$ . Then **C1** shows that  $\exp(G) < N$  and **C2** implies that the  $p$ -rank of  $G$  is bounded above by  $s_p$  for all primes  $p$  dividing  $\exp(G)$ . Thus, the assertion follows.

*Proof of C1.* By Theorem 1.1, there is some  $S = m_1 \cdot \dots \cdot m_\ell \in \mathcal{B}(\mathbb{Z})$  such that  $L = \mathcal{L}_{\mathcal{B}_\pm(\mathbb{Z})}(S)$ . We set  $N = 1 + \sum_{i=1}^\ell |m_i|$ . Let  $n \in \mathbb{N}$  with  $n \geq N$  and let  $\varphi_n: \mathbb{Z} \rightarrow \mathbb{Z}/n\mathbb{Z}$  denote the canonical epimorphism. If  $A \in \mathcal{F}(\mathbb{Z})$  with  $A|S$  in  $\mathcal{F}(\mathbb{Z})$ , then  $A \in \mathcal{B}_\pm(\mathbb{Z})$  if and only if  $\varphi_n(A) \in \mathcal{B}_\pm(\mathbb{Z}/n\mathbb{Z})$ . This implies that  $\mathcal{L}_{\mathcal{B}_\pm(\mathbb{Z})}(S) = \mathcal{L}_{\mathcal{B}_\pm(\mathbb{Z}/n\mathbb{Z})}(\varphi_n(S))$ .

*Proof of C2.* Let  $p$  be a prime and let  $G$  be an elementary  $p$ -group of infinite rank. By Theorem 1.1, there is some  $S \in \mathcal{B}(G)$  such that  $\mathcal{L}_{\mathcal{B}_\pm(G)}(S) = L$ . Clearly,  $G_1 = \langle \text{supp}(S) \rangle \subset G$  is an elementary  $p$ -group of finite rank,  $S \in \mathcal{B}(G_1)$ , and  $L = \mathcal{L}_{\mathcal{B}_\pm(G_1)}(S)$ . Thus, **C2** holds with  $s_p$  being the  $p$ -rank of  $G_1$ .  $\square$

## REFERENCES

- [1] S.D. Adhikari, Y.G. Chen, J.B. Friedlander, S.V. Konyagin, and F. Pappalardi, *Contributions to zero-sum problems*, Discrete Math. **306** (2006), 1 – 10.
- [2] P. Baginski, R. Rodriguez, G.J. Schaeffer, and Y. She, *Finding elements with given factorization lengths and multiplicities*, Amer. Math. Monthly **123** (2016), 849 – 870.
- [3] A. Bashir and A. Reinhart, *On transfer Krull monoids*, Semigroup Forum **105** (2022), 73 – 95.
- [4] S. Boukheche, K. Merito, O. Ordaz, and W.A. Schmid, *Monoids of sequences over finite abelian groups defined via zero-sums with respect to a given set of weights and applications to factorizations of norms of algebraic integers*, Commun. Algebra **50** (2022), 4195 – 4217.
- [5] S.T. Chapman, W.A. Schmid, and W.W. Smith, *On minimal distances in Krull monoids with infinite class group*, Bull. Lond. Math. Soc. **40** (2008), 613 – 618.
- [6] J. Coykendall and R. Hasenauer, *Norms, Normsets, and Factorization*, in Recent Progress in Ring and Factorization Theory, Springer, 2025.
- [7] F. Fabsits, A. Geroldinger, A. Reinhart, and Q. Zhong, *On monoids of plus-minus weighted zero-sum sequences: The isomorphism problem and the characterization problem*, J. Commutative Algebra **16** (2024), 1 – 23.
- [8] V. Fadinger and D. Windisch, *Lengths of factorizations of integer-valued polynomials on Krull domains with prime elements*, Arch. Math. (Basel), to appear (2024).
- [9] V. Fadinger and Q. Zhong, *On product-one sequences over subsets of groups*, Periodica Math. Hung. **86** (2023), 454 – 494.
- [10] V. Fadinger-Held, S. Frisch, and D. Windisch, *Integer-valued polynomials on discrete valuation rings of global fields with prescribed lengths of factorizations*, Monatsh. Math. **202** (2023), 773 – 789.
- [11] S. Frisch, *A construction of integer-valued polynomials with prescribed sets of lengths of factorizations*, Monatsh. Math. **171** (2013), 341 – 350.
- [12] S. Frisch, S. Nakato, and R. Rissner, *Sets of lengths of factorizations of integer-valued polynomials on Dedekind domains with finite residue fields*, J. Algebra **528** (2019), 231 – 249.
- [13] A. Geroldinger and F. Gotti, *On monoid algebras having every nonempty subset of  $\mathbb{N}_{\geq 2}$  as a length set*, <https://arxiv.org/abs/2404.11494>.

- [14] A. Geroldinger and F. Halter-Koch, *Non-Unique Factorizations. Algebraic, Combinatorial and Analytic Theory*, Pure and Applied Mathematics, vol. 278, Chapman & Hall/CRC, 2006.
- [15] A. Geroldinger, F. Halter-Koch, and Q. Zhong, *On monoids of weighted zero-sum sequences and applications to norm monoids in Galois number fields and binary quadratic forms*, Acta Math. Hung. **168** (2022), 144 – 185.
- [16] A. Geroldinger and W.A. Schmid, *A realization theorem for sets of lengths in numerical monoids*, Forum Math. **30** (2018), 1111 – 1118.
- [17] A. Geroldinger, W.A. Schmid, and Q. Zhong, *Systems of sets of lengths: transfer Krull monoids versus weakly Krull monoids*, in Rings, Polynomials, and Modules, Springer, Cham, 2017, pp. 191 – 235.
- [18] A. Geroldinger and Q. Zhong, *The set of minimal distances in Krull monoids*, Acta Arith. **173** (2016), 97 – 120.
- [19] ———, *Factorization theory in commutative monoids*, Semigroup Forum **100** (2020), 22 – 51.
- [20] F. Gotti, *Systems of sets of lengths of Puiseux monoids*, J. Pure Appl. Algebra **223** (2019), 1856 – 1868.
- [21] D.J. Gryniewicz, *Structural Additive Theory*, Developments in Mathematics 30, Springer, Cham, 2013.
- [22] D.J. Gryniewicz and F. Hennecart, *A weighted zero-sum problem with quadratic residues*, Uniform Distribution Theory **10** (2015), 69 – 105.
- [23] D.J. Gryniewicz, L.E. Marchan, and O. Ordaz, *A weighted generalization of two theorems of Gao*, Ramanujan J. **28** (2012), 323 – 340.
- [24] F. Halter-Koch, *Arithmetical interpretation of Davenport constants with weights*, Arch. Math. **103** (2014), 125 – 131.
- [25] F. Kainrath, *Factorization in Krull monoids with infinite class group*, Colloq. Math. **80** (1999), 23 – 30.
- [26] L.E. Marchan, O. Ordaz, D. Ramos, and W.A. Schmid, *Inverse results for weighted Harborth constants*, Int. J. Number Theory **12** (2016), 1845 – 1861.
- [27] L.E. Marchan, O. Ordaz, I. Santos, and W.A. Schmid, *Multi-wise and constrained fully weighted Davenport constants and interactions with coding theory*, J. Comb. Theory, Ser. A. **135** (2015), 237 – 267.
- [28] L.E. Marchan, O. Ordaz, and W.A. Schmid, *Remarks on the plus-minus weighted Davenport constant*, Int. J. Number Theory **10** (2014), 1219 – 1239.
- [29] K. Merito, O. Ordaz, and W.A. Schmid, *On the set of minimal distances of the monoid of plus-minus weighted zero-sum sequences and applications to the characterization problem*, manuscript.
- [30] A. Plagne and W.A. Schmid, *On congruence half-factorial Krull monoids with cyclic class group*, Journal of Combinatorial Algebra **3** (2019), 331 – 400.
- [31] D. Robinson, *A Course in the Theory of Groups*, 2nd ed., Springer, 1996.
- [32] W.A. Schmid, *A realization theorem for sets of lengths*, J. Number Theory **129** (2009), 990 – 999.
- [33] D. Windisch, *On the arithmetic of ultraproducts of commutative cancellative monoids*, <https://arxiv.org/abs/2311.16662>.
- [34] Q. Zhong, *Sets of minimal distances and characterizations of class groups of Krull monoids*, Ramanujan J. **45** (2018), 719 – 737.

UNIVERSITY OF GRAZ, NAWI GRAZ, DEPARTMENT OF MATHEMATICS AND SCIENTIFIC COMPUTING, HEINRICHSTRASSE 36,  
8010 GRAZ

Email address: [alfred.geroldinger@uni-graz.at](mailto:alfred.geroldinger@uni-graz.at), [florian.kainrath@uni-graz.at](mailto:florian.kainrath@uni-graz.at)

URL: <https://imsc.uni-graz.at/geroldinger>