ON THE SPECTRAL EDGE OF NON-HERMITIAN RANDOM MATRICES

ANDREW CAMPBELL[†], GIORGIO CIPOLLONI[‡], LÁSZLÓ ERDŐS[†], AND HONG CHANG JI[†]

ABSTRACT. For general non-Hermitian random matrices X and deterministic deformation matrices A, we prove that the local eigenvalue statistics of A + X close to the typical edge points of its spectrum are universal. Furthermore, we show that under natural assumptions on A the spectrum of A + Xdoes not have outliers at a distance larger than the natural fluctuation scale of the eigenvalues. As a consequence, the number of eigenvalues in each component of Spec(A + X) is deterministic.

Contents

1. Introduction	1
1.1. Main results	1
1.2. History and related results	2
1.3. About the proofs	4
1.4. Notation	6
2. Model and Main Results	6
2.1. Universality at the sharp edges	7
2.2. No outliers and cluster-rigidity for $A + X$	9
2.3. Organization of the proof	11
3. Preliminaries and the proofs of Theorems 2.11–2.13	11
3.1. Preliminary results and the local law for $A + X$	11
3.2. Proof of Theorems 2.11–2.13	13
4. Proof of Theorem 2.7	14
4.1. Proof of Theorem 2.7	14
4.2. Deterministic flow	18
5. Proof of Proposition 4.4, the complex case	22
5.1. General framework	23
5.2. Proof of Proposition 5.2	25
5.3. Proof of Lemmas 5.5–5.7	28
6. Proof of Proposition 4.4, the real case	30
6.1. General framework	33
6.2. Large θ regime	34
6.3. Small θ regime	36
7. Proof of Proposition 4.5	37
8. Zig–zag strategy: Proof of Theorem 3.3	38
8.1. Proof of Proposition 8.3	41
8.2. Proof of Proposition 8.4	43
References	48

1. INTRODUCTION

1.1. Main results. Consider a large $N \times N$ deterministic matrix A (called a *deformation matrix*) and a large random matrix X with independent and identically distributed (i.i.d.) centered entries, which

[†]INSTITUTE OF SCIENCE AND TECHNOLOGY AUSTRIA

[‡] Mathematics and Physics departments, Princeton University

 $^{{\}it E-mail\ addresses:\ and rew.campbell@ist.ac.at,\ gc4233@princeton.edu,\ lerdos@ist.ac.at,\ addresses:\ addres$

hongchang.ji@ista.ac.at.

²⁰²⁰ Mathematics Subject Classification. 15B52, 60B20.

Key words and phrases. Ginibre ensemble, full-rank deformation, universality, cluster-rigidity, outliers, local law. [†] Supported by ERC Advanced Grant "RMTBeyond" No. 101020331.

we refer to as an *IID* matrix. It was established in [53] that the eigenvalue density of the *deformed* non-Hermitian random matrix A+X at the edge of its spectrum exhibits either a sharp cutoff behavior, which we refer to as the *sharp* edge, or there may be a small subset of the spectral edge with quadratic decay, which we refer to as the *quadratic* edge. Motivated by the well known analogous principle in Hermitian random matrix theory, our first main result, Theorem 2.7, establishes that the local eigenvalue statistics at sharp edge points are completely governed only by the symmetry class of the matrix, i.e. real or complex. Specifically, under very mild assumptions on A, we prove full universality for the scaling limit of eigenvalue correlation functions localized around a sharp edge point. These limiting statistics are explicitly given by the well known special case when A = 0 and X has i.i.d. real or complex Gaussian entries with variance $\frac{1}{N}$, known as the real or complex *Ginibre ensembles*, respectively. We additionally show that Theorem 2.7 implies universality for matrices with a non-trivial variance profile or even with certain correlation structure.

In the pure IID case, i.e. when A = 0, this universality principle (both in the bulk and at the edge of the spectrum) of local eigenvalue correlation functions was previously established for matrices whose entries have the same first four moments as the entries of a Ginibre matrix (commonly referred to as four moment matching) by Tao and Vu [90]. Full universality at the spectral edge, i.e. with the four moment matching condition of [90] removed, is due to [37]. The special case of complex Ginibre X and normal A was established only very recently by Liu and Zhang at the quadratic edge in [72] and the sharp edge in [73].

In addition to our first main result determining the local eigenvalue fluctuations, we consider the question of whether there can be any eigenvalues of A + X outside the bulk spectrum beyond the scale of these fluctuations. Answering this question in the negative, our second main result Theorem 2.11 is a *no-outlier* theorem: We establish that, with very high probability, A + X has no eigenvalues beyond the natural fluctuation scale at the edge of the spectrum, under a natural assumption on A that excludes outliers of A itself. Previously Bordenave and Capitaine [21, Theorem 1.4] showed that deformed non-Hermitian random matrices have no outlying eigenvalues of distance on a macroscopic scale from the bulk spectrum, whereas we achieve the same result on the optimal scale.



(a) Eigenvalues (blue) around a base point (red) at the edge (black)

(b) No outlier beyond fluctuation (c) Clusters of eigenvalues (blue) scale (red); see (c) for full spectrum. and fluctuation scale (red)

Figure 1. Illustrations of the main results

Our final main result, Corollary 2.12, concerns the components of Spec(A + X). For general A, Spec(A + X) can concentrate in *clusters* on multiple disconnected domains in \mathbb{C} . Corollary 2.12 asserts that each cluster of Spec(A + X) has a deterministic number of eigenvalues with very high probability. We refer to this phenomenon as *cluster-rigidity*, providing the two-dimensional analogue of the *band rigidity* of Hermitian random matrices proved in [7]. See Figure 1 for an illustration of the main results.

1.2. **History and related results.** For Hermitian random matrices the study of local eigenvalue statistics was initiated by the pioneering work of Wigner, and the universality of these statistics regardless of the exact distribution of the matrix entries is now well understood in many general ensembles. These local statistics of the eigenvalues are universal in the sense that they are determined only by the local density of the eigenvalues and the symmetry class of the matrix, i.e. real symmetric, complex Hermitian, etc. For mean field Hermitian random matrix ensembles it was established

in [6] that there are only three possible local shapes (bulk/edge/cusp) of the eigenvalue density. In all three cases very general universality results have been established over the last decade or so: the bulk [4, 63, 68, 69, 84, 28, 47, 49, 51, 52, 65, 79, 89, 61, 3] (i.e. locally flat spectral density), the edge [7, 26, 27, 66, 67, 85, 87, 78, 62] (i.e. square-root decay of the spectral density), and the cusp [35, 48, 57] (i.e. cube root decay of the spectral density).

For non-Hermitian random matrices, studying the spectra is significantly more difficult, and much less is known about the universality of local eigenvalue statistics. Following the intuition from the Hermitian world, the first step to understanding local statistics is to understand the local shape of the eigenvalue densities of such matrices. For an IID matrix X the limiting eigenvalue density is uniform on the unit disk, known as the *circular law*, and hence a point in the spectrum is either in the bulk, or the density has a sharp cutoff at that point. As discussed in Section 1.1, [53] established the non-Hermitian analogue of [6] classifying the potential eigenvalue densities at the edge. In particular [53, Remark 2.10] shows that the set of quadratic points is an analytic sub-manifold of the entire edge, i.e. typically consists of finitely many points. In this paper we thus focus on universality near the ubiquitous sharp edge points for brevity, the case of quadratic edges is deferred to future work.

The spectra of non-Hermitian operators are unstable with respect to small perturbations, and hence the size of the resolvent $(X - z)^{-1}$ of a non-Hermitian matrix is affected not just by the spectrum of X but also its pseudospectrum. Another obstacle is that for non-normal operators the absence of the spectral theorem makes defining any spectral density more difficult. Fortunately, Brown [29] defined what is now known as the *Brown measure* of an operator in a von Neumann algebra, and this measure serves as our non-normal analogue of the spectral density. By the "edge" of A + X we mean truly the boundary of the support of the Brown measure of A+x, where x is a *circular* element in a von Neumann algebra equipped with a faithful normal trace $\langle \cdot \rangle$. This edge was identified in the realm of free probability by [16] and [92] as the set of solutions to $\langle |A - z|^{-2} \rangle = 1$. For the purposes of interpreting our current work all one needs to know is that x is the appropriate infinite dimensional limit of X, and we refer the interested reader to, for example, [75] for a thorough introduction to free probability.

For Ginibre matrices the local eigenvalue statistics at the spectral edge were explicitly identified for the complex case in [55] and for the more complicated real case in [23, 54]; see also [30, 31] and the references therein for a more complete history on eigenvalue statistics of Ginibre ensembles. Beyond Ginibre ensembles, there are two prominent generalizations; the *invariant* ensembles and matrices with independent, but not necessarily Gaussian entries like IID matrices.

The invariant ensembles are random normal matrices with a density proportional to $e^{-2N \operatorname{tr} Q(M)}$ over the space of $N \times N$ complex normal matrices M for some real potential function Q, whose eigenvalue distributions are also known as 2D Coulomb gases. Hedenmalm and Wennman [58] proved that at the spectral edge the eigenvalues exhibit complex Ginibre edge statistics for a family of real analytic potentials Q, see also the related earlier work of [12]. Local eigenvalue statistics at the edge of other non-Hermitian random matrices with Gaussian entries or from invariant ensembles have also been considered in various works including [5, 24, 32, 71, 12]. However, assuming that the matrix has Gaussian entries or unitarily invariant distribution allows for explicit algebraic formulas on the joint density function and techniques that do not generalize to other entry distributions.

In this paper we consider the generalization of independent entries, for which the lack of algebraic formulas and techniques developed for invariant ensembles makes the situation significantly harder. Beyond [90] and [37] little is known about edge universality of non-Hermitian random matrices. Very recently, the problem of bulk universality for matrices with i.i.d. entries was resolved in the breakthrough papers: [74] for the complex case, by Maltsev and Osman, and [77] for the real case, by Osman. Both works use very different methods compared to this manuscript. More precisely, using the multi–resolvent local laws from [39, 44] as an input, they rely on the partial Schur decomposition to access the k-point correlation functions of the eigenvalues. The derivation of the correlation functions uses supersymmetric (SUSY) techniques that heavily depend on the symmetry class. The real case is much more complicated than the complex case as a consequence of the symmetry of the spectrum with respect to the real axis. Neither these multi-resolvent local laws nor the derivation of the k-point correlation functions in the edge regime are known at the moment. We thus follow a simpler strategy which solely relies on the single resolvent local law. Moreover, our proof allows for the real and complex cases to be treated in parallel with the same effort.

Extending the edge universality from standard IID matrices [37] to arbitrarily deformed ones, A + X, is not only a natural generalization, but also allows us to establish edge universality for other natural non-Hermitian matrices with nontrivial variance profile and even correlations. Thus our present result is the non-Hermitian counterpart of the Tracy-Widom edge universality established for deformed Wigner matrices first in [67] and then in [7] in full generality. Here we stress only that the ability to cover such a wide array of cases with Theorem 2.7 is thanks to our very mild assumptions on A. There is a very general idea behind this approach: the deformed model A + X can be used to study very broad classes of random matrix ensembles, including certain correlations, as long as each matrix element has a sizeable variance. In fact, to prove local laws, central limit theorems for linear eigenvalue statistics or local spectral universality for general matrix ensembles it is basically sufficient to prove the same result for $A + X^{\text{Gin}}$ and then use some version of the *Green function comparison* (GFT). See Section 2.1.1 for further details of this general idea for the current edge universality question.

As a side note we mention that the extremal eigenvalue statistics also exhibit a universality phenomenon; for Ginibre ensembles they were proven to be Gumbel by [80, 81] and extended to full universality for IID matrices X in the series of papers [40, 41, 42, 43].

As far as outliers are concerned, in Section 1.1 we already mentioned [21] which proved that if A does not have any outliers in its spectrum, then A + X also does not have any outliers on a macroscopic scale. This result was an extension of a result of Tao [86, Theorem 1.4] which gave the same no outlier result for the special case of A = 0. Our result Theorem 2.11 improves [21, Theorem 1.4] from an order one result to the optimal scale given by the natural local fluctuations of the eigenvalues. With [21] already describing the existence of outliers inherited from outliers of A, our result Theorem 2.11 provides the no-outlier counterpart on the optimal scale and resolves the question of when A + X does or does not have outliers in the spectrum beyond the natural scale. In an independent work posted on arXiv on the same day as the first version of this manuscript, Alt and Krüger [11] showed that A + X does not have outliers on a macroscopic scale when X has a variance profile and A is diagonal. A much stronger result, showing the non existence of outliers on the optimal fluctuation scale, follows from the techniques developed in this paper if X is *flat* (see Remark 2.15 for a more detailed discussion). We point out that [11] also gives an explicit description of the limiting eigenvalue density by solving the Matrix Dyson Equation (MDE) associated to these models. We also mention that there is a very large literature on outliers (when they exist) that we do not discuss here; we refer the interested readers to [13, 17, 18, 20, 15, 19, 14, 45, 76] and references therein.

1.3. About the proofs. In this section we provide a summary of the proofs and highlight the methodological novelties. As is often the case in the study of non-Hermitian random matrices (with a few notable exceptions such as [74, 77, 22]) we use Girko's Hermitization [56] formula to express the spectral statistics of A + X in terms of traces of the resolvent matrix G of the Hermitization of A + X, which is defined by

(1.1)
$$H^{z} := \begin{pmatrix} 0 & A + X - z \\ (A + X - z)^{*} & 0 \end{pmatrix}, \quad z \in \mathbb{C}.$$

The advantage of considering the Hermitization of a matrix is that it can be related to linear statistics of the non–Hermitian matrix via Girko's celebrated Hermitization formula:

(1.2)
$$\int_{\mathbb{C}} F(z) \mathrm{d}\rho_{A+X}(z) = -\frac{1}{4\pi N} \int_{\mathbb{C}} \Delta F(z) \log |\det H^z| \, \mathrm{d}^2 z,$$

where ρ_{A+X} is the normalized empirical eigenvalue measure of A + X and F is a nice test function. From (1.2), simple functional calculus allows one to express $\int F d\rho_{A+X}$ in terms of the resolvent $G^z = (H^z - i\eta)^{-1}$, and this observation by Girko has become the backbone of non-Hermitian random matrix theory. We then use this formula in conjunction with a *local law*, i.e. an approximation of G^z by the deterministic solution to a *matrix Dyson equation* (MDE). With Girko's formula and this deterministic approximation available we can then compare the eigenvalue statistics of A + X to the known statistics of a Ginibre matrix X^{Gin} through a Green function comparison (GFT).

To prove the optimal local law for the Hermitization H^z we apply a bootstrapping argument, which we refer to as the *iterated zig-zag* method. This is a repeated combination of the characteristic flow method (zig) followed by a GFT (zag). This zig-zag argument has previously been used to prove multi-resolvent local laws in [39, 43, 33, 34, 50]. Crucially these previous works always had the single resolvent local law available as an a priori input, hence they could complete the process in only one or two zig-zag steps. One of the novelties of our paper is the first proof of a single resolvent local law from scratch using the zig-zag approach. However, lacking the a priori input, we need to use a large (but finite) number of zig-zags in the process. We remark the zig step has been previously used for a single resolvent local law in [2, 25, 1, 60]. Our local law covers both possible spectral edges of A + X, and hence so do our no-outlier and cluster-rigidity results. We mention that a clever self-consistent way of proving the GFT part of the isotropic local law has already appeared in [64], where it was used to handle the bulk and edge regimes for deformed Wigner and sample covariance matrices. However, as an input for the GFT argument, this paper still relied on an isotropic local law for the analogous Gaussian model exploiting its unitary invariance property. We use the convenient characteristic flow method, that was not available at the time of [64], to prove local laws directly, using only the trivially available global laws as the sole input. Moreover, the zig-zag method is very robust; here we apply it in the most critical cusp regime without any additional complication.

We next discuss our proof of universality. The comparison between A + X and X^{Gin} can be naturally broken into two steps by inserting the intermediate ensemble $A + X^{\text{Gin}}$, both of which are carried out via GFT. We compare A + X and $A + X^{\text{Gin}}$ by evolving X along an Ornstein–Uhlenbeck flow, and the comparison between $A + X^{\text{Gin}}$ and X^{Gin} follows a continuous flow of the deformation. We focus below on the latter step since the former step is relatively standard. As this step is sensitive to the symmetry class, we first explain the complex case and then handle the additional difficulties in the real case.

In the complex case, to compare $A + X^{\text{Gin}}$ and X^{Gin} , we evolve A - z continuously along a carefully chosen curve $A_t - z_t$ for $t \in [0,1]$ such that this z_t remains a sharp edge of $A_t + X^{\text{Gin}}$ and that $A_0 = A$ and $A_1 = 0$. Along this curve, we show that the time derivatives of the eigenvalue and singular value statistics are sufficiently small. The key technical challenge in this comparison is that the naive estimates of the time derivatives available from the local law are not sufficient, and we need to identify non-trivial cancellations within the derivatives. These cancellations are revealed by introducing and properly matching four time-dependent real parameters; one in order to keep z_t at a sharp edge; another one to keep the (1D) local density of singular values unchanged; and two to keep the (2D) local density of eigenvalues constant. Our approach is reminiscent of Lee and Schnelli's [67] work on deformed Wigner matrices, where they tracked cancellations through two analogous parameters from the evolutions of spectral edge and (1D) eigenvalue density. However, due the non-Hermitian nature of the problem, we have to introduce two extra parameters to get such strong cancellations. In fact, what we show with careful adjustments of these four parameters is that, along any C^1 curve of deformations starting from $A - z_0$ for which the origin remains a sharp edge, the local eigenvalue statistics at the origin remain unchanged. Exploiting the freedom to compare eigenvalue statistics along a fully general curve is a methodological novelty, which allows us to handle the more delicate "inner" edges of the spectrum which do not appear in Ginibre matrices. See Figure 2 in Section 5 for an illustration of how we can use this flexibility.

The real case has an additional, fundamental difficulty due to the symmetry of the spectrum about the real axis. For a mean-field real random matrix, this symmetry results in a one-parameter family of universality classes around a point z, depending on the parameter $\sqrt{N} | \operatorname{Im} z|$. Our proof must accommodate this complication so that the comparison between $A + X^{\operatorname{Gin}}$ and X^{Gin} must be performed along a carefully designed flow $A_t - z_t$ which keeps the universality class. More precisely, the real case requires to handle a few additional terms that are not present in the complex case. These terms stem from correlations between eigenvalues around z and \overline{z} . The key difficulty is to show that these quantities do not affect the evolution of the local statistics along the flow. We divide the proof into two cases: $|\operatorname{Im} z| \leq N^{-1/2}$ and $|\operatorname{Im} z| \gg N^{-1/2}$. In the former case we show that these additional terms have their own cancellations using the construction that A_t keeps the exact universality class. In the latter case the extra terms are small by themselves, as one can expect from the decay of correlation of eigenvalues further apart than the natural scale $N^{-1/2}$. In particular we show that the local statistics for the real ensemble at a distance much larger than $N^{-1/2}$ away from the real axis coincide with the statistics for the complex ensemble.

In both the real and complex cases these eigenvalue comparisons are done in parallel with a comparison on the lower bound of the least singular value of A + X. Control of the least singular value of non-Hermitian matrices is both essential to applying Girko's Hermitization and an interesting question in its own right. We refer the interested reader to the recent survey [91] and the references therein. We focus only on works which are most relevant to eigenvalue universality. For Ginibre matrices, [36, 38] gave an optimal lower tail on the least singular value of $X^{\text{Gin}} - z$ for z at the spectral edge, i.e. |z| = 1. The optimal tail bound in [36] for shifted Ginibre matrices was used in the proof of full edge universality for IID matrices in [37]. Recently, Shcherbina and Shcherbina [82] considered the least singular value tail bound for deformed complex Ginibre matrices $A + X^{\text{Gin}} - z$, however because their result is only for complex matrices and not uniform in the optimal scale of the least singular value we instead use a GFT to the known results of [36].

For our no-outlier and cluster-rigidity results we first show an improvement of the local law for z outside $\text{Spec}(A + \mathbf{x})$. Then, for the proof of cluster-rigidity we consider the interpolation $A + \sqrt{t}X$, for which we demonstrate that the number of eigenvalues of $A + \sqrt{t}X$ in each cluster of A + X is fixed over t with very high probability.

1.4. Notation. In this section we introduce some common notations used throughout the paper. We write \mathbb{C}_+ for the upper half plane $\mathbb{C}_+ := \{z \in \mathbb{C} : \operatorname{Im} z > 0\}$ and \mathbb{S}^1 for the unit circle $\mathbb{S}^1 := \{z \in \mathbb{C} : |z| = 1\}$. For a positive integer $k \in \mathbb{N}$ we write $[k] := \{1, \ldots, k\}$. We denote the Lebesgue measure on $\mathbb{C} \cong \mathbb{R}^2$ by d^2z , and that on $\mathbb{C}^k \cong \mathbb{R}^{2k}$ by $d^{2k}z$. We define the $2N \times 2N$ block constant matrices

(1.3)
$$E_1 := \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad E_2 := \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}.$$

For a generic complex matrix B, we write its real, imaginary, Hermitian, and skew-Hermitian parts as

(1.4)
$$\Re[B] := \frac{B + \overline{B}}{2}, \quad \Im[B] := \frac{B - \overline{B}}{2i}, \quad \operatorname{Re}[B] := \frac{B + B^*}{2}, \quad \operatorname{Im}[B] := \frac{B - B^*}{2i},$$

respectively. For a $d \times d$ (in most cases d = N or 2N) matrix A we use $\langle A \rangle = d^{-1} \operatorname{Tr} A$ to denote the normalized trace. We use $u^* y$ to denote the standard Euclidean inner product of two vectors $u, y \in \mathbb{C}^d$.

We write c and C for generic positive constants (independent of N) whose precise values may vary by lines. For positive quantities f, g we write $f \leq g$ and $f \sim g$ if $f \leq Cg$ and $cg \leq f \leq Cg$, respectively. We say that an N-dependent event holds with very high probability if for any D > 0, the event holds with probability at least $1 - N^{-D}$ for any $N \geq N_0$, where the threshold N_0 depends on D. If this event depends on additional parameters, we say that the event holds with very high probability uniformly over a given set of parameters if a common threshold N_0 can be chosen uniformly for all parameters in this set. Moreover, for two random variables X and Y we say X is stochastically dominated by Y, which we denote by $X \prec Y$, if for every $\xi > 0$ the event $\{|X| \leq N^{\xi}|Y|\}$ holds with very high probability.

2. Model and Main Results

We consider $N \times N$ non–Hermitian random matrices with independent, identically distributed (IID) entries satisfying the following assumption.

Definition 2.1. An N-dependent sequence of complex (resp. real) random matrices $X \equiv X^{(N)} \in \mathbb{C}^{N \times N}$ (resp. $\mathbb{R}^{N \times N}$) is called an IID random matrix if it satisfies the following:

- The entries of X are independent.
- $\mathbb{E}[X] = 0$, $\mathbb{E}[|X_{ij}|^2] = 1/N$. Additionally, $\mathbb{E}[X_{ij}^2] = 0$ in the complex case.
- The entries of $\sqrt{N}X$ have finite moments, i.e. for each $p \in \mathbb{N}$ there exists a constant $c_p > 0$ with

(2.1)
$$\sup_{N \in \mathbb{N}} \max_{i,j} \mathbb{E} |\sqrt{N}X_{ij}|^p \le c_p.$$

A complex (resp. real) IID matrix with Gaussian entries is referred to as complex (resp. real) Ginibre matrix and denoted by $X^{\operatorname{Gin}(\mathbb{C})}$ (resp. $X^{\operatorname{Gin}(\mathbb{R})}$).

Definition 2.2. Let \mathbb{F} be either \mathbb{R} or \mathbb{C} , $X \in \mathbb{F}^{N \times N}$ be an IID matrix, and $A \equiv A^{(N)} \in \mathbb{F}^{N \times N}$ be a general sequence of deterministic matrices such that

(2.2)
$$\sup_{N \in \mathbb{N}} \|A\| \le C_0$$

for a constant $C_0 > 0$. We define the $(N \times N)$ deformed IID matrix as

It is expected that the empirical eigenvalue distribution of A + X concentrates around a deterministic (*N*-dependent) measure denoted by $\rho_{A+\boldsymbol{x}}$ for large *N* (see e.g. [83] for *X* having Gaussian entries). More precisely, $\rho_{A+\boldsymbol{x}}$ is the *Brown measure* of an infinite-dimensional operator $A + \boldsymbol{x}$. We now introduce two definitions to properly define $\rho_{A+\boldsymbol{x}}$.

Definition 2.3 (Brown measure). Let $(\mathcal{M}, \langle \cdot \rangle_{\mathcal{M}})$ be a W^* -probability space, i.e., let \mathcal{M} be a von-Neumann algebra with a faithful, normal, tracial state $\langle \cdot \rangle_{\mathcal{M}}$. For an element $\mathbf{a} \in \mathcal{M}$, we define its Brown measure to be the distributional Laplacian

(2.4)
$$\rho_{\boldsymbol{a}} := \frac{1}{2\pi} \Delta \langle \log |\boldsymbol{a} - \cdot| \rangle_{\mathcal{M}}.$$

For a (random) matrix $B \in \mathbb{C}^{N \times N}$, we consistently write ρ_B for its empirical eigenvalue distribution.

Definition 2.4. Let $(\mathbf{x}_{ij})_{i,j\in[\![N]\!]} \in \mathcal{M}$ be a collection of *-free circular elements¹, and define $\mathbf{x} = (\mathbf{x}_{ij}) \in \mathcal{M}^{N \times N}$. We canonically embed $\mathbb{C}^{N \times N}$ in $\mathcal{M}^{N \times N}$, so that $A + \mathbf{x}$ becomes a sum of two elements of $\mathcal{M}^{N \times N}$. Finally, we consistently extend $\langle \cdot \rangle_{\mathcal{M}} : \mathcal{M}^{N \times N} \to \mathbb{C}^{N \times N}$ to be the partial trace, i.e.

(2.5)
$$\langle \boldsymbol{y} \rangle_{\mathcal{M}} = (\langle \boldsymbol{y}_{ij} \rangle_{\mathcal{M}})_{1 \leq i,j \leq N} \in \mathbb{C}^{N \times N}, \qquad \boldsymbol{y} = (\boldsymbol{y}_{ij})_{1 \leq i,j \leq N} \in \mathcal{M}^{N \times N}.$$

Notice that $(\mathcal{M}^{N \times N}, \langle \langle \cdot \rangle_{\mathcal{M}} \rangle)$ is a W^* -probability space, where the outer bracket $\langle \cdot \rangle$ is the normalized trace on $\mathbb{C}^{N \times N}$.

Remark 2.5. With a slight abuse of notations, when we work with (square) matrices of dimension 2N instead of N, we naturally also denote by $\langle \cdot \rangle_{\mathcal{M}}$ the partial trace $\mathcal{M}^{2N \times 2N} \to \mathbb{C}^{2N \times 2N}$ applied entrywise as in (2.5). Likewise, we often canonically embed $\mathbb{C}^{2N \times 2N}$ in $\mathcal{M}^{2N \times 2N}$. However, we emphasize that we never embed a random matrix in $\mathcal{M}^{d \times d}$, e.g. we never consider X or A + X as an element of $\mathcal{M}^{N \times N}$.

In the bigger W^* -probability space $(\mathcal{M}^{N \times N}, \langle \langle \cdot \rangle_{\mathcal{M}} \rangle)$, the operator \boldsymbol{x} in Definition 2.4 is a circular element and A and \boldsymbol{x} are *-free; to see this, approximate (in *-moments) \boldsymbol{x}_{ij} with independent Ginibre matrices $X^{(ij)}$ of size $N' \gg 1$, so that $X' = (X^{(ij)})_{ij} \in \mathbb{C}^{(NN' \times NN')}$ is a good approximation of \boldsymbol{x} . Then it immediately follows that \boldsymbol{x} is a circular element, and also that $A \otimes I_{N'}$ and X' are asymptotically *-free as $N' \to \infty$ so that A and \boldsymbol{x} are *-free. With Definition 2.3, we define $\rho_{A+\boldsymbol{x}}$ to be the Brown measure of $A + \boldsymbol{x}$ in $(\mathcal{M}^{N \times N}, \langle \langle \cdot \rangle_{\mathcal{M}} \rangle)$.

We now recall the classification of the edge behaviors of ρ_{A+x} from [53]. We say that a point z_0 on the boundary of the support of ρ_{A+x} belongs to the *regular edge* of ρ_{A+x} if $||(A-z_0)^{-1}|| \leq C$ for some (large) constant C > 0. Regular edge points $z_0 \in \mathbb{C}$ are characterized by the condition that $\langle |A-z_0|^{-2} \rangle = 1$. Then [53, Theorem 2.9] asserts that ρ_{A+x} has a density² only admitting one of the two possible asymptotic behaviors around any regular edge z_0 :

i) (Sharp edge) As $z \to z_0$ we have

$$\rho_{A+\boldsymbol{x}}(z) \sim \mathbb{1}(z \in \operatorname{supp}(\rho_{A+\boldsymbol{x}})).$$

ii) (Quadratic edge) As $z \to z_0$ we have

$$\rho_{A+\boldsymbol{x}}(z) \sim |z-z_0|^2 \mathbb{1}(z \in \operatorname{supp}(\rho_{A+\boldsymbol{x}}))$$

In the next two subsections we present our main results. The first main result concerns the universality of the local statistics of the eigenvalues of A + X (see Section 2.1 below) at a sharp edge point. Notice that $d\rho_{\boldsymbol{x}}(z) = \pi^{-1} \mathbb{1}(|z| \leq 1) d^2 z$ by the circular law, hence the sharp edge prevails in $\rho_{\boldsymbol{x}}$. The universality around quadratic edges is deferred to future work. One of our fundamental inputs to prove universality is an a priori control of the spectrum down to any mesoscopic scale, i.e. on any scale that is slightly bigger than the local fluctuations scale of the eigenvalues. As a byproduct of this control on mesoscopic scales, the second main result of this paper proves that with very high probability there are no outliers in the spectrum of A + X at a distance from the edge of the support of $\rho_{A+\boldsymbol{x}}$ which is bigger than the natural local fluctuation scale of the eigenvalues (see Section 2.2 below).

2.1. Universality at the sharp edges. In [53], it is proved that (the density of) ρ_{A+x} has a sharp jump discontinuity around a point $z_0 \in \mathbb{C}$ (called the *base-point*) under the following assumption:

Assumption 2.6. We assume that $z_0 \in \mathbb{C}$ satisfies the following:

(i)
$$\langle |A - z_0|^{-2} \rangle = 1.$$

(ii) for a constant $C_1 > 0$, $z_0 \in \mathcal{D}_{C_1}$ where

(2.6)
$$\mathcal{D}_C := \left\{ z \in \mathbb{C} : \left\| \frac{1}{A-z} \right\| < C, \quad (1+|z|)^3 \left| \left\langle \frac{1}{|A-z|^4} (A-z) \right\rangle \right| > \frac{1}{C} \right\}$$

The last assumption depends on the symmetry class. Let $C_2 > C_1$ be another constant. When $\mathbb{F} = \mathbb{C}$, we assume that

(iiiC) z_0 is in the unique unbounded component of \mathcal{D}_{C_2} .

When $\mathbb{F} = \mathbb{R}$, we fix a small constant $\mathfrak{c} > 0$ and assume the following:

(iii \mathbb{R}) If $|\operatorname{Im} z_0| > N^{-1/2+\mathfrak{c}}$, z_0 is in one of the two unbounded components of $\mathcal{D}_{C_2} \setminus \mathbb{R}$.

(iii \mathbb{R}') If $|\operatorname{Im} z_0| < N^{-1/2+\mathfrak{c}}$, $\operatorname{Re} z_0$ is in the unique unbounded component of $\mathcal{D}_{C_2} \cap \mathbb{R}$.

¹More precisely, $\boldsymbol{x}_{ij} = (\boldsymbol{s}_{ij,1} + i\boldsymbol{s}_{ij,2})/\sqrt{2}$ where $(\boldsymbol{s}_{ij,1}, \boldsymbol{s}_{ij,2})$ is a pair of free semi-circular elements in \mathcal{M} .

²With a slight abuse of notation, we write ρ_{A+x} for both the Brown measure and its density.

The equality $\langle |A - z_0|^{-2} \rangle = 1$ guarantees that the base point z_0 is at the spectral edge of A + X. The first inequality in (2.6) ensures that the associated Dyson equation (see Lemma 3.2 below) is stable around z_0 , and the second excludes the quadratic edges. Assumptions (iii) guarantee that there exists a C^1 path from z_0 to ∞ that purely consists of sharp edge points of $A + \sqrt{t}X$ for some t > 0. In the real case the path has an additional constraint from the symmetry class, hence we divide (iii \mathbb{R}) and (iii \mathbb{R}').

Our first main theorem asserts that the local eigenvalue statistics of A + X around a base point z_0 at a sharp edge is universal in the $N \to \infty$ limit (after proper rescaling). In particular, we show that the corresponding k-point correlation functions coincide with those of Ginibre ensembles in the same symmetry class. To be precise, for a given $z \in S^1$, our reference functions $p_{z,k}^{\operatorname{Gin}(\mathbb{F})}$ are defined by the limiting k-point correlation functions of eigenvalues of $\sqrt{N}(z^{-1}X^{\operatorname{Gin}(\mathbb{F})} - 1)$ near the origin. These are well-known functions; in the complex case they are independent of $z \in S^1$ and given by (see [55])

(2.7)

$$p_{z,k}^{\operatorname{Gin}(\mathbb{C})}(w_1, \dots, w_k) = \det \left[K^{\operatorname{Gin}(\mathbb{C})}(w_i, w_j) \right]_{1 \le i,j \le k},$$

$$K^{\operatorname{Gin}(\mathbb{C})}(w_1, w_2) = \frac{1}{2\pi} \left[1 + \operatorname{erf} \left(-\sqrt{2}(\overline{w_2} + w_1) \right) \right] e^{-\frac{|w_1|^2}{2} - \frac{|w_2|^2}{2} + w_1 \overline{w_2}}.$$

Explicit formulas for $p_{z,k}^{\operatorname{Gin}(\mathbb{R})}$ are known also in the real case [23], but they are much more involved as a consequence of the special role played by the real axis.

The proper rescaling is governed by a single scaling parameter defined as

(2.8)
$$\gamma_0 := -I_4^{-1/2} I_3$$

where we set

(2.9)
$$I_3 := \left\langle \frac{1}{|A - z_0|^4} (A - z_0)^* \right\rangle, \qquad I_4 := \left\langle \frac{1}{|A - z_0|^4} \right\rangle.$$

The parameter $\gamma_0 \in \mathbb{C}$ is chosen so that the density of the Brown measure of $\gamma_0(A + \mathbf{x} - z_0)$ matches that of $(\mathbf{x} - 1)$ near their sharp edge point at the origin (see Lemma 3.1 below for more details).

Theorem 2.7 (Sharp edge universality). Let $A + X \in \mathbb{F}^{N \times N}$ be the deformed IID matrix defined in Definition 2.2, and let the base point $z_0 \in \mathbb{C}$ satisfy Assumption 2.6. Fix any $k \in \mathbb{N}$, and let $p_k^{(N)}$ be the k-point correlation function of the eigenvalues of $\sqrt{N}\gamma_0(A + X - z_0)$ near the origin. Then, for any compactly supported smooth function $F : \mathbb{C}^k \to \mathbb{C}$ we have

(2.10)
$$\int_{\mathbb{C}^k} F(\boldsymbol{w}) \left[p_k^{(N)}(\boldsymbol{w}) - p_{z_1,k}^{\operatorname{Gin}(\mathbb{F})}(\boldsymbol{w}) \right] \mathrm{d}^2 \boldsymbol{w} = O\left(N^{-c} \right),$$

for some small c > 0, where $z_1 \in \mathbb{S}^1$ is chosen³ such that $\operatorname{Im} z_1 = (|\gamma_0| \wedge 1/|\operatorname{Im} z_0|) \operatorname{Im} z_0$. The implicit constant and c > 0 in (2.10) depend only on k, F, and the constants from Definitions 2.1 – 2.2 and Assumption 2.6.

The same edge universality for A + X was proven in [73] for the very special case where X is a Ginibre matrix and A is a normal matrix with finitely supported empirical spectral measure and for A = 0 and IID X in [36].

While Theorem 2.7 is formulated for deterministic A, it is clear that we may assume that A is a random matrix, independent of X, properly chosen so that it satisfies Assumption 2.6 with sufficiently high probability. This simple idea allows us to conclude edge universality for many other natural random matrix ensembles as a direct consequence of Theorem 2.7; we give some examples below. We point out that in all these examples it is fundamental that Theorem 2.7 allows for very general A's, which in particular may not be normal matrices. We remark that the normality of A was a fundamental assumption in the previous works for special $A + X^{\text{Gin}}$ -type matrices in [72, 73].

2.1.1. *Examples.* We now present a few examples of relevant random matrix ensembles for which we can easily prove edge universality by a simple application of our general Theorem 2.7.

We consider an $N \times N$ random matrix X with independent centered entries X_{ij} with a general flat variance profile such that

(2.11)
$$\mathbb{E}|X_{ij}|^2 \coloneqq \frac{S_{ij}}{N}, \qquad \frac{c}{N} \le S_{ij} \le \frac{C}{N},$$

for two N-independent constants c, C > 0. By [8, Theorem 2.5] it follows that the limiting eigenvalue density of such X is supported on a disk of radius $\sqrt{\rho(S)}$, with $\rho(S)$ denoting the spectral radius of the

³This choice of z_1 is relevant only when $\mathbb{F} = \mathbb{R}$, since $p_{z,k}^{\operatorname{Gin}(\mathbb{C})}$ is independent of $z \in \mathbb{S}^1$.

variance matrix S. By the four moment matching comparison theorems [90, Theorems 2 and 12] we can then write⁴

(2.12)
$$X \stackrel{d}{=} X + \epsilon X^{\text{Gin}}, \quad \epsilon \sim 1,$$

with \widetilde{X} IID matrix being independent of X^{Gin} . Whenever \widehat{z}_0 is on the spectral edge of X, i.e. $|\widehat{z}_0| = \sqrt{\rho(S)}$, the pair $(A, z_0) := (\epsilon^{-1} \widetilde{X}, \epsilon^{-1} \widehat{z}_0)$ satisfies Assumption 2.6 with very high probability. To see this, we only need to recall two facts:

- (i) By [47, Theorem 2.1], for any $z \in \mathbb{C}$, on a macroscopic scale the empirical eigenvalue density $\rho_{|\widetilde{X}-z|}$ concentrates around a deterministic probability measure μ_z on $[0, \infty)$ with rate $O(N^{-1})$ and by [47, Corollary 2.3] there is no outlying eigenvalue of $|\widetilde{X} z|$ beyond a distance $N^{-\delta}$, $\delta > 0$, from the support of μ_z .
- (ii) By [8, Proposition 3.2(iii)] and from the easy fact that ρ(S̃) < ρ(S) − cε², where S̃ is the variance profile of X̃, we conclude that the solution of the MDE for the Hermitization of X̃ − ẑ₀ vanishes at zero, moreover it is of order O(η) along imaginary axis. Using the implication (vi) ⇒ (v) of [6, Lemma D.1], we thus conclude that μ_{2̂0} is supported away from zero, i.e., inf supp μ_{2̂0} ~ 1.

These two facts together imply that $|\langle |\epsilon^{-1}(\widetilde{X}-\widehat{z}_0)|^{-2}\rangle -1| \prec N^{-1}$ and $||(\widetilde{X}-\widehat{z}_0)^{-1}|| = O(1)$ with very high probability. Likewise, applying [8, Section 3] to the MDE associated to \widetilde{X} proves that $\{|z| \geq \epsilon^{-1}\sqrt{\rho(S)}\}$ is contained in the domain \mathcal{D}_C from Assumption 2.6 with very high probability, confirming⁵ Assumption 2.6 for $(A, z_0) = (\epsilon^{-1}\widetilde{X}, \epsilon^{-1}\widehat{z}_0)$ with very high probability. Thus, conditioning on \widetilde{X} and working in the probability space of X^{Gin} , we can directly apply Theorem 2.7 to show that the local statistics of Xaround \widehat{z}_0 are universal and that they coincide with the corresponding real/complex Ginibre statistics.

In fact, a very similar argument also shows edge universality for a large class of correlated matrices (e.g. the non–Hermitian counterpart of [48, Assumptions (A)–(F)], as considered in [10]), and for matrices of the form A + X, with X as in (2.11) and A a deterministic matrix satisfying an assumption similar to Assumption 2.6 (see also Remark 2.15 for a related discussion). We omit the details for brevity. We conclude this section pointing out that in particular this implies edge universality for the so called elliptic ensemble, i.e. when $\mathbb{E}X_{ij}X_{ji} = \tau/N$, with $\tau \in [0, 1)$ (see [70] for the special case when the X_{ij} are Gaussians).

2.2. No outliers and cluster-rigidity for A + X. Consider a deterministic matrix A satisfying the following assumption.

Assumption 2.8. There exists a constant $C_3 > 0$ such that for all $z \in \partial \operatorname{supp} \rho_{A+x}$ we have

(2.13)
$$\left\|\frac{1}{A-z}\right\| \le C_3.$$

Remark 2.9. Unlike in Assumption 2.6, we formulated a global assumption on A rather than only at a single point, as we want to exclude the presence of outliers in the whole complement of the support of ρ_{A+x} ; note that now we allow quadratic edges. However, we point out that we assumed the bound (2.13) for all points on the edge of the support of ρ_{A+x} only because in Theorem 2.11 below we exclude outliers globally in the complement of the spectrum. If we are interested in excluding outliers only locally around an edge point z_0 , then it is enough to assume that (2.13) holds for $z = z_0$.

We now define the typical fluctuation scale of the eigenvalues of A + X:

Definition 2.10 (Fluctuation scale). Fix a point $z \in \mathbb{C}$, we say that $\sigma_f(z) = \sigma_f(z, N)$ is the fluctuation scale of the eigenvalues of A + X around the point z if

(2.14)
$$\int_{|w-z| \le \sigma_{\mathbf{f}}(z)} \rho_{A+\boldsymbol{x}}(w) \,\mathrm{d}^2 w = \frac{1}{N}$$

Note that for z's in the bulk of the spectrum or close to sharp edges we have $\sigma_f(z, N) \sim N^{-1/2}$, while for z's close to a quadratic edge $\sigma_f(z, N) \sim N^{-1/4}$. Fix any $\epsilon > 0$, and define the deterministic set

(2.15)
$$\operatorname{Spec}_{\epsilon}(A + \boldsymbol{x}) := \bigcup_{z \in \operatorname{supp}(\rho_{A + \boldsymbol{x}})} D\left(z, N^{\epsilon} \sigma_{\mathrm{f}}(z)\right).$$

⁴We point out that if the entries of X are supported on only two points (e.g. X_{ij} are Bernoulli random variables) then we cannot directly apply [90, Theorem 12]. Instead, we first show that the local statistics of such X are (asymptotically) the same as the ones of another \hat{X} such that \hat{X}_{ij} are independent Gaussians with the same variance profile of X (see Section 7 for the details of this argument). Then we perform the decomposition (2.12) for \hat{X} .

⁵Assumption 2.6 (i) holds with a random error of size $\prec N^{-1}$, which does not affect the local statistics.

Here D(z, r) denotes the open disk of radius r centered at z.

Theorem 2.11 (No outliers). Let A + X be a deformed IID matrix as in Definition 2.2, and suppose that A satisfies Assumption 2.8. Then, for any small $\epsilon > 0$, we have

(2.16)
$$\operatorname{Spec}(A+X) \subset \operatorname{Spec}_{\epsilon}(A+x),$$

with very high probability uniformly in A and X satisfying (2.1), (2.2) and (2.13).

We also prove that the number of eigenvalues in each component (which we often also refer to as "cluster") of the support of ρ_{A+x} is deterministic.

Corollary 2.12 (Cluster-rigidity). Under the assumptions of Theorem 2.11, the following holds. Fix any small $\epsilon, \delta > 0$, and let C be a union of (one or several) components of $\text{Spec}_{\epsilon}(A + \mathbf{x})$ such that $\text{dist}(\mathcal{C}, \text{Spec}_{\epsilon}(A + \mathbf{x}) \setminus \mathcal{C}) \geq \delta$. Then, counting the eigenvalues with multiplicity, we have

(2.17)
$$\left|\operatorname{Spec}(A+X) \cap \mathcal{C}\right| = \left|\operatorname{Spec}(A) \cap \mathcal{C}\right|$$

with very high probability uniformly over A and X satisfying (2.1), (2.2) and (2.13). In particular, the number of eigenvalues in C is a deterministic integer with very high probability.

Both of these results are consequences of the following very general local law close to the edge of the spectrum of A + X. Fix $a \in [0, 1/2]$, consider a smooth compactly supported function $g : \mathbb{C} \to \mathbb{C}$, and define

$$g_{z_0,a}(z) := g(N^a(z-z_0))$$

Denote the eigenvalues of A + X by $\sigma_1, \ldots, \sigma_N$.

Theorem 2.13 (Local law at regular edges). Let A + X be a deformed IID matrix as in Definition 2.2. Fix any $C_4 > 0$, and fix z_0 on the edge of the support of ρ_{A+x} such that $||(A - z_0)^{-1}|| \leq C_4$. Then, for any $\xi > 0$ and $a \in [0, 1/2)$, we have

(2.18)
$$\left| \frac{1}{N} \sum_{i=1}^{N} g_{z_0,a}(\sigma_i) - \int_{\mathbb{C}} g_{z_0,a}(z) \rho_{A+\boldsymbol{x}}(z) \, \mathrm{d}^2 z \right| \le \frac{N^{\xi}}{N^{1-2a}}$$

with very high probability uniformly in A, X, and z_0 satisfying (2.1), (2.2), and $||(A-z_0)^{-1}|| \leq C_4$.

Note that the error term in (2.18) is smaller than leading deterministic term only on mesoscopic scales around the base point z_0 , i.e. when $N^{-a} \gg \sigma_f(z_0)$. In particular, even when z_0 is a quadratic edge, (2.18) remains effective for all mesoscopic scales $a \in [0, 1/4)$. We also point out that to prove (2.18) we do not need Assumption 2.8 for every point at the edge of the support of ρ_{A+x} , but it is enough to assume that (2.13) holds for the base point z_0 . The same result for z_0 in the bulk of the spectrum of A + X follows by the averaged local law in [44, Theorem 2.6]. If z_0 is in an intermediate regime (between bulk and edge) or is outside of the spectrum we can prove a similar local law, but we do not present it here for brevity.

Remark 2.14. Arguing similarly to Section 2.1.1 we immediately conclude that Theorem 2.11, Corollary 2.12, and (2.18) hold for matrices X with a variance profile as in (2.11) or having certain correlations (see e.g. Section 2.1.1) with a deformation A satisfying an assumption similar to Assumption 2.8. In particular, when A = 0 and the entries are independent, combining Theorem 2.11 and (2.18) we immediately conclude

(2.19)
$$\left|\rho(X) - \sqrt{\rho(S)}\right| \le \frac{N^{\xi}}{\sqrt{N}}.$$

with very high probability. Here $\rho(X)$ denotes the spectral radius. When X is centered and has some correlations as in [10] we can also prove⁶ the bound (2.19), but with $\rho(S)$ being replaced with $\rho(\mathscr{S})$, where \mathscr{S} is the covariance operator of X. This extends [9, Theorem 2.1] to include certain correlated matrices as well.

Remark 2.15. Simultaneously with the current paper, an independent work by Alt and Krüger [11] considered the ϵ -pseudospectrum of A + X for some N-independent $\epsilon \sim 1$, where X has a flat variance

 $^{^{6}}$ In these models, by [10, Theorem 2.5], the spectrum consists of one connected component, the limiting density is rotationally symmetric, and there are no critical edges.

profile⁷ as in (2.11) and A is a diagonal matrix. Their main result [11, Theorem 2.4] roughly reads: for each fixed $\epsilon, \delta > 0$, it holds in the large N limit that

(2.20)
$$\Psi \operatorname{Spec}_{\epsilon}(A+X) \subset \Psi \operatorname{Spec}_{\epsilon}(A+x) \subset \Psi \operatorname{Spec}_{\epsilon+\delta}(A+X),$$

where \boldsymbol{x} is a suitable deterministic counterpart of X and $\Psi \operatorname{Spec}_{\epsilon}$ stands for the ϵ -pseudospectrum⁸. In particular, their result implies that A + X has no outliers on macroscopic scale. Notice that the subscript ϵ in $\operatorname{Spec}_{\epsilon}$ from (2.15) and in $\Psi \operatorname{Spec}_{\epsilon}$ from (2.20) plays very different roles.

In the case of constant variance profile, i.e. for $X_{iid} + A$, when X_{iid} is an IID matrix and A is a general deformation satisfying Assumption 2.8, we immediately get a much stronger result: we obtain both inclusions in (2.20) with the order one ϵ, δ replaced with the optimal local fluctuation scale for the smallest singular value, see (3.15) (up to an N^{ϵ} factor). This follows from our optimal local laws both inside the spectrum (3.7) and outside (3.14). Moreover a similar two-sided inclusion with optimal precision also holds for the spectrum instead of the pseudospectrum in the following form:

$$\operatorname{Spec}(A+X) \subset \operatorname{Spec}_{\epsilon}(A+\boldsymbol{x}) \subset \bigcup_{z \in \operatorname{Spec}(A+X)} D(z, N^{\epsilon+\delta}\sigma_{\mathrm{f}}(z)).$$

More precisely, the first inclusion is just (2.16), while the second inclusion follows from the fact that there many eigenvalues close to the edge of the support of ρ_{A+x} , as a consequence of the local law (2.18).

In fact, our method can prove the optimal no-outlier theorem for A + X with X having a flat variance profile using the same decomposition as in Section 2.1.1. More precisely, we consider A + X as a deformed IID matrix, i.e. write $A + X \stackrel{d}{=} A + \tilde{X} + \epsilon X^{\text{Gin}}$ and condition on \tilde{X} . Then we need to guarantee that Assumption 2.8 holds with A replaced by $A + \tilde{X}$ with very high probability, given the same pair of inputs as in Section 2.1.1 for $A + \tilde{X}$ instead of \tilde{X} . Namely, we need to guarantee (i) the macroscopic convergence and no-outlier theorem for $\rho_{|A+\tilde{X}-z|}$ and (ii) inf $\operatorname{supp} \rho_{|A+\tilde{x}-z|} \sim 1$ for $z \in \partial \operatorname{supp} \rho_{A+x}$, where \tilde{x} and x are the operator analogues of \tilde{X} and X. The first input (i) already follows from [47, Theorem 2.1 and Corollary 2.3] since they allow general deformations. The second input (ii), which is a purely deterministic result, needs to be checked by solving the corresponding MDE on the macroscopic scale. For example, this was explicitly done in [11, Proposition 5.15 (iv)] for the special case where A is diagonal.

2.3. Organization of the proof. In Section 3 we gather the gather the free probabilistic and random matrix tools which we use in the proofs of our main results. This includes the statement of the local law (see Theorem 3.3). We then present the proof of Theorem 2.11, Corollary 2.12, and Theorem 2.13. In Section 4 we prove Theorem 2.7 using our main technical comparison result Theorem 4.2, which compares the joint moments of log determinants of various non-Hermitian random matrices. In Sections 5–7 we provide the proof of Theorem 4.2. These three sections divide the proof up into the cases: Section 5 comparing $A + X^{\text{Gin}(\mathbb{C})}$ and $X^{\text{Gin}(\mathbb{C})}$, Section 6 comparing $A + X^{\text{Gin}(\mathbb{R})}$ and $X^{\text{Gin}(\mathbb{R})}$, and finally Section 7 comparing A + X and $A + X^{\text{Gin}(\mathbb{F})}$ for both fields $\mathbb{F} = \mathbb{R}$, \mathbb{C} . Finally, in Section 8, we present the proof of the local law Theorem 3.3.

3. Preliminaries and the proofs of Theorems 2.11-2.13

In this section we present the main technical ingredients. In Section 3.1, we list some properties of the Brown measure ρ_{A+x} and we state the local law close to the cusp regime of the Hermitization of A + X - z, when z is a regular edge point. Then, in Section 3.2 we prove Theorems 2.11–2.13.

3.1. Preliminary results and the local law for A + X. Before we state the local law for A + X (see Theorem 3.3 below), we collect some preliminary results on free probability and matrix Dyson equations.

Lemma 3.1. The Brown measure ρ_{A+x} has a density that satisfies the following:

- (i) $\rho_{A+\boldsymbol{x}}(z) \leq 1/\pi$ for all $z \in \mathbb{C}$.
- (ii) supp $\rho_{A+\boldsymbol{x}} = \overline{\{z \in \mathbb{C} : \langle |A-z|^{-2} \rangle > 1\}}.$
- (iii) If z_0 satisfies Assumption 2.6 (i)-(ii), the density $\rho_{A+x}(z_0+w)$ follows the following asymptotics as $w \to 0$:

(3.1)
$$\rho_{A+\boldsymbol{x}}(z_0+w) = \frac{1}{\pi} |\gamma_0|^2 + O(|w|), \qquad \text{for } z_0+w \in \operatorname{supp} \rho_{A+\boldsymbol{x}}, \\ \rho_{A+\boldsymbol{x}}(z_0+w) \equiv 0, \qquad \text{for } z_0+w \notin \operatorname{supp} \rho_{A+\boldsymbol{x}},$$

 $^{^{7}}$ After we have made them aware of the current work, the authors of [11] informed us that they can relax the flatness assumption.

⁸The ϵ -pseudospectrum of any matrix X is defined by $\Psi \operatorname{Spec}_{\epsilon}(X) := \{z \in \mathbb{C} : ||(X-z)^{-1}|| \ge \epsilon^{-1}\}.$

where γ_0 was defined in (2.8). The implicit constant in (3.1) depends only on the C_0 from Definition 2.2 and C_1 from Assumption 2.6 (ii).

Proof. The first two statements of Lemma 3.1 are due to [16, Theorem 7.10], and the last is from [53, Theorem 2.9 (iii)]. \Box

Lemma 3.2. For each $z \in \mathbb{C}$ and $\hat{z} \in \mathbb{C}_+$, the matrix Dyson equation (MDE)

(3.2)
$$-\frac{1}{M} = -\begin{pmatrix} 0 & A-z \\ (A-z)^* & 0 \end{pmatrix} + \hat{z} + \langle M \rangle$$

has a unique $2N \times 2N$ solution $M \equiv M_{A,z}(\hat{z})$ with $\operatorname{Im} M = \frac{1}{2i}(M - M^*) > 0$. Then we have

(3.3)
$$\left\langle \begin{pmatrix} -\widehat{z} & A+\boldsymbol{x}-z \\ (A+\boldsymbol{x}-z)^* & -\widehat{z} \end{pmatrix}^{-1} \right\rangle_{\mathcal{M}} = M_{A,z}(\widehat{z}) \in \mathbb{C}^{2N \times 2N}, \quad \forall z \in \mathbb{C}, \ \widehat{z} \in \mathbb{C}_+,$$

so that the Brown measure of the sum A + x can be computed as the distributional Laplacian

(3.4)

$$\rho_{A+\boldsymbol{x}}(z) \equiv -\frac{1}{2\pi} \Delta_z \int_0^\infty \left[\left\langle \left\langle \frac{\eta}{|A+\boldsymbol{x}-z|^2+\eta^2} \right\rangle_{\mathcal{M}} \right\rangle - \frac{\eta}{1+\eta^2} \right] \mathrm{d}\eta$$

$$= -\frac{1}{2\pi} \Delta_z \int_0^\infty \left[\mathrm{Im} \langle M_{A,z}(\mathrm{i}\eta) \rangle - \frac{\eta}{1+\eta^2} \right] \mathrm{d}\eta.$$

Uniqueness of solution of the Matrix Dyson Equation in the following lemma is a consequence of [59, Theorem 2.1], see [16] for the form of the solution in terms of circular elements. Note, for $\hat{z} = i\eta$ on the imaginary axis, $\langle M(i\eta) \rangle$ is purely (positive) imaginary and M is the resolvent of the Hermitization of (A - z) evaluated at $i\eta + \langle M \rangle$. Hence for all $\eta > 0$ we have

(3.5)
$$||M(i\eta)|| \le ||(A-z)^{-1}||$$

A key tool in the proof of Theorem 2.7, 2.11–2.13 is an optimal local law for the Hermitization H^z defined in (1.1). We define the resolvent of H^z by

(3.6)
$$G^{A,z}(\hat{z}) = G^{z}(\hat{z}) = (H^{z} - \hat{z})^{-1}, \quad \hat{z} \in \mathbb{C}_{+},$$

although for our local law we will exclusively consider $\hat{z} = i\eta$ for $\eta \ge 0$ on the positive imaginary axis. The dependence of $G^{A,z}$ through H^z defined in (1.1) will often be omitted when A is to be thought of as fixed.

The local law gives an optimal comparison between G^z and the deterministic approximation $M_{A,z}$ near an edge point z_0 . The proof of this local law will be presented in Section 8.

Theorem 3.3 (Local law for H^{z_0} near the edge). Let A + X be a deformed IID matrix as in Definition 2.2. Let z_0 be an edge point of A + X such that $||(A - z_0)^{-1}|| < C_4$ for some $C_4 > 0$. There exists a (small enough) constant c > 0 such that for any deterministic vectors $\boldsymbol{u}, \boldsymbol{y} \in \mathbb{C}^{2N}$ and matrix $B\mathbb{C}^{2N \times 2N}$ (3.7)

$$|\langle B(G^{z}(\mathrm{i}\eta) - M_{A,z}(\mathrm{i}\eta))\rangle| \prec ||B|| \frac{1}{N\eta}, \qquad |\boldsymbol{u}^{*}(G^{z}(\mathrm{i}\eta) - M_{A,z}(\mathrm{i}\eta))\boldsymbol{y}| \prec ||\boldsymbol{u}|| ||\boldsymbol{y}|| \left[\sqrt{\frac{\rho^{A,z}(\mathrm{i}\eta)}{N\eta}} + \frac{1}{N\eta}\right],$$

uniformly in A, X, z_0 satisfying (2.1), (2.2), $||(A - z_0)^{-1}|| \le C_4$, $|z - z_0| < c$ and $N^{-1} \le \eta \le N^{100}$.

For $z \in \mathbb{C} \setminus \text{Spec}(A)$, we define the function $f_{A-z} : \mathbb{C} \to (0, \infty)$ as

(3.8)
$$f_{A-z}(w) := \left\langle \frac{1}{|A-z-w|^2} \right\rangle$$

Lemma 3.4. Let $A \in \mathbb{C}^{N \times N}$ and $z \in \partial \operatorname{supp} \rho_{A+x}$ satisfy $||A|| \leq C_0$ and $||(A-z)^{-1}|| \leq C_1$ for some constants $C_0, C_1 > 0$. As $\eta, |w| \to 0$, we have the following asymptotics:

(3.9)
$$\operatorname{Im}\langle M_{A,z+w}(\mathrm{i}\eta)\rangle \sim \begin{cases} |f_{A-z}(w)-1|^{1/2}+\eta^{1/3}, & z+w \in \operatorname{supp} \rho_{A+x}, \\ \frac{\eta}{|f_{A-z}(w)-1|+\eta^{2/3}}, & z+w \notin \operatorname{supp} \rho_{A+x}. \end{cases}$$

Additionally, if $z \in \mathcal{D}_{C_1}$ is a sharp edge (recall \mathcal{D}_C from (2.6)), then the regularized eigenvalue density of H^z , $v := \operatorname{Im}\langle M_{A,z+w}(i\eta) \rangle + \eta$, satisfies the following asymptotic cubic equation as $|w|, \eta \to 0$:

(3.10)
$$I_4 v^3 - 2 \operatorname{Re}[w I_3] v - \eta = O((|w|^{1/2} + \eta^{1/3})^5).$$

where I_3 and I_4 are the same quantities as in (2.9) but with z_0 replaced by z. The implicit constants in (3.9) and (3.10) depend only on C_0 and C_1 .

Proof. The asymptotic in (3.9) was proven in [53, Lemma 5.1], hence we only prove (3.10) for a sharp edge z. With $v = \eta + \text{Im}\langle M_{A,z+w}(i\eta) \rangle$, we may write the (trace of) the MDE (3.2) as

(3.11)
$$v - \eta = \left\langle \frac{v}{|A - z - w|^2 + v^2} \right\rangle.$$

Then we expand the right-hand side with respect to v and then w, so that

(3.12)
$$\begin{aligned} v - \eta &= v \left\langle \frac{1}{|A - z - w|^2} \right\rangle - v^3 \left\langle \frac{1}{|A - z - w|^4} \right\rangle + O(v^5) \\ &= v \left\langle \frac{1}{|A - z|^2} \right\rangle + 2 \operatorname{Re}[wI_3]v - I_4 v^3 + O(v^5 + v|w|^2 + v^3|w|) \\ &= v + 2 \operatorname{Re}[wI_3]v - I_4 v^3 + O((|w|^{1/2} + \eta^{1/3})^5), \end{aligned}$$

where in the last line we also used (3.9). Rearranging the terms proves the cubic equation.

3.2. **Proof of Theorems 2.11–2.13.** We first present the proof of Theorem 2.13, and then we conclude this section with the proof of Theorem 2.11 and Corollary 2.12. All estimates in this section are uniform over X, A, and z_0 satisfying (2.1), (2.2), and $||(A - z_0)^{-1}|| \leq C_4$.

Proof of Theorem 2.13. Given the local law (3.7) for the Hermitized resolvent, the proof of (2.18) is by now standard (see e.g. [8, Section 5.2]). Here we sketch the main steps for completeness.

This proof relies on Girko's formula (1.2) in the form

(3.13)
$$\int_{\mathbb{C}} F(z) d\rho_{A+X}(z) = -\frac{1}{4\pi N} \int_{\mathbb{C}} \Delta F(z) \int_{N^{-100}}^{N^{100}} \langle \Im G^{z}(i\eta) \rangle d\eta d^{2}z + O(N^{-10})$$

where we used the bound for the smallest singular value in [88, Theorem 3.2] to remove the very small $\eta \leq N^{-100}$ regime and simple explicit calculations to remove the regime $\eta \geq N^{100}$ (see e.g. [8, Eq. (5.27)]). Then using the monotonicity of $\eta \to \eta \langle \Im G^z(i\eta) \rangle$ to extend (3.7) down to $\eta \geq N^{-100}$, and plugging (3.7) into (3.13) we immediately obtain the bound (2.18).

Proof of Theorem 2.11. To prove that the are no outliers in the spectrum of A + X we need an improvement of the averaged local law (3.7) when z_0 is outside of the spectrum (see the proof of [9, Theorem 2.1] for a similar analysis). In particular, for any deterministic $B \in \mathbb{C}^{2N \times 2N}$, we claim that

(3.14)
$$\left| \langle (G^{z}(i\eta) - M_{A,z}(i\eta))B \rangle \right| \prec \frac{1}{N\kappa}$$

with $\kappa := \text{dist}(i\eta, \text{supp}(\rho^z))$; note that if 0 is in $\text{supp}(\rho^z)$, then $\kappa = \eta$. If z is well outside of the support of ρ^z , i.e. more than the typical fluctuation scale, then we can choose even $\eta = 0^+$. The proof of (3.14) is presented at the end of Section 8.1.

Proceeding similarly to the proof of [8, Theorem 2.5 (ii)] we can show that there are no eigenvalues in the region

$$\{z \in \mathbb{C} : \operatorname{dist}(z, \operatorname{supp}(\rho_{A+\boldsymbol{x}})) \ge c\},\$$

with very high probability for some small N-independent constant c > 0. We now strengthen this statement by reducing c to $N^{\epsilon}\sigma_{\rm f}$. More precisely, we show that if we pick $z \notin \operatorname{Spec}_{\epsilon}(A + \boldsymbol{x})$ then $z \notin \operatorname{Spec}(A + X)$. Fix an edge point z_0 , then for $z \notin \operatorname{Spec}_{\epsilon}(A + \boldsymbol{x})$ such that $|z - z_0| \leq c$, we define (cf. [48, Eq. (2.7)]) the Hermitian fluctuation scale

(3.15)
$$\eta_{\rm f} = \eta_{\rm f}(z) := \frac{|f_{A-z_0}(z) - 1|^{1/6}}{N^{2/3}},$$

with f_{A-z_0} being defined in (3.8). Note that if $z \notin \operatorname{Spec}_{\epsilon}(A+\mathbf{x})$ then $|f_{A-z_0}(z)-1| \geq N^{-1/2+c_*\epsilon}$, for some $c_* > 0$. This is a consequence of Lemma 3.1 close to sharp edges and of [53, Theorem 2.9 (iv)] for quadratic edges. Fix $\eta := N^{c_*\epsilon/5}\eta_{\mathrm{f}}$, then, by (3.9) it follows that

(3.16)
$$\langle \Im M_{A,z}(\mathbf{i}\eta) \rangle \le \frac{N^{-c_*\epsilon/5}}{N\eta}$$

Next, using the local law (3.14), together with (3.16), we have

(3.17)
$$\sup\left\{ \langle \Im G^{z}(\mathrm{i}\eta) \rangle : |z - z_{0}| \leq c, |f_{A-z_{0}}(z) - 1| \geq N^{-1/2+c_{*}\epsilon} \right\} \prec \frac{N^{-c_{*}\epsilon/5}}{N\eta}.$$

Here we used a standard grid argument and the Lipschitz continuity of $z \mapsto \langle \Im G^z \rangle$ to show that (3.14) holds simultaneously for all $|z - z_0| \leq c$.

Denote the eigenvalues of H^z by

$$\lambda_{-N}^{z} \leq \dots \leq \lambda_{-1}^{z} \leq 0 \leq \lambda_{1}^{z} \leq \dots \leq \lambda_{N}^{z},$$

then (3.17) implies that as long as z is such that $|f_{A-z_0}(z)-1| \ge N^{-1/2+c_*\epsilon}$ we have $\lambda_1^z \ne 0$ with very high probability (recall that $\lambda_{-i}^z = -\lambda_i^z$). In fact, if $\lambda_1^z = 0$ then we would have $\Im G^z(i\eta) \ge 1/(2N\eta)$, which contradicts (3.17). This concludes the proof showing that $z \notin \operatorname{Spec}(A+X)$, as a consequence of the fact that

$$(3.18) 0 \in \operatorname{Spec}(H^z) \Longleftrightarrow z \in \operatorname{Spec}(A+X).$$

We conclude this section with the proof of cluster-rigidity.

Proof of Corollary 2.12. This proof is similar to the proof of [7, Corollary 2.9]. We thus only highlight the main differences and present a sketch of the proof for the convenience of the reader.

As a consequence of (3.18), similarly to the proof of Theorem 2.11 above, it is enough to show that the Hermitization H^z satisfies "band rigidity", in the sense of [7, Corollary 2.9], when its density has a gap around zero of order one size.

Consider a contour Γ which encircles C and it is such that $\operatorname{dist}(\Gamma, \operatorname{Spec}_{\epsilon}(A+x)) \geq \delta/10$. For any $z \in \Gamma$ we let H^z to be the Hermitization of A+X-z. We now consider a flow that interpolates between H^z and \mathcal{A}^z , the Hermitization of A-z, which is a deterministic matrix, i.e. define H_t^z to be the Hermitization of $A+X_t-z$, where

Note that $H_0^z = H^z$, $H_1^z = \mathcal{A}^z$, and $H_t^z = \sqrt{1-t}W + \mathcal{A}^z$, with W being the Hermitization of X. We now show that for any D > 0 we have

(3.20)
$$\mathbb{P}(0 \in \operatorname{Spec}(H_t^z) \text{ for some } (t, z) \in [0, 1] \times \Gamma) \le N^{-D}.$$

As a consequence of (3.18), this yields the desired result (3.17).

To prove (3.20), we distinguish the analysis into two cases: t close to one and the rest. By assumption, $0 \notin \operatorname{Spec}(\mathcal{A}^z)$ for any $z \in \Gamma$, additionally $||H^0|| \leq C_*$, for some fixed $C_* > 0$, we can thus choose a sufficiently small $c_* > 0$ such that $0 \notin \operatorname{Spec}(H_t^z)$, for any $z \in \Gamma$, for $t \in [1 - c_*, 1]$ with very high probability. In the regime when $t \in [0, 1 - c_*)$, the variance of the entries of X_t is of order 1/N, hence, modulo a simple rescaling, X_t is an IID matrix in the sense of Definition 2.1. Hence, we can use the same argument as in (3.16)–(3.17) to conclude that, with very high probability, $0 \notin \operatorname{Spec}(H_t^z)$ for any $z \in \Gamma$ and $t \in [0, 1 - c_*]$. This proves (3.20).

4. Proof of Theorem 2.7

4.1. **Proof of Theorem 2.7.** The purpose of this section is to reduce the proof of Theorem 2.7 to our main technical result, Theorem 4.2 below, which compares joint moments of log determinants and lower tail of the smallest singular value. All estimates (including asymptotic relations and stochastic dominance) in this section are uniform over A, X, and z_0 satisfying (2.1), (2.2), and Assumption 2.6. We first reduce the test function F, i.e. we claim that Theorem 2.7 follows from the following result:

Theorem 4.1. Let $z_0 \in \mathbb{C}$ satisfy Assumption 2.6, γ_0 be defined in (2.8), and $k \in \mathbb{N}$. Let $F_1, \ldots, F_k \in C_c^{\infty}(\mathbb{C})$, and define their local rescalings

$$\widetilde{F}_j := NF_j(\sqrt{N}\gamma_0(\cdot - z_0)), \qquad \widetilde{F}_j^{\operatorname{Gin}} := NF_j(\sqrt{N}z_1^{-1}(\cdot - z_1)),$$

where z_1 is chosen as in Theorem 2.7. Then, for a constant c > 0 we have

(4.1)
$$\mathbb{E}\prod_{j=1}^{k} \left(\int_{\mathbb{C}} \widetilde{F}_{j}(w) \mathrm{d}(\rho_{A+X} - \rho_{A+x})(w) \right) - \mathbb{E}\prod_{j=1}^{k} \left(\int_{\mathbb{C}} \widetilde{F}_{j}^{\mathrm{Gin}}(w) \mathrm{d}(\rho_{X^{\mathrm{Gin}(\mathbb{F})}} - \rho_{x})(w) \right) = O(N^{-c}),$$

where c > 0 and the implicit constant depend⁹ on k and F_j 's.

Proof of Theorem 2.7. First, we prove (2.10) when F is of the product form $F(\boldsymbol{w}) = \prod_{j=1}^{k} F_j(w_j)$. Given Theorem 4.1, it suffices to check that

(4.2)
$$\left| \int_{\mathbb{C}} \widetilde{F}_{j}(w) \mathrm{d}\rho_{A+\boldsymbol{x}}(w) - \int_{\mathbb{C}} \widetilde{F}_{j}^{\mathrm{Gin}}(w) \mathrm{d}\rho_{\boldsymbol{x}}(w) \right| = O(N^{-1/2}).$$

⁹They depend on F_j 's only via $||F_j||_{L^{\infty}}, ||\Delta F_j||_{L^1}$, and diam(supp F_j).

To see (4.2), we write ρ for the density of $\rho_{A+\boldsymbol{x}}$ and

(4.3)
$$\int_{\mathbb{C}} \widetilde{F}_{j}(w) \mathrm{d}\rho_{A+\boldsymbol{x}}(w) = |\gamma_{0}|^{-2} \int_{\mathbb{C}} F_{j}(w) \rho(z_{0} + \gamma_{0}^{-1} N^{-1/2} w) \mathrm{d}^{2} w.$$

On the other hand, Lemma 3.1 (ii) gives

(4.4)
$$|\gamma_0|^{-2}\rho(z_0 + \gamma_0^{-1}N^{-1/2}w) = \begin{cases} \frac{1}{\pi} + O(N^{-1/2}|w|) & \text{if } z_0 + \gamma_0^{-1}N^{-1/2}w \in \operatorname{supp}\rho, \\ 0 & \text{elsewhere.} \end{cases}$$

and Lemma 3.1 (ii) implies that $\partial \operatorname{supp} \rho_{\gamma_0(A+\boldsymbol{x}-z_0)}$ is tangent to the half-space $\{w : \operatorname{Re} w \leq 0\}$ so that

(4.5)
$$\int_{w \in \operatorname{supp} F_j} |\mathbb{1}(z_0 + \gamma_0^{-1} N^{-1/2} w \in \operatorname{supp} \rho) - \mathbb{1}(\operatorname{Re} w \le 0)| \mathrm{d}^2 w = O(N^{-1/2}).$$

We thus conclude that

(4.6)
$$\left| \int_{\mathbb{C}} \widetilde{F}_j(w) \mathrm{d}\rho_{A+\boldsymbol{x}}(w) - \frac{1}{\pi} \int_{\mathbb{C}} F_j(w) \mathbb{1}(\operatorname{Re} w \le 0) \mathrm{d}^2 w \right| = O(N^{-1/2}).$$

Repeating the same argument for $\rho_{\boldsymbol{x}}$ proves (4.2).

Now it only remains to extend the test function beyond those in the product form $F(\boldsymbol{w}) = \prod_{j=1}^{k} F_j(w_j)$. To this end, we approximate the given smooth test function F in Theorem 2.7 by a linear combination of those in the product form using truncated Fourier series. We omit further details and refer the reader to [37, Theorem 1] or [90, Corollary 7]. This concludes the proof of Theorem 2.7.

We next present the key technical inputs, and use them to prove Theorem 4.1. The main inputs are two parallel comparison results, for the two pairs of regularized functionals¹⁰

(4.7)
$$L_0(w) := \operatorname{Tr}\log(|A + X - z_0 - \gamma_0^{-1}N^{-1/2}w|^2 + \eta_0^2) - N\langle \log(|A + x - z_0 - \gamma_0^{-1}N^{-1/2}w|^2 + \eta_0^2) \rangle,$$

$$L_1^{\operatorname{Gin}(\mathbb{F})}(w) := \operatorname{Tr}\log(|X^{\operatorname{Gin}(\mathbb{F})} - z_1 - z_1N^{-1/2}w|^2 + \eta_1^2) - N\langle \log(|x - z_1 - z_1N^{-1/2}w|^2 + \eta_1^2) \rangle,$$

and

(4.8)
$$N_{0}(w) := \frac{1}{c_{0}} \left\langle \frac{\eta_{0}}{|A + X - z_{0} - \gamma_{0}^{-1} N^{-1/2} w|^{2} + \eta_{0}^{2}} \right\rangle,$$
$$N_{1}^{\operatorname{Gin}(\mathbb{F})}(w) := \left\langle \frac{\eta_{1}}{|X^{\operatorname{Gin}(\mathbb{F})} - z_{1} - z_{1} N^{-1/2} w|^{2} + \eta_{1}^{2}} \right\rangle.$$

Here, the scales η_0 and η_1 are defined by (recall the definition of I_4 from (2.9))

(4.9)
$$\eta_0 := c_0^{-1} \eta_1, \qquad c_0 := I_4^{-1/4}, \qquad \eta_1 := N^{-3/4-\delta},$$

where $0 < \delta < (1 \land \mathfrak{c})/100$ is a small but fixed constant, with¹¹ \mathfrak{c} from Assumption 2.6 (iii \mathbb{R}) and (iii \mathbb{R}'). With our choice of the scale η_1 , we define the fundamental control parameter

(4.10)
$$\Psi := \frac{1}{N\eta_1} = N^{-1/4+\delta}.$$

Our error estimates are typically of the form $O(N^{\epsilon}\Psi^n)$ for some $n \in \mathbb{N}$ and arbitrarily small (but fixed) $\epsilon > 0$. Now we are ready to state our comparison results:

Theorem 4.2. Let $k \in \mathbb{N}$ and C > 0 be fixed. Then there exists a constant c > 0 such that the following hold uniformly in $w \in \mathbb{C}$ and $w = (w_1, \ldots, w_k) \in \mathbb{C}^k$ with $|w|, ||w|| \leq C$:

(4.11)
$$\mathbb{E}\prod_{i=1}^{k} L_0(w_i) = \mathbb{E}\prod_{i=1}^{k} L_1^{\operatorname{Gin}(\mathbb{F})}(w_i) + h^{(0)}(\boldsymbol{w}) + O(N^{-c}),$$

(4.12)
$$\mathbb{E}N_0(w) = \mathbb{E}N_1^{\operatorname{Gin}(\mathbb{F})} + O(N^{-c}\Psi),$$

where $h^{(0)} \equiv h_N^{(0)} \in C^{2k}(\mathbb{C}^k)$ is harmonic in at least one variable, i.e. it satisfies $(\prod_{j=1}^k \Delta_{w_j})h^{(0)} \equiv 0$.

¹⁰The parameter $z_1 \in \mathbb{S}^1$ in (4.7) and (4.8) do not play any role when $\mathbb{F} = \mathbb{C}$ since a complex Ginibre matrix is rotationally invariant. Nonetheless, we keep the current form to facilitate the proof in the real case in Section 6.

¹¹The c-dependent constraint for δ is not needed in the complex case, but we include it here to avoid repetition.

Proof of Theorem 4.1. Using Girko's formula, we express the two statistics in terms of the log potential:

(4.13)

$$\int_{\mathbb{C}} \widetilde{F}_{j}(w) \mathrm{d}\rho_{A+X}(w) = N \int_{\mathbb{C}} F_{j}(w) \mathrm{d}\rho_{N^{1/2}\gamma_{0}(A+X-z_{0})}(w) \\
= -\frac{N}{2\pi} \int_{\mathbb{C}} \Delta_{w} F_{j}(w) \langle \log | N^{1/2}\gamma_{0}(A+X-z_{0}) - w | \rangle \mathrm{d}^{2}w \\
= -\frac{N}{2\pi} \int_{\mathbb{C}} \Delta_{w} F_{j}(w) \langle \log | A+X-z_{0} - \gamma_{0}^{-1}N^{-1/2}w | \rangle \mathrm{d}^{2}w,$$

where in the second line we used integration by parts and in the third line we subtracted $\log(N^{1/2}|\gamma_0|)$ from the integrand using harmonicity. Writing the same expression for $X^{\text{Gin}(\mathbb{F})}$ as well as for the deterministic counterparts, we decompose

(4.14)
$$\int_{\mathbb{C}} \widetilde{F}_{j}(w) \mathrm{d}(\rho_{A+X} - \rho_{A+\boldsymbol{x}})(w) = I_{1}^{(j)} + I_{2}^{(j)},$$
$$\int_{\mathbb{C}} \widetilde{F}_{j}^{\mathrm{Gin}}(w) \mathrm{d}(\rho_{X^{\mathrm{Gin}(\mathbb{F})}} - \rho_{\boldsymbol{x}})(w) = I_{1}^{(j)\mathrm{Gin}(\mathbb{F})} + I_{2}^{(j)\mathrm{Gin}(\mathbb{F})},$$

where

(4.15)
$$I_{1}^{(j)} := -\frac{1}{2\pi} \int_{\mathbb{C}} \Delta_{w} F_{j}(w) L_{0}(w) d^{2}w,$$
$$I_{2}^{(j)} := -\frac{1}{2\pi} \int_{\mathbb{C}} \Delta_{w} F_{j}(w)$$
$$\times \int_{0}^{\eta_{0}} N \left\langle \frac{\eta_{0}}{|A+X-z_{0}-\gamma_{0}^{-1}w|^{2}+\eta_{0}^{2}} - \operatorname{Im} M_{A-z_{0},\gamma_{0}^{-1}w}(\mathrm{i}\eta_{0}) \right\rangle d\eta d^{2}w,$$

and $I_1^{(j)\operatorname{Gin}(\mathbb{F})}$ and $I_2^{(j)\operatorname{Gin}(\mathbb{F})}$ are defined in a similar fashion with (z_0, η_0) replaced by (z_1, η_1) , etc. The next lemma shows that the product of $I_1^{(j)}$'s has the leading contribution to the statistics.

Lemma 4.3. For some constant c > 0 depending on k and F_j 's, we have

(4.16)
$$\mathbb{E}\prod_{j=1}^{k} \left(\int_{\mathbb{C}} \widetilde{F}_{j}(w) \mathrm{d}(\rho_{A+X} - \rho_{A+x})(w) \right) = \mathbb{E}\prod_{j=1}^{k} I_{1}^{(j)} + O(N^{-c}),$$
$$\mathbb{E}\prod_{j=1}^{k} \left(\int_{\mathbb{C}} \widetilde{F}_{j}^{\mathrm{Gin}}(w) \mathrm{d}(\rho_{X^{\mathrm{Gin}(\mathbb{F})}} - \rho_{x})(w) \right) = \mathbb{E}\prod_{j=1}^{k} I_{1}^{(j)\mathrm{Gin}(\mathbb{F})} + O(N^{-c}).$$

Given Lemma 4.3 and (4.11), we conclude the proof of Theorem 4.1 as

$$\mathbb{E} \prod_{j=1}^{k} \left(\int_{\mathbb{C}} \widetilde{F}_{j}(w) \mathrm{d}(\rho_{A+X} - \rho_{A+x})(w) \right) - \mathbb{E} \prod_{j=1}^{k} \left(\int_{\mathbb{C}} \widetilde{F}_{j}^{\mathrm{Gin}}(w) \mathrm{d}(\rho_{X^{\mathrm{Gin}(\mathbb{F})}} - \rho_{x})(w) \right)$$

$$= \left| \mathbb{E} \prod_{j=1}^{k} I_{1}^{(j)} - \mathbb{E} \prod_{j=1}^{k} I_{1}^{(j)\mathrm{Gin}(\mathbb{F})} \right| + O(N^{-c})$$

$$\leq \int_{\mathbb{C}^{k}} \prod_{j=1}^{k} |\Delta F_{j}(w_{j})| \left| \mathbb{E} \prod_{j=1}^{k} L_{0}(w_{j}) - \prod_{j=1}^{k} L_{1}^{\mathrm{Gin}(\mathbb{F})}(w_{j}) \right| \mathrm{d}^{2k} w + O(N^{-c})$$

$$\leq N^{-c},$$

where in the last line we used that $F_j \in C_c^{\infty}(\mathbb{C})$.

Proof of Lemma 4.3. We only prove the first estimate in (4.16), which implies the second by taking A = 0 and $X = X^{\operatorname{Gin}(\mathbb{F})}$.

We first collect the inputs. As a standard corollary of local laws (see e.g. $[37, \text{Lemma 3}]^{12}$), we have

(4.19)
$$|I_1^{(j)}| + |I_2^{(j)}| \prec ||\Delta F_j||_{L^1} \lesssim 1.$$

Also, we can deduce from (4.12) and [37, Eq.(28)] that

(4.20)

$$\mathbb{P}[\lambda_1(A + X - z_0 - \gamma_0^{-1} N^{-1/2} w) \le \eta_0] \le 2c_0 N \eta_0 N_0(w) \\
= 2N \eta_1 N_1^{\operatorname{Gin}(\mathbb{F})} + O(N^{-c}) \\
\lesssim N^{-c},$$

where $\lambda_1(\cdot)$ denotes the smallest singular value. Then (4.20) in turn implies, following the same proof as [37, Lemma 4] and taking small enough $\delta, c > 0$, that (recall δ from (4.9))

(4.21)
$$\mathbb{E}I_2^{(j)} = O(N^{-c}).$$

Finally, we have the non-random, non-asymptotic inequalities

$$(4.22) |I_1^{(j)}| \le \frac{N}{\eta_0^2} \left\| (\|A\|^2 + \|X\|^2 + N^{-1}|w|^2) \Delta F_j(w) \right\|_{L^1(\mathrm{d}^2w)}, |I_2^{(j)}| \le N \|F_j\|_{L^\infty} + |I_1^{(j)}|,$$

which in turn implies

(4.23)
$$|I_1^{(j)}| + |I_2^{(j)}| \lesssim N^{10} (1 + ||X||^2).$$

The first inequality in (4.22) follows from the fact that, for any $Y \in \mathbb{C}^{N \times N}$, $f \in C_c^2(\mathbb{C})$, and $\eta > 0$,

(4.24)
$$\begin{aligned} \left| \int_{\mathbb{C}} \Delta f(w) \left\langle \log(|Y|^{2} + \eta^{2}) \right\rangle d^{2}w \right| &= \left| \int_{\mathbb{C}} \Delta f(w) \left\langle \log\left(1 + \frac{|Y|^{2}}{\eta^{2}}\right) \right\rangle d^{2}w \right| \\ &\leq \int_{\mathbb{C}} |\Delta f(w)| \left(1 + \frac{\left\langle |Y|^{2} \right\rangle}{\eta^{2}}\right) d^{2}w, \end{aligned}$$

and the second inequality in (4.22) is due to (4.14).

Computing the difference of both sides of (4.16) and using (4.14), it suffices to prove that

(4.25)
$$\mathbb{E}\prod_{j=1}^{k} I_{i_j}^{(j)} = O(N^{-c/2}), \quad i_j \in \{1, 2\},$$

whenever $|\{j: i_j = 2\}| \ge 1$. We define the event Ξ on which

(4.26)
$$\max_{1 \le j \le k} (|I_1^{(j)}| + |I_2^{(j)}|) \le N^{c/2k},$$

Then (4.19) implies $\mathbb{P}[\Xi^c] = O(N^{-D})$ for any fixed D > 0.

Assuming $i_1 = 2$ without loss of generality, we now estimate the expectation in (4.25). Restricting on the event Ξ , we have

(4.27)
$$\mathbb{E}\mathbb{1}_{\Xi} \prod_{j=1}^{k} I_{i_j}^{(j)} \le N^{c(k-1)/(2k)} \mathbb{E}I_1^{(2)} \le N^{-c/2}.$$

On the complementary event, we use Cauchy-Schwarz inequality and (4.23) and then choose large enough D > 0 to get

(4.28)
$$\mathbb{E}\mathbb{1}_{\Xi^{c}} \prod_{j=1}^{k} I_{i_{j}}^{(j)} \lesssim \mathbb{P}[\Xi^{c}]^{1/2} N^{10} (1 + \mathbb{E}(\|X\|^{4})^{1/2}) \lesssim N^{10-D/2} \lesssim N^{-c/2}.$$

Combining (4.27) and (4.28) concludes the proof of Lemma 4.3.

(4.18)
$$g \in L^{1+\alpha}(\mathbb{C}), \qquad \|g\|_{1+\alpha} \le N^{\beta}$$

¹²In [37] there is the additional assumption which we do not impose that there exist $\alpha, \beta > 0$ such that the entries of X have density g satisfying

However, as is pointed out in [37, Remark 1] this assumption is to simplify presentation, and could be removed using [88, Theorem 3.2]. Alternatively we could assume X has a very small Gaussian component and use [90, Section 6.1] as was done in [43, Section 6].

The remaining content of this section as well as Sections 5–7 are devoted to the proof of Theorem 4.2. The proof is first divided into two parts by considering $A + X^{\operatorname{Gin}(\mathbb{F})}$ as an intermediate ensemble between A + X and $X^{\operatorname{Gin}(\mathbb{F})}$. To be more precise, we introduce the intermediate functions

(4.29)

$$L_{0}^{\operatorname{Gin}(\mathbb{F})}(w) := \operatorname{Tr} \log(|A + X^{\operatorname{Gin}(\mathbb{F})} - z_{0} - \gamma_{0}^{-1}N^{-1/2}w|^{2} + \eta_{0}^{2}) - \operatorname{Tr} \langle \log(|A + x - z_{0} - \gamma_{0}^{-1}N^{-1/2}w|^{2} + \eta_{0}^{2}) \rangle_{\mathcal{M}},$$

$$N_{0}^{\operatorname{Gin}(\mathbb{F})}(w) := \frac{1}{c_{0}} \left\langle \frac{\eta_{0}}{|A + X^{\operatorname{Gin}(\mathbb{F})} - z_{0} - \gamma_{0}^{-1}N^{-1/2}w|^{2} + \eta_{0}^{2}} \right\rangle,$$

obtained by replacing X with $X^{\operatorname{Gin}(\mathbb{F})}$ in L_0 and N_0 . We now state the two key steps of the proof of Theorem 4.2 as follows, whose combination immediately gives Theorem 4.2:

Proposition 4.4. Let $k \in \mathbb{N}$ and C > 0 be fixed. Then there exists a constant c > 0 such that the following hold uniformly over $w \in \mathbb{C}$ and $w = (w_1, \ldots, w_k) \in \mathbb{C}^k$ with $|w|, ||w|| \leq C$:

(4.30)
$$\mathbb{E}\prod_{i=1}^{k} L_{0}^{\operatorname{Gin}(\mathbb{F})}(w_{i}) = \mathbb{E}\prod_{i=1}^{k} L_{1}^{\operatorname{Gin}(\mathbb{F})}(w_{i}) + h^{(0)}(\boldsymbol{w}) + O(N^{-c}).$$

(4.31)
$$\mathbb{E}N_0^{\operatorname{Gin}(\mathbb{F})}(w) = \mathbb{E}N_1^{\operatorname{Gin}(\mathbb{F})}(w) + O(N^{-c}\Psi),$$

where $h^{(0)} \equiv h_N^{(0)} \in C^{2k}(\mathbb{C}^k)$ satisfies $(\prod_{j=1}^k \Delta_{w_j})h^{(0)} \equiv 0.$

Proposition 4.5. Let $k \in \mathbb{N}$ and C > 0 be fixed. Then there exists a constant c > 0 such that the following hold uniformly over $w \in \mathbb{C}$ and $w = (w_1, \ldots, w_k) \in \mathbb{C}^k$ with $|w|, ||w|| \leq C$:

(4.32)
$$\mathbb{E}\prod_{i=1}^{k} L_0(w_i) = \mathbb{E}\prod_{i=1}^{k} L_0^{\mathrm{Gin}}(w_i) + O(N^{-c}),$$

(4.33)
$$\mathbb{E}N_0(w) = \mathbb{E}N_0^{\operatorname{Gin}(\mathbb{F})}(w) + O(N^{-c}\Psi),$$

We prove Proposition 4.4 for $\mathbb{F} = \mathbb{C}$ and $\mathbb{F} = \mathbb{R}$ respectively in Sections 5 and 6 since the real symmetry class requires a quite different and more involved treatment. The proof of Proposition 4.5 is independent of the symmetry class and it is postponed to Section 7.

4.2. Deterministic flow. In this section, we introduce further notations that are commonly used in Sections 5–6 along the proof of Proposition 4.4. We also present preliminary reductions to the proof. Since the IID matrix X is always a Ginibre ensemble in this section (in fact up to Section 6), we omit the superscripts Gin and $Gin(\mathbb{F})$.

The comparison in Proposition 4.4 is performed along a piecewise C^1 flow of deterministic matrices, denoted by $\mathcal{A}_t : [0,1] \to \mathbb{C}^{N \times N}$. The specific choice of the path \mathcal{A}_t differs by symmetry classes due to an additional constraint in the real case, and we make the actual choice at the beginning of Sections 5 and 6. Nonetheless, they all satisfy the same condition that the origin remains a sharp edge along the entire flow; we state this in the following assumption:

Assumption 4.6. The path $[0,1] \ni t \mapsto \mathcal{A}_t \in \mathbb{C}^{N \times N}$ is piecewise C^1 and it satisfies the following:

(i)
$$\mathcal{A}_0 = A - z_0$$
 and $\mathcal{A}_1 = -z_1 \in \mathbb{S}$

(ii) For all $t \in [0, 1]$, we have

(4.34)
$$\left\langle \frac{1}{|\mathcal{A}_t|^2} \right\rangle = 1$$

(iii) There exists a constant $C_5 > 1$ such that for all but finitely many $t \in [0,1]$ we have

$$(4.35) \|\mathcal{A}_t\| \le C_5, \|\mathcal{A}_t^{-1}\| \le C_5, C_5^{-1} \le |\langle \mathcal{A}_t|\mathcal{A}_t|^{-4}\rangle| \le C_5, \left\|\frac{\mathrm{d}}{\mathrm{d}t}\mathcal{A}_t\right\| \le C_5\log N.$$

For such a choice of \mathcal{A}_t , we will work with the continuous interpolation $\mathcal{A}_t + X$ connecting $A + X - z_0$ to $X - z_1$. The scaling parameters of this interpolation are as follows; we introduce the time-dependent counterpart of the auxiliary quantities I_3, I_4 in (2.9):

(4.36)
$$I_3 \equiv I_3(t) := \left\langle \mathcal{A}_t^* \frac{1}{|\mathcal{A}_t|^4} \right\rangle, \qquad I_4 \equiv I_4(t) := \left\langle \frac{1}{|\mathcal{A}_t|^4} \right\rangle,$$

and define c_t and γ_t as

(4.37)
$$c_t := I_4(t)^{-1/4}, \qquad \gamma_t := -I_4(t)^{-1/2}I_3(t).$$

Next, for $w \in \mathbb{C}$ we introduce the Hermitizations and the regularizing parameter η_t

(4.38)

$$W := \begin{pmatrix} 0 & X \\ X^* & 0 \end{pmatrix},$$

$$H_t^w := \begin{pmatrix} 0 & \mathcal{A}_t + X - \gamma_t^{-1} N^{-1/2} w \\ (\mathcal{A}_t + X - \gamma_t^{-1} N^{-1/2} w)^* & 0 \end{pmatrix},$$

$$\mathbf{h}_t^w := \begin{pmatrix} 0 & \mathcal{A}_t + \mathbf{x} - \gamma_t^{-1} N^{-1/2} w \\ (\mathcal{A}_t + \mathbf{x} - \gamma_t^{-1} N^{-1/2} w)^* & 0 \end{pmatrix},$$

$$J_t^w := \begin{pmatrix} 0 & \gamma_t^{-1} w \\ \overline{\gamma_t}^{-1} \overline{w} & 0 \end{pmatrix},$$

$$\eta_t := c_t^{-1} \eta_1.$$

We denote the resolvent of H_t^w (restricted to the imaginary axis) by

(4.39) $G_t^w(i\eta) = (H_t^w - i\eta)^{-1}, \quad \eta > 0.$

We also introduce the interpolation between $(L_0(w), N_0(w))$ and $(L_1(w), N_1(w))$ as

(4.40)
$$L_t(w) \coloneqq \operatorname{Tr} \log(|H_t^w - i\eta_t|) - \operatorname{Tr} \langle \log(|\mathbf{h}_t^w - i\eta_t|) \rangle_{\mathcal{M}},$$
$$N_t(w) \coloneqq \frac{1}{c_t} \left\langle \frac{\eta_t}{|\mathcal{A}_t + X - \gamma_t^{-1} N^{-1/2} w|^2 + \eta_t^2} \right\rangle,$$

where $\langle \cdot \rangle_{\mathcal{M}}$ is applied entrywise as described in Definition 2.4. Notice the identity

(4.41)
$$N_t(w) = \frac{1}{c_t} \langle \operatorname{Im} G_t^w(i\eta_t) \rangle,$$

and that $L_t(w)$ can be expressed as an integral of $\langle G_t^w(i\eta) - M_t^w(i\eta) \rangle$ in η , i.e.

(4.42)
$$L_t(w) = 2N \int_{\eta_t}^{\infty} \operatorname{Im} \langle G_t^w(i\eta) - M_t^w(i\eta) \rangle \mathrm{d}\eta.$$

Note that the matrix H_t^w as well as the operator h_t^w depend on t only deterministically, i.e.

(4.43)
$$\frac{\mathrm{d}H_t^w}{\mathrm{d}t} = \frac{\mathrm{d}\mathbb{E}H_t^w}{\mathrm{d}t} = \frac{\mathrm{d}\langle \boldsymbol{h}_t^w \rangle_{\mathcal{M}}}{\mathrm{d}t} = \frac{\mathrm{d}\boldsymbol{h}_t^w}{\mathrm{d}t}.$$

The scaling parameters in (4.37) are chosen so that the cubic equation (3.10) for the rescaling of the regularized eigenvalue density of H_t^w , $v_t^w := \text{Im}\langle M_{\mathcal{A}_t, N^{-1/2}\gamma_t^{-1}w}(i\eta_t)\rangle + \eta_t$, becomes time-independent, i.e. we have

(4.44)
$$\left(\frac{v_t^w}{c_t}\right)^3 + 2N^{-1/2}\operatorname{Re}[w]\left(\frac{v_t^w}{c_t}\right) - \eta_1 = O(\Psi^5)$$

thus v_t^w/c_t is essentially constant in t. At the same time, the choice of the scaling parameters also makes the regularized, rescaled, local eigenvalue density

$$w \mapsto \frac{N}{4\pi} \Delta_w \langle \log(|\mathcal{A}_t + X - \gamma_t^{-1} N^{-1/2} w|^2 + \eta_t^2) \rangle$$

almost constant over t for each w; see Lemma 4.8.

Recall from Lemma 3.2 that the $\mathbb{C}^{2N \times 2N}$ -valued Stieltjes transform

(4.45)
$$\left\langle (\boldsymbol{h}_t^w - \hat{z})^{-1} \right\rangle_{\mathcal{M}} \in \mathbb{C}^{2N \times 2N}, \qquad \hat{z} \in \mathbb{C}_+$$

is the unique solution $M\equiv M_{\mathcal{A}_t,N^{-1/2}\gamma_t^{-1}w}(\widehat{z})\in\mathbb{C}^{2N\times 2N}$ to the MDE

(4.46)
$$\frac{1}{M} = \mathbb{E}H_t^w - \hat{z} - \langle M \rangle, \qquad \text{Im}\, M > 0$$

In accordance with (4.38), we write the *w*-dependent solution M

(4.47)
$$M_t^w(\widehat{z}) := M_{\mathcal{A}_t, N^{-1/2} \gamma_t^{-1} w}(\widehat{z})$$

From the properties of \mathcal{A}_t , we have the following time dependent version of Theorem 3.3.

Proposition 4.7. For any fixed large C > 0, we have

(4.48)
$$|\langle B(G_t^w(i\eta) - M_t^w(i\eta))\rangle| \prec \frac{1}{N\eta}, \qquad |\boldsymbol{x}^*(G_t^w(i\eta) - M_t^w(i\eta))\boldsymbol{y}| \prec ||\boldsymbol{x}|| \, ||\boldsymbol{y}|| \, \sqrt{\frac{\rho^{w,t}(i\eta)}{N\eta}},$$

uniformly over \mathcal{A}_t satisfying Assumption 4.6, $t \in [0,1]$, $|w| \leq C$, and $\eta \in [N^{-1}, N^{100}]$. Consequently, the following holds uniformly over $|w| \leq C$ and $t \in [0,1]$:

$$(4.49) |L_t(w)| \prec 1.$$

Proof. For each fixed $t \in [0, 1]$, as the asymptotic of Lemma 3.4 holds for t this is a straightforward generalization of Theorem 3.3 by replacing $A - z_0$ by A_t . Taking a sufficiently large, but still polynomial sized, grid of [0, 1] and union bounding over the very high probability events on which (4.48) holds we see that (4.48) holds uniformly over $t \in [0, 1]$. To see (4.49), we apply (4.48) to (4.42), so that

$$(4.50) \qquad L_t(w) = 2N \int_{\eta_t}^{\infty} \left[\operatorname{Im} \langle G_t^w(i\eta) - M_t^w(i\eta) \rangle \right] \mathrm{d}\eta \prec 2N \int_{\eta_t}^{N^{100}} \frac{1}{N\eta} \mathrm{d}\eta + N \int_{N^{100}}^{\infty} \frac{1}{\eta^2} \mathrm{d}\eta \lesssim \log N.$$

Notice from Lemma 3.4 that $|\langle M_t^w(i\eta)\rangle| = \text{Im}\langle M_t^w(i\eta)\rangle = O(N^{-1/4})$ for |w| = O(1) and $\eta = O(N^{-3/4})$. We often use the abbreviation

(4.51)
$$\mathring{M}_{t}^{w} := (\mathbb{E}H_{t}^{w})^{-1} \equiv (\langle \boldsymbol{h}_{t}^{w} \rangle_{\mathcal{M}})^{-1} \in \mathbb{C}^{2N \times 2N}$$

The matrix \mathring{M}_t^w solves the degenerate MDE $\mathring{M}^{-1} = \mathbb{E}H_t^w$. We may consider \mathring{M} as a perturbation of the solution M of the true MDE, i.e. for |w| = O(1) and $\eta = O(N^{-3/4})$ we have

(4.52)
$$\|M_t^w(i\eta) - \mathring{M}_t^w\| \lesssim \eta + \langle \operatorname{Im} M_t^w(i\eta) \rangle \lesssim N^{-1/4}.$$

Consequently, in the regime $\eta = O(N^{-3/4})$ and |w| = O(1), $G_t^w(i\eta) - \mathring{M}_t^w$ satisfies the same local law Proposition 4.7 as $G_t^w(i\eta) - M_t^w(i\eta)$ since the errors in (4.48) are larger than $N^{-1/4}$.

In the next lemma, we control the time derivative of the deterministic, regularized Brown measure density.

Lemma 4.8. The following holds uniformly over the path A_t satisfying Assumption 4.6 and w in a compact subset of \mathbb{C} .

(4.53)
$$\frac{\mathrm{d}\langle \log |\boldsymbol{h}_t^w - \mathrm{i}\eta_t| \rangle_{\mathcal{M}} \rangle}{\mathrm{d}t} = -\frac{\mathrm{d}\langle \log |\mathring{M}_t^w| \rangle}{\mathrm{d}t} + O(N^{-3/2}).$$

Moreover, the function $w \mapsto \langle \log | \mathring{M}_t^w | \rangle$, and hence $d \langle \log | \mathring{M}_t^w | \rangle / dt$, are harmonic in w over a compact subset of \mathbb{C} .

Proof. We write the time derivative on the left-hand side of (4.53) as

(4.54)
$$\frac{1}{2}\frac{\mathrm{d}}{\mathrm{d}t}\langle\langle \log(|\boldsymbol{h}_t^w|^2 + \eta_t^2)\rangle_{\mathcal{M}}\rangle = \left\langle\left\langle\frac{\mathrm{d}(\boldsymbol{h}_t^w - \mathrm{i}\eta_t)}{\mathrm{d}t}\frac{1}{\boldsymbol{h}_t^w - \mathrm{i}\eta_t}\right\rangle_{\mathcal{M}}\right\rangle = \left\langle\frac{\mathrm{d}(\mathbb{E}H_t^w - \mathrm{i}\eta_t)}{\mathrm{d}t}M_t^w\right\rangle,$$

where we used in the first equality that h_t^w is supported strictly on the off-diagonal blocks, and in the second equality (4.43).

To see (4.53), we write $iv \equiv iv_t^w := i\eta_t + \langle M_t^w \rangle$ so that the MDE (4.46) becomes

(4.55)
$$\frac{1}{M_t^w} = \frac{1}{\mathring{M}_t^w} - \mathrm{i}v$$

Since $|v| = O(N^{-1/4})$ by (3.9), we have the geometric expansion up to the sixth order

(4.56)
$$\left\| M_t^w - \mathring{M}_t^w \sum_{k=0}^5 (iv \mathring{M}_t^w)^k \right\| = O(v^6) = O(N^{-3/2}).$$

Substituting (4.56) into (4.54) gives

(4.57)
$$\left\langle \frac{\mathrm{d}(\mathbb{E}H_t^w - \mathrm{i}\eta_t)}{\mathrm{d}t} M_t^w \right\rangle = -\frac{\mathrm{d}}{\mathrm{d}t} \langle \log|\mathring{M}_t^w|\rangle + \frac{v^2}{2} \frac{\mathrm{d}}{\mathrm{d}t} \langle (\mathring{M}_t^w)^2 \rangle - \frac{v^4}{4} \frac{\mathrm{d}}{\mathrm{d}t} \langle (\mathring{M}_t^w)^4 \rangle + v \frac{\mathrm{d}\eta_t}{\mathrm{d}t} \left\langle (\mathring{M}_t^w)^2 \right\rangle + O(N^{-3/2} \log N),$$

where the contribution of (most) odd powers of v disappears since $\mathbb{E}H$ and $\mathring{M} = (\mathbb{E}H)^{-1}$ are supported on off-diagonal blocks, and we also used $\eta_t = c_t \eta_1$, $\eta_1 = O(N^{-3/4})$. Next, we expand (4.57) around w = 0; (recall the definitions of I_3, I_4, γ_t and J_t^w from (4.36)–(4.38))

$$\langle (\mathring{M}_{t}^{w})^{2} \rangle = \langle (\mathring{M}_{t}^{0})^{2} \rangle + 2 \left\langle J_{t}^{w} (\mathring{M}_{t}^{0})^{3} \right\rangle + O(N^{-1})$$

$$= 1 + 2N^{-1/2} \operatorname{Re} \left[\frac{w}{\gamma_{t}} I_{3} \right] + O(N^{-1}) = 1 + 2N^{-1/2} I_{4}^{1/2} \operatorname{Re}[-w] + O(N^{-1}),$$

$$\langle (\mathring{M}_{t}^{w})^{4} \rangle = \langle (\mathring{M}_{t}^{0})^{4} \rangle + O(N^{-1/2}) = I_{4} + O(N^{-1/2}).$$

Following the same calculations for the time derivatives of the left-hand sides of (4.58), using the extra input from the last estimate of (4.35), we arrive at the time derivatives of the right-hand sides of (4.58) with the same errors except with an extra factor of log N. Combining this with $\eta_t = I_4^{1/4} \eta_1$, we obtain

$$\left\langle \frac{\mathrm{d}(\mathbb{E}H_t^w - \mathrm{i}\eta_t)}{\mathrm{d}t} M_t^w \right\rangle$$

$$= -\frac{\mathrm{d}}{\mathrm{d}t} \langle \log |\mathring{M}_t^w| \rangle + N^{-1/2} v^2 \operatorname{Re}[-w] \frac{\mathrm{d}I_4^{1/2}}{\mathrm{d}t} - \frac{v^4}{4} \frac{\mathrm{d}I_4}{\mathrm{d}t} + v\eta_1 \frac{\mathrm{d}I_4^{1/4}}{\mathrm{d}t} + O(N^{-3/2} \log N)$$

$$= -\frac{\mathrm{d}}{\mathrm{d}t} \langle \log |\mathring{M}_t^w| \rangle - \frac{v}{4I_4} \frac{\mathrm{d}I_4}{\mathrm{d}t} \left(I_4 v^3 + 2N^{-1/2} I_4^{1/2} v \operatorname{Re}[w] - \eta_t \right) + O(N^{-3/2} \log N).$$

To conclude (4.53), notice that the second term of (4.59) is exactly the cubic equation from Lemma 3.4:

(4.60)
$$I_4 v^3 + 2N^{-1/2} I_4^{1/2} v \operatorname{Re}[w] - \eta_t = I_4 v^3 - 2v \operatorname{Re}\left[N^{-1/2} \gamma_t^{-1} w I_3\right] - \eta_t = O(N^{-5/4}).$$

Since $v = O(N^{-1/4})$ and $dI_4/dt = O(1)$, this proves

(4.61)
$$\left\langle \frac{\mathrm{d}(\mathbb{E}H_t^w - \mathrm{i}\eta_t)}{\mathrm{d}t} M_t^w \right\rangle = -\frac{\mathrm{d}}{\mathrm{d}t} \langle \log |\mathring{M}_t^w| \rangle + O(N^{-3/2}\log N),$$

as desired.

Finally, to see the harmonicity, notice from $\|\mathcal{A}_t^{-1}\| \leq C$ that $\min_i |\sigma_i(\mathcal{A}_t)| \geq 1/C$, so that the map

(4.62)
$$w \mapsto \langle \log | \mathring{M}_t^w | \rangle = \frac{1}{N} \sum_{i=1}^N \log |\sigma_i(\mathcal{A}_t) - N^{-1/2} \gamma_t^{-1} w|$$

is harmonic on $\{w: |w| \le C\sqrt{N}|\gamma_t|\}.$

In the last bit of this section, we further reduce the proof of Proposition 4.4 into an estimate of its time derivative. To this end, we introduce the following notation¹³ for simplicity: For an index set $I = \{i_1, \ldots, i_n\} \subset [\![k]\!]$, we write

$$\boldsymbol{w}^{(I)} \equiv \boldsymbol{w}^{(i_1,\dots,i_n)} \coloneqq (w_j)_{j \notin I} \in \mathbb{C}^{||\boldsymbol{k}|| \setminus I}, \qquad \boldsymbol{w} \coloneqq \boldsymbol{w}^{\emptyset},$$

$$(4.63) \qquad \boldsymbol{L}_t^{(I)}(\boldsymbol{w}^{(I)}) \equiv \boldsymbol{L}_t^{(i_1,\dots,i_n)}(\boldsymbol{w}^{(I)}) \coloneqq \prod_{\substack{\ell=1,\\\ell \notin I}}^k L_t(w_\ell), \qquad \boldsymbol{L}_t(\boldsymbol{w}) \coloneqq \boldsymbol{L}_t^{(\emptyset)}(\boldsymbol{w}).$$

Now we compute the time derivative of L_t , with the help of Lemma 4.8, as follows:

Lemma 4.9. For any fixed $\epsilon > 0$, the following holds uniformly over $t \in [0,1]$ and $|w|, ||w|| \leq C$:

(4.64)
$$\mathbb{E}\frac{\mathrm{d}\boldsymbol{L}_t(\boldsymbol{w})}{\mathrm{d}t} = \sum_{j=1}^k \mathbb{E}\operatorname{Tr}\left[\frac{\mathrm{d}(\mathbb{E}H_t^{w_j} - \mathrm{i}\eta_t)}{\mathrm{d}t}G_t^{w_j}(\mathrm{i}\eta_t)\right]\boldsymbol{L}_t^{(j)}(\boldsymbol{w}^{(j)}) + h_t(\boldsymbol{w}) + O(N^{-1/2+\epsilon}),$$

where $\Delta_{w_1} \cdots \Delta_{w_k} h_t \equiv 0$.

Proof. From the first-order perturbation theory

(4.65) $\log \det(1 + dY) = \operatorname{Tr} \log(1 + dY) = \operatorname{Tr} dY + O(||dY||^2), \quad dY \in \mathbb{C}^{2N \times 2N},$ we immediately have

(4.66)
$$\frac{\mathrm{d}L_t(w)}{\mathrm{d}t} = \mathrm{Tr} \, \frac{\mathrm{d}(H_t^w - \mathrm{i}\eta_t)}{\mathrm{d}t} G_t^w - 2N \frac{\mathrm{d}\langle \log | \boldsymbol{h}_t^w - \mathrm{i}\eta_t | \rangle_{\mathcal{M}} \rangle}{\mathrm{d}t} \\ = \mathrm{Tr} \, \frac{\mathrm{d}(\mathbb{E}H_t^{w_j} - \mathrm{i}\eta_t)}{\mathrm{d}t} G_t^w - 2N \frac{\mathrm{d}\langle \log | \mathring{M}_t^w | \rangle}{\mathrm{d}t} + O(N^{-1/2})$$

¹³In Lemma 4.9 we only need |I| = 1, but |I| can reach (at most) five later in Sections 5 and 6.



Figure 2. Eigenvalues of the path $A_t + X$ and the evolution z_t of the base point

where in the second line we used (4.43) and Lemma 4.8. On the other hand, recall from (4.49) that $L_t(w) \prec 1$.

Therefore, combining with Lemma 4.8, we get

(4.67)
$$\mathbb{E}\frac{\mathrm{d}}{\mathrm{d}t}\boldsymbol{L}_{t}(\boldsymbol{w}) = \mathbb{E}\sum_{j=1}^{k} \left(\mathrm{Tr}\,\frac{\mathrm{d}(\mathbb{E}H_{t}^{w_{j}} - \mathrm{i}\eta_{t})}{\mathrm{d}t} G_{t}^{w_{j}} - 2N\frac{\mathrm{d}\langle \log|\mathring{M}_{t}^{w_{j}}|\rangle}{\mathrm{d}t} \right) \boldsymbol{L}_{t}^{(j)}(\boldsymbol{w}^{(j)}) + O(N^{-1/2+\epsilon}).$$

Denote the sum of the second terms in the summation in (4.67) by $h_t(\boldsymbol{w})$. Notice that each of these summands is harmonic in w_j by Lemma 4.8. This proves $\Delta_{w_1} \cdots \Delta_{w_k} h_t \equiv 0$.

5. Proof of Proposition 4.4, the complex case

This section is devoted to the proof of Proposition 4.4 when $\mathbb{F} = \mathbb{C}$. Since we only consider deformed complex Ginibre ensembles, we omit the superscripts Gin and Gin(\mathbb{C}) throughout the section. We start with the construction of the path \mathcal{A}_t .

Proposition 5.1. If z_0 satisfies Assumption 2.6 (i), (ii), and (iii \mathbb{C}), then there exist a path \mathcal{A}_t satisfying Assumption 4.6, with the constant C_5 in (4.35) depending only on the constants from Assumption 2.6.

Proof. Recall the definition of the domain \mathcal{D}_C from Assumption 2.6. Notice from $||A|| \leq C$ that

(5.1)
$$||(A-z)^{-1} + z^{-1}|| = O(|z|^{-2}), \quad \langle (A-z)|A-z|^{-4} \rangle = -z|z|^{-4} + O(|z|^{-4}), \quad \text{as } |z| \to \infty.$$

This in particular implies that for large enough C'' > 0 the complementary ball $\{z : |z| \ge C''\}$ is in $\mathcal{D}_{C'}$, so that $\mathcal{D}_{C'}$ has a unique unbounded component. We define the path $[0,1) \ni t \mapsto \hat{z}_t \in \mathcal{D}_{C'}$ as follows.

- (a) For $0 \le t \le 1/2$, the smooth path \hat{z}_t connects z_0 to $z_{1/2} = C'' z_1$
- (b) For $1/2 \le t < 1$, the path \hat{z}_t is given by $\hat{z}_t = C'' z_1/(1-t)$.

Note that the existence of the path connecting z_0 to 2C'' follows from Assumption 2.6 (iii) and that $\{z : |z| \ge C''\} \subset \mathcal{D}_{C'}$. Finally, define

(5.2)
$$\mathcal{A}_t := \langle |A - \hat{z}_t|^{-2} \rangle^{1/2} (A - \hat{z}_t), \quad t \in (0, 1), \qquad \mathcal{A}_1 := -1$$

The first two conditions in Assumption 4.6 are immediate from the definition of \mathcal{A}_t . For any given (small) $\delta > 0$, the estimates in (4.35) are trivial for $t \in [0, 1 - \delta)$ from the definition of $\mathcal{D}_{C'}$ and $|\hat{z}_t| = O(1)$. When t is sufficiently close to 1, the first three estimates in (4.35) are consequences of the large-|z|-asymptotics (5.1) and

(5.3)
$$||A - z|| = |z| + O(1), \quad \langle |A - z|^{-2} \rangle = |z|^{-2} + O(|z|^{-3}).$$

For the derivative of \mathcal{A}_t when t is close to 1, since \hat{z}_t follows the ray $z_1 \cdot (0, \infty)$ with $|z_1| = 1$, we get

(5.4)
$$\left\| \frac{\mathrm{d}}{\mathrm{d}t} \mathcal{A}_t \right\| = \left| \frac{\mathrm{d}\hat{z}_t}{\mathrm{d}t} \right| \left\| \langle |A - \hat{z}_t|^{-2} \rangle^{-1/2} \operatorname{Re} \left\langle (A - \hat{z}_t)^* \frac{-z_1}{|A - \hat{z}_t|^{-4}} \right\rangle (A - \hat{z}_t) + \langle |A - \hat{z}_t|^{-2} \rangle^{1/2} z_1 \right|$$
$$= \left| \frac{\mathrm{d}\hat{z}_t}{\mathrm{d}t} \right| O(|\hat{z}_t|^{-2}) (1 + ||A||) = O(1).$$

This concludes the proof of Proposition 5.1.

With (4.41), Lemma 4.9, and Proposition 5.1, we immediately find that the next result implies Theorem 4.2 when $\mathbb{F} = \mathbb{C}$.

Proposition 5.2. Let $\mathbb{F} = \mathbb{C}$, $k \in \mathbb{N}$, C > 0, and $\epsilon > 0$ be fixed. Then the following holds uniformly over \mathcal{A}_t satisfying Assumption 4.6, $t \in [0, 1]$, and $|w|, ||w|| \leq C$:

(5.5)
$$\sum_{j=1}^{k} \mathbb{E}\left\langle \frac{\mathrm{d}(\mathbb{E}H_t^{w_j} - \mathrm{i}\eta_t)}{\mathrm{d}t} G_t^{w_j}(\mathrm{i}\eta_t) \right\rangle \boldsymbol{L}_t^{(j)}(\boldsymbol{w}^{(j)}) = h_t(\boldsymbol{w}) + O(N^{\epsilon}\Psi^5)$$

(5.6)
$$\frac{\mathrm{d}}{\mathrm{d}t}\mathbb{E}\frac{\langle G_t^{-}\rangle}{c_t} = O(N^{1+\epsilon}\Psi^6),$$

where $\Delta_{w_1} \cdots \Delta_{w_k} h_t \equiv 0$.

The rest of this section is devoted to the proof of Proposition 5.2, and all estimates depend only on its parameters k, C, ϵ , and C_5 from Assumption 4.6.

5.1. General framework. Recall that in (4.56) we Taylor-expanded the matrix M_t^w in the trace

(5.7)
$$\left\langle \frac{\mathrm{d}\mathbb{E}H_t^w}{\mathrm{d}t} M_t^w \right\rangle$$

around M_t^w with respect to the *scalar* v^2 . The main goal of this section is to emulate the same expansion for (5.5). While all our estimates shall be uniform over $t \in [0, 1]$, we will consider t to be fixed throughout this section. Henceforth we abbreviate $\eta \equiv \eta_t$, $G^{w_j} \equiv G_t^{w_j}$, $M^{w_j} \equiv M_t^{w_j}$, etc. Also, we omit the spectral parameters of G^w and M^w when they are $i\eta$.

Recall the definition of W from (4.38). In what follows, we often use Stein's lemma. For a function $f: \mathbb{C}^{2N \times 2N} \to \mathbb{C}^{2N \times 2N}$, Stein's lemma states

(5.8)
$$\mathbb{E}Wf(W) = \mathbb{E}\mathbb{E}_{\widetilde{W}}W(\partial_{\widetilde{W}}f(W)),$$

where \widetilde{W} is an i.i.d. copy of W (hence has Gaussian entries) and $\partial_{\widetilde{W}}$ denotes the directional partial derivative along \widetilde{W} . As a simple consequence of (5.8) together with the identity

(5.9)
$$G - M = M(M^{-1} - G^{-1})G = M(-W - \langle M \rangle)G$$

we have

(5.10)
$$\mathbb{E}[G-M] = \mathbb{E}[M\mathcal{S}[G-M]G],$$

where $\mathcal{S}: \mathbb{C}^{2N \times 2N} \to \mathbb{C}^{2N \times 2N}$ is defined by

(5.11)
$$\mathcal{S}[B] := \mathbb{E}_{\widetilde{W}} \widetilde{W} B \widetilde{W} = \begin{pmatrix} \langle 2E_2 B \rangle & 0 \\ 0 & \langle 2E_1 B \rangle \end{pmatrix} = \sum_{E=E_1, E_2} E \langle 2\check{E}B \rangle.$$

Here, the block diagonal matrices E_1, E_2 are defined in (1.3) and the check denotes the involution $\check{E}_1 := E_2, \check{E}_2 = E_1$. Also note the identity

(5.12)
$$\mathbb{E}_{\widetilde{W}}\widetilde{W}\operatorname{Tr}[\widetilde{W}B] = \frac{1}{N}\sum_{E=E_1,E_2} EB\check{E}.$$

For the matrix $\mathring{M} = (\mathbb{E}H)^{-1}$, we have the following variant of (5.9);

(5.13)
$$G - \mathring{M} = \mathring{M}(\mathbb{E}H - G^{-1})G = \mathring{M}(i\eta - W)G$$

Note also that

(5.14)
$$\left\| \mathring{M}^{w} - \left(\mathring{M}^{0} + \mathring{M}^{0} J^{w} \mathring{M}^{0} \right) \right\| = O(\|J^{w}\|^{2}) = O(N^{-1}),$$

where J^w was defined in (4.38).

Our 'Taylor expansion' expresses (5.5) as a deterministic linear combination with each term of the form

(5.15)
$$\mathbb{E}\langle GE\cdots GE\rangle\cdots\langle GE\cdots GE\rangle \boldsymbol{L}^{(I)},$$

for $G = G^{w_i}$ with varying *i*'s, $E \in \{E_1, E_2\}$, and $I \subset [\![k]\!]$. We refer to such an expression as 'decoupled', in the sense that it involves only the plain trace $\langle \cdot \rangle$ as opposed to a generic weighted trace $\langle B \cdot \rangle$. To rigorously formulate our method, we introduce a few notations. In what follows, bold-faced symbols w, and F always denote finite tuples (with appropriate dimensions) of elements of

(5.16)
$$\{w_1, \dots, w_k\}$$
 and $\{E_1, E_2\},\$

respectively. Here w_i 's are free (in the literal sense) complex variables (e.g. in Lemma 5.6), unless specified to be in some subset of \mathbb{C} (e.g. in Lemma 5.5).

Definition 5.3. For $n \in \mathbb{N}$, $\boldsymbol{w} = (\widetilde{w}_j)_{j \in [\![n]\!]}$, and $\boldsymbol{F} = (F_j)_{j \in [\![n]\!]}$, define

(5.17)
$$P_n(\boldsymbol{w}; \boldsymbol{F}) := \frac{1}{N^{n-1}} G^{\widetilde{w}_1} F_1 \cdots G^{\widetilde{w}_n} F_n - \mathbb{1}(n=1) \mathring{M}^{\widetilde{w}_1} F_1.$$

Similarly, for a multi-index $\mathbf{m} \equiv (m_j)$ in \mathbb{N} , $\mathbf{w} \equiv (\mathbf{w}_j)$, and $\mathbf{F} \equiv (\mathbf{F}_j)$, define

(5.18)
$$\boldsymbol{Q}_{\boldsymbol{m}}(\boldsymbol{w};\boldsymbol{F}) := \prod_{j} \langle P_{m_j}(\boldsymbol{w}_j;\boldsymbol{F}_j) \rangle, \qquad |\boldsymbol{m}| := \sum_{j} m_j,$$

with the convention $Q_m \equiv 1$ for |m| = 0. For $\ell \in \mathbb{N} \cup \{0\}$ define

(5.19)
$$\boldsymbol{L}_{\ell}(\boldsymbol{w}) := \prod_{j=1}^{\iota} L(\widetilde{w}_j).$$

We often omit the subscript ℓ to write $\widetilde{L} \equiv \widetilde{L}_{\ell}$. Finally, $Q_m^{(I)}$ and $\widetilde{L}^{(I)}$ are defined in the same fashion as (4.63).

Definition 5.4. For $n \in \mathbb{N}$, we define \mathcal{X}_n by the set of expressions

(5.20)
$$\{ \boldsymbol{Q}_{\boldsymbol{m}}(\boldsymbol{w}_1; \boldsymbol{F}_1) \widetilde{\boldsymbol{L}}_{\ell}(\boldsymbol{w}_2) : |\boldsymbol{m}| = n, \ell \ge 0, \boldsymbol{w}_1, \boldsymbol{w}_2, \boldsymbol{F}_1 \}$$

where the components of w_1, w_2 , and F_1 may vary among the symbols in (5.16). For a given matrix $B \in \mathbb{C}^{N \times N}$ and $n \in \mathbb{N}$, we define $\mathcal{Y}_n(B)$ to be the set of expressions

(5.21)
$$\{\langle BP_{n_1}(\boldsymbol{w}_1; \boldsymbol{F}_1) \rangle \boldsymbol{Q}_{\boldsymbol{m}}(\boldsymbol{w}_2; \boldsymbol{F}_2) \boldsymbol{L}_{\ell}(\boldsymbol{w}_3): n_1 + |\boldsymbol{m}| = n, \ell \ge 0, \boldsymbol{w}_1, \boldsymbol{w}_2, \boldsymbol{w}_3, \boldsymbol{F}_1, \boldsymbol{F}_2\},\$$

where the components of w_1, w_2, w_3, F_1 , and F_2 may vary among the symbols in (5.16). Notice that $\mathcal{Y}_n(I) = \mathcal{X}_n$. With a slight abuse of notations, we write \mathfrak{X}_n (resp. $\mathfrak{Y}_n(B)$) for generic finite sums (with the number of summand independent of N) of expressions in \mathcal{X}_n (resp. $\mathcal{Y}_n(B)$), whose exact form may vary by lines.

The notations \mathfrak{X}_n and $\mathfrak{Y}_n(B)$ in Definition 5.4 are introduced purely for the sake of brevity. For all instances of \mathfrak{X}_n and $\mathfrak{Y}_n(B)$, we can track down their exact forms as sums over elements of \mathcal{X}_n or $\mathcal{Y}_n(B)$. However we do not write them down simply because the explicit formulas play no role in the proof while they are excessively lengthy.

We may now present the two inputs that form the basis of our expansion: Their proofs are postponed to the end of this section.

Lemma 5.5. For any fixed $B \in \mathbb{C}^{N \times N}$ and $n \in \mathbb{N}$, we have

(5.22)
$$|\widetilde{\boldsymbol{L}}_{n}(\boldsymbol{w})| \prec 1, \qquad |\langle BP_{n}(\boldsymbol{w};\boldsymbol{F})\rangle| \prec ||B|| \Psi'$$

uniformly over (w_1, \ldots, w_k) in a compact set.

Lemma 5.6 (Decoupling). Let B_d and $B_o \in \mathbb{C}^{N \times N}$ be supported respectively on diagonal and off-diagonal blocks. Then

$$\mathbb{E}\langle B_d P_n \rangle \boldsymbol{Q}_m \widetilde{\boldsymbol{L}} = \langle 2B_d \mathring{M}^{\widetilde{w}_1} \mathring{M}^{\widetilde{w}_n} F_n \rangle \mathbb{E} \langle P_n \rangle \boldsymbol{Q}_m \widetilde{\boldsymbol{L}} + \left(\mathbbm{1}(n=1) \mathrm{i} \eta_t \langle B_d (\mathring{M}^{\widetilde{w}_1})^2 F_1 \rangle + \mathbbm{1}(n=2) \frac{1}{N} \langle B_d \mathring{M}^{\widetilde{w}_1} F_1 \mathring{M}^{\widetilde{w}_2} F_2 \rangle \right) \mathbb{E} \boldsymbol{Q}_m \widetilde{\boldsymbol{L}} + \frac{1}{2} \mathbb{E} \mathfrak{Y}_{|\boldsymbol{m}|+n+1} (B_d \mathring{M}^{\widetilde{w}_1})$$

(5.24)

$$\begin{aligned}
& + i\eta \mathbb{E} \mathfrak{Y}_{|\boldsymbol{m}|+n} (B_d \mathring{M}^{\widetilde{w}_1}) + \frac{1}{N} \mathbb{E} \mathfrak{Y}_{|\boldsymbol{m}|+n-1} (B_d \mathring{M}^{\widetilde{w}_1}), \\
& = \frac{1}{2} \mathbb{E} \mathfrak{Y}_{|\boldsymbol{m}|+n+1} (B_o \mathring{M}^{\widetilde{w}_1}) \\
& + i\eta \mathbb{E} \mathfrak{Y}_{|\boldsymbol{m}|+n} (B_o \mathring{M}^{\widetilde{w}_1}) + \frac{1}{N} \mathbb{E} \mathfrak{Y}_{|\boldsymbol{m}|+n-1} (B_o \mathring{M}^{\widetilde{w}_1}),
\end{aligned}$$

where we abbreviated $P_n \equiv P_n(\widetilde{w}_1, \ldots, \widetilde{w}_n; \mathbf{F})$. The exact forms of the expressions \mathfrak{Y} on right-hand sides do not depend on $B_d(B_o, \text{ resp.})$ except that they are supported on diagonal (off-diagonal, resp.) blocks.

While we can formally recover (5.24) from (5.23) upon plugging in $B_d = B_o$, we separated the two identities since \mathfrak{Y} on the right-hand side depends on the support of B_d or B_o .

As an immediate consequence of Lemma 5.5, we have

(5.25)
$$|\mathbb{E}\mathfrak{X}_n| \lesssim N^{\epsilon}\Psi^n$$
, $|\mathbb{E}\mathfrak{Y}_n(B_d)| \lesssim N^{\epsilon} ||B_d||\Psi^n$, and $|\mathbb{E}\mathfrak{Y}_n(B_o)| \lesssim N^{\epsilon} ||B_o||\Psi^{n+1}$.

In this regard, Lemma 5.6 decouples the expression on the left-hand side to the leading order, but does not provide a useful estimate in terms of a fully deterministic quantity. This in fact is the general theme of our proof; say, we never try to estimate $\mathbb{E}Q_m$ but rather claim that the contribution of each $\mathbb{E}Q_m$ cancels out algebraically.

Notice that $B_d \mathring{M}^{w_1}$ and $B_o \mathring{M}^{w_1}$ are supported respectively on off-diagonal and diagonal blocks. Thus, if necessary, one may iteratively apply Lemma 5.6 to gain an extra precision of Ψ at each step. Indeed, this will be our main task in the next section; we start from the case when $|\mathbf{m}| = 0$ and n = 1, more precisely

(5.26)
$$\mathbb{E}\left\langle \frac{\mathrm{d}\mathbb{E}H_t^w}{\mathrm{d}t}G_t^w \right\rangle \boldsymbol{L}^{(j)},$$

and apply (a more explicit version of) Lemma 5.6 four times so that the resulting decoupled expression has the precision Ψ^5 . Indeed, the result of the last three consecutive applications of Lemma 5.6 can be combined into the following generic lemma. We prove the lemma at the end of this section.

Lemma 5.7. Let $n \in \mathbb{N}$ and m be a multi-index with n + |m| = 2, and $B_d \in \mathbb{C}^{2N \times 2N}$ be supported on diagonal blocks. Then we have, uniformly over (w_1, \ldots, w_k) in a compact set,

$$\mathbb{E}\langle B_d P_n \rangle \boldsymbol{Q_m} \boldsymbol{L} = \langle 2B_d \mathring{M}^{\widetilde{w}_1} \mathring{M}^{\widetilde{w}_n} F_n \rangle \mathbb{E}\langle P_n \rangle \boldsymbol{Q_m} \boldsymbol{L} + \left(\mathbbm{1}(n=1) \mathrm{i}\eta_t \langle B_d (\mathring{M}^{\widetilde{w}_1})^2 F_1 \rangle + \mathbbm{1}(n=2) \frac{1}{N} \langle B_d \mathring{M}^{\widetilde{w}_1} F_1 \mathring{M}^{\widetilde{w}_2} F_2 \rangle \right) \mathbb{E} \boldsymbol{Q_m} \widetilde{\boldsymbol{L}} + \frac{1}{2} \langle B_d (\mathring{M}^0)^4 F_n \rangle \mathbb{E} \mathfrak{X}_4 + O(N^{\epsilon} \| B_d \| \Psi^5),$$

where $P_n = P_n(\widetilde{w}_1, \ldots, \widetilde{w}_n; \mathbf{F})$ and the expression \mathfrak{X}_4 does not depend on B_d .

Taking $B_d = I$, we can solve (5.27) for the unspecified expression $\mathbb{E}\mathfrak{X}_4$. Consequently, we have the following immediate corollary of Lemma 5.7. We omit the proof and only point out that

$$\langle 2(\mathring{M}^0)^4 E_i \rangle = \langle |\mathcal{A}_t|^{-4} \rangle = I_4 \sim 1.$$

Lemma 5.8. Under the same notations as in Lemma 5.7, the following holds uniformly over (w_1, \ldots, w_k) in a compact set: (5.28)

$$\begin{aligned} \mathbb{E}\langle B_{d}P_{n}\rangle \boldsymbol{Q_{m}}\widetilde{\boldsymbol{L}} &= \left(\langle 2B_{d}\mathring{M}^{\widetilde{w}_{1}}\mathring{M}^{\widetilde{w}_{n}}F_{n}\rangle - (\langle 2\mathring{M}^{\widetilde{w}_{1}}\mathring{M}^{\widetilde{w}_{n}}F_{n}\rangle - 1)\frac{\langle 2B_{d}(\mathring{M}^{0})^{4}F_{n}\rangle}{I_{4}}\right)\mathbb{E}\langle P_{n}\rangle \boldsymbol{Q_{m}}\widetilde{\boldsymbol{L}} \\ &+ \mathbb{1}(n=1)\left(\langle B_{d}(\mathring{M}^{\widetilde{w}_{1}})^{2}F_{1}\rangle - \langle (\mathring{M}^{\widetilde{w}_{1}})^{2}F_{1}\rangle\frac{\langle 2B_{d}(\mathring{M}^{0})^{4}F_{n}\rangle}{I_{4}}\right)\mathrm{i}\eta_{t}\mathbb{E}\boldsymbol{Q_{m}}\widetilde{\boldsymbol{L}} \\ &+ \mathbb{1}(n=2)\left(\langle B_{d}\mathring{M}^{\widetilde{w}_{1}}F_{1}\mathring{M}^{\widetilde{w}_{2}}F_{2}\rangle - \langle \mathring{M}^{\widetilde{w}_{1}}F_{1}\mathring{M}^{\widetilde{w}_{2}}F_{2}\rangle\frac{\langle 2B_{d}(\mathring{M}^{0})^{4}F_{n}\rangle}{I_{4}}\right)\frac{1}{N}\mathbb{E}\boldsymbol{Q_{m}}\widetilde{\boldsymbol{L}} \\ &+ O(N^{\epsilon}\|B_{d}\|\Psi^{5}), \end{aligned}$$

5.2. Proof of Proposition 5.2. We first prove (5.5) and then (5.6).

Proof of Proposition 5.2, (5.5). By (5.13) and Stein's lemma, we obtain (recall $\mathring{M} = (\mathbb{E}H)^{-1}$)

$$\begin{aligned}
& \left\{ \frac{\mathrm{d}\mathbb{E}H_{t}^{w_{j}}}{\mathrm{d}t}G^{w_{j}}E_{1} \right\} \mathbf{L}^{(j)} \\
&= \left\langle \frac{\mathrm{d}\mathbb{E}H_{t}^{w_{j}}}{\mathrm{d}t}\mathring{M}^{w_{j}}E_{1} \right\rangle \mathbb{E}\mathbf{L}^{(j)} + \mathrm{i}\eta_{t}\mathbb{E}\left\langle \frac{\mathrm{d}\mathbb{E}H_{t}^{w_{j}}}{\mathrm{d}t}\mathring{M}^{w_{j}}G^{w_{j}}E_{1} \right\rangle \mathbf{L}^{(j)} \\
&= \left\langle E\left\langle \frac{\mathrm{d}\mathbb{E}H_{t}^{w_{j}}}{\mathrm{d}t}\mathring{M}^{w_{j}}WG^{w_{j}}E_{1} \right\rangle \mathbf{L}^{(j)} \\
&= \left\langle B_{d}^{w_{j}}E_{1} \right\rangle \mathbb{E}\mathbf{L}^{(j)} + \mathrm{i}\eta_{t}\mathbb{E}\left\langle B_{d}^{w_{j}}G^{w_{j}}E_{1} \right\rangle \mathbf{L}^{(j)} \\
&+ \mathbb{E}\left\langle G^{w_{j}} \right\rangle \left\langle B_{d}^{w_{j}}G^{w_{j}}E_{1} \right\rangle \mathbf{L}^{(j)} - \frac{1}{N}\sum_{\ell}^{(j)} \mathbb{E}\left\langle B_{d}^{w_{j}}G^{w_{\ell}}E_{2}G^{w_{j}}E_{1} \right\rangle \mathbf{L}^{(j,\ell)},
\end{aligned}$$

where we abbreviated

$$B_d^{w_j} \coloneqq \frac{\mathrm{d}\mathbb{E}H_t^{w_j}}{\mathrm{d}t} \mathring{M}^{w_j}$$

and used for the last term that $B_d^{w_j}$ and E_i commute. Then we directly apply Lemma 5.6 to the second term of (5.29) with the choices $B_d = B_d^{w_j}$, $P_n = P_1(w_j; E_1)$, and $\tilde{\boldsymbol{L}} = \boldsymbol{L}^{(j)}$, so that

$$\mathbb{E}\left\langle 2\frac{\mathrm{d}\mathbb{E}H_{t}^{w_{j}}}{\mathrm{d}t}G^{w_{j}}E_{1}\right\rangle \mathbf{L}^{(j)} \\
=\left\langle 2B_{d}^{w_{j}}E_{1}\right\rangle \mathbb{E}\mathbf{L}^{(j)} + \mathrm{i}\eta\left\langle 2B_{d}^{w_{j}}(\mathring{M}^{w_{j}})^{2}E_{1}\right\rangle \mathbb{E}\langle G^{w_{j}}\rangle \mathbf{L}^{(j)} \\
+ \mathbb{E}\langle G^{w_{j}}\rangle\left\langle 2B_{d}^{w_{j}}G^{w_{j}}E_{1}\right\rangle \mathbf{L}^{(j)} - \frac{1}{N}\sum_{\ell}^{(j)}\mathbb{E}\left\langle 2B_{d}^{w_{j}}G^{w_{\ell}}E_{2}G^{w_{j}}E_{1}\right\rangle \mathbf{L}^{(j,\ell)} + O(N^{\epsilon}\Psi^{5}),$$

where we used $\eta = O(\Psi^3)$ and $N = O(\Psi^4)$.

Likewise, applying Lemma 5.8 to the remaining two undecoupled terms in (5.30), we find that

$$\begin{split} \mathbb{E} \left\langle 2 \frac{\mathrm{d}(\mathbb{E} H_{t}^{w_{j}} - \mathrm{i}\eta_{t})}{\mathrm{d}t} G^{w_{j}} E_{1} \right\rangle \mathbf{L}^{(j)} \\ &= \left\langle 2B_{d}^{w_{j}} E_{1} \right\rangle \mathbb{E} \mathbf{L}^{(j)} \\ &+ \left(\left\langle 2B_{d}^{w_{j}} (\mathring{M}^{w_{j}})^{2} E_{1} \right\rangle - \frac{\left\langle 2B_{d}^{w_{j}} (\mathring{M}^{0})^{4} E_{1} \right\rangle}{I_{4}} \left(\left\langle (\mathring{M}^{w_{j}})^{2} \right\rangle - 1 \right) \right) \mathbb{E} \langle G^{w_{j}} \rangle^{2} \mathbf{L}^{(j)} \\ (5.31) &+ \left(\left\langle 4B_{d}^{w_{j}} (\mathring{M}^{0})^{2} E_{1} \right\rangle - \frac{\left\langle 2B_{d}^{w_{j}} (\mathring{M}^{0})^{4} E_{1} \right\rangle}{I_{4}} + \frac{\mathrm{d}c_{t}}{\mathrm{d}t} \frac{1}{c_{t}} \right) \mathrm{i}\eta \mathbb{E} \langle G^{w_{j}} \rangle \mathbf{L}^{(j)} \\ &+ \sum_{\ell}^{(j)} \left(\left\langle 2B_{d}^{w_{j}} \mathring{M}^{w_{\ell}} \mathring{M}^{w_{j}} E_{1} \right\rangle - \frac{\left\langle 2B_{d}^{w_{j}} (\mathring{M}^{0})^{4} E_{1} \right\rangle}{I_{4}} \left(\left\langle 2\mathring{M}^{w_{\ell}} \mathring{M}^{w_{j}} E_{1} \right\rangle - 1 \right) \right) \mathbb{E} \frac{\left\langle G^{w_{\ell}} E_{2} G^{w_{j}} E_{1} \right\rangle}{N} \mathbf{L}^{(j,\ell)} \\ &+ \sum_{\ell}^{(j)} \frac{1}{2N} \left(\left\langle 2B_{d}^{w_{j}} (\mathring{M}^{w_{j}})^{2} E_{1} \right\rangle - \frac{\left\langle 2B_{d}^{w_{j}} (\mathring{M}^{0})^{4} E_{1} \right\rangle}{I_{4}} \right) \mathbb{E} \mathbf{L}^{(j,\ell)} + O(N^{\epsilon} \Psi^{5}), \end{split}$$

where we also used the definition $\eta_t = c_t^{-1} \eta_1$.

In what follows, we prove that the real part of (5.31) is harmonic (up to some errors) after summing over j = 1, ..., k. We plug in the definition of B_d^w to compute the deterministic prefactors. For the first term, we simply have (recall $\mathring{M} = (\mathbb{E}H)^{-1}$)

(5.32)
$$\operatorname{Re}\langle 2B_d^{w_j}E_1\rangle = \left\langle \frac{\mathrm{d}\mathbb{E}H_t^{w_j}}{\mathrm{d}t}\mathring{M}_t^{w_j}\right\rangle = -\frac{\mathrm{d}}{\mathrm{d}t}\langle \log|\mathring{M}_t^{w_j}|\rangle,$$

which is harmonic in w_j by Lemma 4.8. Secondly, we have from (4.58) that

(5.33)
$$\operatorname{Re}\langle 2B_{d}^{w}(\mathring{M}^{w})^{2}E_{1}\rangle = \operatorname{Re}\left\langle 2\frac{\mathrm{d}\mathbb{E}H_{t}^{w}}{\mathrm{d}t}(\mathring{M}_{t}^{w})^{3}E_{1}\right\rangle = \left\langle \frac{\mathrm{d}\mathbb{E}H_{t}^{w}}{\mathrm{d}t}(\mathring{M}_{t}^{w})^{3}\right\rangle = -\frac{1}{2}\frac{\mathrm{d}}{\mathrm{d}t}\left\langle (\mathring{M}_{t}^{w})^{2}\right\rangle$$
$$= -\frac{1}{2}\frac{\mathrm{d}I_{4}}{\mathrm{d}t}I_{4}^{-1/2}\operatorname{Re}[-N^{-1/2}w] + O(N^{-1}\log N).$$

Likewise we have

(5.34)
$$\operatorname{Re}\langle 2B_d^{w_j}(\mathring{M}^0)^4 E_1 \rangle = \left\langle \frac{\mathrm{d}\mathbb{E}H_t^w}{\mathrm{d}t} \mathring{M}_t^w (\mathring{M}^0)^4 \right\rangle = -\frac{1}{4} \frac{\mathrm{d}}{\mathrm{d}t} I_4 + O(N^{-1/2} \log N).$$

Plugging (4.58), (5.33), and (5.34), we have

so that the second term of (5.31) is $O(\Psi^6)$. For the third term of (5.31), we combine $c_t = I_4^{-1/4}$, (5.33) up to the leading order, and (5.34) to obtain

(5.36)
$$\langle 4B_d^{w_j}(\mathring{M}^0)^2 E_1 \rangle - \frac{\langle 2B_d^{w_j}(\mathring{M}^0)^4 E_1 \rangle}{I_4} + \frac{\mathrm{d}c_t}{\mathrm{d}t} \frac{1}{c_t}$$
$$= O(N^{-1/2}) + \frac{1}{4I_4} \frac{\mathrm{d}I_4}{\mathrm{d}t} - \frac{1}{4I_4} \frac{\mathrm{d}I_4}{\mathrm{d}t} = O(N^{-1/2}) = O(\Psi^2).$$

Thus we conclude that the third term of (5.31) is $O(\Psi^6)$. The last term of (5.31) is constant hence harmonic with respect to the variable w_j . Now it only remains to deal with the fourth term of (5.31). Using the facts that

(5.37)
$$\begin{split} \|B_{d}^{w_{j}} - B_{d}^{0}\| &= O(N^{-1/2}) = O(\Psi^{2}),\\ &\left\langle 2\mathring{M}^{w_{j}}\mathring{M}^{w_{\ell}}E_{1}\right\rangle - 1 = O(N^{-1/2}) = O(\Psi^{2}),\\ &\left\langle 2B_{d}^{w_{j}}\mathring{M}^{w_{\ell}}\mathring{M}^{w_{j}}E_{1}\right\rangle = \left\langle 2\mathring{M}^{w_{j}}\frac{\mathrm{d}\mathbb{E}H_{t}^{w_{j}}}{\mathrm{d}t}\mathring{M}^{w_{j}}\mathring{M}^{w_{\ell}}E_{2}\right\rangle = -\left\langle 2\frac{\mathrm{d}\mathring{M}^{w_{j}}}{\mathrm{d}t}\mathring{M}^{w_{\ell}}E_{2}\right\rangle, \end{split}$$

each summand in the fifth term of (5.31) is equal to, up to an additive error of $O(\Psi^6)$,

(5.38)
$$\operatorname{Re}\left(-\left\langle 2\frac{\mathrm{d}\mathring{M}^{w_{j}}}{\mathrm{d}t}\mathring{M}^{w_{\ell}}E_{2}\right\rangle-\frac{\langle 2B_{d}^{0}(\mathring{M}^{0})^{4}E_{1}\rangle}{I_{4}}(\langle 2\mathring{M}^{w_{j}}\mathring{M}^{w_{\ell}}E_{2}\rangle-1)\right)\mathbb{E}\frac{\langle G^{w_{\ell}}E_{2}G^{w_{j}}E_{1}\rangle}{N}\boldsymbol{L}^{(j,\ell)}.$$

We take the sum of (5.38) over $j \neq \ell$ in the following order; for each fixed pair of indices $j > \ell$, take the sum of (5.38) with the same quantity with j and ℓ interchanged, and then we take the sum over all pairs $j > \ell$. Consequently, the sum becomes

$$2\operatorname{Re}\sum_{j>\ell} \left(-\left\langle \frac{\mathrm{d}(\mathring{M}^{w_j} \mathring{M}^{w_\ell})}{\mathrm{d}t} E_2 \right\rangle - \frac{\langle B^0_d(\mathring{M}^0)^4 \rangle}{I_4} \left(\langle 2\mathring{M}^{w_j} \mathring{M}^{w_\ell} E_2 \rangle - 1 \right) \right) \frac{\langle G^{w_\ell} E_2 G^{w_j} E_1 \rangle}{N} \boldsymbol{L}^{(j,\ell)},$$

where we used that

(5.39)
$$\left\langle \frac{\mathrm{d}\mathring{M}^{w_{\ell}}}{\mathrm{d}t}\mathring{M}^{w_{j}}E_{2} \right\rangle = \left\langle \mathring{M}^{w_{j}}\frac{\mathrm{d}\mathring{M}^{w_{\ell}}}{\mathrm{d}t}E_{2} \right\rangle, \qquad \overline{\langle \mathring{M}^{w_{\ell}}\mathring{M}^{w_{j}}E_{2} \rangle} = \langle \mathring{M}^{w_{j}}\mathring{M}^{w_{\ell}}E_{2} \rangle,$$
$$\overline{\langle B^{0}_{d}(\mathring{M}^{0})^{4}E_{1} \rangle} = \langle B^{0}_{d}(\mathring{M}^{0})^{4}E_{2} \rangle, \qquad \overline{\langle G^{w_{j}}E_{2}G^{w_{\ell}}E_{1} \rangle} = \langle G^{w_{\ell}}E_{2}G^{w_{j}}E_{1} \rangle,$$

due to \mathring{M}^w being Hermitian and $EG\check{E} = EG^*\check{E}$ for $E \in \{E_1, E_2\}$.

Now we notice that the prefactor can be written as

(5.40)
$$\left(-\left\langle \frac{\mathrm{d}(\mathring{M}^{w_j} \mathring{M}^{w_\ell})}{\mathrm{d}t} E_2 \right\rangle - \frac{\langle B^0_d(\mathring{M}^0)^4 \rangle}{I_4} \left(\langle 2\mathring{M}^{w_j} \mathring{M}^{w_\ell} E_2 \rangle - 1 \right) \right) = -\frac{1}{2} I_4^{1/2} \frac{\mathrm{d}}{\mathrm{d}t} \left[I_4^{-1/2} (\langle 2\mathring{M}^{w_j} \mathring{M}^{w_\ell} E_2 \rangle - 1) \right].$$

On the other hand, we have

(5.41)

$$\langle 2\mathring{M}^{w_j}\mathring{M}^{w_\ell}E_1\rangle = \langle 2(\mathring{M}^0)^2\rangle + \langle J^{w_j}(\mathring{M}^0)^3E_2\rangle + \langle J^{w_\ell}(\mathring{M}^0)^3E_1\rangle + O(N^{-1})$$

$$= 1 + N^{-1/2}\left(\overline{\gamma}^{-1}\overline{w}_j\overline{I}_3 + \gamma^{-1}w_\ell I_3\right) + O(N^{-1})$$

$$= 1 - I_4^{1/2}N^{-1/2}(w_j + w_\ell) + O(N^{-1}).$$

As before, the error remains $O(N^{-1} \log N)$ after taking the time derivative, hence we conclude

(5.42)
$$\frac{\mathrm{d}}{\mathrm{d}t} \left[I_4^{-1/2} (\langle 2 \mathring{M}^{w_j} \mathring{M}^{w_\ell} E_2 \rangle - 1) \right] = O(N^{-1} \log N).$$

Plugging this back into (5.40) and then (5.2), we conclude that the contribution of the fifth term of (5.31) is $O(N^{-1}\Psi^2) = O(N^{-1-c})$.

Proof of Proposition 5.2, (5.6). We first make the following two observations about (5.31) in the previous proof for j = 1. Firstly, upon close inspection, the $O(\Psi^5)$ error in (5.31) can be tracked down algebraically. Indeed, we may write the error as $\mathbb{E}\mathfrak{Y}$, where \mathfrak{Y} is a finite sum of terms of the following form:

(5.43)
$$C_{\boldsymbol{w}}N^{-k_1/2} \cdot \eta^{k_2}N^{-k_3}\mathfrak{Y}_n(B).$$

Here k_1 is either 0 or 1, k_2 and k_3 are nonnegative integers, ||B|| = O(1) and $C_w = O(1)$ are deterministic and independent of η , and

$$(5.44) 2k_1 + 3k_2 + 4k_3 + n \ge 5.$$

Secondly, the proof of (5.31) is insensitive to the choice $\eta_1 = N^{-3/4-\delta}$, but only requires that $\eta_1 \sim N^{-3/4-\delta}$.

Differentiating (5.6) in $i\eta$ (more precisely $\partial_{i\eta} = i^{-1}\partial_{\eta}$), we find that

(5.45)
$$\frac{\mathrm{d}}{\mathrm{d}t} \frac{\mathbb{E}\langle G_t \rangle}{c_t} = \partial_{\mathrm{i}\eta_t} \mathbb{E} \left\langle \frac{\mathrm{d}(\mathbb{E}H_t^w - \mathrm{i}\eta_t)}{\mathrm{d}t} G_t \right\rangle.$$

Our goal is to show that the right-hand side of (5.45) is $O(N^{1+\epsilon}\Psi^6)$.

Since (5.31) for j = 1 remains valid over varying η_1 , we may apply $\partial_{i\eta_t}$ so that

$$(5.46) \quad \partial_{\eta_t} \mathbb{E} \left\langle \frac{\mathrm{d}(\mathbb{E}H_t^w - \mathrm{i}\eta_t)}{\mathrm{d}t} G_t^w \right\rangle = \left(\langle B_d^w(\mathring{M}_t^w)^2 \rangle - \frac{\langle B_d^w(\mathring{M}_t^0)^4 \rangle}{I_4} \left(\langle (\mathring{M}_t^w)^2 \rangle - 1 \right) \right) \partial_{\eta} \mathbb{E} \langle G_t^w \rangle^2 \\ + \left(\langle 4B_d^w(\mathring{M}_t^0)^2 \rangle - \frac{\langle B_d^w(\mathring{M}_t^0)^4 \rangle}{I_4} + \frac{\mathrm{d}c_t}{\mathrm{d}t} \frac{1}{c_t} \right) \mathrm{i} \partial_{\eta_t} \left[\eta \mathbb{E} \langle G_t^w \rangle \right] + \partial_{\eta_t} \mathbb{E} \mathfrak{Y},$$

where we used that B_d^w does not depend on η , and \mathfrak{Y} is a finite sum of terms of the form given in (5.43). Then, notice that the derivative $\partial_{\eta_t} \mathfrak{Y}$ is a finite sum of

(5.47)
$$C_{\boldsymbol{w}}N^{-k_1/2}\eta^{k_2-1}N^{-k_3}\mathfrak{Y}_n(B), \quad C_{\boldsymbol{w}}N^{-k_1/2}\eta^{k_2}N^{-k_3+1}\mathfrak{Y}_{n+1}(B).$$

This in turn implies, by Lemma 5.5, that

(5.48)
$$\partial_{\eta_t} \mathbb{E} \mathfrak{Y} = O(N^{1+\epsilon} \Psi^6) = O(N^{-c} \Psi).$$

Finally, the result follows from substituting (5.34) and (5.35) into (5.46).

5.3. **Proof of Lemmas 5.5–5.7.** Throughout this section, we use the following shorthand notation; for a given $\boldsymbol{w} \in \mathbb{C}^n$ and $\boldsymbol{F} \in \{E_1, E_2\}^n$, we write

(5.49)
$$\widetilde{P}_n \equiv \widetilde{P}_n(\boldsymbol{w}, \boldsymbol{F}) \coloneqq N^{-n+1} G^{w_1} F_1 \cdots G^{w_n} F_n \equiv P_n + \mathbb{1}(n=1) \mathring{M}^{w_1} F_1,$$

and for a subset $I \subset \llbracket N \rrbracket$ write

(5.50)
$$P_I := P_{|I|}(\boldsymbol{w}_I; \boldsymbol{F}_i), \quad \widetilde{P}_I := \widetilde{P}_{|I|}(\boldsymbol{w}_I; \boldsymbol{F}_I), \quad \boldsymbol{w}_I = (w_i)_{i \in I}, \quad \boldsymbol{F}_I = (F_i)_{i \in I}.$$

Since $\langle \mathring{M}^w F \rangle = 0$, we have $\langle P_n \rangle = \langle \widetilde{P}_n \rangle$ and similarly $\mathcal{S}[P_n] = \mathcal{S}[\widetilde{P}_n].$

Proof of Lemma 5.5. For n = 1, (5.22) is an immediate consequence of the local law. To prove (5.22) for $n \ge 2$, we write (with the convention $\widetilde{P}_0 = N$)

(5.51)
$$\begin{aligned} |\langle BP_n(\boldsymbol{w};\boldsymbol{F})\rangle| &= \frac{1}{N^2} |\langle G^{w_n} F_n B \widetilde{P}_{[n-2]} G^{w_{n-1}} F_{n-1}\rangle| \\ &\leq \frac{1}{N} ||B \widetilde{P}_{[n-2]}|| \frac{\langle G^{w_n} \rangle^2 + \langle G^{w_{n-1}} \rangle^2}{N}, \end{aligned}$$

where we used the general inequality

(5.52)
$$|\langle B_1 B_2 B_3 \rangle| \le ||B_2||(\langle |B_1|^2 \rangle + \langle |B_2|^2 \rangle), \qquad B_1, B_2, B_3 \in \mathbb{C}^{2N \times 2N}.$$

Then by $\|\widetilde{P}_n\| \leq (N\eta)^{-n+1}\eta^{-1}$, we get

(5.53)
$$\frac{1}{N} \| \langle B\widetilde{P}_{\llbracket n-2 \rrbracket} \rangle \| \le \frac{\|B\|}{(N\eta)^{n-2}}$$

so that by Ward identity and local laws Proposition 4.7 it follows that

(5.54)
$$|\langle BP_n(\boldsymbol{w};\boldsymbol{F})\rangle| \le ||B|| \frac{\mathrm{Im}\langle G^{w_n}\rangle + \mathrm{Im}\langle G^{w_{n-1}}\rangle}{(N\eta)^{n-1}} \prec ||B||\Psi^n.$$

Proof of Lemma 5.6. We only prove (5.23), and (5.24) can be proved analogously by changing the parities of few F's. Expanding the first factor $G^{\tilde{w}_1}$ in $\langle B_d P_n \rangle$, we get

$$\mathbb{E}\langle B_{d}P_{n}\rangle\boldsymbol{Q_{m}}\widetilde{\boldsymbol{L}} = \mathbb{E}\left(-\langle B_{d}\mathring{M}^{w_{1}}W\widetilde{P}_{n}\rangle + \mathbb{1}(n\geq2)\frac{\langle B_{d}\mathring{M}^{w_{1}}F_{1}\widetilde{P}_{\llbracket 2,n \rrbracket}\rangle}{N} + i\eta_{t}\langle B_{d}\mathring{M}^{w_{1}}\widetilde{P}_{n}\rangle\right)\boldsymbol{Q_{m}}\widetilde{\boldsymbol{L}}$$

$$(5.55) = -\mathbb{E}\langle B_{d}\mathring{M}^{w_{1}}W\widetilde{P}_{n}\rangle\boldsymbol{Q_{m}}\widetilde{\boldsymbol{L}}$$

$$+ \left(\mathbb{1}(n=1)i\eta_{t}\langle B_{d}(\mathring{M}^{\widetilde{w}_{1}})^{2}F_{1}\rangle + \mathbb{1}(n=2)\frac{1}{N}\langle B_{d}\mathring{M}^{\widetilde{w}_{1}}F_{1}\mathring{M}^{\widetilde{w}_{2}}F_{2}\rangle\right)\mathbb{E}\boldsymbol{Q_{m}}\widetilde{\boldsymbol{L}}$$

$$+ i\eta\mathbb{E}\mathfrak{Y}_{|\boldsymbol{m}|+n}(B_{d}\mathring{M}^{\widetilde{w}_{1}}) + \frac{1}{N}\mathbb{E}\mathfrak{Y}_{|\boldsymbol{m}|+n-1}(B_{d}\mathring{M}^{\widetilde{w}_{1}}).$$

By Stein's lemma and Leibniz rule, we may write the first term of (5.55) as

$$(5.56) \quad -\mathbb{E}\boldsymbol{Q}_{\boldsymbol{m}}\langle B_{d}\mathring{M}^{\widetilde{w}_{1}}W\widetilde{P}_{n}\rangle\widetilde{\boldsymbol{L}} = -\mathbb{E}\mathbb{E}_{\widetilde{W}}\langle B_{d}\mathring{M}^{\widetilde{w}_{1}}\widetilde{W}\partial_{\widetilde{W}}[\widetilde{P}_{n}]\rangle\boldsymbol{Q}_{\boldsymbol{m}}\widetilde{\boldsymbol{L}} - \mathbb{E}\mathbb{E}_{\widetilde{W}}\langle B_{d}\mathring{M}^{\widetilde{w}_{1}}\widetilde{W}\widetilde{P}_{n}\rangle\partial_{\widetilde{W}}[\boldsymbol{Q}_{\boldsymbol{m}}\widetilde{\boldsymbol{L}}].$$

For the first term of (5.56), we write

(5.57)
$$-\partial_{\widetilde{W}}\widetilde{P}_n = \frac{1}{N}\sum_{i=1}^n \widetilde{P}_{\llbracket i-1 \rrbracket}G^{\widetilde{w}_i}\widetilde{W}\widetilde{P}_{\llbracket i,n \rrbracket},$$

so that

(5.58)
$$-\mathbb{E}\mathbb{E}_{\widetilde{W}}\langle B_d \mathring{M}^{\widetilde{w}_1} \widetilde{W} \partial_{\widetilde{W}}[\widetilde{P}_n] \rangle \boldsymbol{Q}_m \widetilde{\boldsymbol{L}} = \frac{1}{N} \mathbb{E} \sum_{i=1}^n \left\langle B_d \mathring{M}^{\widetilde{w}_1} \mathcal{S}[\widetilde{P}_{[[i-1]]} G^{\widetilde{w}_i}] \widetilde{P}_{[[i,n]]} \right\rangle \boldsymbol{Q}_m \widetilde{\boldsymbol{L}}.$$

For each $i \in [\![1, n]\!]$, we have

(5.59)
$$\frac{\frac{1}{N} \langle B_d \mathring{M}^{\widetilde{w}_1} \mathcal{S}[\widetilde{P}_{\llbracket i-1 \rrbracket} G^{\widetilde{w}_i}] \widetilde{P}_{\llbracket i,n \rrbracket} \rangle \boldsymbol{Q}_m \widetilde{\boldsymbol{L}} = \langle 2B_d \mathring{M}^{\widetilde{w}_1} \widetilde{P}_{\llbracket i,n \rrbracket} \rangle \langle P_i(\widetilde{w}_{\llbracket i \rrbracket}; \boldsymbol{F}_{\llbracket i-1 \rrbracket}, F_n) \rangle \boldsymbol{Q}_m \widetilde{\boldsymbol{L}}}{=\mathbb{1}(i=n) \langle 2B_d \mathring{M}^{\widetilde{w}_1} \mathring{M}^{\widetilde{w}_n} \rangle \langle P_n \rangle \boldsymbol{Q}_m \widetilde{\boldsymbol{L}} + \frac{1}{2} \mathfrak{Y}_{|\boldsymbol{m}|+n+1} (B_d \mathring{M}^{\widetilde{w}_1}).$$

Summing (5.59) over $i \in [1, n]$ and then substituting into (5.56), it only remains to prove

(5.60)
$$-\mathbb{E}_{\widetilde{W}}\langle B_d \mathring{M}^{\widetilde{w}_1} \widetilde{W} \widetilde{P}_n \partial_{\widetilde{W}} [\boldsymbol{Q_m} \widetilde{\boldsymbol{L}}] \rangle = \mathfrak{Y}_{\boldsymbol{m}+n+1} (B_d \mathring{M}^{\widetilde{w}_1}).$$

We first consider the derivative of Q_m . For each factor $\langle P_{m_j} \rangle \equiv \langle P_{m_j}(\tilde{\boldsymbol{w}}', \boldsymbol{F}') \rangle$ in Q_m , we use (5.12) and (5.57) to write

$$(5.61) - \mathbb{E}_{\widetilde{W}} \langle B_d M^{w_1} W P_n \rangle \partial_{\widetilde{W}} [\langle P_{m_j}(\widetilde{\boldsymbol{w}}'; \boldsymbol{F}') \rangle] \boldsymbol{Q}^{(j)} \boldsymbol{L}$$
$$= \sum_i \mathbb{E}_{\widetilde{W}} \langle B_d \mathring{M}^{\widetilde{w}_1} \widetilde{W} \widetilde{P}_n \rangle \langle \widetilde{W} \widetilde{P}_{m_j}(\widetilde{\boldsymbol{w}}'_i; \boldsymbol{F}'_i) G^{\widetilde{w}'_i} \rangle \boldsymbol{Q}^{(j)} \widetilde{\boldsymbol{L}}$$
$$= \frac{1}{2} \sum_i \langle B_d \mathring{M}^{\widetilde{w}_1} P_{m_j+n+1}(\widetilde{\boldsymbol{w}}'_i, \widetilde{\boldsymbol{w}}'_i, \widetilde{\boldsymbol{w}}; \boldsymbol{F}'_i, \check{F}_n, \boldsymbol{F}) \rangle \boldsymbol{Q}^{(j)} \widetilde{\boldsymbol{L}}$$
$$= \frac{1}{2} \mathfrak{Y}_{\boldsymbol{m}+n+1}(B_d \mathring{M}^{\widetilde{w}_1}),$$

for some cyclic rearrangements $\widetilde{\boldsymbol{w}}'_i$ and \boldsymbol{F}'_i of $\widetilde{\boldsymbol{w}}'$ and \boldsymbol{F}' depending on $i \in [m_j]$. Summing (5.61) over j proves

(5.62)
$$-\mathbb{E}_{\widetilde{W}}\langle B_d \mathring{M}^{\widetilde{w}_1} \widetilde{W} \widetilde{P}_n \rangle \partial_{\widetilde{W}} [\boldsymbol{Q}_m] \widetilde{\boldsymbol{L}} = \frac{1}{2} \mathfrak{Y}_{m+n+1} (B_d \mathring{M}^{\widetilde{w}_1}).$$

Similarly, the derivative of \widetilde{L} can be handled using

(5.63)
$$\partial_{\widetilde{W}}L(w) = \operatorname{Tr} G^{w}\widetilde{W}$$

This completes the proof of Lemma 5.6.

Proof of Lemma 5.7. First of all, we replace B_d by $F_n B_d$ using

(5.64)
$$\langle B_d P_n \rangle = \langle B_d P_n F_n \rangle = \langle F_n B_d P_n \rangle$$

Then we apply (5.23) and estimate the last line therein using Lemma 5.5, $\eta \leq \Psi^3$, and $N^{-1} \leq \Psi^4$;

(5.65)
$$\eta|\mathfrak{Y}_2(F_nB_d\mathring{M}^{\widetilde{w}_1})| + N^{-1}|\mathfrak{Y}_1(F_nB_d\mathring{M}^{\widetilde{w}_1})| \prec ||B_d||\Psi^{5}$$

Thus it suffices to express the the third line of (5.23) as \mathfrak{X}_4 , i.e. to prove that

(5.66)
$$\mathbb{E}\mathfrak{Y}_3(F_n B_d \check{M}^{\widetilde{w}_1}) = \langle F_n B_d (\check{M}^0)^4 \rangle \mathfrak{X}_4.$$

Since $F_n B_d \mathring{M}^{\widetilde{w}_1}$ is supported on off-diagonal blocks, we may apply (5.24) to each summand in $\mathbb{E}\mathfrak{Y}_3(F_n B_d \mathring{M}^{\widetilde{w}_1})$ to write it as a finite sum of

(5.67)
$$\mathbb{E}\mathfrak{Y}_4(F_n B_d \mathring{M}^{\widetilde{w}_1} \mathring{M}^{w'_1}) + O(N^{\epsilon} \|B_d\| \Psi^6),$$

where w'_1 may vary in $\{w_1, \ldots, w_k\}$. Likewise, applying (5.23) to each summand in $\mathbb{E}\mathfrak{Y}_4(F_n B_d \mathring{M}^{\widetilde{w}_1} \mathring{M}^{w'_1})$, we conclude that $\mathbb{E}\mathfrak{Y}_3(F_n B_d \mathring{M}^{\widetilde{w}_1})$ is a finite sum of the expressions

(5.68)
$$\langle 2F_n B_d \mathring{M}^{\widetilde{w}_1} \mathring{M}^{w'_1} \mathring{M}^{w'_2} \mathring{M}^{w'_3} E \rangle \mathbb{E}\mathfrak{X}_4 + O(N^{\epsilon} \|B_d\| \Psi^6)$$

where E can be either E_1 or E_2 and w'_1, w'_2, w'_3 may vary in $\{w_1, \ldots, w_k\}$. Furthermore, Lemma 5.6 implies that the exact form of the sum is determined by the expressions P_n, Q_m , and \tilde{L} on the left-hand side of (5.27) that we started from.

Now it only remains to notice that (5.68) can be further expressed as

(5.69)
$$\langle 2F_n B_d \mathring{M}^{\widetilde{w}_1} \mathring{M}^{w'_1} \mathring{M}^{w'_2} \mathring{M}^{w'_3} E \rangle \mathbb{E} \mathfrak{X}_4 = \mathbb{1} (E = F_n) \langle 2B_d (\mathring{M}^0)^4 F_n \rangle + O(N^{\epsilon} \Psi^6),$$

where we used $\|\mathring{M}^w - \mathring{M}^0\| = O(N^{-1/2}) = O(\Psi^2)$ and $|\mathfrak{X}_4| \prec \Psi^4$.

6. Proof of Proposition 4.4, the real case

The main goal of this section is to prove Proposition 4.4 when $\mathbb{F} = \mathbb{R}$. Since we only consider deformed real Ginibre ensembles, we drop the superscripts Gin and $\operatorname{Gin}(\mathbb{F})$.

The fundamental difference in the real case is that the Hermitization

(6.1)
$$W := \begin{pmatrix} 0 & X \\ X^{\mathsf{T}} & 0 \end{pmatrix}$$

is now a real symmetric random matrix. Thus, each application of Stein's lemma results in an additional term as follows (recall the definitions of E_1, E_2 from (1.3)):

(6.2)
$$\mathcal{S}_{\mathbb{R}}[B] := \mathbb{E}_{\widetilde{W}} \widetilde{W} B \widetilde{W} = \mathcal{S}[B] + \frac{1}{N} \sum_{E=E_1, E_2} E B^{\mathsf{T}} \check{E},$$
$$\mathbb{E}_{\widetilde{W}} \widetilde{W} \operatorname{Tr}[\widetilde{W}B] = \frac{1}{N} \sum_{E=E_1, E_2} E(B + B^{\mathsf{T}}) \check{E}.$$

Notice that the additional terms involve the transpose, so that we often encounter G^{\intercal} and its deterministic counterparts M^{\intercal} and M^{\intercal} . In particular, we will shortly see that the size of the quantity

(6.3)
$$\theta_0 := 1 - \left\langle \frac{1}{(A - z_0)(A - z_0)^{\mathsf{T}}} \right\rangle \in \mathbb{C}$$

determines the universality class of our statistics. Notice from Assumption 2.6 (i) that $|1 - \theta_0| \leq 1$, and that if $(A-z_0)^{\dagger}$ is equal to $(A-z_0)^*$, i.e. if $A-z_0$ is real, then θ_0 is identically zero. Furthermore, writing $\mathcal{A}_0 := (A - z_0)$, (recall from (1.4) that $\mathfrak{S}[\mathcal{A}_0]$ stands for entrywise imaginary part of \mathcal{A}_0)

(6.4)

$$0 \leq \operatorname{Re} \theta_{0} = \frac{1}{2} \left\langle \frac{1}{|\mathcal{A}_{0}|^{2}} - \frac{1}{\mathcal{A}_{0}\mathcal{A}_{0}^{\mathsf{T}}} - \frac{1}{\mathcal{A}_{0}^{*}\overline{\mathcal{A}}_{0}} + \frac{1}{|\overline{\mathcal{A}}_{0}|^{2}} \right\rangle = \frac{1}{2} \left\langle \left| \frac{1}{\mathcal{A}_{0}} - \frac{1}{\overline{\mathcal{A}}_{0}} \right|^{2} \right\rangle \sim \langle |\Im[\mathcal{A}_{0}]|^{2} \rangle,$$

$$\operatorname{Im} \theta_{0} = \left\langle \frac{1}{\mathcal{A}_{0}\mathcal{A}_{0}^{\mathsf{T}}} (\Im[\mathcal{A}_{0}]\mathcal{A}_{0}^{\mathsf{T}} + \overline{\mathcal{A}}_{0}}\Im[\mathcal{A}_{0}]^{\mathsf{T}}) \frac{1}{\overline{\mathcal{A}}_{0}\mathcal{A}_{0}^{*}} \right\rangle = 2 \left\langle \Im[\mathcal{A}_{0}]\mathcal{A}_{0}^{*} \frac{1}{|\mathcal{A}_{0}|^{4}} \right\rangle + O(\langle |\Im[\mathcal{A}_{0}]|^{2} \rangle),$$

where we used $\|\mathcal{A}_0^{-1}\|, \|\mathcal{A}_0\| = O(1)$ and Cauchy-Schwarz to get asymptotic relations. In particular, since A is real, the asymptotics (6.4) reads

(6.5)
$$\operatorname{Re} \theta_0 \sim |\operatorname{Im} z_0|^2, \quad \operatorname{Im} \theta_0 = -2I_3 \operatorname{Im} z_0 + O(|\operatorname{Im} z_0|^2)$$

as $|\operatorname{Im} z_0| \to 0$. With this in mind, our proof of Proposition 4.4 in the real case can be divided into two regimes depending on the size of $|\theta_0| \sim |\operatorname{Im} z_0|$ in which we construct a path \mathcal{A}_t differently. When $|\operatorname{Im} z_0| \geq N^{-1/2+\mathfrak{c}}$, we construct a path \mathcal{A}_t that makes the corresponding θ always much larger than $N^{-1/2}$. When $|\operatorname{Im} z_0| \leq N^{-1/2+\mathfrak{c}}$, our choice of \mathcal{A}_t keeps the value of $I_4^{-1/2}\theta$ almost constant. We formalize this intuition in the next assumption.

Assumption 6.1. Assume that A_t is a path satisfying Assumption 4.6. Defining

(6.6)
$$\theta_t := 1 - \left\langle \frac{1}{\mathcal{A}_t \mathcal{A}_t^{\mathsf{T}}} \right\rangle,$$

we further assume that either of the following holds.

- (A1) for some small constants $\mathfrak{c}_1, \mathfrak{c}_2 > 0$, $|\theta_t| \ge N^{-1/2+\mathfrak{c}_1}$ and $|\theta_t^{-2} \mathrm{d}\theta_t/\mathrm{d}t| \le N^{1/2-\mathfrak{c}_2}$ for all $t \in [0, 1]$. (A2) for some small constants $\mathfrak{c}_1, \mathfrak{c}_2 > 0$, $|\theta_t| \le N^{-1/2+\mathfrak{c}_1}$, $|\mathrm{d}[I_4(t)^{-1/2}\theta_t]/\mathrm{d}t| \le N^{-1/2-\mathfrak{c}_2}$, and $\|\mathrm{d}\Im\mathcal{A}_t/\mathrm{d}t\| \leq N^{-\mathfrak{c}_2} \text{ for all } t \in [0,1].$

We next construct such a path \mathcal{A}_t in each of the two cases.

Proposition 6.2. Suppose that A is real and $z_0 \in \mathbb{C}$ satisfies Assumption 2.6. Recall the small constant $\mathfrak{c} > 0$ from Assumption 2.6.

- (i) If $|\operatorname{Im} z_0| \geq N^{-1/2+\mathfrak{c}}$ i.e. if z_0 satisfies Assumption 2.6 (iii \mathbb{R}), then there exists a path \mathcal{A}_t satisfying Assumption 6.1 (A1).
- (ii) If $|\operatorname{Im} z_0| \leq N^{-1/2+\mathfrak{c}}$ i.e. if z_0 satisfies Assumption 2.6 (iii \mathbb{R}'), then there exists a path \mathcal{A}_t satisfying Assumption 6.1 (A2).

Moreover, the constants C_5 in (4.35) and $\mathfrak{c}_1, \mathfrak{c}_2$ in (A1) – (A2) depend only on those in Assumption 2.6.



Figure 3. Case (A2): Leftmost picture shows the path \hat{z}_t , the other pictures show the eigenvalues of $\mathcal{A}_t + X$ and the evolution of the base point $z_t = \langle |A - \hat{z}_t|^{-2} \rangle^{1/2} \hat{z}_t$.

Proof. In what follows, we construct two different paths $\hat{z}_t \in \mathbb{C}$ depending on $\text{Im } z_0$, i.e. the cases (i) and (ii), and show that the matrix-valued path

(6.7)
$$\mathcal{A}_t := \langle |A - \hat{z}_t|^{-2} \rangle^{1/2} (A - \hat{z}_t)$$

satisfies Assumption 6.1, see Figure 3.

We start with the proof of (i), and without loss of generality assume Im $z_0 > 0$. Using Assumption 2.6 (iii \mathbb{R}), we take a smooth path $[0,1] \ni t \to \tilde{z}_t \in \mathcal{D}_{C'} \cap \mathbb{C}_+$ such that $\tilde{z}_0 = z_0$ and $\tilde{z}_1 = Cz_1$ where C > 0 is a large enough constant. Then we notice that, for a small positive constant τ , the path $\tilde{z}_t + i\tau$ is in $\mathcal{D}_{2C'}$. We then define the path \hat{z}_t as follows:

(6.8)
$$\widehat{z}_t := \begin{cases} \operatorname{Re} z_0 + \mathrm{i}(\tau + \operatorname{Im} z_0) \left(\frac{\operatorname{Im} z_0}{\tau + \operatorname{Im} z_0}\right)^{1-3t}, & t \in [0, 1/3]; \\ \widetilde{z}_{3t-2} + \mathrm{i}\tau & t \in [2/3, 2/3]; \\ \frac{2}{3(1-t)}\widetilde{z}_1 + \mathrm{i}\tau & t \in [2/3, 1]. \end{cases}$$

The path \hat{z}_t here is constructed in a similar way to Proposition 5.1, except that in the first part we lift (translate) the path by $i\tau$. Since τ is small but fixed, this guarantees that $|\Im[A - \hat{z}_t]|$ is bounded from below afterwards. We omit verifying Assumption 4.6 since it is analogous to the complex case, but only remark that the last inequality in (4.35), for $t \in [0, 1/3]$, is due to

(6.9)
$$\frac{\mathrm{d}}{\mathrm{d}t}a^t = a^t \log a, \qquad a > 0.$$

Note that by (6.5) we have $|\theta_0| \sim |\operatorname{Im} z_0| \geq N^{-1/2+\mathfrak{c}}$, henceforth we verify that \mathcal{A}_t satisfies (A1). By the definition of \mathcal{A}_t , we have

(6.10)
$$\theta_t = 1 - \langle |A - \hat{z}_t|^{-2} \rangle^{-1} \left\langle \frac{1}{(A - \hat{z}_t)(A - \hat{z}_t)^{\mathsf{T}}} \right\rangle$$

Thus it immediately follows $\theta_t = 2 + O(1-t)$ when t is close enough to 1. Thus we restrict to $t \in [0, 1-\delta)$, in which case $|\hat{z}_t|$ remains bounded so that

(6.11)
$$\operatorname{Re}\theta_t \sim |\operatorname{Im}\widehat{z}_t|^2, \qquad \operatorname{Im}\theta_t = 2\left\langle (A - \widehat{z}_t)^* \frac{1}{|A - \widehat{z}_t|^4} \right\rangle \operatorname{Im}\widehat{z}_t + O(|\operatorname{Im}\widehat{z}_t|^2)$$

by the same argument as in (6.5).

Explicitly computing $d\theta_t/dt$, we find that

$$(6.12) \qquad \begin{aligned} \langle |A - \hat{z}_t|^{-2} \rangle \frac{\mathrm{d}\theta_t}{\mathrm{d}t} \\ = (1 - \theta_t) \frac{\mathrm{d}}{\mathrm{d}t} \left\langle \frac{1}{|A - \hat{z}_t|^2} \right\rangle - \frac{\mathrm{d}}{\mathrm{d}t} \left\langle \frac{1}{(A - \hat{z}_t)(A - \hat{z}_t)^{\mathsf{T}}} \right\rangle \\ = 2(1 - \theta_t) \operatorname{Re}\left[\frac{\mathrm{d}\hat{z}_t}{\mathrm{d}t} \left\langle \frac{1}{(A - \hat{z}_t)^2(A - \hat{z}_t)^*} \right\rangle \right] - 2 \frac{\mathrm{d}\hat{z}_t}{\mathrm{d}t} \left\langle \frac{1}{(A - \hat{z}_t)^2(A - \hat{z}_t)^{\mathsf{T}}} \right\rangle \end{aligned}$$

so that writing down the real and imaginary parts we get

(6.13)
$$\left|\frac{\mathrm{d}\theta_t}{\mathrm{d}t}\right| \lesssim \left|\frac{\mathrm{d}\operatorname{Im}\widehat{z}_t}{\mathrm{d}t}\right| + \left(|\theta| + \operatorname{Im}\widehat{z}_t\right) \left|\frac{\mathrm{d}\operatorname{Re}\widehat{z}_t}{\mathrm{d}t}\right|.$$

From $|\theta_t| \sim |\operatorname{Im} \hat{z}_t|$ due to (6.11) together with $|\operatorname{dRe} \hat{z}_t/\operatorname{d} t| = O(1)$, we thus conclude

(6.14)
$$\left|\frac{1}{\theta_t^2}\frac{\mathrm{d}\theta_t}{\mathrm{d}t}\right| \lesssim \left|\frac{1}{(\mathrm{Im}\,\widehat{z}_t)^2}\frac{\mathrm{d}\,\mathrm{Im}\,\widehat{z}_t}{\mathrm{d}t}\right| + \frac{1}{|\,\mathrm{Im}\,\widehat{z}_t|}$$

Hence (A1) immediately follows from (6.11) and (6.14) since $\operatorname{Im} \hat{z}_t \geq \tau \vee \operatorname{Im} z_0$ and

(6.15)
$$\left|\frac{1}{\operatorname{Im}\widehat{z}_t}\frac{\mathrm{d}\operatorname{Im}\widehat{z}_t}{\mathrm{d}t}\right| \sim \begin{cases} 1+|\log\operatorname{Im}z_0|, & t\in[0,1/3],\\ 1, & t\in[1/3,1], \end{cases}$$

where we used (6.9) for $t \in [0, 1/3]$.

We move on to the proof of (ii), and assume without loss of generality $[\operatorname{Re} z_0, \infty)$ is in $\mathcal{D}_{C'}$. Here we will build a different path \hat{z}_t as follows. Define

(6.16)
$$\operatorname{Re} \widehat{z}_t = \operatorname{Re} z_0 + \frac{t}{1-t}$$

and we take $\operatorname{Im} \widehat{z}_t = \frac{f(\operatorname{Re} \widehat{z}_t)}{f(\operatorname{Re} z_0)} \operatorname{Im} z_0$ where the function $f : [\operatorname{Re} z_0, \infty) \to \mathbb{R} \setminus \{0\}$ is defined by

(6.17)
$$f(x) := \langle |A - x|^{-4} \rangle^{-1/2} \langle |A - x|^{-2} \rangle^{-1} \left\langle \frac{1}{(A - x)^2 (A - x)^{\intercal}} \right\rangle.$$

We then define our matrix-valued path \mathcal{A}_t again by the same formula (6.7). We can easily verify Assumption 4.6 for this path, modulo that the end point is not exactly $-z_1$ but is another point $\mathcal{A}_1 = -z'_1 \in \mathbb{S}^1$ with

(6.18)
$$\operatorname{Im} z_1' = |\gamma| \operatorname{Im} z_0 + o(N^{-1/2}) = \operatorname{Im} z_1 + o(N^{-1/2}).$$

We then concatenate to \mathcal{A}_t the (shorter) arc along the unit circle connecting $-z'_1$ to $-z_1$, which is of length $o(N^{-1/2})$. Henceforth we focus on checking (A2) and (6.18).

Notice that \mathcal{A}_t can be written as

$$\mathcal{A}_t = \Re \mathcal{A}_t + \mathrm{i} k_t \operatorname{Im} z_0$$

where $k_t > 0$ is defined by

(6.19)

(6.20)
$$k_t := \langle |A - \hat{z}_t|^{-2} \rangle^{1/2} \frac{f(\operatorname{Re} \hat{z}_t)}{f(\operatorname{Re} z_0)} = \frac{1}{f(\operatorname{Re} z_0)} \langle |\Re \mathcal{A}_t|^{-4} \rangle^{-1/2} \langle |\Re \mathcal{A}_t|^{-2} \rangle^{-1} \left\langle \frac{1}{(\Re \mathcal{A}_t)^2 \Re \mathcal{A}_t^{\mathsf{T}}} \right\rangle.$$

By direct computation, we easily find that

(6.21)
$$\begin{aligned} \|\Re\mathcal{A}_t\| &= O(1), \qquad \qquad \left\|\frac{1}{\Re\mathcal{A}_t}\right\| = O(1), \qquad \qquad \left\|\frac{\mathrm{d}\Re\mathcal{A}_t}{\mathrm{d}t}\right\| = O(1), \\ \left|\left\langle\frac{1}{(\Re\mathcal{A}_t)^2 \Re\mathcal{A}_t^{\mathsf{T}}}\right\rangle\right| \sim 1, \qquad \qquad |k_t| \sim 1, \qquad \qquad \left|\frac{\mathrm{d}k_t}{\mathrm{d}t}\right| = O(1). \end{aligned}$$

Since \mathcal{A}_t is always of the form (6.19), the asymptotics (6.5) remains true, so that $\operatorname{Im} \theta_t \sim \operatorname{Im} z_0$ and $\operatorname{Re} \theta_t = O(|\operatorname{Im} z_0|^2)$. This immediately proves $|\theta_t| \leq N^{-1/2+\mathfrak{c}_1}$. For the second assertion in (A2), we recall from the definition of θ_t that

(6.22)
$$I_4^{-1/2} \theta_t = I_4^{-1/2} \left\langle \frac{1}{|\mathcal{A}_t|^2} - \frac{1}{\mathcal{A}_t \mathcal{A}_t^{\mathsf{T}}} \right\rangle = \mathrm{i} I_4^{-1/2} k_t \operatorname{Im} z_0 \left\langle \frac{1}{|\mathcal{A}_t|^2 \mathcal{A}_t^{\mathsf{T}}} \right\rangle.$$

By the definition of k_t , we have

(6.23)
$$I_4^{-1/2} \left\langle \frac{k_t}{|\mathcal{A}_t|^2 \mathcal{A}_t^{\mathsf{T}}} \right\rangle = \frac{\langle |\mathcal{A}_t|^{-2} \rangle^2}{\langle |\Re \mathcal{A}_t|^{-2} \rangle^2} \frac{\langle (\Re \mathcal{A}_t^{\mathsf{T}})^{-1} |\Re \mathcal{A}_t|^{-2} \rangle}{\langle (\mathcal{A}_t^{\mathsf{T}})^{-1} |\mathcal{A}_t|^{-2} \rangle} \frac{\langle |\Re \mathcal{A}_t|^{-4} \rangle^{1/2}}{\langle |\mathcal{A}_t|^{-4} \rangle^{1/2}}$$

Thus, after several applications of the resolvent identity

(6.24)
$$\frac{1}{\Re \mathcal{A}_t} - \frac{1}{\mathcal{A}_t} = \mathrm{i} \operatorname{Im} z_0 \frac{1}{\Re [\mathcal{A}_t] \mathcal{A}_t}$$

and (6.21), we arrive at

(6.25)
$$\frac{\mathrm{d}}{\mathrm{d}t} \left\langle \frac{k_t}{|\mathcal{A}_t|^2 \mathcal{A}_t^{\mathsf{T}}} \right\rangle = O(|\operatorname{Im} z_0|)$$

Therefore we conclude

(6.26)
$$\left| \frac{\mathrm{d}\theta_t}{\mathrm{d}t} \right| = O(|\operatorname{Im} z_0|^2) = O(N^{-1+2\mathfrak{c}}) = O(N^{-1/2-\mathfrak{c}_2}),$$

as desired. The last assertion of (A2) immediately follows from (6.21) as

(6.27)
$$\left\|\frac{\mathrm{d}\Im\mathcal{A}_t}{\mathrm{d}t}\right\| = |\operatorname{Im} z_0| \left|\frac{\mathrm{d}k_t}{\mathrm{d}t}\right| \lesssim \operatorname{Im} z_0.$$

To show (6.18), we only need to notice that

$$I_4(0)^{-1/2}\theta_0 = 2i\operatorname{Im} z_0 I_4^{-1/2} \left\langle \frac{1}{(A-z_0)(A-z_0)^{\intercal}(A-z_0)^{\intercal}} \right\rangle = 2i\operatorname{Im} z_0(|\gamma_0| + O(|\operatorname{Im} z_0|))$$

= $I_4(1)^{-1/2}\theta_1 + O(N^{-1/2-\mathfrak{c}_2}) = 1 - \frac{1}{z_1^2} + O(N^{-1/2-\mathfrak{c}_2}) = 2i\operatorname{Im} z_1 + o(N^{-1/2}).$
completes the proof of Proposition 6.2.

1

This completes the proof of Proposition 6.2.

Now, as in the complex case, Proposition 4.4 in the real case is an immediate consequence of the following counterpart of Proposition 5.2. Recall the definition of c_t from (4.37).

Proposition 6.3. Let $\mathbb{F} = \mathbb{R}$, fix $k \in \mathbb{N}$ and C > 0, and suppose that \mathcal{A}_t satisfies Assumption 6.1. Let $0 < \delta < (\mathfrak{c}_1 \wedge \mathfrak{c}_2)/10$, where we recall $\eta_0 = N^{-3/4+\delta}$. Then there exists a constant c > 0 for which the following holds uniformly over \mathcal{A}_t , $t \in [0, 1]$, and $|w|, ||w|| \leq C$.

(6.29)
$$\sum_{j=1}^{k} \mathbb{E}\left\langle \frac{\mathrm{d}(\mathbb{E}H_t^{w_j} - \mathrm{i}\eta_t)}{\mathrm{d}t} G_t^{w_j} \right\rangle \boldsymbol{L}_t^{(j)}(\boldsymbol{w}^{(j)}) = h_t(\boldsymbol{w}) + O(N^{-1-c}),$$

. . 1

(6.30)
$$\frac{\mathrm{d}}{\mathrm{d}t}\mathbb{E}\frac{\langle G_t^w \rangle}{c_t} = O(N^{-c}\Psi),$$

where $h_t(\boldsymbol{w})$ satisfies $(\prod_{j=1}^k \Delta_{w_j})h_t(\boldsymbol{w}) \equiv 0.$

The rest of this section is organized as follows: In Section 6.1, we present a general framework of the proof in the real case, pointing out the main difference from the complex case. In particular, we present the decoupling lemma, Lemma 5.6, adapted to the real case. Then in Sections 6.2 and 6.3 we prove Proposition 6.3 under the assumptions (A1) and (A2), respectively. We emphasize that the rest of Section 6 applies even when the path A_t is complex, as long as it satisfies Assumption 6.1. In particular, all estimates in the rest of this section depend only on $k, C, C_5, \mathfrak{c}_1, \mathfrak{c}_2$.

6.1. General framework. In what follows, we introduce notations parallel to those in Section 5.1 in the complex case. Virtually the only difference is that, on top of symbols \boldsymbol{w} and \boldsymbol{F} in (5.16), we also have another tuple ι whose components vary in $\{1, \intercal\}$ (recall (6.2)). As we will shortly see, the parallel notations allow us to verbatim reuse algebraic expressions appeared in the complex case. To avoid repetition, when we encounter an expression that already appeared in Sections 5.1–5.2, we simply refer to an equation therein and do not rewrite the same expression again; see e.g. (6.46) and (6.54).

The following definitions integrate the newly appeared transposes into the same framework as in Section 5.

Definition 6.4. For a given $n \in \mathbb{N}$ and n-tuples w, F, ι , define

(6.31)
$$P_n(\boldsymbol{w}; \boldsymbol{F}; \boldsymbol{\iota}) \coloneqq (G^{w_1})^{\iota_1} F_1 \cdots (G^{w_n})^{\iota_n} F_n.$$

For a multi-index $\boldsymbol{m} = (m_j)_j$ in \mathbb{N} and $|\boldsymbol{m}|$ -tuples $\boldsymbol{w} = (\boldsymbol{w}_j)_j$, $\boldsymbol{F} = (\boldsymbol{F}_j)_j$, $\boldsymbol{\iota} = (\boldsymbol{\iota}_j)_j$, define

(6.32)
$$\boldsymbol{Q}_{\boldsymbol{m}}(\boldsymbol{w};\boldsymbol{F};\boldsymbol{\iota}) = \prod_{j} \langle P_{m_j}(\boldsymbol{w}_j;\boldsymbol{F}_j;\boldsymbol{\iota}_j) \rangle.$$

We say an n-tuple $\boldsymbol{\iota}$ is mixed if $\boldsymbol{\iota} \neq (1, \ldots, 1), (\boldsymbol{\intercal}, \ldots, \boldsymbol{\intercal})$, in which case we say $P_n(\boldsymbol{w}; \boldsymbol{F}; \boldsymbol{\iota})$ is mixed and write $P_n^{\min} \equiv P_n$. The n-tuple $\boldsymbol{\iota} = (1, \ldots, 1)$ and $P_n(\boldsymbol{w}; \boldsymbol{F}; 1, \ldots, 1)$ are called pure, and we write $P_n^{\text{pure}} \equiv P_n$. Similarly, \boldsymbol{Q}_m is called mixed if at least one of the involved factors of P_{m_j} are mixed, and we write $\boldsymbol{Q}_{\boldsymbol{m}}^{\min} \equiv \boldsymbol{Q}_{\boldsymbol{m}}$ to emphasize this fact. The expression $\boldsymbol{Q}_{\boldsymbol{m}}$ is called pure if all involved factors of P_{m_j} are pure, in which case we write $\boldsymbol{Q}_{\boldsymbol{m}}^{\text{pure}} \equiv \boldsymbol{Q}_{\boldsymbol{m}}$. Notice that $P_n(\boldsymbol{w}; \boldsymbol{F}; \boldsymbol{\uparrow}, \dots, \boldsymbol{\uparrow})$ is not called pure nor mixed.

Definition 6.5. We naturally extend the definitions of \mathcal{X}_n and $\mathcal{Y}_n(B)$ from Definition 5.4 by introducing the ι -variables. We define $\mathcal{X}_n^{\text{mix}}$ (resp. $\mathcal{X}_n^{\text{pure}}$) to be the subset of \mathcal{X}_n consisting of the same expressions as in (5.20) such that Q_m is mixed (resp. pure). Also, we write $\mathcal{Y}_n^{\min}(B)$ (resp. $\mathcal{Y}_n^{pure}(B)$) for the subset of $\mathcal{Y}_n(B)$ such that at least one (resp. both) of

the factors P_{n_1} and Q_m in (5.21) is mixed (resp. are pure). As in Definition 5.4, we have $\mathcal{Y}_n^{\min}(I) = \mathcal{X}_n^{\min}$ and $\mathcal{Y}_n^{\text{pure}}(I) = \mathcal{X}_n^{\text{pure}}$.

Definition 6.6. With a light abuse of notations, we denote generic finite (independent of N) sums of elements of $\mathcal{X}_n, \mathcal{X}_n^{\text{mix}}$, and $\mathcal{X}_n^{\text{pure}}$ as $\mathfrak{X}_n, \mathfrak{X}_n^{\text{mix}}$, and $\mathfrak{X}_n^{\text{pure}}$, respectively. Similarly we define $\mathfrak{Y}_n, \mathfrak{Y}_n^{\text{mix}}$, and $\mathfrak{Y}_n^{\text{pure}}$ to be generic finite sums over $\mathcal{Y}_n, \mathcal{Y}_n^{\text{mix}}$, and $\mathcal{Y}_n^{\text{pure}}$. We now state the decoupling lemma incorporating G^{\intercal} . Its proof is completely analogous to that of Lemma 5.6 hence omitted.

Lemma 6.7. Let B_d and $B_o \in \mathbb{C}^{N \times N}$ be supported respectively on diagonal and off-diagonal blocks. Then we have

$$\mathbb{E}\langle B_d P_n^{\min} \rangle \boldsymbol{Q_m} \boldsymbol{L}$$

$$= \langle 2B_d(\mathring{M}^{\widetilde{w}_1})^{\iota_1}(\mathring{M}^{\widetilde{w}_n})^{\iota_n} F_n \rangle \mathbb{E}\langle P_n^{\min} \rangle \boldsymbol{Q_m} \widetilde{\boldsymbol{L}}$$

$$+ \left(\mathbbm{1}(n=1)i\eta_t \langle B_d((\mathring{M}^{\widetilde{w}_1})^{\iota_1})^2 F_1 \rangle + \mathbbm{1}(n=2) \frac{1}{N} \langle B_d(\mathring{M}^{\widetilde{w}_1})^{\iota_1} F_1(\mathring{M}^{\widetilde{w}_2})^{\iota_2} F_2 \rangle \right) \mathbb{E} \boldsymbol{Q_m} \widetilde{\boldsymbol{L}}$$

$$+ \frac{1}{2} \mathbb{E} \mathfrak{Y}_{|\boldsymbol{m}|+n+1}^{\min} (B_d(\mathring{M}^{\widetilde{w}_1})^{\iota_1})$$

$$+ i\eta \mathbb{E} \mathfrak{Y}_{|\boldsymbol{m}|+n} (B_d(\mathring{M}^{\widetilde{w}_1})^{\iota_1}) + \frac{1}{N} \mathbb{E} \mathfrak{Y}_{|\boldsymbol{m}|+n-1} (B_d(\mathring{M}^{\widetilde{w}_1})^{\iota_1}),$$

and

(6.34)
$$\mathbb{E}\langle B_o P_n^{\min} \rangle \boldsymbol{Q}_{\boldsymbol{m}} \widetilde{\boldsymbol{L}} = \frac{1}{2} \mathbb{E} \mathfrak{Y}_{|\boldsymbol{m}|+n+1}^{\min} (B_o(\mathring{M}^{\widetilde{w}_1})^{\iota_1}) \\ + i\eta \mathbb{E} \mathfrak{Y}_{|\boldsymbol{m}|+n} (B_o(\mathring{M}^{\widetilde{w}_1})^{\iota_1}) + \frac{1}{N} \mathbb{E} \mathfrak{Y}_{|\boldsymbol{m}|+n-1} (B_o(\mathring{M}^{\widetilde{w}_1})^{\iota_1}),$$

where we abbreviated $P_n^{\min} \equiv P_n^{\min}(\widetilde{\boldsymbol{w}}; \boldsymbol{F}; \boldsymbol{\iota})$. The same holds true if we remove all superscripts mix, or if we replace P_n^{\min} on the left-hand sides by P_n^{pure} and $\mathfrak{Y}_{|\boldsymbol{m}+n+1|}^{\min}$ on the right-hand side by $\mathfrak{Y}_{|\boldsymbol{m}|+n+1}^{\text{pure}} + \mathfrak{Y}_{|\boldsymbol{m}|+n+1}^{\min}$.

The counterpart of (5.29) is as follows:

$$\mathbb{E}\left\langle \frac{\mathrm{d}\mathbb{E}H_{t}^{w_{j}}}{\mathrm{d}t}G^{w_{j}}E_{1}\right\rangle \boldsymbol{L}^{(j)} \\
= \left\langle B_{d}^{w_{j}}E_{1}\right\rangle \mathbb{E}\boldsymbol{L}^{(j)} + \mathrm{i}\eta_{t}\mathbb{E}\left\langle B_{d}^{w_{j}}G^{w_{j}}E_{1}\right\rangle \boldsymbol{L}^{(j)} \\
+ \mathbb{E}\left\langle G^{w_{j}}\right\rangle \left\langle B_{d}^{w_{j}}G^{w_{j}}E_{1}\right\rangle \boldsymbol{L}^{(j)} + \frac{1}{N}\mathbb{E}\left\langle B_{d}^{w_{j}}E_{1}(G^{w_{j}})^{\mathsf{T}}E_{2}G^{w_{j}}E_{1}\right\rangle \boldsymbol{L}^{(j)} \\
- \frac{1}{N}\sum_{\ell}^{(j)}\mathbb{E}\left\langle B_{d}^{w_{j}}(G^{w_{\ell}} + (G^{w_{\ell}})^{\mathsf{T}})E_{2}G^{w_{j}}E_{1}\right\rangle \boldsymbol{L}^{(j,\ell)}.$$
(6.35)

In what follows, we take care of the new terms in (6.35) involving G^{\intercal} as well as similar new terms that appears when we expand

(6.36)
$$\mathbb{E}\langle B_d^{w_j}G^{w_j}E_1\rangle \boldsymbol{L}^{(j)}, \qquad \mathbb{E}\langle B_d^{w_j}G^{w_\ell}E_2G^{w_j}E_1\rangle \boldsymbol{L}^{(j,\ell)}.$$

In particular, we will shortly see that their contributions are negligible when $|\theta_t| \gtrsim N^{-1/2+\mathfrak{c}_1}$, and cancel with each other when $|\theta_t| \lesssim N^{-1/2+\mathfrak{c}_1}$. In the latter regime, we use an additional cancellation originating from Assumption 6.1 (A2) that $|d\theta_t/dt|$ is much smaller than $N^{-1/2}$.

6.2. Large θ regime. Throughout this section, we assume that the path \mathcal{A}_t satisfies Assumption 6.1 (A1).

Lemma 6.8. Fix a multi-index m in \mathbb{N} . Then we have

(6.37)
$$|\mathbb{E}\boldsymbol{Q}_{\boldsymbol{m}}^{\min}\widetilde{\boldsymbol{L}}| \lesssim N^{\epsilon}|\boldsymbol{\theta}|^{-1}\Psi^{|\boldsymbol{m}|+2}.$$

uniformly over (w_1, \ldots, w_k) in a compact set.

Proof. Without loss of generality, suppose that the first factor P_{m_1} in Q_m is mixed with $\iota_{11} = 1$ and $\iota_{1m_1} = \intercal$. Taking $B_d = I$ and applying (6.33), we have

(6.38)
$$\mathbb{E}\boldsymbol{Q}_{\boldsymbol{m}}\widetilde{\boldsymbol{L}} = \langle 2\mathring{M}^{w_{11}}(\mathring{M}^{w_{1n}})^{\mathsf{T}}F_n \rangle \mathbb{E}\boldsymbol{Q}_{\boldsymbol{m}}\widetilde{\boldsymbol{L}} + \mathbb{E}\mathfrak{Y}_{|\boldsymbol{m}|+n+1}(\mathring{M}^{w_{11}}) + O(N^{\epsilon}\Psi^{|\boldsymbol{m}|+2}).$$

Notice that

(6.39)
$$1 - \langle 2\mathring{M}^{0}(\mathring{M}^{0})^{\mathsf{T}}F_{n} \rangle = \begin{cases} \overline{\theta} & \text{if } F_{n} = E_{1} \\ \theta & \text{if } F_{n} = E_{2}, \end{cases}$$

so that by (A1)

(6.40)
$$\left|1 - \langle 2\mathring{M}^{w_1}(\mathring{M}^{w_{1n}})^{\mathsf{T}}F_n\rangle\right| = |\theta| + O(N^{-1/2}) = |\theta|(1 + O(N^{-\mathfrak{c}_1})).$$

On the other hand, since $\mathring{M}^{w_{11}}$ is supported on off-diagonal blocks, (6.34) implies

(6.41)
$$\mathbb{E}\mathfrak{Y}_{|\boldsymbol{m}|+n+1}(\dot{M}^{w_{11}}) = O(N^{\epsilon}\Psi^{|\boldsymbol{m}|+2}).$$

Plugging (6.40) and (6.41) into (6.38) proves the result.

As an immediate consequence of Lemmas 6.7 and 6.8, we have that

(6.42)
$$|\mathbb{E}\mathfrak{Y}_n^{\min}(B_d)| \lesssim N^{\epsilon} ||B_d|| |\theta|^{-1} \Psi^{n+2}, \qquad |\mathbb{E}\mathfrak{Y}_n^{\min}(B_o)| \lesssim N^{\epsilon} ||B_o|| |\theta|^{-1} \Psi^{n+3},$$

for any fixed B_d and $B_o \in \mathbb{C}^{2N \times 2N}$ supported respectively on diagonal and off-diagonal blocks. Armed with (6.37) and (6.42), we may now prove the counterpart of Lemma 5.8:

Lemma 6.9. Let n, m, and B_d be as in Lemma 5.7, and assume that $P_n \equiv P_n^{\text{pure}}$ and $Q_m \equiv Q_m^{\text{pure}}$. Then (5.28) remains true, modulo an additional error of $O(N^{\epsilon} ||B_d|| |\theta|^{-1} \Psi^6)$.

Proof. Similarly to Lemma 5.8, we only need to prove the following counterpart of (5.27): The only difference is that the error has an additional factor of $|\theta|^{-1}$.

(6.43)

$$\mathbb{E}\langle B_{d}P_{n}\rangle\boldsymbol{Q_{m}}\widetilde{\boldsymbol{L}} = \langle 2B_{d}\mathring{M}^{\widetilde{w}_{1}}\mathring{M}^{\widetilde{w}_{n}}F_{n}\rangle\mathbb{E}\langle P_{n}\rangle\boldsymbol{Q_{m}}\widetilde{\boldsymbol{L}} + \left(\mathbb{1}(n=1)\mathrm{i}\eta_{t}\langle B_{d}(\mathring{M}^{\widetilde{w}_{1}})^{2}F_{1}\rangle + \mathbb{1}(n=2)\frac{1}{N}\langle B_{d}\mathring{M}^{\widetilde{w}_{1}}F_{1}\mathring{M}^{\widetilde{w}_{2}}F_{2}\rangle\right)\mathbb{E}\boldsymbol{Q_{m}}\widetilde{\boldsymbol{L}} + \frac{1}{2}\langle B_{d}(\mathring{M}^{0})^{4}F_{n}\rangle\mathbb{E}\mathfrak{X}_{4}^{\mathrm{pure}} + O(N^{\epsilon}|\boldsymbol{\theta}|^{-1}||B_{d}||\Psi^{6}).$$

The proof of (6.43) follows that of (5.27) almost verbatim; we apply Lemma 6.7 three times instead of Lemma 5.6. The only difference is that, in the first application of (6.33), we decompose the second order term as

(6.44)
$$\mathbb{E}\mathfrak{Y}_{|\boldsymbol{m}|+n+1}(B_d \mathring{M}^{\widetilde{w}_1}) = \mathbb{E}\mathfrak{Y}_{|\boldsymbol{m}|+n+1}^{\text{pure}}(B_d \mathring{M}^{\widetilde{w}_1}) + \mathbb{E}\mathfrak{Y}_{|\boldsymbol{m}|+n+1}^{\text{mix}}(B_d \mathring{M}^{\widetilde{w}_1}),$$

and estimate the mixed part using (6.42).

Lemma 6.9 shows that the expansions of the 'pure' terms in (6.35) is identical to those of (5.29). Thus, it only remains to show that the 'mixed' terms therein have no contribution. Here we record the application of Lemma 6.7 and (6.42) to them:

Lemma 6.10. Let B_d be supported on diagonal blocks. Then we have

(6.45)
$$\mathbb{E}\langle B_d(G^{\widetilde{w}_1})^{\mathsf{T}} E_2 G^{\widetilde{w}_2} E_1 \rangle \widetilde{\boldsymbol{L}} = \langle 2B_d(\mathring{M}^{\widetilde{w}_1})^{\mathsf{T}} \mathring{M}^{\widetilde{w}_2} E_1 \rangle \mathbb{E}\left(\langle (G^{\widetilde{w}_1})^{\mathsf{T}} E_2 G^{\widetilde{w}_2} E_1 \rangle + \frac{1}{2N}\right) \widetilde{\boldsymbol{L}} + O(N^{\epsilon} \|B_d\| \|\theta\|^{-1} \Psi^6).$$

Proof of Proposition 6.3, (A1). We prove (6.29) assuming that \mathcal{A}_t satisfies Assumption 6.1 (A1). The proof of (6.30) is omitted since it follows from taking the η -derivative of (6.29).

Plugging the result of Lemmas 6.9 and 6.10 into (6.35), we find that

$$\mathbb{E}\left\langle \frac{\mathrm{d}(\mathbb{E}H_t^{w_j} - \mathrm{i}\eta_t)}{\mathrm{d}t} G^{w_j} E_1 \right\rangle \boldsymbol{L}^{(j)} = (\text{right-hand side of } (5.31) \\ + \langle 2B_d^{w_j} (\mathring{M}^{w_j})^{\mathsf{T}} \mathring{M}^{w_j} E_1 \rangle \mathbb{E}\left(\langle (G^{w_j})^{\mathsf{T}} E_2 G^{w_j} E_1 \rangle + \frac{1}{2N}\right) \boldsymbol{L}^{(j)} \\ + \sum_{\ell}^{(j)} \langle 2B_d^{w_j} (\mathring{M}^{w_\ell})^{\mathsf{T}} \mathring{M}^{w_j} E_1 \rangle \mathbb{E}\left(\langle (G^{w_\ell})^{\mathsf{T}} E_2 G^{w_j} E_1 \rangle + \frac{1}{2N}\right) \boldsymbol{L}^{(j,\ell)} \\ + O(N^{\epsilon} |\theta|^{-1} \Psi^6).$$

To estimate the newly appeared terms, note that applying Lemma 6.10 to $B_d = I$ and then using (6.39) gives

(6.47)
$$\mathbb{E}\langle (G^{w_{\ell}})^{\mathsf{T}} E_2 G^{w_j} E_1 \rangle \boldsymbol{L}^{(j,\ell)} = \frac{1+\theta}{\theta} \frac{1}{2N} \mathbb{E} \boldsymbol{L}^{(j,\ell)} + O(N^{\epsilon} |\theta|^{-2} \Psi^6).$$

On the other hand, the deterministic prefactor can be written as

(6.48)
$$\left\langle 2B_d^{w_j}(\mathring{M}^{w_\ell})^{\mathsf{T}}\mathring{M}^{w_j}E_1\right\rangle = \left\langle 2\frac{\mathrm{d}\mathring{M}^{w_j}}{\mathrm{d}t}(\mathring{M}^{w_\ell})^{\mathsf{T}}E_2\right\rangle = \frac{\mathrm{d}\theta}{\mathrm{d}t} + O(N^{-1/2}).$$

Thus the second and third lines of (6.46) can be written as

(6.49)
$$\frac{\mathrm{d}\theta}{\mathrm{d}t}\frac{1+2\theta}{\theta}\frac{1}{2N}\left(\boldsymbol{L}^{(j)}+\sum_{\ell}^{(j)}\boldsymbol{L}^{(j,\ell)}\right)+O\left(N^{\epsilon}|\theta|^{-1}\left(1+\left|\frac{1}{\theta}\frac{\mathrm{d}\theta}{\mathrm{d}t}\right|\right)\Psi^{6}\right).$$

Notice that the first term of (6.49) is independent of w_j hence harmonic. Finally, by Assumption 6.1 (A1) we conclude that

(6.50)
$$|\theta|^{-1} \left(1 + \left| \frac{1}{\theta} \frac{\mathrm{d}\theta}{\mathrm{d}t} \right| \right) \Psi^6 \lesssim (N^{1/2 - \mathfrak{c}_1} + N^{1/2 - \mathfrak{c}_2}) \Psi^6 \lesssim N^{-1 - c}$$

by taking $c < \min(\mathfrak{c}_1, \mathfrak{c}_2) - 6\delta$. This concludes the proof of Proposition 6.3 when \mathcal{A}_t satisfies (A1).

6.3. Small θ regime. In this section we assume that \mathcal{A}_t satisfies Assumption 6.1 (A2).

Lemma 6.11. Let n, \boldsymbol{m} , and B_d be as in Lemma 5.7. Then (5.28) remains true if we replace $\mathring{M}^{\widetilde{w}_i}$ with $(\mathring{M}^{\widetilde{w}_i})^{\iota_i}$ and add an additional error of $O(N^{\epsilon} ||B_d|| |\theta|^{1/2} \Psi^4)$.

Proof. The proof is again three applications of Lemma 6.7. The main difference from those of Lemmas 5.8 and 6.8 is that, in the analogue of (6.43), we keep all $\mathbb{E}\mathfrak{Y}_{|\boldsymbol{m}|+n+1}^{\min}(B_d \mathring{M}^{\widetilde{w}_1})$ terms by expressing it as a finite sum of

(6.51)
$$\langle B_d(\mathring{M}^{\widetilde{w}_1})^{\iota_1}(\mathring{M}^{w'_1})^{\iota'_1}(\mathring{M}^{w'_2})^{\iota'_2}(\mathring{M}^{w'_3})^{\iota'_4}F_n\rangle \mathbb{E}\mathfrak{X}_4,$$

where w' and ι' vary respectively in $\{w_1, \ldots, w_k\}$ and $\{1, \intercal\}$. Then the deterministic prefactor is

(6.52)
$$\langle B_d(\mathring{M}^{\widetilde{w}_1})^{\iota_1}(\mathring{M}^{w'_1})^{\iota'_1}(\mathring{M}^{w'_2})^{\iota'_2}(\mathring{M}^{w'_3})^{\iota'_4}F_n\rangle = \langle B_d(\mathring{M}^0)^4F_n\rangle + O(||B_d||(\sqrt{\operatorname{Re}\theta} + N^{-1/2})),$$

where we used Cauchy-Schwarz and

(6.53)
$$\langle |\mathring{M}^{0} - (\mathring{M}^{0})^{\mathsf{T}}|^{2} \rangle = 2 \operatorname{Re} \theta$$

due to (6.4). This proves the analogue of (5.27) with an additional error of $|\theta|^{1/2}\Psi^4$, from which the result follows by solving for $\mathbb{E}\mathfrak{X}_4$.

Proof of Proposition 6.3, (A2). As before, we focus on the proof of (6.29). Applying Lemma 6.11 to each term of (6.35), we have

$$\mathbb{E}\left\langle \frac{\mathrm{d}(\mathbb{E}H_{t}^{w_{j}}-\mathrm{i}\eta_{t})}{\mathrm{d}t}G^{w_{j}}E_{1}\right\rangle \boldsymbol{L}^{(j)}$$

$$=(\mathrm{right-hand\ side\ of\ (5.31))}$$

$$+\sum_{\ell}\left(\langle 2B_{d}^{w_{j}}(\mathring{M}^{w_{\ell}})^{\intercal}\mathring{M}^{w_{j}}E_{1}\rangle - \frac{\langle 2B_{d}^{w_{j}}(\mathring{M}^{0})^{4}E_{1}\rangle}{I_{4}}(\langle 2(\mathring{M}^{w_{\ell}})^{\intercal}\mathring{M}^{w_{j}}E_{1}\rangle - 1)\right)$$

$$\times \mathbb{E}\frac{\langle (G^{w_{\ell}})^{\intercal}E_{2}G^{w_{j}}E_{1}\rangle}{N}\boldsymbol{L}^{(j,\ell)}$$

$$+\sum_{\ell}\sum_{\ell}\frac{1}{2N}\left(\langle 2B_{d}^{w_{j}}(\mathring{M}^{w_{j}})^{\intercal}\mathring{M}^{w_{j}}E_{1}\rangle - \frac{\langle 2B_{d}^{w_{j}}(\mathring{M}^{0})^{4}E_{1}\rangle}{I_{4}}\right)\mathbb{E}\boldsymbol{L}^{(j,\ell)}$$

$$+O(N^{\epsilon}\Psi^{5}) + O(N^{-1/4-\mathfrak{c}_{1}/2+\epsilon}\Psi^{4}),$$

where we used the first assertion of (A2) to estimate $\sqrt{\operatorname{Re}\theta} \leq N^{-1/4+\mathfrak{c}_1/2}$. By choosing small enough $\delta, c > 0$, we find that the error in (6.54) is $O(N^{-1-c})$. Also, the third term of (6.54) is independent of w_{ℓ} , hence its real part is harmonic.

For the second term of (6.54), we first focus on the case when $\ell = j$. Note that

$$\langle 2B_d^w(\mathring{M}^w)^{\mathsf{T}}\mathring{M}^w E_1 \rangle = -\frac{1}{2} \frac{\mathrm{d}}{\mathrm{d}t} \left\langle \frac{1}{(\mathcal{A} - N^{-1/2}\gamma^{-1}w)(\mathcal{A} - N^{-1/2}\gamma^{-1}w)^{\mathsf{T}}} \right\rangle$$

$$= \frac{1}{2} \frac{\mathrm{d}}{\mathrm{d}t} \theta - N^{-1/2}w \frac{\mathrm{d}}{\mathrm{d}t} \left[\gamma^{-1} \left\langle \frac{1}{\mathcal{A}^2 \mathcal{A}^{\mathsf{T}}} \right\rangle \right] + O(N^{-1} || \mathrm{d}\mathcal{A}_t / \mathrm{d}t ||)$$

$$= \frac{1}{2} \frac{\mathrm{d}}{\mathrm{d}t} \theta + N^{-1/2}w \frac{\mathrm{d}[I_4^{1/2}]}{\mathrm{d}t} + O(N^{-1/2-\mathfrak{c}_2}),$$

where in the third equality we used the last assertion of (A2) and that

(6.56)
$$\left\langle \frac{1}{\mathcal{A}^2 \mathcal{A}^{\mathsf{T}}} \right\rangle - I_3 = \left\langle \frac{1}{\mathcal{A}^*} \Im[\mathcal{A}] \frac{1}{\mathcal{A}^2 \mathcal{A}^{\mathsf{T}}} \right\rangle.$$

Similarly, we have the two estimates

(6.57)
$$\langle 2(\mathring{M}^w)^{\mathsf{T}} \mathring{M}^w E_1 \rangle - 1 = -\theta - 2N^{-1/2} I_4^{1/2} w + O(N^{-1}), \\ \langle 2B_d^w (\mathring{M}^0)^4 E_1 \rangle = -\frac{1}{4} \frac{\mathrm{d}I_4}{\mathrm{d}t} + O(N^{-1/2}).$$

Therefore we have

$$(6.58) \qquad \begin{aligned} \langle 2B_d^{w_j}(\mathring{M}^{w_\ell})^{\mathsf{T}}\mathring{M}^{w_j}E_1 \rangle &- \frac{\langle 2B_d^{w_j}(\mathring{M}^0)^4E_1 \rangle}{I_4} (\langle 2(\mathring{M}^{w_\ell})^{\mathsf{T}}\mathring{M}^{w_j}E_1 \rangle - 1) \\ &= \frac{1}{2}\frac{\mathrm{d}\theta}{\mathrm{d}t} - \frac{1}{4I_4}\frac{\mathrm{d}I_4}{\mathrm{d}t}\theta + N^{-1/2} \left(\frac{\mathrm{d}[I_4^{1/2}]}{\mathrm{d}t} - \frac{1}{2}I_4^{-1/2}\frac{\mathrm{d}I_4}{\mathrm{d}t}\right)w + O(N^{-1/2-\mathfrak{c}_2}) \\ &= \frac{1}{2I_4^{1/2}}\frac{\mathrm{d}(I_4^{-1/2}\theta)}{\mathrm{d}t} + O(N^{-1/2-\mathfrak{c}_2}) = O(N^{-1/2-\mathfrak{c}_2}). \end{aligned}$$

This shows that the second term of (6.54) for $\ell = j$ is bounded by

(6.59)
$$O(N^{-1/2-\mathfrak{c}_2}) \left| \mathbb{E} \frac{\langle (G^{w_j})^{\intercal} E_2 G^{w_j} E_1 \rangle}{N} \mathbf{L}^{(j)} \right| \lesssim N^{-1/2-\mathfrak{c}_2+\epsilon} \Psi^2 = N^{-1-\mathfrak{c}_2+2\delta+\epsilon} \le N^{-1-c}.$$

The sum over $\ell \neq j$ can be dealt with a similar symmetrization as in (5.38) – (5.42), leading to $O(N^{-1-c})$. This concludes the proof of Proposition 6.3 under Assumption 6.1 (A2).

7. Proof of Proposition 4.5

In this section we prove Proposition 4.5, which compares the two deformed non-Hermitian ensembles A + X and $A + X^{\operatorname{Gin}(\mathbb{F})}$. For the sake of brevity we explain only how the proof follows exactly along the same lines as [37]. All estimates in this section are uniform over $z_0 \in \mathcal{D}_C$ defined in (2.6) and X, A satisfying (2.1), (2.2).

Proof of Proposition 4.5. Since the same argument applies to both $\mathbb{F} = \mathbb{R}$ and $\mathbb{F} = \mathbb{C}$, we omit the superscript (\mathbb{F}) throughout the proof.

We consider yet another interpolation $A + X_t$ between A + X and $A + X^{\text{Gin}}$, where $(X_t)_{t>0}$ is the matrix-valued Ornstein–Uhlenbeck process (running from t = 0 to ∞ , unlike $(\mathcal{A}_t)_{t \in [0,1]}$ in Section 4.2)

(7.1)
$$dX_t = -\frac{1}{2}X_t + \frac{1}{\sqrt{N}}dB_t, \qquad X_0 = X_t$$

and the entries of $B_t \in \mathbb{F}^{N \times N}$ are i.i.d. standard Brownian motions. As in Section 4.2, we write

$$G_{t}^{\mathrm{OU},w}(\mathrm{i}\eta) := \begin{pmatrix} -\mathrm{i}\eta & (A + X_{t} - z_{0} - N^{-1/2}w) \\ (A + X_{t} - z_{0} - N^{-1/2}w)^{*} & -\mathrm{i}\eta \end{pmatrix}^{-1},$$

$$(7.2) \qquad L_{t}^{\mathrm{OU}}(w) := \mathrm{Tr}\log(|A + X_{t} - z_{0} - N^{-1/2}w|^{2} + \eta_{0}^{2}) - \mathrm{Tr}\langle\log(|A + x - z_{0} - N^{-1/2}w|^{2} + \eta_{0}^{2})\rangle_{\mathcal{M}},$$

$$N_{t}^{\mathrm{OU}}(w) := \frac{1}{c_{0}} \left\langle \frac{\eta_{0}}{|A + X_{t} - z_{0} - \gamma_{0}^{-1}N^{-1/2}w|^{2} + \eta_{0}^{2}} \right\rangle = \frac{1}{c_{0}} \langle \mathrm{Im} \, G_{t}^{\mathrm{OU},w}(\mathrm{i}\eta_{0})\rangle,$$

where the last equality in the last line is the same identity as in (4.41). Since the first moment $A - z_0 - N^{-1/2}w$ of our ensemble never moves, there is no need to define any dynamics in z, A, γ or c unlike in Section 4.2.

Since the local law Theorem 3.3 immediately applies to $G_t^{OU,w}$ and the first two joint cumulants of $(A + X_t)_{ij}$ remains constant over $t \ge 0$, the proof of Proposition 4.5 follows nearly exactly [37]. To be precise, following the same¹⁴ cumulant expansions as in the proof of [37, Lemma 2], we find that for some constant c > 0 and any $\xi > 0$

(7.3)
$$\left| \frac{\mathrm{d}}{\mathrm{d}t} \mathbb{E} \prod_{j=1}^{k} L_{t}^{\mathrm{OU}}(w_{j}) \right| \lesssim \mathrm{e}^{-3t/2} \frac{N^{\xi}}{N^{5/2} \eta_{0}^{3}}$$

holds uniformly over $t \ge 0$. Similarly, the same proof as [37, Proposition 3] gives for any $\xi > 0$ that

(7.4)
$$\left|\frac{\mathrm{d}}{\mathrm{d}t}\mathbb{E}N_t^{\mathrm{OU}}(w)\right| = \frac{1}{c_0} \left|\frac{\mathrm{d}}{\mathrm{d}t}\mathbb{E}\langle G_t^{\mathrm{OU},w}(\mathrm{i}\eta_0)\rangle\right| \lesssim \mathrm{e}^{-3t/2} \frac{N^{\xi}}{N^{7/2}\eta_0^4}$$

¹⁴The quantity $L_t^{OU}(w)$ is a slight variant of $I_3(t)$ defined in [37, Eq. (36)]; in $I_3(t)$ the variable w is integrated out and that the logarithm is cut-off at N^{100} . These differences do not affect the proof.

uniformly over $t \ge 0$. Integrating (7.3) and (7.4) over t completes the proof.

Remark 7.1 (Universality of the least singular value). The same argument as in the proof of Theorem 4.1 can be used to prove the universality for the smallest singular value λ_1 of $A + X - z_0$. Again, we break the comparison up into:

- (1) Comparing the smallest singular values $A + X^{\text{Gin}}$ and X^{Gin} .
- (2) Comparing the smallest singular values of A + X and $A + X^{Gin}$.

For (1) our proofs of (5.6) and (6.30) can be adapted, beyond the lower-tail estimate as in (4.12). More precisely, we can prove that for each fixed $w = O(N^{-1/2})$ and continuous and compactly supported test function F on \mathbb{R}

(7.5)
$$\mathbb{E}F(\lambda_1(c_0(A+X-z_0-\gamma_0^{-1}w))) = \mathbb{E}F(\lambda_1(X-1-w)) + o(1).$$

A rough sketch of the proof is as follows: Instead of $\mathbb{E}c_0^{-1}\langle G_0(i\eta_0)\rangle$, we take an arbitrary moment (or a more general function) of the regularized counting function

(7.6)
$$\int_0^{E_1} c_0^{-1} \operatorname{Im} \operatorname{Tr}[G_0^w(c_0^{-1}(E+\mathrm{i}\eta_1)] \mathrm{d}E,$$

where $E_1 = O(N^{-3/4+\epsilon})$ lives on the same scale as the typical eigen-spacing around λ_1 . Then it suffices to prove that the time-derivative of this statistic is o(1). To this end, we need to generalize the local law and the content of Section 5 for spectral parameter $\hat{z}_t := c_t^{-1}(E + i\eta_1)$ beyond $\eta_t := c_t^{-1}\eta_1$. Such a generalization does not require a significant changes in Section 5, since \hat{z}_t is close enough to the imaginary axis that the cubic equation (3.10) associated to the MDE remains almost intact. For (2) similarly to the proof of Proposition 4.5 we can compare along an Ornstein–Uhlenbeck process and use the slightly extended local law described above to control the time derivative of moments of the regularized counting function. This same approach can also be applied to the p-th singular value for fixed $p \in \mathbb{N}$.

To adapt the zig-zag approach to Theorem 3.3 to spectral parameters $\hat{z} = E + i\eta$ off the imaginary axis the key is to correctly adjust the zig step for the spectral parameter. The dynamics would

$$\frac{\mathrm{d}\widehat{z}_s}{\mathrm{d}s} = -\frac{\widehat{z}_s}{2} - \langle M(\widehat{z}_s) \rangle$$

Again, because (3.10) remains essentially intact for small E the flow in the zig step will actually only slightly change.

8. ZIG-ZAG STRATEGY: PROOF OF THEOREM 3.3

In this section we prove Theorem 3.3. All estimates in this section are uniform over X, A, z_0 satisfying (2.1), (2.2), and $||(A-z_0)^{-1}|| \le C_4$. It is worth noting, as we only consider the local law for the argument of G and M on the imaginary axis, that from (3.5) we have $||M|| \le C$, for some N-independent C > 0. For any z such that $|z - z_0| < c$ and for every T > 0 we define the following dynamics for $t \in [0, T]$.

Definition 8.1 (Matrix Ornstein–Uhlenbeck flow and the η -flow). Define the matrix-valued Ornstein– Uhlenbeck (OU) process X_t as

(8.1)
$$\mathrm{d}X_t = -\frac{1}{2}X_t\mathrm{d}t + \frac{1}{\sqrt{N}}\mathrm{d}B_t, \qquad X_0 = X,$$

where the entries of $B_t \in \mathbb{C}^{N \times N}$ are *i.i.d.* standard Brownian motions. We define \mathcal{A}_t as ¹⁵

(8.2)
$$\mathcal{A}_t := e^{-\frac{t}{2}} \mathcal{A}_0, \qquad \mathcal{A}_0 = A - z,$$

and define the resolvent

(8.3)
$$G_t(\widehat{z}) := (\mathcal{H}_{\mathcal{A}_t + X_t} - \widehat{z})^{-1}, \qquad \widehat{z} \in \mathbb{C} \setminus \mathbb{R},$$

where for a matrix B

$$\mathcal{H}_B := \begin{pmatrix} 0 & B \\ B^* & 0 \end{pmatrix}.$$

Note then that $A_t + X_t$ satisfies the SDE

(8.4)
$$d(\mathcal{A}_t + X_t) = -\frac{1}{2}(\mathcal{A}_t + X_t)dt + \frac{1}{\sqrt{N}}dB_t.$$

¹⁵The evolutions \mathcal{A}_t and η_t in this section are not related to those in Section 4.2.

For $\eta > 0$, we define $\eta_t \in \mathbb{R}$ to be the solution of the ODE

(8.5)
$$\frac{\mathrm{d}\eta_t}{\mathrm{d}t} = -\frac{\eta_t}{2} - \langle \mathrm{Im}\, M_{\mathcal{A}_t}(\mathrm{i}\eta_t) \rangle, \qquad \eta_0 = \eta.$$

Lemma 8.2. $M_t \equiv M_{\mathcal{A}_t}(i\eta_t)$ satisfies

(8.6)
$$\frac{\mathrm{d}\langle M_t \rangle}{\mathrm{d}t} = \frac{\langle M_t \rangle}{2}$$

Proof. To see (8.6) note that M_t and the Hermitization $\mathcal{H}_{\mathcal{A}_t}$ of \mathcal{A}_t satisfy

(8.7)
$$1 = (\mathcal{H}_{\mathcal{A}_t} - i\eta_t)M_t - \langle M_t \rangle M_t,$$

and then we take the time derivative to find that

(8.8)
$$\left(\frac{\langle M_t \rangle}{2} - \left\langle \frac{\mathrm{d}M_t}{\mathrm{d}t} \right\rangle\right) M_t - \frac{1}{M_t} \left(\frac{M_t}{2} - \frac{\mathrm{d}M_t}{\mathrm{d}t}\right) = 0.$$

Multiplying both sides by M_t , taking the trace, and using $|\langle M_t^2 \rangle| \leq \langle |M_t|^2 \rangle < 1$, we obtain (8.6).

We define the time at which the characteristic curve η_t touch the real axis

(8.9)
$$T^*(\mathcal{A}_0, \eta_0) := \sup\{t : \operatorname{sgn}(\eta_t) = \operatorname{sgn}(\eta_0)\}$$

We will also use the short-hand notation $\rho_t := \rho^{\mathcal{A}_t}(i\eta_t)$.

Proposition 8.3 (Zig step). Fix small constants $c, \epsilon > 0$. Fix $\mathcal{A}_0 \in \mathbb{C}^{N \times N}$ with $\|\mathcal{A}_0^{-1}\| \leq C$ for some C > 0, and pick η_0 such that $0 < |\eta_0|/\rho_0 \leq c$. For $t \leq T^*(\mathcal{A}_0, \eta_0)$, let $\mathcal{A}_t := e^{-t/2}\mathcal{A}_0$, and let η_t be the solution of (8.5) with initial condition η_0 . Define $G_t(i\eta_t) := (\mathcal{H}_{\mathcal{A}_t+X_t} - i\eta_t)^{-1}$. For any $\alpha \geq 0$ and some small $0 < \gamma \leq \epsilon/10$, assume that, with very high probability, we have (8.10)

$$|\langle B(G_0(\mathrm{i}\eta_0) - M_{\mathcal{A}_0}(\mathrm{i}\eta_0))\rangle| \leq \frac{N^{\gamma}(\log N)^{\alpha}}{N\eta_0}, \qquad |\boldsymbol{u}^*\left(G_0(\mathrm{i}\eta_0) - M_{\mathcal{A}_0}(\mathrm{i}\eta_0)\right)\boldsymbol{y}| \leq N^{\gamma}(\log N)^{\alpha}\sqrt{\frac{\rho_0}{N\eta_0}}.$$

uniformly in unit vectors $\boldsymbol{u}, \boldsymbol{y} \in \mathbb{C}^{2N}$ and matrices $B \in \mathbb{C}^{2N \times 2N}$ with $||B|| \leq 1$. Then, (8.11)

$$|\langle B(G_t(i\eta_t) - M_{\mathcal{A}_t}(i\eta_t))\rangle| \leq \frac{N^{\gamma}(\log N)^{\alpha+1}}{N\eta_t}, \qquad |\boldsymbol{u}^*\left(G_0(i\eta_t) - M_{\mathcal{A}_t}(i\eta_t)\right)\boldsymbol{y}| \leq N^{\gamma}(\log N)^{\alpha+1}\sqrt{\frac{\rho_t}{N\eta_t}},$$

with very high probability, uniformly in $t \leq T^*(\mathcal{A}_0, \eta_0)$, $N\eta_t \rho_t \geq N^{\epsilon}$, unit vectors $\boldsymbol{u}, \boldsymbol{y} \in \mathbb{C}^{2N}$ and matrices $B \in \mathbb{C}^{2N \times 2N}$ with $\|B\| \leq 1$.

We remark that, even if stated together, the average local law bounding $|\langle B(G_t(i\eta_t) - M_{\mathcal{A}_t}(i\eta_t))\rangle|$ in Proposition 8.3 can be proven independently of the isotropic local law bounding $|\boldsymbol{u}^*(G_0(i\eta_t) - M_{\mathcal{A}_t}(i\eta_t))\boldsymbol{y}|$.

Fix $\mathcal{A} = A - z \in \mathbb{C}^{N \times N}$, and recall that $\|\mathcal{A}^{-1}\| \leq C$. In the following for given $\eta > 0$ and $r \geq 1$ we define the *scale*

(8.12)
$$\mathfrak{s}(\eta, r) := \frac{\eta}{\rho^{r\mathcal{A}}(\mathrm{i}\eta)} = \left\langle \left\langle \frac{1}{|r\mathcal{A} + \boldsymbol{x}|^2 + \eta^2} \right\rangle_{\mathcal{M}} \right\rangle^{-1}$$

Note that the map $\eta \mapsto \mathfrak{s}(\eta, r)$ is monotone increasing. Thus, given $r \geq 1$, there is a one to one correspondence between η and $\mathfrak{s}(\eta, r)$. In particular, when we say that a local law holds on some given scale \mathfrak{s}^* , we mean that it holds for spectral parameter η determined by $\mathfrak{s}(\eta, r) = \mathfrak{s}^*$. Fix any $\epsilon > 0$, and define the scale $\mathfrak{s}_{\text{fin}} = \mathfrak{s}(\eta_{\text{fin}}, 1)$ so that

(8.13)
$$N\eta_{\rm fin}\rho^{\mathcal{A}}(\mathrm{i}\eta_{\rm fin}) := N^{\epsilon}.$$

Proposition 8.4 (Zag step). Fix a constant $\alpha \geq 0$, let $\epsilon > 0$ from (8.13), fix any small $0 < \gamma \leq \epsilon/10$, and pick $\epsilon_0 > 0$. Fix a scale $\mathfrak{s}_{\text{fin}} < \tilde{\mathfrak{s}}_0 = \mathfrak{s}(\tilde{\eta_0}, r) \leq N^{-\epsilon_0}$, and let $t \leq \tilde{\mathfrak{s}}_0$. Consider an $\tilde{\eta_1}$ such that $\eta_{\text{fin}} \vee \tilde{\eta_0} N^{-\epsilon_0/100} \leq \tilde{\eta_1} \leq \tilde{\eta_0}$ and its corresponding scale $\tilde{\mathfrak{s}}_1 = \mathfrak{s}(\tilde{\eta_1}, r)$ also satisfies $\mathfrak{s}_{\text{fin}} \vee N^{-\epsilon_0/200} \tilde{\mathfrak{s}}_0 \leq \tilde{\mathfrak{s}}_1 \leq \tilde{\mathfrak{s}}_0$. Define $G^t(\mathrm{i}\eta) := (\mathcal{H}_{r,A+X_t} - \mathrm{i}\eta)^{-1}$, and assume that, uniformly in $r \in [1, 2]$, unit vectors $\boldsymbol{u}, \boldsymbol{y} \in \mathbb{C}^{2N}$, and matrices $B \in \mathbb{C}^{2N \times 2N}$ with $\|B\| \leq 1$, the local law

(8.14)
$$|\langle B(G^s(i\eta) - M_{r\mathcal{A}}(i\eta))\rangle| \leq \frac{N^{\gamma}(\log N)^{\alpha}}{N\eta}, \qquad |\boldsymbol{u}^*(G^s(i\eta) - M_{r\mathcal{A}}(i\eta))\boldsymbol{y}| \leq N^{\gamma}(\log N)^{\alpha}\sqrt{\frac{\rho^{r\mathcal{A}}(i\eta)}{N\eta}}$$

holds with very high probability for any time $s \in [0,t]$ on the scale $\tilde{\mathfrak{s}}_0$, and at time s = t on the scale $\tilde{\mathfrak{s}}_1$. Then, (8.14) also holds on the scale $\tilde{\mathfrak{s}}_1$ for any $s \in [0,t]$.

Note that the deformation \mathcal{A} and the spectral parameter $i\eta$ are time-dependent in Proposition 8.3, while they are time-independent in Proposition 8.4 but \mathcal{A} carries an extra rescaling parameter r and the estimate is uniform in r. This important subtlety is expressed by the lower and upper time indices G_t and G^t in the two propositions above. In the application of Proposition 8.4 for proving Theorem 3.3, r will be a frozen time parameter from the previous step.

Our goal is now to prove (8.14) for r = 1 and any η such that $\mathfrak{s}(\eta, 1) \geq \mathfrak{s}_{\text{fin}}$; this would give Theorem 3.3. Fix $\epsilon_0 > 0$, and define $T := N^{-\epsilon_0}$. Then, by standard ODE theory, there exists $\eta_{\text{in}} = \eta_{\text{in}}(\mathcal{A}, T)$ such that $\mathfrak{s}_{\text{in}} = \mathfrak{s}(\eta_{\text{in}}, e^{T/2}) \sim T$ and $\mathfrak{s}(\eta_T, 1) = \mathfrak{s}_{\text{fin}}$, with η_t being the solution of (8.5) with initial condition η_{in} . In particular, $T \leq T^*(e^{T/2}\mathcal{A}, \eta_{\text{in}})$; here we also used that the map $t \mapsto \eta_t$ is decreasing. Note that the η_t -dynamics depends on \mathcal{A}, T through $M_{e^{(T-t)/2}\mathcal{A}}$. Next, we define a decreasing sequence of scales

(8.15)
$$\mathfrak{s}_k := \mathfrak{s}_{\mathrm{in}} N^{-k\epsilon_0/100} \vee \mathfrak{s}_{\mathrm{fin}}$$

Note that $\mathfrak{s}_0 = \mathfrak{s}_{in}$, and let K be the smallest index such that $\mathfrak{s}_K = \mathfrak{s}_{fin}$, i.e. the last step may be a bit smaller than all the previous ones. We define $0 \le t_k \le T$ implicitly via

(8.16)
$$\mathfrak{s}(\eta_{t_k}, e^{(T-t_k)/2}) = \mathfrak{s}_k;$$

in particular, $t_K = T$ as a consequence of $\mathfrak{s}_K = \mathfrak{s}_{in} = \mathfrak{s}(\eta_T, 1)$. Since η_t is decreasing and $\rho^{e^{-t/2}\mathcal{A}}(i\eta_t)$ is increasing in t by (8.5) and (8.6), we find that

(8.17)
$$\mathfrak{s}(\eta_t, \mathrm{e}^{(T-t)/2}) = \frac{\eta_t}{\rho^{\mathrm{e}^{(T-t)/2}\mathcal{A}}(\mathrm{i}\eta_t)}$$

is decreasing in t. In particular, the implicit equation (8.16) has a unique solution and t_k is an increasing sequence. Further, we define domains

(8.18)
$$\mathcal{D}_k := \left\{ (\eta, r) : \eta \ge \eta_{t_k}, \ r \in [1, e^{(T - t_k)/2}] \right\}, \qquad k = 0, \dots, K.$$

Note that

(8.19)
$$\frac{\eta_{t_{k+1}}}{\eta_{t_k}} = \frac{\mathfrak{s}_{k+1}}{\mathfrak{s}_k} \frac{\rho^{\mathrm{e}^{(T-t_k)/2}\mathcal{A}}(\mathrm{i}\eta_{t_k})}{\rho^{\mathrm{e}^{(T-t_{k+1})/2}\mathcal{A}}(\mathrm{i}\eta_{t_{k+1}})} = N^{-\epsilon_0/100} \mathrm{e}^{-(t_{k+1}-t_k)/2} \ge N^{-\epsilon_0/100} \mathrm{e}^{-T}.$$

We say that the local law holds at a tuple (η, r) with error (γ, α) if (8.14) holds. We are now ready to prove Theorem 3.3.

Proof of Theorem 3.3. We fix some z such that $|z - z_0| < c$ and $\epsilon > 0$. As above, we define the scale $\mathfrak{s}_{\text{fin}} = \mathfrak{s}(\eta_{\text{fin}}, 1)$ with $\mathcal{A} = \mathcal{A} - z$. We additionally fix $\epsilon_0 > 0$, and define $T := N^{-\epsilon_0}$, η_{in} , \mathfrak{s}_{in} , \mathfrak{s}_k , K, and \mathcal{D}_k as above. We first establish that (8.14) holds on \mathcal{D}_K . Extending (8.14) to hold uniformly over $|z - z_0| < c$ then follows from a simple union bound, and extending to $\eta \geq N^{-1}$ completely analogously to [37, Appendix A].

From [47, Theorem 2.1] we know that (8.14) holds with very high probability uniformly over \mathcal{D}_0 with error $(\epsilon_0, 0)$. We now want to conclude a local law on \mathcal{D}_k by induction. The main inductive step is formalized in the following proposition, whose proof is presented after we finish the proof of Theorem 3.3.

Proposition 8.5. Fix $k \in \{0, ..., K-1\}$, and assume that (8.14) holds on \mathcal{D}_k with error (ϵ_0, k) . Then (8.14) holds on \mathcal{D}_{k+1} with error $(\epsilon_0, k+1)$.

It then follows from K applications of Proposition 8.5 that (8.14) holds with very high probability uniformly over \mathcal{D}_K with error (ϵ_0, K) . This completes the proof.

Proof of Proposition 8.5. Fix some $(\eta, r) \in \mathcal{D}_{k+1} \setminus \mathcal{D}_k$ and set $\widehat{\eta} = \eta_{t_k}$. We note, by the definition of our domains, for the solution $\widehat{\eta}_t$ to (8.5) with initial condition $\widehat{\eta}_0 = \widehat{\eta}$ satisfies $(\eta, r) = (\widehat{\eta}_t, r)$ and $(\widehat{\eta}, e^{\widehat{t}/2}r) \in \mathcal{D}_k$ for some $0 < \widehat{t} \le t_{k+1} - t_k$. By assumption, (8.14) holds with very high probability on the scale $\mathfrak{s}(\widehat{\eta}, e^{\widehat{t}r})$ with error (ϵ_0, k) . Then, by Proposition 8.3, (8.14) holds on the scale $\mathfrak{s}(\eta, r)$ for $s = \widehat{t}$ with error $(\epsilon_0, k+1)$. Additionally, (8.14) holds on the scale $\mathfrak{s}(\widehat{\eta}, r)$ uniformly over $s \in [0, \widehat{t}]$. We then aim at applying Proposition 8.4 with the choices $(\widetilde{\eta}_0, \widetilde{\eta}_1) = (\widehat{\eta}, \eta)$ and $(\widetilde{\mathfrak{s}}_0, \widetilde{\mathfrak{s}}_1) = (\mathfrak{s}(\widehat{\eta}, r), \mathfrak{s}(\eta, r))$, for which we immediately have

(8.20)
$$1 \ge \frac{\tilde{\eta}_1}{\tilde{\eta}_0} = \frac{\eta}{\hat{\eta}} \ge \frac{\eta_{t_{k+1}}}{\eta_{t_k}} \gtrsim N^{-\epsilon_0/100},$$

where the second and third inequalities are due to $\eta \geq \eta_{t_{k+1}}$ and (8.19). Thus we only need to check $\tilde{\mathfrak{s}}_1 \geq N^{-\epsilon_0/200}\tilde{\mathfrak{s}}_0$. Indeed, we have $1 \geq (\tilde{\mathfrak{s}}_1/\tilde{\mathfrak{s}}_0) \gtrsim (\eta/\tilde{\eta})^{2/3} \geq (\eta_{t_{k+1}}/\eta_{t_k})^{2/3} \gtrsim N^{-\epsilon_0/150}$, where the only

non-trivial inequality $(\tilde{\mathfrak{s}}_1/\tilde{\mathfrak{s}}_0) \gtrsim (\eta/\hat{\eta})^{2/3}$ follows from Lemma 3.4 as

$$(8.21) \qquad \qquad \widetilde{\widetilde{\mathfrak{s}}}_{1} = \frac{\mathfrak{s}(\eta, r)}{\mathfrak{s}(\widehat{\eta}, r)} = \frac{\eta}{\widehat{\eta}} \cdot \frac{\rho^{r\mathcal{A}}(\mathrm{i}\widehat{\eta})}{\rho^{r\mathcal{A}}(\mathrm{i}\eta)} \sim \begin{cases} \frac{\eta}{\widehat{\eta}} \cdot \frac{\langle |r\mathcal{A}|^{-2} - 1\rangle^{1/2} + \widehat{\eta}^{1/3}}{\langle |r\mathcal{A}|^{-2} - 1\rangle^{1/2} + \eta^{1/3}} & \text{if } \langle |r\mathcal{A}|^{-2} \rangle \ge 1, \\ \frac{\langle 1 - |r\mathcal{A}|^{-2} \rangle + \eta^{2/3}}{\langle 1 - |r\mathcal{A}|^{-2} \rangle + \widehat{\eta}^{2/3}} & \text{if } \langle |r\mathcal{A}|^{-2} \rangle < 1. \end{cases}$$

Consequently, (8.14) holds with very high probability on the scale $\mathfrak{s}(\eta, r)$ uniformly over $s \in [0, \hat{t}]$. Union bounding over a sufficiently dense grid of $\mathcal{D}_{k+1} \setminus \mathcal{D}_k$ completes the proof.

8.1. **Proof of Proposition 8.3.** In what follows we will prove both the average local law and the isotropic local law for matrices with some Gaussian component. For simplicity we present first the proof of with B = I and the isotropic law, then explain the short adjustment for general B. Fix $0 < \gamma < \epsilon/10$, $\alpha \ge 0, \eta_0 > 0, A_0 \in \mathbb{C}^{N \times N}$, and unit vectors in $\boldsymbol{u}, \boldsymbol{y} \in \mathbb{C}^{2N}$. Define the stopping time

(8.22)
$$\tau := T^*(\mathcal{A}_0, \eta_0) \wedge \inf \left\{ t > 0 : |\langle G_t(i\eta_t) - M_{\mathcal{A}_t}(i\eta_t) \rangle| \ge C_* \frac{N^{\gamma}(\log N)^{\alpha+1}}{N\eta_t} \right\}$$
$$\wedge \inf \left\{ s > 0 : |\boldsymbol{u}^*(G_t(i\eta_t) - M_{\mathcal{A}_t}(i\eta_t))\boldsymbol{y}| \ge C_* N^{\gamma}(\log N)^{\alpha+1} \sqrt{\frac{\rho^{\mathcal{A}_t}(i\eta_t)}{N\eta_t}} \right\},$$

for some large constant $C_* > 0$. To simplify notation we will write $G_t \equiv G_t(i\eta_t)$, $M_t \equiv M_{\mathcal{A}_t}(i\eta_t)$, and $T \equiv T^*(\mathcal{A}_0, \eta_0)$ throughout this section. By Itô's formula

(8.23)
$$d\langle G_t - M_t \rangle = \left(\frac{1}{2}\langle G_t - M_t \rangle + \langle G_t - M_t \rangle \langle G_t^2 \rangle + \frac{\langle G_t^3 \rangle}{N} \mathbb{1}(\beta = 1)\right) dt$$
$$- \frac{1}{N^{3/2}} \sum_{i,j \in [\![2N]\!]} \left[\left(G_t^2\right)_{ij} + \left(G_t^2\right)_{ji} \mathbb{1}(\beta = 1) \right] d(\mathfrak{B}_t)_{ij}$$
$$=: \mathcal{E}_1(t) \langle G_t - M_t \rangle dt + \frac{\langle G_t^3 \rangle}{N} \mathbb{1}(\beta = 1) dt + d\mathcal{E}_2(t)$$

where $\beta = 1$ if the entries of X are real and $\beta = 2$ if the entries of X are complex, and

$$\mathfrak{B}_t = \begin{pmatrix} 0 & B_t \\ B_t^* & 0 \end{pmatrix}$$

Similarly, we have

(8.24)
$$d\boldsymbol{u}^{*}(G_{t} - M_{t}))\boldsymbol{y} = \left(\frac{1}{2}\boldsymbol{u}^{*}(G_{t} - M_{t})\boldsymbol{y} + \langle G_{t} - M_{t}\rangle\boldsymbol{u}^{*}G_{t}^{2}\boldsymbol{y} + \frac{1}{N}\boldsymbol{u}^{*}G_{t}^{3}\boldsymbol{y}\mathbbm{1}(\beta = 1)\right)dt$$
$$-\frac{1}{N^{1/2}}\sum_{i,j}(G_{t}\boldsymbol{y}\boldsymbol{u}^{*}G_{t})_{ji} + (G_{t}\boldsymbol{y}\boldsymbol{u}^{*}G_{t})_{ij}\mathbbm{1}(\beta = 1)d(\mathfrak{B}_{t})_{ij}$$
$$=:\frac{1}{2}\boldsymbol{u}^{*}(G_{t} - M_{t})\boldsymbol{y}dt + \mathcal{E}_{1}(t, \boldsymbol{u}, \boldsymbol{y})dt + d\mathcal{E}_{2}(t, \boldsymbol{u}, \boldsymbol{y}).$$

Recall the short-hand notation $\rho_t = \rho^{\mathcal{A}_t}(i\eta_t)$. We start with the estimate of the stochastic terms in (8.23)–(8.24). In the following we will often use the simple integration rule

(8.25)
$$\int_0^t \frac{\rho_s}{\eta_s^p} \,\mathrm{d}s \lesssim \frac{1}{\eta_t^{p-1}},$$

with p > 1, even if not stated explicitly. This follows from simple computations using (8.5).

Lemma 8.6. Uniformly over t in [0,T], and uniformly in $\boldsymbol{u}, \boldsymbol{y}$ deterministic unit vectors, we have

(8.26)
$$\left| \int_0^{t\wedge\tau} \mathrm{d}\mathcal{E}_2(s) \right| \prec \frac{1}{N\eta_{t\wedge\tau}},$$

(8.27)
$$\left| \int_0^{t\wedge\tau} \mathrm{d}\mathcal{E}_2(s,\boldsymbol{u},\boldsymbol{y}) \right| \prec \sqrt{\frac{\rho_{t\wedge\tau}}{N\eta_{t\wedge\tau}}}.$$

Proof. From the definition of τ we have that

(8.28)
$$\langle \operatorname{Im} G_t(i\eta_{t\wedge\tau}) \rangle \leq \left(1 + N^{-\epsilon/2}\right) \rho_{t\wedge\tau}.$$

The quadratic variation in (8.23) satisfies

(8.29)
$$\langle \mathrm{d}\mathcal{E}_2, \mathrm{d}\mathcal{E}_2 \rangle = \frac{1}{N^2} \langle G_t^2 (G_t^*)^2 \rangle \mathrm{d}t \le \frac{1}{N^2 \eta_t^2} \frac{\langle \mathrm{Im} \, G_t \rangle}{\eta_t} \mathrm{d}t$$

Fix $t \in [0, T]$, applying Burkholder-Davis-Gundy inequality to (8.29), by (8.28)

(8.30)
$$\max_{\tilde{t}\in[0,t]} \left| \int_{0}^{\tilde{t}\wedge\tau} \mathrm{d}\mathcal{E}_{2}(s) \right| \prec \frac{1}{N} \left(\int_{0}^{t\wedge\tau} \frac{\rho_{s}}{\eta_{s}^{3}} \mathrm{d}s \right) \lesssim \frac{1}{N^{2} \eta_{t\wedge\tau}^{2}}$$

where in the last inequality we used (8.25). This bound can then be extended uniformly over [0, T] by taking a polynomial sized net of times in [0, T] and union bounding over the very high probability events on which (8.30) holds.

We now move on to the proof of (8.27). For fixed $t \in [0, \tau]$

(8.31)
$$\langle \mathrm{d}\mathcal{E}_2(t, \boldsymbol{u}, \boldsymbol{y}), \mathrm{d}\mathcal{E}_2(t, \boldsymbol{u}, \boldsymbol{y}) \rangle = \frac{(\boldsymbol{u}^* \operatorname{Im} G_t \boldsymbol{y})^2}{N\eta_t^2} \mathrm{d}s \leq \frac{1}{N\eta_t^2} \left(\rho_t + N^{-\epsilon/2} \sqrt{\frac{\rho_t}{N\eta_t}}\right)^2 \mathrm{d}t \lesssim \frac{\rho_t}{N\eta_t^2} \mathrm{d}t$$

where we use (3.5) in the second inequality. The proof then follows completely analogously to $\mathcal{E}_2(s)$. \Box

We now move on to controlling the size of the drift terms $\mathcal{E}_1(t)$, $N^{-1}\langle G_t^3 \rangle$, and $\frac{1}{2}\boldsymbol{u}^*(G_t - M_t)\boldsymbol{y} + \mathcal{E}_1(t, \boldsymbol{u}, \boldsymbol{y})$.

We estimate

(8.32)
$$\int_{0}^{t\wedge\tau} \left|\frac{\langle G_{s}^{3}\rangle}{N}\right| \mathrm{d}s \leq \int_{0}^{s} \frac{\langle \mathrm{Im}\,G_{s}\rangle}{N\eta_{s}^{2}} \,\mathrm{d}s \lesssim \frac{1}{N\eta_{t}}.$$

We also note that

$$(8.33) \qquad |\mathcal{E}_1(t)| \le \frac{1}{2} + \frac{\langle \operatorname{Im} G_t \rangle}{\eta_t} \le \frac{1}{2} + \left(1 + N^{-\epsilon/2}\right) \frac{\rho_t}{\eta_t} = -\frac{1 + O(N^{-\epsilon/2})}{\eta_t} \frac{\mathrm{d}\eta_t}{\mathrm{d}t} =: \widetilde{\mathcal{E}}_1(t).$$

Let Ω be the very high probability event on which (8.10) and (8.26)–(8.27) hold. Integrating (8.23) with respect to time, and using (8.10) to estimate the initial condition, we have that on Ω

(8.34)
$$|\langle G_{t\wedge\tau}(i\eta_{t\wedge\tau}) - M_{t\wedge\tau}\rangle| \leq \int_0^{t\wedge\tau} \widetilde{\mathcal{E}}_1(s) |\langle G_{s\wedge\tau}(i\eta_s) - M_{t\wedge\tau}\rangle| \mathrm{d}s + \frac{N^{\gamma}(\log N)^{\alpha}}{N\eta_{t\wedge\tau}}$$

Applying Grönwall's inequality we get

(8.35)
$$|\langle G_{t\wedge\tau}(\mathrm{i}\eta_{t\wedge\tau}) - M_{t\wedge\tau}\rangle| \leq \int_0^{t\wedge\tau} \left(\frac{N^{\gamma}(\log N)^{\alpha}}{N\eta_s}\right) \widetilde{\mathcal{E}}_1(s) \exp\left[\int_s^{t\wedge\tau} \widetilde{\mathcal{E}}_1(u) \mathrm{d}u\right] \mathrm{d}s.$$

Notice from the definition of $\mathcal{E}_1(s)$ that

(8.36)
$$\int_{s}^{t\wedge\tau} \widetilde{\mathcal{E}}_{1}(u) \mathrm{d}u = (1 + O(N^{-\epsilon/2})) \log \frac{\eta_{s}}{\eta_{t\wedge\tau}}$$

and hence combining this with (8.35) we obtain

(8.37)
$$|\langle G_{t\wedge\tau}(\mathrm{i}\eta_{t\wedge\tau}) - M_{t\wedge\tau}\rangle| \lesssim \frac{N^{\gamma}(\log N)^{\alpha+1}}{N\eta_{t\wedge\tau}}$$

On the event Ω we use (8.31) and (8.37) to estimate $\mathcal{E}_1(s, \boldsymbol{u}, \boldsymbol{y})$ uniformly over $s \in [0, \tau]$ as

(8.38)
$$|\mathcal{E}_1(t, \boldsymbol{u}, \boldsymbol{y})| \lesssim \left| \langle G_t - M_t \rangle \boldsymbol{u}^* G_t^2 \boldsymbol{y} \right| \lesssim \frac{N^{\gamma} (\log N)^{\alpha + 1}}{N \eta_t^2} \rho_t$$

We then integrate (8.24) to conclude that

(8.39)
$$|\boldsymbol{u}^*(G_{t\wedge\tau} - M_{t\wedge\tau})\boldsymbol{y}| \lesssim N^{\gamma} (\log N)^{\alpha+1} \sqrt{\frac{\rho_{t\wedge\tau}}{N\eta_{t\wedge\tau}}}$$

Combining (8.37)–(8.39) we thus conclude that $\tau = T$ holds on Ω .

For general B (with $\beta = 2$ for simplicity of presentation) (8.23) becomes

$$(8.40) \quad \mathrm{d}\langle B(G_t - M_t) \rangle = \left(\frac{1}{2} \langle B(G_t - M_t) \rangle + \langle G_t - M_t \rangle \langle BG_t^2 \rangle \right) \mathrm{d}t - \frac{1}{N^{3/2}} \sum_{i,j \in [\![2N]\!]} (G_t BG_t)_{ij} \, \mathrm{d}(\mathfrak{B}_t)_{ij}$$

Comparing the form of (8.40) with (8.24) one can see that the proof is analogous to (8.39), with (8.37) used as an input to control the term $\langle G_t - M_t \rangle \langle BG_t^2 \rangle$.

We conclude this section with the proof of (3.14).

Proof of (3.14). We prove this bound only for matrices with an (almost) order one Gaussian component, as then this Gaussian component can easily be removed with standard arguments similar to Section 8.2; we omit this for brevity. When $\eta/\rho^z(i\eta) \ge N^{-\epsilon}$, for some very small fixed $\epsilon > 0$, the desired result immediately follows from [47, Theorem 2.1]. We now show that this bound can be propagated down to $\eta = 0$ using the flow (8.23). This proof is now completely analogous to the proof of Proposition 8.3, so we only present the main differences. In the estimate of the various terms in the rhs. of (8.23) we often used the bound $||G|| + ||\Im G|| \le 1/\eta$, however, when we are outside of the spectrum these bound can be improved to

(8.41)
$$\|G\| \le \frac{1}{\kappa}, \qquad \|\Im G\| \le \frac{\eta}{\kappa^2},$$

where we recall that $\kappa := \operatorname{dist}(i\eta, \operatorname{supp}(\rho^z))$. The point is that if z is outside of the (non-Hermitian) spectrum, then the density of the Hermitization, ρ^z , has a gap around zero, so κ is much larger than η .

Fix z_0 , η_0 such that $\kappa_0 = \operatorname{dist}(i\eta_0, \operatorname{supp}(\rho^{z_0})) \gtrsim N^{-\epsilon}$. Next, using the bounds (8.41) we estimate the lhs. of (8.26) and the lhs. of (8.32) by $1/(N\kappa_t)$, with $\kappa_t := \operatorname{dist}(i\eta_t, \operatorname{supp}(\rho^{z_t}))$. Here $z_t = e^{-t/2}z_0$, and η_t is the solution of the characteristic equation (8.5) with initial condition η_0 . For completeness, we present the analog of the bound (8.32) obtained using (8.41):

(8.42)
$$\int_0^{t\wedge\tau} \frac{|\langle G_s^3 \rangle|}{N} \,\mathrm{d}s \lesssim \int_0^{t\wedge\tau} \frac{\langle \Im G_s \rangle ||G_s||}{N\eta_s} \,\mathrm{d}s \prec \int_0^{t\wedge\tau} \frac{\rho_s}{N\eta_s\kappa_s} \,\mathrm{d}s \lesssim \frac{1}{N\kappa_t},$$

where we used that $\kappa_s \gtrsim \kappa_t$. Using (8.41), the changes in the estimate of the lhs. of (8.26) are completely analogous and so omitted. We thus get (cf. (8.34))

(8.43)
$$|\langle G_{t\wedge\tau}(\mathrm{i}\eta_{t\wedge\tau}) - M_{t\wedge\tau}\rangle| \leq \int_0^{t\wedge\tau} \widetilde{\mathcal{E}}_1(s) |\langle G_{s\wedge\tau}(\mathrm{i}\eta_s) - M_{s\wedge\tau}\rangle| \mathrm{d}s + \frac{N^\gamma \log N}{N\kappa_{t\wedge\tau}},$$

with $\widetilde{\mathcal{E}}_1(s)$ from (8.33). Finally, using Gronwall inequality and (8.36), together with

$$\frac{\eta_s}{\eta_t} \sim \frac{\eta_s/\rho_s}{\eta_t/\rho_t} \sim \frac{\kappa_s^{2/3}}{\kappa_t^{2/3}} \lesssim \frac{\kappa_s}{\kappa_t}$$

for $s \leq t$, we obtain the desired result.

8.2. Proof of Proposition 8.4. Fix $r \ge 1$ and let $\epsilon > 0$ be as in (8.13). We will prove Proposition 8.4 by first proving the high moment bounds

(8.44)
$$\mathbb{E}\left|\left\langle B\left(G^{0}(\mathrm{i}\eta)-M_{r\mathcal{A}}(\mathrm{i}\eta)\right)\right\rangle\right|^{p} \lesssim \mathbb{E}\left|\left\langle B\left(G^{t}(\mathrm{i}\eta)-M_{r\mathcal{A}}(\mathrm{i}\eta)\right)\right\rangle\right|^{p}+\left(\frac{1}{N\eta}\right)^{r}$$

and

(8.45)
$$\mathbb{E}\left|\boldsymbol{u}^{*}\left(\boldsymbol{G}^{0}(\mathrm{i}\boldsymbol{\eta})-\boldsymbol{M}_{r\mathcal{A}}(\mathrm{i}\boldsymbol{\eta})\right)\boldsymbol{y}\right|^{p} \lesssim \mathbb{E}\left|\boldsymbol{u}^{*}\left(\boldsymbol{G}^{t}(\mathrm{i}\boldsymbol{\eta})-\boldsymbol{M}_{r\mathcal{A}}(\mathrm{i}\boldsymbol{\eta})\right)\boldsymbol{y}\right|^{p}+\left(\sqrt{\frac{\rho^{r\mathcal{A}}(\mathrm{i}\boldsymbol{\eta})}{N\boldsymbol{\eta}}}\right)^{p}$$

on the scale $\tilde{\mathfrak{s}}_1$, i.e. for $\eta = \tilde{\eta}_1$, for $p \in \mathbb{N}$. The proof of these two bounds will be presented shortly.

Proof of Proposition 8.4. By assumption, (8.14) holds for $G^t - M_{rA}$. Using this fact and (8.44)–(8.45) we immediately obtain the desired bound (8.14) for t = 0 as well.

Recall that $\tilde{\eta}_j$ satisfies $\tilde{\mathfrak{s}}_j = \mathfrak{s}(\tilde{\eta}_j, r)$ for j = 0, 1. For the remainder of the section we drop the tilde and write $\eta_j \equiv \tilde{\eta}_j$. Define

(8.46)
$$\Psi(z,\eta) := \sqrt{\frac{\rho^{r\mathcal{A}}(\mathrm{i}\eta)}{N\eta}}.$$

We begin with the following lemma which uses monotonicity to control the entries of $G^{s}(i\eta)$ at η_{1} given a local law at η_{0} . We point out that t will be fixed throughout this section.

Lemma 8.7. Fix $s \in [0, t]$. Then, under the assumptions of Proposition 8.4, we have

(8.47)
$$|\boldsymbol{u}^* \left(G^s(\mathrm{i}\eta_1) - M_{r\mathcal{A}}(\mathrm{i}\eta_1) \right) \boldsymbol{y}| \lesssim \frac{\eta_0}{\eta_1} \rho^{r\mathcal{A}}(\mathrm{i}\eta_0),$$

with very high probability uniformly in $\boldsymbol{u}, \boldsymbol{y} \in \mathbb{C}^{2N}$. Here, Ψ is defined in (8.46).

Proof. By the Cauchy-Schwarz inequality and the Ward identity, we get

(8.48)
$$|\boldsymbol{u}^*(G^s(\mathrm{i}\eta_1) - G^s(\mathrm{i}\eta_0))\boldsymbol{y}| \leq \int_{\eta_1}^{\eta_0} |\boldsymbol{u}^*G^s(\mathrm{i}\widehat{\eta})^2\boldsymbol{y}| \mathrm{d}\widehat{\eta} \leq \frac{\eta_0}{2\eta_1} (\boldsymbol{u}^*\operatorname{Im} G^s(\mathrm{i}\eta_0)\boldsymbol{u} + \boldsymbol{y}^*\operatorname{Im} G^s(\mathrm{i}\eta_0)\boldsymbol{y}).$$

where in the last inequality we used that $\hat{\eta} \mapsto \hat{\eta} \operatorname{Im} G(i\hat{\eta})$ is monotone increasing with respect to the Loewner order. As, by assumption, (8.14) holds on a scale $\tilde{\mathfrak{s}}_0$ we obtain

(8.49)
$$|\boldsymbol{u}^*(G^s(\mathrm{i}\eta_1) - G^s(\mathrm{i}\eta_0))\boldsymbol{y}| \le \frac{\eta_0}{\eta_1} \rho^{r\mathcal{A}}(\mathrm{i}\eta_0) \left(1 + O\left(N^{-\epsilon/5}\right)\right)$$

with very high probability. Similarly, by (8.14), we obtain

(8.50)
$$|\boldsymbol{u}^* \left(G^s(\mathrm{i}\eta_0) - M(\mathrm{i}\eta_0) \right) \boldsymbol{y}| \le \frac{\eta_0}{\eta_1} N^\gamma \left(\log N \right)^\alpha \Psi(z,\eta_0).$$

Combing (8.49) and (8.50) we complete the proof.

For simplicity we will use the short-hand notation $\tilde{\rho} \equiv \rho^{r\mathcal{A}}(i\eta_0)$. Lemma 8.7 will be our main tool for estimating the entries of G below. At times we will keep track of the $\tilde{\rho}$ term, however often we will apply (3.5) and simply use that $\tilde{\rho} \leq 1$. To complete the proofs of (8.44) and (8.45), we can apply Grönwall's inequality to the following propositions. For simplicity of presentation in the remainder of the section we use the short-hand notation $\delta = \frac{\epsilon_0}{100}$ to denote the step size in our scale. We note that Lemma 8.7 is presented as an inequality holding with high probability. This implies that the inequality also holds in the sense of stochastic domination, which will be more convenient to use in the proofs of Propositions 8.8 and 8.9 below.

Proposition 8.8. Define

)
$$R_s := \langle B \left(G^s(i\eta) - M_{r\mathcal{A}}(i\eta) \right) \rangle$$

Let p > 2 and $\delta := \epsilon_0/100$, then

(8.52)
$$\left| \mathbb{E} \frac{\mathrm{d}}{\mathrm{d}s} |R_s|^p \right| \lesssim N^{\frac{1}{2} + 10\delta} \Psi(z, \eta_0) \left[|R_s|^p + \left(\frac{1}{N\eta_1}\right)^p \right],$$

for all $0 \leq s \leq t$.

(8.51)

Proof of (8.44). By (8.52) with $\eta_1 = \eta$, using Gronwall inequality we obtain

(8.53)
$$\mathbb{E}|R_{0}|^{p} \lesssim \exp\left(tN^{\frac{1}{2}+10\delta}\Psi(z,\eta_{0})\right) \left[\mathbb{E}|R_{t}|^{p} + \left(\frac{1}{N\eta}\right)^{p}\right]$$
$$\leq \exp\left(N^{10\delta}\left[\sqrt{\frac{\eta_{0}}{\rho^{r\mathcal{A}}(i\eta_{0})}} + \frac{\eta_{0}}{\sqrt{N}\rho^{r\mathcal{A}}(i\eta_{0})}\right]\right) \left[\mathbb{E}|R_{t}|^{p} + \left(\frac{1}{N\eta}\right)^{p}\right]$$
$$\leq \exp\left(N^{10\delta-\epsilon_{0}/2}\right) \left[\mathbb{E}|R_{t}|^{p} + \left(\frac{1}{N\eta}\right)^{p}\right],$$

where in the first inequality we used that $t \leq \tilde{\mathfrak{s}}_0 = \eta_0 / \rho^{r\mathcal{A}}(i\eta_0)$, and in the second inequality that $\eta_0 / \rho^{r\mathcal{A}}(i\eta_0) \leq N^{-\epsilon_0}$, by assumption. Finally, using $10\delta - \epsilon_0/2 = -2\epsilon_0/5$ we obtain $\exp\left(N^{10\delta - \epsilon_0/2}\right) \leq 1$, which concludes the proof.

Proof. Let H_s be the Hermitization of $r\mathcal{A} + X_s$. Differentiating $R_s = R_s(H_s)$, by Itô's lemma we have

(8.54)
$$\mathbb{E}\frac{\mathrm{d}}{\mathrm{d}s}|R_s|^p = \mathbb{E}\left[-\frac{1}{2}\sum_{a,b}w_{ab}(s)\partial_{ab}|R_s|^p + \frac{1}{2}\sum_{a,b,c,d}\kappa_s(ab,cd)\partial_{ab}\partial_{cd}|R_s|^p\right].$$

where $a, b, c, d \in [2N]$, $w_{ab}(s)$ is the *ab*-th entry of H_s , $\kappa_s(ab, a_1b_1, \ldots)$ are the joint cumulants of $w_{ab}(s), w_{a_1b_1}(s), \ldots$, and $\partial_{ab} = \partial_{w_{ab}(s)}$ denotes the directional derivative in the direction $w_{ab}(s)$. For a smooth function f of the entries of H_s one can expand expectations in the cumulants:

(8.55)
$$\mathbb{E}w_{ab}f(H) = \sum_{k=0}^{L} \sum_{a_1,b_1,\dots,1_k b_k} \frac{\kappa_s(ab,a_1b_1,\dots,a_kb_k)}{k!} \mathbb{E}\partial_{a_1b_1}\cdots \partial_{a_kb_k}f(H) + \Omega_f(L),$$

for some error $\Omega_f(L)$ going to zero as $L \to \infty$. Using this cumulant expansion for the first sum in (8.54) we see that

(8.56)
$$\left| \mathbb{E} \frac{\mathrm{d}}{\mathrm{d}s} |R_s|^p \right| \le \left| \sum_{k=2}^{L-1} \sum_{a,a_1,\dots,a_k,b,b_1,\dots,b_k} \frac{\kappa_s(ab,a_1b_1,\dots,a_kb_k)}{k!} \mathbb{E} \partial_{a_kb_k} \cdots \partial_{ab} |R_s|^p \right| + \Omega(L),$$

where $\Omega(L) \leq \left(\frac{1}{N\eta_1}\right)^p$ for some sufficiently large L independent of N.

We now first present the estimates for the third order cumulants, which are the most delicate, and then present the ones for higher order cumulants. For the third order cumulants we aim to control the term

$$\left| N^{-3/2} \sum_{a,b} \mathbb{E} \partial_{ab}^3 |R_s|^p \right|,$$

which after applying the chain rule can be broken up into terms involving derivatives of the form $(\partial_{ab}R_s)^3|R_s|^{p-3}$, $(\partial_{ab}R_s)(\partial_{ab}^2R_s)|R_s|^{p-2}$, and $(\partial_{ab}^3R_s)|R_s|^{p-1}$. For simplicity of notation, here we neglected the difference between R_s and $\overline{R_s}$ since terms involving one or the other will be estimated in a completely analogous way.

We begin with the term involving $(\partial_{ab}^3 R_s)|R_s|^{p-1}$. First we note

(8.57)
$$\partial_{ab}^3 R_s = \frac{1}{2N} \sum_x \partial_{ab}^3 (BG)_{xx},$$

and $\partial_{ab}^3 G_{xx}$ is a sum of terms of the form $(BG)_{xa}G_{bb}G_{aa}G_{bx}$, $(BG)_{xa}G_{bb}G_{ax}$, $(BG)_{xa}G_{ab}^2G_{bx}$, or similar terms with some transposition of a and b. We will see that there are essentially two cases, when the product of G entries involves an off diagonal term, and when it does not. We now illustrate how to stochastically dominate each of these representative examples. In all three examples, as well as in the case when the derivatives hit more R_s or for higher cumulants, the terms involving $(BG)_x$. will be dealt with by eventually bounding the norm of a column of BG by ||B|| times the identically indexed column of G. To avoid repetition we assume B = I for the remainder of the proof. Applying Cauchy–Schwarz to the sum of x, followed by the Ward identity and Lemma 8.7, we obtain (recalling $\eta_0/\eta_1 \leq N^{3\delta/2}$)

$$(8.58) N^{-5/2} \left| \sum_{a,b,x} G_{xa} G_{ab} G_{ab} G_{bx} \right| \lesssim N^{-5/2} \frac{1}{\eta_1} \sum_{a,b} |G_{ab}|^2 \sqrt{\operatorname{Im} G_{aa}} \sqrt{\operatorname{Im} G_{bb}} \prec N^{-3/2} \frac{1}{\eta_1^2} \left(\frac{\eta_0}{\eta_1} \widetilde{\rho} \right)^2 \lesssim \frac{N^{10\delta} \widetilde{\rho}^2}{\eta_1 \sqrt{N}} \frac{1}{N\eta_1}.$$

Similarly, we bound

(8.59)
$$N^{-5/2} \left| \sum_{a,b,x} G_{xa} G_{ba} G_{bb} G_{ax} \right| \prec \frac{N^{10\delta} \widetilde{\rho}^{5/2}}{\sqrt{\eta_1}} \frac{1}{N\eta_1}$$

For the remaining term, we need to handle the sum over the trace index x more carefully, since we do not have other off-diagonal terms to apply the Ward identity to. Applying Lemma 8.7 and the Ward identity

$$(8.60)$$

$$N^{-5/2} \left| \sum_{a,b,x} G_{xa} G_{aa} G_{bb} G_{bx} \right| \lesssim N^{-5/2} \sum_{a,b} |G_{aa}| |G_{bb}| |(G^2)_{ba}|$$

$$(8.60)$$

$$\prec N^{-5/2} \left(\frac{\eta_0}{\eta_1} \widetilde{\rho} \right)^2 \sum_b \left(N \sum_a |(G^2)_{ab}|^2 \right)^{1/2}$$

$$\lesssim \frac{1}{\sqrt{\eta_1}} \left(\frac{\eta_0}{\eta_1} \widetilde{\rho} \right)^{5/2} \frac{1}{N\eta_1}.$$

Applying Young's inequality, from (8.58), (8.60), and (8.59) we conclude that (8.61)

$$\left| N^{-3/2} \sum_{a,b} (\partial_{ab}^3 R_s) |R_s|^{p-1} \right| \prec \frac{N^{5\delta} \tilde{\rho}^2}{\sqrt{\eta_1}} \left[|R_s|^p + \left(\frac{1}{N\eta_1}\right)^p \right] \le N^{1/2 + 6\delta} \Psi(z,\eta_0) \left[|R_s|^p + \left(\frac{1}{N\eta_1}\right)^p \right]$$

Moving on to terms involving $(\partial_{ab}R_s)(\partial_{ab}^2R_s)|R_s|^{p-2}$, we again use Ward's identity, Cauchy–Schwarz inequality, and Lemma 8.7 to estimate (8.62)

$$N^{-3/2} \left| \sum_{a,b} (\partial_{ab} R_s) (\partial_{ab}^2 R_s) |R_s|^{p-2} \right| \lesssim N^{-3/2} \left| \sum_{a,b} \left[\frac{1}{2N} \sum_x G_{xa} G_{bx} \right] \left[\frac{1}{2N} \sum_x G_{xb} G_{aa} G_{bx} + G_{xa} G_{ba} G_{bx} \right] |R_s|^{p-2} \right|$$
$$\times \left(\frac{\eta_0}{\eta_1} \widetilde{\rho} \right)^{3/2} \left(\frac{1}{N\eta_1} \right)^2 N^{-3/2} \sum_{a,b} |(G^2)_{ab}| |R_s|^{p-2}$$
$$\lesssim \frac{1}{\sqrt{\eta_1}} \left(\frac{\eta_0}{\eta_1} \widetilde{\rho} \right)^2 \left(\frac{1}{N\eta_1} \right)^2 |R_s|^{p-2}.$$

Finally, for the terms involving $(\partial_{ab}R_s)^3 |R_s|^{p-3}$ we apply the same approaches to see

(8.63)
$$\left| N^{-3/2} \sum_{a,b} \left[\partial_{ab} R_s \right]^3 |R_s|^{p-3} \right| \prec \left(\frac{\eta_0}{\eta_1} \tilde{\rho} \right)^2 \left(\frac{1}{N\eta_1} \right)^3 \frac{1}{\eta_1 \sqrt{N}} |R_s|^{p-3}$$

By a simple Young's inequality, together with $\eta_0/\eta_1 \leq N^{3\delta/2}$, we then obtain a bound as in the rhs. of (8.61).

For the higher order terms we aim to estimate

(8.64)
$$\frac{\kappa_n}{n!} \sum_{a,b} \mathbb{E}\partial^n_{ab} |R_s|^p, \qquad n \ge 4.$$

Here we chose the assignment $\kappa_n := \kappa_n(ab, ab, ...)$ for simplicity of notation, all other cases being analogous. We expand the derivative as

(8.65)
$$\frac{\kappa_n}{n!} \sum_{a,b} \partial_{ab}^n |R_s|^p = \frac{\kappa_n}{n!} \sum_{a,b} \sum_{k=1}^n \sum_{\substack{n=(n_1,\dots,n_k) \in \mathbb{N}^k \\ n_1 + \dots + n_k = n}} c_{n,p} \prod_{i=1}^k \partial_{ab}^{n_i} [R_s] |R_s|^{p-k},$$

for some combinatorial numbers $c_{n,p}$ depending on $n \in \mathbb{N}^k$ and $p \in \mathbb{N}$. For any $n_i \geq 1$ the partial derivative $\partial_{ab}R_s$ is the sum over a trace index x over products over $n_i + 1$ entries of G of the form $G_x \cdots G_x$ where the middle terms are entries G_{aa}, G_{ba}, G_{bb} , or G_{bb} . We can apply Lemma 8.7 to the middle terms, and Cauchy–Schwarz followed by two applications of the Ward identity and Lemma 8.7 again to estimate

(8.66)
$$|\partial_{ab}^{n_i} R_s| \prec \frac{1}{N\eta_1} \left(\frac{\eta_0 \widetilde{\rho}}{\eta_1}\right)^{n_i}$$

Hence,

(8.67)
$$N^{-n/2} \sum_{a,b} \prod_{i=1}^{k} |\partial_{ab}^{n_i}[R_s]| \prec N^{2-n/2} \left(\frac{\eta_0 \widetilde{\rho}}{\eta_1}\right)^k \left(\frac{1}{N\eta_1}\right)^k$$

Using $\kappa_n \leq N^{-n/2}$, after applying Young's inequality to (8.65) and (8.67) we see that for $n \geq 4$ we have (8.68)

$$\frac{\kappa_n}{n!} \sum_{a,b} \mathbb{E}\partial_{ab}^n |R_s|^p \prec N^{2-n/2} \left(\frac{\eta_0 \widetilde{\rho}}{\eta_1}\right)^n \left[|R_s|^p + \left(\frac{1}{N\eta_1}\right)^p \right] \le N^{1/2+10\delta} \Psi(z,\eta_0) \left[|R_s|^p + \left(\frac{1}{N\eta_1}\right)^p \right].$$

Proposition 8.9. Let $u, y \in \mathbb{C}^{2N}$ and define

$$(8.69) \qquad S_s(\boldsymbol{u}, \boldsymbol{y}) \coloneqq \boldsymbol{u}^* \left(G^s(\mathrm{i}\eta) - M_{A-z}(\mathrm{i}\eta) \right) \boldsymbol{y}.$$

Let $p > 2$ be even and let $\delta \coloneqq \epsilon_0 / 100$, then
$$(8.70) \qquad \left| \mathbb{E} \frac{\mathrm{d}}{\mathrm{d}s} |S_s(\boldsymbol{u}, \boldsymbol{y})|^p \right| \lesssim N^{\frac{1}{2} + 10\delta} \Psi(z, \eta_0) \left[|S_s(\boldsymbol{u}, \boldsymbol{y})|^p + (\Psi(z, \eta_0))^p \right],$$

for all $0 \leq s \leq t$.

Proof of (8.45). Given (8.70), the proof of (8.45) is completely analogous to the one of (8.44), and so omitted.

Proof of Proposition 8.9. Similar to Proposition 8.9 after applying Itô's lemma and a cumulant expansion we see that

$$(8.71) \left\| \mathbb{E} \frac{\mathrm{d}}{\mathrm{d}s} |S_s(\boldsymbol{u}, \boldsymbol{y})|^p \right\| \leq \left| \sum_{k=2}^{L-1} \sum_{a,a_1,\dots,a_k,b,b_1,\dots,b_k} \frac{\kappa_s(ab,a_1b_1,\dots,a_kb_k)}{k!} \mathbb{E} \partial_{a_kb_k} \cdots \partial_{ab} |S_s(\boldsymbol{u}, \boldsymbol{y})|^p \right| + \Omega(L).$$

For simplicity we will write $G_{\boldsymbol{u}\boldsymbol{y}} = \boldsymbol{u}^*(G^s(i\eta))\boldsymbol{y}$ and $S_s = S_s(\boldsymbol{u}, \boldsymbol{y})$. Here $\Omega(L)$ is a negligible error if L is sufficiently large. We start with the estimate of the third order cumulants, i.e. we present the estimate of $\frac{\kappa_3}{6} \sum_{a,b} \mathbb{E}\partial_{ab}^3 |S_s|^p$, with $\kappa_3 := \kappa_3(ab, ab, ab)$. The estimate of all the other terms coming from third order cumulants is completely analogous and so omitted. Expanding the derivative, we obtain

(8.72)
$$\frac{\kappa_3}{6} \sum_{a,b} \mathbb{E}\partial_{ab}^3 |S_s|^p = \frac{\kappa_3}{6} \sum_{a,b} \mathbb{E}p(\partial_{ab}^3 G_{\boldsymbol{u}\boldsymbol{y}}) |S_s|^{p-1} + \mathbb{E}c_p(\partial_{ab}^2 G_{\boldsymbol{u}\boldsymbol{y}})(\partial_{ab} G_{\boldsymbol{u}\boldsymbol{y}}) |S_s|^{p-2} + \mathbb{E}c_p'(\partial_{ab} G_{\boldsymbol{u}\boldsymbol{y}})^3 |S_s|^{p-3}.$$

Note that $\kappa_3 \leq N^{-3/2}$, and we begin with the term $\mathbb{E}p(\partial_{ab}^3 G_{uy})|S_s(u, y)|^{p-1}$. $\partial_{ab}^3 G_{uy}$ is a linear combination of terms of the form $G_{ua}G_{bb}G_{aa}G_{by}$, $G_{ua}G_{ab}G_{ba}G_{by}$, $G_{ua}G_{ab}^2G_{by}$, or similar terms up to a transposition of a and b. The main observation is that we always have two off-diagonal terms at the beginning and end of the product. All middle terms can be bounded using Lemma 8.7. For the term with two diagonal elements

(8.73)
$$N^{-3/2} \sum_{a,b} G_{ua} G_{bb} G_{aa} G_{by} \cdot |S_s|^{p-1},$$

we estimate the diagonal factors with Lemma 8.7:

(8.74)
$$N^{-3/2} \sum_{a,b} |G_{ua}G_{by}| |G_{aa}G_{bb}| \prec N^{-3/2+2\delta} \left(N \sum_{a} |G_{ua}|^2 \right)^{1/2} \left(N \sum_{b} |G_{by}|^2 \right)^{1/2} \prec N^{-1/2+2\delta} \frac{\eta_0^2}{\eta_1^2} \frac{\widetilde{\rho}}{\eta_0} \leq N^{1/2+8\delta} \Psi(z,\eta_0)^2,$$

and hence we arrive at

(8.75)
$$N^{-3/2} \sum_{a,b} G_{ua} G_{bb} G_{aa} G_{by} \cdot |S_s|^{p-1} \prec N^{1/2+8\delta} \Psi(z,\eta_0)^2 |S_s|^{p-1}.$$

As we can always apply Lemma 8.7 twice to the middle two terms in the product of four $G_{..}$ we conclude

(8.76)
$$N^{-3/2} \sum_{a,b} (\partial_{ab}^3 G_{\boldsymbol{u}\boldsymbol{y}}) |S_s|^{p-1} \prec N^{1/2+8\delta} \Psi(z,\eta_0)^2 |S_s|^{p-1}.$$

Similarly terms of the form $(\partial_{ab}G_{uy})^3$ can also be bounded.

$$N^{-3/2} \sum_{a,b} |G_{\boldsymbol{u}a} G_{b\boldsymbol{y}}|^3 |S_s|^{p-3} \prec N^{-\frac{3}{2}+4\delta} \sum_{a,b} |G_{\boldsymbol{u}a}|^2 |G_{b\boldsymbol{y}}|^2 |S_s|^{p-3} \prec N^{\frac{1}{2}+8\delta} \Psi(z_0,\eta_0)^4 |S_s|^{p-3},$$

and

(8.77)
$$N^{-3/2} \sum_{a,b} \left(\partial_{ab} G_{\boldsymbol{u}\boldsymbol{y}} \right)^3 |S_s|^{p-3} \prec N^{\frac{1}{2} + 8\delta} \Psi(z,\eta_0)^4 |S_s|^{p-3}$$

Again, nearly identical computations yields

(8.78)
$$N^{-3/2} \sum_{a,b} (\partial_{ab} G_{\boldsymbol{u}\boldsymbol{y}}) (\partial_{ab}^2 G_{\boldsymbol{u}\boldsymbol{y}}) |S_s|^{p-2} \prec N^{\frac{1}{2}+10\delta} \Psi(z,\eta_0)^3 |S_s|^{p-2}.$$

Taking expected values of (8.76), (8.77), and (8.78) we conclude

(8.79)
$$\left|\frac{\kappa_3}{6}\sum_{a,b}\mathbb{E}\partial_{ab}^3|S_s|^p\right| \lesssim N^{1/2+10\delta}\Psi(z,\eta_0)\big[\mathbb{E}|S_s|^p + (\Psi(z,\eta_0))^p\big].$$

This concludes the estimate of the third order cumulants.

We now present the bound for cumulants of order four and higher, i.e. our goal is to estimate $\frac{\kappa_n}{n!} \sum_{a,b} \mathbb{E}\partial_{ab}^n |S_s|^p$, for $n \ge 4$, where we again chose the assignment $\kappa_n := \kappa_n(ab, ab, ...)$ for simplicity of notation. We expand the derivative as

(8.80)
$$\frac{\kappa_n}{n!} \sum_{a,b} \partial_{ab}^n (G_{\boldsymbol{u}\boldsymbol{y}})^p = \frac{\kappa_n}{n!} \sum_{a,b} \sum_{k=1}^n \sum_{\substack{\boldsymbol{n}=(n_1,\dots,n_k)\in\mathbb{N}^k\\n_1+\dots+n_k=n}} c_{\boldsymbol{n},p} \prod_{i=1}^k \partial_{ab}^{n_i} [G_{\boldsymbol{u}\boldsymbol{y}}] (S_s)^{p-k},$$

for some combinatorial numbers $c_{n,p}$ depending on $n \in \mathbb{N}^k$ and $p \in \mathbb{N}$. Next, we claim for each $k \in [\![1, n]\!]$ and $n \in \mathbb{N}^k$, with $n_1 + \cdots + n_k = n$, that

(8.81)
$$N^{-n/2} \sum_{a,b} \prod_{i=1}^{k} |\partial_{ab}^{n_i}[G_{xy}]| \prec \left[N^{1/2 + 10\delta} \Psi(z, \eta_0) \right] \cdot (\Psi(z, \eta_0))^k$$

To this end, without loss of generality we assume $n_1 \leq n_2 \leq \cdots \leq n_k$ and divide the proof in two cases, $n_2 = 1$ and $n_2 > 1$.

Firstly when $n_2 = 1$, notice that

$$(8.82) \qquad |\partial_{ab}^{n_1}[G_{uy}]\partial_{ab}^{n_2}[G_{uy}]| = |G_{ua}G_{by} + G_{ub}G_{ay}|^2 \le 2(|G_{ua}|^2|G_{by}|^2 + |G_{ub}|^2|G_{ay}|^2).$$

For the remaining derivatives of order n_3, \ldots, n_k we apply Lemma 8.7 so that

(8.83)
$$|\partial_{ab}^{n}[G_{\boldsymbol{u}\boldsymbol{y}}]| \prec N^{(n+1)\delta}, \qquad n \in \mathbb{N}$$

In conclusion we have

(8.84)
$$N^{-n/2} \sum_{a,b} \prod_{i=1}^{\kappa} |\partial_{ab}^{n_i}[G_{uy}]| \prec N^{-n/2} N^{(n+k-4)\delta} \frac{\operatorname{Im} G_{uu} \operatorname{Im} G_{yy}}{\eta_1^2} \prec N^{-n/2} N^{(n+k-4)\delta} \frac{\eta_0^2}{\eta_1^4} \widetilde{\rho}^2 = N^{-n/2+2} N^{(n+k)\delta} \Psi(z,\eta_0)^4.$$

Finally, we use $N^{-1/2+\epsilon_0/2} \le N^{-1/2} \tilde{\mathfrak{s}}_0^{-1/2} = \Psi(z,\eta_0)$ so that (8.85)

$$N^{-n/2+2}N^{(n+k)\delta}\Psi(z,\eta_0)^4 \le \sqrt{N}N^{2n\delta-(n-4)\epsilon_0/2}\Psi(z,\eta_0) \cdot (\Psi(z,\eta_0))^n \le [N^{1/2+8\delta}\Psi(z,\eta_0)](\Psi(z,\eta_0))^k,$$

where we also used that $n \ge k$ and $\Psi(z, \eta_0) \le 1$. This concludes the proof of (8.81) for $n_2 = 1$.

We next prove (8.81) when $n_2 > 1$. Here we will only use the first factor corresponding to n_1 , and applying using Lemma 8.7 we have that

$$(8.86) \qquad |\partial_{ab}^{n_1}[G_{uy}]| \prec N^{(n_1-1)\delta}(|G_{ua}G_{by}| + |G_{ub}G_{ay}|) \le \frac{1}{2}N^{(n_1-1)\delta}\sum_{i\in\{u,y\},j\in\{a,b\}}|G_{ij}|^2$$

Applying (8.83) to the remaining derivatives, using again $N^{-1/2+\epsilon_0/2} \leq \Psi(z,\eta_0)$, we find

(8.87)
$$N^{-n/2} \sum_{a,b} \prod_{i=1}^{k} |\partial_{ab}^{n_i}[G_{u,y}]| \prec N^{-n/2+2} N^{(n+k)\delta - (n-4)\epsilon_0/2} \Psi(z,\eta_0)^2 \leq N^{1/2+8\delta} \Psi(z,\eta_0)^{n-1} \leq [N^{1/2+8\delta} \Psi(z,\eta_0)^2 \leq N^{1/2+8\delta} \Psi(z,\eta_0)^{n-1}$$

$$\leq [N^{1/2+8\delta}\Psi(z,\eta_0)](\Psi(z,\eta_0))^k$$

We point out that in the last inequality we used that $n_2 > 1$ implies $n \ge 2(k-1) + n_1 \ge 2k - 1$, which together with $n \ge 4$ gives $n \ge (2k+3)/2$ so that $n \ge k+2$. This concludes the proof of (8.81).

References

- A. Adhikari and J. Huang. Dyson Brownian motion for general β and potential at the edge. Probab. Theory Related Fields, 178(3-4):893–950, 2020.
- [2] A. Adhikari and B. Landon. Local law and rigidity for unitary Brownian motion. Probab. Theory Related Fields, 187(3-4):753-815, 2023.
- [3] A. Aggarwal, P. Lopatto, and H.-T. Yau. GOE statistics for Lévy matrices. J. Eur. Math. Soc. (JEMS), 23(11):3707– 3800, 2021.
- [4] O. H. Ajanki, L. Erdős, and T. Krüger. Universality for general Wigner-type matrices. Probab. Theory Related Fields, 169(3-4):667-727, 2017.
- [5] G. Akemann, S.-S. Byun, and N.-G. Kang. Scaling limits of planar symplectic ensembles. SIGMA Symmetry Integrability Geom. Methods Appl., 18:Paper No. 007, 40, 2022.
- [6] J. Alt, L. Erdős, and T. Krüger. The Dyson equation with linear self-energy: spectral bands, edges and cusps. Doc. Math., 25:1421–1539, 2020.
- [7] J. Alt, L. Erdős, T. Krüger, and D. Schröder. Correlated random matrices: band rigidity and edge universality. Ann. Probab., 48(2):963–1001, 2020.
- [8] J. Alt, L. Erdős, and T. Krüger. Local inhomogeneous circular law. Ann. Appl. Probab., 28(1):148–203, 2018.

- [9] J. Alt, L. Erdős, and T. Krüger. Spectral radius of random matrices with independent entries. Probab. Math. Phys., 2(2):221–280, 2021.
- [10] J. Alt and T. Krüger. Inhomogeneous circular law for correlated matrices. J. Funct. Anal., 281(7):Paper No. 109120, 73, 2021.
- [11] J. Alt and T. Krüger. Spectrum occupies pseudospectrum for random matrices with diagonal deformation and variance profile. preprint arXiv:2404.17573, 2024.
- [12] Y. Ameur, N.-G. Kang, and N. Makarov. Rescaling Ward identities in the random normal matrix model. Constr. Approx., 50(1):63–127, 2019.
- [13] J. Baik, G. Ben Arous, and S. Péché. Phase transition of the largest eigenvalue for nonnull complex sample covariance matrices. Ann. Probab., 33(5):1643–1697, 2005.
- [14] A. Basak and O. Zeitouni. Outliers of random perturbations of Toeplitz matrices with finite symbols. Probab. Theory Related Fields, 178(3-4):771–826, 2020.
- [15] S. Belinschi, C. Bordenave, M. Capitaine, and G. Cébron. Outlier eigenvalues for non-Hermitian polynomials in independent i.i.d. matrices and deterministic matrices. *Electron. J. Probab.*, 26:Paper No. 100, 37, 2021.
- [16] S. Belinschi, Z. Yin, and P. Zhong. The Brown measure of a sum of two free random variables, one of which is triangular elliptic. Adv. Math., 441:Paper No. 109562, 2024.
- [17] S. T. Belinschi, H. Bercovici, M. Capitaine, and M. Février. Outliers in the spectrum of large deformed unitarily invariant models. Ann. Probab., 45(6A):3571–3625, 2017.
- [18] S. T. Belinschi and M. Capitaine. Spectral properties of polynomials in independent Wigner and deterministic matrices. J. Funct. Anal., 273(12):3901–3963, 2017.
- [19] F. Benaych-Georges and J. Rochet. Outliers in the single ring theorem. Probab. Theory Related Fields, 165(1-2):313– 363, 2016.
- [20] J. Bigot and C. Male. Freeness over the diagonal and outliers detection in deformed random matrices with a variance profile. Inf. Inference, 10(3):863–919, 2021.
- [21] C. Bordenave and M. Capitaine. Outlier eigenvalues for deformed i.i.d. random matrices. Comm. Pure Appl. Math., 69(11):2131–2194, 2016.
- [22] C. Bordenave, D. Chafaï, and D. García-Zelada. Convergence of the spectral radius of a random matrix through its characteristic polynomial. *Probab. Theory Related Fields*, 182(3-4):1163–1181, 2022.
- [23] A. Borodin and C. D. Sinclair. The Ginibre ensemble of real random matrices and its scaling limits. Comm. Math. Phys., 291(1):177–224, 2009.
- [24] T. Bothner and A. Little. The complex elliptic Ginibre ensemble at weak non-Hermiticity: edge spacing distribution. preprint arXiv:2208.04684, 2022.
- [25] P. Bourgade. Extreme gaps between eigenvalues of Wigner matrices. J. Eur. Math. Soc. (JEMS), 24(8):2823–2873, 2022.
- [26] P. Bourgade, L. Erdős, and H.-T. Yau. Edge universality of beta ensembles. Comm. Math. Phys., 332(1):261–353, 2014.
- [27] P. Bourgade, L. Erdős, H.-T. Yau, and J. Yin. Universality for a class of random band matrices. Adv. Theor. Math. Phys., 21(3):739–800, 2017.
- [28] P. Bourgade, H.-T. Yau, and J. Yin. Random band matrices in the delocalized phase I: Quantum unique ergodicity and universality. Comm. Pure Appl. Math., 73(7):1526–1596, 2020.
- [29] L. G. Brown. Lidskii's theorem in the type II case. In Geometric methods in operator algebras (Kyoto, 1983), volume 123 of Pitman Res. Notes Math. Ser., pages 1–35. Longman Sci. Tech., Harlow, 1986.
- [30] S.-S. Byun and P. Forrester. Progress on the study of the Ginibre ensembles I:GinUE. preprint arXiv:2211.16223, 2022.
- [31] S.-S. Byun and P. Forrester. Progress on the study of the Ginibre ensembles II:GinOE and GinSE. preprint arXiv:2301.05022, 2023.
- [32] S.-S. Byun and K. Noda. Scaling limits of complex and symplectic non-Hermitian Wishart ensembles. preprint arXiv:2402.18257, 2024.
- [33] G. Cipolloni, L. Erdős, and J. Henheik. Eigenstate thermalisation at the edge for Wigner matrices. preprint arXiv:2309.05488, 2023.
- [34] G. Cipolloni, L. Erdős, and J. Henheik. Out-of-time-ordered correlators for Wigner matrices. preprint arXiv:2402.17609, 2024.
- [35] G. Cipolloni, L. Erdős, T. Krüger, and D. Schröder. Cusp universality for random matrices, II: The real symmetric case. Pure Appl. Anal., 1(4):615–707, 2019.
- [36] G. Cipolloni, L. Erdős, and D. Schröder. Optimal lower bound on the least singular value of the shifted Ginibre ensemble. Probab. Math. Phys., 1(1):101–146, 2020.
- [37] G. Cipolloni, L. Erdős, and D. Schröder. Edge universality for non-Hermitian random matrices. Probab. Theory Related Fields, 179(1-2):1–28, 2021.
- [38] G. Cipolloni, L. Erdős, and D. Schröder. On the condition number of the shifted real Ginibre ensemble. SIAM J. Matrix Anal. Appl., 43(3):1469–1487, 2022.
- [39] G. Cipolloni, L. Erdős, and D. Schröder. Mesoscopic central limit theorem for non-Hermitian random matrices. Probab. Theory Related Fields, 188(3-4):1131–1182, 2024.
- [40] G. Cipolloni, L. Erdős, D. Schröder, and Y. Xu. Directional extremal statistics for Ginibre eigenvalues. J. Math. Phys., 63(10):Paper No. 103303, 11, 2022.
- [41] G. Cipolloni, L. Erdős, D. Schröder, and Y. Xu. On the rightmost eigenvalue of non-Hermitian random matrices. Ann. Probab., 51(6):2192–2242, 2023.
- [42] G. Cipolloni, L. Erdős, and Y. Xu. Precise asymptotics for the spectral radius of a large random matrix. preprint arXiv:2210.15643, 2022.
- [43] G. Cipolloni, L. Erdős, and Y. Xu. Universality of extremal eigenvalues of large random matrices. preprint arXiv:2312.08325, 2023.

- [44] G. Cipolloni, L. Erdős, J. Henheik, and D. Schröder. Optimal lower bound on eigenvector overlaps for non-hermitian random matrices. *preprint arXiv:2301.03549*, 2023.
- [45] N. Coston, S. O'Rourke, and P. M. Wood. Outliers in the spectrum for products of independent random matrices. Ann. Inst. Henri Poincaré Probab. Stat., 56(2):1284–1320, 2020.
- [46] L. Erdős, T. Krüger, and D. Renfrew. Randomly coupled differential equations with elliptic correlations. Ann. Appl. Probab., 33(4):3098–3144, 2023.
- [47] L. Erdős, T. Krüger, and D. Schröder. Random matrices with slow correlation decay. Forum Math. Sigma, 7:Paper No. e8, 89, 2019.
- [48] L. Erdős, T. Krüger, and D. Schröder. Cusp universality for random matrices I: local law and the complex Hermitian case. Comm. Math. Phys., 378(2):1203–1278, 2020.
- [49] L. Erdős, S. Péché, J. A. Ramírez, B. Schlein, and H.-T. Yau. Bulk universality for Wigner matrices. Comm. Pure Appl. Math., 63(7):895–925, 2010.
- [50] L. Erdős and V. Riabov. Eigenstate thermalization hypothesis for Wigner-type matrices. preprint arXiv:2403.10359, 2024.
- [51] L. Erdős, B. Schlein, and H.-T. Yau. Universality of random matrices and local relaxation flow. Invent. Math., 185(1):75–119, 2011.
- [52] L. Erdős and K. Schnelli. Universality for random matrix flows with time-dependent density. Ann. Inst. Henri Poincaré Probab. Stat., 53(4):1606–1656, 2017.
- [53] L. Erdős and H. C. Ji. Density of Brown measure of free circular Brownian motion. preprint arXiv:2307.08626, 2023.
- [54] P. J. Forrester and T. Nagao. Eigenvalue statistics of the real ginibre ensemble. *Phys. Rev. Lett.*, 99:050603, Aug 2007.
- [55] J. Ginibre. Statistical ensembles of complex, quaternion, and real matrices. J. Mathematical Phys., 6:440–449, 1965.
- [56] V. L. Girko. The circular law. Teor. Veroyatnost. i Primenen., 29(4):669–679, 1984.
- [57] W. Hachem, A. Hardy, and J. Najim. Large complex correlated Wishart matrices: the Pearcey kernel and expansion at the hard edge. *Electron. J. Probab.*, 21:Paper No. 1, 36, 2016.
- [58] H. Hedenmalm and A. Wennman. Planar orthogonal polynomials and boundary universality in the random normal matrix model. Acta Math., 227(2):309–406, 2021.
- [59] J. W. Helton, T. Mai, and R. Speicher. Applications of realizations (aka linearizations) to free probability. J. Funct. Anal., 274(1):1–79, 2018.
- [60] J. Huang and B. Landon. Rigidity and a mesoscopic central limit theorem for Dyson Brownian motion for general β and potentials. *Probab. Theory Related Fields*, 175(1-2):209–253, 2019.
- [61] J. Huang, B. Landon, and H.-T. Yau. Bulk universality of sparse random matrices. J. Math. Phys., 56(12):123301, 19, 2015.
- [62] H. C. Ji and J. Park. Tracy-Widom limit for free sum of random matrices. preprint arXiv:2110.05147, 2021.
- [63] K. Johansson. Universality of the local spacing distribution in certain ensembles of Hermitian Wigner matrices. Comm. Math. Phys., 215(3):683–705, 2001.
- [64] A. Knowles and J. Yin. Anisotropic local laws for random matrices. Probab. Theory Related Fields, 169(1-2):257–352, 2017.
- [65] B. Landon and H.-T. Yau. Convergence of local statistics of Dyson Brownian motion. Comm. Math. Phys., 355(3):949– 1000, 2017.
- [66] B. Landon and H.-T. Yau. Edge statistics of Dyson Brownian motion. preprint arXiv:1712.03881, 2017.
- [67] J. O. Lee and K. Schnelli. Edge universality for deformed Wigner matrices. Rev. Math. Phys., 27(8):1550018, 94, 2015.
- [68] J. O. Lee and K. Schnelli. Local law and Tracy-Widom limit for sparse random matrices. Probab. Theory Related Fields, 171(1-2):543-616, 2018.
- [69] J. O. Lee, K. Schnelli, B. Stetler, and H.-T. Yau. Bulk universality for deformed Wigner matrices. Ann. Probab., 44(3):2349–2425, 2016.
- [70] S.-Y. Lee and R. Riser. Fine asymptotic behavior for eigenvalues of random normal matrices: ellipse case. J. Math. Phys., 57(2):023302, 29, 2016.
- [71] D.-Z. Liu and L. Zhang. Phase transition of eigenvalues in deformed Ginibre ensembles. preprint arXiv:2204.13171, 2022.
- [72] D.-Z. Liu and L. Zhang. Critical edge statistics for deformed GinUEs. preprint arXiv:2311.13227, 2023.
- [73] D.-Z. Liu and L. Zhang. Repeated erfc statistics for deformed GinUEs. preprint arXiv:2402.14362, 2024.
- [74] A. Maltsev and M. Osman. Bulk Universality for Complex non-Hermitian Matrices with Independent and Identically Distributed Entries. preprint arXiv:2310.11429, 2023.
- [75] J. A. Mingo and R. Speicher. Free probability and random matrices, volume 35 of Fields Institute Monographs. Springer, New York; Fields Institute for Research in Mathematical Sciences, Toronto, ON, 2017.
- [76] S. Nonnenmacher and M. Vogel. Local eigenvalue statistics of one-dimensional random nonselfadjoint pseudodifferential operators. J. Eur. Math. Soc. (JEMS), 23(5):1521–1612, 2021.
- [77] M. Osman. Bulk Universality for Real Matrices with Independent and Identically Distributed Entries. preprint arXiv:2402.04071, 2024.
- [78] L. Pastur and M. Shcherbina. On the edge universality of the local eigenvalue statistics of matrix models. Mat. Fiz. Anal. Geom., 10(3):335–365, 2003.
- [79] L. Pastur and M. Shcherbina. Bulk universality and related properties of Hermitian matrix models. J. Stat. Phys., 130(2):205–250, 2008.
- [80] B. Rider. A limit theorem at the edge of a non-Hermitian random matrix ensemble. J. Phys. A, 36(12):3401–3409, 2003.
- [81] B. Rider and C. D. Sinclair. Extremal laws for the real Ginibre ensemble. Ann. Appl. Probab., 24(4):1621–1651, 2014.
- [82] M. Shcherbina and T. Shcherbina. The least singular value of the general deformed Ginibre ensemble. J. Stat. Phys., 189(2):Paper No. 30, 27, 2022.
- [83] P. Śniady. Random regularization of Brown spectral measure. J. Funct. Anal., 193(2):291–313, 2002.

- [84] S. Sodin. The spectral edge of some random band matrices. Ann. of Math. (2), 172(3):2223–2251, 2010.
- [85] A. Soshnikov. Universality at the edge of the spectrum in Wigner random matrices. Comm. Math. Phys., 207(3):697– 733, 1999.
- [86] T. Tao. Outliers in the spectrum of iid matrices with bounded rank perturbations. *Probab. Theory Related Fields*, 155(1-2):231–263, 2013.
- [87] T. Tao and V. Vu. Random matrices: universality of local eigenvalue statistics up to the edge. Comm. Math. Phys., 298(2):549–572, 2010.
- [88] T. Tao and V. Vu. Smooth analysis of the condition number and the least singular value. Math. Comp., 79(272):2333– 2352, 2010.
- [89] T. Tao and V. Vu. Random matrices: universality of local eigenvalue statistics. Acta Math., 206(1):127-204, 2011.
- [90] T. Tao and V. Vu. Random matrices: universality of local spectral statistics of non-Hermitian matrices. Ann. Probab., 43(2):782–874, 2015.
- [91] K. Tikhomirov. Quantitative invertibility of non-Hermitian random matrices. In ICM—International Congress of Mathematicians. Vol. IV. Sections 5–8, pages 3292–3313. EMS Press, Berlin, 2023.
- [92] P. Zhong. Brown measure of the sum of an elliptic operator and a free random variable in a finite von Neumann algebra. *preprint arXiv:2108.09844*, 2021.