SPECTRUM OF WEIGHTED COMPOSITION OPERATORS PART X THE SPECTRUM AND ESSENTIAL SPECTRA OF WEIGHTED AUTOMORPHISMS OF THE POLYDISC ALGEBRA.

ARKADY KITOVER AND MEHMET ORHON

ABSTRACT. We investigate the spectrum and the essential spectra of weighted automorphisms of the polydisc algebra \mathbb{A}^n . In the case n = 2 we provide a detailed and, in most cases, complete description of these spectra.

1. INTRODUCTION

The spectrum of weighted automorphisms of the disc-algebra was described by Kamowitz in [3]. In [4] the first named author obtained a full description of the spectrum of weighted automorphisms of uniform algebras. The paper [5] contains a description of essential spectra of weighted automorphisms of the algebra C(K). In [6] the authors obtained some general results about the spectrum and essential spectra of weighted isometrical endomorphisms of uniform algebras and applied these results to the problem of describing the essential spectra of weighted automorphisms of the disc-algebra. This description is complete in the case of automorphisms generated by elliptic or parabolic Möbius transformations, and partial in the case of hyperbolic transformations (see [6, Problem 3.7]).

The goal of the current paper is to apply the results obtained in [4, 5, 6] to the problem of describing essential spectra of weighted automorphisms of the polydisc algebra \mathbb{A}^n . While our methods can be applied in the case of any $n \in \mathbb{N}$, we decided to provide as many details as possible in the case n = 2. As the reader will see, even in this case the multitude of distinct possibilities is quite large.

Date: Tuesday 30th April, 2024.

²⁰¹⁰ Mathematics Subject Classification. Primary 47B33; Secondary 47B48, 46B60. Key words and phrases. Weighted composition operators, spectrum, Fredholm spectrum, essential spectra.

2. NOTATIONS AND PRELIMINARIES

The following notations are used throughout the paper.

 \mathfrak{M}_A and ∂A are the space of maximal ideals and the Shilov boundary of a commutative Banach algebra A, respectively.

Let A be a commutative unital Banach algebra and U be an automorphism of A. We will often instead of U write T_{φ} , where φ is the homeomorphism of \mathfrak{M}_A onto itself generated by U.

 \mathbb{N} is the set of all positive integers.

 \mathbb{Z} is the set of all integers.

C is the field of all complex numbers.

We denote by \mathbb{D} and by \mathbb{U} the closed and the open unit disk, respectively.

 $\mathbb T$ is the unit circle.

 $\mathbb{AN}(r, R)$, where $0 < r < R < \infty$, is the annulus $\{\lambda \in \mathbb{C} : r \leq |\lambda| \leq R\}$. For $n = 2, 3, ..., \mathbb{D}^n$, \mathbb{U}^n , and \mathbb{T}^n denote the closed unit polydisc, the open unit polydisc, and the *n*-dimensional unit torus, respectively.

All linear spaces are considered over the field \mathbb{C} .

C(K) is the space of all complex-valued continuous functions on a Hausdorff compact space K endowed with the standard norm.

 \mathbb{A}^n , $n = 2, 3, \ldots$ is the polydisc algebra: the algebra of all functions analytic in \mathbb{U}^n and continuous in \mathbb{D}^n .

It is well known (see e.g [7]) that $\mathfrak{M}_{\mathbb{A}^n} = \mathbb{D}^n$ and $\partial \mathbb{A}^n = \mathbb{T}^n$.

Let K be a compact Hausdorff space and φ be a homeomorphism of K onto itself. Then, $\varphi^0(k) = k, k \in K$, and $\varphi^n = \varphi^{n-1} \circ \varphi, n \in \mathbb{Z}$.

Let K be a Hausdorff compact space, φ be a continuous map of K into itself, and $w \in C(K)$. Then, $w_1 = w$ and $w_n = w_{n-1}(w \circ \varphi^{n-1}), n = 2, 3, \ldots$

Let X be a Banach space and $T : X \to X$ be a continuous linear operator.

 $\sigma(T)$ is the spectrum of T.

 $\rho(T)$ is the spectral radius of T.

$$\rho_{min}(T) = \begin{cases} 1/\rho(T^{-1}, & \text{if the operator } T \text{ is invertible;} \\ 0, & \text{otherwise.} \end{cases}$$

 $\sigma_{a.p.}(T)$ is the approximate point spectrum of T, i.e.

$$\sigma_{a.p.}(T) = \{ \lambda \in \mathbb{C} : \exists x_n \in X, \|x_n\| = 1, Tx_n - \lambda x_n \underset{n \to \infty}{\to} 0 \}.$$

According to this definition the point spectrum of T is a subset of $\sigma_{a.p.}(T)$.

 $\sigma_r(T) = \sigma(T) \setminus \sigma_{a.p.}(T)$ is the residual spectrum of T.

Recall that a continuous linear operator T on a Banach space X is called upper semi-Fredholm (respectively, lower semi-Fredholm) if its image R(T) is closed in X and dim ker $T < \infty$ (respectively, if $codim R(T) < \infty$) ∞).

The operator T is called semi-Fredholm if it is either upper semi-Fredholm or lower semi-Fredholm, and it is called Fredholm if it is both upper semi-Fredholm and lower semi-Fredholm.

The operator T is called Weil operator if dim ker $T = codim R(T) < \infty$. $\sigma_{sf}(T)$ is the semi-Fredholm spectrum of T, i.e. $\sigma_{sf}(T) = \{\lambda \in \mathbb{C} :$ $\lambda I - T$ is not semi-Fredholm}.

 $\sigma_{usf}(T)$ is the upper semi-Fredholm spectrum of T, i.e. $\sigma_{usf}(T) = \{\lambda \in \mathcal{S}_{usf}(T) : \{\lambda \in \mathcal{S}_{usf}(T) \}$ $\mathbb{C}: \lambda I - T$ is not an upper semi-Fredholm operator. It is well known (see e.g. [2]) that

$$\sigma_{usf}(T) = \{\lambda \in \mathbb{C} : \exists x_n \in X, \|x_n\| = 1, Tx_n - \lambda x_n \xrightarrow{} 0, \\ n \to \infty \}$$

and the sequence x_n is singular, i.e. it does not contain

a norm convergent subsequence }.

 $\sigma_{lsf}(T)$ is the lower semi-Fredholm spectrum of T, i.e. $\sigma_{lsf}(T) = \{\lambda \in \mathcal{L}\}$ $\mathbb{C}: \lambda I - T$ is not a lower semi-Fredholm operator}.

$$\sigma_{lsf}(T) = \{ \lambda \in \mathbb{C} : \exists x'_n \in X', \|x'_n\| = 1, T'x'_n - \lambda x'_n \underset{n \to \infty}{\to} 0,$$

and the sequence x'_n is singular.

 $\sigma_f(T)$ is the Fredholm spectrum of T, i.e. $\sigma_f(T) = \{\lambda \in \mathbb{C} : \lambda I - T \text{ is }$ not a Fredholm operator.

 $\sigma_w(T)$ is the Weil spectrum of T, i.e. $\sigma_f(T) = \{\lambda \in \mathbb{C} : \lambda I - T \text{ is not} \}$ a Weil operator}.

It is well known (see e.g [2]) that

- $\sigma_{sf}(T) \subseteq \sigma_{usf}(T) \subseteq \sigma_f(T) \subseteq \sigma_w(T),$ $\sigma_{usf}(T) = \sigma_{lsf}(T'), \ \sigma_{lsf}(T) = \sigma_{usf}(T').$

In the case when operator T is defined and bounded on distinct Banach spaces, $\sigma(T, X)$ means the spectrum of T on the Banach space X. The same agreement relates to all the spectra considered above.

3. AXILLARY RESULTS

Our investigation of essential spectra of weighted automorphisms of the algebra \mathbb{A}^2 is based on several lemmas stated below.

Lemma 3.1. (see [6, Proposition 2.8] Let A be a unital uniform algebra. Assume that ∂A has no isolated points. Let T_{φ} be a periodic automorphism of A, i.e. there is an $n \in \mathbb{N}$ such that $T_{\varphi}^n = I$. Let $w \in A$ and $T = wT_{\varphi}$. For any $t \in \mathfrak{M}_A$ let p(t) be the smallest positive integer such that $\varphi^{p(t)}(t) = t$. Then,

(1)
$$\sigma(T) = \{\lambda \in \mathbb{C} : \exists t \in \mathfrak{M}_A, \lambda^{p(t)} = w_{p(t)}(t)\}.$$

(2) $\sigma_{a.p.}(T) = \sigma_{sf}(t) = \{\lambda \in \mathbb{C} : \exists t \in \partial A, \lambda^{p(t)} = w_{p(t)}(t)\}$

Let us recall the following definition.

Definition 3.2. A uniform algebra A is called analytic if for any nonempty open subset U of ∂A and for any $a \in A$ we have $a|U \equiv 0 \Rightarrow a = 0$.

Lemma 3.3. (see [6, Theorem 2.9]) Let A be a unital analytic uniform algebra, $U = T_{\varphi}$ be a non-periodic automorphism of A, and $T = wT_{\varphi}$. Assume that

- (a) ∂A is a connected set.
- (b) There are periodic automorphisms, $U_n = T_{\varphi_n}$ of A such that for any $t \in \partial A$ we have $\varphi_n(t) \to \varphi(t)$.
- (c) Let $\mu \in (C(\partial A))'$ be a φ -invariant probability measure. Then, μ is φ_n -invariant, $n \in \mathbb{N}$.

Then,

- (1) $\sigma_{a.p.}(T) = \sigma_{sf}(T).$
- (2) The sets $\sigma(T)$ and $\sigma_{sf}(T)$ are connected rotation invariant subsets of \mathbb{C} .
- (3) If either $w \in A^{-1}$ or $\min_{s \in \partial A} |w(s)| = 0$, then $\sigma(T) = \sigma_{sf}(T)$.

Lemma 3.4. (see [4, Lemma 3.6, Theorem 4.2, Corollary 4.4] and [6, Theorem 2.1, Proposition 2.11])

Let A be an analytic uniform unital algebra. Let T_{φ} be a non-periodic automorphism of A, $w \in A$, and $T = wT_{\varphi}$. Then,

- (1) $\sigma(T)$ is a connected rotation invariant subset of \mathbb{C} .
- (2) $\sigma_{a.p.}(T)$ is rotation invariant and $\sigma_{a.p.}(T) = \sigma_{a.p.}(T, C(\partial A))$.
- (3) Let $\lambda \in \mathbb{C}$, then $\lambda \in \sigma_{a.p.}(T)$ if and only if there is a $t \in \partial A$ such that $|w_n(t)| \ge |\lambda|^n$ and $|w_n(\varphi^{-n}(t)) \le |\lambda|^n$ for any $n \in \mathbb{N}$.
- (4) If ∂A has no isolated points, then $\sigma_{usf}(T) = \sigma_{a.p.}(T)$.
- (5) If ∂A has no isolated points and there is a $u \in \partial A$ such that $|w_n(u)| \leq |\lambda|^n$ and $|w_n(\varphi^{-n}(u)) \geq |\lambda|^n$ for any $n \in \mathbb{N}$, then $\lambda \in \sigma_{lsf}(T)$.

Lemma 3.5. (see [5]) Let A be a unital uniform algebra and $T = wT_{\varphi}$ be a weighted automorphism of A. Assume that ∂A contains no isolated points. Assume also that ∂A is the union of φ and φ^{-1} invariant nonempty sets K_1, K_2 and O such that the sets K_1 and K_2 are closed,

4

 $(K_1 \cup K_2) \cap O = \emptyset$, and for any $k \in O$ we have $\bigcap_{m=1}^{\infty} cl\{\varphi^i(k) : i \ge m\} = K_1,$ $\bigcap_{m=1}^{\infty} cl\{\varphi^{-i}(k) : i \ge m\} = K_2.$

Then, the following statements hold.

- (1) If $0 \leq r = \rho(T, C(K_2)) < R = \rho_{min}(T, C(K_1))$, then either $\mathbb{AN}(r, R) \subseteq \sigma_{usf}(T)$, or, if r = 0, $R\mathbb{D} \subseteq \sigma_{usf}(T)$.
- (2) If $0 \leq r = \rho(T, C(K_1)) < R = \rho_{min}(T, C(K_2))$, then either $\mathbb{AN}(r, R) \subseteq \sigma_{lsf}(T)$, or, if r = 0, $\mathbb{RD} \subseteq \sigma_{lsf}(T)$.
- (3) If $\rho_{min}(T, C(K_1)) = \rho(T, C(K_1)) = \rho_{min}(T, C(K_2)) = \rho(T, C(K_2)) = r$, then $r\mathbb{T} \subseteq \sigma_{sf}(T)$.
- (4) Moreover, if there is a $k \in O$ such that w(k) = 0 and the set $\{n \in \mathbb{Z} : w(\varphi^n(k)) = 0\}$ is finite, then $\rho(T, C(K_2))\mathbb{D} \subseteq \sigma_{usf}(T)$ and $\rho(T, C(K_1))\mathbb{D} \subseteq \sigma_{lsf}(T)$.

Lemma 3.6. (see [4]) Let A be a unital uniform algebra and $T = wT_{\varphi}$ be a weighted automorphism of A. Then,

$$\rho(T) = \max_{\mu \in M_{\varphi}} \exp \int \ln |w| d\mu,$$

where $M_{\varphi} = \{\mu \in C'(\partial A) : T'_{\varphi}\mu = \mu, \mu \ge 0, \|\mu\| = 1\}.$

4. Essential spectra of weighted automorphisms of the algebra \mathbb{A}^2

It is well known (see [7]) that every automorphism of the algebra \mathbb{A}^2 is the composition operator T_{Φ} , where the map Φ is on of the following (a) $\Phi(z_1, z_2) = (\varphi(z_1), \psi(z_2)), (z_1, z_2) \in \mathbb{D}^2$, (b) $\Phi(z_1, z_2) = (\psi(z_2), \varphi(z_1)), (z_1, z_2) \in \mathbb{D}^2$,

where φ and ψ are Möbius transformations of \mathbb{D} .

In the case (b) the map Φ^2 is of the form (a) and taking into consideration the spectral mapping theorem for essential spectra (see [1, Corollary 3.61, p. 149]) we can without loss of generality assume that the map Φ is of the form (a).

We also can and will without loss of generality consider only elliptic Möbius transformations of the form $\varphi(z) = \alpha z, z \in \mathbb{D}, \alpha \in \mathbb{T}$.

Proposition 4.1. Let $\varphi(z_1, z_2) = (\alpha_1 z_1, \alpha_2 z_2)$, where α_1 is a primitive p^{th} -root of unity and α_2 is a primitive q^{th} -root of unity, $p, q \in \mathbb{N}$. Let m = l.c.m.(p,q). Then,

Arkady Kitover and Mehmet Orhon

(1)
$$\sigma(T) = \sigma_{lsf}(T) = \{\lambda \in \mathbb{C} : \exists t \in \mathfrak{M}_A, \lambda^m = w_m(t)\}.$$

(2) $\sigma_{a.p.}(T) = \sigma_{sf}(t) = \{\lambda \in \mathbb{C} : \exists t \in \partial A, \lambda^m = w_m(t)\}$

Proof. The proof follows from Lemma 3.1 and the fact that the set of zeros of $w_m - \lambda^m$ in \mathbb{U}^2 is either empty or infinite.

Proposition 4.2. Let $\varphi(z_1, z_2) = (\alpha_1 z_1, \alpha_2 z_2)$, where $\alpha_1, \alpha_2 \in \mathbb{T}$, α_1 is a primitive p^{th} -root of unity and α_2 is not a root of unity. Let $w \in \mathbb{A}^2$ and let $(Tf)(z_1, z_2) = w(z_1, z_2)f(\varphi(z_1, z_2)), (z_1, z_2) \in \mathbb{D}^2, f \in \mathbb{A}^2$. We assume that $w(z_1, z_2) = z_1^s z_2^t \tilde{w}(z_1, z_2)$, where $s, t \ge 0$ and $|\tilde{w}(0, 0)| > 0$

(1) If w is either an invertible element of \mathbb{A}^2 or w is not invertible in $C(\mathbb{T}^2)$, then

$$\sigma(T) = \sigma_{sf}(T) = \{\lambda \in \mathbb{C} : \exists z_1 \in \mathbb{T}, |\lambda|^p = \tilde{w}_p |(z_1, 0)|\}.$$
 (1)

(2) If w is invertible in $C(\mathbb{T}^2)$ but not invertible in \mathbb{A}^2 , then

$$\sigma_{a.p.}(T) = \sigma_{sf}(T) = \{\lambda \in \mathbb{C} : \exists z_1 \in \mathbb{T}, |\lambda| = |\tilde{w}_p(z_1, 0)|\},\$$

$$\sigma(T) = \sigma_{lsf}(T) = \rho(T)\mathbb{D} = \max_{z_1 \in \mathbb{T}} |\tilde{w}_p(z_1, 0)|^{1/p}\mathbb{D}.$$
(2)

Proof. Let us first assume that p = 1, i.e. that $\varphi(z_1, z_2) = (z_1, \alpha z_2)$, where α is not a root of unity. It is immediate to see that if $\mu \in C(\mathbb{T}^2)'$ is an extreme point of the set of all φ -invariant probability measures, then there is a $z_1 \in \mathbb{T}$ such that μ is the normalised Lebesgue measure on the circle $\{(z_1, z) : z \in \mathbb{T}\}$, and therefore

$$\exp \int \ln |w| d\mu = |\tilde{w}(z_1, 0)|. \tag{3}$$

Therefore, in the case p = 1 the statements (1) and (2) follow from [6, Corollary 2.7, Theorem 2.9] and from the obvious observation that in case (2) $def(T) = \infty$.

To finish the proof in general case we consider the operator T^p and apply the spectral mapping theorem for Fredholm and semi-Fredholm spectra (see [1, Corollary 3.61, p.149])

Proposition 4.3. Let $\varphi(z_1, z_2) = (\alpha_1 z_1, \alpha_2 z_2)$, where $\alpha_1, \alpha_2 \in \mathbb{T}$, where both α_1 and α_2 are not roots of unity.

(1) If w is invertible in \mathbb{A}^2 , then

$$\sigma(T) = \sigma_{sf}(T) = \{\lambda \in \mathbb{C} : \rho_{min}(T) \le |\lambda| < \rho(T)\}.$$
(4)

(2) If w is not invertible in $C(\mathbb{T}^2)$, then

$$\sigma(T) = \sigma_{sf}(T) = \rho(T)\mathbb{D}.$$
(5)

 $\mathbf{6}$

Spectrum of weighted automorphisms.

(3) If w is invertible in $C(\mathbb{T}^2)$ but not invertible in \mathbb{A}^2 , then

$$\sigma_{a.p.}(T) = \sigma_{sf}(T) = \{\lambda \in \mathbb{C} : \rho_{min}(T, C(\mathbb{T})) \le |\lambda| \le \rho(T)\}, \\ \sigma(T) = \sigma_{lsf}(T) = \rho(T)\mathbb{D}.$$
(6)

Proof. The proof follows from Lemma 3.3 and the fact that the set of zeros of w in \mathbb{U}^2 is infinite.

By imposing additional conditions on the map φ we can improve the statement of Proposition 4.3.

Proposition 4.4. Let $\varphi(z_1, z_2) = (\alpha_1 z_1, \alpha_2 z_2)$, where $\alpha_1, \alpha_2 \in \mathbb{T}$. Assume that $w(z_1, z_2) = z_1^s z_2^t \tilde{w}(z_1, z_2)$, where $s, t \ge 0$ and $|\tilde{w}(0, 0)| > 0$. Assume additionally that either (A) $\alpha_1^p, \alpha_2^q \ne 1$ for any $p, q \in \mathbb{Z}$, or (B) $\alpha_1^p = \alpha_2^q$ for some $p, q \in \mathbb{N}$.

(1) If w is invertible in \mathbb{A}^2 , then

$$\sigma(T) = \sigma_{sf}(T) = w(0,0)\mathbb{T}.$$
(7)

(2) If w is not invertible in $C(\mathbb{T}^2)$, then

$$\sigma(T) = \sigma_{sf}(T) = \tilde{w}(0,0)\mathbb{D}.$$
(8)

(3) If w is invertible in $C(\mathbb{T}^2)$ but not invertible in \mathbb{A}^2 , then

$$\sigma_{a.p.}(T) = \sigma_{sf}(T) = \tilde{w}(0,0)\mathbb{T},$$

$$\sigma(T) = \sigma_{lsf}(T) = \tilde{w}(0,0)\mathbb{D}.$$
(9)

Proof. (A) The condition (A) of the proposition guarantees that the only φ -invariant probability measure on \mathbb{T}^2 is the normalised Lebesgue measure. Hence $\rho_{min}(T) = \rho(T) = |\tilde{w}(0,0)|$.

(B) Consider the operator T^q . Notice that

$$(T^{q}f)(z_{1}, z_{2}) = w_{q}(z_{1}, z_{2})f(\alpha_{1}^{q}z_{1}, \alpha_{1}^{p}z_{2}).$$
(10)

Let μ be an extreme point of the set of all φ^{q} -invariant probability measures on \mathbb{T}^{2} . It is easy to see that there is $(z_{1}, z_{2}) \in \mathbb{T}^{2}$ such that $supp(\mu) = cl\{(\alpha_{1}^{qn}z_{1}, \alpha_{1}^{pn}z_{2}), n \in \mathbb{N}\}.$

Therefore $\exp \int \log |w_q| d\mu = |\tilde{w}_q(0,0)|$, and the statement of the theorem for the operator T^q follows. It remains to apply the spectral mapping theorem for essential spectra.

As Example 4.5 shows, the statement of Proposition 4.4 is in general not valid in the case when there are $p, q \in \mathbb{Z}$ such that $\alpha_1^p \alpha_2^q = 1$.

Example 4.5. Let $\alpha \in \mathbb{T}$ be not a root of unity. Let $\varphi(z_1, z_2) = (\alpha z_1, \overline{\alpha} z_2), (z_1, z_2) \in \mathbb{T}^2$, and let

$$(Tf)(z_1, z_2) = (2 + z_1 z_2) f(\varphi(z_1, z_2)), f \in \mathbb{A}^2.$$

It is not difficult to see that

$$\sigma_{sf}(T) = \sigma(T) = \{\lambda \in \mathbb{C} : 1 \le |\lambda| \le 3\}.$$

Proposition 4.6. Let $\varphi(z_1, z_2) = (\alpha z_1, \psi(z_2))$, where α is a primitive p^{th} root of unity and ψ is a parabolic Möbius transformation of U. Let ζ be the fixed point of ψ . Let $r = \min_{\varsigma \in \mathbb{T}} |w_p(\varsigma, \zeta)|^{1/p}$ and $R = \max_{\varsigma \in \mathbb{T}} |w_p(\varsigma, \zeta)|^{1/p}$.

(1) If w is invertible in \mathbb{A}^2 , then

$$\sigma(T) = \sigma_{sf}(T) = \mathbb{AN}(r, R).$$
(11)

(2) If w is not invertible in $C(\mathbb{T}^2)$, then

$$\sigma(T) = \sigma_{sf}(T) = R\mathbb{D}.$$
 (12)

(3) If w is not invertible in \mathbb{A}^2 but invertible in $C(\mathbb{T}^2)$, then

$$\sigma_{a.p.}(T) = \sigma_{sf}(T) = \mathbb{AN}(r, R),$$

$$\sigma(T) = \sigma_{lsf}(T) = R\mathbb{D}.$$
(13)

Proof. (1) and (2) follow from Lemma3.4, Lemma 3.6, and [5, Theorem 3.2]. To prove (3) it suffices to notice that $\sigma_r(T) = r\mathbb{U}$ and that $def T = \infty$ (recall that the set of zeros of w in \mathbb{U}^2 is infinite). \Box

Proposition 4.7. Let $\varphi(z_1, z_2) = (\alpha z_1, \psi(z_2))$, where α is not a root of unity and ψ is a parabolic Möbius transformation of \mathbb{U} . Let ζ be the fixed point of ψ . Assume that $w(z_1, z_2) = z_1^p z_2^q \tilde{w}(z_1, z_2)$, where $\tilde{w}(0, 0) \neq 0$

(1) If w is invertible in \mathbb{A}^2 , then

$$\sigma(T) = \sigma_{sf}(T) = w(0,\zeta)\mathbb{T}.$$
(14)

(2) If w is not invertible in $C(\mathbb{T}^2)$, then

$$\sigma(T) = \sigma_{sf}(T) = \begin{cases} \tilde{w}(0,\zeta)\mathbb{D}, & if \int_{0}^{2\pi} ln|w(e^{i\theta},\zeta)|d\theta > -\infty; \\ & 0\\ \{0\}, & if \int_{0}^{2\pi} ln|w(e^{i\theta},\zeta)|d\theta = -\infty. \end{cases}$$
(15)

(3) If w is not invertible in \mathbb{A}^2 but invertible in $C(\mathbb{T}^2)$, then

$$\sigma_{a.p.}(T) = \sigma_{sf}(T) = \tilde{w}(0,\zeta)\mathbb{T},$$

$$\sigma(T) = \sigma_{lsf}(T) = \tilde{w}(0,\zeta)\mathbb{D}.$$
(16)

8

Proof. Similar to the proof of Proposition 4.6.

Proposition 4.8. Let $\varphi(z_1, z_2) = (\phi(z_1), \psi(z_2))$, where ϕ and ψ are parabolic Möbius transformations of U. Let ς and ζ be the fixed points of ϕ and ψ , respectively.

(1) If w is invertible in \mathbb{A}^2 , then

$$\sigma(T) = \sigma_{sf}(T) = w(\varsigma, \zeta)\mathbb{T}.$$
(17)

(2) If w is not invertible in $C(\mathbb{T}^2)$, then

$$\sigma(T) = \sigma_{sf}(T) = w(\varsigma, \zeta)\mathbb{D}.$$
(18)

(3) If w is not invertible in \mathbb{A}^2 but invertible in $C(\mathbb{T}^2)$, then

$$\sigma_{a.p.}(T) = \sigma_{sf}(T) = w(\varsigma, \zeta)\mathbb{T},$$

$$\sigma(T) = \sigma_{lsf}(T) = w(\varsigma, \zeta)\mathbb{D}.$$
(19)

Proof. Similar to the proof of Proposition 4.6.

Proposition 4.9. Let $\varphi(z_1, z_2) = (\alpha z_1, \psi(z_2))$, where α is a p^{th} primitive root of unity and ψ is a hyperbolic Möbius transformation of U. Let ζ_1, ζ_2 be the fixed points of ψ . Assume that $|\psi'(\zeta_1)| < 1$. For any $\varsigma \in \mathbb{T}$ let $a(\varsigma) = |w_p(\varsigma, \zeta_1)|^{1/p}$ and $b(\varsigma) = |w_p(\varsigma, \zeta_2)|^{1/p}$. Let $r = \min_{\varsigma \in \mathbb{T}} \{|a(\varsigma)|, |b(\varsigma)|\}$ and $R = \max_{\varsigma \in \mathbb{T}} \{|a(\varsigma)|, |b(\varsigma)|\}$. Also, let $\mathbb{E} = \{\varsigma \in \mathbb{T} : |a(\varsigma)| \ge |b(\varsigma)|\}$.

(1) If w is invertible in \mathbb{A}^2 , then

$$\sigma(T) = \mathbb{A}\mathbb{N}(r, R).$$

$$\sigma_{usf}(T) = \bigcup_{\varsigma \in \mathbb{E}} \mathbb{A}\mathbb{N}(|b(\varsigma)|, |a(\varsigma)|).$$

$$\sigma_{lsf}(T) \supseteq cl \bigcup_{\varsigma \in \mathbb{T} \setminus \mathbb{E}} \mathbb{A}\mathbb{N}(|a(\varsigma)|, |b(\varsigma)|).$$
(20)

(2) If w is not invertible in \mathbb{A}^2 but invertible in $C(\mathbb{T}^2)$, then

$$\sigma(T) = R\mathbb{D}.$$

$$\sigma_{usf}(T) = \bigcup_{\varsigma \in \mathbb{E}} \mathbb{AN}(|b(\varsigma)|, |a(\varsigma)|).$$

$$\sigma_{lsf}(T) \supseteq r\mathbb{D} \cup cl \bigcup_{\varsigma \in \mathbb{T} \setminus \mathbb{E}} \mathbb{AN}(|a(\varsigma)|, |b(\varsigma)|).$$
(21)

Arkady Kitover and Mehmet Orhon

If w is not invertible in
$$C(\mathbb{T}^2)$$
, then

$$\sigma(T) = R\mathbb{D}.$$

$$\sigma_{usf}(T) = \bigcup_{\varsigma \in \mathbb{E}} \mathbb{AN}(|b(\varsigma)|, |a(\varsigma)|) \cup \bigcup_{\varsigma \in \mathbb{F}} |a(\varsigma)|\mathbb{D},$$
where $\mathbb{F} = \{\varsigma \in \mathbb{T} : \exists z \in \mathbb{T} \text{ such that } w(\varsigma, z) = 0\}.$

$$\sigma_{lsf}(T) \supseteq cl \bigcup_{\varsigma \in \mathbb{T} \setminus \mathbb{E}} \mathbb{AN}(|a(\varsigma)|, |b(\varsigma)|) \cup \bigcup_{\varsigma \in \mathbb{F}} |b(\varsigma)|\mathbb{D}.$$
(22)

Proof. The proof follows from Lemmas 3.3, 3.5, and 3.6.

Proposition 4.10. Let $\varphi(z_1, z_2) = (\alpha z_1, \psi(z_2))$, where α is not a root of unity and ψ is a hyperbolic Möbius transformation of \mathbb{U} . Let ζ_1, ζ_2 be the fixed points of ψ . Assume that $|\psi'(\zeta_1)| < 1$. Let $w(z_1, z_2) = z_1^p z_2^q \tilde{w}(z_1, z_2)$, where $\tilde{w}(0, 0) \neq 0$.

Let $r = \min(|\tilde{w}(0,\zeta_1)|, |\tilde{w}(0,\zeta_2)|)$ and $R = \max(|\tilde{w}(0,\zeta_1)|, |\tilde{w}(0,\zeta_2)|)$. (1) If w is invertible in \mathbb{A}^2 , then

$$\sigma(T) = \mathbb{AN}(r, R). \tag{23}$$

$$\sigma_{usf}(T) = \begin{cases} \mathbb{A}\mathbb{N}(r,R), & \text{if } |\tilde{w}(0,\zeta_2)| \leq |\tilde{w}(0,\zeta_1)| ;\\ r\mathbb{T} \cup R\mathbb{T}, & \text{if } |\tilde{w}(0,\zeta_1)| \leq |\tilde{w}(0,\zeta_2)| . \end{cases}$$
(24)

$$\sigma_{lsf}(T) \supseteq \begin{cases} \mathbb{A}\mathbb{N}(r,R), & \text{if } |\tilde{w}(0,\zeta_1)| \le |\tilde{w}(0,\zeta_2)| ;\\ r\mathbb{T} \cup R\mathbb{T}, & \text{if } |\tilde{w}(0,\zeta_2)| \le |\tilde{w}(0,\zeta_1)| . \end{cases}$$
(25)

(2) If w is not invertible in \mathbb{A}^2 , but invertible in $C(\mathbb{T}^2)$, then

$$\sigma(T) = R\mathbb{D}.\tag{26}$$

$$\sigma_{usf}(T) = \begin{cases} \mathbb{A}\mathbb{N}(r,R), & \text{if } |\tilde{w}(0,\zeta_2)| \leq |\tilde{w}(0,\zeta_1)| ;\\ r\mathbb{T} \cup R\mathbb{T}, & \text{if } |\tilde{w}(0,\zeta_1)| \leq |\tilde{w}(0,\zeta_2)| . \end{cases}$$
(27)

$$\sigma_{lsf}(T) \supseteq \begin{cases} R\mathbb{D}, & \text{if } |\tilde{w}(0,\zeta_1)| \le |\tilde{w}(0,\zeta_2)| ;\\ r\mathbb{D} \cup R\mathbb{T}, & \text{if } |\tilde{w}(0,\zeta_2)| \le |\tilde{w}(0,\zeta_1)| . \end{cases}$$
(28)

(3) If w is not invertible in $C(\mathbb{T}^2)$, then

$$\sigma(T) = R\mathbb{D},$$

$$\sigma_{usf}(T) = |\tilde{w}(0,\zeta_1)|\mathbb{D} \cup |\tilde{w}(0,\zeta_2)|\mathbb{T},$$

$$\sigma_{lsf}(T) \supseteq |\tilde{w}(0,\zeta_2)|\mathbb{D} \cup |\tilde{w}(0,\zeta_1)|\mathbb{T}.$$
(29)

10

(3)

Proof. The proof follows from Lemmas 3.3, 3.5, 3.6, and from the results in [5].

Proposition 4.11. Let $\varphi(z_1, z_2) = (\phi(z_1), \psi(z_2))$, where ϕ and ψ are a parabolic and a hyperbolic Möbius transformations of \mathbb{U} , respectively. Let ζ_1, ζ_2 be the fixed points of ψ . Assume that $|\psi'(\zeta_1)| < 1$. Let ς be the fixed point of φ .

Let $r = \min(|w(\varsigma, \zeta_1)|, |w(\varsigma, \zeta_2)|)$ and $R = \max(|w(\varsigma, \zeta_1)|, |w(\varsigma, \zeta_2)|)$. (1) If w is invertible in \mathbb{A}^2 , then

$$\sigma(T) = \mathbb{AN}(r, R). \tag{30}$$

$$\sigma_{usf}(T) = \begin{cases} \mathbb{A}\mathbb{N}(r,R), & \text{if } |w(\varsigma,\zeta_2)| \le |w(\varsigma,\zeta_1)| ;\\ r\mathbb{T} \cup R\mathbb{T}, & \text{if } |w(\varsigma,\zeta_1)| \le |w(\varsigma,\zeta_2)| . \end{cases}$$
(31)

$$\sigma_{lsf}(T) \supseteq \begin{cases} \mathbb{A}\mathbb{N}(r,R), & \text{if } |w(\varsigma,\zeta_1)| \le |w(\varsigma,\zeta_2)| ; \\ r\mathbb{T} \cup R\mathbb{T}, & \text{if } |w(\varsigma,\zeta_2)| \le |w(\varsigma,\zeta_1)| . \end{cases}$$
(32)

(2) If w is not invertible in \mathbb{A}^2 , but invertible in $C(\mathbb{T}^2)$, then

$$\sigma(T) = R\mathbb{D}.\tag{33}$$

$$\sigma_{usf}(T) = \begin{cases} \mathbb{A}\mathbb{N}(r,R), & \text{if } |w(\varsigma,\zeta_2)| \le |w(\varsigma,\zeta_1)| ;\\ r\mathbb{T} \cup R\mathbb{T}, & \text{if } |w(\varsigma,\zeta_1)| \le |w(\varsigma,\zeta_2)| . \end{cases}$$
(34)

$$\sigma_{lsf}(T) \supseteq \begin{cases} R\mathbb{D}, & \text{if } |w(\varsigma,\zeta_1)| \le |w(\varsigma,\zeta_2)| ;\\ r\mathbb{D} \cup R\mathbb{T}, & \text{if } |w(\varsigma,\zeta_2)| \le |w(\varsigma,\zeta_1)| . \end{cases}$$
(35)

(3) If w is not invertible in $C(\mathbb{T}^2)$, then

$$\sigma(T) = R\mathbb{D}.\tag{36}$$

$$If |w(\varsigma, \zeta_2)| \le |w(\varsigma, \zeta_1)|, \ then$$

$$\sigma_{usf}(T) = R\mathbb{D},$$

$$\sigma_{lsf}(T) \supseteq r\mathbb{D} \cup R\mathbb{T}.$$

$$If |w(\varsigma, \zeta_1)| \le |w(\varsigma, \zeta_2)|, \ then$$
(37)

$$w(\varsigma, \zeta_1)| \le |w(\varsigma, \zeta_2)|, \text{ then}$$

$$\sigma_{lsf}(T) = R\mathbb{D},$$

$$\sigma_{usf}(T) = r\mathbb{D} \cup R\mathbb{T}.$$
(38)

Proof. The proof follows from Lemmas 3.3, 3.5, 3.6 and from the results in [5]. \Box

It remains to consider the case when $\varphi(z_1, z_2) = (\phi(z_1), \psi(z_2))$, where ϕ and ψ are hyperbolic Möbius transformations of U. We denote the attracting fixed points of ϕ and ψ as ζ_1, ζ_1 and the repelling points as ζ_2, ζ_2 , respectively. Let $w \in \mathbb{A}^2$ and let

$$Tf(z_1, z_2) = w(z_1, z_2) f(\phi(z_1), \psi(z_2)), f \in \mathbb{A}^2, (z_1, z_2) \in \mathbb{D}^2.$$
(39)

Let F be the set of all four fixed points of φ and let $r = \min_{(z_1, z_2) \in F} |w(z_1, z_2)|$, $R = \max_{(z_1, z_2) \in F} |w(z_1, z_2)|$. Then,

$$\sigma(T) = \begin{cases} \mathbb{A}\mathbb{N}(r, R), & \text{if } w \text{ is invertible in } \mathbb{A}^2, \\ R\mathbb{D}, & \text{if } w \text{ is not invertible in } \mathbb{A}^2. \end{cases}$$
(40)

The partial description of the essential spectra of T depends on the order of the numbers $|w(z_1, z_2)|, (z_1, z_2) \in F$ on the real line and on invertibility or non-invertibility of the weight w in \mathbb{A}^2 or in $C(\mathbb{T}^2)$, and therefore there are 72 distinct cases. One of these cases is considered in the following proposition.

Proposition 4.12. Let T be the operator of the form (39). Assume that w is invertible in \mathbb{A}^2 . Let $r = |w(\varsigma_1, \zeta_1)| < |w(\varsigma_1, \zeta_2)| < |w(\varsigma_2, \zeta_1)| < |w(\varsigma_2, \zeta_2)| = R$. Then,

$$\sigma_{lsf}(T) = \sigma(T) = \mathbb{A}\mathbb{N}(r, R),$$

$$\sigma_{usf}(T) = r\mathbb{T} \cup |w(\varsigma_1, \zeta_2)|\mathbb{T} \cup |w(\varsigma_2, \zeta_1)|\mathbb{T} \cup R\mathbb{T}.$$
(41)

Proof. The proof follows from Lemmas 3.5 and 3.6.

References

- [1] Aiena P., Fredholm and local spectral theory with applications to multipliers, Kluwer Academic Publishers (2004).
- [2] Edmunds D.E. and Evans W.D., Spectral theory and differential operators, Clarendon Press (1987).
- [3] Kamowitz H., The spectra of a class of operators on the disc algebra., Indiana University Mathematics Journal, 27, No 4, 581-610 (1978).
- [4] Kitover A.K., Spectrum of weighted composition operators: part 1. Weighted composition operators on C(K) and uniform algebras. Positivity, 15, 639 - 659 (2011).
- [5] Kitover A.K., Spectrum of Weighted Composition Operators. Part III: Essential Spectra of Some Disjointness Preserving Operators on Banach Lattices. Ordered Structures and Applications: Positivity VII Trends in Mathematics, Springer International Publishing, 233–261 (2016).
- [6] Kitover A.K., Orhon M., Spectrum of weighted composition operators, part IX. The spectrum and essential spectra of some weighted composition operators on uniform algebras. arXiv:2307.01112v2 [math.SP], submitted to the proceedings of the conference ICMASC'2024, Porto, Portugal, 2024.
- [7] Rudin W. Function theory in polydiscs, W. A. BENJAMIN, INC. (1969)

Community College of Philadelphia, 1700 Spring Garden St., Philadelphia, PA, USA

Email address: akitover@ccp.edu

UNIVERSITY OF NEW HAMPSHIRE, 105 MAIN STREET DURHAM, NH 03824 *Email address:* mo@unh.edu