## Topological Origin of Floquet Thermalization in Periodically Driven Many-body Systems

Hao-Yue Qi,<sup>1,2,\*</sup> Yue Wu,<sup>3,\*</sup> and Wei Zheng<sup>1,2,4,†</sup>

<sup>1</sup>Hefei National Research Center for Physical Sciences at the Microscale and School of Physical Sciences,

University of Science and Technology of China, Hefei 230026, China

<sup>2</sup>CAS Center for Excellence in Quantum Information and Quantum Physics,

University of Science and Technology of China, Hefei 230026, China

<sup>3</sup>Institute for Advanced Study, Tsinghua University, Beijing 100084, China

<sup>4</sup>Hefei National Laboratory, University of Science and Technology of China, Hefei 230088, China

(Dated: April 30, 2024)

Floquet engineering is a powerful manipulation method in modern quantum technology. However, unwanted heating is the main challenge of Floquet engineering, therefore the Floquet thermalization has attracting considerable attentions recently. In this work, we investigate thermalization of periodically driven many-body systems through the lens of Krylov complexity, and find a topological origin of different thermalization behaviors. We demonstrate that If the topology of the Krylov chain is nontrivial, a periodically driven system will reach a state with finite temperature. When the Krylov chain is topologically trivial, the system will be heated to infinite temperature. We further show that the prethermalization can be understood as the tunnelling process of a quasi-edge mode through the local gap on Krylov chain. This picture provides a systematically method to obtain the effective prethermal Hamiltonian.

Quantum systems with periodic driving have attracted much attention in the decades. First, periodic driving can be used to engineer nontrivial Hamiltonians. For example, periodic driving has been used to generate artificial magnetic fields in optical lattice, simulate synthetic lattice gauge theories with ultracold atomic gases [1], and engineer topological bands [2, 3]. Second, periodic driving can induce intrinsically out-of-equilibrium phenomena without counterpart in equilibrium, such as discrete time-crystals [4] and anomalous Floquet topological states [5-8]. However, since periodic driving breaks energy conservation, it was expected that under driving, generic many-body systems would be heated up to featureless infinite temperature ensembles eventually [9–12]. Fortunately, under high-frequency driving, there is a class of systems that exhibit the prethermalization phenomena [13–22]. Such systems have several approximate conservation quantities. Thus, at intermediate time scales, systems will first equilibrate to a finite temperature state governed by a so-called prethermal Hamiltonian. Later, systems will thermalize to infinite temperature ensembles at much longer time scales. Prethermalization has been experimentally observed in one-dimensional Bose gas and spins with dipole interaction [23-25]. However, in what circumstances, a quantum system will be directly heated to infinite temperature or exhibit prethermalization behavior is not clear. There is no simple and clear physical picture of the Floquet thermalization.

Recently, the Krylov complexity of operators has been developed to study the chaos and thermalization in many-body systems [26]. With the help of Krylov basis, the evolution of an operator under Heisenberg equation can be mapped to the spreading of a wavepacket in a one-dimensional tight-binding chain. Krylov complexity has motivated a lot of interesting studies in closed [27–37] [38–42] and open [43–45] quantum many-body systems.

In this work, we found topological origin of different thermalization behavior of periodically driven manybody systems based on Krylov complexity. We first build time-independent Krylov basis for periodically driven systems, and map the Floquet dynamics of an operator to the wavepacket evolution on the Krylov chain. The main results of this work are summarized here:

(1) We construct the unfold Floquet effective Hamiltonian by seeking the zero mode on Krylov chain. Unlike the effective Hamiltonian obtained by taking the logarithm of evolution operator in previous works, which has folded spectrum and does not satisfy the assumption of eigenstate thermalization hypothesis (ETH), our unfold effective Hamiltonian satisfies ETH and can be used to estimate local observables at long time scale.

(2) We further find that it is the topology of Krylov chain that determined the Floquet thermalization behavior. In a finite system, when the Krylov chain is topologically nontrivial, it sustains a localized zero edge mode. Then the system will heated to a state with finite temperature. If topology of the Krylov chain is trivial, the zero mode is extended to the bulk. As a result, the system will be heated to infinite temperature.

(3) In the thermodynamic limit, the Krylov chain is gapless. Therefore there is no eigen zero edge mode, and the system will finally be heated to infinite temperature. However, we find that there could exist a quasi edge mode in high driving frequency regime, which is protected by the local topology and gap of the Krylov chain. We further conjecture that this quasi edge mode represents the Floquet prethermal Hamiltonian. Its life time determines the time scale of intermediate prethermal state. With



FIG. 1: Hopping amplitude  $b_n$  and zero mode  $|H_{\text{eff}}^{\text{zm}}\rangle$  on the Krylov chain. (a)(c) quantum spin chain with system size L = 14 and  $h_z/J = 0.809, h_x/J = 0.9045$ . (b)(d) hard-core Bose-Hubbard model with system size L = 16 and J'/J = 0.2, U/J = 1. For both two models, the hopping amplitude fluctuates around a non-zero value site-by-site in high-frequency regime, and the fluctuation becomes smaller deep in the bulk. The zero mode is an edge mode. In contrast, in the low-frequency regime, the hopping amplitude is nearly constant. The zero mode extends over the bulk.

this picture in mind, one can systematically estimate the prethermal Hamiltonian via Krylov basis.

Krylov Chain. In a periodically driven system,  $\hat{H}(t + T) = \hat{H}(t)$ , the evolution operator during one driving period is given by  $\hat{U}_{\rm F} = \hat{\mathcal{T}} \exp\left[-i\int_0^T dt \hat{H}(t)\right]$ , here we set  $\hbar = 1$ . One can define a time-independent effective Hamiltonian as

$$\hat{U}_{\rm F} = e^{-iH_{\rm eff}T}.$$
(1)

This effective Hamiltonian reproduces the evolution of the driven system in a stroboscopic fashion. In other words, the state evolution under time-independent  $\hat{H}_{\text{eff}}$  is identical to the one under driven Hamiltonian  $\hat{H}(t)$ , only at steps of the driving period t = nT. Thus the effective Hamiltonian well describes the dynamics over time scale much longer than a single driving period. Numerically, we calculate the effective Hamiltonian by taking the logarithm of  $\hat{U}_{\text{F}}$ , as  $\hat{H}_{\text{eff}}^{\text{fold}} = \frac{i}{T} \ln \hat{U}_{\text{F}}$ . Therefore, the long-time evolution of operators can be captured by the following Heisenberg equation in a stroboscopic way,

$$i\partial_t \hat{O} = \left[\hat{O}, \hat{H}_{\text{eff}}^{\text{fold}}\right].$$
 (2)

However, the effective Hamiltonian obtained by this method is not unique. It has a folded energy spectrum. Specifically speaking,  $\hat{H}_{\text{eff}}^{\text{fold}}$  can be diagonalized into  $\hat{H}_{\text{eff}}^{\text{fold}} = \sum_{k} E_{k} |\psi_{k}\rangle \langle \psi_{k}|$ , and its eigen energies are folded in the first Floquet-Brillouin zone,  $-\pi/T < E_{k} < \pi/T$ . As a result, it does not satisfy the assumption of ETH,

and can not predict the expectation values of local observables after a long time evolution. Certainly, we can construct other equivalent effective Hamiltonian by shifting the eigenenergies without altering the eigenstates as  $\hat{H}_{\text{eff}} = \sum_{k} (E_k + m_k \Omega) |\psi_k\rangle \langle \psi_k |$ , where  $m_k$  is an integer and  $\Omega = 2\pi/T$  is the driving frequency. Such Hamiltonians satisfy the definition (1), and thus commute with  $\hat{H}_{\text{eff}}^{\text{fold}}$ .

The Krylov basis of operators is based on the Lanczos algorithm. An operator is denoted by  $\hat{O} = \sum_{ij} O_{ij} |i\rangle \langle j|$ , where  $\{|i\rangle\}$  is a set of orthogonal states and  $O_{ij}$  are corresponding matrix elements. We can map the operator  $\hat{O}$  to a vector in the operator space as  $O \longrightarrow$  $|O\rangle = \sum_{ij} O_{ij} |i\rangle \otimes |j\rangle$ . The corresponding inner product of operators is defined as  $(O_1|O_2) = \text{Tr}(\hat{O}_1^{\dagger}\hat{O}_2)$ . We further introduce a Liouville super-operator  $\mathcal{L}$ , acting on an operator as  $\mathcal{L}\hat{O} \equiv [\hat{H}_{\text{eff}}^{\text{fold}}, \hat{O}]$ . In the operator space, it can be mapped into a matrix,  $\mathcal{L}O \to \mathcal{L}|O\rangle$ . Thus the Heisenberg equation (2) can be expressed into  $i\partial_t |O\rangle = -\mathcal{L} |O\rangle$ . Applying the Baker-Campbell-Hausdorff formula, we expand the evolution of operator into  $|O(t)\rangle = \sum_{n} \frac{(-it)^{n}}{n!} \mathcal{L}^{n} |O\rangle$ . Here  $\{\mathcal{L}^{n} |O\rangle\}$  forms a complete set of operator bases, but it is neither normalized nor orthogonal. We then apply the Gram-Schmidt procedure to obtain a set of orthogonalized and normalized basis, called Krylov basis as,

$$|W_0) = \frac{1}{b_0} |O\rangle, \qquad (3)$$

$$|W_1\rangle = \frac{1}{b_1} \mathcal{L} |W_0\rangle,$$
 (4)

$$|W_n) = \frac{1}{b_n} \left[ \mathcal{L} | W_{n-1} \right] - b_{n-1} \left[ W_{n-2} \right], n \ge 2.$$
 (5)

Here  $b_n$  are the so-called Lanczos coefficients. They are matrix elements of Liouville super-operator in the new basis,  $b_n = (W_n | \mathcal{L} | W_{n+1})$ . We expand the operator in the Krylov basis as  $|O(t)\rangle = \sum_n \varphi_n(t) | W_n\rangle$ . The coefficients  $\varphi_n(t)$  satisfy the following linear equation,

$$i\partial_t\varphi_n = -\left(b_{n-1}\varphi_{n-1} + b_n\varphi_{n+1}\right). \tag{6}$$

Therefore, the operator evolution is mapped to a quantum mechanics problem of one particle hopping on a tight-binding half-infinite chain. Note that the  $b_n$  play the role of the hopping amplitude on this half-infinite Krylov chain. The particle is initially localized at original site, n = 0. During the evolution, it hops into the bulk. The Krylov complexity is defined as  $(O(t)|\mathcal{K}|O(t)) =$  $\sum_n n|\varphi_n(t)|^2$ , where  $\mathcal{K} = \sum_n n |W_n\rangle (W_n|$  is the Krylov super-operator. It is nothing but the center-of-mass of the wave function on the Krylov chain.

*Models.* To illustrate the Floquet thermalization, we study two models with periodical driving. One is a quantum spin chain, the other is a one-dimensional hard-core



FIG. 2: Comparison of zero mode and effective Hamiltonian obtained via Floquet-Magnus expansion in high-frequency regime on Krylov chain. (a) quantum spin chain at TJ = 1, (b) hard-core Bose-Hubbard model at TJ = 1.3. For both models, the zero mode is consistent with the Floquet-Magnus expansion. The simulation parameters are the same as in Fig.1.

Bose-Hubbard Model. The evolution operators of both models in one period are

$$\hat{U}_{\rm F}^{\alpha} = e^{-i\hat{H}_{\alpha}T/2}e^{-i\hat{V}_{\alpha}T/2},\tag{7}$$

where  $\alpha = \text{spin}$ , boson, for spin and Bose-Hubbard models respectively. The corresponding  $\hat{H}_{\alpha}$  are

$$\hat{H}_{\rm spin} = \sum_{i=1}^{L} (-J\sigma_i^z \sigma_{i+1}^z + h_z \sigma_i^z), \qquad (8)$$

$$\hat{H}_{\text{boson}} = -\sum_{i=1}^{L} (J \hat{a}_{i}^{\dagger} \hat{a}_{i+1} + J' \hat{a}_{i}^{\dagger} \hat{a}_{i+2} + \text{H.c.}). \quad (9)$$

And  $\hat{V}_{\alpha}$  are given by  $\hat{V}_{\text{spin}} = \sum_{i=1}^{L} h_x \sigma_i^x$ ,  $\hat{V}_{\text{boson}} = \sum_{i=1}^{L} U \hat{n}_i \hat{n}_{i+1}$ , where  $\hat{n}_i = \hat{a}_i^{\dagger} \hat{a}_i$ . Both models are chaotic.

Unlike previous studies [38–42], we choose the seed operator as the average Hamiltonian in one period,

$$\hat{O} = \hat{H}^{(0)} = \frac{1}{T} \int_0^T dt \hat{H}(t).$$
(10)

For the two models investigated here, we have  $\hat{H}^{(0)}_{\alpha} = (\hat{H}_{\alpha} + \hat{V}_{\alpha})/2$ . We numerically calculate the evolution operator  $\hat{U}_{\rm F}$  and its logarithm, i.e.  $\hat{H}^{\rm fold}_{\rm eff}$  with finite system size. Then we obtain the Krylov basis by the method described above. After that, we expand super-operator  $\mathcal{L}$  on this basis, and obtain the hopping amplitude on the Krylov chain,  $b_n$ , with varying driving frequencies. Typical distributions of  $b_n$  are plotted in Fig.1(a)(b). We find that, for both two models, the hopping amplitude fluctuate around a non-zero value site-by-site in high-frequency driving regime. It reminds us the celebrated Su-Schrieffer-Heeger (SSH) model, which possesses non-trivial topological band structure and can sustain ingap edge states. The fluctuation is large near the origin site, but becomes smaller deep in the bulk. In contrast, in the

low-frequency regime, the fluctuation of hopping amplitude is highly suppressed, and  $b_n$  is nearly constant.

We then numerically seek the zero mode operator of Liouvillian,  $\mathcal{L}|O\rangle = 0$ . That is equivalent to solving the single-particle eigen wave function with zero energy on the Krylov chain, see Fig.1(c)(d). We further conclude that the zero mode obtained by this method is one particular choice of effective Hamiltonian  $\hat{H}_{\text{eff}}^{\text{zm}}$  with unfolded energy spectrum. The reasons are as follows. First, this effective Hamiltonian commutes with  $\hat{H}_{\text{eff}}^{\text{fold}}$ , thus is a zero mode of Liouvillian,  $\mathcal{L}\hat{H}_{\text{eff}}^{\text{zm}} = [\hat{H}_{\text{eff}}^{\text{fold}}, \hat{H}_{\text{eff}}^{\text{sm}}] = 0$ . Second, we analytically calculate the effective Hamiltonian via Floquet-Magnus expansion up to the 5th order in the high-frequency regime [46], and compare it with the zero mode on the Krylov basis. The results are plotted in Fig.2. Note that for both models, the zero mode is consistent with the Floquet-Magnus expansion.

We note that the distribution of  $\hat{H}_{\text{eff}}^{\text{zm}}$  on the Krylov chain varies as driving period changes, see Fig.1(c)(d). In short driving period regime, the effective Hamiltonian is a edge mode, localizing near the origin of Krylov chain. Thus it possesses small Krylov complexity. In long driving period regime,  $\hat{H}_{\text{eff}}^{\text{zm}}$  extends over the bulk, and has larger Krylov complexity. The behaviors of the zero mode can be understood by the band topology of the Krylov chain. Since distribution of  $b_n$  does not possess the translation symmetry, it is hard to investigate the topology in momentum space. We turn to calculate the topological invariance, the winding number, of Krylov chain via a real-space approach as [47–49],

$$\nu = \frac{1}{2N} \operatorname{Tr} \left( \mathcal{SQ} \left[ \mathcal{Q}, \mathcal{K} \right] \right).$$
 (11)

Here  $S = \sum_{n} (|W_{2n}) (W_{2n}| - |W_{2n+1}) (W_{2n+1}|)$  is the chiral-symmetry super-operator on the Krylov chain, and N is the lattice length. One note that  $\mathcal{SLS}^{\dagger} = -\mathcal{L}$ , thus the eigen values of  $\mathcal{L}$  come in pairs. The super-operator  $\mathcal{Q}$  is given by  $\mathcal{Q} = \sum_{i} (|A_i| (A_i) - |A_i|) (A_i')$ , where  $|A_i|$ are the eigen modes of  $\mathcal{L}$  with positive eigen values  $\lambda_i$ ,  $\mathcal{L}|A_i\rangle = \lambda_i |A_i\rangle$ , and  $|A'_i\rangle$  are chiral-symmetry partners of  $|A_i\rangle$ ,  $|A'_i\rangle = S|A_i\rangle$ . Therefore they have negative eigen values,  $\mathcal{L} |A'_i| = -\lambda_i |A'_i|$ . We numerically calculate the winding number  $\nu$ , and compare it with the overlap  $(H^{(0)}|H_{\text{eff}}^{\text{zm}})$  as a function of driving frequency. The latter represents the weight of zero mode on the origin site, and can be used to characterize the edge state. If  $(H^{(0)}|H_{\text{eff}}^{\text{zm}}) \sim 1$ , that indicates the zero mode is well localized on the edge. The results are plotted in Fig.3(d)and Fig.4(d). Note that the band topology of the Krylov chain determines the zero edge mode. In the short driving period regime, winding number  $\nu \approx 1$ , that protects a zero edge mode on the boundary. In long driving period regime, since the gap is closed, the winding number is not well defined, and gives a non-integer. Thus zero mode is extended to bulk.

Unlike the folded effective Hamiltonian  $\hat{H}_{\text{eff}}^{\text{fold}}$ , we found



FIG. 3: Verifying ETH and winding number  $\nu$  of Krylov chain in the quantum spin model. (a-c) Long-time equilibrium values of  $\langle \hat{\sigma}_1^x(mT) \rangle$ ,  $\langle \hat{\sigma}_1^z \hat{\sigma}_2^z(mT) \rangle$  and  $\langle \hat{H}^{(0)}(mT) \rangle$ , as  $m \to \infty$ , are compared to ETH predictions at different driving periods. The dashed line marks the infinite-temperature value. (d) Local winding number  $\nu$  of the Krylov chain is compared with the overlap  $(H^{(0)}|H_{\text{eff}}^{\text{zm}})$ , which exhibits the localization of zero mode on the first Krylov site. We work in zero momentum sector of positive parity. The initial state is chosen as  $Z_2$  state  $\mathcal{P}|\uparrow\downarrow\cdots\uparrow\downarrow\rangle$ , where  $\mathcal{P}$  is the projector onto this sector. The simulation parameters are the same as in Fig.1.

that this zero mode  $\hat{H}_{\text{eff}}^{\text{zm}}$  satisfies the assumption of ETH [?]. We can use it to predict the long-time equilibrium values of local observables as

$$\left\langle \hat{O}_{\text{local}}(mT \to \infty) \right\rangle \approx \text{Tr}\left( \hat{\rho}_{\beta} \hat{O}_{\text{local}} \right),$$
 (12)

where the canonical ensemble density matrix,  $\hat{\rho}_{\beta} =$  $e^{-\beta \hat{H}_{\text{eff}}^{\text{zm}}}/\text{Tr}(e^{-\beta \hat{H}_{\text{eff}}^{\text{zm}}})$ . The inverse temperature  $\beta$  is obtained by solving a single equation,  $\langle \psi(0) | \hat{H}_{\text{eff}}^{\text{zm}} | \psi(0) \rangle =$  $\operatorname{Tr}(\hat{H}_{\text{eff}}^{\text{zm}}\hat{\rho}_{\beta})$ . Here  $|\psi(0)\rangle$  is a generic initial pure state. In Fig.3 and Fig.4, we plot the comparison of local observables after long time evolution with the ETH prediction. All these calculations start from the same initial pure state. Note that, for both models, the long time expectation values are consistent with the ETH prediction not even in high frequency limit, but in the whole frequency regime. By comparing the long time thermalization behavior with  $(H^{(0)}|H_{\text{eff}}^{\text{zm}})$  (Fig.3 and Fig.4), we note that when the zero mode is localized on the edge, system will be driven into an ensemble with finite temperature. If the zero mode is extended to the bulk, the system will be heated to infinite temperature. Thus we conclude that it is the band topology of Krylov chain that determines the long time thermalization of periodically driven manybody systems. When the Krylov chain is topologically non-trivial, the Floquet heating will be bounded to finite energy density. When the Krylov chain is topological trivial or gapless, the heating will be unbounded.

Prethermalization. In the thermodynamic limit, the hopping amplitude  $b_n$  at large n approaches a constant.



FIG. 4: Verifying ETH and winding number  $\nu$  of Krylov chain in the hard-core Bose-Hubbard model. (a-c) Longtime equilibrium values of  $\langle \hat{b}_1^{\dagger} \hat{b}_2(mT) + h.c. \rangle$ ,  $\langle \hat{n}_1 \hat{n}_2(mT) \rangle$  and  $\langle \hat{H}^{(0)}(mT) \rangle$ , as  $m \to \infty$ , are compared to ETH predictions at different driving periods. The dashed line marks the infinitetemperature value. (d) Local winding number  $\nu$  of the Krylov chain is compared with the overlap  $(H^{(0)}|H_{\text{eff}}^{\text{zm}})$ , which exhibits the localization of zero mode on the first Krylov site. We work in zero momentum sector of positive parity. The initial state is chosen as ground state of  $\hat{H}_{\text{boson}}$ . The simulation parameters are the same as in Fig.1.

Therefore, the Krylov chain is gapless, and there is no eigen edge mode. As a result, the system will be finally heated to infinite temperature despite the frequency. However, in the high frequency limit, the configuration of hopping amplitude near the origin site is still similar to the SSH model. That indicates, there exists local gap near the origin. To see this, we plot the local density-of-state  $\rho_n(\omega) = -\frac{1}{\pi} \text{Im} \langle W_n | (\omega - \mathcal{L} + i0^+)^{-1} | W_n \rangle$  on the Krylov chain in Fig.5. Such local gap will sustain a quasi eigen edge mode, denoted by  $\hat{H}_{\text{eff}}^{\text{quasi}}$ . This quasi edge mode has a finite lifetime,  $\tau_*$ . Before this time scale, the initial wavepacket stays localized near the edge. But after  $\tau_*$ , it will tunnel across the local gap, and spread to the bulk.

The damping of this quasi edge mode in the Krylov chain describes the Floquet prethermalization. The lifetime of the quasi edge mode  $\tau_*$  is just the time scale of prethermalization. During this time scale, the system will thermalize to a quasi steady state with finite temperature given by  $\hat{H}_{\text{eff}}^{\text{quasi}}$ . After that, the system will be continuously heated to infinite temperature. To see that, we plot the thermalization process of local observables in a long time scale. We find that their prethermal values are consistent with the prediction of ETH given by  $\hat{H}_{\text{eff}}^{\text{quasi}}$ , but away from the prediction of  $\hat{H}_{\text{eff}}^{\text{zm}}$ , and  $\hat{H}^{(0)}$ see Fig.5(a)(c). Numerically, we approximate the  $\hat{H}_{\text{eff}}^{\text{quasi}}$ by truncating the zero mode to the length of local gap regime. Therefore, we conclude that the Floquet prethermalization process is dominated by the local topological



FIG. 5: Real time thermalization dynamics and local densityof-state on Krylov chains. (a-b) is the quatnum spin model, with L = 14 and TJ = 1.2. (c-d) is the hard-core Bose-Hubbard model with L = 18 and TJ = 3.9. (a)(c) The real time thermalization dynamics of local observable  $\langle \hat{H}_{\alpha}^{(0)}(mT) \rangle$ in a long time scale. The predicted values of ETH given by  $\hat{H}_{\text{eff}}^{(0)}$ ,  $\hat{H}_{\text{eff}}^{\text{quasi}}$  and  $\hat{H}_{\text{eff}}^{\text{zm}}$  are shown by green, yellow and red lines. The dotted gray vertical line marks the value of prethemalization time scale  $\tau_*$ . (b)(d) Local density-of-state on Krylov chains. Note that there exists local gap near the origin for both models.

structure on Krylov chain. With this picture in mind, one can optimize the prethermal Hamiltonian as the proper quasi edge mode  $\hat{H}_{\text{eff}}^{\text{quasi}}$ .

Summary. In summary, we study thermalization of periodically driven many-body quantum systems via Krylov basis. We found that the zero mode on the Krylov chain represents the unfolded Floquet effective Hamiltonian. We demonstrate that the band topology of the Krylov chain dominates the thermalization behavior. We show that the prethermalization phenomena can be understood as the damping process of a quasi edge mode protected by the local gap. Our results bring out the generic connection between Krylov complexity and thermalization of periodically driven systems that deserves further studies.

## Acknowledgments

We thank H. Zhai and Z. Wang for helpful discussion. This work is supported by NSFC (Grant No. GG2030007011 and No. GG2030040453) and Innovation Program for Quantum Science and Technology (No.2021ZD0302004).

- \* These authors contribute equally to this work.
- <sup>†</sup> Electronic address: zw8796@ustc.edu.cn

- C. Schweizer, F. Grusdt, M. Berngruber, L. Barbiero, E. Demler, N. Goldman, I. Bloch, and M. Aidelsburger, Floquet approach to Z2 lattice gauge theories with ultracold atoms in optical lattices, Nat. Phys. 15, 1168 (2019)
- [2] G. Jotzu, M. Messer, R. Desbuquois, M. Lebrat, T. Uehlinger, D. Greif, T. Esslinger, Experimental realization of the topological Haldane model with ultracold fermions, Nature 515, 237 (2014)
- [3] W. Zheng and H. Zhai, Floquet topological states in shaking optical lattices, Phys. Rev. A 89, 061603(R) (2014)
- [4] M. P. Zaletel, M. Lukin, C. Monroe, C. Nayak, F. Wilczek, and N. Y. Yao, Quantum and classical discrete time crystals, Rev. Mod. Phys. 95, 031001 (2023)
- [5] T. Kitagawa, E. Berg, M. Rudner, and E. Demler, Topological characterization of periodically driven quantum systems, Phys. Rev. B 82, 235114 (2010)
- [6] M. S. Rudner, N. H. Lindner, E. Berg, and M. Levin, Anomalous Edge States and the Bulk-Edge Correspondence for Periodically Driven Two-Dimensional Systems, Phys. Rev. X, 3, 031005 (2013)
- [7] K. Wintersperger, C. Braun, F. Nur Ünal, A. Eckardt, M. Di Liberto, N. Goldman, I. Bloch, M. Aidelsburger, Realization of an anomalous Floquet topological system with ultracold atoms, Nat. Phys. 16, 1058 (2020)
- [8] P. Xu, W. Zheng and H. Zhai, Topological micromotion of Floquet quantum systems, Phys. Rev. B 105, 045139 (2022)
- [9] A. Lazarides, A. Das, and R. Moessner, Equilibrium states of generic quantum systems subject to periodic driving, Phys. Rev. E 90, 012110 (2014)
- [10] L. D.lessio and M. Rigol, Long-time Behavior of Isolated Periodically Driven Interacting Lattice Systems, Phys. Rev. X 4, 041048 (2014)
- [11] H. Kim, T. N. Ikeda, and D. A. Huse, Testing whether all eigenstates obey the eigenstate thermalization hypothesis, Phys. Rev. E 90, 052105 (2014)
- [12] P. Ponte, A. Chandran, Z. Papić, and D. A. Abanin, Periodically driven ergodic and many-body localized quantum systems, Annals of Physics **353**, 196 (2015)
- [13] D. A. Abanin, W. De Roeck, and F. Huveneers, Exponentially Slow Heating in Periodically Driven Many-Body Systems, Phys. Rev. Lett. **115**, 256803 (2015)
- [14] T. Mori, T. Kuwahara, and K. Saito, Rigorous Bound on Energy Absorption and Generic Relaxation in Periodically Driven Quantum Systems, Phys. Rev. Lett. 116, 120401 (2016)
- [15] T. Kuwahara, T. Mori, K. Saitoc, Floquet-Magnus theory and generic transient dynamics in periodically driven many-body quantum systems, Annals of Physics 367, 96 (2016)
- [16] D. Abanin, W. De Roeck, W. W. Ho, and F. Huveneers, A Rigorous Theory of Many-Body Prethermalization for Periodically Driven and Closed Quantum Systems. Comm. Math. Phys. **354**, 809 (2017).
- [17] M. Bukov, M. Heyl, D. A. Huse, and A. Polkovnikov, Heating and many-body resonances in a periodically driven two-band system, Phys. Rev. B 93, 155132 (2016)
- [18] S. A. Weidinger and M. Knap, Floquet prethermalization and regimes of heating in a periodically driven, interacting quantum system, Scientific Reports,
- [19] E. Canovi, M. Kollar, and M. Eckstein, Stroboscopic prethermalization in weakly interacting periodi-

cally driven systems, Phys. Rev. E 93, 012130 (2016)

- [20] D. A. Abanin, W. De Roeck, W. W. Ho, and F. Huveneers, Effective Hamiltonians, prethermalization, and slow energy absorption in periodically driven many-body systems Phys. Rev. B 95, 014112 (2017)
- [21] C. Fleckenstein and M. Bukov, Thermalization and prethermalization in periodically kicked quantum spin chains, Phys. Rev. B 103, 144307 (2021)
- [22] W. W. Ho, T. Mori, D. A. Abanin, E. G. Dalla Torre, Quantum and classical Floquet prethermalization, Annals of Physics 454, 169297 (2023)
- [23] A. Rubio-Abadal, M. Ippoliti, S. Hollerith, D. Wei, J. Rui, S. L. Sondhi, V. Khemani, C. Gross, and I. Bloch, Floquet Prethermalization in a Bose-Hubbard System, Phys. Rev. X 10, 021044 (2020)
- [24] P. Peng, C. Yin, X. Huang, C. Ramanathan, and P. Cappellaro, Floquet prethermalization in dipolar spin chains, Nat. Phys. 17, 444 (2021)
- [25] W. Beatrez, O. Janes, A. Akkiraju, A. Pillai, A. Oddo, P. Reshetikhin, E. Druga, M. McAllister, M. Elo, B. Gilbert, D. Suter, and A. Ajoy Floquet Prethermalization with Lifetime Exceeding 90 s in a Bulk Hyperpolarized Solid, Phys. Rev. Lett. **127**, 170603 (2021)
- [26] D. E. Parker, X. Cao, A. Avdoshkin, T. Scaffidi, and E. Altman, A universal operator growth hypothesis, Phys. Rev. X 9, 041017 (2019).
- [27] C. Lv, R. Zhang, and Q. Zhou, Building Krylov complexity from circuit complexity, arXiv:2303.07343
- [28] R. Zhang and H. Zhai, Universal Hypothesis of Autocorrelation Function from Krylov Complexity, arXiv:2305.02356
- [29] A. Dymarsky and M. Smolkin, Krylov complexity in conformal field theory, arXiv:2104.09514
- [30] E. Rabinovici, A. Sánchez-Garrido, R. Shira and J. Sonnerb, Krylov localization and suppression of complexity, JHEP 03, 211 (2022)
- [31] P. Caputa, J. M. Magan, and D. Patramanis, Geometry of Krylov complexity, Phys. Rev. Res. 4, 013041 (2022)
- [32] Universal chaotic dynamics from Krylov space, J. Erdmenger, S.-K. Jian, and Z.-Y. Xian, arXiv:2303.12151
- [33] A. Dymarsky and A. Gorsky, Quantum chaos as delocalization in Krylov space, Phys. Rev. B 102, 085137 (2020)
- [34] D. J. Yates, A. G. Abanov, and A. Mitra, Lifetime of Almost Strong Edge-Mode Operators in One-Dimensional, Interacting, Symmetry Protected Topological Phases, Phys. Rev. Lett. **124**, 206803 (2020)

- [35] H.-C. Yeh, G. Cardoso, L. Korneev, D. Sels, A. G. Abanov, and A. Mitra, Slowly decaying zero mode in a weakly nonintegrable boundary impurity model, Phys. Rev. B 108, 165143 (2023)
- [36] P. Zhang and Z. Yu, Dynamical transition of operator size growth in quantum systems embedded in an environment, Phys. Rev. Lett. 130, 250401 (2023)
- [37] L. Chen, B. Mu, H. Wang, and P. Zhang, Dissecting Quantum Many-body Chaos in the Krylov Space, arXiv:2404.08207
- [38] D. J. Yates and A. Mitra, Strong and almost strong modes of Floquet spin chains in Krylov subspaces Phys. Rev. B 104, 195121 (2021)
- [39] D, J. Yates, A. G. Abanov, and A. Mitra Long-lived period-doubled edge modes of interacting and disorderfree Floquet spin chains, Communications Physics 5, 10.1038 (2022).
- [40] A. A. Nizami and A. W. Shrestha, Krylov construction and complexity for driven quantum systems, arXiv:2305.00256
- [41] P. Suchsland, R. Moessner, and P. W. Claeys, Krylov complexity and Trotter transitions in unitary circuit dynamics, arXiv:2308.03851
- [42] H.-C. Yeh and A. Mitra A Universal Model of Floquet Operator Krylov Space, arXiv:2311.15116
- [43] C. Liu, H. Tang, and H. Zhai, Krylov complexity in open quantum systems, Phys. Rev. Res. 5, 033085 (2023)
- [44] A. Bhattacharya, P. Nandy, P. P. Nath, and H. Sahua, On Krylov complexity in open systems: an approach via bi-Lanczos algorithm, arXiv:2303.04175
- [45] N. S. Srivatsa and C. von Keyserlingk, The operator growth hypothesis in open quantum systems, arXiv:2310.15376
- [46] A. Eckardt and E. Anisimovas, High-frequency approximation for periodically driven quantum systems from a Floquet-space perspective, New J. Phys. 17, 093039 (2015)
- [47] A. Kitaev, Anyons in an exactly solved model and beyond, Ann. Phys. (Amsterdam) 321, 2 (2006).
- [48] R. Bianco and R. Resta, Mapping topological order incoordinate space, Phys. Rev. B 84, 241106(R) (2011).
- [49] F. Song, S. Yao, and Z. Wang, Non-Hermitian Topological Invariants in Real Space, Phys. Rev. Lett. 123, 246801 (2019)