

Asymptotically safe – canonical quantum gravity junction

T. Thiemann^{1*}

¹ Inst. for Quantum Gravity, FAU Erlangen – Nürnberg,
Staudtstr. 7, 91058 Erlangen, Germany

April 30, 2024

Abstract

The canonical (CQG) and asymptotically safe (ASQG) approach to quantum gravity share to be both non-perturbative programmes. However, apart from that they seem to differ in several aspects such as: 1. Signature: CQG is Lorentzian while ASQG is mostly Euclidian. 2. Background Independence (BI): CQG is manifestly BI while ASQG is apparently not. 3. Truncations: CQG is apparently free of truncations while ASQG makes heavy use of them.

The purpose of the present work is to either overcome actual differences or to explain why apparent differences are actually absent. Thereby we intend to enhance the contact and communication between the two communities. The focus of this contribution is on conceptual issues rather than deep technical details such as high order truncations. On the other hand the paper tries to be self-contained in order to be useful to researchers from both communities.

The point of contact is the path integral formulation of Lorentzian CQG in its reduced phase space formulation which yields the formal generating functional of physical (i.e. gauge invariant) either Schwinger or Feynman N-point functions for (relational) observables. The corresponding effective actions of these generating functionals can then be subjected to the ASQG Wetterich type flow equations which serve in particular to find the rigorous generating functionals via the inverse Legendre transform of the fixed pointed effective action.

1 Introduction

It is widely believed that only a non-perturbative approach can yield a predictive theory of quantum gravity (QG). The asymptotic safety approach (ASQG) [1] and the canonical approach (CQG) [2], the most modern incarnation of which is Loop Quantum Gravity (LQG) [3], are two such programmes. These have been developed quite independently of each other and it is therefore an interesting challenge to try to bring them into contact in order to investigate their differences and their hopefully complementary strengths and weaknesses.

At first sight, this seems to be an easy task: Both theories start from a Lagrangian which can then be subjected to either canonical (CQG) or path integral (ASQG) quantisation. The connection between those types of quantisations is that the path integral formulation allows to compute correlation functions of field operator valued distributions of the canonical formulation. This relation is quite well understood, both conceptually and technically, for quantum field theories (QFT) in Minkowski space without gauge symmetries. On more general backgrounds with or without gauge symmetries a rigorous implementation of this relation is more challenging but at a formal level one can still derive one formulation from the other via the reduced phase space path integral [4].

However, apart from these mostly technical challenges, there appear to be unsurmountable differences between the two programmes:

1. Signature:

Apart from few exceptions [5], ASQG is a path integral formulation for Euclidian signature metrics while CQG in its LQG incarnation is explicitly for Lorentzian signature [6]. One cannot simply pass between the signatures using analytic continuation in time as a quantum metric that one integrates over has quite arbitrary (generically

*thomas.thiemann@gravity.fau.de

non-analytic) time dependence.

2. Background dependence (BI):

ASQG makes heavy use of background metric dependent constructions such as the very definition of the averaging kernel that enters the Wetterich equation. Sometimes this background is kept arbitrary [7], sometimes it is chosen to be maximally symmetric [8] and it is at least confusing to the newcomer in what sense the predictions of ASQG can be interpreted background independently. CQG on the other hand is manifestly background independent by construction.

3. Truncations:

The Wetterich equation of ASQG is an exact and non-perturbative renormalisation flow equation for the so-called effective average action (EAA), however, to actually solve this equation one must restrict that action to a finite number of terms and truncate the Wetterich equations to those terms. Such kind of truncations are absent in LQG.

The purpose of this paper is two-fold:

A.
On the one hand we want to understand to what extent one can derive ASQG from CQG using the reduced phase space path integral. This should then ease the comparison between the two formulations.

B.
On the other hand, such a derivation will shed light in what sense the above three differences between the two programmes can be overcome.

The architecture of this contribution is as follows:

To be useful for both communities, in section 2 we will review the canonical quantisation programme as well as the reduced phase space path integral and in section 3 we will review the main ingredients of the asymptotic safety programme. Special attention will be paid on how to implement manifest gauge invariance in both programmes and how these can be translated into each other. This ensures that the correlation functions that one eventually computes are for (relational) Dirac observables only.

While the review material laid out in sections 2, 3 is quite general, in section 4 we pick a concrete class of theories within the Einstein-Scalar sector. While the matter content of that theory is not very physical, it serves the purpose to showcase for a model that is technically not too involved, that a connection between the two programmes can indeed be established. We develop and prepare that model up to the point at which an ASQG treatment is immediately possible. The ASQG treatment itself is carried out in the companion paper [9].

In section 5 we summarise and conclude.

In the appendices we collect some background material:

In appendix A we establish the non-trivial relation between effective actions whose preimages under Legendre transformation are related by restriction. In appendix B we show how to get rid of square root actions in the path integral by a method which is inspired by Polyakov's trick to switch from the Nambu-Goto to the Polyakov action for the closed bosonic string [11]. In appendix C we report a side result of our analysis, namely that the so-called optimised cut-off, which are much used in the Euclidian version of ASQG, is not in the image of the Laplace transform even of a positive measure and thus raises the question whether it is in the image of the Laplace transform of some well defined mathematical object at all. Likewise some smoother cut-offs used in Euclidian ASQG are not in the Laplace transform image of functions but rather distributions so that the heat kernel time integrals that are required to compute the flow are ill-defined as they stand. This kind of analysis leads to appendix D where we introduce and explore a new class of cut-off functions both for Euclidian and Lorentzian signature which safely are in the image of the Laplace and Fourier transform respectively of a Schwartz function. In the Lorentzian regime the flow defined by this cut-off is necessarily in *complex coupling constant space* but among the possible trajectories running into the fixed point we only allow *admissible ones* which have real valued, finite $k = 0$ limit for the physical dimensionful couplings.

As far as the obstructions I.-III. are concerned, we learn the following:

I. Signature

One can formulate the reduced phase space path integral either for Schwinger N-point functions or for

Feynman time ordered N-point functions. This is possible because in the reduced phase space formulation one has access to a Hamiltonian operator H and Wick rotation of e^{itH} with respect to t has a well-defined meaning. However, the physical Hamiltonian H is not of standard Schrödinger type, it typically involves a square root, so that integrating out the momenta is not immediately possible in exact form. The square root Hamiltonian is generic for generally covariant theories and arises already for the relativistic particle [10]. By the method provided in appendix B one can get rid of the square root and in that case, one can indeed integrate out the momenta and implement Euclidian signature ASQG. The resulting theory is however spatially very non-local. One can implement a more local version. Another method to get rid of the square root and to avoid non locality is to unfold the reduced phase space path integral to the unreduced one but then one must choose the Lorentzian route. One is then able to integrate out the momenta exactly and relate the path integral to the Einstein-Hilbert action for Lorentzian signature. At this point one may think that comparing the programmes is no longer possible. However, as noted e.g. in [5], it is also possible to give a Lorentzian version of the Wetterich equation for the EAA although its interpretation is less clear.

II. Background Independence

The derivation of the reduced phase space path integral results in the Einstein-Hilbert action with metric dependent cosmological constant term and a non-standard gauge fixing structure that arises both from integrating out the momenta and the gauge fixing conditions that we imposed. If the corresponding generating functional was well-defined one could compute all (connected) Feynman N-point functions from it. However, it is not and thus one may use the proposal of the ASQG programme to modify the action by quadratic term that involves a so-called averaging kernel which depends on a parameter $k \in \mathbb{R}$ and a background metric \bar{g} of Lorentzian signature and k^2 can locally be related to the norm of a vector k with respect to \bar{g} . The averaging kernel vanishes when $k = 0$.

One may now pass from the modified generating functional to the EAA by Legendre transform which obeys a Wetterich like exact equation. From the viewpoint of CQG, the Wetterich equation serves the sole purpose to select a fixed point EAA (solution of the Wetterich in terms of the dimensionless couplings with finite limit as $k \rightarrow \infty$ fixing all but a finite number of those) which also has a limit as $k \rightarrow 0$ with respect to the running dimensionful couplings which then serves as the rigorous definition of the ordinary effective action of the gauge invariant theory that we are interested in. Note that in the usual ASQG flow equation one considers the flow with respect to all degrees of freedom while actual effective action is restricted the gauge invariant degrees of freedom. This restriction requires an additional step after the solution of the Wetterich equation has been obtained.

Now the reason why the end result of this analysis is background independent is due to the use of the background method [12] that one employs: We keep the background arbitrary and leave it unspecified aside from assuming that it has Lorentzian signature. This not only allows to unambiguously monitor the flow of the couplings that enter the EAA, it also leads to an effective action as follows: The Wetterich equation yields a functional $\bar{\Gamma}_k(\hat{g}, \bar{g})$ where \hat{g} is the field that arises by Legendre transforming the generating functional $\bar{C}_k(f, \bar{g})$ of connected functions with respect to the current f using the background field method. If $\bar{\Gamma}_0(f, \bar{g})$ exists then $\Gamma(g) := \Gamma_0(0, g)$ is the standard effective action and its Legendre transform defines $C(g)$. Thus, the background method applied in this way is in fact background independent! This is of course well known in field theory and in fact it was DeWitt, one of the founding fathers of CQG, who co-invented this method.

• III. Truncations

In order to solve the Wetterich equation, we make an Ansatz for $\bar{\Gamma}_k$ that involves a small set of k dependent coupling constants. The left hand side of the Wetterich equation only depends on the derivatives of those coupling constants with respect to k . To match these terms to those that arise on the right hand side of the Wetterich equation one can make use of the Lorentzian version of the heat kernel method [13] which is based on DeWitt's seminal work on the Synge world function [14] and which plays a fundamental role in microlocal analysis [15] and QFT in curved spacetime (CST) [13]. We may call this the "Schrödinger kernel" method. The properties used for the Synge world function are signature independent and can be used for our purposes as well. While in ASQG one often matches the gauge fixing in order to avoid certain

“off-diagonal” terms (basically polynomials of covariant background derivatives that are not polynomials of the background d’Alambertian) the “heat” kernel method can be extended to off-diagonal situations [16] and one does not even need to perform a transverse-traceless decomposition by suitable organisation of the truncation scheme (e.g. by orders of derivatives rather than orders of curvature polynomials). One is then left with performing integrals with respect to the “heat” kernel parameter which in Lorentzian signature are more difficult to perform than in Euclidian signature.

After all of this has been done, one must discard on the right hand side all terms that have no counter part in the Ansatz (truncation), then perform the fixed point analysis and finally take $k \rightarrow 0$ (if possible within the limits of the chosen truncation). In principle one can make the system of equations close iteratively by refining the Ansatz step by step by the terms that are produced by the Wetterich equation not already contained in the Ansatz.

One may argue that in CQG this truncation is not necessary. However, the physical Hamiltonian in LQG suffers from many quantisation ambiguities which must be fixed using renormalisation methods [17], e.g. Hamiltonian renormalisation [18]. In any renormalisation scheme one usually is forced to perform some form of approximation (here: truncations) due to practical limitations (in case of ASQG it is the limits in complexity that symbolic computation programmes are able to handle in reasonable time). Thus renormalisation is present in both approaches and truncations are in principle not necessary except for reasons of practicability. What is missing, to the best of our knowledge in every approach, is an error control analysis, i.e. a mathematical proof that a certain truncation does not miss out a relevant coupling.

2 Relational observables, reduced phase space, canonical quantisation, path integral formulation

In this section we review elements of canonical quantisation of constrained systems. CQG practitioners can safely skip this section.

2.1 Classical phase space of constrained systems

Consider a phase space coordinatised by two sets of canonical pairs $z = ((p_a, q^a), (y_j, x^j))$. We define the coordinate functions $Z = (P_a, Q^a, Y_j, X^j)$ on the phase space by $P_a(z) = p_a$ etc. The non vanishing Poisson brackets are

$$\{P_a, Q^b\} = \delta_a^b, \{Y_j, X^k\} = \delta_j^k \quad (2.1)$$

The labels $a, b, c, \dots = 1, 2, \dots$ and $j, k, l, \dots = -1, 2, \dots$ can be taken from a discrete index set. In classical mechanics these sets are finite, in classical field theory they are countably infinite. In the latter case we think of them as obtained as $P_a = \langle e_a, P \rangle$, $Q^a = \langle f^a, Q \rangle$, $Y_j = \langle b_j, Y \rangle$, $X^j = \langle c^j, X \rangle$ where $\langle \dots \rangle$ denotes the inner product of the appropriate 1-particle Hilbert space including summation over gauge group representation labels (tensor representations of the diffeomorphism group, irreducible representations of Yang-Mills type gauge groups etc.) which we suppress, P, Q, Y, X are the continuum fields and (e_a, f^a) , (b_j, c^j) are pairs of bases and co-bases respectively of the one particle Hilbert spaces, i.e. $\langle e_a, f^b \rangle = \delta_a^b$, $\langle b_j, c^k \rangle = \delta_j^k$.

These labels are adapted to a system of first class constraints

$$\{C_j, C_k\} = f_{jk}{}^l C_l \quad (2.2)$$

where the structure functions $f_{jk}{}^l$ need not be constant on the phase space which is indeed the case for canonical GR. Again in field theory the constraints $C_j = \langle e_j, C \rangle$ are smeared versions of the continuum constraints C . There may also be a true Hamiltonian H already (not the case for generally covariant systems such as GR) which is gauge invariant i.e.

$$\{C_j, H\} = f_{j0}{}^k C_k \quad (2.3)$$

in which case we expand the phase space by an additional canonical pair (y_0, x^0) which has vanishing Poisson brackets with all others and we add an additional constraint

$$C_0 = y_0 + H \Rightarrow \{C_j, C_0\} = f_{j0}{}^k C_k \quad (2.4)$$

The augmented set of constraints is thus again first class so that w.l.g. we may restrict to the case without true Hamiltonian.

The construction of the reduced phase space can now be described in two equivalent ways, either using so called relational Dirac observables which are gauge invariant or by using gauge fixing. Both descriptions take the following steps: First we solve the constraints $C_j = 0$ for the momenta y_j to arrive at equivalent sets of constraints

$$\hat{C}_j = y_j + h_j(p, q; x) \quad (2.5)$$

In general this involves picking a square root because typically the constraints depend quadratically on the momenta (recall the case of the relativistic particle). Also the dependence of the functions h_j on x is in general non-trivial, if it is trivial one says that the system is deparametrised. This happens only in very fortunate cases. In field theory to write the constraints as (2.5) often is not possible algebraically but one must also solve PDE's or equivalently invert non-diagonal infinite dimensional matrices which is technically challenging. One will therefore select the split of the canonical coordinates into the two sets in such a way as to avoid these complications as much as possible.

The virtue of the equivalent set of constraints (2.5) is that they are Abelian, i.e. they have vanishing Poisson brackets among each other. To see this we note that $\{\hat{C}_j, \hat{C}_j\}$ must vanish when the constraints hold since $\hat{C}_j = M_j^k C_k$ for some matrix M . On the other hand, $\{\hat{C}_j, \hat{C}_j\}$ does not depend on y_j , it may therefore be evaluated at any y_j and in particular at $y_j = -h_j$. Thus $\{\hat{C}_j, \hat{C}_j\}$ vanishes identically, not only when the constraints hold.

2.2 Relational Dirac observables

This observation is the mechanism behind relational observables [19]. Consider "coordinate functions" $k^j : \mathbb{R} \rightarrow \mathbb{R}; t \mapsto c^j(t)$ which are constants on the phase space and a one parameter set of gauge fixing conditions

$$G^j(t) := X^j - k^j(t) \quad (2.6)$$

Then for any function F on the phase space the quantity

$$O_F(t) := [\exp(s^j V_j \cdot) F]_{s=-G(t)} \quad (2.7)$$

is gauge invariant, i.e. it has vanishing Poisson brackets with the constraints \hat{C}_j . Here $V_j \cdot F := \{\hat{C}_j, F\}$ is the Hamiltonian vector field of \hat{C}_j and in (2.6) one first computes the Hamiltonian flow for s a constant function on the phase space and after that evaluates at $s = -G(t)$. To verify that

$$\{\hat{C}_j, O_F(t)\} = V_j \cdot O_F(t) = 0 \quad (2.8)$$

it is enough to use the Taylor expansion of (2.7), that $V_j \cdot G^k = \delta_j^k$ and that $[V_j, V_k] = X_{\{\hat{C}_j, \hat{C}_k\}} = 0$.

The observables $O_F(t)$ satisfy a number of interesting properties which are not difficult to prove [4, 19]. We mention two of them

$$\{O_{F_1}(t), O_{F_2}(t)\} = O_{\{F_1, F_2\}^*}(t), \quad O_{F_1 \circ F_2}(t) = F_1(O_{F_2}(t)) \quad (2.9)$$

where

$$\{F_1, F_2\}^* = \{F_1, F_2\} + \{F_1, \hat{C}_j\} \{G^j(t), F_2\} - \{F_2, \hat{C}_j\} \{G^j(t), F_1\} \quad (2.10)$$

is the Dirac bracket defined by $\hat{C}_j, G^j(t)$ considered as a second class set of constraints. The second relation in (2.9) implies that

$$[O_F(t)](z) = F([O_Z(t)](z)) \quad (2.11)$$

i.e. to compute $O_F(t)$ we just need to compute the observables corresponding to the coordinate functions $Z(z)$. We easily see $O_{X^j}(t) = k^j(t)$ is just a constant function on the phase space and thus trivially has vanishing Poisson brackets with the constraints. Next, since \hat{C}_j is already gauge invariant

$$O_{\hat{C}_j}(t) = \hat{C}_j = O_{Y_j}(t) + h_j(O_P(t), O_Q(t); k(t)) \quad (2.12)$$

Thus on the constraint surface $O_{Y_j}(t)$ can be expressed in terms of the $O_{P_a}(t), O_{Q^a}(t)$ which are therefore, at any time, the coordinates of the gauge invariant reduced phase space. Since $\{P_a, G^j(t)\} = \{Q^a, G^j(t)\} = 0$ from the first relation in (2.10)

$$\{O_{P_a}(t), O_{Q^b}(t)\} = O_{\{P_a, Q^b\}}(t) = \delta_a^b \quad (2.13)$$

thus the observables $O_{P_a}(t), O_{Q^a}(t)$ are canonically conjugate at all times t . This is important: If (2.13) would be a complicated expression, it would be hard to find representations of the canonical commutation relations in the quantum theory.

Let us focus on a function F which just depends on p, q . Then

$$\frac{d}{dt} O_F(t) = O_{\dot{k}^j(t)\{\hat{C}_j, F\}} = O_{\dot{k}^j(t)\{h_j, F\}} = O_{\dot{k}^j(t)\{h_j, F\}^*} = \{O_{\dot{k}^j(t) h_j}(t), O_F(t)\} \quad (2.14)$$

where we used $\{F, G^j(t)\} = \{h_k, G^j(t)\} = 0$ in the step before the last and then (2.9). It follows that the gauge invariant, physical Hamiltonian that drives the one parameter time evolution of the those observables O_F is given by

$$H(t) = \dot{k}^j(t) h_j(O_P(0), O_Q(0); k(t)), \{H(t), O_F\}([O_P(t), O_Q(t)]) = \frac{d}{dt} O_F(t) \quad (2.15)$$

It is explicitly time dependent unless \dot{k}^j is a constant and if the dependence on $k(t)$ in h_j vanishes (for example if the dependence on x of h_j is only through spatial derivatives). Of course a time independent physical Hamiltonian is much preferred (conservative system).

2.3 Gauge fixing

The gauge fixing condition (2.6) has already appeared in the relational approach but there it served to extend functions $F(q, p)$ off the gauge cut $G^j = 0$ of the constraint surface $C_j = 0$ to the full phase space. In the gauge fixing approach we consider the reduced phase space as coordinatised by the so called ‘‘true degress of freedom’’ P, Q rather than $O_F(0), O_P(0)$. The ‘‘gauge degrees of freedom’’ Y, X are discarded from this description altogether. This gives an equivalent description as follows: First we note that the canonical brackets among the P, Q and the $O_P(0), O_Q(0)$ are identical. Next consider a gauge transformation generated by the constrained Hamiltonian $\hat{C}(s) = s^j \hat{C}_j$. There is a residual 1-parameter set family of gauge transformations allowed that stabilise the gauge condition $G^j(t) = 0$

$$\frac{d}{dt} G^j(t) = \partial_t G_j(t) + \{\hat{C}(s), G^j(t)\} = -\dot{k}^j(t) + s^j = 0 \quad (2.16)$$

which fixes the Lagrange multiplier s^j . We note the fixed values

$$s_*^j = \dot{k}^j(t) \ 1, \ X_*^j = k^j(t) \ 1, \ Y_j^* = -h_j(Q, P; k(t)) \quad (2.17)$$

Let F be a function of the true degrees of freedom q, p only. Then the reduced Hamiltonian $H_*(t) = H_*(q, p; t)$ is defined by

$$\{H_*(t), F\}(q, p) := (\{\hat{C}(s), F\}_{s=s_*, x=x_*, y=y_*})(q, p) = \dot{k}^j(t) \{h_j(\cdot, \cdot; c(t)), F\}(q, p) \quad (2.18)$$

whence

$$H_*(q, p; t) = \dot{k}^j(t) h_j(q, p; k(t)) \quad (2.19)$$

Comparing (2.19) and (2.19) we see that the reduced Hamiltonian is the same function of the true degrees of freedom as is the physical Hamiltonian of the relational Dirac degrees of freedom. The two descriptions are truly isomorphic. However one should keep in mind that the physical interpretation of the true degrees of freedom and the reduced Hamiltonian depends on the choice of gauge fixing: Q, P are those observables which on the gauge cut defined by $G^j(0) = 0$ coincide with the gauge invariant observables $O_P(0), O_Q(0)$ and their evolution is generated by the reduced Hamiltonian induced by $G^j(t) = 0$.

2.4 Reduced phase space quantisation: Hamiltonian formulation

Since the gauge invariant content of the theory can be described in terms of the true degrees of freedom q, p and the reduced Hamiltonian $H_*(t)$ we consider canonical and path integral quantisation of this description. We restrict to the case of a reduced Hamiltonian which is not explicitly time dependent. This will be sufficient for our application in section 4. An extension to the explicitly time dependent case is possible but more complicated.

Canonical quantisation involves finding a representation of the canonical commutation relations and $*$ -relations (we set $\hbar = 1$ and just consider bosonic variables for simplicity)

$$[P_a, Q^b] = i \delta_a^b 1, \quad P_a^* = P^a, (Q^a)^* = Q^a \quad (2.20)$$

The algebra generated by the generators $P_a, Q^a, 1$ is called the Heisenberg algebra and has the disadvantage that representations ρ of P_a, Q^a are unbounded operators. Thus one passes to the Weyl elements

$$W(f, g) := e^{i [f_a Q^a + g^a P_a]} \quad (2.21)$$

which obey the corresponding Weyl relations induced by (2.20) [19]. The Weyl elements generated the corresponding Weyl algebra \mathfrak{A} which is a $*$ -algebra (even a C^* -algebra).

We consider cyclic representations of \mathfrak{A} . These are in 1-1 correspondence with states ω on \mathfrak{A} i.e. positive, normalised, linear functionals on \mathfrak{A} : $\omega(a^*a) \geq 0$, $\omega(1) = 1$, $\omega(za + z'a') = z\omega(a) + z'\omega(a')$; $z, z' \in \mathbb{C}$; $a, a' \in \mathfrak{A}$. Moreover ω gives rise to GNS data $(\mathcal{H}, \Omega, \rho)$ where \mathcal{H} is a Hilbert space, Ω a unit vector in \mathcal{H} and ρ a $*$ -representation of \mathfrak{A} as bounded operators on \mathcal{H} . The vector Ω is cyclic for $\rho(\mathfrak{A})$ i.e. $\mathcal{D} := \rho(\mathfrak{A})\Omega$ is a dense, common, invariant domain for the operators $\rho(a)$. Note the representation properties $\rho(a)\rho(b) = \rho(ab)$, $\rho(a) + \rho(b) = \rho(a + b)$, $\rho(a)^\dagger = \rho(a^*)$. The correspondence between ω and the GNS data is via the formula

$$\omega(a) = \langle \Omega, \rho(a)\Omega \rangle_{\mathcal{H}} \quad (2.22)$$

i.e. correlation functions among the operators $\rho(a)$ with respect to the cyclic vector Ω are computed by evaluating ω on them. One shows that the GNS data are determined by ω up to unitary transformations between representations, see e.g. [20] for all the details.

Cyclic representations of, or equivalently states on, \mathfrak{A} are not difficult to find, any Fock representation is of that form. The real challenge is to find states ω that support a quantum version of the reduced Hamiltonian H_* . We will drop the subscript $*$ in what follows in order not to get confused with the $*$ -operation and thus just denote it by H . It is here where all the difficulties of interacting QFT enter and Haag's theorem [21] tells us that such representations must be carefully adapted to H if they exist at all. On the other hand, this physical selection criterion on ω is welcome as otherwise we would be confronted with the question which unitary equivalence class of representation to choose (recall that spectra, i.e. measurement values, of operators depend on the unitary equivalence class).

The construction of a quantum version of H faces many issues such as: 1. operator ordering choices, 2. ultra-violet (UV) singularities (summation over an infinite number of mode labels a) and 3. infra-red (IR) singularities (working in non-compact spatial manifolds). It is a common theme in constructive QFT [22] to tame these issues by imposing a mode (UR) cut-off $a < M$, an IR cut-off (compactification of the spatial manifold e.g. on a torus of size R with periodic boundary conditions) and to pick some truncation and ordering H_M of H with respect to the P_a, Q^a , $a \leq M$ such that H_M is a self-adjoint operator on \mathcal{H}_M where $(\mathcal{H}_M, \Omega_M, \rho_M)$ are the GNS data of a state ω_M on \mathfrak{A}_M where \mathfrak{A}_M is the Weyl algebra generated by the Weyl elements (2.21) subject to the condition $f_a = 0$, $a > M$, $g^a = 0$, $a > M$. As we are now in a situation with a finite number of degrees of freedom the problem of finding suitable ω_M is trivial as the von Neumann theorem [20] says that there is only one unitary equivalence class available if one asks that the representation is regular (i.e. we can differentiate the Weyl elements to obtain self-adjoint operators $\rho_M(P_a), \rho_M(Q^a)$, $a \leq M$ by Stone's theorem). The remaining task, at fixed M , is then to find a suitable operator $\rho_M(H_M)$. We assume that this task can be performed at any finite M, R . We will abuse notation and write H_M for $\rho_M(H_M)$ and $W(f, g)$ for $\rho_M(W(f, g))$ in what follows in order not to clutter the formulae.

We now have constructed a family of theories labelled by M, R . They can be interpreted as a compactified version of the theory that we actually want at finite resolution M . The idea of renormalisation is to impose

that all those theories labelled by M at fixed R descend from a common theory at the same fixed R by coarse graining, i.e. by restricting observables to mode labels $a \leq M$. In other words we want that

$$\omega(a) = \omega_M(a) \quad \forall a \in \mathfrak{A}_M, \quad P_M H P_M = H_M \quad (2.23)$$

where $P_M : \mathcal{H} \rightarrow \mathcal{H}_M$ is the orthogonal projection. These conditions are not automatically met using some choice of ω_M, H_M that we pick at any given M . The idea of Hamiltonian renormalisation [17] is now to define a renormalisation flow on the family of theories (ω_M, H_M) starting with some initial family $(\omega_M^{(0)}, H_M^{(0)})$. For this one picks coarse graining maps $J_{MM'}^{(n)}$, $M < M'$, $n \in \mathbb{N}_0$ with $J_{MM'}^{(n)} : \mathcal{H}_M^{(n+1)} \rightarrow \mathcal{H}_{M'}^{(n)}$ and picks the states $\omega_M^{(n+1)}$ such that these are isometric injections. Then one defines $H_M^{(n+1)} := J_{MM'}^\dagger H_{M'}^{(n)} J_{MM'}^{(n)}$. This defines a flow of families $n \mapsto (\omega_M^{(n)}, H_M^{(n)})$ a fixed point of which then accomplishes the task (2.23). Namely a fixed point (ω, H) of this flow results in an inductive limit Hilbert space [20] \mathcal{H} with isometric injections $J_M : \mathcal{H}_M \rightarrow \mathcal{H}$ and where $P_M = J_M J_M^\dagger$ is the corresponding projection together with a quadratic form H on \mathcal{H} such that $H_M = J_M^\dagger H J_M$. If the quadratic form is closable and is bounded from below, we do get a self-adjoint operator via the Friedrichs extension [23].

The details of the flow depend on the choice of the coarse graining maps which are typically of the form

$$J_{MM'}^{(n)} W(f, g) \Omega_M^{(n+1)} = W(I_{MM'}(f, g)) \Omega_{M'}^{(n)} \quad (2.24)$$

where $I_{MM'}$ is a coarse graining map on the smearing functions which maps those at coarse resolution M into those at finer resolution M' . In order that the inductive limit construction works one must pick those $I_{MM'}$ such that $I_{M_2 M_3} \circ I_{M_1 M_2} = I_{M_1 M_3}$ ("cylindrical consistency") for $M_1 < M_2 < M_3$ and such that the label set \mathcal{M} of cut-offs is partially ordered and directed. This can be ensured for example using multi resolution analysis familiar from wavelet theory (see e.g. [24] and references therein).

If one has managed to remove the cut-off M in this manner, one still must take the thermodynamic limit $R \rightarrow \infty$ which meets new technical challenges and can give rise to different "phases" of the theory [20].

Summary:

The reduced phase space quantisation can be carried out entirely within the Hamiltonian or canonical framework but in QFT renormalisation is typically required. Since the flow equations can typically not be solved in closed form, some approximation scheme is required. This is the canonical or Hamiltonian equivalent of truncations performed in ASQG.

2.5 Reduced phase space quantisation: Path integral formulation

We now turn to a path integral framework. We start again from some ω_M, H_M with GNS data $(\mathcal{H}_M, \Omega_M, \rho_M)$. The cyclic vector Ω_M is generically not an eigenstate of H_M . Suppose that H_M is bounded from below (w.l.g. by zero) and has a unique ground state $\Omega_{0,M}$ i.e. $H_M \Omega_{0,M} = 0$. For sufficiently complicated H_M this vector will not be computable in closed form. Let $U_M(t) = \exp(-itH_M)$ $t \in \mathbb{R}$; $C_M(s) = \exp(-sH_M)$, $s \geq 0$ be the unitary group and contraction semi-group generated by H_M respectively and consider the following normalised vectors

$$\Omega_{\pm, M}^T := U_M(\pm T) \Omega_M, \quad \Omega_M^T := \frac{C_M(T) \Omega_M}{\|C_M(T) \Omega_M\|} \quad (2.25)$$

If the strong limits of (2.25) exist, denoted by Ω_M^\pm, Ω_M^C respectively, then these obey $U_M(t) \Omega_M^\pm = \Omega_M^\pm$, $C_M(s) \Omega_M^C = \Omega_M^C$, i.e. they are ground states of H_M . By the assumed uniqueness

$$\Omega_M^\pm = e^{i\alpha_\pm} \Omega_{0,M}, \quad \Omega_M^C = e^{i\alpha_C} \Omega_{0,M} \quad (2.26)$$

for some phases α_\pm, α_C . We have

$$\lim_T \langle U_M(T) \Omega_M, U_M(-T) \Omega_M \rangle = e^{-i[\alpha_+ - \alpha_-]}, \quad \lim_T \|C_M(t) \Omega_M\|^2 = \lim_T \langle \Omega_M, C_M(-2T) \Omega_M \rangle \quad (2.27)$$

whence

$$\begin{aligned} \langle \Omega_{0,M}, \cdot \Omega_{0,M} \rangle &= \lim_{T \rightarrow \infty} \langle U_M(T) \Omega_M, \cdot U_M(-T) \Omega_M \rangle e^{i[\alpha_+ - \alpha_-]} = \lim_{T \rightarrow \infty} \frac{\langle U_M(T) \Omega_M, \cdot U_M(-T) \Omega_M \rangle}{\langle \Omega_M, \cdot U_M(-2T) \Omega_M \rangle} \\ \langle \Omega_{0,M}, \cdot \Omega_{0,M} \rangle &= \lim_T \frac{\langle C_M(T) \Omega_M, \cdot C_M(T) \Omega_M \rangle}{\langle \Omega_M, \cdot C_M(2T) \Omega_M \rangle} \end{aligned} \quad (2.28)$$

2.5.1 Lorentzian formulation

In the Lorentzian formulation we are interested in the time ordered correlation functions (Feynman N-point functions)

$$F_{N,M}((t_1, f^1), \dots, (t_N, f^N)) := \langle \Omega_{0,M}, \mathcal{T}[W_{t_N}(f^N, 0) \dots W_{t_1}(f^1, 0)] \Omega_{0,M} \rangle_{\mathcal{H}_M} \quad (2.29)$$

where $W_t(f, g) := U_M(t)W(f, g)U_M(t)^{-1}$ and f_a, g^a vanish for $a > M$. The time ordering symbol \mathcal{T} orders the time dependence from left to right with largest time to the outmost left. Upon derivation by $f_{a_k}^k$, $k = 1, \dots, N; a \leq M$ at $f^1 = \dots = f^N = 0$ we obtain

$$F_{N,M}((t_1, a_1), \dots, (t_N, a_N)) := \langle \Omega_{0,M}, \mathcal{T}[Q_M^{a_N}(t_N) \dots Q_M^{a_1}(t_1)] \Omega_{0,M} \rangle_{\mathcal{H}_M} \quad (2.30)$$

with $Q_M^a(t) := U_M(t)Q^a U_M(t)^{-1}$, $a \leq M$. This can also be written as

$$F_{N,M}((t_1, a_1), \dots, (t_N, a_N)) = i^{-N} \left[\frac{\delta^N}{\delta f_{a_N}(t_N) \dots \delta f_{a_1}(t_1)} \chi_M(f) \right]_{f=0}, \quad (2.31)$$

with the generating functional

$$\chi_M(f) := \langle \Omega_{0,M}, \mathcal{T}[e^{i \int_{\mathbb{R}} dt f_a(t) Q_M^a(t)}] \Omega_{0,M} \rangle_{\mathcal{H}_M} \quad (2.32)$$

The way in which (2.32) is written is not practically useful as we have not explicit access to $\Omega_{0,M}$ but rather to Ω_M . Suppose that f has compact time support in $[-\tau, \tau]$ then we have explicitly using (2.28)

$$\begin{aligned} \chi_M(f) &= \lim_{T \rightarrow \infty} \frac{\langle \Omega_M, e^{iTH_M} \mathcal{T}[e^{i \int_{-\tau}^{\tau} dt f_a(t) Q_M^a(t)}] e^{iTH_M} \Omega_M \rangle}{\langle \Omega_M, e^{2iTH_M} \Omega_M \rangle} \\ &= \lim_{T \rightarrow \infty} \lim_{N \rightarrow \infty} \frac{\langle \Omega_M, e^{i(T-t_{N-1})H_M} e^{i\Delta_N f_a(t_{N-1})Q^a} e^{i\Delta_N H_M} \dots e^{i\Delta_N H_M} e^{i\Delta_N f_a(t_N)Q^a} e^{i(t_N - (-T))H_M} \Omega_M \rangle}{\langle \Omega_M, e^{2iTH_M} \Omega_M \rangle} \end{aligned} \quad (2.33)$$

where $\Delta_N = \frac{\tau}{N}$, $t_k = k\Delta_N$, $k = -N, \dots, N-1$. For sufficiently large T we have $T \geq \tau$ and by extending f by zero to $\mathbb{R} - [\tau, -\tau]$ we can identify $\tau = T$. Then

$$\begin{aligned} \chi_M(f) &= \lim_{T \rightarrow \infty} \lim_{N \rightarrow \infty} \frac{Z_{M,T,N}(f)}{Z_{M,T,N}(0)}, \\ Z_{M,T,N}(f) &= \langle \Omega_M, e^{i\Delta_N H_M} e^{i\Delta_N f_a(t_{N-1})Q^a} e^{i\Delta_N H_M} \dots e^{i\Delta_N H_M} e^{i\Delta_N f_a(t_N)Q^a} e^{i\Delta_N H_M} \Omega_M \rangle \end{aligned} \quad (2.34)$$

We consider the integral kernel of the unitary propagator

$$\int [dq']_M U_{M,T,N}(q, q') \psi_M(q') := (e^{i\Delta_N H_M} \psi_M)(q) \quad (2.35)$$

where $[dq]_M$ is the Lebesgue measure of the q^a , $a \leq M$. As at finite M the representation is in the unitarity class of the Schrödinger representation we consider Q^a as multiplication by q^a and P_a as $i\partial/\partial q^a$. Then one has to argue that for $N \rightarrow \infty$ at finite T

$$\begin{aligned} U_{M,T,N}(q, q') &= \int [d\hat{q}]_M U_{M,T,N}(q, \hat{q}) \delta_{q'}(\hat{q}) = [e^{i\Delta_N H_M(Q,P)} \delta_{q'}](q) = \int [d\frac{p}{2\pi}]_M e^{i\Delta_N H_M(q,p)} e^{-i \sum_{a \leq M} p_a (q - q')^a} \\ &= \int [d\frac{p}{2\pi}]_M e^{i\Delta_N H_M(q,p)} e^{-i \sum_{a \leq M} p_a (q - q')^a} \end{aligned} \quad (2.36)$$

Without going into the details of H_M this step is hard to justify, for Schrödinger type of operators one may use Feynman-Kac arguments [23].

Using this heuristics we obtain

$$Z_{M,T,N}(f) = \prod_{k=-N}^N [dq_k]_M \prod_{l=-N}^{N-1} \left[\frac{dp^l}{2\pi} \right]_M \overline{\Omega_M(q_N)} \Omega_M(q_{-N}) e^{-i \sum_{k=-N}^{N-1} \Delta_N [\sum_{a \leq M} p_a^k \frac{q_{k+1}^a - q_k^a}{\Delta_N} - H(p^k, q_k)]} \times e^{i \sum_{k=-N}^{N-1} \sum_{a \leq M} f_a(t_k) q_k^a} \quad (2.37)$$

In what follows we mean by the symbolic expression

$$Z(f) = \int [dp] [dq] \overline{\Omega(q(\infty))} \Omega(q(-\infty)) e^{i \int dt f_a(t) q^a(t)} e^{-i \int dt [p_a(t) \dot{q}^a(t) - H(q(t), p(t))]} \quad (2.38)$$

some form of limit of (2.37) as $N \rightarrow \infty, T \rightarrow \infty, M \rightarrow \infty, R \rightarrow \infty$ in that precise order, if it exists. $Z(f)$ is the generating functional of all Feynman functions and in $Z(f)/Z(0)$ the vacuum to vacuum correlations are removed.

It is important to note that the path integral (2.38) is over phase space and not over configuration space to begin with. Integrating out the momenta explicitly is only possible when H has a sufficiently simple dependence on p and even then it may modify the ‘‘Lebesgue measure’’ $[dq]$. We will see this explicitly for the case of GR in section 4. Note also the dependence of $Z(f)$ on the cyclic vector Ω or equivalently the corresponding state ω .

2.5.2 Euclidian formulation

There is also an Euclidian version of (2.38). We obtain it by going back to (2.32) and now using the second way in (2.27) to express $\Omega_{0,M}$ in terms of Ω_M

$$\chi_M(f) = \lim_{T \rightarrow \infty} \lim_{N \rightarrow \infty} \frac{\langle \Omega_M, e^{-TH_M} e^{-it_{N-1}H_M} e^{i\Delta_N f_a(t_{N-1})Q^a} e^{i\Delta_N H_M} \dots e^{i\Delta_N f_a(t_{-N})Q^a} e^{it_{-N}H_M} e^{-TH_M} \Omega_M \rangle}{\langle \Omega_M, e^{-(T-(-T))H_M} \Omega_M \rangle} \quad (2.39)$$

We define $2N$ smearing functions f_a^k by $f_a^k := f_a(t_k)$ with respect to the label a (which are considered independent of t). Then we analytically continue $\tau \rightarrow i\sigma, \sigma > 0$ which means $t_k = k\tau/N \rightarrow is_k, s_k = k\sigma/N$. Then we define, abusing the notation $f_a(s_k) := -f_a^k$. We take T sufficiently large $T \geq \sigma$ and extend f trivially to $\mathbb{R} - [-\sigma, \sigma]$ so that we can set $T = \sigma$. Then (2.39) becomes with $\Delta_N = \frac{T}{N}$

$$\chi_M(f) = \lim_{T \rightarrow \infty} \lim_{N \rightarrow \infty} \frac{Z_{M,T,N}^E(f)}{Z_{M,T,N}^E(0)}, \quad Z_{M,T,N}^E(f) = \langle \Omega_M, e^{-\Delta_N H_M} e^{\Delta_N f_a(s_{N-1})Q^a} \dots e^{-\Delta_N f_a(s_{-N})} e^{-\Delta_N H_M} \Omega_M \rangle \quad (2.40)$$

The rest of the analysis is the same as in the ‘‘Lorentzian’’ case and results in the symbolic formula

$$Z^E(f) \int [dp] [dq] \overline{\Omega(q(\infty))} \Omega(q(-\infty)) e^{\int ds f_a(s) q^a(s)} e^{-\int ds [i p_a(s) \dot{q}^a(s) + H(q(s), p(s))]} \quad (2.41)$$

which is the generating functional of Schwinger functions which are obtained as (without vacuum correlators)

$$S_N((s_N, a_N), \dots, (s_1, a_1)) = \left[\frac{\delta^N}{\delta f_{a_N}(s_N) \dots \delta f_{a_1}(s_1)} \frac{Z^E(f)}{Z^E(0)} \right]_{f=0} \quad (2.42)$$

2.5.3 Unconstrained phase space path integral

We now want to relate these reduced phase space path integrals to path integrals over the unconstrained phase space. This can be accomplished by recalling that y_j is eliminated by using the constraint $\hat{C}_j = y_j + h_j = 0$ while x^j is eliminated using the gauge fixing condition $G^j = x^j - k^j$. Consider again first the Lorentzian version. Then

$$Z(f) = \int [dq] [dp] [dy] [dx] \delta[\hat{C}] \delta[G] \overline{\Omega(q(\infty))} \Omega(q(-\infty)) e^{i \langle f \cdot q \rangle} e^{-i \langle p \cdot \dot{q} - H \rangle} \quad (2.43)$$

with $\delta[\hat{C}] = \prod_t \delta(\hat{C}_j(z(t)))$ and $\delta[G] = \prod_t \delta(x^j(t) - k^j(t))$ and $\langle (\cdot) \rangle = \int dt(\cdot)$. Using the relation $C_j = M_j^k \hat{C}_k$ between the original constraints C and the constraints \hat{C} solved for the y we have $\{C_j, G^k\} = M_j^k$ when $\hat{C}_k = 0$ whence

$$\delta[\hat{C}] = |\det[M]| \delta[C], \quad \det[M] = \prod_t \det(M(t)) \quad (2.44)$$

Strictly speaking, formula (2.44) is not entirely correct: If C_j labels constraints quadratic in the momenta y then the full constraint surface $\bar{\Gamma}$ defined by $C_j = 0$ for all j of the phase space Γ has in fact many branches corresponding, for each j , to a choice of the two possible roots $y_j = -h_j^\pm$ where in the canonical quantisation we only quantised the sector $y_j = -h_j$, $h_j := h_j^+$ corresponding to the positive root for all j . Since typically h_j^- is negative that sector is selected if we restrict the integral over y_j to the negative half-axis but this would prevent us from using Gaussian integral techniques to integrate out y . To analyse this issue, let $\sigma_j = \pm 1$ and $\bar{\Gamma}_\sigma = \{(q, p, x, y) \mid y_j = -h_j^{\sigma_j}(q, p, x)\}$ then we have a disjoint (up to measure zero sets) union $\bar{\Gamma} = \cup_\sigma \bar{\Gamma}_\sigma$. Accordingly, we may pick on each sector $\bar{\Gamma}_\sigma$ an individual gauge fixing condition $G_\sigma^j = x^j - k_\sigma^j$ which defines the corresponding sector $\hat{\Gamma}_\sigma = \{(p, q, x, y), y = -h^\sigma(q, p, x), x = k_\sigma\}$ of the reduced phase space. We assume now 1. that we can write the original constraints C_j , i.e. before solving them for y_j , in the form

$$C_j = \sum_k P_j^k \hat{C}_k^+ \hat{C}_k^-, \quad \hat{C}_k^\pm = y_k + h_k^\pm \quad (2.45)$$

for some invertible matrix P . This is not the most general situation but it will be satisfied for the concrete theory that we study later on. Then the gauge stability condition on the sector σ is given by

$$\dot{k}_\sigma^j = \{f^k C_k, G_\sigma^j\}_{\bar{\Gamma}_\sigma} = -f^k P_k^j \sigma_j (h_j^+ - h_j^-) =: f^k [M_\sigma]_k^j \quad (2.46)$$

which can be solved for $f^j = f_\sigma^j$. The reduced Hamiltonian on the sector σ is determined by the condition that for any function F of q, p

$$\{H_\sigma, F\} = \{f^k C_k, F\}_{\hat{\Gamma}_\sigma, f=f_\sigma} = -f_\sigma^k \sum_j P_k^j \sigma_j (h_j^+ - h_j^-) \{h_j^{\sigma_j}, F\} = \sum_j \dot{k}_\sigma^j \{h_j^{\sigma_j}, F\} \quad (2.47)$$

thus

$$H_\sigma = \sum_j \dot{k}_\sigma^j h_j^{\sigma_j}(q, p, x = k_\sigma) = -[\sum_j \dot{x}^j y_j]_{\hat{\Gamma}_\sigma} \quad (2.48)$$

which returns the earlier result for the "totally positive" sector. Suppose now that in fact 2. $h_j^\sigma = \sigma h_j$ and that 3. $h_j(q, p, x) = h_j(q, p, -x)$. Then for the choice $k_\sigma^j := \sigma_j k^j$ for k^j the choice for the totally positive sector we find

$$H_\sigma = \sum_j \dot{k}^j h_j(q, p, x = k) = H \quad (2.49)$$

is *sector independent* and agrees with the value of the totally positive sector. Next we have (subscript σ denotes sector restriction)

$$\begin{aligned} \delta(G) |\det(\{C, G\})| \delta(C) &= \sum_\sigma \delta(G)_\sigma |\det(\{C, G\})|_\sigma \delta(C)_\sigma \\ &= \sum_\sigma \delta(G)_\sigma |\det(M_\sigma)| \prod_j \delta(C_j)_\sigma = \sum_\sigma \delta(G)_\sigma |\det(M_\sigma)| \prod_j \delta(-\sum_k P_j^k (h_k^+ - h_k^-) \sigma_k \hat{C}_k^{\sigma_k}) \\ &= \sum_\sigma \delta(G)_\sigma \delta(\hat{C}_\sigma), \quad \delta(\hat{C}_\sigma) = \prod_j \delta(\hat{C}_j^{\sigma_j}) \end{aligned} \quad (2.50)$$

Therefore using (2.47)

$$\begin{aligned} [\sum_\sigma 1] \int [dq dp] e^{i \langle f, q \rangle - i [\langle p, \dot{q} \rangle - H]} &= \sum_\sigma \int [dq dp dx dy] \delta(G_\sigma) \delta(\hat{C}_\sigma) e^{i \langle f, q \rangle - i [\langle p, \dot{q} \rangle - H_\sigma]} \\ &= \int [dq dp dx dy] |\det(\{G, C\})| \delta(G) \delta(C) e^{i \langle f, q \rangle - i [\langle p, \dot{q} \rangle + \langle y, \dot{x} \rangle]} \end{aligned} \quad (2.51)$$

It follows that under the assumptions 1.-3. formula (2.44) is correct up to the constant factor $\sum_{\sigma} 1$ which drops out in the quotient $Z(f)/Z(0)$ that removes pure vacuum to vacuum amplitudes.

We also can bring the delta distribution for the constraints to the exponent using a Lagrange multiplier N^j . This results in (again constant factors e.g. powers of 2π are dropped as we consider the fraction $Z(f)/Z(0)$)

$$Z(f) = \int [dq] [dp] [dy] [dx] [dN] |\det(M)| \delta[G] \overline{\Omega(q(\infty))} \Omega(q(-\infty)) e^{i\langle f \cdot q \rangle} e^{-i\langle p \cdot \dot{q} + y \cdot \dot{x} - N \cdot C \rangle} \quad (2.52)$$

If we introduce independent ghost and anti-ghost fields η^j, ρ_j together with their ‘‘Berezin measures’’ $[d\eta][d\rho]$ one can also bring the Dirac matrix determinant into the exponent. The δ distribution of G can be brought to the exponent by introducing an additional Lagrange multiplier integral $\delta[G] = \int [dl] \exp(i \langle l \cdot G \rangle)$. An argument often used is that $Z(f)$ is unchanged when replacing G by $G - l$ as when $Z(f)$ is gauge invariant. If that was true in our case we could integrate both numerator and denominator with $\int [dl] \exp(- \langle l \cdot \kappa l \rangle)$ where κ is a positive integral kernel which would then replace $\delta[G]$ by $\exp(- \langle l \cdot \kappa l \rangle)$. However, this is manifestly not the case because unwinding what we have done would then replace the physical Hamiltonian by $H = \dot{k} \cdot h(P, Q, ; k) \rightarrow [\dot{k} + \dot{l}] \cdot h(Q, P; k + l) \neq H$ unless h is independent of k (which is true in the case considered here (no explicit time dependence)) and if $\dot{l} = 0$. However, the latter condition would imply that l is the same at all times and thus we could not solve $\delta[G] = \prod_{t,j} \delta(G^j(t) - l^j)$ for all t . To obtain no explicit time dependence we need $\dot{k} = \text{const.}$ but not $\dot{k} = 0$ (then we would have $H = 0$). The reason why this happens is due to the different notions of gauge transformations in the Lagrangian and Hamiltonian setting [30]: The Lagrangian gauge transformations are kinematical, i.e. they come from a Lie algebra (in GR it is the Lie algebra of vector fields, the Lie algebra of the diffeomorphism group). In the Hamiltonian setting the gauge transformations are generated by the constraints and they do not form a Lie algebra if the structure functions are not constant on the phase space (which is precisely the case in GR). The two notions coincide only on shell, when the classical equations of motion hold but of course in the path integral the set of classical paths has measure zero.

We will therefore leave $\delta[G]$ untouched and obtain the final expression

$$Z(f) = \int [dq] [dp] [dy] [dx] [dN] \int [d\eta] [d\rho] e^{-i\langle \eta \cdot M \cdot \rho \rangle} |\overline{\Omega(q(\infty))} \Omega(q(-\infty)) \delta[G] e^{i\langle f \cdot q \rangle} e^{-i\langle p \cdot \dot{q} + y \cdot \dot{x} - N \cdot C \rangle} \quad (2.53)$$

where again $M = (\{C, G\})$ is the Dirac matrix between constraints and gauge fixing conditions. Note that the ghost integral produces $\det(M)$ rather than $|\det(M)|$ (times an infinite power of i which drops out in the modulus) which is why we keep the modulus, a fact often ignored.

The expression (2.53) is as close as one can get to the usual formula $\int [d\Phi] e^{-iS(\Phi)}$, where Φ denotes the collection of all configuration fields, without further specifying the constraints. Indeed one would like to integrate out p, y in (2.53) which would turn the Hamiltonian action that has appeared in the exponent into the Lagrangian action. Whether this is possible explicitly depends on the y, p dependence of C . Even when possible, this may result in an additional measure contribution coming from a corresponding Jacobean. We will see that this precisely happens in GR where for our model it turns out that the dependence of $\langle N \cdot C \rangle$ and $\langle \eta, M \cdot \rho \rangle$ is at most quadratic and linear in y respectively and that the appearing Jacobean can be absorbed into a redefinition of the Dirac matrix M . That this is possible in the Lorentzian framework rests on the following elementary property of the Gaussian integral:

$$\int_{\mathbb{R}} du e^{z u^2} \quad (2.54)$$

It exists for all $z \in \mathbb{C}$ such that either $\Re(z) < 0$ or $\Re(z) = 0, \Im(z) \neq 0$. It does not exist for $\Re(z) > 0$. Finally note also that correctly there is only a smearing function f for the true configuration degrees of freedom q and not for x .

We now repeat this analysis for the Euclidian formulation. Almost all the work has been done already, we just need to remove two factors of i . We find

$$Z^E(f) = \int [dq] [dp] [dy] [dx] [dN] \int [d\eta] [d\rho] e^{\langle -\eta \cdot M \cdot \rho \rangle} |\overline{\Omega(q(\infty))} \Omega(q(-\infty)) \delta[G] e^{\langle f \cdot q \rangle} e^{-i\langle p \cdot \dot{q} + y \cdot \dot{x} - N \cdot C \rangle} \quad (2.55)$$

This looks innocent but it is actually disastrous for the Euclidian programme: When integrating over y one does not invert the Legendre transform between the Lagrangean and the Hamiltonian formulation because the symplectic term $y \cdot x$ comes with a relative factor of i as compared to the constrained Hamiltonian $N \cdot C$. This is the reason why we prefer the Lorentzian version of the Wetterich equation. We will see this in more detail in section 4. One may argue that the Euclidian path integral is better defined than the Lorentzian version because the Lorentzian integral is oscillatory while the Euclidian one is damped. This is in fact true when the Hamiltonian is bounded from below. However, exactly for GR the situation is opposite because the Euclidian action is not bounded from below due to the famous negative conformal mode [25]. This corresponds to case $\Re(z) > 0$ for which (2.54) does not exist.

3 Asymptotically safe quantum gravity: Summary and logic from point of view of CQG

As was stressed and reviewed in the previous section one can go back and forth between the Hamiltonian and path integral formulation of the QFT of a gauge theory. However, whatever formulation one prefers, the rigorous mathematical formulation is a challenge in interacting QFT and renormalisation methods are an essential ingredient in order to arrive at a mathematically well defined theory with predictive power. From that perspective we consider ASQG as a particular incarnation of that general theme.

The starting point of the asymptotic safety approach for a general theory is the generating functional $Z(f)$ of time ordered N-point functions (Lorentzian version) or $Z^E(f)$ of N-point Schwinger functions (Euclidian version) from which one can obtain effective actions by the usual methods. As we stressed in the derivation of the previous section, the symbolic expressions (2.38) and (2.41) (reduced phase space path integral) and (2.53) and (2.55) (full phase space path integral), while well motivated, they and their effective actions are ill-defined as they stand. The idea is to assume that the effective actions are well-defined, to derive a renormalisation group equation from them which by itself is well defined (Wetterich equation) and then to forget about the derivation and rather take the renormalisation group equation (RGE) as the fundamental guiding principle for the construction of the theory. A solution of the RGE then *defines* the theory. In the subsequent subsections we sketch elements of that programme.

ASQG practitioners can mostly skip this section, except that we develop both the Lorentzian and Euclidian versions which leads to adaptations by powers of i in the Wetterich equation, the heat kernel calculus and the Laplace versus Fourier transform with respect to heat kernel time. There are also some remarks concerning the treatment of the ghost part of the effective action and the functional derivatives at zero of the Wetterich equation with respect to the field on which the EAA depends, which are not often mentioned in the ASQG literature, see however [26].

3.1 Effective action without gauge invariance

Given the generating functional $Z_L(f), Z_E(f)$ of time ordered or Schwinger N-point functions respectively, formally defined by

$$Z_L(f) := \int [dq] e^{-iS_L(q)} e^{i\langle f, q \rangle}, \quad Z_E(f) := \int [dq] e^{-S_E(q)} e^{\langle f, q \rangle}, \quad \langle f, q \rangle := \langle f \cdot q \rangle \quad (3.1)$$

the functional

$$C_L(f) := i^{-1} \ln(Z_L(f)), \quad C_E(f) := \ln(Z_E(f)) \quad (3.2)$$

is the generating functional of “connected” such functions and its Legendre transform

$$\Gamma_L(\hat{q}) := \text{extr}_f [-C_L(f) + \langle f, \hat{q} \rangle], \quad \Gamma_E(\hat{q}) := \text{extr}_f [-C_E(f) + \langle f, \hat{q} \rangle] \quad (3.3)$$

the effective action or generating functional of “one particle irreducible” (1PI) such functions. This terminology comes from their graphical interpretation, see e.g. [27]. If any of these objects is well defined in the sense that its functional derivatives yield non-singular tempered distributions, so are the others and one has “solved the theory”. The effective action is the most compact definition of the theory because the 1-PI functions are the “atoms” of all correlation functions.

3.2 Background field method without gauge invariance

It maybe convenient for various purposes to introduce a background field \bar{q} , especially in GR where it can be used to define preferred covariant derivatives with respect to a background metric in order to construct additional structures (such as those used in asymptotic safety). The method has been invented in [12] in fact first in the Lorentzian signature setting and is naturally used in QFT in CST for example in order to define the Hadamard condition on the 2-point function [15, 13].

The background objects will carry an additional overbar and are defined by (Euclidian: $s = 0$; Lorentzian: $s = 1$)

$$\begin{aligned}\bar{Z}_s(f, \bar{q}) &:= \int [dh] e^{-i^s S_s(\bar{q}+h)} e^{i^s \langle f \cdot h \rangle} \\ \bar{C}_s(f, \bar{q}) &:= i^{-s} \ln(\bar{Z}_s(f, \bar{q})) \\ \bar{\Gamma}_s(\hat{q}, \bar{q}) &:= \text{extr}_f [-\bar{C}_s(f, \bar{q}) + \langle f \cdot \hat{q} \rangle]\end{aligned}\tag{3.4}$$

They are explicitly dependent on \bar{q} . The relation to the background independent objects are as follows: Introducing a new integration variable $q = \bar{q} + h$ in the first line we find

$$\begin{aligned}\bar{Z}_s(f, \bar{q}) &= e^{-i^s \langle f, \bar{q} \rangle} Z_s(f) \\ \bar{C}_s(f, \bar{q}) &= C_s(f) - \langle f \cdot \bar{q} \rangle \\ \bar{\Gamma}_s(\hat{q}, \bar{q}) &= \Gamma_s(\hat{q} + \bar{q})\end{aligned}\tag{3.5}$$

Since what we are interested in is really the background independent object $\Gamma_s(\hat{q})$ we may obtain it from the background dependent object $\bar{\Gamma}_s(\hat{q}, \bar{q})$ by the simple formula

$$\Gamma_s(\hat{q}) = [\bar{\Gamma}_s(\hat{q}', \bar{q})]_{\hat{q}'=0, \bar{q}=\hat{q}}\tag{3.6}$$

The reason for not setting instead $\hat{q}' = \hat{q}$, $\bar{q} = 0$ is because a zero background may jeopardise intermediate constructions that one performs with it (e.g. a zero background metric cannot be inverted).

3.3 Effective action with gauge invariance

We now consider a gauge theory with gauge group \mathfrak{G} and Lie algebra \mathfrak{g} . The group \mathfrak{G} acts on the configuration space Ω of the fields q by gauge transformations

$$\alpha : \mathfrak{G} \times \Omega \rightarrow \Omega; q \mapsto \alpha_g(q), \alpha_g \circ \alpha_{g'} = \alpha_{gg'}\tag{3.7}$$

for $g, g' \in \mathfrak{G}$. A gauge fixing condition

$$G : \Omega \rightarrow \mathfrak{g}; q \mapsto G(q)\tag{3.8}$$

is a Lie algebra valued function on the configuration space subject to the condition that for each $q \in \Omega$ there exists a unique $g(q) \in \mathfrak{G}$ such that $G(\alpha_{g(q)}(q)) = 0$. If $\exp : \mathfrak{g} \rightarrow \mathfrak{G}$ is the exponential map then one can instead require that for each $q \in \Omega$ there exists a unique $u(q) \in \mathfrak{g}$ such that $G(\alpha_{\exp(u(q))}(q)) = 0$. The conditions are not equivalent because the exponential map is not a bijection, typically it is neither injective (e.g. the Lie algebra of $U(1)$ is \mathbb{R} but all $u = v + 2\pi n, n \in \mathbb{Z}, v \in [0, 2\pi)$ map to $g = e^{iu} = e^{iv}$) nor surjective (general $g \in \mathfrak{G}$ are not in the component of the identity). We will, as is customary ignore these complications and work with the Lie algebra condition on the gauge fixing condition. Then we have the Fadeev-Popov identity

$$1 = \int_{\mathfrak{g}} [du] \delta[G(\alpha_{\exp(u)}(q))] \left| \det \left[\frac{\delta}{\delta u} G(\alpha_{\exp(u)}(q)) \right] \right|\tag{3.9}$$

for any $q \in \Omega$.

Let now $F : \Omega \rightarrow \mathbb{C}$ be a gauge invariant function $F \circ \alpha_g = F \forall g \in \mathfrak{G}$. Then the symbolic functional integral

$$I_F := \int_{\Omega} [dq] F(q)\tag{3.10}$$

can also be written

$$I_F := \int_{\mathfrak{g}} [du] \int_{\Omega} [dq] F(q) \delta[G(\alpha_{\exp(u)}(q))] \left| \det \left[\frac{\delta}{\delta u} G(\alpha_{\exp(u)}(q)) \right] \right| \quad (3.11)$$

We introduce a new integration variable $q' = \alpha_{\exp(u)}(q)$. If α_g is a bijection for any $g \in G$ (e.g. no fixed points) then $\alpha_{\exp(u)}$ is a permutation on Ω so that the Jacobean or Radon-Nikodym derivative is

$$J(q, u) := \frac{d[\alpha_{\exp(u)}(q)]}{[dq]} = \left| \det \left[\frac{\delta \alpha_{\exp(u)}(q)}{\delta q} \right] \right| = 1 \quad (3.12)$$

It would be enough to require that $J(q, u) = J(u)$ just depends on u . Clearly $F(q) = F(q')$ by gauge invariance and provided that

$$\det \left[\frac{\delta}{\delta} G(\alpha_{\exp(v+u)}(q)) \right] - \det \left[\frac{\delta}{\delta v} G(\alpha_{\exp(v)} \exp(u)(q)) \right] = O(v) \quad (3.13)$$

we have

$$\det \left[\frac{\delta}{\delta u} G(\alpha_{\exp(u)}(q)) \right] = \left\{ \det \left[\frac{\delta}{\delta v} G(\alpha_{\exp(v+u)}(q)) \right] \right\}_{v=0} = \left\{ \det \left[\frac{\delta}{\delta v} G(\alpha_{\exp(v)}(q')) \right] \right\}_{v=0} \quad (3.14)$$

Then after relabelling q' by q

$$I_F = \int_{\mathfrak{g}} [du] J(u) \int [dq] F(q) \delta[G(q)] \left| \det \left[\frac{\delta}{\delta v} G(\alpha_{\exp(v)}(q)) \right] \right|_{v=0} \quad (3.15)$$

or for any fixed gauge invariant functional F_0

$$\frac{I_F}{I_{F_0}} = \frac{\int [dq] F(q) \delta[G(q)] \left| \det \left[\frac{\delta}{\delta v} G(\alpha_{\exp(v)}(q)) \right] \right|_{v=0}}{\int [dq] F_0(q) \delta[G(q)] \left| \det \left[\frac{\delta}{\delta v} G(\alpha_{\exp(v)}(q)) \right] \right|_{v=0}} \quad (3.16)$$

If G meets the stated conditions then so does $G - w$ where $w \in \mathfrak{g}$ and since (3.16) does not depend on the choice of this class of G we can integrate both numerator and denominator against $\int [dw] \exp(-\langle w, \kappa \cdot w \rangle)$ with G replaced by $G - w$ where $\kappa : \mathfrak{g} \rightarrow \mathfrak{g}$ is an integral kernel. Finally introducing ghost and anti-ghost η, ρ Berezin integrals one finds

$$\frac{I_F}{I_{F_0}} = \frac{\int [dq] [d\eta] [d\rho] F(q) e^{-\langle G(q), \kappa G(q) \rangle} \exp(\langle \eta, [\frac{\delta}{\delta v} G(\alpha_{\exp(v)}(q))]_{v=0} \cdot \rho \rangle)}{\int [dq] [d\eta] [d\rho] F_0(q) e^{-\langle G(q), \kappa G(q) \rangle} \exp(\langle \eta, [\frac{\delta}{\delta v} G(\alpha_{\exp(v)}(q))]_{v=0} \cdot \rho \rangle)} \quad (3.17)$$

The application of (3.17) consists in picking a gauge invariant action $S_s(q)$ and $F_0 = e^{-i^s S_s}$ while $F = e^{-i^s S_s} O$ where O is any gauge invariant functional that one may obtain by functionally differentiating $e^{i^s \langle f, q \rangle}$ at $f = 0$ which thus gives rise to the generating functional (rescaling κ and the ghost term by e^{i^s})

$$Z_s(f) = \int [dq] [d\eta] [d\rho] e^{-i^s S_s(q)} e^{-e^{i^s} \langle G(q), \kappa G(q) \rangle} \exp(e^{i^s} \langle \eta, [\frac{\delta}{\delta v} G(\alpha_{\exp(v)}(q))]_{v=0} \cdot \rho \rangle) e^{i^s \langle f, q \rangle} \quad (3.18)$$

for the case of a gauge theory. From here on the definition of $C_s(f), \Gamma_s(\hat{q})$ is identical to the case without gauge invariance.

3.4 Background field method with gauge invariance

The background dependent generating functional in the case with gauge invariance is defined by

$$\bar{Z}_s(f, \bar{q}) = \int [dh] [d\eta] [d\rho] e^{-i^s S_s(\bar{q}+h)} e^{-e^{i^s} \langle \bar{G}_{\bar{q}}(h), \kappa \bar{G}_{\bar{q}}(h) \rangle} \exp(e^{i^s} \langle \eta, [\frac{\delta}{\delta v} \bar{G}_{\bar{q}}(\alpha_{\exp(v)}^{\bar{q}}(h))]_{v=0} \cdot \rho \rangle) e^{i^s \langle f, h \rangle} \quad (3.19)$$

while $\bar{C}_s(f, \bar{q}), \bar{\Gamma}(\hat{q}, \bar{q})$ are defined as in the case without gauge invariance. Here $\bar{G}_{\bar{q}}(h)$ is a background dependent gauge fixing condition which is as admissible as any other one subject to the conditions spelled out in the previous subsection and

$$\alpha_{\exp(u)}^{\bar{q}}(h) := \alpha_{\exp(u)}(\bar{q} + h) - \bar{q} \quad (3.20)$$

If we now introduce $q = \bar{q} + h$ as a new integration variable we find again

$$\bar{Z}_s(f, \bar{q}) = Z_s(f) e^{-i^s \langle f, \bar{q} \rangle} \quad (3.21)$$

provided that we choose in Z_s the gauge fixing condition

$$G(q) := \bar{G}_{\bar{q}}(q - \bar{q}) \quad (3.22)$$

In particular we have again (3.6). Thus although we may use a background heavily in order to define $\bar{Z}_s(f, \bar{q})$, we can use the corresponding $\bar{\Gamma}_s(\hat{q}, \bar{q})$ to define the background independent $\Gamma_s(q)$.

The real power of the background method in usual QFT arises due to the existence of distinguished, so-called "background gauges". These are particular gauge conditions with the property that $Z_s(f, \bar{q})$ is in fact invariant under simultaneous gauge transformations of both f and \bar{q} for suitable choice of kernel κ (see [12]). Here a gauge transformation of \bar{q} is defined to be the same as for q while the gauge transformation of f is defined by duality

$$\langle f, \alpha_g(q) \rangle =: \langle \alpha_{g^{-1}}^*(f), q \rangle \quad (3.23)$$

In case of GR these background gauge transformations are just the ordinary transformations of f, \bar{q} viewed as ordinary tensor fields and thus suitable background gauges have to be diffeomorphism covariant in the usual sense, e.g. the deDonder gauge would be such a gauge. This has the advantage that the terms that appear in the effective action are restricted to be gauge invariant functionals of \hat{q} under the usual action α_g which implies a huge reduction in the number of all possible terms.

3.5 Effective average action

We now modify the generating functional $\bar{Z}_s(f, \bar{q})$ by

$$\begin{aligned} \bar{\Gamma}_{s,k}(f, \bar{q}) := & \int [dh] [d\eta] [d\rho] e^{-i^s S_s(\bar{q}+h)} e^{-e^{i^s} \langle \bar{G}_{\bar{q}}(h), \kappa \bar{G}_{\bar{q}}(h) \rangle} \exp(e^{i^s} \langle \eta, [\frac{\delta}{\delta v} \bar{G}_{\bar{q}}(\alpha_{\exp(v)}^{\bar{q}}(h))]_{v=0} \cdot \rho \rangle) \times \\ & e^{i^s \langle f, h \rangle} e^{-\frac{i^s}{2} \langle h, R_k(\bar{q}) h \rangle} \end{aligned} \quad (3.24)$$

If we are given a system without gauge invariance, the terms involving G and the ghost integrals have to be dropped.

The one parameter family $k \mapsto R_k(\bar{q})$ of possibly background dependent and coupling dependent (coupling dependence increases the non-linearity in the flow of couplings as derived below) integral kernels is supposed to obey a number of properties. Consider harmonic analysis on the spacetime determined by 1. the manifold on which the given QFT is defined and 2. the background metric determined by \bar{q} as well as the Fourier modes $\hat{R}_k(p)$ of $\hat{R}_k(\bar{q})$ where p are the mode labels with respect to a generalised eigenbasis of the background d'Alembertian ($s = 1$) or Laplacian ($s = 0$) respectively. Then:

1. $\lim_{k \rightarrow 0} \hat{R}_k(p) = 0$ at fixed p . This property will ensure that the k -dependent effective average action that we obtain from it as below reduces to the background dependent effective action which in turn determines the actual object of interest, the background independent effective action.
2. $\lim_{k \rightarrow \infty} \hat{R}_k(p) = \infty$ at fixed p . This property makes the additional term suppress non-zero values of h either due to oscillations ($s = 1$) or due to damping so that the effective average action becomes almost exactly equal to S_s in the saddle point approximation.
3. $\lim_{p \rightarrow 0} \hat{R}_k(p) = \text{const.}$ at fixed k . This property ensures that the additional term acts like an additional mass term for $p \leq k$ which acts suppressingly either due to oscillations or damping.
4. $\lim_{p \rightarrow \infty} \hat{R}_k(p) = 0$ at fixed k . This property ensures that the momenta $p \geq k$ are unsuppressed by the additional term and thus are integrated over in the usual way.

Altogether $\hat{R}_k(p) \approx \theta(k - p) \hat{R}(k)$ with $\hat{R}(0) = 0$, monotonously increasing as $k \rightarrow \infty$ where it diverges. As $R_k(\bar{q})$ appears in the Wetterich equation (see below) in the form $\partial_k R_k(\bar{q})$ inside a trace (integral over p), that trace receives essential contributions only from a neighbourhood of k which improves convergence of the trace. Following the intuition that $R_k(\bar{q})$ suppresses integrating out modes $p \leq k$, it follows that $\bar{Z}_{s,k}(f, \bar{q})$ has

the more momenta integrated out the lower k , i.e. lowering k is similar to, but different from, coarse graining in a strict Wilsonian renormalisation group sense.

We define now in almost the usual pattern the k -dependent functionals

$$\begin{aligned}\bar{C}_{s,k}(f, \bar{q}) &= i^{-s} \ln(\bar{Z}_{s,k}(f, \bar{q})) \\ \bar{\Gamma}_{s,k}(\hat{q}, \bar{q}) &= \text{extr}_f [-\bar{C}_{s,k}(f, \bar{q}) + \langle f, \hat{q} \rangle] - \frac{1}{2} \langle \hat{q}, R_k(\bar{q}) \hat{q} \rangle\end{aligned}\quad (3.25)$$

Note the additional $R_k(\bar{q})$ dependent term that we subtract in the second line that is non-standard as compared to the previous sections. Again, due to $R_0(\bar{q}) = 0$ that additional term does not spoil that

$$\bar{\Gamma}_{s,k=0}(\hat{q}, \bar{q}) = \bar{\Gamma}_s(\hat{q}, \bar{q}) \quad (3.26)$$

3.6 Wetterich equation

To show that the Wetterich equation really is just a consequence of the definition of the EAA, we give here an elementary derivation for completeness. We will suppress the argument \bar{q} of the functions that depend on it in order not to clutter the formulae and because it plays no role in the following analysis.

By definition of the Legendre transform, the extremal condition

$$\frac{\delta \bar{C}_{s,k}(f)}{\delta f} - \hat{q} = 0 \quad (3.27)$$

must be solved for $f = f_{s,k}(\hat{q})$. We assume that the solution is unique and thus find explicitly

$$\bar{\Gamma}_{s,k}(\hat{q}) = [-\bar{C}_{s,k}(f) + \langle f, \hat{q} \rangle]_{f=f_{s,k}(\hat{q})} - \frac{1}{2} \langle \hat{q}, R_k \hat{q} \rangle \quad (3.28)$$

Since (3.27) becomes an identity when evaluated at $f = f_{s,k}(\hat{q})$

$$\hat{q} \equiv \left[\frac{\delta \bar{C}_{s,k}(f)}{\delta f} \right]_{f=f_{s,k}(\hat{q})} \quad (3.29)$$

it follows

$$\frac{\delta \bar{\Gamma}_{s,k}(\hat{q})}{\delta \hat{q}} = f_{s,k}(\hat{q}) - R_k \cdot \hat{q} \quad (3.30)$$

The functional derivative of (3.29) with respect to \hat{q} yields (1 denotes the identity integral kernel on the space Ω)

$$\begin{aligned}1 &= \left\langle \left[\frac{\delta^2 \bar{C}_{s,k}(f)}{\delta f \delta f} \right]_{f=f_{s,k}(\hat{q})}, \frac{\delta f_{s,k}(\hat{q})}{\delta \hat{q}} \right\rangle \\ &= \left\langle \left[\frac{\delta^2 \bar{C}_{s,k}(f)}{\delta f \delta f} \right]_{f=f_{s,k}(\hat{q})}, \left[\frac{\delta^2 \bar{\Gamma}_{s,k}(\hat{q})}{\delta \hat{q} \otimes \delta \hat{q}} + R_k \right] \right\rangle\end{aligned}\quad (3.31)$$

where in the second step we inserted the functional derivative of (3.30).

Next we write

$$\begin{aligned}\bar{Z}_{s,k}(f) &= \int d\mu[h] e^{i^s \langle f, h \rangle} e^{-\frac{i^s}{2} \langle h, R_k h \rangle}, \quad \frac{d\mu[h]}{d[h]} = \int [d\eta] [d\rho] e^{-i^s S_s(\bar{q}+h)} e^{-e^{i^s} \langle \bar{G}_{\bar{q}}(h), \kappa \bar{G}_{\bar{q}}(h) \rangle} \times \\ &\exp(e^{i^s} \langle \eta, \left[\frac{\delta}{\delta v} \bar{G}_{\bar{q}}(\alpha_{\text{exp}(v)}^{\bar{q}}(h)) \right]_{v=0} \cdot \rho \rangle)\end{aligned}\quad (3.32)$$

and find

$$\begin{aligned}\frac{\delta \bar{C}_{s,k}(f)}{\delta f} &= \frac{1}{\bar{Z}_{s,k}(f)} \int d\mu[h] h e^{i^s \langle f, h \rangle} e^{-\frac{i^s}{2} \langle h, R_k h \rangle} =: \hat{q}_{s,k}(f) \\ \frac{\delta^2 \bar{C}_{s,k}(f)}{\delta f \otimes \delta f} &= i^s \left[\frac{1}{\bar{Z}_{s,k}(f)} \int d\mu[h] h \otimes h e^{i^s \langle f, h \rangle} e^{-\frac{i^s}{2} \langle h, R_k h \rangle} - \hat{q}_{s,k}(f) \otimes \hat{q}_{s,k}(f) \right]\end{aligned}\quad (3.33)$$

and

$$\begin{aligned} & \frac{\partial \bar{\Gamma}_{s,k}(\hat{q})}{\partial k} = -\frac{\partial \bar{C}_{s,k}(f)}{\partial k} \Big|_{f=f_{s,k}(\hat{q})} - \frac{1}{2} \langle \hat{q}, [\partial_k R_k] \cdot \hat{q} \rangle \\ & = \frac{1}{2} \left[-\langle \hat{q}, [\partial_k R_k] \hat{q} \rangle + \left(\frac{i^s}{\bar{Z}_{s,k}(f)} \int d\mu[h] \langle h, [\partial_k R_k] h \rangle e^{i^s \langle f, h \rangle} e^{-\frac{i^s}{2} \langle h, R_k h \rangle} \right) \Big|_{f=f_{s,k}(\hat{q})} \right] \end{aligned} \quad (3.34)$$

We see that in (3.33) the combination $h \otimes h$ appears at general f while in (3.34) the combination

$$\langle h, [\partial_k R_k] h \rangle = \text{Tr}([\partial_k R_k] \cdot h \otimes h) \quad (3.35)$$

appears at $f = f_{s,k}(\hat{q})$. We can therefore relate (3.33) to (3.34) if we trace (3.34) with $\partial_k R_k$ at $f = f_{s,k}(\hat{q})$. Note the identity

$$[\hat{q}_{s,k}(f)]_{f=f_{s,k}(\hat{q})} = \hat{q} \quad (3.36)$$

Then

$$\begin{aligned} & \text{Tr} \left(\frac{\delta^2 \bar{C}_{s,k}(f)}{\delta f \otimes \delta f} [\partial_k R_k] \right)_{f=f_{s,k}(\hat{q})} \quad (3.37) \\ & = i^s \left[\left(\frac{1}{\bar{Z}_{s,k}(f)} \int d\mu[h] \langle h, [\partial_k R_k] h \rangle e^{i^s \langle f, h \rangle} e^{-\frac{i^s}{2} \langle h, R_k h \rangle} \right) \Big|_{f=f_{s,k}(\hat{q})} - \langle \hat{q}, [\partial_k R_k] \hat{q} \rangle \right] \\ & = 2 i^s \partial_k \bar{\Gamma}_{s,k}(\hat{q}) \\ & = \text{Tr} \left(\left[\frac{\delta^2 \bar{\Gamma}_{s,k}(\hat{q})}{\delta \hat{q} \otimes \delta \hat{q}} + R_k \right]^{-1} [\partial_k R_k] \right) \end{aligned} \quad (3.38)$$

where in the last step we solved (3.31) for $\left[\frac{\delta^2 \bar{C}_{s,k}(f)}{\delta f \otimes \delta f} \right]_{f=f_{s,k}(\hat{q})}$. This is the celebrated *Wetterich equation* [31] for both signatures (where we display the dependence on the background again)

$$\partial_k \bar{\Gamma}_{s,k}(\hat{q}, \bar{q}) = \frac{i^{-s}}{2} \text{Tr} \{ [\bar{\Gamma}_{s,k}^{(2)}(\hat{q}, \bar{q}) + R_k(\bar{q})]^{-1} [\partial_k R_k](\bar{q}) \} \quad (3.39)$$

This is an exact and non-perturbative identity which has to be obeyed by the EAA if it comes from a well defined $\bar{Z}_{s,k}(f, \bar{q})$. The idea is now to consider (3.28) as the fundamental equation, to solve it and try to find interesting solutions. By a solution we mean a solution of (3.39) which exists *for all values of k* , in particular for $k \rightarrow \infty$ and for $k \rightarrow 0$, in the sense described in the next subsection, so that our object of interest $\lim_{k \rightarrow 0} \bar{\Gamma}_{s,k}(\hat{q}, \bar{q})$ exists. The requirement that the solution has a suitable limit as $k \rightarrow \infty$ is the selection criterion of *asymptotic safety* for a viable and predictive theory because it typically dictates that all but a finite number of couplings have to be fine tuned to their fixed point values while the remaining couplings run with k and approach finite values at $k = 0$ in their dimensionful form. We stress that from the point of view of CQG the value $k = 0$ is the only one for which the EAA has a clear physical meaning, namely it becomes the standard effective action, because k has no physical interpretation.

Before closing this subsection we note that in the above derivation the ghost action was considered to be integrated out and thus gives a ghost determinant $\det(M) = \exp(\ln(\det(M))) =: \exp(-i^s S^g)$. The ghost action S^g and the gauge fixing action $S^{gf} = \langle \bar{G}_{\bar{q}}, \kappa \cdot \bar{G}_{\bar{q}} \rangle$ are thereby considered as corrections of the classical action. Therefore all three parts flow according to the above Wetterich equation. It is sometimes convenient to treat the ghost part individually by not integrating it out thereby avoiding the exponentiated logarithms. Thus one generalises the EAA by introducing also anticommuting test fields e_j, r^j for the ghosts and a similar R_k term bilinear in η, ρ . Then we introduce into the integrand of $\bar{Z}_{s,k}(f, \bar{q})$ the additional factor

$$\exp(i^s [\langle e, \eta \rangle + \langle r, \rho \rangle]) \exp(-i^s \langle \eta, R_k^g(\bar{q}) \rho \rangle) \quad (3.40)$$

where $R_k^g(\bar{q})$ is a ghost kernel with the same properties as $R_k(\bar{q})$. Then $\bar{Z}_{s,k}(f, \bar{q})$ becomes $\bar{Z}_{s,k}(f, e, r, \bar{q})$ and $\bar{C}_{s,k}(f, e, r, \bar{\rho}) = i^{-s} \ln(\bar{Z}_{s,k}(f, e, r, \bar{\rho}))$. Then

$$\bar{\Gamma}_{s,k}(\hat{q}, \hat{\eta}, \hat{\rho}, \bar{q}) := \text{extr}_{f,e,r} [\langle f, \hat{q} \rangle + \langle e, \hat{\eta} \rangle + \langle r, \hat{\rho} \rangle - \bar{C}_{s,k}(f, e, r, \bar{q})] - \left[\frac{1}{2} \langle \hat{q}, R_k(\bar{q}) \hat{q} \rangle + \langle \hat{\eta}, R_k^g(\bar{q}) \hat{\rho} \rangle \right] \quad (3.41)$$

Then a repetition of the above chain of steps gives the generalised Wetterich equation (we collect $\hat{\phi} := (\hat{q}, \hat{c}), \hat{c} = (\hat{\eta}, \hat{\rho})$) and suppress the dependence on \bar{q} for notational simplicity)

$$\partial_k \bar{\Gamma}_{s,k}(\hat{\phi}) = \frac{i^{-s}}{2} \text{Tr} \left\{ \left[\frac{\delta^2 \bar{\Gamma}_{s,k}^{(2)}(\hat{\phi})}{[\delta \hat{q}]^2} + R_k \right]^{-1} [\partial_k R_k] \right\} - i^{-s} \text{Tr} \left\{ \left[\frac{\delta^2 \bar{\Gamma}_{s,k}^{(2)}(\hat{\phi})}{[\delta \hat{c}]^2} + R_k^g \right]^{-1} [\partial_k R_k^g] \right\} \quad (3.42)$$

where the relative factor of -2 between bosonic and fermionic term comes from the anticommutativity of Grassman valued derivatives and the independence of $\hat{\rho}, \hat{\eta}$.

In view of lemma A.1 one must however pay attention to the following: It is not true that one recovers the original EAA (ghosts integrated out) by simply setting $\hat{c} = 0$. Rather we must apply lemma A.1 of the appendix to find the exact relation between (3.42) and (3.39).

3.7 Solving the Wetterich equation in principle and asymptotic safety prescription

The EAA is supposed to yield well defined tempered distributions when taking functional derivatives of any order and thus can be written as

$$\bar{\Gamma}_{s,k}(\hat{q}, \bar{q}) = \sum_{N=0}^{\infty} \bar{\Gamma}_{s,k}^N(\hat{\phi}, \bar{q}) \quad (3.43)$$

where $\bar{\Gamma}_{s,k}^N(\hat{q}, \bar{q})$ is a monomial in $\hat{\phi}$ of order N . One can use a geometric series expansion

$$[\bar{\Gamma}_{s,k}^{(2)} + R_k]^{-1} = R_k^{-1} \sum_{M=0}^{\infty} (-1)^M [\bar{\Gamma}_{s,k}^{(2)} R_k^{-1}]^M \quad (3.44)$$

and derive an infinite recursion among the $\bar{\Gamma}_{s,k}^N$. In practice one often combines R_k with a piece of $\bar{\Gamma}_{s,k}^{(2)}(\hat{q} = 0, \bar{q})$ in the geometric series expansion, see below. To make this practically useful one will try to find a basis of distributions such that the recursion closes. After such a basis $T_\alpha(\bar{q})$ of distributions has been identified, the coefficients $g_\alpha(k)$ of that basis have to be determined from the Wetterich equation where α runs through a countably infinite index set \mathcal{A} . These coefficients are in general dimensionful of mass dimension d_α and it is convenient to pass to dimensionfree objects $\hat{g}_\alpha(k) = k^{-d_\alpha} g_\alpha(k)$. Setting $t = \ln(k/k_0)$ for some arbitrary k_0 and collecting $\hat{g}(t) = \{g_\alpha(k_0 e^t)\}_{\alpha \in \mathcal{A}}$ one obtains an infinite first order ODE system

$$\frac{d}{dt} \hat{g}(t) = \beta(\hat{g}(t)) \quad (3.45)$$

which is autonomous, i.e. β does not depend explicitly on t if k is the only scale in the problem, but it is highly non-linear. By the general theory of finite dimensional ODE systems, maximal solutions exist and are unique given initial conditions at k_0 or equivalently integration constants c . We assume that we can also find here maximal solutions whose allowed domain D_c with respect to t depends on the choice c . Let then $\hat{g}_c(t), t \in D_c$ be the unique maximal solution with initial data c at $t = 1$ or $k = k_0$.

Such a solution of the flow equation (3.44) is an integral curve of the vector field β . Let \hat{g}_* be a zero of the vector field β , if it exists, $\beta(\hat{g}_*) = 0$. Now consider a general solution $\hat{g}_c(t)$. If we have to fine tune all but a finite number of the initial data c in order that $\lim_{t \rightarrow \infty} \hat{g}_c(t) = \hat{g}_*$ then the zero is called *predictive* for it says that we have only a finite dimensional space of admissible solutions (labelled by the non fine tuned, so called relevant, initial data) whose domain includes $(1, +\infty)$ and then $\lim_{t \rightarrow \infty} \hat{g}_c(t) = \hat{g}_*$ i.e. on that space of solutions \hat{g}_* is an UV fixed point of the flow. We may write $c = (c_r, c_i^*)$ where c_i^* is a point in an infinite dimensional space corresponding to the fine tuned parameters while c_r may still range in a finite dimensional space. The value c_i^* is considered a prediction of the theory given the zero \hat{g}_* .

We now return to the corresponding dimensionful couplings $g_{c_r, c_i^*}(k)$ which have a physical meaning at $k = 0$ and can be measured there. Only a finite number of them need to be measured to determine the remaining initial data c_r , after that the theory is completely fixed. We call the zero \hat{g}_* *complete* if the $g_{c_r, c_i^*}(0)$ are also finite. Note that this is not automatically the case.

To have an example in mind for such a scenario suppose that we have a basis consisting of only two distributions which come with dimensionful constants $g_1(k), g_2(k)$ of dimension cm^2 and cm^{-2} respectively. The dimensionless

objects are $\hat{g}_1(k) = k^2 g_1(k)$, $\hat{g}_2(k) = k^{-2} g_2(k)$. Suppose that $\hat{g}_1(k) = \frac{k^2}{k^2 + L_1^{-2}} \hat{g}_1^*$, $\hat{g}_2(k) = \frac{k^2 + L_2^{-2}}{k^2} \hat{g}_2^*$ with integration constants L_1, L_2 of dimension of length and some numerical values \hat{g}_1^*, \hat{g}_2^* . Then both couplings are relevant, both run into the zero of the vector field as $k \rightarrow \infty$ where they reach their fixed point values \hat{g}_1^*, \hat{g}_2^* . On the other hand $g_1(k) = \frac{\hat{g}_1^*}{k^2 + L_1^{-2}}$, $g_2(k) = \hat{g}_2^* (k^2 + L_2^{-2})$. Thus $g_1(0) = L_1^2 \hat{g}_1^*$, $g_2(0) = L_2^{-2} \hat{g}_2^*$ are supposed to equal measured values g_{10}, g_{20} which thus fixes L_1, L_2 . We see that it is important to require the finiteness of the *dimensionless* couplings at $k \rightarrow \infty$ and of the *dimensionful* couplings at $k \rightarrow 0$: If the couplings run, i.e. have non-trivial k dependence then typically the running will be monotonous and thus to have finiteness of say the dimensionful coupling at both $k = 0, \infty$ would be impossible.

The application to GR in n dimensions is as follows: We expect the EAA to involve curvature polynomials of order $N = 0, 1, 2, \dots$ such as $g_N(k) R^N$ where R is the Ricci scalar and $g_N(k)$ is its dimensionful coupling. As $d^n x g_N R^N$ is dimensionless, $g_N(k)$ has dimension cm^{2N-n} , hence $\hat{g}_N(k) = k^{2N-n} g_N(k)$. Accordingly a running with k of the form $g_N(k) = \frac{L_N^{2N-n}}{1 + (L_N k)^{2N-n}} \hat{g}_N^*$ would be compatible with both requirements where L_N is a constant of length dimension. Note that we do not require that all but a finite number of dimensionful couplings vanish, rather that they depend only on a finite number of such free length parameters L_N .

3.8 Solving the Wetterich equation in practice

It is of course out of question to complete the programme of the previous subsection for all couplings. One has to resort to the so-called truncations in the space of couplings. It would be nice to have some kind of error control at one's disposal which would grant that given some error ϵ and a fixed point discovered by a certain truncation space, all extensions of that truncation space exceeding some dimension depending on ϵ lead to corrections of the couplings of the given truncation of less than ϵ . It is of course very hard to come up with such an error control.

The still exact and non-polynomial Wetterich equation (we just display the non-ghost version)

$$\partial_k \bar{\Gamma}_{s,k}(\hat{q}, \bar{q}) = \frac{i^{-s}}{2} \text{Tr} \{ [\bar{\Gamma}_{s,k}^{(2)}(\hat{q}, \bar{q}) + R_k(\bar{q})]^{-1} [\partial_k R_k(\bar{q})] \} \quad (3.46)$$

is equivalent to an infinite hierarchy of non-polynomial equations among the 1-PI distributions

$$T_{s,k}^n(\bar{q}) := \left\{ \frac{\delta^n \bar{\Gamma}_{s,k}(\hat{q}, \bar{q})}{\otimes^n \delta \hat{q}} \right\}_{\hat{q}=0} \quad (3.47)$$

with $n = 0, 1, 2, \dots$. A compact way to write the n -th such equation is

$$\frac{\delta^n}{\otimes^n \delta \hat{q}} \{ \partial_k \bar{\Gamma}_{s,k}(\hat{q}, \bar{q}) - \frac{i^{-s}}{2} \text{Tr} ([\bar{\Gamma}_{s,k}(\hat{q}, \bar{q}) + R_k]^{-1} [\partial_k R_k(\bar{q})]) \}_{\hat{q}=0} = 0 \quad (3.48)$$

which no longer depends on \hat{q} . For $n = 0$ we simply get

$$\partial_k T_{s,k}^0(\bar{q}) = \frac{i^{-s}}{2} \text{Tr} ([T_{s,k}^2(\bar{q}) + R_k(\bar{q})]^{-1} [\partial_k R_k(\bar{q})]) \quad (3.49)$$

which is the equation most studied in the literature. For $n = 1$ we get

$$\partial_k T_{s,k}^1(\bar{q})(\cdot) = -\frac{i^{-s}}{2} \text{Tr} ([T_{s,k}^2(\bar{q}) + R_k(\bar{q})]^{-1} T_{s,k}^3(\bar{q})(\cdot) [T_{s,k}^2(\bar{q}) + R_k(\bar{q})]^{-1} [\partial_k R_k(\bar{q})]) \quad (3.50)$$

etc. where (\cdot) denotes the point at which one performs the functional derivatives. The *first truncation parameter* is therefore the number N of those first equations that one considers, $n = 0, 1, \dots, N - 1$.

Each of these equations can be expanded into a geometric series by splitting

$$T_{s,k}^2(\bar{q}) = D_{s,k}^2(\bar{q}) + N_{s,k}^2(\bar{q}) \quad (3.51)$$

where $D_{s,k}^2(\bar{q})$ ("diagonal part") can be easily treated by the heat kernel methods sketched below while $N_{s,k}^2(\bar{q})$ ("off-diagonal part") requires more work, and then expanding around $R'_{s,k}(\bar{q}) = R_k(\bar{q}) + D_{s,k}^2(\bar{q})$

$$[T_{s,k}^2(\bar{q}) + R_k(\bar{q})]^{-1} = \lim_{M \rightarrow \infty} R'_{s,k}(\bar{q})^{-1} \sum_{m=0}^M (-1)^m [N_{s,k}^2(\bar{q}) R'_{s,k}(\bar{q})^{-1}]^m \quad (3.52)$$

A *second truncation parameter* consists in the number M of terms $m = 0, 1, \dots, M - 1$ that one keeps in that expansion (for the n -th equations there are $n + 1$ such series and for each of them we keep all M terms).

Having decided on the numbers N, M one now makes an Ansatz for $\bar{\Gamma}_{s,k}(\hat{q}, \bar{q})$. The Ansatz is usually motivated by the action functional $S_s(q)$ that one started from. This is justified in the following sense: Recall that we have

$$\bar{Z}_s(f, \bar{q}) = \int [dh] J[\bar{q}+h] e^{i^s \langle f, h \rangle - S_s^t(\bar{q}+h)}, \quad \bar{C}_s(f, \bar{q}) = i^{-s} \ln(\bar{Z}_s(f, \bar{q})), \quad \bar{\Gamma}_s(\hat{q}, \bar{q}) = \text{extr}_f [\langle f, \hat{q} \rangle - \bar{C}_s(f, \bar{q})] \quad (3.53)$$

where we have formally integrated out the ghosts giving rise to the measure factor $J = \det(M)$, i.e. the functional determinant of the ghost matrix M and we have combined the classical action S_s and the gauge fixing action S^{gf} into $S_s^t = S_s + S^{gf}$.

Let $F = F(\hat{q}, \bar{q})$ be the solution of $\bar{C}_s^{(1)}(f, \bar{q}) = \hat{q}$ where we denote the n -th order functional derivatives of the functional F with respect to its argument by $F^{(n)}$. Then $\bar{\Gamma}_s^{(1)} = F$ and suppressing the dependence on \bar{q}

$$\begin{aligned} e^{-i^s \bar{\Gamma}_s(\hat{q})} &= \exp(i^s [\bar{C}_s(F) - \langle F, \hat{q} \rangle]) = \bar{Z}_s(F) e^{-i^s \langle F, \hat{q} \rangle} = \int [dh] J[\bar{q} + h] e^{i^s \langle F, h - \hat{q} \rangle - S_s^t(h + \bar{q})} \\ &= \int [dh] J[\bar{q} + h] e^{i^s \langle \bar{\Gamma}_s^{(1)}(\hat{q}), h - \hat{q} \rangle - S_s^t(\bar{q} + h)} \end{aligned} \quad (3.54)$$

which is a functional integro differential equation for $\bar{\Gamma}_s(\hat{q})$. We Taylor expand

$$S_s^t(\bar{q} + h) = \sum_{n=0}^{\infty} \frac{1}{n!} \langle S_s^{t(n)}(\bar{q} + \hat{q}), \otimes^n [h - \hat{q}] \rangle \quad (3.55)$$

and similar for J and shift the integration variable to obtain to second order

$$\begin{aligned} e^{-i^s [\bar{\Gamma}_s(\hat{q}, \bar{q}) - S_s^t(\bar{q} + \hat{q})]} &= \int [dh] e^{-i^s [\frac{1}{2} \langle S_s^{t(2)}(\bar{q} + \hat{q}), \otimes^2 h \rangle + \langle S_s^{t(1)}(\bar{q} + \hat{q}) - \bar{\Gamma}_s^{(1)}(\hat{q}, \bar{q}), h \rangle]} \times \\ &\quad \{ J[\bar{q} + \hat{q}] + \langle J^{(1)}(\bar{q} + \hat{q}), h \rangle + \frac{1}{2} \langle J^{(2)}(\bar{q} + \hat{q}), h \otimes h \rangle \} \end{aligned} \quad (3.56)$$

The Gaussian integrals can be performed. It yields an iteration for $\bar{\Gamma}_s - S_s$. If one truncates the iteration at the lowest order (1-loop approximation) one obtains

$$\bar{\Gamma}_s(\hat{q}, \bar{q}) = S_s^t(\bar{q} + \hat{q}) + \frac{1}{2 i^s} \text{Tr}(\ln(S_s^{t(2)}(\bar{q} + \hat{q}) - \frac{1}{i^s} \text{Tr}(\ln(J(\bar{q} + \hat{q})))) \quad (3.57)$$

Thus a well motivated starting Ansatz for the EAA is always the classical action plus gauge fixing action plus the above additional logarithmic term with argument split into background \bar{q} and argument \hat{q} . Note that for $s = 1$ the EAA, like the usual effective action, is necessarily complex valued rather than real valued.

The ingredients $S_s(q), S_{gf}(q), \text{Tr}(\ln(M))$ can be decomposed into several terms $g_\alpha S_{s,\alpha}(q)$ differing by the order of the derivatives of the field q each of which comes with its own coupling constant g_α . We make these couplings k dependent by hand, $g_\alpha \rightarrow g_\alpha(k)$ and write $S_{s,\alpha}(\hat{q}, \bar{q}) := S_{s,\alpha}(\bar{q} + \hat{q})$ by expansion where we use the covariant differential defined by \bar{q} . Accordingly we now step with

$$T_{s,k}^n(\bar{q}) := \sum_{\alpha=1}^A g_\alpha(k) S_{s,\alpha}^n(\bar{q}), \quad S_{s,\alpha}^n(\bar{q}) = \left[\frac{\delta^n S_{s,\alpha}(\hat{q}, \bar{q})}{\otimes^n \delta \hat{q}} \right]_{\hat{q}=0} \quad (3.58)$$

into the above hierarchy. Thus a *third truncation parameter* consists in the number A of terms in the above Ansatz (assuming they can be enumerated in some way).

Next one must isolate on the r.h.s. of the Wetterich equation truncated at N, M as detailed above the contributions $S_{s,\alpha}^n(\bar{q})$ and compare coefficients. For this isolation of terms we require the heat kernel techniques as detailed below. The reason for this is that the l.h.s., when written in terms of the covariant differential ∇ compatible with \bar{q} , generates distributions which are of the form of polynomials of ∇ applied to products of δ

distributions which generates curvature polynomials. The heat kernel technique is an elegant method for rewriting also the r.h.s. in terms of such distributions.

Some terms that appear on the r.h.s. of the Wetterich equation are not of the form $S_{s,\alpha}^n(\bar{q})$, $\alpha \leq A$ (they correspond to some $\alpha > A$) and thus cannot be compared to the l.h.s. These are discarded by hand. One must tune A to N, M such that after discarding those unmatchable terms one gets as many independent equations as we have couplings, that is, A . After this, one gets a closed, finite system of equations $\partial_k g(k) = \beta(g(k))$ which is then treated along the logic laid out in the previous subsection.

One can then improve the analysis and increase the numbers N, M, A . A systematic way to do this is concisely described in [16, 1]. The hope is of course that the fixed points and (finite!) number of relevant couplings stabilises beyond certain values of N, A, M .

3.9 Comparing coefficients: Sygne world function, Schrödinger kernel, heat kernel

The material contained in this subsection is mostly well known in the Euclidian setting, we include it here because it allows to pin point differences with the Lorentzian version and to make this review section self contained.

3.9.1 Heat kernel: definition

Let (M, m) be an n -dimensional manifold of Lorentzian ($s = 1$) or Euclidian ($s = 0$) signature, ∇ the Levi Civita covariant differential determined by m and $\Delta := m^{\mu\nu} \nabla_\mu \nabla_\nu$ the Laplacian (for $s = 1$ it is better called the d'Alembertian). The heat kernel of Δ (for $s = 1$ it is better called the Schrödinger kernel) is the solution to the initial value problem

$$\left[i^{-s} \frac{d}{dt} - \Delta_x \right] H_t^s(x, y) = 0, \quad H_0(x, y) = \delta(x, y) \quad (3.59)$$

In flat space $m_{\mu\nu} = \eta_{\mu\nu}$ (Minkowski metric $s = 1$) or $m_{\mu\nu} = \delta_{\mu\nu}$ (Euclidian metric $s = 0$) this equation can be explicitly solved by Fourier transformation

$$H_t(x, y) = H_t(x - y) = e^{i^s t \Delta_x} \delta(x, y) = \int_{\mathbb{R}^n} \frac{d^n l}{[2\pi]^n} e^{-i k_\mu (x-y)^\mu} e^{-i^s t m^{\mu\nu} k_\mu k_\nu} \quad (3.60)$$

Using the elementary integrals

$$\int_{\mathbb{R}} du e^{-a u^2} = \sqrt{\frac{\pi}{a}} \quad \forall a > 0; \quad \int_{\mathbb{R}} du e^{i a u^2} = \sqrt{\frac{\pi}{|a|}} e^{i \frac{\pi}{4} \text{sgn}(a)} \quad \forall a \in \mathbb{R} - \{0\} \quad (3.61)$$

one finds

$$H_t^s(x, y) = [4\pi|t|]^{-n/2} e^{\frac{i^s}{4t} m_{\mu\nu} (x-y)^\mu (x-y)^\nu} e^{i^s \frac{\pi}{4} \text{sgn}(t)(2-n)} \quad (3.62)$$

where for $s = 0$ only $t \geq 0$ is allowed. For $s = 1$ this is a one parameter group of unitary operators, for $s = 0$ a one parameter semi-group of contraction operators.

Irrespective of signature one observes that the x, y dependence is determined by the square of the geodesic distance $m(x - y, x - y)$ between the points x, y . It is thus reasonable to expect that at least for sufficiently close points x, y in a general spacetime the heat kernel is also largely determined by the square of the geodesic distance.

3.9.2 Sygne world function

In flat spacetime the geodesic between any two points is uniquely given by the straight line through them. In curved spacetime the geodesics through two points are unique only within a convex normal neighbourhood. Consider then two such points x, y in (M, m) and let $\gamma_{x,y}(s)$ be the unique geodesic with starting and end point $\gamma_{x,y}(0) = x$, $\gamma_{x,y}(1) = y$ respectively (we may assume that the convex neighbourhood is contained in a chart, otherwise restrict the neighbourhood). We know that the geodesic is the solution of the Euler Lagrange equations following from variation of the action

$$S[\gamma] := \int_0^1 ds w_\gamma(s), \quad w_\gamma(s) = \sqrt{|m_{\mu\nu}(\gamma(s)) \dot{\gamma}^\mu(s) \dot{\gamma}^\nu(s)|} \quad (3.63)$$

subject to the boundary condition $\gamma(0) = x$, $\gamma(1) = y$. When evaluated on the solution $\gamma_{x,y}$ (3.63) becomes the Hamilton-Jacobi function $S(x, y)$ that equals the absolute value of the geodesic distance. Using the Euler Lagrange equations and the boundary conditions one finds

$$\partial_{y^\mu} S(x, y) = \delta \frac{m_{\mu\nu}(\gamma_{x,y}(1)) \dot{\gamma}_{x,y}^\nu(1)}{w_{\gamma_{x,y}}(1)} \quad (3.64)$$

where $\delta = 1$ for $s = 0$ and $\delta = \pm 1$ if the geodesic is spacelike or timelike respectively when $s = 1$. When the geodesic is null then $S(x, y) = 0$ of course and the r.h.s. of (3.64) is to be replaced by zero. Thus one finds the Hamilton-Jacobi equation (replace r.h.s. by zero in the null case).

$$m^{\mu\nu}(y) S_{,y^\mu}(x, y) S_{,y^\nu}(x, y) = \delta \quad (3.65)$$

We now define the *Synge world function*

$$\sigma(x, y) := \delta \frac{1}{2} S(x, y)^2 \quad (3.66)$$

as the *signed* half of the square of the geodesic distance. Note that in flat space $\sigma = \delta |m(x - y, x - y)|/2 = m(x - y, x - y)/2$ is the function on which heat kernel exponent depends. Then we obtain the *master equations*

$$m^{\mu\nu}(y) [\nabla_\mu^y \sigma(x, y)] [\nabla_\nu^y \sigma(x, y)] = 2\sigma(x, y), \quad \sigma(x, x) = 0 \quad (3.67)$$

Remarkably, this equation, and all identities that follow from it, are completely insensitive to the signature and in the Lorentzian signature case holds for all three types of geodesics. One shows that (3.67) also holds if we replace $\nabla^y, m(y)$ by $\nabla^x, m(x)$.

The Synge world function is the simplest example of a bi-scalar. In general a bi-tensor $T(x, y)$ of type $(a, b), (a', b')$ transforms as a tensor of type (a, b) with respect to diffeomorphisms of x and as a tensor of type (a', b') with respect to diffeomorphisms of y . The coincidence limit of a bi-tensor is denoted as $\bar{T}(x) = T(x, x)$ and is a tensor of type $(a + a', b + b')$. In the Euclidian case the coincidence is unambiguously defined, for the Lorentzian case we take $y \rightarrow x$ along a spacelike path for definiteness. For our purposes it will be sufficient to consider bi-tensors of type $(a, b), (0, 0)$ i.e. it transforms as a scalar w.r.t. y .

We have seen above that $\nabla_\mu^x \sigma(x, y) = \propto S(x, y) \partial_\mu^x S(x, y)$ is either zero for a null geodesic or $\nabla_\mu^x S(x, y)$ is a (timelike or spacelike) unit vector. Therefore

$$\overline{\nabla_\mu^x \sigma(x, \cdot)} = 0 \quad (3.68)$$

We obtain complete information about the coincidence limits of all covariant derivatives of σ

$$s_{\mu_1 \dots \mu_n}(x) := \overline{\nabla_{\mu_1}^x \dots \nabla_{\mu_n}^x \sigma(x, \cdot)} \quad (3.69)$$

as follows: We compute successively the covariant derivatives of $\sigma(x, y)$ of order $n = 1, 2, \dots$ w.r.t. x and obtain relations from them using the master equations. The n -th derivative relation depends on the r.h.s. on covariant derivatives of σ up to order $n + 1$ where that $(n + 1)$ -th derivative is multiplied with a first derivative. Assuming inductively that the coincidence limit of the $(n + 1)$ -th derivative is regular one obtains an identity for the coincidence limits of the n -th derivative due to (3.68).

For instance (we drop the arguments x, y from now on and understand that all derivatives are taken at x at fixed $y \neq x$ and all contractions and curvature tensors are evaluated at x)

$$\begin{aligned} \nabla_\mu \sigma &= m^{\alpha\beta} [\nabla_\mu \nabla_\alpha \sigma] [\nabla_\beta \sigma] \\ \nabla_\mu \nabla_\nu \sigma &= m^{\alpha\beta} \{ [\nabla_\mu \nabla_\nu \nabla_\alpha \sigma] [\nabla_\beta \sigma] + [\nabla_\nu \nabla_\alpha \sigma] [\nabla_\mu \nabla_\beta \sigma] \} \end{aligned} \quad (3.70)$$

Assuming $s_{\mu\nu\rho}$ to be regular we find from the second relation that

$$s_{\mu\nu} = m^{\alpha\beta} s_{\mu\alpha} s_{\nu\beta} \quad (3.71)$$

which means that $s_{\mu}{}^{\nu}$ is a projection. Since σ for sufficiently close x, y approaches the flat space expression in suitable coordinates, the projection has maximal rank, therefore

$$s_{\mu\nu} = m_{\mu\nu} \quad (3.72)$$

Continuing like this one finds

$$s_{\mu\nu\rho} = 0, \quad s_{\mu\nu\rho\sigma} = \frac{2}{3} R_{\mu(\rho\sigma)\nu} \quad (3.73)$$

etc. Thus we can consider the coincidence limits (3.69) as known tensors constructed from m .

Next we consider an arbitrary bi-tensor of type $(0, b), (0, 0)$ (we can use $m(x)$ to reduce the case $(b - a, a), (0, 0)$; $b \geq a$ to this case). We are interested in the coincidence limit of its covariant derivatives at x . To that end we expand it with respect to its y dependence which due to (3.68) is equivalent to an expansion in $\nabla\sigma$

$$T_{\mu_1 \dots \mu_b}(x, y) = \sum_{l=0}^{\infty} \frac{1}{l!} [T^n]_{\mu_1 \dots \mu_b}^{\nu_1 \dots \nu_l}(x) \prod_{k=1}^l [\nabla_{\nu_k}^x \sigma(x, y)] \quad (3.74)$$

where the coefficients are automatically completely symmetric in $\nu_1 \dots \nu_l$. Then one finds using the above results for the coincidence limit of the σ derivatives

$$\overline{T_{\mu_1 \dots \mu_b}} = [T^0]_{\mu_1 \dots \mu_b}, \quad \overline{\nabla_{\nu} T_{\mu_1 \dots \mu_b}} = \nabla_{\nu} [T^0]_{\mu_1 \dots \mu_b} + [T^1]_{\mu_1 \dots \mu_b}{}^{\rho} m_{\rho\nu} \quad (3.75)$$

etc., i.e. we can relate all expansion coefficients in (3.74) to coincidence limits of covariant derivatives of T .

3.9.3 Heat kernel: computation

The Ansatz for the heat kernel in general (M, m) is therefore well motivated to be

$$H_t^s(x, y) = [4\pi|t|]^{-n/2} e^{\frac{i^s}{2t}\sigma(x, y)} e^{i^s \frac{\pi}{4} \text{sgn}(t)(2-n)} \Omega_t(x, y) \quad (3.76)$$

which contains a correction factor $\Omega_t(x, y)$ that captures the curvature effects. We know already

$$\overline{\Omega_{t=0}} = 1 \quad (3.77)$$

as the other factor in (3.76) reduces to $\delta(x, y)$ at $t = 0$. Once we have determined Ω_t to sufficient accuracy we know the heat kernel to sufficient accuracy. To determine Ω_t we plug (3.76) into the heat equation. We obtain for $t \neq 0$ using (3.67) (all derivatives and contractions at x)

$$\frac{[\Delta\sigma] - n}{2t} \Omega_t + \partial_t \Omega_t + \frac{1}{t} m^{\mu\nu} [\nabla_{\mu} \Omega_t] [\nabla_{\nu} \sigma] - i^s [\Delta \Omega_t] = 0 \quad (3.78)$$

We Taylor expand with respect to the heat time t

$$\Omega_t(x, y) = \sum_{k=0}^{\infty} (i^s t)^k \Omega_k(x, y) \quad (3.79)$$

and obtain for $k = 0, 1, \dots$

$$\left(\frac{[\Delta\sigma] - n}{2} + k \right) \Omega_k + m^{\mu\nu} [\nabla_{\mu} \Omega_k] [\nabla_{\nu} \sigma] - [\Delta \Omega_{k-1}] = 0 \quad (3.80)$$

with the convention $\Omega_{-1}(x, y) = 0$. Also $\Omega_{t=0}(x, x) = \Omega_{k=0}(x, x) = 1 = \overline{\Omega_0}$. Equation (3.80) is independent of signature and as all coefficients are real valued we can pick Ω_k real valued.

We can explicitly determine the Ω_k using the Taylor expansion (3.75) together with (3.80)

$$\Omega_k(x, y) = \sum_l \frac{1}{l!} [\Omega_{k,l}]^{\mu_1 \dots \mu_l}(x) [\nabla_{\mu_1}^x \sigma(x, y)] \dots [\nabla_{\mu_l}^x \sigma(x, y)] \quad (3.81)$$

where $\Omega_{k,l}^{\mu_1 \dots \mu_l}(x)$ is determined in terms of the coincidence limits of the $\nabla_{\mu_1} \dots \nabla_{\mu_r} \Omega_k$, $0 \leq r \leq l$ via (3.75). To obtain the coincidence limits of the $\nabla^l \Omega_k$ we take the ∇^l derivative of (3.80) and then take the coincidence limit. This involves only $\nabla^l \Omega_k$ and $\nabla^{l+2} \Omega_{k-1}$ because the $\nabla^{l+1} \Omega_k$ term has a coefficient $\nabla \sigma$ which vanishes in the coincidence limit. Thus, as $\Omega_{-1} \equiv 0$ we can compute all $\Omega_{k,l}$ to any desired order.

We exhibit this for $l = 0, 1$ and general k :

$$[\Omega_{k,0}] = \overline{\Omega_k}, \quad [\Omega_{k,1}]^\mu = \overline{\nabla^\mu \Omega_k} - \nabla_\mu \overline{\Omega_k} \quad (3.82)$$

The coincidence limit of (3.80) is

$$k \overline{\Omega_k} - \overline{\Delta \Omega_{k-1}} = 0 \quad (3.83)$$

and the first derivative of (3.80) is

$$\left[k - \frac{n}{2} + \frac{1}{2} \Delta \sigma \right] [\nabla_\mu \Omega_k] + \frac{1}{2} [\nabla_\mu \Delta \sigma] \Omega_k + [\nabla_\mu \nabla_\nu \Omega_k] [\nabla^\nu \sigma] + [\nabla_\nu \Omega_k] [\nabla_\mu \nabla^\nu \sigma] - \nabla_\mu \Delta \Omega_{k-1} = 0 \quad (3.84)$$

whose coincidence limit is

$$(k+1) \overline{\nabla_\mu \Omega_k} - \overline{\nabla_\mu \Delta \Omega_{k-1}} = 0 \quad (3.85)$$

For $k = 0$ (3.83) is identically satisfied while (3.85) gives $\overline{\nabla_\mu \Omega_1} = 0$. Thus by (3.82) $[\Omega_{0,0}] = 1$, $[\Omega_{0,1}]^\mu = 0$ and taking this one step further one finds $\overline{\nabla_\mu \nabla_\nu \Omega_0} = \frac{1}{6} R_{\mu\nu}$ (Ricci tensor). Thus

$$\Omega_0 = 1 + \frac{1}{12} R^{\mu\nu} \sigma_{,\mu} \sigma_{,\nu} \quad (3.86)$$

up to second order. Inserted into (3.83) for $k = 1$ gives $\overline{\Omega_1} = \frac{1}{6} R$ (Ricci scalar). To compute (3.85) for $k = 1$ requires to know $\nabla^3 \Omega_0$ etc.

Concluding, we can consider the heat kernel to be known to any desired order k, l in $t, [\nabla \sigma]$. Also arbitrary covariant derivatives of the heat kernel can be obtained from again computing derivatives of (3.80) and taking coincidence limits as well as the known coincidence limits of σ .

3.9.4 Functions of the heat kernel and traces

Let for, $s = 0$, $f(z)$ be a complex valued function defined for $\Re(z) > 0$ and $\hat{f}(t)$ its pre-image under the Laplace transform

$$f(z) = \int_{\mathbb{R}^+} dt e^{-tz} \hat{f}(t), \quad \hat{f}(t) = [2\pi i]^{-1} \int_{x+i\mathbb{R}} dz e^{tz} f(z); \quad x > 0 \quad (3.87)$$

Let, for $s = 1$, $f(z)$ be a complex valued function defined on the real line and $\hat{f}(t)$ its pre-image under the Fourier transform

$$f(z) = \int_{\mathbb{R}} dt e^{-itz} \hat{f}(t), \quad \hat{f}(t) = [2\pi]^{-1} \int_{\mathbb{R}} dz e^{itz} f(z); \quad x > 0 \quad (3.88)$$

For a function $f = f(-\Delta)$ we may use the spectral theorem

$$f(-\Delta) = \begin{cases} \int_0^\infty dt \hat{f}(t) e^{t\Delta} & s = 0 \\ \int_{-\infty}^\infty \hat{f}(t) dt e^{it\Delta} & s = 1 \end{cases} \quad (3.89)$$

Then the trace of $f(-\Delta)$ with respect to the volume form of m is given by

$$\int dt \int d^n x |\det(m)|^{1/2} H_t(x, x) \quad (3.90)$$

which demonstrates why we are interested in the coincidence limit and justifies the assumption about the convex normal neighbourhood above. Obviously $H_t(x, x) \propto \overline{\Omega_t}(x)$ which is an expansion in terms of curvature invariants.

As laid out in the previous subsection, on the r.h.s of the Wetterich equation what we are interested in are expressions of the form (recall (3.51), (3.52))

$$\text{Tr}(P_0(-\Delta)) Q_1(\nabla) P_2(-\Delta) \dots Q_m(\nabla) P_m(-\Delta) \quad (3.91)$$

where $m \leq M - 1$ and where the P_k just depend on Δ ("diagonal operators") while the Q_k may depend on all combinations of ∇ . We may write these expressions as

$$\begin{aligned} & \int d^{m+1}t \hat{P}_0(t_0) \dots \hat{P}_m(t_m) \text{Tr}(e^{i^s t_0 \Delta} Q_1(\nabla) e^{i^s t_1 \Delta} \dots Q_m(\nabla) e^{i^s t_m \Delta}) \\ &= \int d^{m+1}t \hat{P}_0(t_0) \dots \hat{P}_m(t_m) \text{Tr}([e^{i^s r_1 \Delta} Q_1(\nabla) e^{-i^s r_1 \Delta}] e^{i^s r_2 \Delta} Q_1(\nabla) e^{-i^s r_2 \Delta} \dots [e^{i^s r_m \Delta} Q_m(\nabla) e^{-i^s r_m \Delta}] e^{i^s r_{m+1} \Delta}) \end{aligned} \quad (3.92)$$

where $r_k = t_0 + \dots t_{k-1}$, $k = 1, \dots, m + 1$. Then we may use

$$e^{i^s r \Delta} Q(\nabla) e^{-i^s \Delta} = \sum_{n=0}^{\infty} \frac{(i^s r)^n}{n!} [\Delta, Q(\nabla)]_{(n)}, [A, B]_{(0)} := B, [A, B]_{(n+1)} := [A, [A, B]_{(n)}] \quad (3.93)$$

and discard in the sums over n all terms above a certain order order that is dictated by the chosen values of N, M, A . Then working out the multiple commutators we end up, inside the integrals over t_0, \dots, t_m with an expression of the form

$$\text{Tr}(Q(\nabla) H_r(\Delta)) \quad (3.94)$$

which can be evaluated by the tools of the previous subsection. Also here we discard terms above a certain order in the expression of $\overline{\Omega}_r$ above a certain order in terms of k, l as dictated by N, M, A .

3.9.5 Performing the Laplace or Fourier integrals

It remains to perform the integrals over the heat kernel times t_0, \dots, t_m ; $m \leq M - 1$. Their convergence is of course much determined by the properties of the Laplace or Fourier transform of R_k . As an example consider the truncation $N = 1$ for the case $N_{s,k}^2(\bar{q}) = 0$ and $D_{s,k}^2(\bar{q}) = \Delta + R_k(\Delta)$ then it suffices to consider the case $M = 1$ and we need

$$\text{Tr}([\Delta + R_k(\Delta)]^{-1} [\partial_k R_k(\Delta)]) = \int dt \hat{f}_k(t) \text{Tr}(e^{i^s t \Delta}) = \int dt \hat{f}_k(t) [4\pi |t|]^{-n/2} e^{i^s \text{sgn}(t) \pi/4} \int d^n x \left[\sum_{k'=0}^{\infty} (i^s t)^{k'} \overline{\Omega}_{k'}(x) \right] \quad (3.95)$$

i.e. we require the integrals for $k' = 0, 1, 2, \dots$

$$q_k(n, k') := \int \hat{f}_k(t) t^{k'} |t|^{-n/2} e^{i^s \text{sgn}(t) \frac{\pi}{4}} \quad (3.96)$$

over \mathbb{R}_+ ($s = 0$) and \mathbb{R} ($s = 1$) respectively where $\hat{f}_k(t)$ is the *inverse* of the Laplace respectively Fourier transform of

$$f_k(x) = \frac{\partial_k R_k(x)}{x + R_k(x)} \quad (3.97)$$

Strictly speaking, what we are given is $f_k(x)$ and we assume that there exists $\hat{f}_k(t)$ such that $f_k(x) = \int_{\mathbb{R}_+} dt e^{-xt} \hat{f}_k(t)$ for $s = 0$ and $f_k(x) = \int_{\mathbb{R}} dt e^{ixt} \hat{f}_k(t)$ for $s = 1$ respectively. This formula can be inverted for $\hat{f}_k(t)$ for $s = 1$ on the space of Schwartz functions and tempered distributions while for $s = 0$ the existence of \hat{f}_k is not granted given f_k (there exists an inverse if instead \hat{f}_k is given). This fact has important consequences for some of the results quoted in the literature in particular in connection with the so called *optimised cut-off* [1]. As we show in appendix C, the optimised cut-off is not granted (and is likely not) to lie in the image of the Laplace transform of a meaningful mathematical object (function, distribution, measure,...) which would mean that the formulas that one quotes for its q_k integrals do not hold and have to be revisited.

Whether the integrals (3.96) exist depends on the chosen shape of the suppressing function $R_k(x)$ which may depend on the chosen signature s . For the case $s = 1$ we may pick $f_k(x)$ as the the Fourier transform of \hat{f}_k where besides the already stated conditions 1.-4. we require \hat{f}_k to be smooth of rapid decrease at $t = 0, \pm\infty$. A controllable class of such functions and its integrals was studied recently in [28] and we showcase their usefulness in appendix D. Again one will discard terms above a maximal value of k' in agreement with the chosen values of M, N, A .

In conclusion for suitable choices of N, M, A , upon discarding, we may evaluate both the l.h.s. and the r.h.s. of the Wetterich equation in closed form producing a closed system of equations $\partial_k g(k) = \beta(g(k))$.

4 Contact between CQG and ASQG in a concrete model

In this section we combine the two frameworks and study a concrete model for Lorentzian GR coupled to some form of matter. In the first subsection we introduce the classical Lagrangian of the model and carry out the steps of section 2 to pass from the Hamiltonian formulation to the path integral. In the second subsection we release the asymptotic safety machinery to the model. To avoid confusion note that the meaning of the indices $a, b, c, \dots, j, k, l, \dots$ in this section is different from section 2: There they took infinite range labelling test smearing functions while here they take finite range $1, \dots, n-1$ labelling tensor components and field species respectively.

4.1 CQG treatment of the model

4.1.1 Classical analysis

The classical Lagrangian is given by

$$L = \frac{1}{G} |g|^{1/2} [R^{(n)}(g) - 2\Lambda] - \frac{1}{2} |g|^{1/2} g^{\mu\nu} S_{IJ} \phi_{,\mu}^I \phi_{,\nu}^J \quad (4.1)$$

where g is a Lorentzian signature metric on an n -dimensional manifold M diffeomorphic to $\mathbb{R} \times \Sigma$ and ϕ^I , $I = 0, \dots, n-1$ are n scalar fields, $\mu, \nu = 0, \dots, n-1$ are tensor indices. The real valued, constant matrix S_{IJ} is positive definite, G, Λ are Newton's and the cosmological constant respectively. Thus (4.1) is just the Einstein-Hilbert Lagrangian minimally coupled to n massless Klein-Gordon fields without potential. This model was also treated by the methods of LQG in [29]. To avoid discussions about boundary terms we assume that Σ is compact without boundary.

We note that since S is positive we find non-singular matrices s_J^I such that $M_{IJ} = \delta_{KL} s_L^K s_J^I$ and thus could replace S_{IJ} by δ_{IJ} by the field redefinition $s_J^I \phi^J \mapsto \phi^I$. We will however keep S_{IJ} general as it does not blow up the formalism. The above model can be considered as a dark matter model as the scalars couple only gravitationally, albeit not a very realistic one as there is no mass term.

The classical Hamiltonian formulation of this system is well known [30]. An arbitrary foliation of M , using time t and spatial coordinates x^a on Σ and as parametrised by lapse N and shift N^a functions $a, b, \dots = 1, \dots, n-1$ leads to canonical pairs $(P, N); (P_a, N^a); (p^{ab}, q_{ab}), (\pi_I, \phi^I)$ where P, P_a are constrained to vanish (primary constraints) while

$$p^{ab} = \sqrt{\det(q)} [q^{ac} q^{bd} - q^{ab} q^{cd}] k_{cd}, \quad k_{ab} = \frac{1}{2N} [\dot{q}_{ab} - [L_{\vec{N}} q]_{ab}], \quad \pi_I = \sqrt{\det(q)} S_{IJ} [L_n \phi^J] \quad (4.2)$$

where L_u denotes the Lie derivative with respect to u and where $n = N^{-1}(\partial_t - \vec{N})$, $\vec{N} = N^a \partial_{x^a}$. Here q_{ab} is the intrinsic spatial metric of Euclidian signature on the leaves of the foliation and k_{ab} their extrinsic curvature and n is a unit timelike normal to the leaves while \vec{N} is tangential.

The secondary constraints are (all spatial indices are moved with q_{ab} , q^{ab} ; $q_{ac} q^{cb} = \delta_a^b$ and $Q := [\det(q)]^{1/2}$)

$$\begin{aligned} C_a &= -2D_b p_a^b + \pi_I \phi_{,a}^I; & C &= C^g + C^s; \\ C^g &= Q^{-1} [p^{ab} p_{ab} - \frac{1}{n-2} (p^a_a)^2] - Q [R^{(n-1)}(q) - \Lambda], & C^s &= \frac{1}{2} [Q^{-1} S^{IJ} \pi_I \pi_J + Q S_{IJ} q^{ab} \phi_{,a}^I \phi_{,b}^J] \end{aligned} \quad (4.3)$$

known as spatial diffeomorphism and Hamiltonian constraint respectively. Here D is the Levi-Civita differential of q (note that P^{ab} is a spatial tensor density of weight one). It is straightforward but tedious to check that the $2n$ constraints are first class, i.e. their mutual Poisson brackets vanish when all constraints hold.

The constrained Hamiltonian density is

$$H = v P + v^a P_a + N^a C_a + N C \quad (4.4)$$

where v, v^a are undetermined Lagrange multipliers as the Legendre transform does not determine the velocities of N, N^a . This Hamiltonian is a sum of constraints and its Hamiltonian flow is to be interpreted as gauge transformations.

We now consider the reduced phase space of the system. We assume that the n scalar fields serve as a material reference system in the sense that

$$\det([\partial\phi/\partial(t, x)]) \neq 0 \quad (4.5)$$

so that ϕ in fact defines a diffeomorphism. It is then natural to adopt the gauge fixing condition

$$G^I(t, x) = \phi^I(t, x) - k^I(t, x), G^0 = N - c(t, x), G^a = N^a - c^a(t, x) \quad (4.6)$$

where k^I, c, c^a are coordinate functions independent of the phase space variables subject to the constraint $\det(\partial k / \partial(t, x)) \neq 0$. That is, we consider ϕ^I, N, N^a and their conjugate momenta as gauge degrees of freedom while q_{ab}, p^{ab} are the true degrees of freedom. The correspondence with the general theory laid out in section 2 is that (x, y) are given by $(P, P_a, \phi^I), (N, N^a, \pi_I)$ and (p, q) by (p_{ab}, q^{ab}) . The model thus has $\frac{1}{2}n(n+1) + n - 2n = \frac{1}{2}n(n-1)$ physical degrees of freedom, i.e. n more than in vacuum due to the presence of the n scalar fields. That we choose the true degrees of freedom purely in terms of q_{ab} rather than say the $\frac{1}{2}n(n+1) - 2n = \frac{1}{2}n(n-3)$ gravitational wave polarisations and n scalar fields is because the description of the reduced phase space is much simpler then, as we can solve the constraints algebraically (otherwise we would need to solve PDE's). This is similar to the Higgs mechanism where rather than 2 polarisations for the W_+, W_-, Z boson we have 3 because 3 of 4 degrees of freedom of the complex Higgs doublet are gauge fixed. The additional polarisation degrees of freedom could be called Goldstone bosons.

In detail, let $j, k, \dots = 1, \dots, n-1$ then we may solve algebraically

$$\begin{aligned} \hat{C}_j &:= \pi_j + h_j, \quad \hat{C} = \pi_0 + h_0 & (4.7) \\ h_j &= \phi_j^a [-2D_b P_a^b + \pi_0 \phi_{,a}^0] =: h_j^g + \pi_0 \phi_j^0; \quad \phi_{,a}^j \phi_j^b = \delta_a^b \\ h_0 &= \frac{B}{A} \pm \sqrt{-\frac{Q}{A} \left[\frac{B^2}{A} + C_g + \frac{1}{2}(Q^{-1} S^{jk} h_j^g h_k^g + Q S_{IJ} q^{ab} \phi_{,a}^I \phi_{,b}^J) \right]} \\ A &= S^{00} - 2S^{0j} \phi_j^0 + S^{jk} \phi_j^0 \phi_k^0 \\ B &= (S^{0j} + S^{jk} \phi_k^0) h_j^g \end{aligned} \quad (4.8)$$

where the solution $\pi_0 = -h_0$ is to be inserted in the first line.

Note that as detailed in section 2 there are two possible roots for π_0 . Note also that the solutions h_I drastically simplify at the gauge cut $\phi^I = k^I$ when $k_{,a}^0 = 0$ so that $k_j^0 = k_j^a k_{,a}^0 = 0$ and if $S^{0j} = 0$ i.e. $S_{0j} = 0$ i.e. S_{IJ} is block diagonal in which case $B = 0, A = S^{00} = S_{00}^{-1}$. In fact, in order to apply the results of section 2 we must adopt this choice because in this case 1. the quadratic constraints are of the type $C_j = \sum_k P_j^k \hat{C}_k^+ C_k^-$ with $\hat{C}_k^\sigma = y_k + h_k^\sigma$, 2. $h_k^\pm = \pm h_k$ and 3. h_k depends only quadratically on x^j .

The reduced Hamiltonian density is according to the general theory of section 2 given by

$$H_*(q, p; t) = \dot{k}^I(t) h_I(q, p; \phi = k(t)) \quad (4.9)$$

It is not explicitly time dependent iff \dot{k} is a constant and the particular way in which k appears in h_I loses its time dependence. Since the square root in h_0 depends on the term $S_{IJ} q^{ab} k_{,a}^I k_{,b}^J$ we require that $k_{,a}^I$ is time independent. The terms A, B in h_0 are then also time independent since $k_j^0 = k_j^a k_{,a}^0$ and k_j^a is the inverse of $k_{,a}^j$. Accordingly the gauge fixing condition is of the form $k^I(t, x) = k_0^I t + k_1^I(x)$ where k_0^I are n constants and $k_1^I(x)$ are functions of the spatial coordinates only, subject to the condition that

$$\det(\partial k / \partial(t, x)) = \frac{1}{(n-1)!} k_0^{I_0} \epsilon_{I_0 \dots I_{n-1}} \epsilon^{a_1 \dots a_{n-1}} \prod_{l=1}^{n-1} k_{1,a_l}^{I_l} \neq 0 \quad (4.10)$$

A particularly simple choice is $k_0^I = \kappa_0 \delta_0^I$ in which case (4.10) reduces to the condition that $\det(\partial k_1 / \partial x) \neq 0$ which is for example achieved for $k_1^j(x) = [\kappa_1]_a^j x^a$ and κ_1 is an invertible matrix. In this case

$$S_{IJ} q^{ab} k_{,a}^I k_{,b}^J = q^{ab} S_{ab}, \quad S_{ab} = S_{jk} [\kappa_1]_a^j [\kappa_1]_b^k \quad (4.11)$$

Note that we could keep H_* free of explicit time dependence in this gauge if we would add a mass and potential term which only depend on ϕ^j but not on ϕ^0 .

At this point we have declared q_{ab}, p^{ab} as the independent physical degrees of freedom of the model. However, to define an entire spacetime metric it is not enough to know the time evolution of $g_{ab} = q_{ab}$, we also need to

know $g_{tt} = -N^2 + q_{ab}N^b$ and $g_{ta} = q_{ab}N^b$ to have access to the full spacetime metric, i.e. we need to know N, N^a as functions of q_{ab}, p^{ab} . This dependence is provided by the stability condition on the gauge condition, i.e.

$$[\partial_t G^I + \{C(N), G^I\}]_{\phi=k, \pi=-h} = -\dot{k}^I + N^a k^I_a - \frac{N}{Q} S^{IJ} h_J = 0 \quad (4.12)$$

which can be solved for N, N^a . For the choice discussed above which avoids explicit time dependence of the reduced Hamiltonian one finds

$$N = -\kappa_0 S_{00} \frac{Q}{h_0}, \quad N^a = -\kappa_0 S_{00} ([\kappa_1]^{-1})_j^a S^{jk} \frac{k_k}{h_0} \quad (4.13)$$

4.1.2 Canonical quantisation

In the canonical quantisation of this model one now considers the Weyl algebra generated by

$$W(f, g) = \exp(i \int_{\Sigma} d^{n-1}x [f^{ab} q_{ab} + g_{ab} p^{ab}]) \quad (4.14)$$

and representations thereof which allow a quantisation of H_* . This is quite a challenging task due to the square root involved and all the other non-linearities that are involved in C^g which depends on inverse powers of Q . To make this task as simple as possible, one will resort to the simplest gauge condition $k^0 = \kappa_0 t, k^j = [\kappa_1]_a^j x^a$ and block diagonal S so that for the choice of the positive root

$$H_*(q, p) = \frac{\kappa_0}{S_{00}^{1/2}} \sqrt{-Q [C_g + \frac{1}{2}(Q^{-1} S^{ab} h_a^g h_b^g + Q S_{ab} q^{ab})]} \quad (4.15)$$

where we have used

$$S^{jk} h_j^g h_k^g = S^{ab} h_a^g h_b^g, \quad h_a^g = -2D_b p_a^b \quad (4.16)$$

with $S^{ac} S_{cb} = \delta_b^a$ and S_{ab} is just a constant matrix. One possibility to proceed is to notice that the expression $-E$ under the square root is classically constrained to be positive and therefore classically

$$H_* = \frac{\kappa_0}{S_{00}^{1/2}} \sqrt{|-E|} = \frac{\kappa_0}{S_{00}^{1/2}} [E^2]^{1/4}, \quad E = Q [C_g + \frac{1}{2}(S^{ab} h_a^g h_b^g + Q S_{ab} q^{ab})] \quad (4.17)$$

which now at least frees us of making sure that the expression under the square root is positive after quantisation. The function E^2 can now be ordered symmetrically and one can implement this as an honest symmetric operator on a finite lattice (recall that σ was chosen compact) and then study its Hamiltonian renormalisation flow.

4.1.3 Path integral formulation

In the path integral approach outlined in section 2 one can now formally obtain the generating functional of Feynman ($s = 1$) or Schwinger ($s = 0$) functions

$$Z_s(f) = \int [dq] [dp] \overline{\Omega_0(q(t = -\infty))} \Omega_0(q(t = \infty)) e^{-\int d^n x [i p_{ab} \dot{q}^{ab} + (-1)^s i^s H_*(q, p)]} e^{i^s \int d^n x f^{ab} q_{ab}} \quad (4.18)$$

This formula is rigorous as long as one understands that the integrals involved are in fact finite Riemann sums, and accordingly $[dp], [dq]$ finite product Lebesgue measures, in a discretisation of the phase space on the compact spacetime $[-T, T] \times \sigma$ upon which $t = \pm\infty$ in the cyclic state is to be replaced by $t = \pm T$. As usual we keep a continuum notation but the steps that follow can only be strictly justified with that discretised understanding.

There are now two possibilities. Either one stays within this strictly reduced phase space and tries to integrate out the momenta p^{ab} . This is not possible directly because of the square root involved in $H_*(q, p)$. However, one can get rid of the square root using an exact integral transform based on a Lagrange multiplier field. Or one passes to an unreduced phase space path integral along the lines of section 2 which is another way to get rid of the square root. We will describe both methods below. For both methods we obtain a problem in the

case $s = 0$ which is specific to GR. The problem of the first approach is that while the argument of the square root involved in H_* is classically constrained to be positive, in the integral over all paths that condition cannot be maintained, the Hamiltonian is unbounded from below and the momentum integral diverges which is partly caused by the conformal mode. If one ignores the infinite factors that appear in carrying out the momentum integrals because they formally drop out in the non vacuum correlation functions one still faces the problem that the resulting expression is spatially rather non-local. We leave it for future investigations. The problem of the second approach is that while there is no convergence issue in integrating out the momenta, we face the problem already mentioned in section 2, namely that the ‘‘Euclidian’’ action becomes complex valued rather than real valued. In the second approach we are therefore unambiguously directed to consider the $s = 1$ formulation (Feynman functions rather than Schwinger functions). In both formulations, the configuration space measure receives non-trivial corrections to the ‘‘naive’’ Lebesgue measure.

Reduced configuration space path integral

The square root Hamiltonian reminds of the square root action for the relativistic particle or the Nambu-Goto action for closed bosonic string and one may recall [11] that in a lowest order saddle point approximation

$$\int d\lambda e^{\frac{\epsilon}{2}[\lambda f + \lambda^{-1}]} \propto e^{c\sqrt{f}} \quad (4.19)$$

(the infinite constant involved would drop out in $Z_s(f)/Z_s(0)$) would let us pass to the analog of the Polyakov action without square root. If (4.19) was exact one could now integrate out p as H_* depends only quadratically on p yielding a Gaussian integral. Unfortunately (4.19) is not exact. However, there exists a more complicated version of (4.19) which is based on lemma B.1 proved in the appendix and which to the best of our knowledge has not been reported before.

We write $H_* =: Q\sqrt{h}$ which displays h as a scalar with zero density weight while H_*, Q have density weight unity. Let $V_I := \epsilon_I Q_I$ be a discretisation of $\int_{c_I} d^4x Q$, the volume measured by Q of an n -cell c_I with centre p_I into which $M = [-T, T] \times \sigma$ is partitioned (the partition is finite) with ϵ_I its coordinate volume and $Q_I = Q(p_I)$. Then with our understanding of the symbolic continuum expression with $h_I = h(p_I)$

$$\int d^4x H = \sum_I V_I \sqrt{h_I} \quad (4.20)$$

By lemma B.1 we have with $d = V_I, z = h_I$

$$e^{(-1)^{1+s} i^s \sum_I V_I \sqrt{h_I}} = \prod_I \int_{\mathbb{R}_+} \frac{d\lambda_I}{\lambda_I^{3/2}} \prod_I \frac{e^{-i s \frac{\pi}{4} V_I}}{2\sqrt{\pi}} e^{(-1)^{1+s} i^s \sum_I [\lambda_I h_I + \frac{V_I^2}{4\lambda_I}]} \quad (4.21)$$

We substitute $\lambda_I \rightarrow \lambda_I V_I$ to obtain

$$e^{(-1)^{1+s} i^s \sum_I V_I \sqrt{h_I}} = \prod_I \int_{\mathbb{R}_+} \frac{d\lambda_I}{\lambda_I^{3/2}} \prod_I \frac{e^{-i s \frac{\pi}{4} V_I^{1/2}}}{2\sqrt{\pi}} e^{(-1)^{1+s} i^s \sum_I V_I [\lambda_I h_I + \frac{1}{4\lambda_I}]} \quad (4.22)$$

Accordingly, dropping the power of $e^{-is\pi/4}/(2\sqrt{\pi})$ we have

$$Z_s(f) = \int [Q^{1/2} dq] [dp] [\lambda^{-3/2} d\lambda] \overline{\Omega_0(q(t = -\infty))} \Omega_0(q(t = \infty)) e^{-\int d^n x [i p_{ab} \dot{q}^{ab} + (-1)^s i^s (\lambda H(q,p) + \frac{Q}{4\lambda})]} e^{i^s \langle f, q \rangle} \quad (4.23)$$

where it is understood that one integrates over $\lambda \in \mathbb{R}_+$ $q_{ab}, p^{ab} \in \mathbb{R}$ and

$$H = | - [C_g + \frac{1}{2} (S^{ab} h_a^g h_b^g + Q S_{ab}; q^{ab})] | \quad (4.24)$$

The expression in (4.24) between the absolute values is constrained to be positive in the classical theory (this is possible because the gravitational contribution C_g is not bounded from below). This can be ensured in the path integral only if one does not integrate unconditionally over q_{ab}, p^{ab} . But then integrating out p_{ab} is not just reduced to Gaussian integrals and therefore not possible explicitly.

To explore (4.24) we assume without proof that the paths in phase for which one can drop the modulus dominate. This means that we have to perform the Gaussian integral

$$\int [dp] e^{-[i\langle p, \dot{q} \rangle - (-1)^s i^s \langle p, O(q, \lambda) p \rangle + \langle Q, \lambda(\Lambda - R + \frac{1}{2}S \cdot q) + \frac{1}{4\lambda} \rangle]} \quad (4.25)$$

where $O(q, \lambda)$ is the differential operator

$$[O(q) p]_{ab} := Q^{-1} G_{abcd}(q, \lambda) p^{cd} - 2D_{(a} q_{b)c} \frac{\lambda S^{cd}}{Q} q_{de} D_f p^{fe}, \quad G_{abcd} = \lambda(q_a(cq_d)b - \frac{1}{n-2} q_{ab}q_{cd}) \quad (4.26)$$

This differential operator is known to be indefinite with respect to the first term (conformal mode problem) and the second term, although it looks like minus a Laplacian, is likely not to repair this. This is the reason why $s = 1$ is preferred because oscillatory Gaussians $x \mapsto e^{ix^2}$ can be integrated. Completing the square we obtain after dropping (infinite) constant factors

$$Z_1(f) = \int [Q^{1/2} dq] [\lambda^{-3/2} d\lambda] \overline{\Omega_0(q(t = -\infty))} \Omega_0(q(t = \infty)) |\det[O(q, \lambda)]|^{-1/2} \times e^{-i[\frac{1}{4}\langle \dot{q}, O(q, \lambda)^{-1} \dot{q} \rangle - \langle \lambda Q, \Lambda - R + S \cdot q/2 + (2\lambda)^{-2} \rangle]} e^{i\langle f, q \rangle} \quad (4.27)$$

This expression has no resemblance any more with the naive expression (Lebesgue measure times exponential of the classical action). It has an exponential part but that part involves the inverse of the full differential operator O and not just the DeWitt metric term G_{abcd} and thus is spatially non-local. There is also a measure part which involves the determinant of O .

In principle one can now release the ASQG machinery on (4.27) where no gauge fixing term and no ghost term appear because we have reduced the phase space prior to quantisation, i.e. this would be the background formalism without gauge invariance where a background is introduced just for q_{ab} and not the full spacetime metric. There is no current for the Lagrange multiplier field λ because its whole purpose was to get rid of the square root and thus plays a role similar to the ghosts in presence of gauge invariance which are just there in order to write the Fadeev-Popov determinant factor as an exponential. One can see this also from the fact that no time derivatives of λ appear.

In order to solve the Wetterich equation one needs to make an Ansatz for the EAA which is usually the logarithm of the integrand of $Z(0)$. If we write this integrand as a measure $d\mu[q]$ by formally integrating out λ then this Ansatz would be $\bar{\Gamma}(\hat{q}, \bar{q}) = i \ln(\frac{d\mu[q]}{[dq]})_{q=\bar{q}+\hat{q}}$. Since it is practically not possible to carry out the integral over λ we instead artificially multiply the integrand by $\exp(i \langle \iota, \lambda \rangle)$ where ι is a current for λ thus obtaining $Z'_1(f, g)$ where of course we are only interested in $Z_1(f) = Z'_1(f, g = 0)$. We can now define in the usual way $\bar{Z}'_1(f, g; \bar{q})$, $\bar{C}'_1(f, g; \bar{q})$, $\bar{\Gamma}'_1(f, g; \bar{q})$ but while $\bar{Z}_1(f, \bar{q}) = \bar{Z}'_1(f, g = 0, \bar{q})$ and $\bar{C}_1(f, \bar{q}) = \bar{C}'_1(f, g = 0, \bar{q})$ we do not have $\bar{\Gamma}_1(\hat{q}, \bar{q}) = \bar{\Gamma}'_1(\hat{q}, \hat{\lambda} = 0, \bar{q})$. Instead, $\bar{\Gamma}_1(\hat{q}, \bar{q})$ can and must be retrieved from $\bar{\Gamma}'_1(\hat{q}, \hat{\lambda}, \bar{q})$ via lemma A.1 proved in the appendix. The advantage of proceeding like this is that it frees us from integrating out λ and thus the Ansatz for the EAA is now given by $\bar{\Gamma}(\hat{q}, \hat{\lambda}; \bar{q}) = i \ln(\frac{d\mu[q, \lambda]}{[dq][d\lambda]})_{q=\bar{q}+\hat{q}, \hat{\lambda}=\lambda}$ if we write the integrand of $Z(0)$ as $d\mu[q, \lambda]$. In a first approximation one may discard the contribution from the determinant, the cyclic state and $[Q^{1/2}][\lambda^{-3/2}]$. We leave the investigation of this route for future publications.

Unreduced configuration space path integral

If one wants to avoid the above spatial non-localities one may proceed as in section 2 to obtain a path integral over the unconstrained phase space given by

$$Z_s(f) = \int [dq] [dp] [d\phi] [d\pi] [dN] \left| \int [d\eta] [d\rho] e^{-i^s \int d^n x \eta^\mu \{C_\mu, G^I\} \rho_I} \overline{\Omega_0(q(t = -\infty))} \Omega_0(q(t = \infty)) \right. \delta[G] \times \left. e^{-i \int d^n x (p^{ab} \dot{q}_{ab} + (-1)^s i^s \pi_I \dot{\phi}^I + N^\mu C_\mu)} e^{i^s \int d^n x f^{ab} q_{ab}} \right. \quad (4.28)$$

where C_μ are the original constraints (4.3) and

$$\{C_\mu, G^I\} = \frac{\partial C_\mu}{\partial \pi_I} = \delta_\mu^a \phi_{,a}^I + \delta_\mu^0 Q^{-1} S^{IJ} \pi_J \quad (4.29)$$

Note that while the exponent now depends at most quadratically on p, π there is again an obstacle for $s = 0$: The symplectic term $\pi_I \dot{\phi}^I$ has a relative factor of i as compared to C_μ . Then performing the Gaussian integral would not undo the Legendre transform (4.2) but would rather replace π_I by the complex valued rather than real valued object

$$S_{IJ} \frac{1}{N} [i \dot{\phi}^J - N^a \phi_{,a}^J] \quad (4.30)$$

This is the problem already spelled out in the general case in section 2. We thus fix $s = 1$ for what follows.

To integrate out p^{ab} we collect the terms that depend on it using (4.2)

$$\begin{aligned} & p^{ab} \dot{q}_{ab} - N Q^{-1} G_{abcd} p^{ab} p^{cd} - p^{ab} [L_{\bar{N}} q]_{ab} = N Q^{-1} (2 p^{ab} k_{ab} - G_{abcd} p^{ab} p^{cd}) \\ & = N \{ Q G^{abcd} k_{ab} k_{cd} - Q^{-1} G_{abcd} [p^{ab} - Q G^{abef} k_{ef}] [p^{cd} - Q G^{cdgh} k_{gh}] \} \end{aligned} \quad (4.31)$$

where

$$G_{abcd} = q_{a(c} q_{d)b} - \frac{1}{n-2} q_{ab} q_{cd}, \quad G^{abcd} = q^{a(c} q^{d)b} - q^{ab} q^{cd}, \quad G_{abef} G^{efcd} = \delta_{(a}^c \delta_{b)}^d \quad (4.32)$$

The Gaussian integral over p^{ab} can now be performed by shifting p^{ab} and producing a factor of $|\det(NQ^{-1}G)|^{-1/2}$ per spacetime point up to a global phase that drops out in $Z_s(f)/Z_s(0)$ where the determinant is for square matrices of type G which has rank $r = \frac{1}{2}n(n-1)$. There is no conformal mode problem [25] because the integral of an oscillatory Gaussian e^{iax^2} , $a \in \mathbb{R} - \{0\}$ exists while it does not for an undamped Gaussian e^{ax^2} , $a > 0$. Note that there would also not be a conformal mode problem if we had kept $s = 0$ because the exponent $G_{abcd}(q)p^{ab}p^{cd}$ always has a prefactor i as it comes from implementing $\delta[C] = \int [dN] e^{i\langle N, C \rangle}$.

To compute this determinant, we note that

$$G_{abcd} = \hat{G}_{ijkl} E_{ab}^{ij} E_{cd}^{kl}, \quad \hat{G}_{ijkl} = \delta_{i(k} \delta_{l)j} - \frac{1}{n-2} \delta_{ij} \delta_{kl}, \quad E_{ab}^{ij} = e_{(a}^i e_{b)}^j \quad (4.33)$$

where $q_{ab} = \delta_{ij} e_a^i e_b^j$ defines an $(n-1)$ -Bein for q . The matrix \hat{G} viewed as a matrix on symmetric tensors has signature $(r-1, 1)$ i.e. $r-1$ positive eigenvalues $+1$ and one negative eigenvalue $-1/(n-2)$ corresponding to tracefree and pure trace tensors respectively. Let u_α^a be an eigenvector of e_a^i i.e. $e_a^i u_\alpha^a = \lambda_\alpha \delta_a^i u_\alpha^a$. Then $u_{\alpha\beta}^{ab} = u_\alpha^{(a} u_\beta^{b)}$ is an eigenvector of E_{ab}^{ij} i.e. $E_{ab}^{ij} u_{\alpha\beta}^{ab} = \lambda_\alpha \lambda_\beta \delta_{(a}^i \delta_{b)}^j u_{\alpha\beta}^{ab}$. Thus

$$\det(E) = \prod_{1 \leq \alpha \leq \beta \leq n-1} \lambda_\alpha \lambda_\beta = \left[\prod_{1 \leq \alpha \leq n-1} \lambda_\alpha \right]^n = \det(e)^n \quad (4.34)$$

and

$$\det(G) = -\frac{1}{n-2} \det(E)^2 = -\frac{1}{n-2} Q^{2n}, \quad Q = \sqrt{\det(q)} \quad (4.35)$$

Accordingly dropping overall factors

$$\begin{aligned} Z_1(f) &= \int [dq] [d\phi] [d\pi] [dN] \left| \int [d\eta] [d\rho] e^{-i \int d^n x \eta^\mu \{C_\mu, G^I\} \rho_I} \overline{\Omega_0(q(t = -\infty))} \Omega_0(q(t = \infty)) \right. \times \\ & \left. [(Q^{2n} [|N| Q^{-1}]^r)^{-1/2}] \delta[G] e^{-i \int d^n x (|g|^{1/2} [R^{(n)}(g) - \Lambda] + \pi_I \dot{\phi}^I + N^\mu C_\mu)} e^{i \int d^n x f^{ab} q_{ab}} \right. \end{aligned} \quad (4.36)$$

where we used the Gauss-Coddacci relation and $|g|^{1/2} = |N|Q$.

To perform the integral over π_I we note that π_I also appears in the ghost term. The ghost integral appears in a modulus and contains a factor of i in the exponent, but the ghost matrix is even dimensional and thus yields a real valued determinant. Thus the modulus just multiplies the the ghost integral without modulus by a factor ± 1 depending on the non-ghost variables. We collect the π_I dependent terms

$$\begin{aligned} & \pi_I [\dot{\phi}^I - N^a \phi_{,a}^I - Q^{-1} S^{IJ} \eta^0 \rho_J] - \frac{1}{2} N Q^{-1} S^{IJ} \pi_I \pi_J \\ &= -\frac{N}{2Q} [\pi_I - [Q S_{IK} L_n \phi^K + N^{-1} \eta_0 \rho_I]] S^{IJ} [\pi_I - (Q S_{JL} L_n \phi_L + N^{-1} \eta_0 \rho_J)] \\ & \quad + \frac{1}{2Q} (Q S_{IK} L_n \phi^K + N^{-1} \eta_0 \rho_I) S^{IJ} (Q S_{JL} L_n \phi_L + N^{-1} \eta_0 \rho_J) \end{aligned} \quad (4.37)$$

The term $\eta_0 \rho_I$ is bilinear in the ghosts and thus a commuting number, we can shift it away in the integral together with the $L_n \phi$ term and then perform the n Gaussians which up to a global factor gives $[|N|/Q]^{-n/2}$ per spacetime point. Note that the surviving term quartic in the ghosts is ultra-local and thus drops out as η_0 is nil-potent. Thus we obtain

$$\begin{aligned} Z_1(f) &= \int [dq] [d\phi] [dN] \int [d\eta] [d\rho] e^{-i \int d^n x [\eta^a \phi_{,a}^I + \eta_0 (L_n \phi^I)] \rho_I} \overline{\Omega_0(q(t = -\infty))} \Omega_0(q(t = \infty)) \times \\ & \quad [(Q^{2n} [|N|Q^{-1}]^r)^{-1/2} (|N|/Q)^{-n/2}] \delta[G] e^{-i \int d^n x |g|^{1/2} [R^{(n)}(g) - \Lambda - \frac{1}{2} S_{IJ} g^{\mu\nu} \phi_{,\mu}^I \phi_{,\nu}^J]} e^{i \int d^n x f^{ab} q_{ab}} \end{aligned} \quad (4.38)$$

The local measure factor can be written

$$[(Q^{2n} [|N|Q^{-1}]^r)^{-1/2} (|N|/Q)^{-n/2}] = [Q^{-1} (|N|/Q)^{-(n+1)/4}]^n = (|g|^{-(n+1)/8} Q^{(n+1)/2})^n =: f_1(g)^n \quad (4.39)$$

Instead of using $[dq][dN]$ we may want to use $[dg]$. We have $g_{tt} = -N^2 + q_{ab} N^a N^b$, $g_{ta} = q_{ab} N^b$, $g_{ab} = q_{ab}$. The change of variables gives the Jacobean

$$|\det(\partial g / \partial(N, \vec{N}, q))| = |-2N \det(q)| = 2[(-\det(g) \det(q))^{-2/n}]^n =: 2f_2(g)^{-n} \quad (4.40)$$

To see this note that the matrix $\partial g / \partial(N, \vec{N}, q)$ is a rank $n(n+1)/2$ square matrix with rank 1, $n-1$, $1/2n(n-1)$ diagonal blocks given by $-2N$, q_{ac} , $\delta_{(a}^c \delta_{b)}^d$ respectively and which is upper triangular with respect to this block structure. Therefore its determinant is equal to the product of the determinant of its diagonal blocks.

We absorb f_1, f_2 into the ghost action and also carry out the gauge fixing δ distribution and find with $f(g) := f_1(g)f_2(g)$

$$\begin{aligned} Z_1(f) &= \int [dg] \int [d\eta] [d\rho] e^{-i \int d^n x \eta^\mu M_\mu^I(g) \rho_I} \overline{\Omega_0(q(t = -\infty))} \Omega_0(q(t = \infty)) \times \\ & \quad e^{-i \int d^n x |g|^{1/2} [R^{(n)}(g) - \Lambda - \frac{1}{2} S_{IJ} \kappa_\mu^I \kappa_\nu^J g^{\mu\nu}]} e^{i \int d^n x f^{ab} q_{ab}} \end{aligned} \quad (4.41)$$

with the ghost matrix

$$M_\mu^I(g) = f(g) [\delta_\mu^0 (L_n k)^I + \delta_\mu^a k_{,a}^I] = f(g) [N^{-1} \delta_\mu^0 (\kappa_0 \delta_0^I - N^a [\kappa_1]_a^j \delta_j^I + \delta_\mu^a [\kappa_1]_a^j \delta_j^I] \quad (4.42)$$

Note that the gauge fixing condition requires $\phi^I(x) = \sigma(x) k^I(x)$ on the sector σ but since the action depends only quadratically on ϕ^I and because the ghost determinant is formally a product over x of determinants of 4×4 matrices depending linearly on ϕ^I the signs $\sigma(x)$ drop out of $Z(f)$ so that the final expression only contains k^I . We have regularised $\partial_\mu \sigma \equiv 0$ by first replacing by piecewise constant σ and taking one sided derivatives at jumps.

One could of course integrate out the ghosts at the price of picking up a measure factor $|\det[M](g)|$ per spacetime point. Conversely we may formally absorb the state dependence into the ghost matrix by multiplying (4.42) by the factor

$$e^{h/n}, h(x^0, \vec{x}) = \delta(x^0, \infty) I(x^0, \vec{x}) + \delta(x^0, -\infty) \overline{I(x^0, \vec{x})}, \Omega_0(q(x^0)) =: \exp\left(\int d^{n-1} x I(x^0, \vec{x})\right) \quad (4.43)$$

It contributes only at $x^0 = \pm\infty$ and is ultra-local in time in the sense that $I(x^0, \vec{x})$ depends only on $q(x^0)$. We may pick for instance $I(x^0, \vec{x}) = w(\vec{x}) \ln(\Omega_0(q(x^0)))$ where w is any normalised weight function $\int d^{n-1}x w(\vec{x}) = 1$. If $\Omega_0[q] =_{\infty} e^{-\langle q, l \cdot q \rangle / 2}$ is a Fock state for some spatial integral kernel l it is natural to pick

$$I(x^0, \vec{x}) = -1/2 q_{ab}(x^0, \vec{x}) \int d^{n-1}y l^{ab;cd}(\vec{x}, \vec{y}) q_{cd}(x^0, \vec{y}) \quad (4.44)$$

Remarkably the final result is very close to the Einstein-Hilbert path integral for vacuum GR with a cosmological constant which has been intensively studied in ASQG, the scalar field has disappeared completely. However, there are several *key* differences:

1. There is no longer a (say de Donder) gauge fixing term for the metric and no corresponding Fadeev-Popov determinant (after integrating out the ghosts). This is because the model has n additional degrees of freedom as compared to vacuum GR. These are the n scalar fields “eaten by the gauge bosons” $g_{\mu\nu}$ similar to the Higgs Kibble mechanism for the electroweak interaction. The gauge fixing term and FP determinant respectively have been replaced by the “reduction term” that corresponds to the κ dependence in the exponent and respectively a more general determinant that depends on the Dirac matrix, the deWitt metric and ADM to metric variable transformations. While ghost and Dirac matrix are closely related, the reduction term and the gauge fixing term are only related in the sense that both originate from gauge fixing, however, otherwise they are logically independent because we have solved the gauge fixing condition exactly via the δ distribution while the gauge fixing term just suppresses field configurations violating the gauge condition. In this suppressed version, the reduction term would still be there but with k^I replaced by ϕ^I together with the ϕ integral.
2. Accordingly in application of the background field method to the above generating functional we are in the situation of a theory with no gauge invariance left over any more, that is, we apply the method of section 2.3 rather than 2.4 and therefore do not need to worry about adapting background dependent and independent gauge conditions (3.22).
3. The generating functional $Z_1(f)$ has a current f only for the true degrees of freedom q . In principle one could therefore integrate out N, N^a and obtain a path integral over q only just as for a system without gauge symmetry at all. All correlators derivable from (4.42) are directly observable, one does not need to worry about the gauge invariant meaning of correlators among the $g_{\mu\nu}$ as in the usual setting.
4. The contribution from the cyclic state $\Omega_0(q(t = \pm\infty))$ reminds us that (4.40) was truly derived from a canonical quantisation time ordered correlation function. Note that the very notion of time ordering and a Hamiltonian is only meaningful in this relational observable setting where one has a distinguished notion of time and Hamiltonian.
5. The cosmological constant term is augmented by a field dependent contribution (reduction term) $g^{\mu\nu} S_{IJ} \kappa_{\mu}^I \kappa_{\nu}^J$ which, like the measure factor $[\det(M(g))]$, breaks spacetime diffeomorphism or coordinate transformation invariance. This is no problem because the coordinates have been fixed by the gauge condition $\phi^I = \kappa_{\mu}^I x^{\mu}$ and thus acquire a physical, operational meaning.
6. We found that one can obtain a closed expression in terms of a configuration path integral only for the generating functional of time ordered functions, rather than the Schwinger functions. Thus we are forced to consider the Lorentzian version of ASQG in what follows.
7. The path integral is directly for Lorentzian signature QG, no Wick rotation (a questionable notion in a theory in which the metric is not a background) from a Euclidian version is necessary.

4.2 ASQG treatment of the model

We consider $Z_1(f)$ as given in (4.40) as our starting point with the ghost action not integrated out. We define in the usual way as reviewed in section 3 the functionals

$$C_1(f) = i^{-1} \ln(Z_1(f)), \quad \Gamma_1(\hat{q}) = \text{extr}_f[\langle f, \hat{q} \rangle - C(f)] \quad (4.45)$$

If everything would be well defined the definition of the theory would be complete at this point. As this is not the case we formally introduce additional terms to the current and extend with $F^{ab} =: f^{ab}$, $F^{t\mu} =: u^\mu$

$$\langle f, q \rangle = \int d^n x f^{ab} q_{ab} \rightarrow \langle F, g \rangle = \int d^n x F^{\mu\nu} g_{\mu\nu} = \langle f, q \rangle + \int d^n x u^\mu g_{t\mu} \quad (4.46)$$

Thus we obtain $Z'_1(F) = Z'_1(f, u)$ with the property that $Z'_1(f, u)_{u=0} = Z_1(f)$ and consequently also $C'_1(F) = i^{-1} \ln Z'_1(F)$ satisfies $C'_1(f, u)_{u=0} = C_1(f)$. Next we introduce a background \bar{g} , write the dependence of the integrand of (4.40) as $g = \bar{g} + h$ (i.e. in the cyclic state, the ghost matrix and Einstein-Hilbert like action term), replace $[dg]$ by $[dh]$ and $\langle F, g \rangle$ by $\langle F, h \rangle$, thereby obtaining $\bar{Z}'_1(F, \bar{g}) = Z'_1(F) e^{-i\langle F, \bar{g} \rangle}$ so that with $F = (f, u)$

$$\bar{Z}'_1(f, u, \bar{g})_{u=0} = Z'_1(f, u)_{u=0} e^{-i\langle f, \bar{g} \rangle} = Z_1(f) e^{-i\langle f, \bar{g} \rangle} = \bar{Z}_1(f, \bar{g}) \quad (4.47)$$

where $\bar{Z}_1(f, \bar{g})$ results from $Z_1(f)$ by just writing the spatial metric as $q = \bar{q} + \tilde{h}$ and replace just $[dq]$ by $[d\tilde{h}]$ and $\langle f, q \rangle$ by $\langle f, \tilde{h} \rangle$, i.e. the background method is just applied to q , not to the full g .

Next we introduce into $Z'_1(F, \bar{g})$ currents e, f for the ghosts and the R_k, R_k^g terms as displayed in section 3 thereby producing $\bar{Z}'_{1,k}(F, e, r, \bar{g})$ with $\bar{Z}'_{1,k=0}(F, e=0, r=0, \bar{g}) = Z'_1(F, \bar{g})$ where we can set $e = r = 0$ before or after setting $k = 0$. From here on one is now precisely in the situation to which we can apply the ASQG machinery and obtain a Wetterich equation for $\bar{\Gamma}'_{1,k}(\hat{g}, \hat{\eta}, \hat{\rho}; \bar{g})$. Note that the Wetterich equation applies because its derivation does not make use of any details of the measure $d\mu(h, \eta, \rho)$, the only structural elements that it refers to are the R_k, R_k^g terms and the current terms. After solving it we compute $\bar{\Gamma}'_1(\hat{g}, \hat{\eta}, \hat{\rho}; \bar{g}) = \bar{Z}'_{1,k=0}(\hat{g}, \hat{\eta}, \hat{\rho}; \bar{g})$. Now we apply lemma A.1 to get rid of $\hat{\eta}, \hat{\rho}$ and obtain $\bar{\Gamma}'_1(\hat{g}, \bar{g})$ from $\bar{\Gamma}'_1(\hat{g}, \hat{\eta}, \hat{\rho}; \bar{g})$. Then $\Gamma'_1(\hat{g}) = \bar{\Gamma}'_1(\hat{g}, \bar{g})_{\hat{g}'=0, \bar{g}=\hat{g}}$. Finally, splitting $\hat{g} = (\hat{q}, \hat{N})$ we apply lemma A.1 once more to get rid of \hat{N} to finally obtain the desired answer $\Gamma_1(\hat{q})$.

Alternatively, one could refrain from working with the extended primed objects but then would need to adapt the asymptotic safety machinery and in particular heat kernel methods to a split treatment of spatial and temporal derivatives, see [5]. Also alternatively one can integrate out the ghosts i.e. refrain from introducing R_k^g, e, f which has the advantage that we do not need to invoke lemma A.1 but then the Ansatz for $\bar{\Gamma}'$ needs to be generalised to incorporate the logarithm of the ghost determinant as detailed in (3.57).

This analysis is carried out in our companion paper [9].

5 Conclusion and Outlook

The aim of this paper was twofold: On the one hand, to review, in a hopefully not too technical fashion, the formalism of CQG in its reduced phase space incarnation for ASQG practitioners and vice versa the formalism of ASQG for CQG practitioners. On the other, to show that the two approaches can be brought into contact.

From the point of ASQG, the CQG input is to provide a starting point or starting action for the analysis of the Wetterich equation which is based on a clear physical interpretation of the system: First, there is a clear distinction between true i.e. physical and gauge degrees of freedom. Next, if one uses the unitary group generated by the reduced Hamiltonian to obtain the generating functional of physical Feynman distributions then the generating functional *can* be formulated as a path integral over Lorentzian signature metrics involving the Lorentzian Einstein-Hilbert action by unfolding the reduced phase space integral to the unreduced phase space. If one uses instead the contraction semi-group generated by the reduced Hamiltonian to obtain the generating functional of physical Schwinger distributions then the generating functional *cannot* be formulated as a path integral over Euclidian signature metrics involving the Euclidian Einstein-Hilbert action by unfolding the reduced phase space integral to the unreduced phase space. In this case the best one can do is to keep the reduced phase

space path integral and integrate out the momenta. We have sketched how to do this but focussed mostly on the Lorentzian interpretation. Furthermore, the process of integrating out the momenta gives precise measure corrections to the naive “Ansatz Lebesgue measure times exponential of the sum of classical plus gauge fixing plus ghost action”. Finally there is a non-trivial dependence of that measure on the cyclic state on which the QFT underlying the Hamiltonian formulation is based.

From the point of CQG, the ASQG input is to provide a powerful renormalisation scheme for the reduced Hamiltonian since from the exponential of the Legendre transform of the effective action that one obtains from the ASQG framework at $k = 0$ one can reconstruct the matrix elements of the reduced Hamiltonian (in the Euclidian setting this is known as Osterwalder-Schrader reconstruction [22]). We emphasised that at least in the Lorentzian regime from the point of CQG the only object of physical interest is the effective action which is the $k = 0$ limit of the effective average action. The effective average action and its fixed point analysis serves merely as a tool to derive the effective action.

We also had the opportunity to address a number of reservations that practitioners from other approaches to QG sometimes articulate concerning ASQG:

1.

Signature:

We have seen that the ASQG framework can also be applied to Lorentzian GR although parts of the framework have to be adapted. Mathematically this concerns the analytic properties of oscillatory versus damped integrals [23]. The suppression kernels R_k of the Euclidian framework can at best be interpreted as oscillation kernels in the Lorentzian framework.

2.

Background independence:

We showed that there is in fact no background dependence in ASQG: In ASQG one simply uses the background field method as a convenient tool to construct the background independent (effective) average action which is a standard instrument in QFT. It was invented by DeWitt, one of the founding fathers of CQG. The important point is to keep the background unspecified because only then can one unambiguously reconstruct the background independent effective action from the background dependent one.

3.

Truncations:

Truncations in ASQG should be viewed as an approximation method of the renormalisation scheme, unfortunately without error control at present. To the best of our knowledge, there is no renormalisation scheme without such an error uncontrolled approximation due to the sheer complexity of renormalisation group equations. Although in LQG, an incarnation of CQG, one can give a concrete UV finite definition of the reduced theory [32, 33] which in principle is complete, that definition is not free from quantisation ambiguities and thus also requires renormalisation methods. In that sense the issue of truncations exists for both CQG and ASQG, in fact, any approach to QG.

To see both frameworks in their interaction we have studied a concrete model for Lorentzian GR minimally coupled to scalar fields which is not taken to be a serious dark matter model but rather as a proof of principle or showcase example. Starting from CQG quantisation we have cast that model into a form to which a Lorentzian ASQG treatment is immediately applicable. We reserve that analysis for our companion paper [9]. The Lorentzian version of ASQG follows in spirit all the steps of the Euclidian version but some modifications are necessary in order that the flow induced by the Wetterich equation be well defined. This concerns in particular the class of cut-off functions which are now defined by Fourier transforms rather than Laplace transforms and thus have to come equipped with different analytical behaviour due to the fact that the Lorentzian heat kernel is a group rather than a semi-group. We have proposed such an adapted class of cut-off functions and as a spin-off found that some of the cut-off functions used in the Euclidian regime are in fact not granted to be in the image of the Laplace transform of any meaningful mathematical object which likely requires revisiting some results reported in the literature based on the existence of such a pre-image.

The analysis performed in this contribution and our companion paper [9] leaves of course many interesting open questions such as the amount of freedom in the choice of Lorentzian cut-off functions, gauge reduction by other than Klein-Gordon matter, exploration of the Euclidian interpretation of the reduced phase space path integral and its ASQG treatment. We leave this for future research.

Acknowledgements

We thank Renata Ferrero for in depth discussions about the details of Euclidian ASQG.

A Relations between effective actions upon restriction

We consider a generating function $Z_2(f_1, f_2)$ in two variables f_1, f_2 such that $Z_1(f_1) = Z_2(f_1, f_2 = 0)$ is a generating function in one variable. Then of course also $C_1(f_1) = C_2(f_1, f_2 = 0)$ for $C_1(f_1) = i^{-s} \ln(Z_1(f_1))$, $C_2(f_1, f_2) = i^{-s} \ln(Z_2(f_1, f_2))$. The question arises how the Legendre transforms $\Gamma_1(q_1)$ and $\Gamma_2(q_1, q_2)$ of these two functionals are related. The naive guess $\Gamma_1(q_1) = \Gamma_2(q_1, q_2 = 0)$ turns out to be completely false.

Lemma A.1.

Let $C_1(f_1), C_2(f_1, f_2)$ be any functionals such that $C_1(f_1) = C_2(f_1, f_2 = 0) = (R_2 \cdot C_1)(f_1)$ where R_2 denotes restriction to zero in the second argument. Denote by $\Gamma_1(q_1) = [T_1 \cdot C_1](q_1)$, $\Gamma_2(q_1, q_2) = [T_2 \cdot C_2](q_1, q_2)$ their non-singular Legendre transforms w.r.t. only the first and both variables respectively. Then

$$\Gamma_1 = T_1 \circ R_2 \circ T_2 \circ \Gamma_2 \quad (\text{A.1})$$

Proof. :

The identity (A.1) follows from the fact that the Legendre transform is idempotent $T_2 \circ T_2 = \text{id}_2$ when non-singular. \square

That generically $\Gamma_1 \neq R_2 \circ \Gamma_2$ even if $C_1 = R_2 \circ C_2$ can already be seen for a generic polynomial $C_2(\vec{f})$ of second order in 2 variables $\vec{f} = (f_1, f_2)$ for which we define $C_1(f_1) := C_2(f_1, f_2 = 0)$. If one writes $C_2 = A + \vec{B}^T \vec{f} + \frac{1}{2} \vec{f}^T C \vec{f}$ then we see that $\Gamma_1 = R_2 \circ \Gamma_2$ iff $B_2 = 0 = C_{12}$, i.e. there is no ‘‘condensate’’ and no coupling between the subsystems.

B Avoiding a square root Hamiltonian

Lemma B.1.

Let $d, z > 0$ and $c_d^s = \frac{d e^{-is\frac{\pi}{4}}}{2\sqrt{\pi}}$ with $s = 0, 1$. Then

$$[c_d^s]^{-1} e^{(-1)^{s+1} i^s d \sqrt{z}} = \int_{\mathbb{R}^+} \frac{dl}{l^{3/2}} e^{(-1)^{1+s} i^s [z l + \frac{d^2}{4l}]} \quad (\text{B.1})$$

Note that the integral converges absolutely for $s = 0$ both at infinity and zero. One idea of proof is based on the observation of [28] that integrals of exponentials of $-(x^2 + x^{-2})$ are in fact analytically performable and the saddle point approximation of (B.1) which is proportional to the exponential of $(-1)^{1+s} i^s \sqrt{z} d$ in leading order. Another idea is obtained for $s = 0$ by solving the equation $f_d(z) = e^{-d\sqrt{z}} = c_d^0 \int dl e^{-zl} \hat{f}_d(l)$ for $\hat{f}_d(l)$ using the inverse formula for the Laplace transform $\hat{f}_d(l) = \int_C \frac{dz}{2\pi i} e^{lz} f_d(z)$ where C is any path in $\mathbb{C}_+ := \{z \in \mathbb{C}; \Re(z) > 0\}$ between end points z_{\pm} , $\Im(z_{\pm}) = \pm\infty$. Given such a path is given by $u \in \mathbb{R} \mapsto \sqrt{z}(u) = \frac{d}{2l} + |u| \exp(i \text{sgn}(u) \frac{\pi}{4})$. The resulting integral over u is then Gaussian which yields two contributions, one is the integrand of (B.1), the other is proportional to $e^{-\frac{d^2}{4l}} \delta(l) \equiv 0$. In what follows we verify (B.1) directly using the methods of [28] because the Laplace method does not carry over to $s = 1$ (the case $s = 1$ is formally obtained from $s = 0$ by analytic continuation).

Proof. :

The claim is equivalent to

$$\begin{aligned}
[c_d^s]^{-1} &= 2 \int_0^\infty \frac{du}{u^2} e^{(-1)^{s+1} i^s [\frac{d}{2u} - \sqrt{z}u]^2} \\
&= 2 \int_0^\infty dv e^{(-1)^{s+1} i^s [\frac{dv}{2} - \frac{\sqrt{z}}{v}]^2} \\
&= \frac{4}{d} \int_0^\infty dv e^{(-1)^{s+1} i^s [v - \frac{r}{v}]^2}
\end{aligned} \tag{B.2}$$

where we succesively changed variables $l = u^2$, $u = v^{-1}$, $v d/2 \rightarrow v$ and set $r := \frac{d\sqrt{z}}{2} \in \mathbb{R}_+$. Now set $x(v) := v - \frac{r}{v}$ which is a bijection $\mathbb{R}_+ \rightarrow \mathbb{R}$ since $x'(v) = 1 + \frac{r}{v^2} > 0$ with inverse $v(x) = \frac{x}{2} + w(x)$, $w(x) = \sqrt{r + [\frac{x}{2}]^2}$. It follows

$$\begin{aligned}
[c_d^s]^{-1} &= \frac{4}{d} \int_{-\infty}^\infty dx \frac{\frac{x}{2} + w(x)}{2w(x)} e^{(-1)^{s+1} i^s x^2} \\
&= \frac{2}{d} \int_{-\infty}^\infty dx e^{(-1)^{s+1} i^s x^2} \\
&= \frac{2\sqrt{\pi}}{d} e^{i s \frac{\pi}{4}}
\end{aligned} \tag{B.3}$$

where we used that $\frac{x}{w(x)} e^{(-1)^{s+1} i^s x^2}$ is odd under $x \rightarrow -x$. \square

C Issues with the inverse Laplace transform

In the ASQG literature one introduces certain cut-off functions R_k and assumes that they and their products with other functions are in the image of the Laplace transform. Assuming this, the required heat kernel time ‘‘Q’’ integrals over the pre-image that appear in the computation of the flow equations can be reduced to properties of the cut-off function itself, thus one never needs to know the pre-image explicitly. In the first subsection we show that there are issues with the existence of the pre-image for selected proposed cut-offs and in the second we verify for a function that i^s is in the image of the Laplace transform, that it qualifies as a Euclidian cut-off function.

C.1 Inverse Laplace transform of selected proposed cut-offs

A standard reference on the (one sided) Laplace transform is [23]. Recall that given a function $\hat{f} : \mathbb{R}_+ \rightarrow \mathbb{C}$ one defines its Laplace transform $f_k : \mathbb{R}_+ \rightarrow \mathbb{C}$ by

$$f(z) = [L \cdot \hat{f}](z) := \int_0^\infty dt e^{-zt} \hat{f}(t) \tag{C.1}$$

if the integral exists. A class of functions for which this is true are for example the piecewise continuous functions of at most exponential growth, i.e. there exist constants $A, B > 0$ such that $|\hat{f}(t)| \leq A e^{Bt}$ so that f exists for $z > B$.

For this domain of functions for (C.1), the image is a holomorphic function in the half plane $\Re(z) > B$ and by the theorem of Lerch, the Laplace transform is an injection. Thus given two holomorphic functions in the image of this domain we can uniquely reconstruct \hat{f}_k by the Bromwich formula

$$\hat{f}(t) = [L^{-1} \cdot f](t) = \int_{a+i\mathbb{R}} \frac{dz}{2\pi i} e^{zt} f(z) \tag{C.2}$$

as a piecewise constant function of at most exponential growth, where $a > B$ is some constant chosen such that the path in (C.2) is to the right of all singularities of f .

It is quite difficult to determine the exact image of the Laplace transform given a domain. For instance we may extend (C.1) to a distribution which is the weak derivative of a piecewise constant function of compact support in $(0, \infty)$, say $\hat{f}(t) = [\frac{d}{dt}]^n \chi_{[a,b]}(t)$ with $0 < a < b < \infty$ and χ is the characteristic function. Then

$[L \cdot T](z) := z^n [L \cdot \chi]_{[a,b]}(z)$ exists, it is even a holomorphic function, but it is not clear whether the pre-image is unique if we consider the domain of tempered distributions of compact support in $(0, \infty)$.

Therefore it is not clear whether we can turn the logic around, i.e. given a function or distribution $f(z)$ whether we can find a function or distribution \hat{f} and from what precise space such that $f = L \cdot \hat{f}$ holds and whether \hat{f} is unique. This is however the situation we are confronted with in asymptotic safety: We are given a certain function $f_k(\square)$ of the Laplacian and wish to write it as a Laplace transform.

The concrete application in asymptotic safety are the functions

$$f_{k,N}(z) = \frac{k \partial_k R_k(x)}{[z + R_k(z)]^N}, \quad n \geq 1 \quad (\text{C.3})$$

where $R_k(x)$ is the chosen cut-off function. A popular choice, applied in the Euclidian signature regime, is the *optimised cut-off*

$$R_k(z) = (k^2 - z) \theta(k^2 - z) \Rightarrow f_{k,N}(z) = 2 k^{2(1-N)} \theta(k^2 - z) \quad (\text{C.4})$$

Accordingly we are interested in the question whether the function $f_k(x) = \theta(k^2 - x)$ is in the image of the Laplace transform. We will be satisfied to show that there exists at least one pre-image, even if it is not unique.

A possible Ansatz is to apply the Bromwich integral but we expect trouble because $\theta(k^2 - z)$ is obviously not a holomorphic function. At best it is a holomorphic distribution. In fact, we can make use of the identity

$$\theta(z) = \frac{1}{2}(\text{sgn}(z) + 1) = \frac{1}{2} + \int_{\mathbb{R}} \frac{dp}{2\pi} \frac{e^{ipz}}{ip} \quad (\text{C.5})$$

which can be proved using the residue theorem. Formula (C.5) applied to $f_k(z) = \theta(k^2 - z)$ however cannot be analytically extended to a half space $\Re(z) > a$ as the integral over p extends over the whole real axis. If we nevertheless formally apply the Bromwich integral we obtain an ill-defined expression. On the other hand (C.5) demonstrates that $f_{k,n}$ for the optimised cutoff formally applied to the Lorentzian regime can be written as a *Fourier transform* and thus has a Lorentzian heat kernel expression.

Returning to the Euclidian regime we will now rigorously prove:

Proposition C.1.

The function $f_k(z) = \theta(k^2 - z)$, $k \neq 0$ on \mathbb{R}_+ is not in the image of the Laplace transform of a positive measure on the positive real line.

Proof. :

Suppose that a given $f_k(z)$ had a pre-image $\hat{f}_k(t)$ under the Laplace transform. Then we are interested in the heat kernel time integrals

$$I_{k,n} := \int_0^\infty dt \hat{f}_k(t) t^n = \begin{cases} (-1)^n ([\frac{d}{dz}]^n f_k)(z)_{z=0} & n \geq 0 \\ \frac{1}{(|n|-1)!} \int_0^\infty dz z^{|n|-1} f_k(z) & n \leq -1 \end{cases} \quad (\text{C.6})$$

This yields for $f_k(z) = \theta(k^2 - z)$ the explicit expression

$$I_{n,k} = \theta(n) \delta_{n,0} + \theta(-n) \frac{1}{|n|!} k^{2|n|} \quad (\text{C.7})$$

We interpret $\hat{f}_k(t) dt = d\mu_k(t)$ as the symbolic notation of a possibly signed measure on \mathbb{R}^+ . Then for $n \geq 0$

$$\mu_{k,n} := \int d\mu_k(t) t^n = \delta_{n,0} \quad (\text{C.8})$$

is a Stieltjes moment problem [23] (see vol. 2, ch. X, problem 25). It has a solution in the class of positive measures if and only if

$$\sum_{m,n=0}^{\infty} \bar{z}_m z_n \mu_{k,m+n} \geq 0, \quad \sum_{m,n=0}^{\infty} \bar{z}_m z_n \mu_{k,m+n+1} \geq 0 \quad (\text{C.9})$$

and it is unique if there exist $A, B > 0$ such that $\mu_{k,n} \leq A B^n (2n)!$. It is not difficult to show that both criteria are satisfied. The unique positive measure solving (C.7) is the Dirac measure supported at zero, formally $d\mu_k(t) = \delta(t) dt$. But then for any $n > 1$

$$\frac{k^{2n}}{n!} = \int d\mu_k(t) t^{-n} = +\infty \quad (\text{C.10})$$

which is a contradiction. \square

A more formal way to see that (C.7) is contradictory is to Taylor expand the exponential in the relation $\theta(k^2 - x) = \int dt \hat{f}_k(t) e^{-zt}$ and to use (C.7) for $n \geq 0$ which gives $\theta(k^2 - z) \equiv 1$ for all z , i.e. there is no jump at $z = k^2$.

While we cannot exclude that there exists a distribution of a different kind than the ones discussed at the beginning of this section with non-holomorphic images under the Laplace transform, the claimed properties C.7 based on the existence of the pre-image $\hat{f}_k(t)$ of the optimised cut-off are very doubtful.

In the Lorentzian regime we infer from (C.5) that

$$\theta(k^2 - z) = \int_{-\infty}^{\infty} \frac{dt}{2\pi} e^{itz} \left[\frac{e^{-itk^2}}{it} + 2\pi\delta(t) \right] =: \int_{-\infty}^{\infty} dt \hat{f}_k(t) \quad (\text{C.11})$$

which means that indeed a heat kernel expansion exists but the heat kernel time integrals $\int_{-\infty}^{\infty} dt \hat{f}_k(t) t^n$ badly diverge for all $n \in \mathbb{Z}$. Hence the optimised cut-off is also not practical in the Lorentzian regime.

There are also issues with suggested other ‘‘holomorphic’’ cut-offs such as $R_k(z) = k^2 r(z/k^2)$, $r(y) = y [e^y - 1]^{-1}$ which implies $k\partial_k R_k/(z + R_k) = 2 r(y) = 2 R_k(z)/k^2$. The function r has poles on the imaginary axis at $y = 2\pi in$, $n \in \mathbb{Z}$, thus for any $a > 0$ the path in the Bromwich integral is to the right of those poles. Suppose $r(y)$ is the Laplace transform of $\hat{r}(t)$. Then pick a Bromwich contour in the right complex plane starting and ending at $- \mp i(2n + 1)\pi$ respectively and take $n \rightarrow \infty$. We can close that contour in the left complex plane with a semi-circle of radius $r_n = (2n + 1)\pi$ so that $e^y, e^{yt}; t > 0$ decay to zero exponentially as $n \rightarrow \infty$ on that semi-circle. Thus by the residue theorem the Bromwich integral returns the infinite series

$$\hat{r}(t) = \sum_{k \in \mathbb{Z}} (2\pi ik) e^{2\pi ikt} = \frac{d}{dt} \delta_{S^1}(t) \quad (\text{C.12})$$

which is a distribution, namely the derivative of the δ distribution on the circle, and no function as it diverges pointwise in t . In particular it is not piecewise continuous and of exponential type thus the Bromwich integral is not granted to apply. Indeed we can check whether r is the Laplace transform of (C.12) by integrating term-wise for $y > 0$ to obtain

$$\int_0^{\infty} dt e^{-yt} \hat{r}(t) = - \sum_{k \in \mathbb{Z}} \frac{2\pi ik}{2\pi ik - y} = -8\pi^2 \sum_{k=1}^{\infty} \frac{k^2}{[2\pi k]^2 + y^2} \quad (\text{C.13})$$

which diverges for all $y > 0$. Thus if \hat{r} exists, then certainly not as a function. However, we may interpret $\delta_{S^1}(t)$ as the periodisation of the δ distribution on the real axis, i.e.

$$\delta_{S^1}(t) = \sum_{n \in \mathbb{Z}} \delta(t, n) \quad (\text{C.14})$$

Then for $t > 0$ we may try the definition (dropping the $n \leq 0$ contributions by hand)

$$\hat{r}(t) := \frac{d}{dt} \sum_{n=1}^{\infty} \delta(t, n) \quad (\text{C.15})$$

which yields for $y > 0$ upon formally integrating by parts

$$\int_0^{\infty} dt \hat{r}(t) e^{-yt} = y \sum_{n=1}^{\infty} e^{-ny} = y \frac{e^{-y}}{1 - e^{-y}} = r(y) \quad (\text{C.16})$$

since the geometric series converges for $y > 0$.

Accordingly, $r(y)$ can indeed be interpreted as the Laplace transform of the distribution (C.15). However, with this interpretation, the claimed identities (C.6) a priori fail. We have for $N \in \mathbb{Z}$

$$\int_0^\infty t^N \hat{r}(t) = \begin{cases} 0 & N = 0 \\ -N \sum_{n=1}^\infty n^{N-1} = -N \zeta(1-N) & N \neq 0 \end{cases} \quad (\text{C.17})$$

where ζ is the Riemann ζ function. As it stands, the r.h.s. of (C.17) is zero for $N = 0$, diverges to $-\infty$ for $N > 0$ and converges to a positive number for $N < 0$. To make (C.6) hold we interpret the ill defined r.h.s. of (C.17) in terms of the analytic continuation of the ζ function. We define $\zeta(1) = +\infty$ such that $0 \cdot \zeta(1) := 1$ so that (C.17) produces $r(0)$. Then for $N < 0$ (C.6) and (C.17) agree as the integral in (C.6) coincides with the integral definition

$$\zeta(z) = \frac{\int_0^\infty dy y^{z-1} [e^y - 1]^{-1}}{\int_0^\infty dy y^{z-1} e^{-y}} \quad (\text{C.18})$$

of the ζ function for $\Re(z) > 1$. Finally the analytic continuation yields for $N < 0$

$$\zeta(-N) = (-1)^N \frac{B_{N+1}}{N+1} \quad (N \geq 0), \quad (\text{C.19})$$

where B_N denotes the N -th Bernoulli number.

C.2 Cut-off functions in the image of the Laplace transform

The lesson learnt from the previous subsection is that for the cut-off functions proposed in ASQG for the Euclidian regime, a non-trivial task is to show that they are in fact in the image of a function, distribution or measure such that the important relations (C.7) may be applied to compute and show the convergence of the heat kernel time integrals. This issue is open for the optimal cut-off while for the ‘‘holomorphic’’ cut-off we found a distribution in the pre-image. The integrals (C.7) are ill-defined for that distribution, only a ‘‘reinterpretation’’ (ζ function regularisation) makes (C.7) work. On the other hand, the fact that $\hat{r}(t)$ is a distribution rather than a smooth function carries the danger of producing further ill-defined expressions in multiple integrals over heat kernel times as products of distributions are ill-defined.

A safer route is to define the cut-off function as the Laplace transform of a given function. But then it is not a priori clear whether the image of the Laplace transform satisfies the desired properties of a cut-off function. This touches on the branch of mathematics known as Paley-Wiener theory [23]. For instance it is known that $L \cdot L_2(\mathbb{R}_+, dt) = H_2(\mathbb{C}_+)$, i.e. the image of the one sided Laplace transform of L_2 functions \hat{r} on the positive real line are the entire functions r on the upper half plane (with complex variable $z = x + iy$) such that the integrals of $|r|^2$ along the real axis at constant y are uniformly bounded in y (Hardy space). This is certainly not the case for the above $r(z) = -iz/(e^{-iz} - 1)$ whose corresponding integrals over x diverge for any $y > 0$. We leave this interesting question for future research.

The source of the problem is that according to (C.6) the function $\hat{f}_k(t)$ must be able to integrate *all positive and negative powers of t* . This suggests that $\hat{f}_k(t)$ decays rapidly at both $t = 0, = \infty$ and is the reason why we were lead to consider functions such as

$$\hat{f}_k(t) = k^4 \exp(-([k^2 t]^2 + [k^2 t]^{-2})/2) \quad (\text{C.20})$$

in the next section in application to the Lorentzian sector. The integrals of this function multiplied by $t^n, n \in \mathbb{Z}$ exist and are computed in [28]. But then we define $f_k(z)$ as the Laplace transform of $\hat{f}_k(t)$ with *given, existing* $\hat{f}_k(t)$ and $f_k(z)$ is no longer a prescribed but a derived function.

We now estimate the the Laplace transform (C.20) to demonstrate that it has the usually desired properties. We have for $z \geq 0$

$$f_k(z) = k^2 f(z/k^2), \quad f(y) = \int_0^\infty dt e^{-t^2 - t^{-2}} e^{-yt} \quad (\text{C.21})$$

For fixed $k > 0$ we are interested in $z \ll k^2$ and $z \gg k^2$ i.e. $y \rightarrow 0+$ and $y \rightarrow +\infty$. We have [28]

$$\lim_{y \rightarrow 0+} f(y) = f(0) = \frac{e^{-2} \sqrt{\pi}}{2}, \quad f(y) \leq \int_0^\infty e^{-ty} = \frac{1}{y} \quad (\text{C.22})$$

Thus at fixed k the function f_k indeed approaches a constant for $z \ll k^2$ and decays to zero at least as $1/z$ for $z \gg k^2$. Next, for fixed $z > 0$ we are interested in $k \rightarrow 0+$ and $k \rightarrow +\infty$ i.e. $y \rightarrow +\infty$ and $y \rightarrow 0+$ respectively. Using (C.22) we see that $f_k(z)$ decays as k^4/z as $k \rightarrow 0$ and $f_k(z) \rightarrow k^2 f(0)$ as $k \rightarrow \infty$. Thus (C.20) has all the properties that are usually required in the *Euclidian regime* and now it is secured that $f(y)$ is in the image of the Laplace transform. This also enables one to check that the product of two functions f, g is in the image of the Laplace transform when f, g are. This will be the case if e.g. the convolution integral $\int_0^t ds \hat{f}(s) \hat{g}(t-s)$ exists for all $t \geq 0$ and grows at most polynomially in t .

Finally we compute the ‘‘Q’’ integrals for (C.20)

$$Q_n = \int_0^\infty dt t^n \hat{f}(t) = \int_0^\infty dt t^n e^{-t^2-t^{-2}} \quad (\text{C.23})$$

for $n \in \mathbb{Z}$ which cannot be treated by (C.6) because we do not know $f(y)$ in closed form, thus we must compute directly. We note that $Q_n = Q_{-(n+2)}$ which enables us to concentrate on $n \geq -1$. Introducing the bijection $\mathbb{R}_+ \rightarrow \mathbb{R}$ defined by $u = t - t^{-1}, t = u/2 + w, w := \sqrt{1 + [u/2]^2}$ we have $dt = t/(2w) du$ and thus

$$Q_n = \frac{e^{-2}}{2} \int_{-\infty}^\infty \frac{du}{w} e^{-u^2} (w + u/2)^{n+1} \quad (\text{C.24})$$

It is immediate to see that Q_{2n} is computable in closed form [28] in terms of elementary Gaussian integrals. These integrals are also related to modified Bessel functions of the second kind. For $Q_{2n+1}, n \geq -1$ we can make use of the elementary estimate $w^{-1} < 1 < w$ to see that Q_{2n+1} has analytically computable upper and lower bounds [28]. These integrals will be used in our companion paper [9].

Accordingly (C.20) qualifies as a Euclidian cut-off function.

D Lorentzian cut-off functions

In the Lorentzian signature case the Wetterich equation

$$k \partial_k \Gamma_k = \frac{1}{2i} \text{Tr}([\Gamma_k^{(2)} + R_k]^{-1} [k \partial_k R_k]) \quad (\text{D.1})$$

can be usefully expanded as follows (we use the background formalism): We split

$$\Gamma_k = F_k + U_k \quad (\text{D.2})$$

where $F_k^{(2)}$ has the property to be independent of k and to depend just on the background Laplacian, also having its dimension. If both numerator and denominator in (D.1) contain a common prefactor f_k that depends on the dimensionless couplings g_k then $\partial_k R_k$ is to be replaced by $k \partial_k R_k + \eta_k R_k, \eta_k = k \partial_k \ln(f_k)$, the additional ‘‘anomalous dimension’’ term increases the non-linearity of the flow as one must eventually solve the system for $k \partial_k g_k$. We refrain from displaying this term as it can be treated by the same methods as described below.

We then expand (D.1) into a geometric series

$$k \partial_k \Gamma_k = (2i)^{-1} \sum_{n=0}^\infty (-1)^n \text{Tr}([k \partial_k R_k] [F_k^{(2)} + R_k]^{-1} (U_k^{(2)} [F_k^{(2)} + R_k]^{-1})^n) \quad (\text{D.3})$$

which can be rewritten as

$$k \partial_k [\Gamma_k - (2i)^{-1} \text{Tr}(\ln([F_k^{(2)} + R_k])] = (2i)^{-1} \sum_{n=1}^\infty (-1)^n \text{Tr}([k \partial_k R_k] [F_k^{(2)} + R_k]^{-1} (U_k^{(2)} [F_k^{(2)} + R_k]^{-1})^n) \quad (\text{D.4})$$

The logarithmic term on the l.h.s converges to zero as $k \rightarrow 0$ if $R_k \rightarrow 0$. One can therefore ignore it in that limit. We also ignore it for the moment and come back to it at the end of this section. We just remark that since it can be written in terms of the functional determinant $\det(F_k^{(2)} + R_k)$ one may envisage also independent

regularisation schemes such as zeta function regularisation. Note also that if $F_k = F$ is also independent of k and not only $F_k^{(2)}$ then $k\partial_k\Gamma_k = k\partial_k U_k$ and (D.4) becomes a flow equation entirely in terms of the effective potential U_k .

The right hand side of (D.4) can be written (we formally use cyclicity of the trace)

$$(2i)^{-1} \sum_{n=0}^{\infty} (-1)^n \text{Tr}([k\partial_k(F_k^{(2)} + R_k)^{-1}] U_k^{(2)} [(F_k^{(2)} + R_k)^{-1}] U_k^{(2)})^n \quad (\text{D.5})$$

which displays the fundamental role of the operator

$$(F_k^{(2)} + R_k)^{-1} =: [F_k^{(2)}]^{-1} + G_k \quad (\text{D.6})$$

where we assume that G_k is an operator just depending on \square and $G_k \rightarrow 0$ as $k \rightarrow 0$ which is consistent with $R_k \rightarrow 0$ as $k \rightarrow 0$. As an explicit example consider

$$G_k(z) = k^{-2} e^{-k_0^2/k^2} \int_{-\infty}^{\infty} dt e^{-t^2-t^{-2}} e^{i t z/k^2} \quad (\text{D.7})$$

which is a smooth, real valued function of rapid decrease at $|z| = \infty$. The fixed constant k_0 is there just for dimension reasons and does not run with k . Instead of k_0^2/k^2 one could also use $\lambda_k = \Lambda_k/k^2$ the dimensionless cosmological constant where $k_0^2 = \Lambda_{k=0}$. For both choices $G_k \rightarrow 0$ rapidly at $k = 0$ while G_k has the correct dimension of $1/R_k$.

Since the integrand is smooth at $t = 0$ where it and all of its derivatives vanish we can extend the integrand smoothly to zero for all $t < 0$ and therefore may also consider

$$G_k(z) = k^{-2} e^{-k_0^2/k^2} G(z/k^2), \quad G(y) = \int_0^{\infty} dt e^{-t^2-t^{-2}} e^{i t y} \quad (\text{D.8})$$

which however is no longer real valued but still vanishes at $k = 0$. We may also consider the case $k_0 = k$ which has the advantage that this cut-off function leads to autonomous flow equations.

Note that

$$k\partial_k(F_k^{(2)} + R_k)^{-1} = k\partial_k G_k \quad (\text{D.9})$$

just depends on G_k . The reason for choosing (D.7) or (D.8) is that in the heat kernel expansion we are therefore interested in the heat kernel time integrals

$$\int dt \hat{G}_k(t) e^{i\pi/4 (2-d)\text{sgn}(t)} |t|^{-d/2} t^l, \quad l \in \mathbb{N}_0 \quad (\text{D.10})$$

where \hat{G}_k is the integrand of the integral defining G_k , i.e. its Fourier transform and d the spacetime dimension. These integrals all converge due to the properties of \hat{G}_k at $t = 0$.

We consider the first two orders on the right hand side of (D.5) which already display the essential features of the expansion

$$i^{-1} \text{Tr}([k\partial_k G_k] U_k^{(2)} [1 - ((F_k^{(2)})^{-1} + G_k) U_k^{(2)}]) \quad (\text{D.11})$$

which requires us to know also the heat kernel expansion of $(F_k^{(2)})^{-1}$. If $F_k^{(2)}(z)$ can be written as $Az + B$ with $A, B \neq 0$ real constants independent of k we have the *Schwinger proper time expression*

$$(Az + B)^{-1} = -\frac{i}{A} \left[\int_0^{\infty} dt e^{i t z - t C} \right]_{C \rightarrow -iB/A} \quad (\text{D.12})$$

where the integral is performed at $C > 0$ and then analytically extended to $C = -iB/A$. By rescaling one can assume $A = 1$ and it is natural to identify $B = k_0^2 = \Lambda_{k=0}$.

If we now use the version (D.8) we will be confronted in (D.11) with heat kernel expressions of the form

$$\begin{aligned} & \int_0^{\infty} dt_1 \int_0^{\infty} dt_2 \hat{G}'_k(t_1) \tilde{G}_k(t_2) \text{Tr}(U_k^{(2)} e^{it_1 \square} U_k^{(2)} e^{it_2 \square}) \\ &= \int_0^{\infty} dt_1 \int_0^{\infty} dt_2 \hat{G}'_k(t_1) \tilde{G}_k(t_2) \text{Tr}(U_k^{(2)} [e^{it_1 \square} U_k^{(2)} e^{-it_1 \square}] e^{i(t_1+t_2)\square}) \end{aligned} \quad (\text{D.13})$$

where $\hat{G}'_k(t) = \hat{G}_k(t)$ or $\hat{G}'_k(t) = \dot{\hat{G}}_k(t)$ and $\tilde{G}_k(t) = \hat{G}_k(t)$ or $\tilde{G}_k(t) = e^{-Ct}$. As indicated in the main body of the paper we now expand

$$e^{it_1\Box} U_k^{(2)} e^{-it_1\Box} = \lim_N \sum_{n=0}^N \frac{(it_1)^n}{n!} [\Box, U_k^{(2)}]_{(n)} \quad (\text{D.14})$$

and truncate at some finite N . The heat kernel expansion now results in time integrals of the type (note that the sign of $t_1, t_2, t_1 + t_2$ is positive)

$$= \int_0^\infty dt_1 \int_0^\infty dt_2 \hat{G}_k(t_1) \tilde{G}_k(t_2) e^{i\pi/4(2-d)} t_1^r (t_1 + t_2)^s, \quad r \geq 0, s \geq -d/2 \quad (\text{D.15})$$

which all converge due to the estimate $(t_1 + t_2)^{-s} \leq t_1^{-s}$, $s \geq 0$ and the fact that all times are positive. Since there is always at least one factor of $\hat{G}_k(t_1)$ also at higher orders, all those integrals converge. Note that this would not be the case had we used (D.7). Note however that with R_k being complex valued we expect the coupling constants g_k also to be complex valued for $k > 0$. Thus the number of real coupling constants is in principle doubled. We confine to the original number of real couplings by considering only *admissible* trajectories which are those such that all dimensionful couplings are real valued as $k \rightarrow 0$. This halves the number of admissible initial conditions and presents some form of fine tuning.

Coming back to the logarithmic term in (D.4) we note that with the above choice of R_k, F_k we have

$$\begin{aligned} k\partial_k \ln(F_k^{(2)} + R_k) &= -k\partial_k \ln([F_k^{(2)}]^{-1} + G_k) \\ &= -2 \frac{F_k^{(2)} G_k}{1 + F_k^{(2)} G_k} = \sum_{n=1}^{\infty} (-1)^n [F_k^{(2)} G_k]^n \\ F_k^{(2)}(z) G_k(z) &= k^2 k_0^2 \int dt e^{itz} [-i A \frac{d}{dt} + B] e^{-(t k_0)^2 - (t k_0)^{-2}} \end{aligned} \quad (\text{D.16})$$

and if the integration domain of t is positive (note that due to rapid decay there are no boundary terms picked up by the integration by parts) then the heat kernel traces and time integrals of all powers of (D.16) converge. Thus in this case we can even tame the logarithmic term if we truncate the geometric series.

The above choice of R_k in terms of G_k has the advantage to enable a simple series expansion of the Wetterich equation just in terms of U_k . However, the resulting equation is not autonomous as we were forced to introduce k_0 . Had we replaced $B = k_0^2$ by Λ_k then the heat kernel time integrals would not always contain a factor of \hat{G}_k and therefore in general diverge. We therefore now consider a more complicated expansion which however is autonomous.

This is obtained by refraining from introducing G_k and instead we use

$$R_k(z) = k^2 r(z/k^2), \quad r(y) = \int_0^\infty dt \hat{r}(t) e^{ity}, \quad \hat{r}(t) = e^{-t^2 - t^{-2}} \Rightarrow k\partial_k R_k = 2k^2 [r - yr'(y)] = 2k^2 [r - ir_1], \quad \hat{r}_1(t) = \dot{\hat{r}}(t) \quad (\text{D.17})$$

Consider again (D.2) where now $F_k^{(2)} = z + \Lambda_k$. Here Λ_k is the dimensionful cosmological constant. Then we expand the r.h.s. of (D.1) with $u_k = U_k^{(2)}/k^2$; $\lambda_k = \Lambda_k/k^2$, $y = \Box/k^2$

$$\begin{aligned} 2 \text{Tr}[(r - yr') [(y + \lambda_k) + (r + u_k)]^{-1}] &= 2 \sum_{n=0}^{\infty} (-1)^n \text{Tr}\left(\frac{r - ir_1}{y + \lambda_k} [(r + u_k) [y + \lambda_k]^{-1}]^n\right) \\ [y + \lambda_k]^{-1} &= -i \left[\int_0^\infty dt e^{t(iy - \epsilon)} \right]_{\epsilon \rightarrow -i\lambda_k} \end{aligned} \quad (\text{D.18})$$

where all traces and t integrals are computed at finite $\epsilon > 0$ and at the end are analytically continued as displayed. The essential mechanism that makes all t integrals converge namely that at least one factor of $\hat{r}, \dot{\hat{r}}$ appears is preserved, the system is autonomous but the expansion is more complicated than before.

References

- [1] R. Percacci. An introduction to covariant quantum gravity and asymptotic safety. World Scientific, Singapore, 2017.
M. Reuter, F. Saueressig. Quantum gravity and the functional renormalization group. Cambridge monographs on mathematical physics, Cambridge, 2019.
- [2] P.A.M Dirac, Phys. Rev. **73** (1948) 1092; Rev. Mod. Phys. **21** (1949) 392
J. A. Wheeler, "Geometrodynamics", Academic Press, New York, 1962
B. S. DeWitt, Phys. Rev. **160** (1967) 1113; Phys. Rev. **162** (1967) 1195; Phys. Rev. **162** (1967) 1239
A. Komar, "General Relativistic Observables via Hamilton Jacobi Functionals", Phys. Rev. **D4** (1971) 923-927; "Commutator Algebra of General Relativistic Observables", Phys. Rev. **D9** (1974) 885-888; "Generalized Constraint Structure for Gravitation Theory", Phys. Rev. **D27** (1983) 2277-2281; "Consistent Factor Ordering of General Relativistic Constraints", Phys. Rev. **D20** (1979) 830-833
P. G. Bergmann, A. Komar, "The Coordinate Group Symmetries of General Relativity", Int. J. Theor. Phys. **5** (1972) 15; "The Phase Space Formulation of General Relativity and Approaches Towards its Canonical Quantization", Gen. Rel. Grav., **1** (1981) 227-254
K. Kuchař, in : "Quantum Gravity II : A second Oxford Symposium", C.J. Isham, R. Penrose, D. W. Sciama (eds.), Clarendon Press, Oxford, 1981
C.J. Isham, K. Kuchař, Ann. Phys. **164** (1985) 288; Ann. Phys. **164** (1985) 316
- [3] C. Rovelli, "Quantum Gravity", Cambridge University Press, Cambridge, 2004.
T. Thiemann, "Modern Canonical Quantum General Relativity", Cambridge University Press, Cambridge, 2007
J. Pullin, R. Gambini, "A first course in Loop Quantum Gravity", Oxford University Press, New York, 2011
C. Rovelli, F. Vidotto, "Covariant Loop Quantum Gravity", Cambridge University Press, Cambridge, 2015
K. Giesel, H. Sahlmann, "From Classical To Quantum Gravity: Introduction to Loop Quantum Gravity", PoS QGGS2011 (2011) 002, [arXiv:1203.2733]
- [4] M. Henneaux, C. Teitelboim. "Quantisation of Gauge Systems", Princeton University Press, Princeton, 1992
- [5] E. Manrique, S. Rechenberger, F. Saueressig. Asymptotically Safe Lorentzian Gravity. Phys. Rev. Lett. **106** (2011) 251302. arXiv:1102.5012 [hep-th]
E. D'Angelo, K. Rejzner. A Lorentzian renormalisation group equation for gauge theories. e-Print: 2303.01479 [math-ph]
E. D'Angelo. Asymptotic safety in Lorentzian quantum gravity. Phys. Rev. **D 109** (2024) 066012
- [6] T. Thiemann, "Anomaly-free Formulation of non-perturbative, four-dimensional Lorentzian Quantum Gravity", Physics Letters **B380** (1996) 257-264, [gr-qc/9606088]
T. Thiemann, "Quantum Spin Dynamics (QSD)", Class. Quantum Grav. **15** (1998) 839-73, [gr-qc/9606089]; "Quantum Spin Dynamics (QSD) : II. The Kernel of the Wheeler-DeWitt Constraint Operator", Class. Quantum Grav. **15** (1998) 875-905, [gr-qc/9606090]; "Quantum Spin Dynamics (QSD) : IV. 2+1 Euclidean Quantum Gravity as a model to test 3+1 Lorentzian Quantum Gravity", Class. Quantum Grav. **15** (1998) 1249-1280, [gr-qc/9705018]; "Quantum Spin Dynamics (QSD) : V. Quantum Gravity as the Natural Regulator of the Hamiltonian Constraint of Matter Quantum Field Theories", Class. Quantum Grav. **15** (1998) 1281-1314, [gr-qc/9705019]
- [7] K. Falls. Background independent exact renormalisation. Eur. Phys. J.C **81** (2021) 2, 121. e-Print: 2004.11409 [hep-th]
- [8] M. Demmel, F. Saueressig, O. Zanusso. RG flows of Quantum Einstein Gravity on maximally symmetric spaces. JHEP **6** (2014) 26. arXiv:1401.5495 [hep-th]
J. Daas, W. Oosters, F. Saueressig, J. Wang. Asymptotically Safe Gravity-Fermion systems on curved backgrounds. Universe **7** (2021) 306. arXiv:2107.01071 [hep-th]
- [9] R. Ferreoro, T. Thiemann. Relational Lorentzian asymptotically safe quantum gravity: showcase model
- [10] E. S. Fradkin, D. M. Gitman, S. M. Shvartsman. Path integral for a relativistic particle theory. Europhys. Lett. **15** (1991) 241-244
- [11] M. B. Green, J. H. Schwarz, E. Witten. Superstring Theory. Cambridge University Press, Cambridge, 2012.
- [12] L.F. Abbott. Introduction to the Background Field Method. Acta Phys. Polon. **B 13** (1982) 33
- [13] S. Fulling. Aspects of Quantum Field Theory in Curved Spacetime. London Math. Society Student Texts, vol. 17, 1989.

- [14] B. S. DeWitt, R. W. Brehme. Radiation damping in a gravitational field. *Ann. Phys.* **9** (1960) 220-259.
- [15] M. J. Radzikowski. Micro-local approach to the Hadamard condition in quantum field theory on curved spacetime. *Comm. Math. Phys.* **179** (1996) 529-553
- [16] D. Benedetti, K. Groh, P. F. Machado, F. Saueressig. The universal RG machine. *JHEP* **1106** (2011) 079. arXiv:1012.3081 [hep-th]
- [17] T. Thiemann. Canonical quantum gravity, constructive QFT and renormalisation. *Front. in Phys.* **8** (2020) 548232, *Front. in Phys.* **0** (2020) 457. e-Print: 2003.13622 [gr-qc]
- [18] T. Lang, K. Liegener, T. Thiemann. Hamiltonian renormalisation I: derivation from Osterwalder–Schrader reconstruction. *Class. Quant. Grav.* **35** (2018) 24, 245011. e-Print: 1711.05685 [gr-qc]
- [19] C. Rovelli, “Time in Quantum Gravity: Physics Beyond the Schrodinger Regime”, *Phys. Rev.* **D43** (1991) 442-456
 A. S. Vytheeswaran, “Gauge unfixing in second class constrained systems”, *Annals Phys.* **236** (1994) 297-324
 P. Mitra, R. Rajaraman, “Gauge-invariant reformulation of an anomalous gauge theory”, *Physics Letters* **B225** (1989) 267–271
 R. Anishetty, A. S. Vytheeswaran, “Gauge invariance in second-class constrained systems”, *Journal of Physics A: Mathematical and General* **26** (1993) 5613–5619
 B. Dittrich, “Partial and complete observables for canonical general relativity”, *Class. Quant. Grav.* **23** (2006) 6155-6184, [gr-qc/0507106]
 T. Thiemann, “Reduced phase space quantization and Dirac observables” *Class. Quant. Grav.* **23** (2006) 1163-1180 [gr-qc/0411031]
- [20] O. Bratteli, D. W. Robinson, “Operator Algebras and Quantum Statistical Mechanics”, vol. 1,2, Springer Verlag, Berlin, 1997
- [21] R. Haag, “Local Quantum Physics”, Springer Verlag, Berlin, 1984
- [22] J. Glimm and A. Jaffe, “Quantum Physics”, Springer Verlag, New York, 1987
- [23] M. Reed, B. Simon, “Methods of modern mathematical physics”, vol. I-IV, Academic Press, 1980
- [24] T. Thiemann. Renormalization, wavelets, and the Dirichlet-Shannon kernels. *Phys. Rev.* **D 108** (2023) 12, 125008. e-Print: 2207.08294 [hep-th]
- [25] P. A. Mazur, E. Mottola. The path integral measure, conformal factor problem and stability of the ground state of quantum gravity. *Nucl. Phys.* **B341** (1990) 187-212
- [26] J. M. Pawłowski, M. Reichert. Quantum gravity from dynamical metric fluctuations. e-Print: 2309.10785 [hep-th]
- [27] C. Itzykson and J.-M. Drouffe, *Statistical Field Theory*, Cambridge University Press, Cambridge, 1989
- [28] T. Thiemann. Properties of a smooth, dense, invariant domain for singular potential Schroedinger operators. e-Print: 2305.06718 [quant-ph]
 J. Neuser, T. Thiemann. Smooth, invariant orthonormal basis for singular potential Schroedinger operators. e-Print: 2308.07059 [quant-ph]
- [29] K. Giesel, A. Vetter. Reduced loop quantization with four Klein–Gordon scalar fields as reference matter. *Class. Quant. Grav.* **36** (2019) 14, 145002 e-Print: 1610.07422 [gr-qc]
- [30] R. M. Wald, “General Relativity”, The University of Chicago Press, Chicago, 1989
- [31] C. Wetterich. Exact evolution equation for the effective potential. *Phys.Lett.* **B 301** (1993) 90-94. e-Print: 1710.05815 [hep-th]
- [32] K. Giesel, T. Thiemann, “Scalar Material Reference Systems and Loop Quantum Gravity”, *Class. Quant. Grav.* **32** (2015) 135015, [arXiv:1206.3807]
- [33] K. Giesel, T. Thiemann. Hamiltonian theory: dynamics. In: *Handbook of Quantum Gravity*. C. Bambi, L. Modesto, I. Shapiro (eds.), Springer Verlag, Berlin, 2023.