A Note on Asynchronous Challenges: Unveiling Formulaic Bias and Data Loss in the Hayashi—Yoshida Estimator

Evangelos Georgiadis*

April 30, 2024

Abstract

The Hayashi-Yoshida (H-Y)-estimator exhibits an intrinsic, telescoping property that leads to an often overlooked computational bias, which we denote, *formulaic or intrinsic bias*. This *formulaic bias* results in data loss by cancelling out potentially relevant data points, the *nonextant data points*. This paper attempts to formalize and quantify the data loss arising from this bias. In particular, we highlight the existence of *nonextant data points* via a concrete example, and prove necessary and sufficient conditions for the telescoping property to induce this type of *formulaic bias*.

Since this type of bias is nonexistent when inputs, i.e., observation times, $\Pi^{(1)} := (t_i^{(1)})_{i=0,1,\ldots}$ and $\Pi^{(2)} := (t_j^{(2)})_{j=0,1,\ldots}$, are synchronous, we introduce the (a, b)-asynchronous adversary. This adversary generates inputs $\Pi^{(1)}$ and $\Pi^{(2)}$ according to two independent homogenous Poisson processes with rates a > 0 and b > 0, respectively. We address the foundational questions regarding cumulative minimal (or least) average data point loss, and determine the values for a and b.

We prove that for equal rates a = b, the minimal average cumulative data loss over both inputs is attained and amounts to 25%. We present an algorithm, which is based on our theorem, for computing the *exact number of nonextant data points* given inputs $\Pi^{(1)}$ and $\Pi^{(2)}$, and suggest alternative methods. Finally, we use simulated data to empirically compare the (cumulative) average data loss of the (H-Y)-estimator.

1 Introduction

In [1], Hayashi and Yoshida proposed an estimator for the (cumulative) covariance of two diffusion processes when those are observed at only discrete times in an asynchronous setting. More specifically, Hayashi and Yoshida proposed an estimator that natively computes on irregular and asynchronous observations.

While the novelty of the Hayashi-Yoshida (H-Y)-estimator is manifold, the two key value propositions of their estimator, are, its ability to

- compute natively on asynchronous data (or observations) avoiding any synchronization, preprocessing that might introduce bias (along with potential data loss)
- utilize all available data (or observations)¹.

^{*}Theory Group, CQuant Technologies Limited, The Hong Kong Science & Technology Park (HK-STP), Hong Kong. Email:egeorg@cquant.xyz

¹assuming no sampling process is employed, and all available data points are used.

Chronologically and conceptually, the (H-Y)-estimator represents a third generation statistical toolkit for analyzing asynchronous high-frequency data, e.g., financial market data. This estimator, radically, supersedes, its first and second generation predecessors, which were largely based on conceptually simplistic, if not merely flawed, model assumptions. The first generation laid its foundation on the (implicit and very strong) assumptions that data was generated by an underlying process exhibiting properties of synchronicity and equidistant regularity. The second generation, attempted to rectify these strong assumptions imposed on real-world data by providing an intermediate computational preprocessing layer between raw data and estimator; namely, an artificial way to synchronize raw data. Arguably, this attempt turned out to be a quick fix solution, introducing a variety of different types of biases, known as *synchronization bias*, *imputation bias*, *etc...*; types which we refer to as belonging to the class of *extrinsic bias*.²

The first attempt towards a third generation statistical estimator that natively computes on asynchronous data was devised by Jong and Nijman in [5]. The estimator avoids arbitrary imputation methods and utilizes all available transaction data to compute consistent covariance estimates. Their approach follows a regression based methodology, and can be viewed as a generalization of results in [6], which discusses a related problem of *intervalling effect bias*. A non-regression based approach with a continuous time setting, unlike [5], was then proposed by Hayashi and Yoshida in [1], following further investigations and generalizations in [3], [2] and [4]. Despite its nice properties, i.e., being consistent, asymptotically normally distributed, and, in particular, absent from extrinsic bias, the H–Y estimator suffers from an intrinsic bias,³ which can be attributed to the telescoping property of its summation formula. This *formulaic bias*, under certain conditions, leads to data points being completely cancelled out during computation, which we call *nonextant data points*. These are data points that do not impact the output of the estimator. Or more concretely, the output of the estimator does not depend on the value of the functions at those points.

The study of *nonextant data points* for an estimator is of pivotal significance. For one, it provides insights into the inherent limitations of an estimator's ability to compute on inputs, i.e., addressing the foundational question of whether an estimator can make use of all available data points that it is provided with, without (intrinsically) discarding data points in way that do not influence the output – leading to potential information loss. This, in turn, opens up avenues to formalize an alternative information-theoretic metric of an estimator's efficiency (to obtaining the true value) with respect to its intrinsic (or formulaic) ability to utilize all inputs. For another, it provides additional insight of an estimator's breakdown point analysis along with its robustness properties. The breakdown point of an estimator, "is, roughly, the smallest amount of contamination that may cause an estimator to take on arbitrarily large aberrant values" [10]; or more abstractly, "the natural notion that quantifies the effect (or influence) of the outliers on its performance" [11]. In fact, understanding how nonextant data points occur and whether they follow some form of pattern, or occur within specific intervals, links to the notion of stochastic breakdown, as outlined in (Donoho and Huber) [10, Section 5.1 on page 178, where contamination arranged in certain patterns is not effective at all at disturbing or impacting the output of an estimator, than, let's say, contamination that is randomly placed among the data. This illustrates a deficiency, or, mildly put, property,

²i.e., bias induced via preprocessing methods that previous generations of estimators suffered from.

³i.e., bias directly associated with the evaluation or computation of the estimator.

of an estimator that an adversary could use to compromise (or mislead) the estimator. For instance, by enforcing a configuration (or situation) where certain data points, that need to be obfuscated, fall into patterns that do not impact the output of estimator. In the context of a naive, direct application of the (H-Y)-estimator (2.2), which is used to quantify a lead-lag relationship, the existence of *nonextant data points* highlights inherent limitations in its ability to quantify or capture granular (lead-lag) variations. This property, also opens up potential avenues for an adversary to compromise the estimator, by rendering its output to be misleading. For example, an adversary that wants to obfuscate certain data points in the presence of a (H-Y)-analysis, might enforce a configuration in which these data points cancel out without impacting the output of the estimator. To this end, a thorough and rigorous analysis of this estimator is of interest to regulators and regulatory institutions, such as the U.S. Securities and Exchange Commission (SEC), whose goal is to detect and prevent market manipulation. We note that this estimator has extensively been deployed as the chosen statistical toolkit to consult the SEC on certain market activities of exchanges, due to its characteristics of being (supposedly) free from any bias and able to handle asynchronicity, as seen from recent SEC related filings [13, page 5536]. Last but not least, this study provides a reference point to compare average data loss arising from classical, extrinsic bias, such as synchronization and imputation bias, to average data loss arising from our intrinsic or formulaic bias.

Hence, this note serves as a natural first step towards understanding this *formulaic bias*, which involves analyzing the conditions that induce *nonextant data points*, providing an algorithm to compute the *exact* number of nonextant data points, along with addressing the question of estimating the least (cumulative) average data loss under real world type inputs, i.e. inputs that are asynchronous.

1.1 Structure of paper

We highlight via concrete example an arguably counterintuitive bias in the (H-Y)estimator which can be attributed to the intrinsic, telescoping property of its summation formula. We show that this *formulaic bias* induces data points (or observations) to be cancelled out completely during the computation, which we denote as *nonextant data points*. Furthermore, we observe that these data points do not influence the output of the estimator. We note that *formulaic bias* exhibits similar characteristics to those resulting from *synchronization or imputation bias*.

Second, we characterize *nonextant data points*, illustrate and prove the conditions needed for *nonextant data points* to arise. Our theorem is sufficiently conducive to employ as decision procedure for deciding whether a data point is nonextant.

Third, we derive an expression for the expected number of *nonextant data points* assuming that inputs are generated by our (a, b)-asynchronous adversary.

Finally, we present two different algorithms to compute the exact number of *nonex*tant data points. We employ a Monte Carlo style exploration to empirically investigate (cumulative) data loss as the number of time units for the Poisson processes increases, over varying rates. For each instance of varying rates, we generate a fixed number of random samples and compute the data loss. The results obtained provide a reference point for future iterations of our work and provide a basis for comparison to our theoretically obtained results.

2 Preliminaries: definitions, remarks and examples

2.1 Our version

In this section, we are building up definitions to construct expression (2.1), the equivalent, original H–Y formula⁴. For interval notation, we adopt the minimalized variant of that pioneered by C. A. R. Hoare in his note [7, p. 337]. The original notation had three dots; it was then modified by Lyle H. Ramshaw, and popularized in [8].

Let $[a \dots b)$ be a half-open interval, denoting the set of real numbers x in the range $a \leq x < b$. Similar notations apply to open intervals $(a \dots b)$, closed intervals $[a \dots b]$ and half-open intervals that include the right endpoint but not the left, $(a \dots b]$.

We fix $T \in (0..\infty)$ to be an arbitrary terminal time for observing⁵ prices of two securities, $P^{(l)}$, where l = 1, 2. Accordingly, we define, $\Pi^{(1)} := (t_i^{(1)})_{i=0,1,\ldots,M^{(1)}}$ and $\Pi^{(2)} := (t_j^{(2)})_{j=0,1,\ldots,M^{(2)}}$, to be increasing sequences of random observation times such that $t_0^{(1)}, t_0^{(2)} \ge 0$ and $t_{M^{(1)}}^{(1)}, t_{M^{(2)}}^{(2)} \le T$. We will sometimes refer to observation times as points. We note that the estimator cannot natively compute on multiplicities without preprocessing. Additionally, for our work, we assume that with probability 1 all points in both $\Pi^{(1)}$ and $\Pi^{(2)}$ are distinct, by invoking our *asynchronous adversary* to generate completely asynchronous inputs. Additionally, we define the intervals between successive observation times as $I_i^{(1)} := \left(t_{i-1}^{(1)} \dots t_i^{(1)}\right]$ with $I^{(1)} := \bigcup_{i=1}^{(M^{(1)})} (I_i^{(1)})$, and accordingly $I_j^{(2)} := \left(t_{j-1}^{(2)} \dots t_j^{(2)}\right]$ with $I^{(2)} := \bigcup_{j=1}^{(M^{(2)})} (I_j^{(2)})$. For convenience, we let $t_0^{(1)} \wedge t_0^{(2)} := \min(t_0^{(1)}, t_0^{(2)})$ and $t_0^{(1)} \vee t_0^{(2)} := \max(t_0^{(1)}, t_0^{(2)})$. Thus the covariance between $P^{(1)}$ and $P^{(2)}$ over period $|I^{(1)} \cap I^{(2)}| = (t_0^{(1)} \vee t_0^{(2)} \dots t_{M^{(1)}}^{(1)} \wedge t_{M^{(2)}}^{(2)}]$, is provided via

$$\left\langle P_{\Pi^{(1)}}^{(1)}, P_{\Pi^{(2)}}^{(2)} \right\rangle_{\mathrm{H-Y}} \coloneqq \sum_{j=1}^{\left(M^{(2)}\right)} \sum_{i=1}^{\left(M^{(1)}\right)} \left(P_{t_{i}^{(1)}}^{(1)} - P_{t_{i-1}^{(1)}}^{(1)} \right) \left(P_{t_{j}^{(2)}}^{(2)} - P_{t_{j-1}^{(2)}}^{(2)} \right) \mathbb{1}_{\left\{ \left(t_{i-1}^{(1)} \dots t_{i}^{(1)} \right] \cap \left(t_{j-1}^{(2)} \dots t_{j}^{(2)} \right] \neq \emptyset \right\}}$$
$$= \sum_{i,j} \left(\Delta P_{i}^{(1)} \right) \left(\Delta P_{j}^{(2)} \right) \mathbb{1}_{\left\{ I_{i}^{(1)} \cap I_{j}^{(2)} \neq \emptyset \right\}}, \tag{1}$$

where $\Delta P_i^{(1)} := P_{t_i^{(1)}}^{(1)} - P_{t_{i-1}^{(1)}}^{(1)}$ and $\Delta P_j^{(2)} := P_{t_j^{(2)}}^{(2)} - P_{t_{j-1}^{(2)}}^{(2)}$.

2.2 Original version

The original definition of the (cumulative) covariance estimator is presented in [1, Definition 3.1, on page 368] and outlined for completeness and convenience below. Note that our expression in (1) is equivalent to (2).

$$\sum_{i,j} \Delta P^{(1)}(I^i) \Delta P^{(2)}(J^j) \mathbb{1}_{\{I^i \cap J^j \neq \emptyset\}},\tag{2}$$

where, $P^{(1)}(I^i)$ is essentially our $P_i^{(1)}$, and $P^{(2)}(J^j)$ is essentially our $P_j^{(2)}$; additionally, we use $I_i^{(1)}$ and $I_j^{(2)}$ in place of I^i and J^j , respectively.

⁴without the sampling mechanism, as we compute on all available data

⁵i.e., we record changes in prices instantaneously.

Remark 1. At first inspection, (1) reveals that contributions to the sum depend on the indicator function. That is, the only contributions of the product of any pair of (successive) differences $\Delta P_i^{(1)} \Delta P_j^{(2)}$ to the sum occur only when the respective observation intervals $I_i^{(1)}$ and $I_j^{(2)}$ overlap.

A deeper inspection of (1) reveals that due to the telescoping nature of the summation not all of the observations or data points actually end up counting towards the sum. Cancellation enters the computation, resulting in observations or data points being intrinsically cancelled out. The cancelled out data points are our nonextant data points. To illustrate instances that generate nonextant data points, we provide the following concrete example below 2.3.

Remark 2. Further, a word on interval type. The H-Y formula is based on half-open intervals that include right endpoints, in particular, $\mathbb{1}_{\left\{ \begin{pmatrix} t^{(1)}_{i-1} \dots t^{(1)}_i \end{bmatrix} \cap \begin{pmatrix} t^{(2)}_{j-1} \dots t^{(2)}_j \end{bmatrix} \neq \emptyset \right\}}$. Since the phenomenon of cancellation and thus nonextant data points is invariant to whether intervals are half-open to the right or left, and since half-open intervals that include the left endpoints are slightly more common and slightly more convenient to compute with, we could conduct our analysis and computation on half-open intervals that include the left endpoint.

2.3 Concrete example

Suppose two securities have the following observation times (in units of time, e.g., seconds) and prices (in USD denomination). We let terminal time, T = 12, and have observation times $\Pi^{(1)} = (2, 3, 4, 5, 7, 8, 11.5)$, with respectively observed prices $P^{(1)} = (10, 15, 25, 10, 5, 1, 5)$, and $\Pi^{(2)} = (1, 6, 9, 10, 11, 12)$, with respectively observed prices $P^{(2)} = (5, 10, 15, 20, 25, 20)$. To put our notation into use, the third⁶ observation time of $P^{(1)}$ is $t_2^{(1)} = 4$, and thus, the price of $P^{(1)}$ at the third observation time is $P_{t_2^{(1)}}^{(1)} = 25$. The third⁷ observation interval of $P^{(1)}$ is $I_3^{(1)}$. Hence, $I_3^{(1)} = (t_2^{(1)} \dots t_3^{(1)}] = (4 \dots 5]$. The following figure illustrates observation times $\Pi^{(1)}$ and $\Pi^{(2)}$.

Figure 1 illustrates observation times $\Pi^{(1)}$ and $\Pi^{(2)}$. Figure 1 illustrates that interval $I_1^{(2)}$ overlaps with intervals $I_1^{(1)}, I_2^{(1)}, I_3^{(1)}$, and $I_4^{(1)}$, respectively; interval, $I_2^{(2)}$, overlaps with intervals $I_4^{(1)}, I_5^{(1)}$ and $I_6^{(1)}$, respectively; and intervals $I_3^{(2)}$ overlaps with interval $I_6^{(1)}$. Interval, $I_4^{(2)}$, also overlaps with interval $I_6^{(1)}$; and $I_5^{(2)}$ also overlaps with $I_6^{(1)}$. We could have summarized the last three overlaps more succinctly,namely, interval $I_6^{(1)}$ also overlaps with intervals $I_3^{(2)}, I_4^{(2)}$ and $I_5^{(2)}$.

 $^{^{6}\}mathrm{Note},$ the index starts at 0.

⁷Note, the index starts at 1.



Figure 1: Visualizing notation in use for $\left\langle P_{\Pi^{(1)}}^{(1)}, P_{\Pi^{(2)}}^{(2)} \right\rangle_{\mathrm{H-Y}}$. Illustrating the relevant interval overlaps where $\mathbb{1}_{\left\{ \left(t_{i-1}^{(1)} \dots t_{i}^{(1)}\right] \cap \left(t_{j-1}^{(2)} \dots t_{j}^{(2)}\right] \neq \emptyset \right\}}$ holds. There are ten corresponding interval overlaps. Note, this visualization is not sufficiently granular for highlighting half-open intervals. We further note that the input sequence of observation times is equivalent to *BAAAABAABBBAB*, assuming points in $\Pi^{(2)}$ have been labelled with *B*, and points in $\Pi^{(1)}$ have been labelled with an *A*.

$$\left\langle P_{\Pi^{(1)}}^{(1)}, P_{\Pi^{(2)}}^{(2)} \right\rangle_{\mathrm{H-Y}} := \sum_{j=1}^{5} \sum_{i=1}^{6} \left(P_{t_{i}^{(1)}}^{(1)} - P_{t_{i-1}^{(1)}}^{(1)} \right) \left(P_{t_{j}^{(2)}}^{(2)} - P_{t_{j-1}^{(2)}}^{(2)} \right) \mathbb{1}_{\left\{ \left(t_{i-1}^{(1)} \dots t_{i}^{(1)} \right] \cap \left(t_{j-1}^{(2)} \dots t_{j}^{(2)} \right] \neq \emptyset \right\}}$$
(3)

Evaluating only where 1 holds,

$$\begin{split} &= \left(P_{t_{1}^{(1)}}^{(1)} - P_{t_{0}^{(1)}}^{(1)}\right) \left(P_{t_{1}^{(2)}}^{(2)} - P_{t_{0}^{(2)}}^{(2)}\right) \\ &+ \left(P_{t_{2}^{(1)}}^{(1)} - P_{t_{1}^{(1)}}^{(1)}\right) \left(P_{t_{1}^{(2)}}^{(2)} - P_{t_{0}^{(2)}}^{(2)}\right) \\ &+ \left(P_{t_{3}^{(1)}}^{(1)} - P_{t_{2}^{(1)}}^{(1)}\right) \left(P_{t_{1}^{(2)}}^{(2)} - P_{t_{0}^{(2)}}^{(2)}\right) \\ &+ \left(P_{t_{4}^{(1)}}^{(1)} - P_{t_{3}^{(1)}}^{(1)}\right) \left(P_{t_{2}^{(2)}}^{(2)} - P_{t_{0}^{(2)}}^{(2)}\right) \\ &+ \left(P_{t_{4}^{(1)}}^{(1)} - P_{t_{3}^{(1)}}^{(1)}\right) \left(P_{t_{2}^{(2)}}^{(2)} - P_{t_{1}^{(2)}}^{(2)}\right) \\ &+ \left(P_{t_{5}^{(1)}}^{(1)} - P_{t_{4}^{(1)}}^{(1)}\right) \left(P_{t_{2}^{(2)}}^{(2)} - P_{t_{1}^{(2)}}^{(2)}\right) \\ &+ \left(P_{t_{6}^{(1)}}^{(1)} - P_{t_{5}^{(1)}}^{(1)}\right) \left(P_{t_{2}^{(2)}}^{(2)} - P_{t_{2}^{(2)}}^{(2)}\right) \\ &+ \left(P_{t_{6}^{(1)}}^{(1)} - P_{t_{5}^{(1)}}^{(1)}\right) \left(P_{t_{3}^{(2)}}^{(2)} - P_{t_{2}^{(2)}}^{(2)}\right) \\ &+ \left(P_{t_{6}^{(1)}}^{(1)} - P_{t_{5}^{(1)}}^{(1)}\right) \left(P_{t_{4}^{(2)}}^{(2)} - P_{t_{3}^{(2)}}^{(2)}\right) \\ &+ \left(P_{t_{6}^{(1)}}^{(1)} - P_{t_{5}^{(1)}}^{(1)}\right) \left(P_{t_{4}^{(2)}}^{(2)} - P_{t_{3}^{(2)}}^{(2)}\right) \\ &+ \left(P_{t_{6}^{(1)}}^{(1)} - P_{t_{5}^{(1)}}^{(1)}\right) \left(P_{t_{5}^{(2)}}^{(2)} - P_{t_{4}^{(2)}}^{(2)}\right) \end{split}$$

Factoring sequentially and pairwise encircling cancellations, we obtain the following.

(4)

$$= \Delta P_1^{(2)} \left[\left(\begin{pmatrix} P_{t_1}^{(1)} - P_{t_0}^{(1)} \end{pmatrix} + \left(\begin{pmatrix} P_{t_1}^{(1)} \end{pmatrix} - \begin{pmatrix} P_{t_1}^{(1)} \end{pmatrix} + \left(\begin{pmatrix} P_{t_1}^{(1)} \end{pmatrix} - \begin{pmatrix} P_{t_1}^{(1)} \end{pmatrix} + \left(\begin{pmatrix} P_{t_1}^{(1)} \end{pmatrix} + \left(\begin{pmatrix} P_{t_1}^{(1)} \end{pmatrix} + \begin{pmatrix} P_{t_1}^{(1)} \end{pmatrix} + \left(\begin{pmatrix} P_{t_1}^{(1)} \end{pmatrix} + \begin{pmatrix} P_{t_1}^{(1)} \end{pmatrix} + \left(\begin{pmatrix} P_{t_1}^{(1)} \end{pmatrix} + \begin{pmatrix} P_{t_1}^{(1)} \end{pmatrix} + \begin{pmatrix} P_{t_1}^{(1)} \end{pmatrix} + \left(\begin{pmatrix} P_{t_1}^{(1)} \end{pmatrix} + \begin{pmatrix} P_{t_1}^{(1)} \end{pmatrix} + \begin{pmatrix} P_{t_1}^{(1)} - \begin{pmatrix} P_{t_1}^{(1)} \end{pmatrix} + \begin{pmatrix} P_{t_1}^{(1)} \end{pmatrix} + \begin{pmatrix} P_{t_1}^{(1)} - \begin{pmatrix} P_{t_1}^{(1)} \end{pmatrix} + \begin{pmatrix} P_{t_1}^{(1)} \end{pmatrix} + \begin{pmatrix} P_{t_1}^{(1)} - \begin{pmatrix} P_{t_1}^{(1)} \end{pmatrix} + \begin{pmatrix} P_{t_1}^{($$

simplifying after cancellations,

$$= \left(P_{t_1^{(2)}}^{(2)} - P_{t_0^{(2)}}^{(2)}\right) \left(P_{t_4^{(1)}}^{(1)} - P_{t_0^{(1)}}^{(1)}\right) + \left(P_{t_2^{(2)}}^{(2)} - P_{t_1^{(2)}}^{(2)}\right) \left(P_{t_6^{(1)}}^{(1)} - P_{t_3^{(1)}}^{(1)}\right) + \left(P_{t_6^{(1)}}^{(1)} - P_{t_5^{(1)}}^{(1)}\right) \left(P_{t_5^{(2)}}^{(2)} - P_{t_2^{(2)}}^{(2)}\right)$$
(5)

Remark 3. Ten product summands are reduced to three product summands. Given the sequence of computation which involves sequential factorization in step (4) of (3), four cancellations occur. In particular, data points $P_{t_1}^{(1)}, P_{t_2}^{(1)}, P_{t_3}^{(2)}, P_{t_4}^{(2)}$ are cancelled. These are the four nonextant data points.

Remark 4. Since observation times, $\Pi^{(1)}$ and $\Pi^{(2)}$, rather than prices, $P^{(1)}$ and $P^{(2)}$, dictate the cancellation phenomenon, for simplicity, some of the analysis will be reduced to considering $\Pi^{(1)}$ and $\Pi^{(2)}$ as inputs.

Remark 5. Finally, the sequence of computations involving sequential factorization needs to be examined more carefully. In particular, if we factor out $\left(P_{t_4^{(1)}}^{(1)} - P_{t_3^{(1)}}^{(1)}\right)$, we end up with four (instead of three) product summands. However, we further note, (8) is equivalent to (5) in value but not in minimal representation; for the actual computation, refer to the appendix in (.1). Through the lens of symbolic computation, this illustrates a canonicalization challenge that requires attention and opens avenues for further exploration with respect to whether a standard, minimal representation exists that best exemplifies the maximal cancellation⁸.

2.4 (a,b)-Asynchronous Adversary, asynchronous inputs

Our asynchronous adversary is a statistical adversary that assures that inputs $\Pi^{(1)}$ and $\Pi^{(2)}$ are generated by two independent homogenous Poisson processes with rates a > 0 and b > 0, respectively, guaranteeing that inputs maintain a certain property. In our case, this property is that with probability 1, inputs are asynchronous; additionally, our adversary assures that initial overlaps occur at $I_1^{(1)}$ with $I_1^{(2)}$, and terminal overlaps occur analogously. This property is necessary for Remark 7 to hold.

Our concrete example in (2.3) illustrates an instance of asynchronous inputs, $\Pi^{(1)} = (2, 3, 4, 5, 7, 8, 11.5)$, and $\Pi^{(2)} = (1, 6, 9, 10, 11, 12)$. On the other hand, an instance of (complete) synchronous inputs would look like $\Pi^{(1)} = (2, 3, 4, 5, 7, 8, 11.5)$, and $\Pi^{(2)} = (2, 3, 4, 5, 7, 8, 11.5)$. This type of inputs do not trigger the formulaic bias, hence do not cause any nonextant data points. An instance of partially synchronous inputs is $\Pi^{(1)} = (2, 3, 4, 5, 7, 8, 11.5)$, and $\Pi^{(2)} = (1, 3, 5, 6, 7, 9, 12)$.

⁸If so, what is an efficient procedure to compute such a representation. In what context is such a representation unique?

2.4.1 Asynchronicity and boundary effects

For the asynchronous setting, the minimal possible initial and terminal configurations for $\Pi_j^{(2)}$ are as follows. Minimal in the sense that a configuration captures with minimal amount of data points when *nonextant data* points can arise due to asynchronous boundary condition. In our case, determining whether $t_1^{(2)}$ and $t_{M^{(2)}-1}^{(2)}$ are *nonextant*. Further, we label points in $\Pi^{(1)}$ with A and points in $\Pi^{(2)}$ with B and observe the input sequences in the following configurations.



Figure 2: Minimal initial and terminal configuration for $\Pi^{(2)}$ in asynchronous setting. Note that points plotted as O represent *nonextant data points*, and will be cancelled during the computation, whereas other points survive in the computation.

We observe a combination of configurations 2a and 2b, that yields the following sequence BABBAB in the closed observation time interval, which contains subsequence BABBA and subsequence ABBAB.



Figure 3: Combination of configurations 2a and 2b in closed interval, forming sequence BABBAB, which contains subsequences BABBA and ABBAB. This configuration yields to consecutive *nonextant data points* in $\Pi^{(2)}$.

2.4.2 Asynchronicity and absence of boundary effects

For the remaining $t_j^{(2)}$ in $\Pi^{(2)}$ that are not impacted by boundary effects, the following minimal configuration induces *nonextant data points*, giving rise to sequence *BBB* in this case. Analogously, *AAA* for $t_i^{(1)}$ in $\Pi^{(1)}$.



Figure 4: For $t_j^{(2)}$ in Π^2 not affected by boundary condition.

2.5 Overlaps

We let m be the maximal number of overlaps between $I^{(1)}$ and $I^{(2)}$.

Lemma 6. Label points in $\Pi^{(1)}$ with A and points in $\Pi^{(2)}$ with B. Then the number of overlaps, m, is "almost" equal to the total number of points. So asymptotically, for large T, this is T(a + b).

Proof. Add points of both types at 0 and T; this will add one interval of each type at each end, and, in expectation, O(1) intersections. (It should be easy to calculate this error precisely, at least for large T.) If we have double points at 0 and T, then the number of intersections equals 1 plus the number of interior points, by induction: with no interior points we have just two intervals and one overlap; if you add one point, it will divide one existing interval into two, and the total number of overlaps of these with intervals of the opposite class will increase by 1. (We assume that with probability 1, all interior points are distinct.)

Remark 7. Note that inputs generated by our asynchronous adversary are completely asynchronous. Additionally, we can assure that the number of intersections or overlaps, m, equals the number of data points in $\Pi^{(1)}$ and $\Pi^{(2)}$ minus three, or, m = (a + b) - 3. Reviewing our concrete example 2.3, helps to clarify this.

2.6 Telescoping, nonextant data points, average data point loss

While there are various ways to formalize nonextant data points, i.e., the output of the estimator does not depend on the value of the functions at those points, or, points that completely cancel out during the computation without influencing the output of the estimator, a mathematically precise definition appears to be estimator specific. For the (H-Y)-estimator, we note necessary and sufficient conditions to trigger the formulaic bias which results from an interaction of telescoping property and indicator function, is input specific. Completely synchronous input does not induce this bias; asynchronous input is necessary. One way to obtain a clearer mathematical definition is by analyzing the summation formula, 1. A key insight to the formulaic bias, follows from considering what happens when we fix one of the index variables while summing over the other, as seen in our concrete example 2.3. In particular, observe which endpoints get cancelled due to telescoping while others survive when the indicator function, $\mathbb{1}_{\left\{ \begin{pmatrix} t_{i-1}^{(1)} \dots t_i^{(2)} \\ t_{i-1}^{(2)} \dots t_j^{(2)} \end{bmatrix} \neq \emptyset \right\}}$, holds true. Using this insight, we can formalize the nonextant intervals, intervals that might contain nonextant points. Following Remark 4, for simplicity, we focus on observation times, rather than prices.

Proposition 1. Let $\Pi^{(1)} := (t_i^{(1)})_{i=0,1,\dots,M^{(1)}}$ and $\Pi^{(2)} := (t_j^{(2)})_{j=0,1,\dots,M^{(2)}}$, be the inputs to the (H-Y)-estimator, and let $t_{i^+}^{(2)} := \min\{t_v^{(2)} : t_v^{(2)} > t_i^{(1)}\}$ and $t_{i^-}^{(2)} := \max\{t_v^{(2)} : t_v^{(2)} < t_i^{(1)}\}$, then given endpoints, $t_{i-1}^{(1)}$ and $t_i^{(1)}$ in $\Pi^{(1)}$ for $1 < i < M^{(1)}$, nonextant data points are precisely the data points in $\Pi^{(2)}$ that occur between $(t_{(i-1)^+}^{(2)} ... t_{(i)^-}^{(2)})$.

Proof. Note that $\begin{pmatrix} t_{i-1}^{(1)} \dots t_i^{(1)} \end{bmatrix} \cap \begin{pmatrix} t_{j-1}^{(2)} \dots t_j^{(2)} \end{bmatrix} \neq \emptyset$ if and only if $t_i^{(1)} > t_{j-1}^{(2)}$ and $t_{i-1}^{(1)} < t_j^{(2)}$. This is equivalent to $t_{j-1}^{(2)} \leq t_{i-}^{(2)}$ and $t_j^{(2)} \geq t_{(i-1)+}^{(2)}$. Hence, for the actual inputs with prices, the terms $\Delta P_i^{(1)} \cdot \Delta P_j^{(2)}$ for a fixed i are those with such $t_j^{(2)}$. Then, telescoping the sum of them cancels the terms with $t_j^{(2)}$ for $t_j^{(2)} \leq t_{i-}^{(2)}$ and $t_j^{(2)} \geq t_{(i-1)+}^{(2)}$. However, these are not necessarily the nonextant ones, since the smallest of them will appear in a term with $\Delta P_{i-1}^{(1)}$ (unless i = 1), and similarly for the largest.

Remark 8. Given asynchronous inputs, separate consideration for the initial case when i = 1, and terminal case when $i = M^{(1)}$, is needed. Let i = 1, then, given endpoints $t_0^{(1)}$ and $t_1^{(1)}$ in $\Pi^{(1)}$, the nonextant interval is simply $\left(\min\{t_{(0)}^{(2)}, t_{(0)}^{(2)}\}, t_{(1)}^{(2)}\right)$. We note since endpoint $\min\{t_{(0)}^{(2)}, t_{(0)}^{(2)}\}$, which is the initial endpoint of the initial overlap, and hence, not a nonextant data point, can occur prior to $t_0^{(1)}$, therefore, we can no longer simply use $t_{(0)}^{(2)}$. Analogous consideration for the terminal case, when $i = M^{(1)}$.

Let us now characterize *nonextant data points* in a more succinct and algorithmically conducive way. Our procedure decides whether a data point is *nonextant* based on whether the union of its (two) immediate adjacent intervals forms a subset over the corresponding overlapping interval. For boundary effects potentially impacting second and penultimate data point, an additional condition is employed.

We abbreviate the cumulative number of nonextant data points in both $\Pi^{(1)}$ and $\Pi^{(2)}$ as f. With regard to quantifying and defining average data loss, we observe that, asymptotically, for large T, our metric converges to the theoretical values (as evidenced in Table 1 of section (2.8)). A natural definition for average data loss, agnostic to how inputs are generated, can be based on overlaps. Since overlaps form a necessary but not sufficient condition for nonextant data points, defining data loss as the ratio of the number of total nonextant data points over the number of total overlaps might seem reasonable, i.e., f/m. As we noted in the above Remark 7, we can compute the number of overlaps from the number of data points.

In this note we specify the underlying stochastic process that our inputs are generated from and *estimate the average proportion of data loss* as shown in (10) and (11).

Theorem 9. Let $\Pi^{(1)} := (t_i^{(1)})_{i=0,1,..,M^{(1)}}$ and $\Pi^{(2)} := (t_j^{(2)})_{j=0,1,..,M^{(2)}}$, be the inputs to the (H-Y)-estimator. Then, $t_j^{(2)}$, for $1 < j < M^{(2)} - 1$, is nonextant if and only if $I_j^{(2)} \cup I_{j+1}^{(2)} \subseteq I_i^{(1)}$ for some *i*.

Proof. $t_j^{(2)}$ is nonextant, if and only if, for exactly one i in $I_i^{(1)}$, $I_i^{(1)} \cap I_j^{(2)} \neq \emptyset \iff I_i^{(1)} \cap I_{j+1}^{(2)} \neq \emptyset$ because this will make all terms with $t_j^{(2)}$ (or, $P_{t_j}^{(2)}$, if we had used $P_{\Pi^{(1)}}^{(1)}, P_{\Pi^{(2)}}^{(2)}$ as inputs) cancel. This, in turn, happens, if and only if, $I_j^{(2)} \cup I_{j+1}^{(2)} \subseteq I_i^{(1)}$ for $1 < j < M^{(2)} - 1$. For j = 1, and $j = M^{(2)} - 1$, if the condition $I_j^{(2)} \cup I_{j+1}^{(2)} \subseteq I_i^{(1)}$ does not hold, we need test whether $I_j^{(2)} \cup I_{j+1}^{(2)}$ intersects $I_i^{(1)}$ for exactly one i.

With the above out of the way, we address the questions of what the minimal average data point loss is given inputs $\Pi^{(1)}$ and $\Pi^{(2)}$ generated by our (a, b)-asynchronous adversary, and for what values of a and b can we guarantee this.

⁹Perhaps alternatively, $t_j^{(2)}$ is *nonextant*, if and only if, $\left(I_j^{(2)} \cup I_{j+1}^{(2)}\right) \cap I_i^{(1)} \neq \emptyset$ and $\forall_{k \neq i} \left((I_j^{(2)} \cup I_{(j+1)}^{(2)}) \cap I_k^{(1)} = \emptyset \right)$.

Theorem 10. Assume that $\Pi^{(1)}$ and $\Pi^{(2)}$ are generated by our (a, b)-asynchronous adversary, and are inputs to the (H-Y)-estimator. Additionally, label points in $\Pi^{(1)}$ with A and points in $\Pi^{(2)}$ with B. Further, let T be large, and ignore border effects. Then the proportion of all points, i.e., $\Pi^{(1)} \cup \Pi^{(2)}$, that are in $\Pi^{(1)}$ and are nonextant there¹⁰, is equivalent to the expected proportion of occurrence of substring AAA which is $(a/(a + b))^3$.

Proof. Using language of and construction in Lemma 6, a point labelled A is nonextant \iff it is preceded and followed by A. In that case, it borders two intervals that are subsets of the same class 2-interval, but if not, then, say the point is followed by B, and thus the A-interval to the right is the first A-interval that intersects the next B-interval, and there the point is not nonextant. Hence, ignoring border effects of the order O(1), the number of nonextant points should be the same as the number of substrings AAA. The number of triples equals the number of total data points minus two which is approximately equal to the number of overlaps, m. Each point has probability a/(a+b) to be an A; thus the expected proportion of AAA is $(a/(a+b))^3$.

Corollary 11. Let $\Pi^{(1)}$ and $\Pi^{(2)}$ be inputs to the (H-Y)-estimator, which were generated by our (a,b)-asynchronous adversary. Further, assume T is large, and ignore border effects. Then the (cumulative) average proportion of nonextant data points in both $\Pi^{(1)}$ and $\Pi^{(2)}$ is provided by $f(a,b) = (\frac{a}{a+b})^3 + (\frac{b}{a+b})^3$, where a > 0 and b > 0.

Theorem 12. Let $f(a,b) = (\frac{a}{a+b})^3 + (\frac{b}{a+b})^3$, where a > 0, b > 0. Then f(a,b) attains the global minimum value f(a,b) = 1/4 for equal rates, a = b.

Proof. Given a > 0 and b > 0, we easily obtain f(a, b) > 0. After algebraic manipulation, we can rewrite $f(a, b) = \frac{(a^2 - ab + b^2)}{(a + b)^2} = 1 - \frac{3ab}{(a + b)^2}$. Now, express f(a, b) as g(t) via the following substitution $t = \frac{a}{a+b}$; hence, $(1-t) = \frac{b}{a+b}$. We obtain g(t) = 1 - 3t(1-t). Further, after completing the square, we obtain $g(t) = 3t^2 - 3t + 1$, which is a parabola with positive coefficient of t^2 . Therefore, its vertex represents the global minimum at $t_{\text{vertex}} = \frac{1}{2}$. Further, this value implies that the global minimum is attained when a = b, since $t = 1/2 = \frac{a}{(a+b)}$ and $1 - t = 1 - 1/2 = \frac{b}{(a+b)}$. Evaluating g(t) at t_{vertex} , we obtain the global minimum value of f(a, b) via $g(\frac{1}{2}) = 1 - 3\left(\frac{1}{2}\right)\left(1 - \frac{1}{2}\right) = \frac{1}{4}$.

2.7 Algorithms

This section includes the relevant material for the algorithms. We outline the implementation for counting the exact number of nonextant data points, which is based on proof of theorem 9. The next section proposes a mergesort algorithm.

2.7.1 Algorithm counting nonextant data points

The algorithm 1 computes nonextant data points excluding boundary points, $t_1^{(2)}$ and $t_{M^{(2)}-1}^{(2)}$. To determine whether these two points are *nonextant*, we have for j = 1,

 $\left(I_1^{(2)} \cup I_2^{(2)}\right) \cap I_i^{(1)} \neq \emptyset \text{ and } \forall_{k \neq i} \left((I_1^{(2)} \cup I_2^{(2)}) \cap I_k^{(1)} = \emptyset\right), \text{ for some } i. \text{ Analogous consideration for } j = M^{(2)} - 1.$

¹⁰NB: The expected proportion of points that are in $\Pi^{(1)}$ is a/(a+b), and thus the expected proportion of nonextant points in $\Pi^{(1)}$ is $(a/(a+b))^2$.

Algorithm 1 Counting Nonextant Data Points in $\Pi^{(2)}$ given inputs: $\Pi^{(1)}$ and $\Pi^{(2)}$

1: procedure CountNonextantDataPoints($\Pi^{(2)}, \Pi^{(1)}$) // initialize counter for nonextant data points in $\Pi^{(2)}$ $c \leftarrow 0$ 2: $m \leftarrow length.(\Pi^{(1)}) - 1$ // # of intervals in $I_i^{(1)}$; $\Pi^{(1)}$ indexed from 0 to $M^{(1)}$ 3: // # of intervals in $I_i^{(2)}$; $\Pi^{(2)}$ indexed from 0 to $M^{(2)}$ $n \leftarrow length.(\Pi^{(2)}) - 1$ 4: for $j \leftarrow 2$ to n-2 do 5:
$$\begin{split} I_{j} &\leftarrow 2 \text{ to } n-2 \text{ do} \\ I_{j}^{(2)} &\leftarrow \left(\Pi_{j-1}^{(2)}, \Pi_{j}^{(2)}\right) \\ I_{(j+1)}^{(2)} &\leftarrow \left(\Pi_{j}^{(2)}, \Pi_{j+1}^{(2)}\right) \\ \text{for } i &\leftarrow 1 \text{ to } m \text{ do} \\ I_{i}^{(1)} &\leftarrow \left(\Pi_{i-1}^{(1)}, \Pi_{i}^{(1)}\right) \\ \end{split}$$
6: 7: 8: 9: if $\left(I_{j}^{(2)} \cup I_{(j+1)}^{(2)}\right) \subseteq I_{i}^{(1)}$ then 10:11: 12:return c

2.7.2 Counting nonextant data points via mergesorting observation times

This algorithm harnesses the property that inputs need to possess to generate *nonextant* data points. It is equivalent to counting the number of subsequences 'AAA' or 'BBB' (apart from initial boundary conditions, 'ABAA' or 'BABB', and terminal boundary condition, 'AABA' and 'BBAB') while mergesorting $\Pi^{(1)}$ and $\Pi^{(2)}$, whose inputs are naturally sorted in ascending order and have accordingly been labelled with A (for inputs in $\Pi^{(1)}$) and B (for inputs in $\Pi^{(2)}$), respectively. This can be done in $\mathcal{O}(n)$ time, where $n = |\Pi^{(1)}| + |\Pi^{(2)}|$.

2.7.3 Counting nonextant data points given that entire observation sequence is provided

Alternatively, if we are provided with the entire input sequence in terms of A's and B's, then counting nonextant data points appears to be a routine application of the Knuth-Morris-Pratt or KMP algorithm [12].

2.7.4 Simulation Algorithm for (a,b)-asynchronous adversary

In order to generate inputs via our (a, b)-asynchronous adversary, we need to implement a simple algorithm that generates data points for observation times from a homogeneous Poisson process with intensity or rate a. Our implementation harnesses the fact that times between successive events for a homogeneous Poisson process are independent exponential random variables each with rate a. We devise a variant of Ross's implementation for generating the first T time units of two independent homogenous Poisson processes with rates a and b, respectively, as illustrated in [9, p. 72].

2.8 Computational Exploration

We employ the algorithm in 2.7.4 to generate observation times for $\Pi^{(1)}$ and $\Pi^{(2)}$ for two independent homogenous Poisson processes with varying rates, r. Table 1 shows computed values for the *data loss*, which we have previously defined to be the ratio of the cumulative, ($\Pi^{(1)}$ and $\Pi^{(2)}$), *number of nonextant points* over the *number of* overlaps. All entries are provided in the form $\mu \pm \sigma$, where μ is the sample mean and σ is an estimate of the standard deviation. The number of test runs for each instance or experiment was 1000.

Table 1. Empirical data point loss, f/m, for inputs $\Pi^{(1)}$ and $\Pi^{(2)}$ generated by (a, b)-asynchronous adversary over $(0 \dots T]$, compared to theoretical values via f(a, b)(excluding boundary points.)

	r = 1, 1	$r = 1, \frac{1}{2}$	$r = 1, \frac{1}{4}$	$r = 1, \frac{1}{10}$
T = 100	0.249 ± 0.040	0.330 ± 0.055	0.512 ± 0.072	0.742 ± 0.074
T = 1000	0.250 ± 0.012	0.333 ± 0.018	0.519 ± 0.023	0.751 ± 0.020
T = 10000	0.250 ± 0.004	0.333 ± 0.006	0.520 ± 0.007	0.752 ± 0.007
T = 100000	0.250 ± 0.001	0.333 ± 0.002	0.520 ± 0.002	0.752 ± 0.002
Theoretical:	f(1,1) = 1/4	$f(1, \frac{1}{2}) = 1/3$	$f(1, \frac{1}{4}) = \frac{13}{25}$	$f(1, \frac{1}{10}) = \frac{91}{121}$

Acknowledgement

I heartily thank Svante Janson for extensive comments, insightful feedback, assistance in formalizing the proposition as well as correcting errors in earlier drafts, and I am greatly indebted to Daniel M. Kane for helping narrow the initial scope of the problem statement, sharpening the precision of definitions and proposition, as well as extensive comments that helped enhance the clarity and readability of previous versions.

References

- Takaki Hayashi and Nakahiro Yoshida, On covariance estimation of nonsynchronously observed diffusion processes, Bernoulli, 11 (2), (2005), 359-379. https://projecteuclid.org/download/pdf_1/euclid.bj/1116340299
- Takaki Hayashi and Nakahiro Yoshida, Asymptotic normality of a covariance estimator for nonsynchronously observed diffusion processes, Annals of the Institute of Statistical Mathematics, 60, (2008), 367–406. https://doi.org/10.1007/s10463-007-0138-0
- [3] Takaki Hayashi and Shigeo Kusuoka, Consistent estimation of covariation under nonsynchronicity, Stat Infer Stoch Process, 11, (2008), 93-106. https://doi.org/10.1007/s11203-007-9009-9
- [4] Takaki Hayashi and Nakahiro Yoshida, Nonsynchronous covariation process and limit theorems, Stochastic Processes and their Applications, 121 (10), (2011), 2416– 2454.
 https://www.sciencedirect.com/science/article/pii/S0304414910002905
- [5] Frank de Jong and Theo Nijman, High frequency analysis of lead-lag relationships between financial markets, Journal of Empirical Finance, 4, (1997), 259-277. https://www.sciencedirect.com/science/article/abs/pii/S0927539897000091

- [6] Kalman J. Cohen, Gabriel A. Hawawini, Steven F. Maier, Robert A.Schwartz and David K. Whitcomb, Friction in the trading process and the estimation of systematic risk, Journal of Financial Economics, 12 (2), (1983), 263-278. https://www.sciencedirect.com/science/article/pii/0304405X83900387
- [7] C. A. R. Hoare, A note on the for statement, BIT 12, (1972), 334-341. https://doi.org/10.1007/BF01932305
- [8] Ronald L. Graham, Donald E. Knuth, and Oren Patashnik, Concrete Mathematics, 2nd Ed., (Reading, Massachusetts: Addison-Wesley, 1994), xiii–657.
- [9] Sheldon M. Ross, A Course in Simulation, (MacMillan Publishing Company: NY, 1990), vii–202.
- [10] David L. Donoho and Peter J. Huber, The Notion of Breakdown Point, In A Festschrift for Erich L. Lehmann in honor of his 65th birthday (eds P. J. Bickel,K. A. Doksum and J. L. Hodges, Jr.), (Belmont: Wadsworth, 1983), 157–184.
- [11] Ilias Diakonikolas and Daniel M. Kane, Algorithmic High-Dimensional Robust Statistics, (Cambridge University Press: Cambridge, 2023), vii–284. https://doi.org/10.1017/9781108943161
- [12] Donald E. Knuth, James H. Morris, Jr., and Vaughan R. Pratt, Fast Pattern Matching in Strings, SIAM Journal on Computing, 6 (2), (1977), 323–350. https://doi.org/10.1137/0206024
- [13] Federal Register Volume 87, Issue 21 (February 1, 2022), Regulatory Information, Federal Register, Notices, Securities and Exchange Commission, 87 FR 5527, Release No. 34-94080, File No. SR-CboeBZX-2021-039, (Office of the Federal Register, National Archives and Records Administration, 2022), 5527–5541. https://www.govinfo.gov/content/pkg/FR-2022-02-01/pdf/2022-02001.pdf

Appendix

.1 Computation I

$$\begin{split} \left\langle P_{\Pi^{(1)}}^{(1)}, P_{\Pi^{(2)}}^{(2)} \right\rangle_{\mathrm{H-Y}} &:= \sum_{j=1}^{5} \sum_{i=1}^{6} \left(P_{t_{i}^{(1)}}^{(1)} - P_{t_{i-1}^{(1)}}^{(1)} \right) \left(P_{t_{j}^{(2)}}^{(2)} - P_{t_{j-1}^{(2)}}^{(2)} \right) \mathbb{1}_{\left\{ \left(t_{i-1}^{(1)} \dots t_{i}^{(1)} \right] \cap \left(t_{j-1}^{(2)} \dots t_{j}^{(2)} \right] \neq \emptyset \right\}} \tag{6} \\ & \text{factoring sequentially, via } \left(P_{t_{4}^{(1)}}^{(1)} - P_{t_{3}^{(1)}}^{(1)} \right), \text{ and pairwise encircling cancellations} \\ &= \left(P_{t_{1}^{(2)}}^{(2)} - P_{t_{0}^{(2)}}^{(2)} \right) \left[\left(\left(P_{t_{1}^{(1)}}^{(1)} - P_{t_{0}^{(1)}}^{(1)} \right) + \left(\left(P_{t_{3}^{(1)}}^{(1)} - \left(P_{t_{3}^{(1)}}^{(1)} \right) \right) + \left(P_{t_{3}^{(1)}}^{(1)} - \left(P_{t_{3}^{(1)}}^{(1)} \right) \right) \right] \\ &+ \left(P_{t_{4}^{(1)}}^{(1)} - P_{t_{0}^{(1)}}^{(1)} \right) \left[\left(\left(P_{t_{2}^{(2)}}^{(2)} - P_{t_{0}^{(2)}}^{(2)} \right) + \left(P_{t_{2}^{(2)}}^{(2)} - \left(P_{t_{1}^{(2)}}^{(2)} \right) \right) \right] \\ &+ \left(P_{t_{2}^{(2)}}^{(1)} - P_{t_{1}^{(2)}}^{(1)} \right) \left[\left(\left(P_{t_{3}^{(1)}}^{(1)} - P_{t_{0}^{(1)}}^{(1)} \right) + \left(P_{t_{2}^{(2)}}^{(2)} - \left(P_{t_{1}^{(2)}}^{(2)} \right) \right) \right] \\ &+ \left(P_{t_{2}^{(1)}}^{(1)} - P_{t_{1}^{(1)}}^{(1)} \right) \left[\left(\left(P_{t_{3}^{(1)}}^{(2)} - P_{t_{0}^{(2)}}^{(2)} \right) + \left(\left(P_{t_{2}^{(2)}}^{(2)} - \left(P_{t_{1}^{(2)}}^{(2)} \right) \right) \right] \\ &+ \left(P_{t_{2}^{(1)}}^{(1)} - P_{t_{1}^{(1)}}^{(1)} \right) \left[\left(\left(P_{t_{3}^{(2)}}^{(2)} - P_{t_{2}^{(2)}}^{(2)} \right) + \left(\left(P_{t_{2}^{(2)}}^{(2)} - \left(P_{t_{3}^{(2)}}^{(2)} \right) \right) \right] \\ &+ \left(P_{t_{6}^{(1)}}^{(1)} - P_{t_{1}^{(1)}}^{(1)} \right) \left[\left(\left(P_{t_{3}^{(2)}}^{(2)} - P_{t_{2}^{(2)}}^{(2)} \right) + \left(\left(P_{t_{3}^{(2)}}^{(2)} - \left(P_{t_{3}^{(2)}}^{(2)} \right) \right) \right) + \left(P_{t_{5}^{(2)}}^{(2)} - \left(P_{t_{3}^{(2)}}^{(2)} \right) \right) \right] \end{aligned}$$

Further tidying up, we obtain,

$$\begin{pmatrix} P_{t_1}^{(2)} - P_{t_0}^{(2)} \end{pmatrix} \begin{pmatrix} P_{t_1}^{(1)} - P_{t_0}^{(1)} \end{pmatrix} + \begin{pmatrix} P_{t_1}^{(1)} - P_{t_1}^{(1)} \end{pmatrix} \begin{pmatrix} P_{t_2}^{(2)} - P_{t_0}^{(2)} \end{pmatrix} + \begin{pmatrix} P_{t_2}^{(2)} - P_{t_0}^{(2)} \end{pmatrix} + \begin{pmatrix} P_{t_1}^{(2)} - P_{t_1}^{(2)} \end{pmatrix} \begin{pmatrix} P_{t_1}^{(1)} - P_{t_1}^{(1)} \end{pmatrix} + \begin{pmatrix} P_{t_1}^{(1)} - P_{t_1}^{(1)} \end{pmatrix} \begin{pmatrix} P_{t_2}^{(2)} - P_{t_2}^{(2)} \end{pmatrix} \begin{pmatrix} P_{t_1}^{(2)} - P_{t_1}^{(2)} \end{pmatrix} \begin{pmatrix} P_{t_1}^{(1)} - P_{t_1}^{(1)} \end{pmatrix} + \begin{pmatrix} P_{t_1}^{(1)} - P_{t_1}^{(1)} \end{pmatrix} \begin{pmatrix} P_{t_2}^{(2)} - P_{t_2}^{(2)} \end{pmatrix} \begin{pmatrix} P_{t_1}^{(2)} - P_{t_1}^{(2)} \end{pmatrix} \begin{pmatrix} P_{t_1}^{(2)} - P_{t_1}^{$$

Via this sequence of computation we note four cancellations or nonextant data points. However, there are 4 product summands. Obvious algebraic manipulation shows that we can derive (5).