Interface motion from Glauber-Kawasaki dynamics of non-gradient type

Tadahisa Funaki

April 30, 2024

Abstract

We consider the Glauber-Kawasaki dynamics on a d-dimensional periodic lattice of size N, that is, a stochastic time evolution of particles performing random walks with interaction subject to the exclusion rule (Kawasaki part), in general, of non-gradient type, together with the effect of the creation and annihilation of particles (Glauber part) whose rates are set to favor two levels of particle density, called sparse and dense. We then study the limit of our dynamics under the hydrodynamic space-time scaling, that is, 1/N in space and a diffusive scaling N^2 for the Kawasaki part and another scaling K = K(N), which diverges slower, for the Glauber part in time. In the limit as $N \to \infty$, we show that the particles autonomously make phase separation into sparse or dense phases at the microscopic level, and an interface separating two regions is formed at the macroscopic level and evolves under an anisotropic curvature flow.

In the present article, we show that the particle density at the macroscopic level is well approximated by a solution of a reaction-diffusion equation with a nonlinear diffusion term of divergence form and a large reaction term. Furthermore, by applying the results of Funaki, Gu and Wang [arXiv:2404.12234] for the convergence rate of the diffusion matrix approximated by local functions, we obtain a quantitative hydrodynamic limit as well as the upper bound for the allowed diverging speed of K = K(N).

The above result for the derivation of the interface motion is proved by combining our result with that in a companion paper by Funaki and Park [arXiv:2403.01732], in which we analyzed the asymptotic behavior of the solution of the reaction-diffusion equation obtained in the present article and derived an anisotropic curvature flow in the situation where the macroscopic reaction term determined from the Glauber part is bistable and balanced.

1 Introduction — model and results

The present article studies the hydrodynamic behavior of the Glauber-Kawasaki dynamics with a diverging scaling parameter K in the Glauber part. The Kawasaki part governs the

Beijing Institute of Mathematical Sciences and Applications, No. 544 Hefangkou, Huairou District, Beijing 101408, China. e-mail: funaki@ms.u-tokyo.ac.jp

Abbreviated title: Interface motion from Glauber-Kawasaki dynamics

MSC2020: 60K35, 82C22, 74A50.

Keywords: Kawasaki dynamics, exclusion process, Glauber dynamics, hydrodynamic limit, non-gradient model, anisotropic curvature flow, interface problem, phase separation.

time evolution of particles moving as interacting random walks subject to the exclusion rule. It is generally of non-gradient type and we assume its reversibility under Bernoulli measures. The Glauber part prescribes the law of creation and annihilation of particles. Here we quote the original papers [16] by Glauber and [18] by Kawasaki initially designed for stochastic dynamics corresponding to the Ising model. We show that, in particular in the situation that the particles have two favorable stable phases called 'sparse' and 'dense' with different mean densities, they autonomously undergo phase separation into one of these two phases at the microscopic level, and an interface separating the two regions is formed at the macroscopic level and it evolves under an anisotropic curvature flow; see Theorem 1.4.

To show this, in the present article, we establish that the particle density at the macroscopic level is well approximated by the solution of a reaction-diffusion equation with nonlinear diffusion and a diverging reaction term; see Theorem 1.1. The asymptotic behavior of the solution of this equation is studied in a separate article [12] based on a method in partial differential equations (PDEs); see Theorems 1.2 and 1.3. Combining these results, we can complete the derivation of the interface motion from our particle systems. Note that the PDE is used only secondarily.

The present article extends the results obtained in [10], [3] for the Glauber-Kawasaki dynamics or the Glauber-Zero range process of gradient type to a model of non-gradient type. Moreover, applying the results in [9], we obtain a quantitative hydrodynamic limit with its convergence rate and also the upper bound for the allowed diverging speed of K = K(N) in the Glauber part.

1.1 Model at microscopic level

Let us formulate our model. We consider the Glauber-Kawasaki dynamics on a *d*-dimensional periodic square lattice $\mathbb{T}_N^d = (\mathbb{Z}/N\mathbb{Z})^d \equiv \{1, 2, \ldots, N\}^d$ of large size $N \in \mathbb{N} \equiv \{1, 2, \ldots\}$. The generator \mathcal{L}_N of our dynamics is given by the sum of those of Kawasaki and Glauber dynamics with time change factors N^2 and $K = K(N) \geq 1$, respectively:

(1.1)
$$\mathcal{L}_N = N^2 L_K + K L_G.$$

The Kawasaki part is the same as in [14], [6]. It is of non-gradient type and reversible under Bernoulli measures; see (1.2) and the conditions (1)-(3) below. The hydrodynamic scaling limit for the Kawasaki dynamics reversible under Gibbs measures was studied by [29]. In [10], [11], the so-called gradient condition was assumed for the Kawasaki part but here we discuss without assuming it.

To define the operators L_K and L_G precisely, we introduce several notations. The configuration space of the dynamics is $\mathcal{X}_N = \{0,1\}^{\mathbb{T}_N^d}$ whose element is denoted by $\eta = \{\eta_x; x \in \mathbb{T}_N^d\}$ where $\eta_x = 0$ or 1 indicates that the site x is vacant or occupied, respectively. We denote \mathcal{F}_N the set of all functions on \mathcal{X}_N . Let $\tau_x, x \in \mathbb{T}_N^d$, be the shift operators acting on \mathcal{X}_N by $(\tau_x \eta)_y = \eta_{y+x}, y \in \mathbb{T}_N^d$, where addition is modulo N. They also act on \mathcal{F}_N by $(\tau_x f)(\eta) = f(\tau_x \eta), f \in \mathcal{F}_N$.

For $x, y \in \mathbb{T}_N^d$ and $\eta \in \mathcal{X}_N$, $\eta^{x,y}$ denotes the element of \mathcal{X}_N , obtained from η by

exchanging the values of η_x and η_y , that is,

$$(\eta^{x,y})_z = \begin{cases} \eta_y, & \text{if } z = x, \\ \eta_x, & \text{if } z = y, \\ \eta_z, & \text{if } z \neq x, y. \end{cases}$$

For $x \in \mathbb{T}_N^d$ and $\eta \in \mathcal{X}_N$, η^x denotes the element of \mathcal{X}_N , obtained from η by flipping the value of η_x , that is,

$$(\eta^x)_z = \begin{cases} 1 - \eta_x, & \text{if } z = x, \\ \eta_z, & \text{if } z \neq x. \end{cases}$$

The notations τ_x , $\eta^{x,y}$ and η^x , $x, y \in \mathbb{Z}^d$ also indicate the corresponding ones for $\mathcal{X} = \{0,1\}^{\mathbb{Z}^d}$, the configuration space on the whole lattice \mathbb{Z}^d . For $\Lambda \subset \mathbb{T}^d_N$ or $\subset \mathbb{Z}^d$, $(\Lambda)^*$ denotes the set of all (undirected) bonds $b = \{x, y\}$ inside Λ , i.e., $x, y \in \Lambda$ and |x-y| = 1. Throughout the paper, we use the norm |z| for $z = (z_i)_{i=1}^d \in \mathbb{Z}^d$ in ℓ^1 -sense: $|z| = \max_{1 \le i \le d} |z_i|$. We sometimes write η^b instead of $\eta^{x,y}$ for bonds $b = \{x, y\}$.

The generator L_K of the Kawasaki part on \mathbb{T}_N^d is defined as

(1.2)
$$L_K = \sum_{b \in (\mathbb{T}_N^d)^*} c_b(\eta) \pi_b = \frac{1}{2} \sum_{x,y \in \mathbb{T}_N^d : |x-y|=1} c_{x,y}(\eta) \pi_{x,y}$$

where $\pi_b \equiv \pi_{x,y}, b = \{x, y\}$, is the exchange operator on \mathcal{F}_N defined by

$$\pi_b f(\eta) = f(\eta^b) - f(\eta), \quad f \in \mathcal{F}_N.$$

The functions $\{c_b(\eta) \equiv c_{x,y}(\eta); b = \{x, y\} \in (\mathbb{Z}^d)^*\}$ are defined on \mathcal{X} and determine the jump (or exchange) rates of particles between two neighboring sites x and y. We assume that they satisfy the following three conditions (1)–(3):

- (1) Non-degeneracy and locality: $c_{x,y}(\eta) > 0$ and it is local, that is, it depends only on $\{\eta_z; |z-x| \leq r\}$ for some r > 0.
- (2) Spatial homogeneity: $c_{x,y} = \tau_x c_{0,y-x}$ for every $\{x, y\} \in (\mathbb{Z}^d)^*$.
- (3) Detailed balance under Bernoulli measures: $c_{x,y}(\eta)$ does not depend on $\{\eta_x, \eta_y\}$.

In view of (1), the jump rate $c_b(\eta)$ is naturally regarded as a function on \mathcal{X}_N for $b \in (\mathbb{T}_N^d)^*$, at least if N is large enough such that N > 2r. The third condition (3) is equivalent to the symmetricity of L_K with respect to the Bernoulli measures $\nu_{\rho}^N, \rho \in [0, 1]$, that is, the product probability measures on \mathcal{X}_N such that $\nu_{\rho}^N(\eta_x = 1) = \rho$ for every $x \in \mathbb{T}_N^d$. We will denote the Bernoulli measures on \mathcal{X} by ν_{ρ} , under which the operator L_K considered on \mathbb{Z}^d is symmetric; see [8], [10]. Note that we do not assume the gradient condition; cf. [8], [10].

On the other hand, the generator L_G of the Glauber part is given by

$$L_G = \sum_{x \in \mathbb{T}_N^d} c_x(\eta) \pi_x,$$

where $\pi_x, x \in \mathbb{T}_N^d$, is the flip operator on \mathcal{F}_N defined by

$$\pi_x f(\eta) = f(\eta^x) - f(\eta), \quad f \in \mathcal{F}_N.$$

For the flip rates $c_x(\eta)$ defined on \mathcal{X} , we assume the non-negativity, the locality and the spatial homogeneity, that is, $c_x(\eta) = \tau_x c(\eta)$ for some local function $c(\eta) = c_0(\eta) \ge 0$ on \mathcal{X} (regarded as that on \mathcal{X}_N for N large enough). Since η_0 takes values only in $\{0, 1\}$, $c(\eta)$ can be decomposed as

(1.3)
$$c(\eta) = c^{+}(\eta)(1 - \eta_0) + c^{-}(\eta)\eta_0$$

for some local functions $c^{\pm}(\eta)$ which do not depend on η_0 . We interpret $c^+(\eta)$ and $c^-(\eta)$ as the rates of creation and annihilation of a particle at x = 0, respectively.

Let $\eta^N(t) = \{\eta^N_x(t); x \in \mathbb{T}_N^d\}$ be the Markov process on \mathcal{X}_N governed by the infinitesimal generator \mathcal{L}_N in (1.1) with properly taken K such that $1 \leq K = K(N) \nearrow \infty$ as $N \to \infty$. We are interested in the asymptotic behavior as $N \to \infty$ of its macroscopic empirical mass distribution, that is, the measure-valued process defined by

(1.4)
$$\rho^N(t,dv) = N^{-d} \sum_{x \in \mathbb{T}_N^d} \eta^N_x(t) \delta_{x/N}(dv), \quad v \in \mathbb{T}^d,$$

where $\mathbb{T}^d = \mathbb{R}^d / \mathbb{Z}^d$ is a *d*-dimensional continuous torus identified with $[0, 1)^d$ and δ_v is the δ -measure at v.

Our main result can be stated as follows. We consider the situation, under a proper choice of rates $\{c_{x,y}, c_x\}$, that the particles favor two levels of mean densities ρ_+ and $\rho_- \in (0, 1)$ with the same degree of stability. We then prove that $\rho^N(t, dv)$ converges to $\rho_+ dv$ or $\rho_- dv$ on two regions separated by a hypersurface Γ_t called the interface and Γ_t evolves under the anisotropic curvature flow; see Theorem 1.4. We also give the rate of convergence in probability.

The proof is divided into two parts, that is, a probabilistic part which is developed in this article and a PDE part summarized in Theorems 1.2 and 1.3, shown in [12].

1.2 Diffusion matrix and reaction term at macroscopic level

To state our result for the hydrodynamic limit, which is the main contribution of this article, first corresponding to the Kawasaki part, we introduce a quadratic form $(\theta, \hat{c}(\rho)\theta), \theta \in \mathbb{R}^d$, for each $\rho \in [0, 1]$ called the conductivity via the variational formula

(1.5)
$$(\theta, \hat{c}(\rho)\theta) = \inf_{F \in \mathcal{F}_0^d} \left(\theta, \hat{c}(\rho; F)\theta\right),$$

where (\cdot, \cdot) is the inner product of \mathbb{R}^d , \mathcal{F}_0 denotes the class of all local functions on \mathcal{X} , $\mathcal{F}_0^d = (\mathcal{F}_0)^d$,

$$(\theta, \widehat{c}(\rho; F)\theta) = \frac{1}{2} \sum_{|x|=1} \left\langle c_{0,x} \left(\theta, x(\eta_x - \eta_0) - \pi_{0,x} \left(\sum_{y \in \mathbb{Z}^d} \tau_y F \right) \right)^2 \right\rangle_{\rho},$$

and $\langle \cdot \rangle_{\rho} \equiv E^{\nu_{\rho}}[\cdot]$ stands for the expectation with respect to the Bernoulli measure ν_{ρ} on \mathcal{X} . Note that $\pi_{0,x}(\sum_{y \in \mathbb{Z}^d} \tau_y F)$ is well-defined as a finite sum $\sum_{y \in \mathbb{Z}^d} \pi_{0,x}(\tau_y F)$. We choose a $d \times d$ symmetric matrix $\widehat{c}(\rho) = \{\widehat{c}_{ij}(\rho)\}_{1 \leq i,j \leq d}$, which is written in the same notation, corresponding to the quadratic form introduced above, especially to apply results in partial differential equations; cf. *Remark* after Theorem 1.1 of [14].

We also introduce the compressibility:

(1.6)
$$\chi(\rho) = \rho - \rho^2,$$

and the diffusion matrix $D(\rho) = \{D_{ij}(\rho)\}_{1 \le i,j \le d}$ by the Einstein relation:

(1.7)
$$D(\rho) = \frac{\widehat{c}(\rho)}{2\chi(\rho)}, \quad \rho \in [0, 1],$$

see Proposition 2.2 of [25], p. 180 for the relation to the Green-Kubo formula. It is known that $D(\rho)$ is a C^{∞} -function of $\rho \in [0, 1]$:

$$(1.8) D \in C^{\infty}([0,1]),$$

see [1]. Furthermore, the diffusion matrix $D(\rho)$ is uniformly positive and bounded:

(1.9)
$$c_*|\theta|^2 \le (\theta, D(\rho)\theta) \le c^*|\theta|^2, \quad \theta \in \mathbb{R}^d, \ \rho \in [0, 1].$$

where $c_*, c^* > 0$ are constants defined by

$$0 < c_* := \min_{\eta \in \mathcal{X}, x \in \mathbb{Z}^d : |x|=1} c_{0,x}(\eta) \le c^* := \max_{\eta \in \mathcal{X}, x \in \mathbb{Z}^d : |x|=1} c_{0,x}(\eta) < \infty,$$

which follows from the condition (1); see [26] and Lemma 8.1 below for the proof of (1.9).

Next, corresponding to the Glauber part, we introduce a function $f = f(\rho)$ as an ensemble average under ν_{ρ} as follows:

(1.10)
$$f(\rho) = \langle (1-2\eta_0)c(\eta) \rangle_{\rho}$$
$$= (1-\rho)\langle c^+ \rangle_{\rho} - \rho \langle c^- \rangle_{\rho}$$

Note that f is a polynomial of ρ and, in particular, a C^{∞} -function of $\rho \in [0, 1]$. Note also that f(0) > 0 and f(1) < 0, which is important to show the comparison theorem; see Section 7.

1.3 Quantitative and non-gradient hydrodynamic limit

We show that, asymptotically as $N \to \infty$, the macroscopic empirical mass distribution $\rho^N(t, dv)$ is close to the solution $\rho(t, v) = \rho_K(t, v)$ of the following reaction-diffusion equation with a nonlinear diffusion term and a diverging factor K = K(N) in a reaction term

(1.11)
$$\partial_t \rho(t,v) = \sum_{i,j=1}^d \partial_{v_i} \left\{ D_{ij}(\rho(t,v)) \partial_{v_j} \rho(t,v) \right\} + K f(\rho(t,v)), \quad t \ge 0, \ v \in \mathbb{T}^d,$$

where $v = (v_i)_{i=1}^d$. We assume that its initial value $\rho_0(v)$ satisfies the condition:

(1.12)
$$\rho_0 \in C^5(\mathbb{T}^d) \text{ and } 0 < \rho_0(v) < 1$$

Then, the equation (1.11) has a unique classical solution $\rho(t, v) \in C^{1,3}([0, \infty) \times \mathbb{T}^d)$ by applying the results in [20]; see the beginning of Section 6.

To state our result, let $\nu^N \equiv \nu_{1/2}^N$ be the Bernoulli measure on \mathcal{X}_N with $\rho = 1/2$, that is, the uniform probability measure on \mathcal{X}_N : $\nu^N(\eta) = 2^{-N}, \eta \in \mathcal{X}_N$. For two probability densities f and g with respect to ν^N , define the relative entropy $H(f|g) \equiv H_N(f|g)$ by

(1.13)
$$H(f|g) = \int_{\mathcal{X}_N} f \log(f/g) \, d\nu^N$$

We set

(1.14)
$$\bar{\rho}(\lambda) = e^{\lambda}/(e^{\lambda}+1), \quad \lambda \in \mathbb{R},$$

and denote its inverse function by $\overline{\lambda}(\rho)$:

(1.15)
$$\bar{\lambda}(\rho) = \log\{\rho/(1-\rho)\}, \quad \rho \in (0,1).$$

Let $f_0 = f_0(\eta), \eta \in \mathcal{X}_N$, be the (initial) density of the distribution of $\eta^N(0)$ on \mathcal{X}_N with respect to ν^N and, for $\rho_0 = \rho_0(v)$ satisfying (1.12), let

(1.16)
$$\widetilde{\psi}_0(\eta) = Z_N^{-1} \exp\left\{\sum_{x \in \mathbb{T}_N^d} \bar{\lambda}(\rho_0(x/N))\eta_x\right\}$$

with $\overline{\lambda}$ defined by (1.15) and a normalization constant Z_N with respect to ν^N . In other words, $\widetilde{\psi}_0 d\nu^N$ denoted also by $P^{\widetilde{\psi}_0}$ is a product measure on \mathcal{X}_N with a marginal distribution $P^{\widetilde{\psi}_0}(\eta_x = 1) = \rho_0(x/N)$ for every $x \in \mathbb{T}_N^d$.

Then, the difference between $\rho^N(t, dv)$ and $\rho_K(t, v)dv$ with K = K(N) is estimated in probability as in the following theorem. We call it a quantitative hydrodynamic limit, since it gives the convergence rate. Note that ρ_K is moving in N. To set the upper threshold for the allowed K = K(N), define

(1.17)
$$\overline{K}(N) \equiv \overline{K}_{\delta}(N) := \delta \log N,$$

for $\delta > 0$.

Theorem 1.1. For each $K \geq 1$, let $\rho_K(t, v)$ be the solution of the equation (1.11) with an initial value $\rho_0(v)$ that satisfies the condition (1.12). We assume that f_0 and ψ_0 defined above satisfy $H(f_0|\tilde{\psi}_0) \leq C_1 N^{d-\kappa_1}$ for some $C_1 > 0$ and $\kappa_1 > 0$. Then, there exist $\kappa > 0$ and $\mathfrak{c} > 0$ small enough such that for any T > 0, taking $\delta = \delta_T := \mathfrak{c}/T$ in $\overline{K}_{\delta}(N)$, if K = K(N) satisfies $1 \leq K(N) \leq \overline{K}_{\delta}(N)$ and $K(N) \to \infty$, we have

$$P(|\langle \rho^N(t), \phi \rangle - \langle \rho_K(t), \phi \rangle| > \varepsilon) \le CN^{-\kappa}$$

for all $t \in [0,T]$, $\varepsilon > 0$ and $\phi \in C^{\infty}(\mathbb{T}^d)$ for some $C = C_T(\varepsilon,\phi) > 0$, where $\langle \rho, \phi \rangle = \int_{\mathbb{T}^d} \phi(v) \rho(dv)$ for a measure $\rho = \rho(dv)$ on \mathbb{T}^d and $\rho_K(t)$ is identified with $\rho_K(t,v)dv$.

The constants $\kappa > 0$, $\mathfrak{c} > 0$ and $\delta > 0$ in $\overline{K}_{\delta}(N)$ are chosen in Section 5.2. In fact, to make $\kappa > 0$, the product $\delta \cdot T$ must be small enough and this determines $\mathfrak{c} > 0$. Once \mathfrak{c} is determined, we choose $\delta = \delta_T = \mathfrak{c}/T$. In this sense, κ depends on \mathfrak{c} , but it is independent of δ and T. Especially if $f_0 = \tilde{\psi}_0$, the assumption for $H(f_0|\tilde{\psi}_0)$ in this theorem is fulfilled. In [3], [10], a similar theorem was shown for models of gradient type; see also [11]. In particular, it was shown that the theorem holds taking $\overline{K}(N) = \delta(\log N)^{\sigma/2}$ for some small $\delta > 0$ and some $\sigma \in (0, 1)$. The bound for $\overline{K}(N)$ in Theorem 1.1 is better than this. The reason is that our estimates are more accurate for several error terms. Moreover, for our non-gradient model, new results obtained in [9] based on the method of the quantitative homogenization theory are essential to derive the above decay rate $CN^{-\kappa}$ and also to determine the upper bound $\overline{K}(N)$.

Theorem 1.1 reduces the study of the limit of $\rho^N(t, dv)$ as $N \to \infty$ to that of $\rho_K(t, v)$ as $K \to \infty$. The latter is a pure PDE problem and is discussed separately in a companion paper [12]. The results are summarized in the next subsection.

1.4 Interface motion from nonlinear Allen-Cahn equation

We now consider the case that $d \ge 2$ and assume that the reaction term $f(\rho)$ in (1.11) is bistable, i.e. f has three zeros $0 < \rho_- < \rho_* < \rho_+ < 1$ such that $f'(\rho_{\pm}) < 0$ and $f'(\rho_*) > 0$. In addition, we assume the balance condition:

$$\int_{\rho_-}^{\rho_+} f(\rho) D(\rho) d\rho = 0.$$

This means that two stable phases with densities ρ_{\pm} , sparse and dense, have the same degree of stability. Recalling the condition (1.12) for the initial value $\rho_0(v)$, we define Γ_0 by

$$\Gamma_0 := \{ v \in \mathbb{T}^d : \rho_0(v) = \rho_* \}$$

and assume that Γ_0 is a C^5 -hypersurface in \mathbb{T}^d without boundary such that

(1.18)
$$\nabla \rho_0(v) \cdot n(v) \neq 0 \text{ for } v \in \Gamma_0,$$

where n(v) is the normal vector to Γ_0 . Two regions Ω_0^{\pm} surrounded by Γ_0 are defined by $\Omega_0^+ = \{\rho_0 > \rho_*\}$ and $\Omega_0^- = \{\rho_0 < \rho_*\}$, respectively.

Then, one can show that

$$\rho_K(t,v) \to \Xi_{\Gamma_t}(v) \quad \text{as} \ K \to \infty.$$

Here, Γ_t is a smooth closed hypersurface in \mathbb{T}^d and Ξ_{Γ_t} is a step function taking two values ρ_{\pm} defined by

$$\Xi_{\Gamma_t}(v) = \begin{cases} \rho_+, & v \in \Omega_t^+, \\ \rho_-, & v \in \Omega_t^-, \end{cases}$$

where Ω_t^{\pm} are two regions surrounded by Γ_t . The sides of these regions are determined initially at t = 0 and then continuously for t > 0.

The evolution of Γ_t starting from Γ_0 is governed by the equation

(1.19)
$$V = -\operatorname{Tr}(\mu(n)\partial_v n) \equiv -\sum_{i,j=1}^d \mu_{ij}(n)\partial_{v_i} n_j \quad \text{on} \ \Gamma_t$$

where V is the normal velocity of Γ_t from the side of Ω_t^- to Ω_t^+ and $n = (n_i)$ denotes the unit normal vector to Γ_t of the same direction. This describes an anisotropic (direction dependent) curvature flow. It is a mean curvature flow in the special case that $\mu_{ij}(e) = \mu \delta_{ij}$ with a constant μ .

The matrix $\mu(e) = \{\mu_{ij}(e)\}_{1 \le i,j \le d}$ is defined for $e \in \mathbb{R}^d$: |e| = 1 as

$$\mu_{ij}(e) = \frac{1}{\lambda(e)} \int_{\rho_{-}}^{\rho_{+}} \left[D_{ij}(\rho) \sqrt{W_e(\rho)} - \frac{1}{2} \partial_{e_i}(W_e(\rho)) \partial_{e_j} \left(\frac{a_e(\rho)}{\sqrt{W_e(\rho)}} \right) \right] d\rho,$$

$$\lambda(e) = \int_{\rho_{-}}^{\rho_{+}} \sqrt{W_e(\rho)} d\rho, \quad W_e(\rho) = -2 \int_{\rho_{-}}^{\rho} a_e(\rho) f(\rho) d\rho, \quad a_e(\rho) = e \cdot D(\rho) e.$$

Note that $W_e(\rho) \ge 0$ for $\rho \in [\rho_-, \rho_+]$ by the conditions for $f(\rho)$.

The local-in-time well-posedness on a certain time interval [0, T] with T > 0 of the equation (1.19) follows from the non-degeneracy of $\mu(e)$ in the tangential direction to the interface Γ : For some c > 0,

 $(\theta, \mu(e)\theta) \ge c|\theta|^2$ for $\theta \in \mathbb{R}^d$ such that $(\theta, e) = 0$.

Note that (1.19) can be rewritten in an equivalent PDE for the signed distance function d(t, v) from Γ_t ; see [12].

The first result is for the generation of the interface, that is, $\rho_K(t, v)$ reaches the neighborhood of ρ_- or ρ_+ in a very short time of order $K^{-1} \log K$.

Theorem 1.2. (Theorem 1.1 of [12] on $\Omega = \mathbb{T}^d$) Let $\rho_K(t, v)$ be the solution of the equation (1.11) and let ϵ be such that $0 < \epsilon < \overline{\rho}$, where $\overline{\rho} := \min\{\rho_+ - \rho_*, \rho_* - \rho_-\}$. We assume the conditions given at the beginning of this subsection. Then, there exist $K_0 > 0$ and $M_0 > 0$ such that, for all $K \ge K_0$, the following holds at $t = t_K := K^{-1} \log K/(2f'(\rho_*))$.

- (1) $\rho_{-} \epsilon \leq \rho_K(t_K, v) \leq \rho_{+} + \epsilon \text{ for all } v \in \mathbb{T}^d$,
- (2) $\rho_K(t_K, v) \ge \rho_+ \epsilon$ for $v \in \mathbb{T}^d$ such that $\rho_0(v) \ge \rho_* + M_0/K^{1/2}$,
- (3) $\rho_K(t_K, v) \leq \rho_- + \epsilon$ for $v \in \mathbb{T}^d$ such that $\rho_0(v) \leq \rho_* M_0/K^{1/2}$.

The second result is for the propagation of the interface, that is, the derivation of the interface motion Γ_t as long as it remains smooth. The size of the transition layer is $O(K^{-1/2})$.

Theorem 1.3. (Theorem 1.2 of [12]) Under the conditions of Theorem 1.2, for any $0 < \epsilon < \bar{\rho}$, there exist $K_0 > 0, C > 0$ and T > 0 (within the local-in-time well-posedness of (1.19)) such that for every $K \ge K_0$ and $t \in [t_K, T]$ we have

$$\rho_{K}(t,v) \in \begin{cases} [\rho_{-}-\epsilon,\rho_{+}+\epsilon] & \text{for all } v \in \mathbb{T}^{d}, \\ [\rho_{+}-\epsilon,\rho_{+}+\epsilon] & \text{if } v \in \Omega_{t}^{+} \setminus \mathcal{N}_{C/K^{1/2}}(\Gamma_{t}), \\ [\rho_{-}-\epsilon,\rho_{-}+\epsilon] & \text{if } v \in \Omega_{t}^{-} \setminus \mathcal{N}_{C/K^{1/2}}(\Gamma_{t}), \end{cases}$$

where $\mathcal{N}_r(\Gamma_t) := \{ v \in \mathbb{T}^d, \operatorname{dist}(v, \Gamma_t) \leq r \}$ is the r-neighborhood of Γ_t .

The proofs of these two theorems are given based on the comparison theorem for the PDE (1.11). We construct its super and sub solutions. If the PDE (1.11) is a gradient flow of a certain functional, one can apply the method of Γ -convergence to derive the motion of Γ_t . In particular, if the energy in the limit is a total surface tension, one would obtain an evolution of Wulff shape. But our equation is different from this class.

1.5 Main result

The following theorem is obtained for the particle system $\eta^N(t)$ by combining Theorems 1.1 and 1.3.

Theorem 1.4. Assume the condition (1.12) for the initial value $\rho_0(v)$ and $H(f_0|\psi_0) \leq C_1 N^{d-\kappa_1}$ for some $C_1 > 0$ and $\kappa_1 > 0$ in Theorem 1.1 and those in Theorem 1.3, that is, $d \geq 2$, $f(\rho)$ determined in (1.10) from the flip rate of the Glauber part and the initial value $\rho_0(v)$ satisfy the conditions given at the beginning of Section 1.4. Then, we have

$$P(|\langle \rho^N(t), \phi \rangle - \langle \Xi_{\Gamma_t}, \phi \rangle| > \varepsilon) \le C N^{-\kappa}$$

for all $t \in (0,T]$, $\varepsilon > 0$ and $\phi \in C^{\infty}(\mathbb{T}^d)$ for some $C = C_T(\varepsilon, \phi)$, where T > 0 is determined as in Theorem 1.3 and $\kappa > 0$ as in Theorem 1.1.

Proof. Noting that $\Xi_{\Gamma_t} = \rho_+ \mathbf{1}_{\Omega_t^+} + \rho_- \mathbf{1}_{\Omega_t^-}$ and $|\mathbb{T}^d| = 1$, for any $0 < \epsilon < \bar{\rho}$, we have by Theorem 1.3

$$|\langle \rho_K(t), \phi \rangle - \langle \Xi_{\Gamma_t}, \phi \rangle| \le \left(\epsilon + (C_\epsilon / K^{1/2})^{d-1} \cdot |\Gamma_t|\right) \|\phi\|_{\infty}$$

for $K \geq K_0 = K_{0,\epsilon}$ and $t \in [t_K, T]$. Choosing $\epsilon > 0$ small enough and $K \geq K_0$ large enough, since $|\Gamma_t|$ is bounded, one can make the right-hand side smaller than $\varepsilon/2$. Note also that t_K is small for K large. Thus, taking $K = \overline{K}_{\delta_T}(N) \vee K_0$ and $C = C_T(\varepsilon/2, \phi)$, the concluding estimate follows from Theorem 1.1 at least for large N.

This theorem establishes the autonomous phase separation directly for the particle system. The PDE (1.11) was used only secondarily. We set the flip rate c_0 in the Glauber part as the corresponding f determined in (1.10) to satisfy the bistability and balance conditions. This means that the particles prefer two phases with mean densities ρ_{-} or ρ_{+} .

From gradient models, we derived the mean curvature flow, i.e., $\mu_{ij}(e) = \mu \delta_{ij}$; see [3], [4], [10] and also [17]. In the unbalanced case for gradient models assuming that $D(\rho) \equiv D$ (constant), on a shorter time scale such as $O(1/K^{1/2})$, Huygens' principle was derived. In other words, the stronger phase region expands with a constant speed; see [11]. See [7] for other approaches.

1.6 Outline of the article

The article is organized as follows. In Section 2, following [14], we introduce a local equilibrium state $\psi_t(\eta)$ of second order approximation with a leading order term determined by the hydrodynamic equation (1.11). We then prove that Theorem 1.1 is shown once one can prove the bound $N^{-d}H(f_t|\psi_t) \leq CN^{-\kappa}$ for some $C, \kappa > 0$ for the relative entropy per volume of the density $f_t(\eta)$ of the distribution of the process $\eta^N(t)$ with respect to $\psi_t(\eta)$ with a properly chosen second order term $F(\eta)$. This bound is formulated in Proposition 2.1.

The proof of Proposition 2.1 is given in Section 3. Differently from [14], our model has the Glauber part and it includes a diverging factor K = K(N). Moreover, in order to establish a quantitative hydrodynamic limit, we need a decay estimate for the relative entropy per volume. This forces us to derive sufficiently strong error estimates at all steps of the proof. We first calculate the time derivative of $H(f_t|\psi_t)$ for general ψ_t along with obtaining careful error estimates; see Lemma 3.1. Second, we show the refined one-block estimate, that is, the replacement of microscopic functions (even those with a diverging factor K) by their ensemble averages with fine error estimates based on the argument in [10] and the equivalence of ensembles with precise convergence rate; see Theorem 3.3. Third, we show the Boltzmann-Gibbs principle for the gradient replacement, by which one can replace a microscopic function looking of order O(N) but with vanishing ensemble average by a linear function of η_x of gradient form with a properly chosen coefficient; see Theorem 3.5. This theorem, which gives good error estimates for the replacement, is shown by using the decay estimates for the CLT variances given in Section 4. We then need Lemma 3.7 to replace the linear function of η_x of gradient form with a function of order O(1).

The estimates obtained so far are summarized in Lemma 3.8. To give a further bound for an expectation under f_t (which is an unknown distribution) by that under ψ_t (which is a well-understood distribution), we apply the entropy inequality. Then, we are required to show the large deviation type upper bound under ψ_t with fine error estimates; see Theorem 3.9 and Theorem 3.11 as its consequence.

One can observe that the leading term in the estimate in Theorem 3.11 is cancelled if we determine the leading term of ψ_t according to the hydrodynamic equation (1.11); see Lemma 3.13. The equation (1.11) is used only for this lemma. Summarizing these and choosing K = K(N) and other parameters properly as indicated in Section 5, one can conclude the proof of Proposition 2.1. We apply multiscale analysis with different scaling parameters $1 \ll n \ll \ell \ll N$. The proof of Theorem 3.9 is given in Section 3.8.

Section 4 concerns the CLT variances related to the gradient replacement and deals with quantities determined only by the localized Kawasaki generator and under the Bernoulli measures $\{\nu_{\rho}\}_{\rho}$. In particular, the Glauber part plays no role in this section. Proposition 4.1 is a refinement of Proposition 5.1 of [14]. We provide a decay estimate for the error terms in the CLT variances, which was not given in [14]. We use the equivalence of ensembles with a precise convergence rate. Then, for the decay rate for the CLT variance of the term A_{ℓ} , we have Theorem 4.2 as a refinement of Theorem 5.1 of [14]. This is the key to the gradient replacement and is obtained by applying new results of [9]. We note that Theorem 5.1 of [14] gave only the convergence without rate and was shown based on the so-called characterization of the closed forms originally due to Varadhan [28]; cf. [19]. This is however not sufficient for our purpose, instead we use the results of [9], which were shown by inspired by the recent progress in quantitative homogenization theory. The result of this section is used in the proof of Theorem 3.5.

The choice of K = K(N) together with the second order term $F_N = \Phi_{n(N)}$ of ψ_t and mesoscopic scaling parameters n = n(N) and $\ell = \ell(N)$, $1 \ll n \ll \ell \ll N$, is given in Section 5. We use the results in [9] for the convergence rate of the diffusion matrix approximated from the finite volume and shown uniformly in the density, and an L^{∞} estimate on the local function Φ_n with support size n, which is close to the optimizer of the variational formula (1.5) in F. The Schauder estimates for the solution of the hydrodynamic equation (1.11) are essentially used only in this section.

Sections 6, 7 and 8 are complementary and are devoted to the proofs of the Schauder estimates, the comparison theorem and the non-degeneracy of the diffusion matrix $D(\rho)$, respectively.

2 Proof of Theorem 1.1

To prove Theorem 1.1, we apply the relative entropy method as in [14] comparing the distribution of $\eta^N(t)$ with a properly taken local equilibrium state of second order approximation. Its leading term is determined from the solution $\rho_K(t, v)$ of the hydrodynamic equation (1.11). The second order term is determined by $F = F(\eta)$ which appears in the variational formula (1.5), and plays a similar role to the corrector in the theory of homogenization. Note that, for gradient models, the second order approximation is unnecessary; see [3], [10], [11].

Given a function $\lambda = \lambda(t, v) \in C^{1,3}([0, T] \times \mathbb{T}^d)$ and a function $F = F(\eta) \in \mathcal{F}_0^d$, we define a local equilibrium state $\psi_t(\eta) d\nu^N$ of second order approximation by

(2.1)
$$\psi_t(\eta) \equiv \psi_{\lambda(t,\cdot),F}(\eta) \\ = Z_t^{-1} \exp\bigg\{\sum_{x \in \mathbb{T}_N^d} \lambda(t, x/N)\eta_x + \frac{1}{N} \sum_{x \in \mathbb{T}_N^d} \left(\partial\lambda(t, x/N), \tau_x F(\eta)\right)\bigg\},$$

for $\eta \in \mathcal{X}_N$, where Z_t is the normalization constant with respect to $\nu^N \equiv \nu_{1/2}^N$ and $\partial \lambda \equiv \partial_v \lambda = \{\partial_{v_i} \lambda\}_{1 \leq i \leq d}$. We also write $\dot{\lambda} = \partial \lambda / \partial t$, $\partial^2 \lambda \equiv \partial_v^2 \lambda = \{\partial_{v_i} \partial_{v_j} \lambda\}_{1 \leq i,j \leq d}$ and $\partial^3 \lambda \equiv \partial_v^3 \lambda = \{\partial_{v_i} \partial_{v_j} \partial_{v_k} \lambda\}_{1 \leq i,j,k \leq d}$ for $\lambda = \lambda(t, v)$.

Denote by $f_t(\eta) = f_t^N(\eta)$ the density of the distribution of $\eta^N(t)$ on \mathcal{X}_N with respect to ν^N and consider the relative entropy per volume defined by

(2.2)
$$h_N(t) = N^{-d} H(f_t | \psi_t).$$

We will show that $h_N(t) \leq CN^{-\kappa}$, $t \in [0, T]$ with some $C = C_T > 0$ and $\kappa > 0$ by choosing $\lambda(t, \cdot)$ and F in $\psi_t = \psi_t^N(\eta)$ properly as in the following proposition.

Proposition 2.1. Determine $\lambda(t, v) \equiv \lambda_K(t, v) := \overline{\lambda}(\rho(t, v))$ from the solution $\rho(t, v) = \rho_K(t, v)$ of the hydrodynamic equation (1.11) with K = K(N) and the initial value ρ_0 satisfying (1.12); recall (1.15) for $\overline{\lambda}(\rho)$. Suppose a sequence $K = K(N) \to \infty$ ($N \to \infty$) is given and satisfies $1 \leq K \leq \overline{K}_{\delta}(N) = \delta \log N$. Let $\delta = \delta_T > 0$ and functions $F_N = F_N(\eta) \in \mathcal{F}_0^d$ be chosen as in Section 5.2 below, namely, $\delta_T = \mathfrak{c}/T > 0$ with sufficiently small $\mathfrak{c} > 0$ and $F_N = \Phi_{n(N)}$ with Φ_n satisfying (5.10) and $n(N) = n^{a_1}$ where $a_1 \in (0, 1/(2d + 5))$ is chosen sufficiently small. Consider $h_N(t)$ defined by (2.2) with $\psi_t = \psi_{\lambda(t,\cdot),F_N}$, and assume that $h_N(0) \leq C_1 N^{-\kappa_1}$ for some $C_1 > 0$ and $\kappa_1 > 0$.

Then, there exist some $C = C_T > 0$ and $\kappa > 0$ such that $h_N(t) \leq CN^{-\kappa}$ holds for every $t \in [0,T]$.

The proof of Proposition 2.1 will be given in Section 3. We will calculate the time derivative of $h_N(t)$, in which derivatives of $\lambda_K(t, v)$ appear. For the solution $\rho = \rho_K(t, v)$ of the hydrodynamic equation (1.11) with smooth coefficients D and Kf, we have the Schauder estimates in terms of $K \geq 1$:

(2.3)
$$\|\partial^k \rho\|_{\infty}, \ \|\partial_t \rho\|_{\infty} \|\partial\partial_t \rho\|_{\infty} \le CK^{\alpha_0}, \quad k = 1, 2, 3,$$

for some $\alpha_0 > 0$ and $C = C_T > 0$, where $\|\cdot\|_{\infty} = \|\cdot\|_{L^{\infty}([0,T]\times\mathbb{T}^d)}$; see Section 6. In particular, for $\lambda = \lambda_K(t, v) = \overline{\lambda}(\rho_K(t, v))$, recalling the definition (1.15) of $\overline{\lambda}$ and noting that $\rho_K(t, v)$ is uniformly away from 0 and 1 by Lemma 7.1 (the comparison theorem) below, we have

(2.4)
$$\|\partial\lambda\|_{3,\infty}, \|\dot{\lambda}\|_{\infty}, \|\partial\dot{\lambda}\|_{\infty} \le CK^{\alpha_0}$$

by changing $C = C_T > 0$, where we set

$$\|\partial\lambda\|_{3,\infty} := \|\partial\lambda\|_{\infty} + \|\partial^2\lambda\|_{\infty} + \|\partial^3\lambda\|_{\infty}.$$

The Schauder estimates (2.3), (2.4) are actually used only in Section 5 except for (2.7) and Corollary 3.10 below. But, as these estimates suggest and also as e^{Kt/δ_*} appears in (3.64) after applying Gronwall's inequality at the end of the proof in Section 3.7, at all steps in the proof of Proposition 2.1, we need to obtain error estimates that are strong enough to control diverging factors caused by K. This is useful to show the quantitative version of the hydrodynamic limit as well.

Note that, as the hydrodynamic equation, a discrete PDE was used in [3], [10], [11] to ensure an exact cancellation for the leading term in the entropy computation, but here we use a continuous PDE (1.11), since our error estimates are strong enough to cover the difference.

Proof of Theorem 1.1. Theorem 1.1 is deduced from Proposition 2.1 in the same way as in [3], [14], [30]. In fact, for every $\phi \in C^{\infty}(\mathbb{T}^d)$, $\varepsilon > 0$ and $t \in [0,T]$, consider a subset $\mathcal{A} \equiv \mathcal{A}_{N,\phi,\varepsilon,t}$ of \mathcal{X}_N defined by

(2.5)
$$\mathcal{A} = \left\{ \eta \in \mathcal{X}_N; |\langle \rho^N(t), \phi \rangle - \langle \rho_K(t), \phi \rangle| > \varepsilon \right\}.$$

Then, determining $\psi_t = \psi_t^N$ from $\lambda = \lambda_K$ and $F = F_N$ as in Proposition 2.1, since the assumption of Corollary 3.10 below is satisfied (in particular, the contribution of F_N is negligible), the large deviation estimate holds for $\psi_t d\nu^N$ and thus we have

(2.6)
$$N^{-d}\log P^{\psi_t}(\mathcal{A}) \le -C(\varepsilon),$$

for some $C(\varepsilon) > 0$ and every $N \in \mathbb{N}$. We will sometimes identify ψ_t and f_t with $\psi_t d\nu^N$ and $f_t d\nu^N$, denoted by P^{ψ_t} and P^{f_t} , and the expectations under them by E^{ψ_t} and E^{f_t} , respectively. Note that the condition for the relative entropy at t = 0 in Theorem 1.1 implies that for $h_N(0)$ in Proposition 2.1. Indeed, we have

(2.7)
$$|H(f_0|\psi_0) - H(f_0|\widetilde{\psi}_0)| = |\int f \log(\widetilde{\psi}_0/\psi_0) \, d\nu^N|$$

 $\leq C_2 N^{d-1} ||\partial\lambda||_{\infty} ||F_N||_{\infty} \leq C_3 N^{d-\kappa_2},$

for some $\kappa_2 > 0$, by the Schauder estimates (2.4) and (5.10) noting that $F_N = \Phi_{n(N)}$ with $n(N) = N^{a_1}$, $a_1 < 1/(2d+5)$. Thus, the assumption $h_N(0) \leq C_4 N^{-\kappa_3}$ of Proposition 2.1 (where κ_1, C_1 are replaced by κ_3, C_4) holds taking $\kappa_3 = \kappa_1 \wedge \kappa_2 > 0$, where κ_1 is the constant in Theorem 1.1.

Thus, by applying the entropy inequality and then by Proposition 2.1 and (2.6), we get

$$P^{f_t}(\mathcal{A}) \le \frac{\log 2 + H(f_t|\psi_t)}{\log\{1 + 1/P^{\psi_t}(\mathcal{A})\}} \le \frac{\log 2 + CN^{d-\kappa}}{\log\{1 + e^{C(\varepsilon)N^d}\}} \le C_5 N^{-\kappa}.$$

This completes the proof of Theorem 1.1.

The goal is to show Proposition 2.1, that is, $h_N(t) \leq CN^{-\kappa}$, $t \in [0,T]$ for some $C = C_T > 0$ and $\kappa > 0$.

3 Proof of Proposition 2.1

3.1 Time derivative of the relative entropy

Let us begin with the following lemma in which we calculate the time derivative of $h_N(t)$. If we take K = 0 (i.e. no Glauber part), this lemma coincides with Lemma 3.1 of [14], but the error estimate of order o(1) given there is not sufficient for our purpose and we need to make it more precise.

For $\ell \in \mathbb{N}$ and $x \in \mathbb{T}_N^d$ (or \mathbb{Z}^d), define

(3.1)
$$\Lambda_{\ell,x} := \{ y \in \mathbb{T}_N^d (\text{or } \mathbb{Z}^d); |y - x| \le \ell \}$$

and $\ell_* := 2\ell + 1$. Note that, in case of \mathbb{T}_N^d , by embedding it in \mathbb{Z}^d , the ℓ^1 -norm |y - x| and therefore $\Lambda_{\ell,x}$ are well-defined at least if $\ell_* < N$. Note that $|\Lambda_{\ell,x}| = \ell_*^d$.

For $F = F(\eta) \equiv (F_i(\eta))_{i=1}^d \in \mathcal{F}_0^d$, we set

(3.2)
$$|||F|||_{k,\infty} = r(F)^{d+k} ||F||_{\infty}, \quad k = 0, 1, 2,$$

where $r(F) \geq 1$ denotes the radius (in ℓ^1 -sense, centered at 0) of the support of $F = F(\eta)$, that is,

 $r(F) = \min \{r > 0; \text{supp } F \subset \Lambda_{r,0} \}.$

We use an abbreviation $\sum_{x,y}$ for $\sum_{x,y \in \mathbb{T}_N^d: |x-y|=1}$.

Lemma 3.1. Assume $\lambda \in C^{1,3}([0,T] \times \mathbb{T}^d)$ and $F \in \mathcal{F}_0^d$ for ψ_t in (2.1). Then, setting $\rho(t,v) := \bar{\rho}(\lambda(t,v))$, we have

$$\partial_t h_N(t) \le E^{f_t} [\Omega_1 + \Omega_2] + N^{-d} \sum_{x \in \mathbb{T}_N^d} \dot{\lambda}(t, x/N) \rho(t, x/N) + Q_N^{En}(\lambda, F),$$

where $\Omega_1 = \Omega_1(\eta)$ and $\Omega_2 = \Omega_2(\eta)$ are defined respectively by

$$\begin{split} \Omega_{1} &= -\frac{N^{1-d}}{2} \sum_{x,y} c_{x,y} \Omega_{x,y}, \\ \Omega_{2} &= -N^{-d} \sum_{x \in \mathbb{T}_{N}^{d}} \dot{\lambda}(t, x/N) \eta_{x} + \frac{N^{-d}}{4} \sum_{x,y} c_{x,y} \Omega_{x,y}^{2} \\ &- \frac{N^{-d}}{4} \sum_{x,y} c_{x,y} \sum_{i,j} \partial_{v_{i}} \partial_{v_{j}} \lambda(t, x/N) (y_{i} - x_{i}) (y_{j} - x_{j}) (\eta_{y} - \eta_{x}) \\ &+ \frac{N^{-d}}{2} \sum_{x,y} c_{x,y} \sum_{i,j} \partial_{v_{i}} \partial_{v_{j}} \lambda(t, x/N) \pi_{x,y} (\sum_{z \in \mathbb{T}_{N}^{d}} (z_{j} - x_{j}) \tau_{z} F_{i}) \\ &+ N^{-d} K \sum_{x \in \mathbb{T}_{N}^{d}} \Big\{ \frac{c_{x}^{+}(\eta)}{\rho(t, x/N)} - \frac{c_{x}^{-}(\eta)}{1 - \rho(t, x/N)} \Big\} \{\eta_{x} - \rho(t, x/N)\}, \end{split}$$

and

$$\Omega_{x,y} \equiv \Omega_{x,y}(\eta) = \left(\partial\lambda(t, x/N), (y-x)(\eta_y - \eta_x) - \pi_{x,y}\left(\sum_{z \in \mathbb{T}_N^d} \tau_z F\right)\right).$$

The error term $Q_N^{En}(\lambda, F)$ has the estimate

(3.3)
$$|Q_N^{En}(\lambda, F)| \leq CN^{-1} \left(1 + \|\partial\lambda\|_{3,\infty} \right)^3 \left(1 + \|F\|_{2,\infty} \right)^3 \\ + CN^{-1} K e^{\|\lambda\|_{\infty}} \|\partial\lambda\|_{\infty} \|F\|_{0,\infty} \\ + CN^{-1} \|\partial\dot{\lambda}\|_{\infty} \|F\|_{\infty} + CN^{-1} \|\dot{\lambda}\|_{\infty} \|\partial\lambda\|_{\infty} \|F\|_{0,\infty},$$

as long as λ and F satisfy the bounds

(3.4)
$$N^{-1} \|\partial\lambda\|_{\infty} \le 1, \quad N^{-1} r(F)^d \|\partial\lambda\|_{\infty} \|F\|_{\infty} \le 1,$$

(the constant 1 in the right-hand side may be replaced by any other constant).

Proof. We apply the estimate

(3.5)
$$\partial_t h_N(t) \le N^{-d} \int \psi_t^{-1} \left(\mathcal{L}_N^* \psi_t - \partial_t \psi_t \right) \cdot f_t \, d\nu^N$$

which holds for a large class of Markovian models, where \mathcal{L}_N^* denotes the dual operator of the generator \mathcal{L}_N with respect to the measure ν^N ; see [30] Lemma 1, [23] Lemma 3.1.

The term in the integrand on the right-hand side of (3.5) derived from the Kawasaki part was already computed in the proof of Lemma 3.1 of [14], but only with the error estimate o(1). Let us record the computation (3.2) in [14] here to make the error term clear. Noting that $L_K^* = L_K$,

$$(3.6) \qquad N^{-d}\psi_{t}^{-1}N^{2}L_{K}\psi_{t} \\ = \frac{N^{2-d}}{2}\sum_{x,y}c_{x,y}\left[\exp\left\{\left(\lambda(t,x/N) - \lambda(t,y/N)\right)(\eta_{y} - \eta_{x})\right. + \frac{1}{N}\pi_{x,y}\sum_{z\in\mathbb{T}_{N}^{d}}\left(\partial\lambda(t,z/N),\tau_{z}F\right)\right\} - 1\right] \\ = -\frac{N^{1-d}}{2}\sum_{x,y}c_{x,y}\Omega_{x,y} + \frac{N^{-d}}{4}\sum_{x,y}c_{x,y}\Omega_{x,y}^{2} \\ - \frac{N^{-d}}{4}\sum_{x,y}c_{x,y}\sum_{i,j}\partial_{v_{i}}\partial_{v_{j}}\lambda(t,x/N)(y_{i} - x_{i})(y_{j} - x_{j})(\eta_{y} - \eta_{x}) \\ + \frac{N^{-d}}{2}\sum_{x,y}c_{x,y}\sum_{i,j}\partial_{v_{i}}\partial_{v_{j}}\lambda(t,x/N)\pi_{x,y}(\sum_{z\in\mathbb{T}_{N}^{d}}(z_{j} - x_{j})\tau_{z}F_{i}) + Q_{1,N}, \end{cases}$$

where the error term $Q_{1,N} \equiv Q_{1,N}(\lambda, F)$ can be estimated as

(3.7)
$$|Q_{1,N}| \leq CN^{-1} \left(1 + \|\partial\lambda\|_{3,\infty}\right)^3 \left(1 + \|F\|_{2,\infty}\right)^3,$$

if the condition (3.4) and accordingly $|\varepsilon_N| \leq C$ below are satisfied for some C > 0.

Indeed, to show (3.7), we expand the second to the third lines of (3.6) as

$$\exp\{\varepsilon_N\} - 1 = \varepsilon_N + \frac{1}{2}\varepsilon_N^2 + O(\varepsilon_N^3), \quad |\varepsilon_N| \le C,$$

for $\varepsilon_N \equiv \varepsilon_{N,x,y}$ given in the braces. Then, by Taylor's formula, we have

$$\begin{split} \lambda(t, x/N) - \lambda(t, y/N) &= -\frac{1}{N} \left(\partial \lambda(t, x/N), (y - x) \right) \\ &- \frac{1}{2N^2} (y - x) \cdot \partial^2 \lambda(t, x/N) (y - x) + O\left(N^{-3} \| \partial^3 \lambda \|_{\infty} \right), \end{split}$$

and therefore

$$\varepsilon_N = -\frac{1}{N}\Omega_{xy} + \frac{1}{N}\bar{\Omega}_{xy} - \frac{1}{2N^2}(y-x)\cdot\partial^2\lambda(t,x/N)(y-x)(\eta_y-\eta_x) + O\left(N^{-3}\|\partial^3\lambda\|_{\infty}\right),$$

where

$$\begin{split} \bar{\Omega}_{x,y} &\equiv \bar{\Omega}_{x,y}(\eta) = \pi_{x,y} \sum_{z \in \mathbb{T}_N^d} \left(\partial \lambda(t, z/N) - \partial \lambda(t, x/N), \tau_z F \right) \\ &= N^{-1} \Big(\partial^2 \lambda(t, x/N) \pi_{x,y} \sum_{z \in \mathbb{T}_N^d} (z - x), \tau_z F \Big) + O\Big(N^{-2} \| \partial^3 \lambda \|_{\infty} \Big| \pi_{xy} \sum_{z \in \mathbb{T}_N^d} (z - x)^{\otimes 2}, \tau_z F \Big| \Big). \end{split}$$

Thus, $Q_{1,N} \equiv Q_{1,N}(\lambda, F)$ is given by

$$Q_{1,N} = \frac{N^{2-d}}{2} \sum_{x,y} c_{x,y} O\left(N^{-3} \|\partial^{3}\lambda\|_{\infty} (1+\|\|F\|\|_{2,\infty})\right) + \frac{N^{2-d}}{4} \sum_{x,y} c_{x,y} (\varepsilon_{N}^{2} - N^{-2}\Omega_{x,y}^{2}) + O\left(N^{2-d} \sum_{x,y} c_{x,y} \varepsilon_{N}^{3}\right).$$

The first term is $O(N^{-1} \| \partial^3 \lambda \|_{\infty} (1 + \| F \|_{2,\infty}))$. By noting that $|\Omega_{xy}| \leq \| \partial \lambda \|_{\infty} (1 + \| F \|_{0,\infty})$, the second is bounded by

$$\begin{split} \left| \frac{N^{2-d}}{4} \sum_{x,y} c_{x,y} \left(\varepsilon_N + \frac{1}{N} \Omega_{x,y} \right) \left(\varepsilon_N - \frac{1}{N} \Omega_{x,y} \right) \right| \\ &\leq C N^2 \left(\frac{1}{N^2} \| \partial^2 \lambda \|_{\infty} \| \|F\| \|_{1,\infty} + \frac{1}{N^3} \| \partial^3 \lambda \|_{\infty} \| \|F\| \|_{2,\infty} + \frac{1}{N^2} \| \partial^2 \lambda \|_{\infty} + \frac{1}{N^3} \| \partial^3 \lambda \|_{\infty} \right) \\ &\times \left(\frac{2}{N} \| \partial \lambda \|_{\infty} \left(1 + \| \|F\| \|_{0,\infty} \right) + \frac{1}{N^2} \| \partial^2 \lambda \|_{\infty} \| \|F\| \|_{1,\infty} \right) \\ &\quad + \frac{1}{N^3} \| \partial^3 \lambda \|_{\infty} \| \|F\| \|_{2,\infty} + \frac{1}{N^2} \| \partial^2 \lambda \|_{\infty} + \frac{1}{N^3} \| \partial^3 \lambda \|_{\infty} \right) \\ &\leq C N^{-1} \left(1 + \| \partial \lambda \|_{3,\infty} \right)^2 \left(1 + \| \|F\| \|_{2,\infty} \right)^2. \end{split}$$

Similarly, the third is bounded by

$$CN^{2} \left\{ \frac{1}{N^{3}} \sup_{x,y} |\Omega_{xy}|^{3} + \left(\frac{1}{N^{2}} \|\partial^{2}\lambda\|_{\infty} \|F\|_{1,\infty} + \frac{1}{N^{3}} \|\partial^{3}\lambda\|_{\infty} \|F\|_{2,\infty} \right)^{3} + \left(\frac{1}{N^{2}} \|\partial^{2}\lambda\|_{\infty} \right)^{3} + \left(\frac{1}{N^{3}} \|\partial^{3}\lambda\|_{\infty} \right)^{3} \right\}$$
$$\leq CN^{-1} \left(1 + \|\partial\lambda\|_{3,\infty} \right)^{3} \left(1 + \|F\|_{2,\infty} \right)^{3}.$$

Summarizing these estimates for three terms of $Q_{1,N}(\lambda, F)$, we obtain (3.7) for $Q_{1,N}(\lambda, F)$.

Next, we compute the contribution from the Glauber part. First we note that the dual operator L_G^* of L_G with respect to ν^N is given by

$$L_G^*g(\eta) = \sum_{x \in \mathbb{T}_N^d} \left\{ c_x(\eta^x) g(\eta^x) - c_x(\eta) g(\eta) \right\}.$$

Therefore, we have

(3.8)

$$N^{-d}K\psi_t^{-1}L_G^*\psi_t$$
$$= N^{-d}K\sum_{x\in\mathbb{T}_N^d} \left[c_x(\eta^x) \exp\left\{\pi_x\left(\sum_z \lambda(t,z/N)\eta_z + \frac{1}{N}\sum_{z\in\mathbb{T}_N^d} \left(\partial\lambda(t,z/N), \tau_z F(\eta)\right)\right)\right\} - c_x(\eta) \right]$$

$$= N^{-d}K \sum_{x \in \mathbb{T}_{N}^{d}} \left[c_{x}(\eta^{x}) \exp\left\{\lambda(t, x/N)(1 - 2\eta_{x}) + \frac{1}{N} \sum_{z \in \mathbb{T}_{N}^{d}} \left(\partial\lambda(t, z/N), \pi_{x}\tau_{z}F(\eta)\right)\right\} - c_{x}(\eta) \right]$$
$$= N^{-d}K \sum_{x \in \mathbb{T}_{N}^{d}} \left[c_{x}(\eta^{x}) \exp\left\{\lambda(t, x/N)(1 - 2\eta_{x})\right\} - c_{x}(\eta) \right] + Q_{2,N},$$

However, the leading term can be rewritten as the fifth term of Ω_2 noting (1.3) and (1.14). Indeed, writing $\lambda = \lambda(t, x/N)$, $\rho = \rho(t, x/N)$ and $c_x^{\pm} = c_x^{\pm}(\eta)$ for simplicity, the term inside the brackets in the last sum is rewritten as

$$\left(c_x^+ e^{-\lambda} \mathbf{1}_{\{\eta_x=1\}} + c_x^- e^{\lambda} \mathbf{1}_{\{\eta_x=0\}}\right) - \left(c_x^+ \mathbf{1}_{\{\eta_x=0\}} + c_x^- \mathbf{1}_{\{\eta_x=1\}}\right).$$

Then, we may use $e^{-\lambda} = (1 - \rho)/\rho$, $e^{\lambda} = \rho/(1 - \rho)$, $1_{\{\eta_x = 1\}} = (\eta_x - \rho) + \rho$ and $1_{\{\eta_x = 0\}} = (1 - \rho) - (\eta_x - \rho)$; see Lemma 2.3 of [10].

The error term $Q_{2,N} \equiv Q_{2,N}(\lambda, F)$ in (3.8) can be estimated as

$$(3.9) |Q_{2,N}| \le CN^{-1} K e^{\|\lambda\|_{\infty}} \|\partial\lambda\|_{\infty} \|F\|_{0,\infty},$$

if the second bound in the condition (3.4) is satisfied. Indeed,

$$\exp\{\varepsilon_N\} = 1 + O(\varepsilon_N), \quad |\varepsilon_N| \le C,$$

for $\varepsilon_N \equiv \varepsilon_{N,x} = \frac{1}{N} \sum_{z \in \mathbb{T}_N^d} (\partial \lambda(t, z/N), \pi_x \tau_z F(\eta))$. Thus,

$$Q_{2,N} = N^{-d}K \sum_{x \in \mathbb{T}_N^d} c_x(\eta^x) \exp\left\{\lambda(t, x/N)(1 - 2\eta_x)\right\}$$
$$\times O\left(\frac{1}{N} \sum_{z \in \mathbb{T}_N^d} \left(\partial\lambda(t, z/N), \pi_x \tau_z F(\eta)\right)\right)$$

and therefore we obtain (3.9).

Finally, in (3.5),

$$(3.10) \qquad N^{-d}\psi_t^{-1}\partial_t\psi_t = -N^{-d}Z_t^{-1}\partial_t Z_t + N^{-d}\partial_t \left\{ \sum_{x\in\mathbb{T}_N^d} \lambda(t,x/N)\eta_x + \frac{1}{N}\sum_{x\in\mathbb{T}_N^d} \left(\partial\lambda(t,x/N),\tau_x F\right) \right\}.$$

From (3.6), (3.8) and (3.10), it follows that

(3.11)
$$N^{-d}\psi_t^{-1}\left(\mathcal{L}_N^*\psi_t - \partial_t\psi_t\right) = \Omega_1(\eta) + \Omega_2(\eta) + a(t) + \sum_{i=1}^3 Q_{i,N},$$

where

$$Q_{3,N} \equiv Q_{3,N}(\lambda,F) = -N^{-d-1} \sum_{x \in \mathbb{T}_N^d} (\partial \dot{\lambda}(t,x/N), \tau_x F),$$

and

(3.12)
$$a(t) := N^{-d} Z_t^{-1} \partial_t Z_t = E^{\psi_t} \left[N^{-d} \sum_{x \in \mathbb{T}_N^d} \dot{\lambda}(t, x/N) \eta_x \right] + O\left(N^{-1} \| \partial \dot{\lambda} \|_{\infty} \|F\|_{\infty} \right).$$

The last equality for a(t) is seen from (3.10) by noting that

$$E^{\psi_t} \left[N^{-d} \psi_t^{-1} \partial_t \psi_t \right] = 0$$

and

$$(3.13) |Q_{3,N}| \le N^{-1} \|\partial \dot{\lambda}\|_{\infty} \|F\|_{\infty}$$

Thus,

$$a(t) = N^{-d} \sum_{x \in \mathbb{T}_N^d} \dot{\lambda}(t, x/N) \rho(t, x/N) + Q_{4,N} + O\left(N^{-1} \|\partial \dot{\lambda}\|_{\infty} \|F\|_{\infty}\right),$$

where the error $Q_{4,N} \equiv Q_{4,N}(\lambda, F)$ is given by

$$Q_{4,N} = N^{-d} \sum_{x \in \mathbb{T}_N^d} \dot{\lambda}(t, x/N) \Big(E^{\psi_t} [\eta_x] - \rho(t, x/N) \Big).$$

We only need to estimate $|E^{\psi_t}[\eta_x] - \rho(t, x/N)|$.

To give an estimate for $Q_{4,N} \equiv Q_{4,N}(\lambda, F)$, we note the *r*-Markov property of ψ_t (or P^{ψ_t}). In the next lemma, we denote ψ_t by ψ for simplicity by omitting t and $\frac{1}{N}$ in front of the sum involving F. For $\Lambda \subset \mathbb{T}_N^d$ and $r \in \mathbb{N}$, set $\partial_r \Lambda = \{y \in \mathbb{T}_N^d \setminus \Lambda; \operatorname{dist}(y, \Lambda) \leq r\}$, $\overline{\Lambda} = \Lambda \cup \partial_r \Lambda$, where $\operatorname{dist}(y, \Lambda) = \min\{|x - y|; x \in \Lambda\}$ is defined in the ℓ^1 -sense. We consider Λ with a sufficiently small radius so that $\operatorname{dist}(y, \Lambda)$ is well defined; actually we take $\Lambda = \{x\}$ in (3.18) later. For $\eta \in \{0, 1\}^{\Lambda}$ and $\omega \in \{0, 1\}^{\partial_r \Lambda}$, $\eta \cdot \omega \in \{0, 1\}^{\overline{\Lambda}}$ is defined by $(\eta \cdot \omega)_x = \eta_x$ for $x \in \Lambda$ and $= \omega_x$ for $x \in \partial_r \Lambda$. Define

$$\begin{split} H_{\bar{\Lambda}}(\eta \cdot \omega) &:= \sum_{y \in \bar{\Lambda}} \lambda(y/N)(\eta \cdot \omega)_y + \sum_{y \in \mathbb{T}_N^d: \, \text{supp } \tau_y F \subset \bar{\Lambda}} (\partial \lambda(y/N), \tau_y F(\eta \cdot \omega)), \\ P_{\Lambda}^{\omega}(\eta) &:= Z_{\omega}^{-1} e^{H_{\bar{\Lambda}}(\eta \cdot \omega)}, \quad Z_{\omega} := \sum_{\eta \in \{0,1\}^{\Lambda}} e^{H_{\bar{\Lambda}}(\eta \cdot \omega)}. \end{split}$$

Lemma 3.2. Let $F \in \mathcal{F}_0^d$ and r = r(F). Then, ψ has an r-Markov property, that is, for $\Lambda \subset \mathbb{T}_N^d$ and $\eta \in \{0,1\}^{\Lambda}$,

(3.14)
$$P^{\psi}([\eta]|\mathcal{F}_{\Lambda^c})(\omega) = P^{\bar{\omega}}_{\Lambda}(\eta), \quad P^{\psi}\text{-a.s. }\omega$$

holds, where $[\eta] = \{\zeta \in \{0,1\}^{\mathbb{T}_N^d}; \zeta|_{\Lambda} = \eta\}$ and $\bar{\omega} = \omega|_{\partial_r \Lambda}$. Recall $P^{\psi} = \psi \, d\nu^N$.

Proof. The lemma is well-known and the proof is elementary, but we give it for completeness. Since the right-hand side of (3.14) is \mathcal{F}_{Λ^c} -measurable in ω , it is enough to show

(3.15)
$$E^{\psi}[f(\eta)g(\omega)] = E^{\psi}[g(\omega)E^{P_{\Lambda}^{\omega}}[f]]$$

for every \mathcal{F}_{Λ} -measurable function f and \mathcal{F}_{Λ^c} -measurable function g. The left-hand side of (3.15) is given by

(3.16)
$$Z^{-1} \sum_{\eta \in \{0,1\}^{\Lambda}, \omega \in \{0,1\}^{\Lambda^c}} f(\eta) g(\omega) e^{H_N(\eta \cdot \omega)},$$

while the right-hand side is rewritten as

$$(3.17) Z^{-1} \sum_{\eta \in \{0,1\}^{\Lambda}, \omega \in \{0,1\}^{\Lambda^c}} g(\omega) e^{H_N(\eta \cdot \omega)} Z_{\bar{\omega}}^{-1} \sum_{\zeta \in \{0,1\}^{\Lambda}} f(\zeta) e^{H_{\bar{\Lambda}}(\zeta \cdot \bar{\omega})},$$

where Z is a normalization constant (including the factor $\nu^{N}(\eta \cdot \omega) \equiv 2^{-N}$), and

$$\begin{split} H_N(\eta \cdot \omega) &:= \sum_{y \in \mathbb{T}_N^d} \lambda(y/N)(\eta \cdot \omega)_y + \sum_{y \in \mathbb{T}_N^d} (\partial \lambda(y/N), \tau_y F(\eta \cdot \omega)) \\ &= H_{\bar{\Lambda}}(\eta \cdot \bar{\omega}) + \widetilde{H}_{\bar{\Lambda}^c}(\eta \cdot \omega), \end{split}$$

with $H_{\bar{\Lambda}}(\eta \cdot \bar{\omega})$ defined above and

$$\widetilde{H}_{\bar{\Lambda}^c}(\eta \cdot \omega) = \sum_{y \in \bar{\Lambda}^c} \lambda(y/N)(\eta \cdot \omega)_y + \sum_{y \in \mathbb{T}_N^d : \operatorname{supp} \tau_y F \cap \bar{\Lambda}^c \neq \emptyset} (\partial \lambda(y/N), \tau_y F(\eta \cdot \omega)).$$

However, $(\eta \cdot \omega)_y = \omega_y$ for $y \in \overline{\Lambda}^c$ in the first sum and $\tau_y F(\eta \cdot \omega) = \tau_y F(\omega)$ for $y \in \mathbb{T}_N^d$ such that $\operatorname{supp} \tau_y F \cap \overline{\Lambda}^c \neq \emptyset$ in the second sum, since the radius of the support of F is rso that $\operatorname{supp} \tau_y F \cap \Lambda = \emptyset$. Thus, $\widetilde{H}_{\overline{\Lambda}^c}(\eta \cdot \omega) \equiv \widetilde{H}_{\overline{\Lambda}^c}(\omega)$ is a function of ω only. Therefore, in (3.17),

$$\sum_{\eta \in \{0,1\}^{\Lambda}} e^{H_N(\eta \cdot \omega)} = Z_{\bar{\omega}} \cdot e^{\tilde{H}_{\bar{\Lambda}^c}(\omega)}.$$

Accordingly, the right-hand side is equal to

$$Z^{-1}\sum_{\omega\in\{0,1\}^{\Lambda^c}}g(\omega)\sum_{\zeta\in\{0,1\}^{\Lambda}}f(\zeta)e^{H_{\bar{\Lambda}}(\zeta\cdot\bar{\omega})}e^{\tilde{H}_{\bar{\Lambda}^c}(\omega)},$$

which coincides with (3.16), that is the left-hand side of (3.15). This completes the proof of the lemma.

We now return to the estimate for $Q_{4,N} \equiv Q_{4,N}(\lambda, F)$. Taking $\Lambda = \{x\}$, r = r(F) and $\overline{\Lambda} = \Lambda_{r(F),x}$ in Lemma 3.2, we see

(3.18)
$$E^{\psi_t}[\eta_x] = E^{\psi_t} \left[E^{P_{\{x\}}^{\psi,F}}[\eta_x] \right]$$

and

$$(3.19) \qquad E^{P_{\{x\}}^{\omega,F}}[\eta_x] := \widetilde{Z}_{\omega,F}^{-1} \sum_{\eta_x \in \{0,1\}} \eta_x e^{\lambda(t,x/N)\eta_x + \frac{1}{N}\sum_{y \in \mathbb{T}_N^d: \operatorname{supp} \tau_y F \subset \overline{\Lambda}}(\partial\lambda(t,y/N), \tau_y F(\eta_x \cdot \omega))}$$
$$=: \widetilde{Z}_{\omega,F}^{-1} A_{\omega,F},$$

for $\omega \in \{0,1\}^{\bar{\Lambda} \setminus \{x\}}$, where

(3.20)
$$\widetilde{Z}_{\omega,F} := \sum_{\eta_x \in \{0,1\}} e^{\lambda(t,x/N)\eta_x + \frac{1}{N}\sum_{y \in \mathbb{T}_N^d: \operatorname{supp} \tau_y F \subset \bar{\Lambda}} (\partial\lambda(t,y/N), \tau_y F(\eta_x \cdot \omega))}.$$

Note that $\sum_{y \in \bar{\Lambda} \setminus \{x\}} \lambda(y/N) \omega_y$ in $H_{\bar{\Lambda}}(\eta \cdot \omega)$ cancels with that in Z_{ω} and gives the formula (3.19) with $\tilde{Z}_{\omega,F}$ determined by (3.20).

Noting that $E^{P_{\{x\}}^{\omega,0}}[\eta_x] = \rho(t, x/N)$ taking $F \equiv 0$, we have

$$E^{P_{\{x\}}^{\omega,F}}[\eta_x] - \rho(t,x/N) = \widetilde{Z}_{\omega,F}^{-1}A_{\omega,F} - \widetilde{Z}_{\omega,0}^{-1}A_{\omega,0}$$

A simple estimate shows

$$\begin{aligned} \widetilde{Z}_0 e^{-\frac{c}{N}r(F)^d} \|\partial\lambda(t,\cdot)\|_{\infty} \|F\|_{\infty} &\leq \widetilde{Z}_{\omega,F} \leq \widetilde{Z}_0 e^{\frac{c}{N}r(F)^d} \|\partial\lambda(t,\cdot)\|_{\infty} \|F\|_{\infty}, \\ A_0 e^{-\frac{c}{N}r(F)^d} \|\partial\lambda(t,\cdot)\|_{\infty} \|F\|_{\infty} &\leq A_{\omega,F} \leq A_0 e^{\frac{c}{N}r(F)^d} \|\partial\lambda(t,\cdot)\|_{\infty} \|F\|_{\infty}, \end{aligned}$$

for $A_0 = A_{\omega,0}$, $\widetilde{Z}_0 = \widetilde{Z}_{\omega,0}$ and some c > 0. Therefore, taking the expectation in ω under P^{ψ_t} , we obtain by (3.18)

$$|E^{\psi_t}[\eta_x] - \rho(t, x/N)| = \left| E^{\psi_t} \left[\frac{A_0}{\widetilde{Z}_{\omega,F}} \left(\frac{A_{\omega,F}}{A_0} - \frac{Z_{\omega,F}}{\widetilde{Z}_0} \right) \right] \right|$$
$$\leq C_1 \left| e^{\frac{c}{N}r(F)^d \|\partial\lambda(t,\cdot)\|_{\infty} \|F\|_{\infty}} - 1 \right|$$
$$\leq C_2 N^{-1} r(F)^d \|\partial\lambda\|_{\infty} \|F\|_{\infty},$$

if F satisfies the condition (3.4): $N^{-1}r(F)^d \|\partial\lambda\|_{\infty} \|F\|_{\infty} \leq 1$; note that $0 \leq A_0/\widetilde{Z}_{\omega,F} = A_0/\widetilde{Z}_0 \cdot \widetilde{Z}_0/\widetilde{Z}_{\omega,F} \leq e^c$ under this condition. This leads to the bound on $Q_{4,N} \equiv Q_{4,N}(\lambda,F)$:

$$(3.21) |Q_{4,N}| \le CN^{-1} \|\dot{\lambda}\|_{\infty} \|\partial\lambda\|_{\infty} \|F\|_{0,\infty}.$$

By (3.7), (3.9), (3.13) and (3.21), we obtain (3.3), and this concludes the proof of Lemma 3.1. $\hfill \Box$

3.2 Refined one-block estimate

We now show the one-block estimate, that is, the error estimate for the replacement of the microscopic function Ω_2 of order O(1) (first four terms) and O(K) (last term) by its ensemble average with mean $\rho = \bar{\eta}_x^{\ell}$ (see (3.22) below) under the integral with respect to P^{f_t} and also in $t \in [0, T]$. As we pointed in Section 2, we need to prepare an error estimate strong enough for later use. The classical one-block estimate can be shown even with the diverging Glauber part as discussed in Remark 3.1 below, but it is not sufficient for our purpose.

Indeed, we will show a refined one-block estimate in Theorem 3.3 below in a similar manner to Theorem 1.4 of [10], but, differently from it, we don't need the relative entropy on the right-hand side of the estimate (3.25).

For $h \in \mathcal{F}_0$, set

$$f_x(\eta) \equiv f_x^{\ell}(\eta) = \tau_x h(\eta) - \langle h \rangle (\bar{\eta}_x^{\ell})$$

where we denote $\langle \cdot \rangle(\rho)$ for $\langle \cdot \rangle_{\rho}$, $\rho \in [0,1]$. Define a local sample average of η over $\Lambda_{\ell,x}$

(3.22)
$$\bar{\eta}_x^{\ell} \equiv \bar{\eta}_{\Lambda_{\ell,x}} := \ell_*^{-d} \sum_{y \in \Lambda_{\ell,x}} \eta_y,$$

recall (3.1) for $\Lambda_{\ell,x}$ and ℓ_* (< N).

Theorem 3.3. Let $a_{t,x}$, $t \in [0,T]$ and $x \in \mathbb{T}_N^d$, be deterministic coefficients which satisfy

$$(3.23) |a_{t,x}| \le M for all \ t \in [0,T] \ and \ x \in \mathbb{T}_N^d.$$

 $We \ assume$

(3.24)
$$K^{3/2}\ell_*^{(d+2)/2} \le \delta N$$

for some $\delta > 0$ small enough (see Lemma 3.4 below for the choice of δ) and for all N large enough. Then, there exists C > 0 such that for all N large enough,

(3.25)
$$E\left[\left|\int_{0}^{T} N^{-d} \sum_{x \in \mathbb{T}_{N}^{d}} a_{t,x} f_{x}(\eta^{N}(t)) dt\right|\right] \leq C(M \|h\|_{\infty} + 1) \left(N^{-1} K^{1/2} \ell^{(d+2)/2} + \ell^{-d} r(h)^{d}\right).$$

Recall that r(h) denotes the radius of the support of h.

We will apply this theorem for Ω_2 in the proof of Lemma 3.8 later.

Proof. We follow the argument in Section 3 of [10], where the same Glauber-Kawasaki dynamics $\eta^{N}(t)$ was considered. Note that although the gradient condition was assumed in [10], it was not used in Section 3. We first decompose as

$$(3.26) \qquad E\left[\left|\int_{0}^{T}N^{-d}\sum_{x\in\mathbb{T}_{N}^{d}}a_{t,x}f_{x}dt\right|\right] \leq E\left[\left|\int_{0}^{T}N^{-d}\sum_{x\in\mathbb{T}_{N}^{d}}a_{t,x}m_{x}dt\right|\right] + E\left[\left|\int_{0}^{T}N^{-d}\sum_{x\in\mathbb{T}_{N}^{d}}a_{t,x}E^{\nu_{1/2}}[f_{x}\mid\bar{\eta}_{x}^{\ell}]dt\right|\right],$$

see (3.1) of [10], where $\nu_{1/2}$ is the Bernoulli measure with $\rho = 1/2$ on \mathcal{X} and

$$m_x := f_x - E^{\nu_{1/2}}[f_x \mid \bar{\eta}_x^{\ell}]$$

Then, for the first term in (3.26), Lemma 3.1 of [10] holds; we take $\Theta = 0, \theta = 1, \varepsilon(N) \downarrow 0$ (so that $\varepsilon^{1-\Theta} = 0, \varepsilon^{2\Theta} = 1$) and get the next Lemma 3.4. Note that the proof is the same, except that at the last point we estimate

$$\sup_{t,x,\eta} |a_{t,x}m_x|^2 \le (2M)^2 ||h||_{\infty}^2$$

Lemma 3.4. Let $\gamma = \gamma(N) > 0$. If $\gamma K \ell_*^{d+2} \leq \delta N^2$ for some $\delta > 0$ small enough, then for all N large enough

$$E\left[\left|\int_0^T N^{-d} \sum_{x \in \mathbb{T}_N^d} a_{t,x} m_x dt\right|\right] \le C\left(\frac{K}{\gamma} + \frac{\gamma \ell_*^{d+2}}{N^2} M^2 \|h\|_\infty^2\right),$$

for some C > 0. In particular, choosing $\gamma = NK^{1/2}\ell_*^{-(d+2)/2}/(M\|h\|_{\infty}+1)$, the right-hand side is bounded by

$$2CN^{-1}K^{1/2}\ell_*^{(d+2)/2}(M||h||_{\infty}+1).$$

For this choice of γ , the above condition for γ is satisfied if (3.24) stated above holds.

Note that the condition $\gamma K \ell_*^{d+2} \leq \delta N^2$ for some $\delta > 0$ small enough' was used for the denominator in (3.10) of [10] (with $\theta = 1, \varepsilon^{-\Theta} = 1$), derived by Rayleigh estimate, to stay uniformly positive, say $\geq 1/2$. This determines how small $\delta > 0$ needs to be.

Next, by Theorem 4.1 of [2] which provides the equivalence of ensembles for general Gibbs measures with precise convergence speed (cf. Lemma 3.2 of [10]), we have the following estimate. There exists C > 0 such that for all N sufficiently large

(3.27)
$$\sup_{\rho \in [0,1]} \max_{x \in \mathbb{T}_N^d} \sup_{\eta \in \mathcal{X}_N} \left| E^{\nu_{\rho}}[\tau_x h \mid \bar{\eta}_x^\ell] - \langle h \rangle(\bar{\eta}_x^\ell) \right| \le \frac{C}{\ell^d} |\Delta| \, \|h\|_{\infty},$$

where the local function h has a support in Δ such that $|\Delta| \leq \ell^{d(1-\delta)}$. Note that one may take h for f_x in Lemma 3.2 of [10]. Combining the two estimates given in Lemma 3.4 and (3.27), the proof of Theorem 3.3 is concluded.

Remark 3.1. In the estimate (4.10) of [8], taking $\nu_t \equiv \nu^N (= \nu_{1/2}^N)$, the second term on the right-hand side vanishes and also, in the first term $-\mathcal{D}(\sqrt{f_t}; \nu^N)$, one can drop the contribution of the Glauber part (note that this term is not really the Dirichlet form associated with the Glauber generator in the sense that ν^N is not in general a reversible measure of the Glauber part), since it is non-positive in view of (4.9) of [8]. Since a classical one-block estimate (and also two-blocks estimate) was shown relying only on the estimate obtained in this way, we have the same result as Theorem 2.7 of [8] or Theorem 3.1 of [14] even with a diverging Glauber part in our generator \mathcal{L}_N . In other words, one can show that the expectation in (3.25) with $a_{t,x} \equiv 1$ tends to 0 as $N \to \infty$ and then $\ell \to \infty$. But, this is not enough for our purpose.

3.3 Boltzmann-Gibbs principle for gradient replacement

For the microscopic function Ω_1 which is given in Lemma 3.1 and looks O(N), we need the following Boltzmann-Gibbs principle for the gradient replacement. This is a refinement of Theorem 3.2 of [14] adding the Glauber part and giving the necessary error estimates.

Theorem 3.5. There exists C > 0 such that

$$\int_0^T E^{f_t} \left[\Omega_1 + N^{1-d} \sum_{x \in \mathbb{T}_N^d} \left(D(\bar{\eta}_x^\ell) \partial \lambda(t, x/N), \ell_*^{-d} \tau_x A_\ell \right) \right]$$

$$-\frac{\beta N^{-d}}{2} \sum_{x \in \mathbb{T}_N^d} \left(\partial \lambda(t, x/N), R(\bar{\eta}_x^\ell; F) \partial \lambda(t, x/N) \right) \right] dt$$
$$\leq \frac{C}{\beta} K + \beta^2 Q_{N,\ell}^{(1)}(\lambda, F) + \beta Q_\ell^{(2)}(\lambda, F)$$

for every $\beta > 0$, where

(3.28)
$$A_{\ell} = \{A_{\ell,i}(\eta)\}_{i=1}^d := \sum_{b=\{x,y\}\in(\Lambda(\ell))^*} (\eta_y - \eta_x)(y-x),$$

denoting the box $\Lambda_{\ell,0}$ with the center at 0 by $\Lambda(\ell)$, and

(3.29)
$$R(\rho; F) = \widehat{c}(\rho; F) - \widehat{c}(\rho).$$

The error $Q_{N,\ell}^{(1)} \equiv Q_{N,\ell}^{(1)}(\lambda,F)$ is estimated as

(3.30)
$$|Q_{N,\ell}^{(1)}| \le CN^{-1}\ell^{(2d+4)} \|\partial\lambda\|_{\infty}^{3} (1+\||F|\|_{0,\infty}^{3}).$$

The other error $Q_{\ell}^{(2)} \equiv Q_{\ell}^{(2)}(\lambda, F)$ appears in the computation of the central limit theorem variance (see Section 4) and has a bound

$$(3.31) |Q_{\ell}^{(2)}| \le \|\partial\lambda\|_{\infty}^2 Q_{\ell}(F)$$

with $Q_{\ell}(F)$ bounded in Corollary 4.3 below.

Before giving a proof of Theorem 3.5, we prepare a key lemma by which one can reduce a non-equilibrium problem into a static problem under the canonical equilibrium measure, sometimes called the Kipnis-Varadhan estimate or the Itô-Tanaka trick. We introduce some notation. For $\Lambda \in \mathbb{Z}^d$ (i.e., finite subset) and $m \in [0, |\Lambda|] \cap \mathbb{Z}$,

$$\mathcal{X}_{\Lambda} = \{0,1\}^{\Lambda}, \quad \mathcal{X}_{\Lambda,m} = \{\eta \in \mathcal{X}_{\Lambda}; \sum_{x \in \Lambda} \eta_x = m\},$$

 $\nu_{\Lambda,m}$ = uniform probability measure on $\mathcal{X}_{\Lambda,m}$,

and $\langle \cdot \rangle_{\Lambda,m}$ denotes the expectation with respect to $\nu_{\Lambda,m}$. Then, we have a result similar to Lemmas 3.3 (the case $M(\rho) = 0$) and 6.1 of [14], but the last term $\frac{C}{\beta}$ in their estimates is replaced by $\frac{C}{\beta}K$ due to the contribution from the Glauber part. We also show an estimate for the error term instead of taking the limit in N as in [14].

Given a positive integer ℓ , we write $G_{\zeta} = G_{\zeta}(\xi)$ for a function $G = G(\eta)$ when we consider G as a function of $\xi = \eta|_{\Lambda(\ell)}$ regarding $\zeta = \eta|_{\Lambda(\ell)^c}$ as a parameter. The following lemma will be used for the proof of Theorem 3.5 taking n = d and for the proof of Lemma 3.7 below taking n = 1.

Lemma 3.6. Let $n \in \mathbb{N}$, $J(t, v) = \{J_i(t, v)\}_{i=1}^n \in C^{\infty}([0, T] \times \mathbb{T}^d, \mathbb{R}^n)$, $G(\eta) = \{G_i(\eta)\}_{i=1}^n \in \mathcal{F}_0^n \text{ and } M(\rho) = \{M_{ij}(\rho)\}_{1 \leq i,j \leq n} \in C([0, 1], \mathbb{R}^n \otimes \mathbb{R}^n)$. Let $1 \leq \ell \leq N/2$ (in fact,

 $\max_{|y|=1} r(c_{0,y}) < N - \ell$ is enough). Suppose that $\langle G_{\zeta} \rangle_{\Lambda(\ell),m} = 0$ for every ζ and $m: 0 \leq m \leq \ell_*^d$. Then, there exists C > 0 such that for every $\beta > 0$

$$\begin{split} \int_0^T E^{f_t} & \left[N^{1-d} \sum_{x \in \mathbb{T}_N^d} J(t, x/N) \cdot \tau_x G(\eta) - \beta N^{-d} \sum_{x \in \mathbb{T}_N^d} J(t, x/N) \cdot M(\bar{\eta}_x^\ell) J(t, x/N) \right] dt \\ & \leq \beta T \sup_{|\theta| \leq \|J\|_{\infty}} \sup_{m, \zeta \in \mathcal{X}_{\Lambda(\ell)^c}} \left[d\,\ell_*^d \big\langle \theta \cdot G_\zeta \left(-L_{\Lambda(\ell), \zeta} \right)^{-1} \theta \cdot G_\zeta \big\rangle_{\Lambda(\ell), m} - \theta \cdot M(m/\ell_*^d) \theta \right] \\ & \quad + \frac{C}{\beta} K + \beta^2 Q_{N, \ell}^{(1)}(J, G), \end{split}$$

where $L_{\Lambda(\ell),\zeta}$ is the operator defined by

(3.32)
$$L_{\Lambda(\ell),\zeta}f(\xi) = \sum_{b \in (\Lambda(\ell))^*} c_b(\xi \cdot \zeta) \pi_b f(\xi),$$

for $\xi \in \mathcal{X}_{\Lambda(\ell)}$, $\zeta \in \mathcal{X}_{\Lambda(\ell)^c}$, f is a function on $\mathcal{X}_{\Lambda(\ell)}$ and $\xi \cdot \zeta$ denotes the configuration $\eta \in \mathcal{X}$ such that $\eta|_{\Lambda(\ell)} = \xi$ and $\eta|_{\Lambda(\ell)^c} = \zeta$. The error term $Q_{N,\ell}^{(1)} \equiv Q_{N,\ell}^{(1)}(J,G)$ has a bound

(3.33)
$$|Q_{N,\ell}^{(1)}| \le CN^{-1}\ell^{(2d+4)} ||G||_{\infty}^{3} ||J||_{\infty}^{3}.$$

Proof. The proof is given by combining that of Lemma 6.1 of [14] and the calculation developed in that of Lemma 3.1 of [10] for Glauber-Kawasaki dynamics (see also Lemma 7.6 of [3] for Glauber-Zero range process).

Setting

$$W_{N,t}(\eta) = N \sum_{x \in \mathbb{T}_N^d} J(t, x/N) \cdot \tau_x G - \beta \sum_{x \in \mathbb{T}_N^d} J(t, x/N) \cdot M(\bar{\eta}_x^\ell) J(t, x/N),$$

by the entropy inequality (see (6.2) of [14]) noting that $H(P_{f_0}|P_{\nu^N}) \leq CN^d$ (P_{f_0} denotes the distribution on the path space $D([0,T], \mathcal{X}_N)$ of $\eta^N(\cdot)$ with the initial distribution $f_0 d\nu^N$) at the process level, we have

$$\int_{0}^{T} E^{f_{t}} [N^{-d}W_{N,t}(\eta)] dt \leq \frac{1}{\beta} N^{-d} \log E_{\nu^{N}} \left[e^{\int_{0}^{T} \beta W_{N,t} dt} \right] + \frac{C}{\beta},$$

for every $\beta > 0$; recall $\nu^N = \nu_{1/2}^N$. Then, similar to (3.6) (with $\varepsilon = 0, \varepsilon^{1-\Theta} = 0$) of [10] shown for Glauber-Kawasaki dynamics, the right-hand side is bounded by

$$\frac{1}{\beta}N^{-d}\int_0^T \Omega_{N,\beta}(t)dt + \frac{C}{\beta},$$

where

$$\Omega_{N,\beta}(t) = \sup_{\psi: \int \psi^2 d\nu^N = 1} \Big\{ E^{\nu^N} [\beta W_{N,t} \psi^2] - E^{\nu^N} [\psi(-\mathcal{L}_N \psi)] \Big\}.$$

Note that

$$E^{\nu^{N}}[\psi(-\mathcal{L}_{N}\psi)] = N^{2}E^{\nu^{N}}[\psi(-L_{K}\psi)] + KE^{\nu^{N}}[\psi(-L_{G}\psi)]$$

For the Glauber part, it was shown in the proof of Lemma 3.1 of [10] that

$$E^{\nu^N}[\psi(-L_G\psi)] \ge -CN^d.$$

This bound gives $\frac{C}{\beta}K$ in the final form of the estimate.

The contribution of the Kawasaki part is computed in [14] and the same bound holds in the present setting until the very end of the proof of Lemma 6.1 of [14]. In fact, (6.3) of [14] is the same as our $\Omega_{N,\beta}(t)$ without the contribution of the Glauber part. The only difference is that in (6.7) of [14], the limsup was taken as $N \to \infty$, but we have to derive the error estimate. The error is given by the second term in the estimate given in Theorem 6.1 of [14] (the Rayleigh-Schrödinger bound):

$$Q_{N,\ell,\beta}(V) := \frac{N^2 T}{\beta |(\Lambda(\ell))^*|} \times 4 ||V||_{\infty}^3 \left(\sup_{f:\langle f \rangle_{\Lambda(\ell),m}=0} \frac{\langle f^2 \rangle_{\Lambda(\ell),m}}{\langle -fL_{\Lambda(\ell),\zeta}f \rangle_{\Lambda(\ell),m}} \right)^2,$$

where $V = N^{-1}\beta |(\Lambda(\ell))^*| \theta \cdot G$; note that the front factor $\frac{N^2T}{\beta |(\Lambda(\ell))^*|}$ appears as in ℓ . 4, p. 28 of [14] and because of the time integral. Since we have a spectral gap estimate for the Kawasaki generator:

(3.34)
$$\langle -fL_{\Lambda(\ell),\zeta}f\rangle_{\Lambda(\ell),m} \ge C_0 \ell^{-2} \langle f^2 \rangle_{\Lambda(\ell),m},$$

with a constant $C_0 > 0$ independent of ζ, m, ℓ (see [22] or Appendix A of [14]), we have a bound for the error term:

$$\begin{aligned} |Q_{N,\ell,\beta}(V)| &\leq 4T N^{-1} \beta^2 |\Lambda(\ell)^*|^2 \|\theta \cdot G\|_{\infty}^3 \left(C_0^{-1} \ell^2\right)^2 \\ &\leq C N^{-1} \beta^2 \ell^{(2d+4)} \|G\|_{\infty}^3 \|J\|_{\infty}^3, \end{aligned}$$

for $\theta: |\theta| \leq ||J||_{\infty}$. This completes the proof of the lemma.

When K = 0 (i.e. no Glauber part), the proof of Theorem 3.5 without error estimates was given in Sections 4-6 of [14]. Since only the Kawasaki part appears in the first term of the estimate in Lemma 3.6 above, the arguments in Sections 4 and 5 of [14] are applicable in our setting; see Section 4 below for more details and refined estimates.

To rewrite the function in the expectation in Theorem 3.5, we introduce the following three \mathbb{R}^d -valued functions A_ℓ , B_ℓ , and H_ℓ of η ; in fact, A_ℓ is the same as in (3.28). We denote $\Psi_i = \eta_{e_i} - \eta_0$ and $W_{x,y}(\eta) = c_{x,y}(\eta)(\eta_y - \eta_x)$ for the microscopic current, where $e_i \in \mathbb{Z}^d$ stands for the unit vector in the *i*-th direction.

(3.35)
$$A_{\ell}(\eta) \equiv A_{\ell}(\xi) = \frac{1}{2} \sum_{x,y \in \Lambda(\ell): |x-y|=1} (\xi_{y} - \xi_{x})(y-x)$$
$$= \sum_{i=1}^{d} \sum_{x:\tau_{x}e_{i}^{*} \in (\Lambda(\ell))^{*}} \tau_{x}\Psi_{i}(\xi) e_{i},$$
(3.36)
$$B_{\ell}(\eta) \equiv B_{\ell,\zeta}(\xi) = \frac{1}{2} \sum_{x,y \in \Lambda(\ell): |x-y|=1} W_{x,y}(\eta)(y-x)$$

$$= -L_{\Lambda(\ell),\zeta}(\sum_{x \in \Lambda(\ell)} x\xi_x),$$

and

(3.37)
$$H_{\ell}(\eta) \equiv H_{\ell,\zeta,F}(\xi) = \sum_{x \in \Lambda(\ell-n)} \tau_x(L^N F)(\xi \cdot \zeta)$$
$$= L_{\Lambda(\ell),\zeta} \left(\sum_{x \in \Lambda(\ell-n)} \tau_x F\right)(\xi),$$

for a given function $F = (F_i)_{i=1}^d \in \mathcal{F}_0^d$ which is $\mathcal{F}_{\Lambda(n-1)}$ -measurable with n = r(F) + 1.

Proof of Theorem 3.5. We apply Lemma 3.6 with $n = d, J(t, v) = \partial \lambda(t, v)$ and

(3.38)

$$G(\eta) = \ell_*^{-d} \left\{ D(\bar{\eta}_\ell) A_\ell - B_\ell + H_\ell \right\},$$

$$M(\rho) = \frac{1}{2} \left\{ \widehat{c}(\rho; F) - \widehat{c}(\rho) \right\} \left(= \frac{1}{2} R(\rho; F) \right)$$

Then, we have $||G||_{\infty} \leq C(1 + |||F|||_{0,\infty})$. Indeed, the function $D = D(\rho)$ is bounded from (1.9), $||A_{\ell}||_{\infty} \leq C\ell_*^d$, $||B_{\ell}||_{\infty} \leq C\ell_*^d$ since $W_{x,y}$ is bounded, and from

$$H_{\ell} = \sum_{\{x,y\} \in \Lambda(\ell)^*} c_{xy}(\xi \cdot \zeta) \pi_{xy}\Big(\sum_{z \in \Lambda(\ell-n)} \tau_z F\Big),$$

with n = r(F) + 1, we have

$$||H_{\ell}||_{\infty} \le C\ell_*^d r(F)^d ||F||_{\infty} = C\ell_*^d ||F||_{0,\infty}.$$

Thus, by (3.33), the error $Q_{N,\ell}^{(1)}(\lambda, F) := Q_{N,\ell}^{(1)}(J,G)$ with J and G determined as above has the estimate (3.30).

Then the proof is concluded by observing that

$$Q_{\ell}^{(2)}(\|J\|_{\infty},F) := \sup_{|\theta| \le \|J\|_{\infty}} \sup_{m,\zeta \in \mathcal{X}_{\Lambda(\ell)^{c}}} \left\{ \ell_{*}^{d} \langle \theta \cdot G_{\zeta}, (-L_{\Lambda(\ell),\zeta})^{-1} \theta \cdot G_{\zeta} \rangle_{\Lambda(\ell),m} - \theta \cdot M(m/\ell_{*}^{d}) \theta \right\}$$

is bounded by $||J||_{\infty}^{2}|Q_{\ell}(F)|$ with $Q_{\ell}(F)$ given in Corollary 4.3. Taking $J = \partial \lambda$ and denoting $Q_{\ell}^{(2)}(\lambda, F) := Q_{\ell}^{(2)}(||\partial \lambda||_{\infty}, F)$, we obtain (3.31).

Next, we show the following lemma, in which we evaluate the error to rewrite the term appearing in Theorem 3.5 from Ω_1 into another term of order O(1).

Lemma 3.7. Set

$$\begin{aligned} Q_{N,\ell}^{(3)}(\lambda) &= \int_0^T E^{f_t} \bigg[N^{1-d} \sum_{x \in \mathbb{T}_N^d} \left(D(\bar{\eta}_x^\ell) \partial \lambda(t, x/N), \ell_*^{-d} \tau_x A_\ell(\eta) \right) \\ &+ N^{-d} \sum_{x \in \mathbb{T}_N^d} \sum_{i,j=1}^d \partial_{v_i} \partial_{v_j} \lambda(t, x/N) P_{ij}(\bar{\eta}_x^\ell) \bigg] \, dt, \end{aligned}$$

where $P(\rho) \equiv \{P_{ij}(\rho)\}_{1 \leq i,j \leq d} := \int_0^{\rho} D(\rho') d\rho', \ \rho \in [0,1].$ Then, we have the following upper bound for $Q_{N,\ell}^{(3)} \equiv Q_{N,\ell}^{(3)}(\lambda)$:

$$Q_{N,\ell}^{(3)} \le C(\ell^{-1} \|\partial^2 \lambda\|_{\infty} + N^{-1} \|\partial^3 \lambda\|_{\infty}) + C\beta\ell^{-1} \|\partial \lambda\|_{\infty}^2 + \frac{C}{\beta}K + C\beta^2 N^{-1}\ell^{2d-2} \|\partial \lambda\|_{\infty}^3$$

for every $\beta > 0$.

Proof. We essentially follow the proof of Lemma 3.4 of [14], but make it more precise. Denote the first and second terms in the expectation $E^{f_t}[\cdots]$ by $I_1^N = I_1^N(t,\eta)$ and $I_2^N = I_2^N(t,\eta)$, respectively. Then, similar to the proof of Lemma 3.4 of [14] (but, with the - sign), we have the decomposition of I_1^N :

(3.39)
$$I_1^N = J_+^N - J_-^N,$$

where

$$J^N_{\pm} = N^{1-d} \ell^{-d}_* \sum_{x \in \mathbb{T}^d_N} \sum_{i,j=1}^d D_{ij}(\bar{\eta}^\ell_x) \partial_{v_j} \lambda(t, x/N) \tau_{x \pm \ell e_i} \widetilde{A}_{\ell,i}(\eta),$$

and

$$\widetilde{A}_{\ell,i} = \sum_{x \in \mathbb{Z}^d: x_i = 0, |x_j| \leq \ell \text{ for } j \neq i} \eta_x$$

However, J_{-}^{N} can be rewritten as

$$J_{-}^{N} = N^{1-d} \ell_{*}^{-d} \sum_{x \in \mathbb{T}_{N}^{d}} \sum_{i,j=1}^{d} D_{ij}(\bar{\eta}_{x-e_{i}}^{\ell}) \partial_{v_{j}} \lambda(t, (x-e_{i})/N) \tau_{x-(\ell+1)e_{i}} \widetilde{A}_{\ell,i}$$
$$= N^{1-d} \ell_{*}^{-d} \sum_{x \in \mathbb{T}_{N}^{d}} \sum_{i,j=1}^{d} D_{ij}(\bar{\eta}_{x-e_{i}}^{\ell}) \partial_{v_{j}} \lambda(t, x/N) \tau_{x-(\ell+1)e_{i}} \widetilde{A}_{\ell,i} + Q_{1,\ell},$$

with an error $Q_{1,\ell} \equiv Q_{1,\ell}(\lambda)$ bounded as

$$|Q_{1,\ell}| \le C_1 \ell^{-1} \|\partial^2 \lambda\|_{\infty},$$

by noting $\partial_{v_j}\lambda(t, (x-e_i)/N) = \partial_{v_j}\lambda(t, x/N) + O(\|\partial^2\lambda\|_{\infty}/N)$ and $\tau_{x-(\ell+1)e_i}\widetilde{A}_{\ell,i} = O(\ell^{d-1})$. Therefore, by (3.39), we obtain

$$\begin{split} I_{1}^{N} = & N^{1-d} \ell_{*}^{-d} \sum_{x \in \mathbb{T}_{N}^{d}} \sum_{i,j=1}^{d} \left\{ D_{ij}(\bar{\eta}_{x}^{\ell}) \tau_{x+\ell e_{i}} \widetilde{A}_{\ell,i} \right. \\ & \left. - D_{ij}(\bar{\eta}_{x-e_{i}}^{\ell}) \tau_{x-(\ell+1)e_{i}} \widetilde{A}_{\ell,i} \right\} \partial_{v_{j}} \lambda(t, x/N) - Q_{1,\ell}. \end{split}$$

On the other hand, since $\partial_{v_i}\partial_{v_j}\lambda(t, x/N) = N\{\partial_{v_j}\lambda(t, (x+e_i)/N) - \partial_{v_j}\lambda(t, x/N)\} + O(\|\partial^3\lambda\|_{\infty}/N),$

$$I_2^N = -N^{1-d} \sum_{x \in \mathbb{T}_N^d} \sum_{i,j=1}^d \left\{ P_{ij}(\bar{\eta}_x^\ell) - P_{ij}(\bar{\eta}_{x-e_i}^\ell) \right\} \partial_{v_j} \lambda(t, x/N) + Q_{2,N},$$

with an error $Q_{2,N} \equiv Q_{2,N}(\lambda)$ bounded as

$$|Q_{2,N}| \le C_2 N^{-1} \|\partial^3 \lambda\|_{\infty}.$$

Note that $P_{ij}(\rho)$ is bounded.

From these two equalities for I_1^N and I_2^N , $Q_{N,\ell}^{(3)}(\lambda)$ in the statement of the lemma is rewritten as

(3.40)
$$\int_{0}^{T} E^{f_{t}} \left[N^{1-d} \sum_{x \in \mathbb{T}_{N}^{d}} \sum_{i,j=1}^{d} \partial_{v_{j}} \lambda(t, x/N) Q_{ij;\ell}(x, \eta) - Q_{1,\ell} + Q_{2,N} \right] dt,$$

where

$$Q_{ij;\ell}(x,\eta) = -P_{ij}(\bar{\eta}_x^{\ell}) + P_{ij}(\bar{\eta}_{x-e_i}^{\ell}) + \ell_*^{-d} \left\{ D_{ij}(\bar{\eta}_x^{\ell}) \tau_{x+\ell e_i} \widetilde{A}_{\ell,i} - D_{ij}(\bar{\eta}_{x-e_i}^{\ell}) \tau_{x-(\ell+1)e_i} \widetilde{A}_{\ell,i} \right\}.$$

Now, we apply Lemma 3.6 to estimate (3.40) (except $Q_{1,\ell}$ and $Q_{2,N}$) for each fixed iand j from the above; namely, take n = 1, $J(t, v) = \partial_{v_j}\lambda(t, v)$, $G(\eta) \equiv G_{ij}(\eta) = Q_{ij;\ell}(0, \eta)$, $M(\rho) = 0$ and apply Lemma 3.6 with $\ell + 1$ instead of ℓ by noting that $\langle G \rangle_{\Lambda(\ell+1),m} = 0$ for any $0 \leq m \leq (\ell + 1)^d_* \equiv |\Lambda(\ell + 1)|$. Then, noting $||J||_{\infty} \leq ||\partial\lambda||_{\infty}$, we see that (3.40) (except for $Q_{1,\ell}$ and $Q_{2,N}$) is bounded by the sum in i, j (note $G = G_{ij}$) of

(3.41)
$$\beta T \|\partial\lambda\|_{\infty}^{2} d(\ell+1)_{*}^{d} \sup_{m,\zeta \in \mathcal{X}_{\Lambda(\ell+1)^{c}}} \langle G, (-L_{\Lambda(\ell+1),\zeta})^{-1}G \rangle_{\Lambda(\ell+1),m} + \frac{C}{\beta} K + \beta^{2} Q_{N,\ell+1}^{(1)}(\partial\lambda,G).$$

Therefore, recalling the spectral gap of $L_{\Lambda(\ell),\zeta}$ given in (3.34), that is, $(-L_{\Lambda(\ell+1),\zeta})^{-1} \leq C_0^{-1}(\ell+1)^2$, the first term of (3.41) is bounded by

(3.42)
$$C\beta \|\partial\lambda\|_{\infty}^2 \ell^{d+2} \langle G^2 \rangle_{\Lambda(\ell+1),m}$$

However, by applying Taylor's formula to P_{ij} , we have

$$P_{ij}(\bar{\eta}_0^{\ell}) - P_{ij}(\bar{\eta}_{-e_i}^{\ell}) = D_{ij}(\bar{\eta}_0^{\ell})\varphi_{i,\ell}(\eta) + \frac{1}{2}D'_{ij}(\rho^*)\varphi_{i,\ell}(\eta)^2$$

with some $\rho^* \in [0, 1]$, where

(3.43)
$$\varphi_{i,\ell}(\eta) = \bar{\eta}_0^\ell - \bar{\eta}_{-e_i}^\ell$$
$$= \ell_*^{-d} \left\{ \tau_{\ell e_i} \widetilde{A}_{\ell,i} - \tau_{-(\ell+1)e_i} \widetilde{A}_{\ell,i} \right\}.$$

Therefore, after cancelling common terms $\ell_*^{-d} D_{ij}(\bar{\eta}_0^\ell) \tau_{\ell e_i} \widetilde{A}_{\ell,i}$, we have

(3.44)
$$G(\eta) = \ell_*^{-d} \left\{ D_{ij}(\bar{\eta}_0^\ell) - D_{ij}(\bar{\eta}_{-e_i}^\ell) \right\} \tau_{-(\ell+1)e_i} \widetilde{A}_{\ell,i} - \frac{1}{2} D'_{ij}(\rho^*) \varphi_{i,\ell}(\eta)^2 \\ = \ell_*^{-d} D'_{ij}(\rho^{**}) \varphi_{i,\ell}(\eta) \tau_{-(\ell+1)e_i} \widetilde{A}_{\ell,i} - \frac{1}{2} D'_{ij}(\rho^*) \varphi_{i,\ell}(\eta)^2$$

$$= \ell_*^{-d} D'_{ij}(\rho^{**}) \varphi_{i,\ell}(\eta) \big(\tau_{-(\ell+1)e_i} \widetilde{A}_{\ell,i} - \bar{A}_{\ell,m} \big) \\ + \ell_*^{-d} D'_{ij}(\rho^{**}) \varphi_{i,\ell}(\eta) \bar{A}_{\ell,m} - \frac{1}{2} D'_{ij}(\rho^{*}) \varphi_{i,\ell}(\eta)^2,$$

with some $\rho^{**} \in [0,1]$, where $\bar{A}_{\ell,m} := \langle \widetilde{A}_{\ell,i} \rangle_{\Lambda(\ell+1),m} \in [0, \ell_*^{d-1}]$. We have used Taylor's formula again. Since $D'_{ij}(\rho)$ is bounded due to (1.8) and

(3.45)
$$\langle \{\tau_{-(\ell+1)e_i} \widetilde{A}_{\ell,i} - \bar{A}_{\ell,m}\}^4 \rangle_{\Lambda(\ell+1),m} \le C_3 \ell^{2(d-1)},$$

(3.46)
$$\langle \varphi_{i,\ell}(\eta)^4 \rangle_{\Lambda(\ell+1),m} \le C_3 \ell^{-2(d+1)},$$

we obtain that (3.42) is bonded by

$$C\beta \|\partial\lambda\|_{\infty}^{2} \ell^{d+2} \{\ell^{-2d} \sqrt{\ell^{-2(d+1)}\ell^{2(d-1)}} + \ell^{-d-3} + \ell^{-2(d+1)} \} \le 3C\beta \|\partial\lambda\|_{\infty}^{2} \ell^{-1}.$$

Here, (3.45) (i.e. in CLT scaling) follows by using Lemma A.2 in Appendix of [14] taking $n = (\ell + 1)^d_*$ (indeed, the sum of the terms of the form $\langle (\eta_1 - m/n)^2 (\eta_2 - m/n)^2 \rangle (\leq 1)$ gives the leading order), (3.46) follows from (3.43) and (3.45), and the expectation under $\nu_{\Lambda(\ell+1),m}$ of the square of the middle term on the right-hand side of (3.44) is $O(\ell^{-d-3})$ from (3.46).

For $Q_{N,\ell+1}^{(1)} \equiv Q_{N,\ell+1}^{(1)}(\partial\lambda, G)$ in (3.41), since $||G||_{\infty} \leq C_4 \ell^{-2}$ from (3.44), $0 \leq \widetilde{A}_{\ell,i} \leq \ell_*^{d-1}$ and $|\varphi_{i,\ell}(\eta)| \leq \ell_*^{-1}$, we have by (3.33) in Lemma 3.6

$$|Q_{N,\ell+1}^{(1)}| \le CN^{-1}\ell^{(2d+4)}\ell^{-6} \|\partial\lambda\|_{\infty}^3 = CN^{-1}\ell^{(2d-2)} \|\partial\lambda\|_{\infty}^3$$

Summarizing these, we obtain the desired upper bound for $Q_{N,\ell}^{(3)}(\lambda)$.

3.4 Summary of estimate for $h_N(t)$ obtained in above three subsections

Lemma 3.1 (calculation for $\partial_t h_N(t)$), Theorem 3.3 (refined one-block estimate) and Theorem 3.5 (Boltzmann-Gibbs principle for gradient replacement, postponing the estimates for the CLT variances to Section 4) are summarized in the following lemma. The constant $C = C_T > 0$ may depend on T.

Lemma 3.8. Assume $\lambda \in C^{1,3}([0,T] \times \mathbb{T}^d)$, $F \in \mathcal{F}_0^d$ and the conditions (3.4) and (3.24). Then, setting $\rho(t,v) = \bar{\rho}(\lambda(t,v))$, we have

$$h_{N}(t) \leq h_{N}(0) + \int_{0}^{t} E^{f_{t}}[W] dt + Q_{N,\ell,\beta,K}^{\Omega_{1}}(\lambda,F) + C(\beta+1) \|\partial\lambda\|_{\infty}^{2} \sup_{\rho} \|R(\rho;F)\| + Q_{N}^{En}(\lambda,F) + Q_{N,\ell}^{\Omega_{2}}(\lambda,F)$$

for every $\beta > 0$ and $1 \leq \ell \leq N/2$, where ||R|| denotes the operator norm of matrix R (recall (3.29) for R), $Q_N^{En}(\lambda, F)$ is in Lemma 3.1 and

(3.47)
$$Q_{N,\ell,\beta,K}^{\Omega_1}(\lambda,F) := \frac{C}{\beta}K + \beta^2 Q_{N,\ell}^{(1)}(\lambda,F) + \beta Q_{\ell}^{(2)}(\lambda,F) + Q_{N,\ell}^{(3)}(\lambda),$$

see Theorem 3.5 for $Q_{N,\ell}^{(1)}$, $Q_{\ell}^{(2)}$ and Lemma 3.7 for $Q_{N,\ell}^{(3)}$. The function W is defined by

$$\begin{split} W(\eta) &= -N^{-d} \sum_{x \in \mathbb{T}_N^d} \dot{\lambda}(t, x/N) \{ \bar{\eta}_x^\ell - \rho(t, x/N) \} \\ &+ N^{-d} \sum_{x \in \mathbb{T}_N^d} \operatorname{Tr} \left(\partial^2 \lambda(t, x/N) \{ P(\bar{\eta}_x^\ell) - P(\rho(t, x/N)) \} \right) \\ &+ \frac{N^{-d}}{2} \sum_{x \in \mathbb{T}_N^d} \left(\partial \lambda(t, x/N), \{ \hat{c}(\bar{\eta}_x^\ell) - \hat{c}(\rho(t, x/N)) \} \partial \lambda(t, x/N) \right) \\ &+ N^{-d} K \sum_{x \in \mathbb{T}_N^d} \sigma_G(\bar{\eta}_x^\ell; t, x/N), \end{split}$$

where

(3.48)
$$\sigma_G(u;t,v) = \left(\frac{\langle c^+ \rangle_u}{\rho(t,v)} - \frac{\langle c^- \rangle_u}{1 - \rho(t,v)}\right) \{u - \rho(t,v)\}$$

The error $Q_{N,\ell}^{\Omega_2}(\lambda,F)$ is estimated as

$$(3.49) |Q_{N,\ell}^{\Omega_2}(\lambda,F)| \leq C \left(N^{-1} K^{1/2} \ell^{(d+2)/2} + \ell^{-d} (1+r(F)^d) \right) \\ \times \left(\|\dot{\lambda}\|_{\infty} + \|\partial\lambda\|_{\infty}^2 (1+\|F\|_{0,\infty})^2 + \|\partial^2\lambda\|_{\infty} (1+\|F\|_{1,\infty}) + K \right) \\ + C N^{-1} (\|\partial^3\lambda\|_{\infty} + \|\partial^2\lambda\|_{\infty} \|\partial\lambda\|_{\infty} + \|\partial\lambda\|_{\infty}^2 \|\partial\lambda\|_{\infty}).$$

In the above estimate, for each $\lambda_* > 0$, the constant $C = C_{\lambda_*}$ can be taken uniformly for $\lambda = \lambda(t, v)$ such that $\|\lambda\|_{\infty} \leq \lambda_*$.

When we apply this lemma later in Lemma 3.13, we take $\lambda(t, v) = \overline{\lambda}(\rho_K(t, v))$ with a solution ρ_K of (1.11). Under the condition (1.12) for the initial value ρ_0 , by Lemma 7.1 (the comparison theorem for ρ_K) and recalling (1.15), λ_* can be taken to be uniform in K.

Proof. The proof is similar to Lemma 3.2 of [14], but compared to it, we have the Glauber part and also require to derive an appropriate error estimate. For Ω_1 , we apply Theorem 3.5 (with an error $\frac{C}{\beta}K + \beta^2 Q_{N,\ell}^{(1)} + \beta Q_{\ell}^{(2)}$) and Lemma 3.7 (with an error $Q_{N,\ell}^{(3)}$), and obtain a similar replacement of $\int_0^t E^{f_t}[\Omega_1] dt$ by the following integral shown on p. 15 of [14]:

$$\begin{split} &\int_0^t \bigg\{ E^{f_t} \bigg[N^{-d} \sum_{x \in \mathbb{T}_N^d} \operatorname{Tr} \left(\partial^2 \lambda(t, x/N) P(\bar{\eta}_x^\ell) \right) \bigg] \\ &+ E^{f_t} \bigg[\frac{\beta N^{-d}}{2} \sum_{x \in \mathbb{T}_N^d} \left(\partial \lambda(t, x/N), R(\bar{\eta}_x^\ell; F) \partial \lambda(t, x/N) \right) \bigg] \bigg\} \, dt \end{split}$$

with an error replacing $\frac{Ct}{\beta} + o(1)$ by $Q_{N,\ell,\beta,K}^{\Omega_1}(\lambda, F)$ defined in (3.47). The second term is bounded by $C\beta \|\partial\lambda\|_{\infty}^2 \sup_{\rho} \|R(\rho; F)\|$ on the right-hand side of the desired estimate for $h_N(t)$. For Ω_2 , we apply (3.25) in Theorem 3.3 (refined one-block estimate) to replace $\int_0^t E^{f_t}[\Omega_2] dt$ by

(3.50)
$$\int_{0}^{t} E^{f_{t}} \left[N^{-d} \sum_{x \in \mathbb{T}_{N}^{d}} \left\{ -\dot{\lambda}(t, x/N) \bar{\eta}_{x}^{\ell} + \frac{1}{2} \left(\partial \lambda(t, x/N), \widehat{c}(\bar{\eta}_{x}^{\ell}; F) \partial \lambda(t, x/N) \right) + K \sigma_{G}(\bar{\eta}_{x}^{\ell}; t, x/N) \right\} \right] dt.$$

Note that the ensemble averages of the third and fourth terms of Ω_2 vanish: $\langle c_{xy}(\eta_x - \eta_y) \rangle_{\rho} = 0$, $\langle c_{xy}\pi_{xy}G \rangle_{\rho} = 0$ for all $G = G(\eta) \in \mathcal{F}_0$. The error for this replacement is estimated by

$$C\left(N^{-1}K^{1/2}\ell^{(d+2)/2} + \ell^{-d}(1+r(F)^{d})\right) \\ \times \left(\|\dot{\lambda}\|_{\infty} + \|\partial\lambda\|_{\infty}^{2}(1+\|F\|_{0,\infty})^{2} + \|\partial^{2}\lambda\|_{\infty}(1+\|F\|_{1,\infty}) + K\right)$$

by applying Theorem 3.3 taking $M = \|\dot{\lambda}\|_{\infty}$ for the first term of Ω_2 , $M\|h\|_{\infty} \leq C\|\partial\lambda\|_{\infty}^2(1+\|F\|_{0,\infty})^2$ for the second term, $M\|h\|_{\infty} \leq C\|\partial^2\lambda\|_{\infty}$ for the third term, $M\|h\|_{\infty} \leq C\|\partial^2\lambda\|_{\infty}\|F\|_{1,\infty}$ for the fourth term and $M\|h\|_{\infty} \leq CK$ for the fifth term noting that $\rho(t,v)$ is uniformly away from 0 and 1: $e^{-\lambda_*}/(e^{-\lambda_*}+1) \leq \rho(t,v) \leq e^{\lambda_*}/(e^{\lambda_*}+1)$ for λ such that $\|\lambda\|_{\infty} \leq \lambda_*$. This gives the first term in the error estimate of $|Q_{N,\ell}^{\Omega_2}(\lambda, F)|$.

Note that $\widehat{c}(\overline{\eta}_x^{\ell}; F)$ in (3.50) can be replaced by $\widehat{c}(\overline{\eta}_x^{\ell})$ with an error $C \|\partial\lambda\|_{\infty}^2 \sup_{\rho} \|R(\rho; F)\|$.

The reason that we have the second term in the error estimate of $|Q_{N,\ell}^{\Omega_2}(\lambda, F)|$ in (3.49) is as follows. Recalling that $\rho(t, v) = \bar{\rho}(\lambda(t, v))$, we have the identity

(3.51)
$$-\int_{\mathbb{T}^d} \operatorname{Tr}\left(\partial^2 \lambda(t,v) P(\rho(t,v))\right) dv = \frac{1}{2} \int_{\mathbb{T}^d} \left(\partial \lambda(t,v), \widehat{c}(\rho(t,v)) \partial \lambda(t,v)\right) dv,$$

This is shown by integration by parts and noting

(3.52)
$$\frac{\partial \bar{\rho}}{\partial \lambda} = \chi(\rho)$$

The terms with $\rho(t, x/N)$ in the second and third terms of W cancels by this identity with an error bounded by

$$CN^{-1}(\|\partial^{3}\lambda\|_{\infty} + \|\partial^{2}\lambda\|_{\infty}\|\partial\rho\|_{\infty} + \|\partial^{2}\lambda\|_{\infty}\|\partial\lambda\|_{\infty} + \|\partial\lambda\|_{\infty}^{2}\|\partial\rho\|_{\infty}),$$

which appears when we discretize the above two integrals in (3.51); note that P, \hat{c} are bounded and $P, \hat{c} \in C^{\infty}((0, 1))$. Note that $\|\partial \rho\|_{\infty}$ is uniformly bounded by $\|\partial \lambda\|_{\infty}$ for λ such that $\|\lambda\|_{\infty} \leq \lambda_*$.

3.5 Entropy inequality and large deviation error estimate

In Lemma 3.8, $E^{f_t}[W]$ is an expectation under $P^{f_t} = f_t d\nu^N$. We apply the entropy inequality to reduce it to the expectation under $P^{\psi_t} = \psi_t d\nu^N$:

(3.53)
$$E^{f_t}[W] \le \frac{K}{\delta N^d} \log E^{\psi_t}[e^{\delta N^d K^{-1}W}] + \frac{K}{\delta} h_N(t),$$

for every $\delta > 0$ (different from $\delta > 0$ in $\overline{K}_{\delta}(N)$). We study the asymptotic behavior of the first term on the right-hand side via a large deviation type upper bound with an appropriate error estimate under $\psi_t d\nu^N$; see Theorem 3.9 below.

For $\lambda(\cdot) \in C^1(\mathbb{T}^d)$ and $F = F(\eta) \in \mathcal{F}_0^d$, dropping the *t*-dependence in (2.1), the local equilibrium state $\psi_{\lambda(\cdot),F}^N(\eta)d\nu^N$ of second order approximation is a probability measure on \mathcal{X}_N defined by

(3.54)
$$\psi_{\lambda(\cdot),F}^{N}(\eta) = Z^{-1} \exp\bigg\{\sum_{x \in \mathbb{T}_{N}^{d}} \lambda(x/N)\eta_{x} + \frac{1}{N} \sum_{x \in \mathbb{T}_{N}^{d}} \left(\partial\lambda(x/N), \tau_{x}F(\eta)\right)\bigg\},$$

for $\eta \in \mathcal{X}_N$, where $Z = Z_{\lambda(\cdot),F,N}$ is the normalization constant with respect to ν^N . Then we have the following large deviation type upper bound for $\psi_{\lambda(\cdot),F}^N d\nu^N$. A similar result is shown in Lemmas 5 and 7 of [30] in the case of $F \equiv 0$ and Theorem 3.3 of [14]. But here, we give its error estimate.

The Bernoulli measure on \mathcal{X} associated with the chemical potential $\lambda \in \mathbb{R}$ is denoted by ν_{λ} , that is, $\nu_{\lambda} = \bar{\nu}_{\lambda}^{\otimes \mathbb{Z}^d}$, where $\bar{\nu}_{\lambda}$ is a probability measure on $\{0, 1\}$ defined by $\bar{\nu}_{\lambda}(\eta) := e^{\lambda \eta}/(e^{\lambda} + 1), \eta = 0, 1$. We abuse the notation ν_{λ} and ν_{ρ} which have different parametrizations but are clearly distinguishable. Indeed, $\nu_{\lambda} = \nu_{\bar{\rho}(\lambda)}$, i.e. ν_{ρ} with $\rho = \bar{\rho}(\lambda)$ defined by (1.14).

The rate function for the large deviation principle for the Bernoulli measure ν_{λ} is denoted by $I(u; \lambda)$, namely, for $u \in [0, 1]$,

(3.55)
$$I(u;\lambda) = -\lambda u - q(u) + p(\lambda),$$
$$p(\lambda) = \log(e^{\lambda} + 1), \quad q(u) = -\{u \log u + (1-u) \log(1-u)\},$$

see Section 3.8 for more details.

When we apply the following theorem later, we take K = K(N), but in this theorem, we consider K as a parameter in the function $G(v, \rho)$, such that $\frac{1}{K} \in (0, 1]$, and derive an estimate which is uniform in $\frac{1}{K}$.

Theorem 3.9. For every $G_1(v,\rho), G_2(v,\rho) \in C^1(\mathbb{T}^d \times [0,1])$, we have

$$N^{-d} \log E^{\psi_{\lambda(\cdot),F}^{N}}[\exp \widetilde{G}(\eta)]$$

$$\leq \sup_{\rho(v)\in C(\mathbb{T}^{d};[0,1])} \int_{\mathbb{T}^{d}} \{G(v,\rho(v)) - I(\rho(v);\lambda(v))\} dv + Q_{N,\ell}^{LD}(\lambda,F;G),$$

where $G(v, \rho) = G_1(v, \rho) + \frac{1}{K}G_2(v, \rho)$ and

(3.56)
$$\widetilde{G}(\eta) \equiv \widetilde{G}_{N,\ell}(\eta) = \sum_{x \in \mathbb{T}_N^d} G(x/N, \bar{\eta}_x^\ell).$$

The error term $Q_{N,\ell}^{LD}(\lambda, F; G)$ has an estimate which is uniform in $\frac{1}{K} \in (0, 1]$:

$$\begin{aligned} |Q_{N,\ell}^{LD}(\lambda,F;G)| &\leq CN^{-1} \|\partial\lambda\|_{\infty} \|F\|_{\infty} + CN^{-1}\ell \Big(\|\partial\lambda\|_{\infty} + \|\partial G_1\|_{\infty} + \|\partial G_2\|_{\infty} \Big) \\ &+ C\ell^{-d} \Big(\log\ell + \|\lambda\|_{\infty} + \|\partial_{\rho}G_1\|_{\infty} + \|\partial_{\rho}G_2\|_{\infty} \Big), \end{aligned}$$

where $\partial G_i = \partial_v G_i$ for i = 1, 2.

This theorem will be shown in Section 3.8. We will apply this theorem for the first term on the right-hand side of (3.53), which determines G_1 and G_2 by

(3.57)
$$\delta N^{d} K^{-1} W(\eta) = \sum_{x \in \mathbb{T}_{N}^{d}} \left\{ G_{1}(x/N, \bar{\eta}_{x}^{\ell}) + K^{-1} G_{2}(x/N, \bar{\eta}_{x}^{\ell}) \right\},$$

see Theorem 3.11 below. The definition of \widetilde{G} was slightly different from (3.56) in Theorem 3.3 of [14], but (3.56) is more convenient for our application.

The following corollary was used in Section 2; see (2.6). The contribution of F_N is negligible because of the property $||F_N||_{\infty} \leq CN^{1-\epsilon}$ for some $C, \epsilon > 0$.

Corollary 3.10. Choose $\lambda = \lambda_N$ and $F = F_N$ as in Section 5.2 below, namely, $\lambda = \lambda_{K(N)}$ with $K(N) \leq \overline{\delta} \log N$ for some $\overline{\delta} > 0$ and λ_K satisfying $\|\partial \lambda_K\|_{3,\infty} \leq CK^{\gamma}$ for some $\gamma > 0$, $F = \Phi_{n(N)}$ with Φ_n and $n(N) = N^{a_1}$ satisfying (5.10) and (5.11). Then, we have

$$\overline{\lim}_{N \to \infty} N^{-d} \log P^{\psi_{\lambda(\cdot),F}^{N}}(\mathcal{A}_{N,\phi,\varepsilon}) < 0, \quad \varepsilon > 0, \phi \in C^{\infty}(\mathbb{T}^{d}),$$

where $\mathcal{A}_{N,\phi,\varepsilon}$ is the subset of \mathcal{X}_N defined by (2.5) with $\rho^N(t)$ and $\rho_K(t,v)$ replaced by ρ^N defined by (1.4) with η_x instead of $\eta_x^N(t)$ and $\bar{\rho}(\lambda(v))$, respectively.

Proof. We follow the proof of Corollary B.1 of [14]. Consider

$$g(\eta) \equiv g_{\ell}(\eta) = N^{-d} \sum_{x \in \mathbb{T}_N^d} \phi(x/N) \{ \bar{\eta}_x^{\ell} - \bar{\rho}(\lambda(x/N)) \}.$$

Then, from $\phi \in C^{\infty}(\mathbb{T}^d)$ and the conditions for λ and $\ell = N^{a_2} = N^{1/(2d+5)}$, $\mathcal{A}_{N,\phi,\varepsilon} \subset \{|g(\eta)| > \varepsilon/2\}$ for large enough N. Furthermore, taking $G(v,\rho) = \phi(v)\{\rho - \bar{\rho}(\lambda(v))\} - \varepsilon/2$, by (3.56)

$$\widetilde{G}(\eta) \equiv \widetilde{G}_{N,\ell}(\eta) = \sum_{x \in \mathbb{T}_N^d} [\phi(x/N)\{\overline{\eta}_x^\ell - \overline{\rho}(\lambda(x/N))\} - \varepsilon/2] = N^d \{g(\eta) - \varepsilon/2\}.$$

Therefore, by Theorem 3.9, for any $\delta > 0$

$$N^{-d} \log P^{\psi_{\lambda(\cdot),F}^{N}}(g(\eta) > \varepsilon/2) \leq N^{-d} \log E^{\psi_{\lambda(\cdot),F}^{N}}[e^{\delta \widetilde{G}_{N,\ell}}]$$
$$\leq \sup_{\rho(v)\in C(\mathbb{T}^{d};[0,1])} \int_{\mathbb{T}^{d}} \{\delta G(v,\rho(v)) - I(\rho(v);\lambda(v))\} \, dv + Q_{N,\ell}^{LD}(\lambda,F;\delta G).$$

However, for the large deviation rate function $I(\rho; \lambda)$, we have

(3.58)
$$I(\rho;\lambda(v)) \ge C(\rho - \bar{\rho}(\lambda(v)))^2, \quad \rho \in [0,1],$$

for some C > 0, since $\frac{\partial^2 I}{\partial u^2}(u; \lambda) = \frac{1}{\chi(u)} > 0$ and $I(u; \lambda) = 0$ if and only if $u = \bar{\rho}(\lambda)$. From this, we see that there exists $\delta > 0$ small enough such that

$$\delta G(v,\rho) - I(\rho;\lambda(v)) \le -\varepsilon \delta/4$$

for all $\rho \in [0, 1]$.

For the error term, noting $\phi \in C^{\infty}(\mathbb{T}^d)$, we have

$$|Q_{N,\ell}^{LD}(\lambda,F;\delta G)| \le CN^{-1} \|\partial\lambda\|_{\infty} \|F\|_{\infty} + CN^{-1}\ell (1+\|\partial\lambda\|_{\infty}) + C\ell^{-d} (\log \ell + \|\lambda\|_{\infty}),$$

which tends to 0 as $N \to \infty$ under the choice of $\lambda = \lambda_N, F = F_N$ and $\ell = \ell(N)$ stated above. This concludes

$$\overline{\lim}_{N \to \infty} N^{-d} \log P^{\psi_{\lambda(\cdot),F}^N}(g(\eta) > \varepsilon/2) < 0.$$

A similar result can be shown for the event $\{g(\eta) < -\varepsilon/2\}$. This completes the proof of the corollary.

3.6 Further estimate for $h_N(t)$

The next theorem is immediate by combing Lemma 3.8, (3.53) and Theorem 3.9. A corresponding result was given for $h(t) = \overline{\lim}_{N\to\infty} h_N(t)$ in Theorem 2.1 of [14] (the error estimate of o(1) was sufficient in [14]). But, in the present setting, for the reason already noted, we need a bound for $h_N(t)$ with an appropriate error estimate.

To state the theorem, let us define g(t) as follows:

$$g(t) \equiv g_{\delta,K}(t) = \sup_{u(v) \in C(\mathbb{T}^d; [0,1])} \int_{\mathbb{T}^d} \left\{ \frac{\delta}{K} \cdot \sigma(u(v); t, v) - I(u(v); \lambda(t, v)) \right\} dv,$$

$$\sigma(u; t, v) \equiv \sigma_K(u; t, v) = -\dot{\lambda}(t, v) \{u - \rho(t, v)\} + \operatorname{Tr} \left(\partial^2 \lambda(t, v) \{P(u) - P(\rho(t, v))\}\right)$$

$$+ \frac{1}{2} \left(\partial \lambda(t, v), \{\widehat{c}(u) - \widehat{c}(\rho(t, v))\} \partial \lambda(t, v)\right) + K \sigma_G(u; t, v).$$

Recall $\rho(t, v) = \bar{\rho}(\lambda(t, v))$, Lemma 3.7 for $P(\rho)$ and (3.48) for $\sigma_G(u; t, v)$.

In view of (3.57) and W given in Lemma 3.8, we take

(3.59)
$$G_{1}(v,\rho) = \delta \cdot \sigma_{G}(\rho;t,v),$$
$$G_{2}(v,\rho) = \delta \Big[-\dot{\lambda}(t,v) \{\rho - \rho(t,v)\} + \operatorname{Tr} \left(\partial^{2}\lambda(t,v) \{P(\rho) - P(\rho(t,v))\}\right) \\ + \frac{1}{2} \left(\partial\lambda(t,v), \{\widehat{c}(\rho) - \widehat{c}(\rho(t,v))\}\partial\lambda(t,v)\right) \Big].$$

The corresponding error $\int_0^T Q_{N,\ell}^{LD}(\lambda(s), F; G) ds$ integrated in s will be denoted by $Q_{N,\ell,\delta}^{LD}(\lambda, F)$.

Theorem 3.11. Assume $\lambda \in C^{1,3}([0,T] \times \mathbb{T}^d)$, $F \in \mathcal{F}_0^d$ and the conditions (3.4) and (3.24). Then, there exist $\delta_0, C > 0$ such that for any $0 < \delta < \delta_0$, $\beta > 0$ and $1 \le \ell \le N/2$, we have

(3.60)
$$h_{N}(t) \leq h_{N}(0) + \frac{K}{\delta} \int_{0}^{t} g(s) \, ds + \frac{K}{\delta} \int_{0}^{t} h_{N}(s) \, ds \\ + Q_{N,\ell,\beta,K}^{\Omega_{1}}(\lambda,F) + Q_{N,\ell,\delta}^{LD}(\lambda,F) \\ + C(\beta+1) \, \|\partial\lambda\|_{\infty}^{2} \sup_{\rho \in [0,1]} \|R(\rho;F)\| + Q_{N}^{En}(\lambda,F) + Q_{N,\ell}^{\Omega_{2}}(\lambda,F).$$

The estimate is uniform in λ such that $\|\lambda\|_{\infty} \leq \lambda_*$ for each $\lambda_* > 0$ as we noted in Lemma 3.8.

See Section 5.1 for a summary of several error terms that appear in the estimate (3.60). In this theorem, it is not necessary that λ is determined from the solution $\rho_K(t, v)$ of (1.11) as $\lambda(t, v) = \overline{\lambda}(\rho_K(t, v))$. We assume this in Lemma 3.13 below.

Remark 3.2. Compared to Theorem 2.1 of [14], the term $K\sigma_G$, which results from the Glauber part, is new. Since $K\sigma_G$ and therefore σ contains a diverging factor K, we change $\delta \cdot \sigma$ to $\frac{\delta}{K} \cdot \sigma$ in the definition of g. From this (and when applying the entropy inequality), we have $\frac{K}{\delta}$ in front of $\int_0^t gds$ in the estimate in Theorem 3.11. The reason for having $\frac{K}{\delta}$ in front of $\int_0^t h_N ds$ in the estimate in Theorem 3.11 is that, when we apply the entropy inequality, we need to pay for K in $K\sigma_G$ by the entropy factor; recall (3.53).

3.7 Proof of Proposition 2.1

The following lemma was shown in Lemma 2.1 of [14].

Lemma 3.12.

$$\inf_{F \in \mathcal{F}_0^d} \sup_{\rho \in [0,1]} \| R(\rho; F) \| = 0.$$

In fact, [9] improved the result of this lemma giving the decay rate in n of $R(\rho; F)$ approximated by F satisfying $r(F) \leq n$; see (5.10) below.

The next lemma is shown in (2.6) in [14] when K = 0. We can easily extend it to the case where $K \ge 1$, since we put $\frac{\delta}{K}$ in front of σ in the definition of g(t).

Lemma 3.13. Let $\rho(t, v) \equiv \rho_K(t, v)$ be the solution of the hydrodynamic equation (1.11) with ρ_0 satisfying the condition (1.12), and define $\lambda(t, v)$ as $\lambda(t, v) = \overline{\lambda}(\rho(t, v))$. Then, we have

$$(3.61) g(t) \equiv g_{\delta_*,K}(t) \le 0$$

by making $\delta_* > 0$ small enough.

Proof. We first recall (3.58) to have

$$(3.62) I(u; \lambda(t, v)) \ge C\{u - \rho(t, v)\}^2$$

for some C > 0. Note that the large deviation rate function $I(u; \lambda)$ is determined only from ν_{λ} and especially it doesn't depend on K.

On the other hand, we have $\sigma(\rho(t, v); t, v) = 0$ and moreover, noting that $\frac{\partial}{\partial u}\sigma_G(\rho(t, v); t, v) = \frac{1}{\chi(\rho(t, v))}f(\rho(t, v)),$

$$(3.63) \qquad \frac{\partial \sigma}{\partial u}(\rho(t,v);t,v) = -\dot{\lambda}(t,v) + \operatorname{Tr}\left(\partial^2 \lambda(t,v) D(\rho(t,v))\right) + \frac{1}{2}\left(\partial \lambda(t,v), \hat{c}'(\rho(t,v)) \partial \lambda(t,v)\right) + \frac{K}{\chi(\rho(t,v))}f(\rho(t,v)) = 0,$$

where \hat{c}' denotes the matrix obtained by differentiating each element of \hat{c} in ρ . In fact, one can easily recognize that (3.63) is equivalent to the hydrodynamic equation (1.11) itself by noting (3.52). Therefore, by Taylor's formula

$$\sigma(u;t,v) = \frac{1}{2} \frac{\partial^2 \sigma}{\partial u^2} (u_0;t,v) \{u - \rho(t,v)\}^2$$

for some $u_0 \in [0, 1]$; note that (1.8) implies $\sigma \in C^{\infty}$ as a function of u. However, $\frac{\partial^2}{\partial u^2}(\sigma - K\sigma_G)(u; t, v)$ is bounded on $[0, 1] \times [0, T] \times \mathbb{T}^d$ and also independent of K, while

$$\left| K \frac{\partial^2 \sigma_G}{\partial u^2}(u;t,v) \right| = \left| K \frac{\partial^2}{\partial u^2} \left[\left(\frac{\langle c^+ \rangle_u}{\rho(t,v)} - \frac{\langle c^- \rangle_u}{1 - \rho(t,v)} \right) \{u - \rho(t,v)\} \right] \right| \le C_1 K_2$$

for some $C_1 > 0$, since $0 < c \le \rho(t, v) \le 1 - c < 1$ for some $c \in (0, 1/2)$ by Lemma 7.1 below; also recall the comment below Lemma 3.8. Thus we obtain

$$|\sigma(u;t,v)| \le (C_1 K + C_2) \{u - \rho(t,v)\}^2.$$

This combined with (3.62) completes the proof of (3.61) recalling $K \ge 1$.

Proof of Proposition 2.1. Proposition 2.1 follows from Theorem 3.11, Lemma 3.13 and a refined version of Lemma 3.12 given in (5.10) below. In fact, by (3.60) in Theorem 3.11 taking $F = F_N$ as in Proposition 2.1 (i.e., as in Section 5.2) (and $\lambda_* = \log(1-c)/c > 0$ with c in the proof of Lemma 3.13), and by Lemma 3.13 with $\delta_* > 0$ sufficiently small, we have

$$h_{N}(t) \leq h_{N}(0) + \frac{K}{\delta_{*}} \int_{0}^{t} h_{N}(s) \, ds + Q_{N,\ell,\beta,K}^{\Omega_{1}}(\lambda,F_{N}) + Q_{N,\ell,\delta_{*}}^{LD}(\lambda,F_{N}) + C(\beta+1) \|\partial\lambda\|_{\infty}^{2} \sup_{\rho \in [0,1]} \|R(\rho;F_{N})\| + Q_{N}^{En}(\lambda,F_{N}) + Q_{N,\ell}^{\Omega_{2}}(\lambda,F_{N}).$$

Therefore, with the help of Gronwall's inequality, we have

(3.64)
$$0 \le h_N(t) \le e^{Kt/\delta_*} \Big(h_N(0) + Q_{N,\ell,\beta,K}^{\Omega_1}(\lambda, F_N) + Q_{N,\ell,\delta_*}^{LD}(\lambda, F_N) + C(\beta + 1) \|\partial\lambda\|_{\infty}^2 \sup_{\rho \in [0,1]} \|R(\rho; F_N)\| + Q_N^{En}(\lambda, F_N) + Q_{N,\ell}^{\Omega_2}(\lambda, F_N) \Big).$$

Choosing $\beta = \beta(N), \ell = \ell(N), F_N, K = K(N)$ as in Section 5.2 below, we can show that each of the nine terms (recall that $Q_{N,\ell,\beta,K}^{\Omega_1}(\lambda, F_N)$ consists of four terms as in (3.47)) multiplied by e^{KT/δ_*} on the right-hand side of (3.64) is bounded by $\bar{C}N^{-\kappa}$ for some $\bar{C} = \bar{C}_T > 0$ and $\kappa > 0$. Therefore, we conclude $h_N(t) \leq CN^{-\kappa}, t \in [0,T]$ for $C = 9\bar{C}$.

3.8 Proof of Theorem 3.9

First we recall some known facts on thermodynamic functions; see [15] and Appendix B of [14] with $J \equiv 0$, $H_{\Lambda} \equiv 0$ and $R_1 = 0$. For a finite domain $\Lambda \Subset \mathbb{Z}^d$ ($\Lambda \neq \emptyset$), a chemical potential $\lambda \in \mathbb{R}$ and particle number $k : 0 \le k \le |\Lambda|$, define grand canonical and canonical partition functions by

(3.65)
$$Z_{\Lambda,\lambda} = \sum_{\eta_{\Lambda} \in \mathcal{X}_{\Lambda}} \exp\left\{\lambda \sum_{x \in \Lambda} \eta_x\right\} = e^{p(\lambda)|\Lambda|},$$

(3.66)
$$Z_{\Lambda,k} = \sum_{\eta_{\Lambda} \in \mathcal{X}_{\Lambda}: \sum_{x \in \Lambda} \eta_{x} = k} 1 = \begin{pmatrix} |\Lambda| \\ k \end{pmatrix},$$

respectively, where $\mathcal{X}_{\Lambda} = \{0, 1\}^{\Lambda}$ and recall (3.55) for $p(\lambda)$. Then,

(3.67)
$$\log Z_{\Lambda,k} = |\Lambda|q(k/|\Lambda|) + \frac{1}{2}\log\frac{|\Lambda|}{2\pi k(|\Lambda|-k)} + o(1),$$

as $|\Lambda|, k \to \infty$; recall (3.55) for q(u). (Note, for k = 0 or $|\Lambda|, Z_{\Lambda,0} = Z_{\Lambda,|\Lambda|} = 1$ and q(0) = q(1) = 0.) Indeed, by Stirling's formula $n! \sim n^n e^{-n} \sqrt{2\pi n}$ $(a_n \sim b_n \text{ means } a_n/b_n \to 1 \text{ as } n \to \infty)$, we have

$$\begin{split} \log Z_{\Lambda,k} &= \log \left(|\Lambda|!/k! (|\Lambda| - k)! \right) \\ &= |\Lambda| \log |\Lambda| - |\Lambda| + \frac{1}{2} \log 2\pi |\Lambda| - \left\{ k \log k - k + \frac{1}{2} \log 2\pi k \right\} \\ &- \left\{ (|\Lambda| - k) \log(|\Lambda| - k) - (|\Lambda| - k) + \frac{1}{2} \log 2\pi (|\Lambda| - k) \right\} + o(1) \\ &= |\Lambda| \log |\Lambda| - k \log k - (|\Lambda| - k) \log(|\Lambda| - k) + \frac{1}{2} \log \frac{|\Lambda|}{2\pi k(|\Lambda| - k)} + o(1) \\ &= |\Lambda| \left\{ -\frac{k}{|\Lambda|} \log \frac{k}{|\Lambda|} - \left(1 - \frac{k}{|\Lambda|}\right) \log(1 - \frac{k}{|\Lambda|}) \right\} + \frac{1}{2} \log \frac{|\Lambda|}{2\pi k(|\Lambda| - k)} + o(1), \end{split}$$

as $|\Lambda|, k \to \infty$, and this shows (3.67).

The functions $p(\lambda)$ and $-q(\rho)$ are continuous and convex, and satisfy

(3.68)
$$p(\lambda) = \sup_{\rho \in [0,1]} \{q(\rho) + \lambda\rho\},$$

(3.69)
$$q(\rho) = \inf_{\lambda \in \mathbb{R}} \{ p(\lambda) - \lambda \rho \}.$$

For each $\rho \in [0, 1]$, there exists a unique $\lambda = \overline{\lambda}(\rho) \in \mathbb{R} := \mathbb{R} \cup \{\pm \infty\}$ such that

(3.70)
$$p(\lambda) = q(\rho) + \lambda \rho.$$

Before proceeding to the proof of Theorem 3.9, we prepare a similar statement for finite volume Bernoulli measures with constant chemical potentials. For $\Lambda \Subset \mathbb{Z}^d$ $(\Lambda \neq \emptyset)$, $\lambda \in \mathbb{R}$, the finite volume (grand canonical) Bernoulli measure $\nu_{\Lambda,\lambda}$ on Λ , which is a probability measure on \mathcal{X}_{Λ} , is defined by

(3.71)
$$\nu_{\Lambda,\lambda}(\eta) = Z_{\Lambda,\lambda}^{-1} \exp\left\{\lambda \sum_{x \in \Lambda} \eta_x\right\} \equiv \bar{\nu}_{\lambda}^{\otimes \Lambda}(\eta), \quad \eta \in \mathcal{X}_{\Lambda}.$$

Proposition 3.14. For every $\lambda \in \mathbb{R}$ and $G(\rho) = G_1 + \frac{1}{K}G_2 \in C^1([0,1])$,

$$\ell_*^{-d} \log E^{\nu_{\Lambda(\ell),\lambda}} [\exp\{\ell_*^d G(\bar{\eta}_0^\ell)\}] = \sup_{\rho \in [0,1]} \{G(\rho) - I(\rho;\lambda)\} + Q_\ell(\lambda,G),$$

recall $\Lambda(\ell) = \Lambda_{\ell,0}$ in (3.1) and (3.22) for $\bar{\eta}_0^{\ell}$. We have an estimate for the error term $Q_{\ell}(\lambda, G)$, which is uniform in $\frac{1}{K} \in (0, 1]$:

(3.72)
$$|Q_{\ell}(\lambda, G)| \le C\ell^{-d} (\log \ell + |\lambda| + ||G'||_{\infty}),$$

where $G' \equiv \partial_{\rho} G$.

Proof. By (3.65), setting $\bar{\eta}_{\Lambda} = \frac{1}{|\Lambda|} \sum_{x \in \Lambda} \eta_x$ for $\Lambda \subset \mathbb{Z}^d$ in general, we have

$$I := \frac{1}{|\Lambda|} \log E^{\nu_{\Lambda,\lambda}} [\exp\{|\Lambda|G(\bar{\eta}_{\Lambda})\}]$$

= $\frac{1}{|\Lambda|} \log \left[\sum_{\eta \in \mathcal{X}_{\Lambda}} \exp\{\lambda \sum_{x \in \Lambda} \eta_x + |\Lambda|G(\bar{\eta}_{\Lambda})\}\right] - \frac{1}{|\Lambda|} \log Z_{\Lambda,\lambda}$
= $\frac{1}{|\Lambda|} \log \left[\sum_{m=0}^{|\Lambda|} Z_{\Lambda,m} \exp\{\lambda m + |\Lambda|G(m/|\Lambda|)\}\right] - p(\lambda).$

Then we apply (3.67) for $Z_{\Lambda,m}$. First note that

(3.73)
$$\frac{4}{|\Lambda|} \le \frac{|\Lambda|}{m(|\Lambda| - m)} \le 2$$

for $1 \leq m \leq |\Lambda| - 1$, $|\Lambda| \geq 2$. Therefore, by (3.67), $Z_{\Lambda,m} \leq e^{|\Lambda|q(m/|\Lambda|) + \frac{1}{2}\log\frac{2}{2\pi} + C_1}$ for $0 \leq m \leq |\Lambda|, |\Lambda| \geq 2$ (including $m = 0, |\Lambda|$). Thus, denoting $a^* := \sup_{\rho \in [0,1]} \{G(\rho) - I(\rho; \lambda)\} + p(\lambda)$ and estimating all terms in the sum in m by using a^* , we have an upper bound of I:

$$I \leq \frac{1}{|\Lambda|} \log \left[(|\Lambda|+1) \exp\{|\Lambda|a^* + C_2\} \right] - p(\lambda)$$
$$= \frac{1}{|\Lambda|} \left[\log(|\Lambda|+1) + |\Lambda|a^* + C_2 \right] - p(\lambda)$$
$$\leq a^* - p(\lambda) + \frac{C}{|\Lambda|} \log |\Lambda|.$$

(Note that the constant C is uniform in G; indeed, C = 2 is enough for large $|\Lambda|$.)

Next, we show the lower bound for I. The supremum for a^* is attained at some $\rho_* \in [0,1]$. (Note that q is concave and q(0) = q(1) = 0, and $q'(\rho) = \log(1-\rho)/\rho$ so that $q'(0) = +\infty, q'(1) = -\infty$.) Since $G \in C^1([0,1])$, choosing m such that $\frac{m}{|\Lambda|}$ is close to ρ_* in the sense that $|\frac{m}{|\Lambda|} - \rho_*| \leq \frac{1}{|\Lambda|}$ and $\frac{m}{|\Lambda|} \leq \rho_*$ (if $\rho_* \leq \frac{1}{2}$) or $\frac{m}{|\Lambda|} \geq \rho_*$ (if $\rho_* > \frac{1}{2}$), we have

$$\left| \{ G(\rho_*) - I(\rho_*; \lambda) \} - \{ G(m/|\Lambda|) - I(m/|\Lambda|; \lambda) \} \right| \le \frac{\|G'\|_{\infty} + |\lambda|}{|\Lambda|} + |q(\rho_*) - q(m/|\Lambda|)|.$$

Here, $|q(\rho_*) - q(m/|\Lambda|)|$ is estimated as

$$|q(\rho_*) - q(m/|\Lambda|)| \le \frac{C_3}{|\Lambda|}$$

for some $C_3 = C_3(\varepsilon) > 0$ if $0 < \varepsilon \le \rho_* \le 1 - \varepsilon < 1$. If $0 < \rho_* \le \varepsilon$, since $\frac{m}{|\Lambda|} \le \rho_*$, we have $|q'(\frac{m}{|\Lambda|})| \ge |q'(\rho)|$ for every $\rho \in [m/|\Lambda|, \rho_*]$ and therefore

$$|q(\rho_*) - q(m/|\Lambda|)| \le \frac{1}{|\Lambda|} \left| q'\left(\frac{m}{|\Lambda|}\right) \right| = \frac{1}{|\Lambda|} \log \frac{|\Lambda| - m}{m} \le \frac{1}{|\Lambda|} \log |\Lambda|.$$

The case $1 > \rho_* \ge 1 - \varepsilon$ is similar.

Then, taking only the term given by this m in the sum of I and using the lower bound in (3.73) for $Z_{\Lambda,m}$, we have

$$\begin{split} I &\geq \frac{1}{|\Lambda|} \log \left[\exp\left\{ |\Lambda| \left(a^* - \frac{\|G'\|_{\infty} + |\lambda| + C_3 + \log|\Lambda|}{|\Lambda|} \right) + \frac{1}{2} \log \frac{4}{2\pi|\Lambda|} \right\} \right] - p(\lambda) \\ &= a^* - \frac{\|G'\|_{\infty} + |\lambda| + C_3 + \log|\Lambda|}{|\Lambda|} - \frac{\log|\Lambda| - \log\frac{2}{\pi}}{2|\Lambda|} - p(\lambda) \\ &\geq a^* - p(\lambda) - \frac{C}{|\Lambda|} (\log|\Lambda| + |\lambda| + \|G'\|_{\infty}). \end{split}$$

We therefore have the conclusion with the estimate (3.72) by taking $\Lambda = \Lambda(\ell)$.

We are now ready to give the proof of Theorem 3.9. As we noted, this theorem is an extension of Theorem 3.3 of [14] and gives a further error estimate. Because the definition of \tilde{G} is different, the proof is modified by applying Hölder's inequality.

Proof of Theorem 3.9. To apply Proposition 3.14 by localizing in space, finally to obtain the estimate in terms of a spatial integral and also to match with the scale ℓ in \widetilde{G} , we divide \mathbb{T}_N^d into disjoint boxes $\{\Lambda_{\ell,a}\}_a$ with side length $\ell_* = 2\ell + 1$ and centered at $a \in$ $\ell_*\mathbb{T}_{N/\ell_*}^d = \{\ell_*, 2\ell_*, \ldots, N\}^d$, assuming $N/\ell_* \in \mathbb{N}$ for simplicity (if not, we may make the side lengths of some of the boxes smaller or larger by 1).

Since $x \in \mathbb{T}_N^d$ is uniquely decomposed as x = y + a with $y \in \Lambda(\ell) = \Lambda_{\ell,0}$ and $a \in \ell_* \mathbb{T}_{N/\ell_*}^d$ (we write a(x) for a) according to this division, writing the sum and the product in $a \in \ell_* \mathbb{T}_{N/\ell_*}^d$ by \sum_a^* and \prod_a^* , respectively, we have

$$\begin{split} I &:= E^{\nu_{\lambda(\cdot),F}^{N}} [\exp \widetilde{G}(\eta)] \\ &= Z_{\lambda(\cdot),F,N}^{-1} E^{\nu^{N}} \bigg[\exp \bigg\{ \sum_{y \in \Lambda(\ell)} \sum_{a}^{*} \big[G(y+a)/N, \bar{\eta}_{y+a}^{\ell} \big] + \lambda(a/N) \bar{\eta}_{y+a}^{\ell} \big] + R(\eta) \bigg\} \bigg] \\ &= Z_{\lambda(\cdot),F,N}^{-1} E^{\nu^{N}} \bigg[\prod_{y \in \Lambda(\ell)} \exp \bigg\{ \sum_{a}^{*} \big[G((y+a)/N, \bar{\eta}_{y+a}^{\ell}) + \lambda(a/N) \bar{\eta}_{y+a}^{\ell} \big] \bigg\} \cdot e^{R(\eta)} \bigg] \\ &\leq e^{\|R\|_{\infty}} Z_{\lambda(\cdot),F,N}^{-1} \prod_{y \in \Lambda(\ell)} E^{\nu^{N}} \bigg[\exp \bigg\{ \ell_{*}^{d} \sum_{a}^{*} \big[G((y+a)/N, \bar{\eta}_{y+a}^{\ell}) + \lambda(a/N) \bar{\eta}_{y+a}^{\ell} \big] \bigg\} \bigg]^{1/\ell_{*}^{d}} , \end{split}$$

where $R(\eta) = R_1 + R_2$ and

$$R_1 = \frac{1}{N} \sum_{x \in \mathbb{T}_N^d} \left(\partial \lambda(x/N), \tau_x F(\eta) \right),$$

$$R_2 = \sum_{x \in \mathbb{T}_N^d} \left(\lambda(x/N) \eta_x - \lambda(a(x)/N) \bar{\eta}_x^\ell \right).$$

For the last line, we have first estimated $e^R \leq e^{||R||_{\infty}}$ and then applied Hölder's inequality under ν^N . Two error terms R_1 and R_2 are estimated as follows:

(3.74)
$$|R_1(\eta)| \le N^{d-1} ||\partial\lambda||_{\infty} ||F||_{\infty}, \quad |R_2(\eta)| \le \ell_* N^{d-1} ||\partial\lambda||_{\infty}.$$

In fact, for R_2 , note that the sum of the second term of R_2 can be rewritten as

$$\sum_{x \in \mathbb{T}_N^d} \lambda(a(x)/N) \ell_*^{-d} \sum_{z \in \Lambda(\ell)} \eta_{x+z} = \sum_{x' \in \mathbb{T}_N^d} \eta_{x'} \ell_*^{-d} \sum_{z \in \Lambda(\ell)} \lambda(a(x'-z)/N).$$

Since $\bar{\eta}_{y+a}^{\ell}$ are functions of $\eta|_{\Lambda_{\ell,y+a}}$, they are independent for different *a* under ν^N . Therefore, the last expectation in the bound for *I* is factorized and is rewritten as

$$E^{\nu^{N}} \left[\exp\left\{ \ell_{*}^{d} \sum_{a}^{*} \left[G((y+a)/N, \bar{\eta}_{y+a}^{\ell}) + \lambda(a/N) \bar{\eta}_{y+a}^{\ell} \right] \right\} \right]$$

= $\prod_{a}^{*} E^{\nu^{N}} \left[\exp\left\{ \ell_{*}^{d} \left[G((y+a)/N, \bar{\eta}_{y+a}^{\ell}) + \lambda(a/N) \bar{\eta}_{y+a}^{\ell} \right] \right\} \right]$
= $\prod_{a}^{*} \left(Z_{\Lambda_{\ell,y+a},\lambda(a/N)} \cdot 2^{-\ell_{*}^{d}} \right) E^{\nu_{\Lambda_{\ell,y+a},\lambda(a/N)}} \left[\exp\left\{ \ell_{*}^{d} G((y+a)/N, \bar{\eta}_{y+a}^{\ell}) \right\} \right].$

In the second line, ν^N can be replaced by $\bar{\nu}_{1/2}^{\otimes \Lambda_{\ell,y+a}}$ and, recalling (3.71), we obtain the third line. Thus, noting the shift-invariance of $Z_{\Lambda,\lambda}$ and $\nu_{\Lambda,\lambda}$ in Λ , and $\prod_y \prod_a^* = \prod_x$, we have

$$I \le e^{\|R\|_{\infty}} Z_{\lambda(\cdot),F,N}^{-1} \prod_{a}^{*} Z_{\Lambda(\ell),\lambda(a/N)} \cdot 2^{-\ell_{\ast}^{d}} \times \prod_{x \in \mathbb{T}_{N}^{d}} E^{\nu_{\Lambda(\ell),\lambda(a(x)/N)}} \bigg[\exp\big\{\ell_{\ast}^{d} G(x/N, \bar{\eta}_{0}^{\ell})\big\} \bigg]^{1/\ell_{\ast}^{d}},$$

and therefore,

$$(3.75) N^{-d} \log I \le N^{-d} ||R||_{\infty} + N^{-d} \log \left(Z_{\lambda(\cdot),F,N}^{-1} \prod_{a}^{*} Z_{\Lambda(\ell),\lambda(a/N)} \cdot 2^{-\ell_{*}^{d}} \right) + N^{-d} \sum_{x \in \mathbb{T}_{N}^{d}} \ell_{*}^{-d} \log E^{\nu_{\Lambda(\ell),\lambda(a(x)/N)}} \left[\exp \left\{ \ell_{*}^{d} G(x/N, \bar{\eta}_{0}^{\ell}) \right\} \right].$$

For the partition functions, recalling that

$$Z_{\Lambda,\lambda} \cdot 2^{-|\Lambda|} = E^{\bar{\nu}_{1/2}^{\otimes \Lambda}} \Big[\exp\left\{\lambda \sum_{x \in \Lambda} \eta_x\right\} \Big], \quad \lambda \in \mathbb{R},$$
$$Z_{\lambda(\cdot),F,N} = E^{\nu^N} \Big[\exp\left\{\sum_{x \in \mathbb{T}_N^d} \lambda(x/N)\eta_x + R_1(\eta)\right\} \Big],$$

and changing $\Lambda(\ell)$ to $\Lambda_{\ell,a}$ in $Z_{\Lambda(\ell),\lambda(a/N)}$, we obtain

$$\prod_{a}^{*} Z_{\Lambda_{\ell,a},\lambda(a/N)} \cdot 2^{-\ell_{*}^{d}} = Z_{\widetilde{\lambda}(\cdot),0,N},$$

where $\widetilde{\lambda}(v) = \sum_{a}^{*} \lambda(a/N) \mathbf{1}_{\Lambda_{\ell,a}}(Nv)$ is a step function on \mathbb{T}^{d} and F = 0 on the right-hand side. To replace the step function $\widetilde{\lambda}(\cdot)$ with the original function $\lambda(\cdot)$, note that

$$\frac{Z_{\widetilde{\lambda}(\cdot),0,N}}{Z_{\lambda(\cdot),0,N}} = \frac{E^{\nu_N} [\exp\{\sum_{x \in \mathbb{T}_N^d} \widetilde{\lambda}(x/N)\eta_x\}]}{E^{\nu_N} [\exp\{\sum_{x \in \mathbb{T}_N^d} \lambda(x/N)\eta_x\}]}$$

$$=\frac{E^{\nu_N}[\exp\{\sum_{x\in\mathbb{T}_N^d}\lambda(x/N)\eta_x+\sum_{x\in\mathbb{T}_N^d}(\widetilde{\lambda}(x/N)-\lambda(x/N))\eta_x\}]}{E^{\nu_N}[\exp\{\sum_{x\in\mathbb{T}_N^d}\lambda(x/N)\eta_x\}]}$$
$$\leq e^{\ell N^{d-1}\|\partial\lambda\|_{\infty}}.$$

To compare two partition functions with and without F, we have

$$\frac{Z_{\lambda(\cdot),0,N}}{Z_{\lambda(\cdot),F,N}} \le e^{\|R_1\|_{\infty}}.$$

Summarizing these and noting (3.74), we have shown for first two terms in (3.75) that

(3.76)
$$N^{-d} \|R\|_{\infty} + N^{-d} \log \left(Z_{\lambda(\cdot),F,N}^{-1} \prod_{a}^{*} Z_{\Lambda(\ell),\lambda(a/N)} \cdot 2^{-\ell_{*}^{d}} \right)$$
$$\leq C N^{-1} \|\partial\lambda\|_{\infty} (\ell + \|F\|_{\infty}).$$

For the last term of (3.75), one can apply Proposition 3.14 to see that

$$\begin{split} N^{-d} & \sum_{x \in \mathbb{T}_N^d} \ell_*^{-d} \log E^{\nu_{\Lambda(\ell),\lambda(a(x)/N)}} \Big[\exp \left\{ \ell_*^d G(x/N, \bar{\eta}_0^\ell) \right\} \Big] \\ &= N^{-d} \sum_{x \in \mathbb{T}_N^d} \left\{ \sup_{\rho \in [0,1]} \{ G(x/N, \rho) - I(\rho; \lambda(a(x)/N)) \} + Q_\ell(\lambda(a(x)/N), G(x/N, \cdot)) \right\}, \end{split}$$

and

(3.77)
$$|Q_{\ell}(\lambda(a(x)/N), G(x/N, \cdot))| \leq \frac{C}{\ell^d} (\log \ell + \|\lambda\|_{\infty} + \|\partial_{\rho}G_1\|_{\infty} + \|\partial_{\rho}G_2\|_{\infty}).$$

Moreover, one can estimate

$$(3.78) N^{-d} \sum_{x \in \mathbb{T}_N^d} \sup_{\rho \in [0,1]} \{ G(x/N,\rho) - I(\rho; \lambda(a(x)/N)) \} \le \sup_{\rho: \text{step on } \mathbb{T}^d} \mathbb{G}(\rho) + Q_{\ell,N},$$

where the supremum on the right-hand side is taken over all step functions $\rho(v)$ on \mathbb{T}^d ,

(3.79)
$$Q_{\ell,N} \equiv Q_{\ell,N}(\lambda;G) = \frac{2\ell}{N} \left(\|\partial\lambda\|_{\infty} + \|\partial_v G_1\|_{\infty} + \|\partial_v G_2\|_{\infty} \right)$$

and

$$\mathbb{G}(\rho) = \int_{\mathbb{T}^d} \{ G(v, \rho(v)) - I(\rho(v); \lambda(v)) \} \, dv,$$

for $\rho \in L^1(\mathbb{T}^d; [0, 1])$. Indeed, we first replace $G(x/N, \rho)$ in (3.78) by $G(a(x)/N, \rho)$ with an error within $\ell N^{-1}(\|\partial_v G_1\|_{\infty} + \|\partial_v G_2\|_{\infty})$, and then the supremum is attained at some $\rho(a(x)/N)$ for each a = a(x). This defines a step function $\rho(v) = \sum_a^* \rho(a/N) \mathbf{1}_{\Lambda_{\ell,a}}(Nv)$, and leads to the functional $\mathbb{G}(\rho)$, but with an discretization also for G and λ in the variable v. The error of removing these discretization in G and λ is estimated by using

$$\sup_{|v_1-v_2| \le \ell/N} |\lambda(v_1) - \lambda(v_2)| \le \ell N^{-1} \|\partial\lambda\|_{\infty},$$

$$\sup_{\substack{|v_1 - v_2| \le \ell/N \ \rho \in [0,1]}} \sup_{\rho \in [0,1]} |G(v_1,\rho) - G(v_2,\rho)| \le \ell N^{-1} \left(\|\partial_v G_1\|_{\infty} + \|\partial_v G_2\|_{\infty} \right),$$
$$\left| \int_{\mathbb{T}^d} p(\lambda(v)) \, dv - N^{-d} \sum_{x \in \mathbb{T}^d_N} p(\lambda(a(x)/N)) \right| \le \ell N^{-1} \|p'\|_{\infty} \|\partial\lambda\|_{\infty} \le \ell N^{-1} \|\partial\lambda\|_{\infty}.$$

and obtain (3.78) with $Q_{\ell,N}$ given by (3.79).

Since $\mathbb{G}(\rho)$ is continuous in $\rho \in L^1(\mathbb{T}^d; [0, 1])$ and $C(\mathbb{T}^d; [0, 1])$ is dense in $L^1(\mathbb{T}^d; [0, 1])$, we obtain the conclusion of the theorem from (3.76), (3.77), (3.78) and (3.79).

4 Error estimates for the CLT variances

We now give the error estimates for the variational quantity appearing in Lemma 3.6, which is sometimes called the central limit theorem (CLT) variance; recall also the error term $Q_{\ell}^{(2)}(\lambda, F)$ in Theorem 3.5. Recall that this quantity is defined by means of the localized Kawasaki generator $L_{\Lambda(\ell),\zeta}$ given in (3.32) and defined on the box $\Lambda(\ell) = \Lambda_{\ell,0}$ centered at 0, where we denoted $\zeta = \eta|_{\Lambda(\ell)^c}$ and $\xi = \eta|_{\Lambda(\ell)}$ for $\eta \in \mathcal{X}$. In particular, the Glauber part plays no role and the arguments in Sections 4 (characterization of closed forms) and 5 (computation of variances) of [14] are valid as is in our setting.

However, these are not sufficient for our purpose to derive appropriate decay rates for the error terms. In particular, for the CLT variance of A_{ℓ} , a qualitative convergence result as $\ell \to \infty$ (see (4.8) below) is known and shown based on the characterization of the closed forms, initiated by Varadhan [28] for the Ginzburg-Landau type model. But, this is weak for our purpose and, to fill the gap, Theorem 4.2 below was shown in [9] by a new technique inspired by the method of the quantitative homogenization. This theorem provides an appropriate decay rate of the CLT variance of A_{ℓ} and, in particular, the characterization of the closed forms was not used for the proof of this theorem.

For $f, g: \mathcal{X}_{\Lambda(\ell),m} \to \mathbb{R}$ which satisfy $\langle f \rangle_{\Lambda(\ell),m} = \langle g \rangle_{\Lambda(\ell),m} = 0$, we define

$$\Delta_{\ell,m,\zeta}(f,g) = \langle f(-L_{\Lambda(\ell),\zeta})^{-1}g \rangle_{\Lambda(\ell),m},$$

$$\Delta_{\ell,m,\zeta}(f) = \Delta_{\ell,m,\zeta}(f,f),$$

where $m \in [0, \ell_*^d] \cap \mathbb{Z}$ and $\zeta \in \mathcal{X}_{\Lambda(\ell)^c}$. Recall three \mathbb{R}^d -valued functions A_ℓ, B_ℓ and H_ℓ , which are defined by (3.35), (3.36), and (3.37), respectively, and n = r(F) + 1 in H_ℓ .

Proposition 4.1. For $F \in \mathcal{F}_0^d$ and $\theta, \tilde{\theta} \in \mathbb{R}^d : |\theta| = |\tilde{\theta}| = 1$, set

$$Q_{\ell}^{(4)}(F) \equiv Q_{\ell}^{(4)}(F;\theta,m,\zeta) = \ell_{*}^{-d} \Delta_{\ell,m,\zeta}(\theta \cdot (B_{\ell,\zeta} - H_{\ell,\zeta,F})) - \frac{1}{2} \theta \cdot \hat{c}(m/\ell_{*}^{d};F)\theta,$$

$$Q_{\ell}^{(5)}(F) \equiv Q_{\ell}^{(5)}(F;\theta,\tilde{\theta},m,\zeta) = \ell_{*}^{-d} \Delta_{\ell,m,\zeta}(\tilde{\theta} \cdot A_{\ell},\theta \cdot (B_{\ell,\zeta} - H_{\ell,\zeta,F})) - (\theta \cdot \tilde{\theta})\chi(m/\ell_{*}^{d}).$$

Then, we have

(4.1)
$$|Q_{\ell}^{(4)}(F)| \le Cr(F)^{d}(1 + ||F||_{0,\infty}^{2})\ell^{-1},$$

(4.2) $|Q_{\ell}^{(5)}(F)| \le Cr(F)^{2d} ||F||_{\infty} \ell^{-d} + C\ell^{-1},$

uniformly in $\theta, \tilde{\theta} \in \mathbb{R}^d$: $|\theta| = |\tilde{\theta}| = 1$, $m \in [0, \ell^d_*] \cap \mathbb{Z}$ and $\zeta \in \mathcal{X}_{\Lambda(\ell)^c}$, where $\theta \cdot \tilde{\theta}$ stands for the inner product of θ and $\tilde{\theta} \in \mathbb{R}^d$ which has been denoted by $(\theta, \tilde{\theta})$ in the previous sections.

Proof. We use the following two identities shown in the proof of Proposition 5.1 of [14]:

(4.3)
$$\ell_*^{-d} \Delta_{\ell,m,\zeta} \Big(\theta \cdot (B_\ell - H_\ell) \Big) \\= \frac{1}{2} \ell_*^{-d} \sum_{b = \tau_x e_i^* \in (\Lambda(\ell))^*} \left\langle c_b \Big[\theta \cdot e_i (\xi_{x+e_i} - \xi_x) - \pi_b \Big(\sum_{y \in \Lambda(\ell-n)} \tau_y F \cdot \theta \Big) \Big]^2 \right\rangle_{\Lambda(\ell),m},$$

and

$$(4.4) \quad \ell_*^{-d} \Delta_{\ell,m,\zeta} \big(\tilde{\theta} \cdot A_\ell, \theta \cdot (B_\ell - H_\ell) \big) \\ = \ell_*^{-d} \Big\langle \Big\{ \sum_{i=1}^d \tilde{\theta}_i \sum_{y:\tau_y e_i^* \in (\Lambda(\ell))^*} \tau_y \Psi_i \Big\} \Big\{ \sum_{x \in \Lambda(\ell)} (\theta \cdot x) \xi_x + \sum_{x \in \Lambda(\ell-n)} \tau_x(\theta \cdot F) \Big\} \Big\rangle_{\Lambda(\ell),m} \\ = \ell_*^{-d} \sum_{i=1}^d \tilde{\theta}_i \sum_{y:\tau_y e_i^* \in (\Lambda(\ell))^*} \sum_{x \in \Lambda(\ell)} \Big\langle \tau_y \Psi_i \big\{ (\theta \cdot x) \xi_x + 1_{x \in \Lambda(\ell-n)} \tau_x(\theta \cdot F) \big\} \Big\rangle_{\Lambda(\ell),m}.$$

To show (4.1), we first replace the expectation on the right-hand side of (4.3) under the canonical equilibrium measure $\nu_{\Lambda(\ell),m}$ by that under the grandcanonical equilibrium measure ν_{ρ} with $\rho = m/\ell_*^d$:

(4.5)
$$\frac{1}{2}\ell_*^{-d} \sum_{b=\tau_x e_i^* \in (\Lambda(\ell))^*} \left\langle c_b \Big[\theta \cdot e_i(\xi_{x+e_i} - \xi_x) - \pi_b \Big(\sum_{y \in \Lambda(\ell-n)} \tau_y F \cdot \theta \Big) \Big]^2 \right\rangle (m/\ell_*^d).$$

By the equivalence of ensembles (3.27) (with $|\Delta| \leq r(F)^d$) and recalling $|\theta| = 1$, the error for this replacement is bounded by

$$Cr(F)^{d}(1 + |||F|||_{0,\infty}^{2})\ell^{-d}$$

Then, if $b = \tau_x e_i^* \in (\Lambda(\ell - 2n))^*$, the sum in $y \in \Lambda(\ell - n)$ in (4.5) can be replaced by the sum in $y \in \mathbb{Z}^d$, Thus, by the translation invariance of ν_{ρ} , (4.5) is equal to

$$\frac{1}{2}\ell_*^{-d}\sum_{i=1}^d \sharp\{b \in (\Lambda(\ell-2n))^* : i\text{th directed}\} \\ \times \left\langle c_{e_i^*} \Big[\theta \cdot e_i(\xi_{e_i}-\xi_0) - \pi_{e_i^*}\Big(\sum_{y \in \mathbb{Z}^d} \tau_y F \cdot \theta\Big)\Big]^2 \right\rangle (m/\ell_*^d),$$

with an error bounded by

$$C(1+|||F|||_{0,\infty}^2)\frac{|(\Lambda(\ell))^* \setminus (\Lambda(\ell-2n))^*|}{\ell_*^d} \le C'(1+|||F|||_{0,\infty}^2)n\ell^{-1}.$$

Finally, noting that $\sharp \{b \in (\Lambda(\ell - 2n))^* : i$ th directed $\}/\ell_*^d = \frac{(2(\ell - 2n))^d}{(2\ell + 1)^d} = 1 + O(\frac{n}{\ell})$ and recalling n = r(F) + 1, the above expression can be replaced by

(4.6)
$$\frac{1}{2} \sum_{i=1}^{d} \left\langle c_{e_i^*} \left[\theta \cdot e_i(\xi_{e_i} - \xi_0) - \pi_{e_i^*} \left(\sum_{y \in \mathbb{Z}^d} \tau_y F \cdot \theta \right) \right]^2 \right\rangle (m/\ell_*^d),$$

with an error bounded by

$$Cr(F)(1 + ||F|||_{0,\infty}^2)\ell^{-1}.$$

However, by the identity at the bottom of p. 22 of [14], (4.6) is equal to $\frac{1}{2}\theta \cdot \hat{c}(m/\ell_*^d; F)\theta$ and we obtain the estimate (4.1).

To show (4.2), we first note the exchangeability under the canonical measure $\nu_{\Lambda(\ell),m}$, that is, η and $\eta^{x,y}$ have the same distributions under $\nu_{\Lambda(\ell),m}$ for $x, y \in \Lambda(\ell), x \neq y$. This is easily seen from the uniformity of $\nu_{\Lambda(\ell),m}$ or the invariance of $\sum_{z \in \Lambda(\ell)} \eta_z$ under the transform $\eta \mapsto \eta^{x,y}$. In the expression (4.4), the right-hand side looks like $O(\ell^{2d}/\ell^d)$, but due to the exchangeability under $\eta \mapsto \eta^{y,y+e_i}$, we have

$$\langle \tau_y \Psi_i \xi_x \rangle_{\Lambda(\ell),m} = \langle (\eta_{y+e_i} - \eta_y) \xi_x \rangle_{\Lambda(\ell),m} = 0$$

if $x \neq y, y + e_i$ and

$$\langle \tau_y \Psi_i \, \tau_x(\theta \cdot F) \rangle_{\Lambda(\ell),m} = 0$$

if $y, y + e_i \notin \tau_x(\text{supp}F)$, i.e., $y - x, y - x + e_i \notin \text{supp}F$, especially, if $|y - x| \ge r(F) + 1$. In particular, the first part of the sum (4.4) (related to 5.) is equal to

In particular, the first part of the sum (4.4) (related to ξ_x) is equal to

$$\ell_*^{-d} \sum_{i=1}^d \tilde{\theta}_i \sum_{y:\tau_y e_i^* \in (\Lambda(\ell))^*} \left\langle \tau_y \Psi_i \{ (\theta \cdot y) \xi_y + (\theta \cdot y + e_i) \xi_{y+e_i} \} \right\rangle_{\Lambda(\ell),m}$$
$$= \ell_*^{-d} \sum_{i=1}^d \tilde{\theta}_i \sum_{y:\tau_y e_i^* \in (\Lambda(\ell))^*} \left\{ -(\theta \cdot y) + (\theta \cdot y + e_i) \right\} \chi_{\Lambda(\ell),m}$$
$$= \frac{(2\ell)^d}{(2\ell+1)^d} (\tilde{\theta} \cdot \theta) \chi_{\Lambda(\ell),m},$$

by noting that both x = y and $y + e_i$ satisfy $x \in \Lambda(\ell)$ and then recalling the uniformity of $\nu_{\Lambda(\ell),m}$, where

$$\chi_{\Lambda(\ell),m} := -\langle \Psi_i \xi_0 \rangle_{\Lambda(\ell),m} \left(= -\langle (\xi_{e_i} - \xi_0) \xi_0 \rangle_{\Lambda(\ell),m} \right).$$

However, by the equivalence of ensembles (3.27), we have

$$|\chi_{\Lambda(\ell),m} - \chi(m/\ell_*^d)| \le C\ell^{-d}.$$

Therefore, the first part of the sum (4.4) behaves as

$$(\hat{\theta} \cdot \theta)\chi(m/\ell_*^d) + O(1/\ell).$$

On the other hand, as we pointed out above, the second part of the sum (4.4) (related to F) can be restricted to the terms of $x \in \Lambda(\ell)$: $|x - y| \leq r(F)$. Then, applying the equivalence of ensembles (3.27) (with $|\Delta| \leq r(F)^d$), we can replace it by

(4.7)
$$\ell_*^{-d} \sum_{i=1}^d \tilde{\theta}_i \sum_{y:\tau_y e_i^* \in (\Lambda(\ell))^*} \sum_{x \in \Lambda(\ell): |x-y| \le r(F)} \left\langle \tau_y \Psi_i \, \mathbf{1}_{x \in \Lambda(\ell-n)} \tau_x(\theta \cdot F) \right\rangle (m/\ell_*^d)$$

with an error bounded by

$$Cr(F)^{2d} \|F\|_{\infty} \ell^{-d}.$$

Denoting $\rho = m/\ell_*^d$, since $\langle \tau_y \Psi_i \ \tau_x(\theta \cdot F) \rangle_{\rho} = 0$ if $|x - y| \ge r(F) + 1$, we can drop the condition $|x - y| \le r(F)$ in the sum (4.7) and obtain for each *i*

$$\ell_*^{-d} \sum_{y:\tau_y e_i^* \in (\Lambda(\ell))^*} \sum_{x \in \Lambda(\ell-n): |x-y| \le r(F)} \langle \tau_y \Psi_i \ \tau_x(\theta \cdot F) \rangle_\rho$$

=
$$\ell_*^{-d} \sum_{y:\tau_y e_i^* \in (\Lambda(\ell))^*} \sum_{x \in \Lambda(\ell-n)} \langle \tau_y \Psi_i \ \tau_x(\theta \cdot F) \rangle_\rho = 0.$$

Here, to see that the last sum vanishes, we first observe

$$\sum_{y:\tau_y e_i^* \in (\Lambda(\ell))^*} \tau_y \Psi_i = \sum_{y:\tau_y e_i^* \in (\Lambda(\ell))^*} (\eta_{y+e_i} - \eta_y) = \sum_{z_+ \in \partial_+^i \Lambda(\ell)} \eta_{z_+} - \sum_{z_- \in \partial_-^i \Lambda(\ell)} \eta_{z_-},$$

where $\partial_{\pm}^{i}\Lambda(\ell) = \{z = (z_{j})_{j=1}^{d}; z_{i} = \pm \ell, |z_{j}| \leq \ell (j \neq i)\}$, and $\tau_{x}(\theta \cdot F)$ is $\Lambda(\ell - 1)$ -measurable for $x \in \Lambda(\ell - n)$. In particular, these variables are independent under ν_{ρ} . Since $\langle \eta_{z_{+}} - \eta_{z_{-}} \rangle_{\rho} = 0$, we see that the last sum vanishes.

Summing up these calculations and bounds, we get (4.2).

For the CLT variance of A_{ℓ} , it was shown in Theorem 5.1 of [14] that

(4.8)
$$\lim_{\ell,m\to\infty,m/\ell^d_*\to\rho} \ell^{-d}_* \Delta_{\ell,m,\zeta}(\theta \cdot A_\ell) = 2(\theta \cdot \widehat{c}^{-1}(\rho)\theta)\chi^2(\rho),$$

uniformly in $\rho \in [0,1]$ and $\eta \in \mathcal{X}$, where $\zeta = \eta|_{\Lambda(\ell)^c}$. This is the key in the gradient replacement and shown based on a characterization of closed forms given in Corollary 4.1 of [14]. However, this is not sufficient for our purpose. Then, recently more detailed analysis has been developed by applying the method in the quantitative homogenization theory and, as a by-product, a decay rate in the limit (4.8) was obtained as in the following theorem; see (7.12) in [9].

Theorem 4.2. ([9]) For $m \in [0, \ell^d_*] \cap \mathbb{Z}$, $\zeta \in \mathcal{X}_{\Lambda(\ell)^c}$ and $\theta \in \mathbb{R}^d : |\theta| = 1$, set

$$Q_{\ell}^{(6)}(m,\zeta,\theta) = \ell_{*}^{-d} \Delta_{\ell,m,\zeta}(\theta \cdot A_{\ell}) - 2(\theta \cdot \hat{c}^{-1}(m/\ell_{*}^{d})\theta)\chi^{2}(m/\ell_{*}^{d}).$$

Then, there exist C > 0 and $\alpha_1 > 0$ such that

$$|Q_{\ell}^{(6)}(m,\zeta,\theta)| \le C\ell^{-\alpha_1}$$

uniformly in $m \in [0, \ell^d_*] \cap \mathbb{Z}, \zeta \in \mathcal{X}_{\Lambda(\ell)^c}$ and $\theta \in \mathbb{R}^d : |\theta| = 1$.

The following corollary is immediate from Proposition 4.1 and Theorem 4.2, and gives a quantitative refinement of Corollary 5.1 of [14].

Corollary 4.3. Taking the supremum over all $m \in [0, \ell^d_*] \cap \mathbb{Z}$, $\zeta \in \mathcal{X}_{\Lambda(\ell)^c}$ and $\theta \in \mathbb{R}^d$: $|\theta| = 1$, set

$$Q_{\ell}(F) := \sup \left| \ell_*^{-d} \Delta_{\ell,m,\zeta} (\theta \cdot \{ D(m/\ell_*^d) A_{\ell} - (B_{\ell,\zeta} - H_{\ell,\zeta,F}) \}) - \frac{1}{2} \Big\{ \theta \cdot \widehat{c}(m/\ell_*^d; F) \theta - \theta \cdot \widehat{c}(m/\ell_*^d) \theta \Big\} \right|,$$

where $D(\rho)$ is the diffusion matrix defined in (1.7). Then, $Q_{\ell}(F)$ is estimated as

$$Q_{\ell}(F) \le C(1 + r(F)^{2d})(1 + |||F|||_{0,\infty}^2)\ell^{-1} + C\ell^{-\alpha_1}$$

for some C > 0 and $\alpha_1 > 0$.

Proof. Apply (4.1), (4.2) and Theorem 4.2; see the proof of Proposition 7.1 of [9] for more details.

5 Choice of β, ℓ, F, K

5.1 Summary of the error estimates

Here we summarize the estimates for the error terms appearing in (3.64). Note that $\lambda(t, v)$ is now determined from the hydrodynamic equation (1.11) as in Lemma 3.13 and, in particular, the Schauder estimate (2.4) is applicable. We use the exponent

$$\gamma := \max\{3\alpha_0, \alpha_0 + 1\} > 0$$

determined from $\alpha_0 > 0$ in (2.4). In the following, the constant $C = C_T > 0$ changes from line to line and depends on T.

The term $Q_{N,\ell,\beta,K}^{\Omega_1}(\lambda,F)$ is the error for Ω_1 with the non-gradient microscopic current function. It is given in (3.47) as

(5.1)
$$Q_{N,\ell,\beta,K}^{\Omega_1}(\lambda,F) = \frac{C}{\beta}K + \beta^2 Q_{N,\ell}^{(1)}(\lambda,F) + \beta Q_{\ell}^{(2)}(\lambda,F) + Q_{N,\ell}^{(3)}(\lambda),$$

where the first three terms are obtained in Theorem 3.5 and the last one in Lemma 3.7. Here, the term $Q_{N,\ell}^{(1)} \equiv Q_{N,\ell}^{(1)}(\lambda, F)$ is estimated in (3.30), and together with the Schauder estimate (2.4), we have

(5.2)
$$|Q_{N,\ell}^{(1)}| \le CN^{-1}K^{\gamma}\ell^{2d+4}(1+|||F|||_{0,\infty}^3).$$

The term $Q_{\ell}^{(2)} \equiv Q_{\ell}^{(2)}(\lambda, F)$ is for the gradient replacement and is estimated in (3.31) together with (2.4) as

$$(5.3) |Q_{\ell}^{(2)}| \le CK^{\gamma}Q_{\ell}(F)$$

and by Corollary 4.3,

(5.4)
$$Q_{\ell}(F) \le C(1 + r(F)^{2d})(1 + |||F|||_{0,\infty}^2)\ell^{-1} + C\ell^{-\alpha_1}.$$

for some $\alpha_1 > 0$. The term $Q_{N,\ell}^{(3)} \equiv Q_{N,\ell}^{(3)}(\lambda)$ given in Lemma 3.7 is independent of F and it is estimated from above as

(5.5)
$$Q_{N,\ell}^{(3)} \le CK^{\gamma}(\ell^{-1} + N^{-1}) + C\beta K^{\gamma}\ell^{-1} + \frac{C}{\beta}K + C\beta^2 K^{\gamma} N^{-1}\ell^{2d-2}$$

The term $Q_{N,\ell,\delta_*}^{LD}(\lambda,F)$ appears in the large deviation estimate (see Theorems 3.11 and 3.9) after applying the entropy inequality and, by noting that we have $\|\partial_v G_i\|_{\infty} \leq CK^{\gamma}$, $\|\partial_\rho G_i\|_{\infty} \leq CK^{\gamma}$ for i = 1, 2 by Schauder estimates (2.3), (2.4) and the comparison theorem (Lemma 7.1) used for $\|\lambda\|_{\infty}$, it is estimated as

(5.6)
$$|Q_{N,\ell,\delta_*}^{LD}(\lambda,F)| \le CN^{-1}K^{\gamma}(\ell+\|F\|_{\infty}) + C\ell^{-d}(\log\ell+K^{\gamma}).$$

The term $Q_N^{En}(\lambda, F)$ is the error in the entropy calculation and is estimated as in (3.3). Indeed, together with the Schauder estimate (2.4) and the comparison theorem (Lemma 7.1), we have

(5.7)
$$|Q_N^{En}(\lambda, F)| \le CN^{-1}K^{\gamma}(1 + |||F|||_{2,\infty})^3,$$

if the condition (3.4), in particular,

(5.8)
$$N^{-1}K^{\gamma} |||F|||_{0,\infty} \le 1$$

is satisfied.

The term $Q_{N,\ell}^{\Omega_2}(\lambda, F)$ estimated in (3.49) is the error for the one-block estimate (3.25) applied for the gradient term Ω_2 , adding the discretization error of (3.51). Applying the Schauder estimate (2.4), it is bounded as

(5.9)
$$|Q_{N,\ell}^{\Omega_2}(\lambda,F)| \leq CK^{\gamma} \left(N^{-1} K^{1/2} \ell^{(d+2)/2} + \ell^{-d} (1+r(F)^d) \right) \\ \times (1+ |||F|||_{0,\infty}^2 + |||F|||_{1,\infty}) + CK^{\gamma} N^{-1}.$$

5.2 Choice of β , ℓ , F, K

By Lemma 4.4 and Proposition 6.9 of [9] (see also Theorem 1.2 and Section 7 of [9]), one can find a sequence of local functions $\Phi_n = \Phi_n(\eta) \in (\mathcal{F}_0)^d$ on \mathcal{X} such that

(5.10)
$$r(\Phi_n) \le n, \quad \sup_{\rho \in [0,1]} \|R(\rho; \Phi_n)\| \le C_2 n^{-\alpha_2}, \quad \|\Phi_n\|_{\infty} \le C_2 n^2 \log n,$$

for some $C_2 > 0$ and $\alpha_2 > 0$. We will choose $n = n(N), \ell = \ell(N), K = K(N)$ and $\beta = \beta(n)$, from which we can determine $\beta(N) := \beta(n(N))$ and $F_N := \Phi_{n(N)}$ (i.e. Φ_n with n = n(N)), respectively. In fact, we will choose $1 \le n = N^{a_1} \ll \ell = N^{a_2} \ll N$ with exponents

(5.11)
$$0 < a_1 < a_2 := 1/(2d+5) < 1.$$

In particular, the radius $r(\Phi_n)$ of the support of Φ_n is much smaller than ℓ so that the arguments in Section 4 work smoothly for $F = F_N$; note that n in $H_{\ell}(\eta)$ in (3.37) is taken as n = r(F) + 1 so that it is n + 1 with n given here.

For $K = K(N) \ge 1$, we assume

(5.12)
$$1 \le K \le \overline{K}_{\delta}(N) := \delta \log N$$

and will make $\delta > 0$ sufficiently small. In particular, by (5.11) and (5.12), the condition (3.24) holds (note that $\delta > 0$ in (3.24) is different from that in (5.12)). Moreover, also noting (5.10), the condition (5.8) holds. We will take $\beta = n^{\alpha_2/2}$.

In (3.64), setting $c_* := T/\delta_*$ with $\delta_* > 0$ chosen in Lemma 3.13, the factor e^{Kt/δ_*} is bounded by $e^{c_*K} \leq N^{c_*\delta}$ for $t \in [0,T]$. We also bound $K, K^{\gamma} \leq CN^{c_*\delta}$ for simplicity. Below we give a bound for each of the nine terms in (3.64) (recall that $Q_{N,\ell,\beta,K}^{\Omega_1}(\lambda, F_N)$ consists of four terms as in (3.47) or equivalently (5.1)) first choosing $a_1 > 0$ small enough and then $c_*\delta = T\delta/\delta_*$, so $\mathfrak{c} = T\delta$ in Proposition 2.1 and Theorem 1.1, further small. The constants \overline{C} and κ below may change from line to line but they can always be taken as $\overline{C} = \overline{C}_T > 0$ and $\kappa > 0$ (\overline{C} large enough depending on T, while κ small enough depending on \mathfrak{c} but independently of T and δ).

1. The term $\beta^2 Q_{N,\ell}^{(1)}(\lambda, F) \cdot e^{c_* K}$ in (5.1) with $e^{c_* K}$ is bounded, from (5.2) and (5.10), by

$$Cn^{\alpha_2}N^{-1}\ell^{2d+4}n^{3d+7}N^{2c_*\delta} \le \bar{C}N^{-\kappa}.$$

Note that $\ell^{2d+4} = N^{(2d+4)/(2d+5)}$ by (5.11).

2. The term $\beta Q_{\ell}^{(2)}(\lambda, F) \cdot e^{c_* K}$ in (5.1) is bounded, from (5.3), (5.4) and (5.10), by

$$Cn^{\alpha_2/2} (n^{4d+5}\ell^{-1} + \ell^{-\alpha_1}) N^{2c_*\delta} \le \bar{C}N^{-\kappa}.$$

3. The upper bound of $Q_{N,\ell}^{(3)}(\lambda) \cdot e^{c_*K}$ in (5.1) is given, from (5.5), by

$$C\Big\{\ell^{-1} + N^{-1} + n^{\alpha_2/2}\ell^{-1} + n^{-\alpha_2/2} + n^{\alpha_2}N^{-1}\ell^{2d-2}\Big\}N^{2c_*\delta} \le \bar{C}N^{-\kappa}.$$

4. The term $|Q_{N\ell \delta_*}^{LD}(\lambda, F)| \cdot e^{c_*K}$ is bounded, from (5.6) and (5.10), by

$$C\left\{N^{-1}(\ell+n^3) + \ell^{-d}\log\ell\right\}N^{2c_*\delta} \le \bar{C}N^{-\kappa}.$$

5. Noting that (5.8) is satisfied, the term $Q_N^{En}(\lambda, F) \cdot e^{c_*K}$ is bounded, from (5.7) and (5.10), by

$$CN^{-1}n^{3d+13}N^{2c_*\delta} \le \bar{C}N^{-\kappa}.$$

6. The term $|Q_{N,\ell}^{\Omega_2}(\lambda,F)| \cdot e^{c_*K}$ is bounded, from (5.9) and (5.10), by

$$C(N^{-1}\ell^{(d+2)/2} + \ell^{-d}n^d)n^{2d+5}N^{3c_*\delta} + CN^{2c_*\delta}N^{-1} \le \bar{C}N^{-\kappa}.$$

7. The term $(\beta + 1)K^{\gamma} \sup_{\rho \in [0,1]} ||R(\rho; F)|| \cdot e^{c_*K}$ is bounded, from (5.10), by

$$Cn^{\alpha_2/2}N^{2c_*\delta}n^{-\alpha_2} \le \bar{C}N^{-\kappa},$$

by choosing $\delta > 0$ small, once $a_1 > 0$ is determined in the above procedure.

8. The term $\frac{C}{\beta}K \cdot e^{c_*K}$ in (5.1) with e^{c_*K} is bounded by

$$Cn^{-\alpha_2/2}N^{2c_*\delta} \le \bar{C}N^{-\kappa},$$

by choosing $\delta > 0$ small enough.

9. Finally, by the assumption $h_N(0) \leq C_1 N^{-\kappa_1}, C_1, \kappa_1 > 0$ of Proposition 2.1, we have

$$h_N(0)e^{c_*K} \le \bar{C}N^{-\kappa}.$$

Nine terms in (3.64) are all bounded by $\bar{C}N^{-\kappa}$ and therefore one can conclude the proof of Proposition 2.1 as already stated at the end of Section 3.7.

6 Schauder estimates

Here we give the proof of the Schauder estimates (2.3) for the solution $\rho = \rho(t, v) \equiv \rho_K(t, v)$ of (1.11). We apply the results given in [21] for linear second order parabolic partial differential equations (PDEs). See also [20], but an explicit dependence of the constants in the estimates on several data of the equation is not clearly indicated as in (6.6), (6.7) below.

Before showing the Schauder estimates, we briefly note that the equation (1.11) has a unique classical solution $\rho(t, v) \in C^{1,3}([0, \infty) \times \mathbb{T}^d)$ under the condition $\rho_0 \in C^4(\mathbb{T}^d)$ for its initial value; recall (1.12) in which $\rho_0 \in C^5(\mathbb{T}^d)$ is assumed. In fact, noting that (1.11) is a quasi-linear equation in divergence form (0.1) in Chapter V of [20], p. 417, by Theorem 6.1 of [20], p. 452, we see that it has a unique classical solution $\rho(t, v) \in C^{1,2}([0, \infty) \times \mathbb{T}^d)$ if $\rho_0 \in C^3(\mathbb{T}^d)$. Note that the consistency condition (6.3) assumed in Theorem 6.1 is unnecessary in our setting, since \mathbb{T}^d has no boundary. To obtain $C^{1,3}$ -property, we consider the equation for the derivative $\rho_{(k)} := \partial_{v_k}\rho(t, v)$ for each fixed $k : 1 \leq k \leq d$. By a simple calculation, we see that $\rho_{(k)}$ satisfies the equation (0.1) of [20] with

$$a_{i}(v,t,p) = \sum_{j=1}^{d} D_{ij}(\rho(t,v))p_{j} + \sum_{j=1}^{d} D'_{ij}(\rho(t,v))\partial_{v_{k}}\rho(t,v)\partial_{v_{j}}\rho(t,v),$$

$$a(v,t) = -f'(\rho(t,v))\partial_{v_{k}}\rho(t,v),$$

where we regard that $\rho(t, v)$, $\partial_{v_k}\rho(t, v)$ and $\partial_{v_j}\rho(t, v)$ are all given. Then, applying Theorem 6.1 of [20] again, we see that $\rho_{(k)} \in C^{1,2}([0,\infty) \times \mathbb{T}^d)$ if $\partial_{v_k}\rho_0 \in C^3(\mathbb{T}^d)$ for each k. This shows $\rho(t, v) \in C^{1,3}([0,\infty) \times \mathbb{T}^d)$ under the condition $\rho_0 \in C^4(\mathbb{T}^d)$.

Now we show the Schauder estimates. To apply the results of [21], the first step is to show the Hölder continuity of the solution of the nonlinear PDE (1.11).

For this purpose, first in (1.11) dropping the term $Kf(\rho(t, v))$ and replacing $D(\rho(t, v))$ by $D(t, v) = \{D_{ij}(t, v)\}_{1 \le i,j \le d}$ which is symmetric matrix-valued Borel measurable function on $[0, T] \times \mathbb{T}^d$ satisfying a bound similar to (1.9): $c_* |\theta|^2 \le (\theta, D(t, v)\theta) \le c^* |\theta|^2$ for $\theta \in \mathbb{R}^d$ with $c_*, c^* > 0$, we consider a solution $u^{(0)} = u^{(0)}(t, v)$ of the linear parabolic PDE

(6.1)
$$\partial_t u^{(0)} = \sum_{i,j=1}^d \partial_{v_i} \{ D_{ij}(t,v) \partial_{v_j} u^{(0)} \}, \quad t \ge 0, v \in \mathbb{T}^d,$$

with initial value u_0 . Let $p(t, v; s, \xi)$, $v, \xi \in \mathbb{T}^d$, $0 \leq s < t \leq T$ be the associated fundamental solution. Then, we have the following Nash Hölder estimates; cf. [27], [24] and related estimates can also be found in [20].

Lemma 6.1. There exist $\sigma \in (0,1]$ and C > 0 such that

(6.2)
$$|u^{(0)}(t_1, v_1) - u^{(0)}(t_2, v_2)| \le C ||u_0||_{\infty} \Big(\frac{|t_1 - t_2|^{1/2} \vee |v_1 - v_2|}{(t_1 \wedge t_2)^{1/2}} \Big)^{\sigma},$$

for $t_1, t_2 \in (0, T], v_1, v_2 \in \mathbb{T}^d$ and

(6.3)
$$|p(t_1, v_1; 0, \xi_1) - p(t_2, v_2; 0, \xi_2)| \le \frac{C}{(t_1 \wedge t_2)^{d/2}} \left(\frac{|t_1 - t_2|^{1/2} \vee |v_1 - v_2| \vee |\xi_1 - \xi_2|}{(t_1 \wedge t_2)^{1/2}}\right)^{\sigma},$$

for $t_1, t_2 \in (0, T], v_1, v_2, \xi_1, \xi_2 \in \mathbb{T}^d$.

Proof. To show (6.2), we separate into two cases: $(t_1 \vee t_2) \ge 2(t_1 \wedge t_2)$ and $(t_1 \vee t_2) < 2(t_1 \wedge t_2)$. In the case $(t_1 \vee t_2) \ge 2(t_1 \wedge t_2)$, since $|t_1 - t_2| = (t_1 \vee t_2) - (t_1 \wedge t_2) \ge (t_1 \wedge t_2)$, (6.2) is obvious with C = 2, noting $||u^{(0)}(t)||_{\infty} \le ||u_0||_{\infty}$ by the maximum principle; see Lemma 7.1 below with $f \equiv 0$.

In the other case $(t_1 \vee t_2) < 2(t_1 \wedge t_2)$, we take $t_0 = 2(t_1 \wedge t_2)$, $R = \sqrt{2}(t_1 \wedge t_2)^{1/2}$, $\delta = \sqrt{2/3}$ and $x_0 = v_1$ in Theorem 4.1 of [24] (or Theorem II.1.8 of [27], although it is stated only for the time-homogeneous case). Note that $t_0 - R^2 = 0$ and $t_1, t_2 \in (t_0 - (\delta R)^2, t_0) = (2/3(t_1 \wedge t_2), 2(t_1 \wedge t_2))$. Then, by Theorem 4.1 of [24], (6.2) holds if $|v_1 - v_2| \le \sqrt{4/3}(t_1 \wedge t_2)^{1/2}$. If $|v_1 - v_2| \ge \sqrt{4/3}(t_1 \wedge t_2)^{1/2}$, (6.2) is obvious with C = 2.

To show (6.3), taking $\delta = (t_1 \wedge t_2)^{1/2}$ in Corollary 4.2 of [24] (or Corollary II.1.9 of [27]), this holds if $|v_1 - v_2| \vee |\xi_1 - \xi_2| \leq (t_1 \wedge t_2)^{1/2}$. Conversely, if $|v_1 - v_2| \vee |\xi_1 - \xi_2| \geq (t_1 \wedge t_2)^{1/2}$, by Aronson's Gaussian upper bound (see Theorem 1.2 of [24]), we have

$$|p(t_1, v_1; 0, \xi_1) - p(t_2, v_2; 0, \xi_2)| \le p(t_1, v_1; 0, \xi_1) + p(t_2, v_2; 0, \xi_2)$$
$$\le Ct_1^{-d/2} + Ct_2^{-d/2} \le 2C(t_1 \wedge t_2)^{-d/2}$$

and this is bounded by the right-hand side of (6.3) (with C replaced by 2C).

This lemma is used to show the Hölder continuity of the solution $\rho(t, v) \equiv \rho_K(t, v), K \geq 1$ of (1.11) with $\rho(0) = \rho_0$. The regularity of ρ_0 allows to remove the singularity of $\rho(t, v)$ near t = 0.

Lemma 6.2. Assume the condition (1.12) for ρ_0 . Then, we have

(6.4)
$$|\rho(t_1, v_1) - \rho(t_2, v_2)| \le CK (|t_1 - t_2|^{\sigma/2} + |v_1 - v_2|^{\sigma}), \quad t_1, t_2 \in [0, T], v_1, v_2 \in \mathbb{T}^d,$$

for some $\sigma \in (0,1]$. Since $D(\rho) = \{D_{ij}(\rho)\}_{1 \le i,j \le d} \in C^{\infty}([0,1])$, we have a similar Hölder estimate for $D_{ij}(\rho(t,v))$.

Proof. The proof is similar to that of Corollary 2.6 of [13] (in the discrete setting). First, let D(t, v) be as above and let g(t, v) be a bounded Borel measurable function. We consider the linear PDE

$$\partial_t u = \sum_{i,j=1}^d \partial_{v_i} \{ D_{ij}(t,v) \partial_{v_j} u \} + g(t,v), \quad t \ge 0, v \in \mathbb{T}^d,$$

with $u(0) = u_0$. Then, by Duhamel's formula, we have

$$u(t,v) = u^{(0)}(t,v) + \int_0^t ds \int_{\mathbb{T}^d} g(s,\xi) p(t,v;s,\xi) d\xi,$$

where $u^{(0)}(t, v)$ is the solution of (6.1). Then, by Lemma 6.1 (actually only by (6.2)), similar to the proof of Theorem 2.2 of [13], we get

(6.5)
$$|u(t_1, v_1) - u(t_2, v_2)| \le C(||u_0||_{\infty} + ||g||_{\infty}) \Big(\frac{|t_1 - t_2|^{1/2} + |v_1 - v_2|}{(t_1 \wedge t_2)^{1/2}}\Big)^{\sigma},$$

for $t_1, t_2 \in (0, T], v_1, v_2 \in \mathbb{T}^d$.

To improve the regularity near t = 0, we apply a trick similar to the one used in the proof of Theorem 2.5 of [13]. We supplementarily consider the heat equation on \mathbb{T}^d

$$\partial_s U = \Delta U, \quad s \in (0, 1]$$

with $U(0) = \rho_0$, and set $\hat{U}(t) := U(1-t)$ and $\hat{h}(t,v) := -\Delta \hat{U}(t,v)$ for $t \in [0,1)$. Then, $\hat{U}(t,v)$ satisfies the equation

$$\partial_t \hat{U} = -\Delta \hat{U} = \Delta \hat{U} + 2\hat{h}$$

Since $h(t) := \hat{h}(1-t) = -\Delta U(t)$ satisfies $\partial_t h(t) = \Delta h(t)$ with $h(0) = -\Delta U(0) = -\Delta \rho_0$, we see by the maximum principle that $||h(t)||_{\infty} \le ||\Delta \rho_0||_{\infty} < \infty$ by (1.12).

To apply (6.5) for our equation (1.11), define $\hat{\rho}(t,v)$, $\{\hat{D}_{ij}(t,v)\}$ and $\hat{g}(t,v)$, respectively, by

$$\hat{\rho}(t,v) = \begin{cases} \rho(t-1,v), & t \in [1,T+1], \\ \hat{U}(t,v), & t \in [0,1). \end{cases}$$
$$\hat{D}_{ij}(t,v) = \begin{cases} D_{ij}(\rho(t-1,v)), & t \in [1,T+1] \\ \delta_{ij}, & t \in [0,1), \end{cases}$$
$$\hat{g}(t,v) = \begin{cases} Kf(\rho(t-1,v)), & t \in [1,T+1], \\ 2\hat{h}(t,v), & t \in [0,1), \end{cases}$$

where $\rho(t, v) \equiv \rho_K(t, v)$ is the solution of (1.11) with initial value ρ_0 . Then, these functions satisfy

$$\partial_t \hat{\rho} = \sum_{i,j=1}^d \partial_{v_i} (\hat{D}_{ij}(t,v) \partial_{v_i} \hat{\rho}) + \hat{g}(t,v),$$

with $\hat{\rho}(0, v) = U(1)$. Note that $||U(1)||_{\infty} \leq ||\rho_0||_{\infty}$. So, noting that $||\hat{g}||_{\infty} \leq K ||f||_{\infty} + 2||\Delta\rho_0||_{\infty}$ and $\{\hat{D}_{ij}\}$ satisfies the uniform positive-definiteness and boundedness conditions, from (6.5) with T replaced by T + 1, we have

$$|\hat{\rho}(t_1, v_1) - \hat{\rho}(t_2, v_2)| \le C(\|\rho_0\|_{\infty} + \|\hat{g}\|_{\infty}) \Big(\frac{|t_1 - t_2|^{1/2} + |v_1 - v_2|}{(t_1 \wedge t_2)^{1/2}}\Big)^{\sigma},$$

for $t_1, t_2 \in (0, T+1]$, $v_1, v_2 \in \mathbb{T}^d$. The desired bound (6.4) is obtained by restricting the above estimate to [1, T+1] and shifting it by 1 in time.

Now we apply Theorem 4.8 of [21], in particular the detailed estimate given on p. 59 for a solution u of the linear equation (4.22) of [21] cited below. Before explaining it, let us first recall several Hölder norms for functions on $\Omega = [0, T] \times \mathbb{T}^d$ from pp. 46–47 of [21]. The sup-norm on Ω is denoted by $\|\cdot\|_{\infty} = |\cdot|_0 \equiv |\cdot|_0^{(0)}$ and let $\alpha \in (0, 1]$. The (semi) norms $[u]_{\alpha}^{(b)}, b = 0, 1, |u|_{\alpha}^{(1)}$ and $|u|_{1+\alpha}^* \equiv |u|_{1+\alpha}^{(0)}$ are defined for functions u on Ω which may have a singularity like $t^{-1/2}$ or others near t = 0, respectively, by

$$\begin{split} &[u]_{\alpha}^{(b)} = \sup_{X \neq Y} (t \wedge s)^{(\alpha+b)/2} \frac{|u(X) - u(Y)|}{|X - Y|^{\alpha}}, \quad b = 0, 1, \\ &|u|_{\alpha}^{(1)} = |u|_{0}^{(1)} + [u]_{\alpha}^{(1)}, \quad |u|_{0}^{(1)} = \sup_{\Omega} t^{1/2} |u(X)|, \\ &|u|_{1+\alpha}^{(0)} := ||u||_{\infty} + |\partial_{v}u|_{0}^{(1)} + [u]_{1+\alpha}^{(0)} + \langle u \rangle_{1+\alpha}^{(0)} \\ &\equiv ||u||_{\infty} + \sup_{\Omega} t^{1/2} |\partial_{v}u(X)| + \sup_{X \neq Y} (t \wedge s)^{(1+\alpha)/2} \frac{|\partial_{v}u(X) - \partial_{v}u(Y)|}{|X - Y|^{\alpha}} \\ &+ \sup_{X \neq Y, x = y} (t \wedge s)^{(1+\alpha)/2} \frac{|u(X) - u(Y)|}{|X - Y|^{1+\alpha}}, \end{split}$$

where $X = (t, v), Y = (s, v') \in \Omega$ and $|X - Y| = \max\{|x - y|, |t - s|^{1/2}\}$. Note that $d(X, Y) = \min\{d(X), d(Y)\} = \min\{t^{1/2}, s^{1/2}\} = (t \wedge s)^{1/2}$ and $\langle u \rangle_{\alpha}^{(1)} = 0$ in the norm $|u|_{\alpha}^{(1)}$ on p. 47 of [21]. The norm $|u|_{1+\alpha}$ for functions u without singularity near t = 0 is defined on p. 46 of [21] as

$$|u|_{1+\alpha} := ||u||_{\infty} + ||\partial_{v}u||_{\infty} + \sup_{X \neq Y} \frac{|\partial_{v}u(X) - \partial_{v}u(Y)|}{|X - Y|^{\alpha}} + \sup_{X \neq Y, x = y} \frac{|u(X) - u(Y)|}{|X - Y|^{1+\alpha}}.$$

The detailed estimate mentioned above clarifies the dependence of the constants in the estimate on various data; see (6.6) below. The equation (4.22) of [21] is defined as follows on Ω and written by dropping the sums in i, j as

(4.22)
$$-\partial_t u + \partial_{v_i} \left(a^{ij} \partial_{v_j} u + b^i \partial_{v_i} u \right) + c^i \partial_{v_i} u + c^0 u = \partial_{v_i} f^i + g,$$

where a^{ij}, b^i, f^i are α -Hölder continuous and c^i, c^0, g are bounded measurable functions on Ω that satisfy

(4.20a)
$$\lambda |\xi|^2 \le a^{ij} \xi_i \xi_j \le \Lambda |\xi|^2, \quad [a^{ij}]^{(0)}_{\alpha} \le A,$$

$$(4.20b) |b^i|^{(1)}_{\alpha} \le B,$$

(4.20b)
$$|b^{i}|_{\alpha}^{(1)} \leq B,$$

(4.20c) $||c^{i}||_{\infty} \leq c_{1}, \quad ||c^{0}||_{\infty} \leq c_{2},$
(4.21) $|f^{i}|_{\alpha}^{(1)} \leq F, \quad ||g||_{\infty} \leq G,$

(4.21)
$$|f^i|_{\alpha}^{(1)} \le F, \quad ||g||_{\infty} \le G$$

for some positive constants $\lambda, \Lambda, A, B, c_1, c_2, F$ and G. Then, we have

(6.6)
$$|u|_{1+\alpha}^* \left(\equiv |u|_{1+\alpha}^{(0)} \right) \le C(d,\alpha,\lambda,\Lambda)(M||u||_{\infty} + F + G)$$

with

(6.7)
$$M = [1 + A + c_1]^{(1+\alpha)/\alpha} + \sum_i (|b^i|_0^{(1)})^{1+\alpha} + c_2 + \sum_i [b^i]_\alpha^{(1)}.$$

In our applications, we will take $b^i = c^i = 0$. Note that the above conditions (4.20c) and (4.21) are simplifications of those in [21] by noting that $\|\cdot\|_{p,\delta}^{(b)} \leq C \|\cdot\|_{\infty}$ for b = 1, 2, but these will suffice for our purpose.

First, we consider (1.11) to be a linear equation (4.22) for $u = \rho$ with $a^{ij} = D_{ij}(\rho(t, v))$, $g = -Kf(\rho(t, v))$ where $\rho(t, v)$ is already given and $b^i = c^i = c^0 = f^i = 0$. Then, since (6.4) implies $[a^{ij}]^{(0)}_{\sigma} \leq CK$ (note that $(t \wedge s)^{\alpha/2} \leq T^{\alpha/2}$ is bounded), one can take $\alpha = \sigma$, A = CK in the estimate (4.20a) and G = CK in (4.21). So, noting $M = [1 + A]^{(1+\sigma)/\sigma} \leq C K^{(1+\sigma)/\sigma}$ and $||u||_{\infty} \leq 1$, we obtain by (6.6) the estimate:

(6.8)
$$|\rho|_{1+\sigma}^* \le CK^{(1+\sigma)/\sigma},$$

for the solution ρ of (1.11). Here and in the following, we use the comparison theorem for (1.11), that is, under our assumption (1.12) for the initial value ρ_0 , we have $0 < c \leq c$ $\rho(t, v) \leq 1 - c < 1$ for some $c \in (0, 1/2)$; see Lemma 7.1 below.

Considering the extended system in time similar to the proof of Lemma 6.2 (see also Theorem 4.2 of [13]), the regularity of the initial value $\rho_0(v)$ improves the estimate (6.8) into that without singularity near t = 0. In fact, we have the following lemma.

Lemma 6.3. For the norm without singularity, we have

(6.9)
$$|\rho|_{1+\sigma} \le CK^{(1+\sigma)/\sigma}.$$

Proof. Since we need to extend a^{ij} to \hat{a}^{ij} while preserving its σ -Hölder property, we modify the choice of \hat{D}_{ij} in the proof of Lemma 6.2. We take $\hat{b}^i = \hat{c}^i = \hat{c}^0 = \hat{f}^i = 0$ and define $\hat{u}(t,v), \{\hat{a}^{ij}(t,v)\}\$ and $\hat{g}(t,v),$ respectively, by

(6.10)
$$\hat{u}(t,v) = \begin{cases} \rho(t-1,v), & t \in [1,T+1], \\ \hat{U}(t,v), & t \in [0,1). \end{cases}$$
$$\hat{a}^{ij}(t,v) = \begin{cases} D_{ij}(\rho(t-1,v)), & t \in [1,T+1], \\ D_{ij}(\rho_0(v)), & t \in [0,1), \end{cases}$$

$$\hat{g}(t,v) = \begin{cases} -Kf(\rho(t-1,v)), & t \in [1,T+1], \\ -2\hat{h}(t,v), & t \in [0,1), \end{cases}$$

where $\hat{U}(t) := U(1-t)$ is defined from the solution $U(s), s \in [0,1]$ of the equation

$$\partial_s U = L_{\rho_0} U \equiv \partial_{v_i} (D_{ij}(\rho_0(v)) \partial_{v_j} U), \quad s \in (0, 1],$$

with initial value $U(0) = \rho_0$, and $\hat{h}(t, v) := -L_{\rho_0}\hat{U}(t, v), t \in [0, 1]$. Then, $\hat{U}(t)$ satisfies the equation

$$\partial_t \hat{U} = -L_{\rho_0} \hat{U} = L_{\rho_0} \hat{U} + 2\hat{h}.$$

For \hat{g} , the measurability is sufficient.

Note that $h(t) := \hat{h}(1-t) = -L_{\rho_0}U(t)$ satisfies the equation $\partial_t h(t) = L_{\rho_0}h(t)$ and $h(0) = \hat{h}(1) = -L_{\rho_0}U(0) = -L_{\rho_0}\rho_0$. Thus, according to the maximum principle noting

$$L_{\rho_0}u = D_{ij}(\rho_0)\partial_{v_i}\partial_{v_j}u + D'_{ij}(\rho_0)\partial_{v_i}\rho_0 \cdot \partial_{v_j}u,$$

we see

$$\|h(t)\|_{\infty} \le \|L_{\rho_0}\rho_0\|_{\infty} \le C(\|\partial^2 \rho_0\|_{\infty} + \|\partial \rho_0\|_{\infty}^2) < \infty,$$

since $\rho_0 \in C^2(\mathbb{T}^d)$ by our condition (1.12). So we have $\|\hat{g}\|_{\infty} \leq CK$.

Since $[\hat{a}^{ij}]^{(0)}_{\sigma} \leq CK$, (6.8) holds for the extended solution $\hat{u}(t,v)$ on [0, T+1]. Restricting this estimate to [1, T+1], (6.9) follows.

Second, we take the derivative of (1.11) in v_k and obtain by dropping the sum in i, j

(6.11)
$$\partial_t \rho_{v_k} = \partial_{v_i} \left(\partial_{v_k} \left\{ D_{ij}(\rho) \partial_{v_j} \rho \right\} \right) + K f'(\rho) \rho_{v_k} \\ = \partial_{v_i} \left\{ D_{ij}(\rho) \partial_{v_j} \rho_{v_k} \right\} - \partial_{v_i} f^{i,(1)}(\rho) + K f'(\rho) \rho_{v_k}$$

where

$$f^{i,(1)}(\rho) := -\sum_{j=1}^{d} D'_{ij}(\rho) \rho_{v_k} \rho_{v_j}.$$

We consider this equation to be a linear equation (4.22) for $u = \rho_{v_k}$ with $a^{ij} = D_{ij}(\rho)$, $b^i = c^i = 0$, $f^i = f^{i,(1)}(\rho)$ (here both $\rho_{v_k} = \rho_{v_k}(t, v)$ and $\rho_{v_j} = \rho_{v_j}(t, v)$ are considered already given) and $c^0 = Kf'(\rho)$. We may add a bounded function g (actually, \hat{g}) to apply for the proof of the next Lemma 6.4. Then, one can take A = CK, $F = CK(K^{(1+\sigma)/\sigma})^2$ by using (6.4) (for Hölder estimate for $D'(\rho)$) and (6.9) (noting that we can drop $t^{1/2}$ and $(t \wedge s)^{(1+\alpha)/2}$ in the norm $|f^i|^{(1)}_{\alpha}$, since these are bounded) and $c_2 = CK$. Thus, by (6.6) with G for g (for later use, currently g = 0), we obtain the estimate:

(6.12)
$$|u|_{1+\sigma}^* \le C(M||u||_{\infty} + K^{2(1+\sigma)/\sigma+1} + G)$$

with $M = [1 + A]^{(1+\sigma)/\sigma} + K$. Therefore, noting $||u||_{\infty} = ||\rho_{v_k}||_{\infty} \leq CK^{(1+\sigma)/\sigma}$ by (6.9) and G = 0, we have

$$(6.13) \qquad \qquad |\partial\rho|_{1+\sigma}^* \le CK^{2(1+\sigma)/\sigma+1}.$$

We can apply a similar argument to the proof of Lemma 6.3 to remove the singularity near t = 0.

Lemma 6.4. For the norm without singularity, we have

$$(6.14) \qquad \qquad |\partial\rho|_{1+\sigma} \le CK^{2(1+\sigma)/\sigma+1}$$

Proof. We take $\{\hat{a}^{ij}(t,v)\}$ as in (6.10) and $\hat{b}^i = \hat{c}^i = 0$. Then, for each k, we define $\hat{u}(t,v)$, $\{\hat{f}^i(t,v)\}, \hat{c}^0(t,v)$ and $\hat{g}(t,v)$, respectively, by

$$\hat{u}(t,v) = \begin{cases} \rho_{v_k}(t-1,v), & t \in [1,T+1], \\ \hat{V}(t,v), & t \in [0,1). \end{cases}$$
(6.15)
$$\hat{f}^i(t,v) = \begin{cases} f^{i,(1)}(\rho(t-1,v)), & t \in [1,T+1], \\ f^{i,(1)}(\rho_0(v)), & t \in [0,1), \end{cases}$$

$$\hat{c}^0(t,v) = \begin{cases} Kf'(\rho(t-1,v)), & t \in [1,T+1], \\ 0, & t \in [0,1), \end{cases}$$

$$\hat{g}(t,v) = \begin{cases} 0, & t \in [1,T+1], \\ -2\hat{h}(t,v), & t \in [0,1), \end{cases}$$

where $\hat{V}(t) \equiv \hat{V}_k(t) := V(1-t)$ is defined from the solution $V(s), s \in [0,1]$ of the equation

$$\partial_s V = L_{\rho_0} V - \partial_{v_i} f^{i,(1)}(\rho_0), \quad s \in (0,1],$$

with initial value $V(0) = \partial_{v_k} \rho_0$, and $\hat{h}(t, v) := -L_{\rho_0} \hat{V}(t, v), t \in [0, 1]$, where L_{ρ_0} is the same as in the proof of Lemma 6.3. Then, $\hat{V}(t)$ satisfies the equation

$$\partial_t \hat{V} = -L_{\rho_0} \hat{V} + \partial_{v_i} f^{i,(1)}(\rho_0) = L_{\rho_0} \hat{V} + 2\hat{h} + \partial_{v_i} f^{i,(1)}(\rho_0)$$

Note that $h(t) := \hat{h}(1-t) = -L_{\rho_0}V(t)$ satisfies the equation

$$\partial_t h(t) = L_{\rho_0} h(t) + L_{\rho_0} \partial_{v_i} f^{i,(1)}(\rho_0)$$

and $h(0) = -L_{\rho_0} \partial_{v_k} \rho_0$. Thus, by applying Duhamel's formula

$$h(t) = e^{tL_{\rho_0}} h(0) + \int_0^t e^{(t-s)L_{\rho_0}} L_{\rho_0} \partial_{v_i} f^{i,(1)}(\rho_0) ds$$

with the semigroup $e^{tL_{\rho_0}}$ generated by L_{ρ_0} , which is a contraction under the sup-norm by the maximum principle, we see

$$\|h(t)\|_{\infty} \le \|L_{\rho_0}\partial_{v_k}\rho_0\|_{\infty} + \|L_{\rho_0}\partial_{v_i}f^{i,(1)}(\rho_0)\|_{\infty} < \infty, \quad t \in [0,1],$$

since $\rho_0 \in C^4(\mathbb{T}^d)$ by the condition (1.12). So $G = \|\hat{g}\|_{\infty} \leq C$.

This shows (6.14) by applying (6.12) for the extended solution defined on [0, T+1].

These two lemmas also imply an estimate for $\partial_t \rho$. Indeed, expanding the right-hand side of (1.11) by noting $D_{ij}(\rho) \in C^{\infty}([0,1])$, and then applying (6.9) and (6.14), we obtain

(6.16)
$$\|\partial_t \rho\|_{\infty} \le C K^{2(1+\sigma)/\sigma+1}$$

Third, we take the derivative of (6.11) in v_{ℓ} and obtain

(6.17)
$$\partial_t \rho_{v_k v_\ell} = \partial_{v_i} \left\{ D_{ij}(\rho) \partial_{v_j} \rho_{v_k v_\ell} \right\} - \partial_{v_i} f^{i,(2)}(\rho) + K f'(\rho) \rho_{v_k v_\ell} + K f''(\rho) \rho_{v_k} \rho_{v_\ell},$$

where

$$f^{i,(2)}(\rho) = -\sum_{j=1}^{d} \left\{ D'_{ij}(\rho)\rho_{v_{\ell}}\rho_{v_{j}v_{k}} + D''_{ij}(\rho)\rho_{v_{\ell}}\rho_{v_{k}}\rho_{v_{j}} + D'_{ij}(\rho)\partial_{v_{\ell}}(\rho_{v_{k}}\rho_{v_{j}}) \right\}$$

We consider this equation to be a linear equation (4.22) for $u = \rho_{v_k v_\ell}$ with $a^{ij} = D_{ij}(\rho)$, $b^i = c^i = 0$, $f^i = f^{i,(2)}$ (regarding all terms in $f^{i,(2)}$ already given), $c^0 = Kf'(\rho)$ and $g = -Kf''(\rho)\rho_{v_k}\rho_{v_\ell}$ (with all terms already given). Then, one can take A = CK, $F = CK^2(K^{(1+\sigma)/\sigma})^3$ by using (6.4), (6.9) and (6.14), $c_2 = CK$ and $G = CK \cdot K^{2(1+\sigma)/\sigma}$. Thus, by (6.6), we obtain the estimate:

$$|u|_{1+\sigma}^* \le C(M ||u||_{\infty} + K^{3(1+\sigma)/\sigma+2})$$

with $M = [1 + A]^{(1+\sigma)/\sigma} + K$. Therefore, noting $||u||_{\infty} = ||\rho_{v_k v_\ell}||_{\infty} \leq C K^{2(1+\sigma)/\sigma+1}$ by (6.14), we have

(6.18)
$$|\partial^2 \rho|_{1+\sigma}^* \le C K^{3(1+\sigma)/\sigma+2}.$$

We can again use the same argument as above to remove the singularity near t = 0.

Lemma 6.5. For the norm without singularity, we have

$$(6.19) \qquad \qquad |\partial^2 \rho|_{1+\sigma} \le C K^{3(1+\sigma)/\sigma+2}$$

Proof. Take $\{\hat{a}^{ij}(t,v)\}$ as in (6.10), $\hat{c}^0(t,v)$ as in (6.15) and $\hat{b}^i = \hat{c}^i = 0$. Then, for each k, ℓ , we define $\hat{u}(t,v)$, $\{\hat{f}^i(t,v)\}$ and $\hat{g}(t,v)$, respectively, by

$$\hat{u}(t,v) = \begin{cases} \rho_{v_k v_\ell}(t-1,v), & t \in [1,T+1], \\ \hat{V}(t,v), & t \in [0,1). \end{cases}$$
(6.20)
$$\hat{f}^i(t,v) = \begin{cases} f^{i,(2)}(\rho(t-1,v)), & t \in [1,T+1], \\ f^{i,(2)}(\rho_0(v)), & t \in [0,1), \end{cases}$$

$$\hat{g}(t,v) = \begin{cases} Kf''(\rho(t-1,v))\rho_{v_k}(t-1,v)\rho_{v_\ell}(t-1,v), & t \in [1,T+1], \\ -2\hat{h}(t,v), & t \in [0,1), \end{cases}$$

where $\hat{V}(t) \equiv \hat{V}_{k,\ell}(t) := V(1-t)$ is defined from the solution $V(s), s \in [0,1]$ of the equation

$$\partial_s V = L_{\rho_0} V - \partial_{v_i} f^{i,(2)}(\rho_0), \quad s \in (0,1],$$

with initial value $V(0) = \partial_{v_k} \partial_{v_\ell} \rho_0$, and $\hat{h}(t, v) := -L_{\rho_0} \hat{V}(t, v), t \in [0, 1]$. Then, $\hat{V}(t)$ satisfies the equation

$$\partial_t \hat{V} = -L_{\rho_0} \hat{V} + \partial_{v_i} f^{i,(2)}(\rho_0) = L_{\rho_0} \hat{V} + 2\hat{h} + \partial_{v_i} f^{i,(2)}(\rho_0).$$

Note that $h(t) := \hat{h}(1-t) = -L_{\rho_0}V(t,v)$ satisfies the equation

$$\partial_t h(t) = L_{\rho_0} h(t) + L_{\rho_0} \partial_{v_i} f^{i,(2)}(\rho_0)$$

and $h(0) = -L_{\rho_0} \partial_{v_k} \partial_{v_\ell} \rho_0$. Thus, by Duhamel's formula similarly as before, we see

$$\|h(t)\|_{\infty} \le \|L_{\rho_0}\partial_{v_k}\partial_{v_\ell}\rho_0\|_{\infty} + \|L_{\rho_0}\partial_{v_i}f^{i,(2)}(\rho_0)\|_{\infty} < \infty,$$

since $\rho_0 \in C^5(\mathbb{T}^d)$ by the condition (1.12). Thus, $\|\hat{g}\|_{\infty}$ has the same bound by G stated above. This shows (6.19) by applying (6.18) for the extended solution.

Finally for $\partial_{v_k} \partial_t \rho$, using (6.11), expanding the terms on the right-hand side and recalling $D(\rho), f(\rho) \in C^{\infty}([0,1])$, we obtain from (6.9), (6.14) and (6.19)

(6.21)
$$\|\partial_{v_k}\partial_t\rho\|_{\infty} \le CK^{3(1+\sigma)/\sigma+2}$$

The estimate (2.3) follows from (6.9), (6.14), (6.16), (6.19), and (6.21).

7 Comparison theorem

The following lemma is standard and follows directly from the maximum principle. Recall that f(0) > 0 and f(1) < 0 for f in the PDE (1.11) and K > 0.

Lemma 7.1. For the solution $\rho(t, v)$ of (1.11), if $0 < \rho_{-} \le \rho_{0}(v) \le \rho_{+} < 1$, then we have $\rho_{-} \land \alpha_{-} \le \rho(t, v) \le \rho_{+} \lor \alpha_{+}$ for all $t \ge 0$, where $\alpha_{-} = \min\{\alpha \in [0, 1]; f(\alpha) = 0\} > 0$ and $\alpha_{+} = \max\{\alpha \in [0, 1]; f(\alpha) = 0\} < 1$.

Proof. To show the upper bound, set $w(t, v) := \rho_+ \vee \alpha_+ + \varepsilon - \rho(t, v)$ for $\varepsilon > 0$ small enough such that $\rho_+ \vee \alpha_+ + \varepsilon < 1$. Noting w(0, v) > 0, assume that there exist $t_0 > 0$ and $v_0 \in \mathbb{T}^d$ such that w(t, v) > 0 for $t \in [0, t_0)$ and $w(t_0, v_0) = 0$. Then, we have

(7.1)
$$\partial_t w(t_0, v_0) - \sum_{i,j=1}^d \partial_{v_i} \{ D_{ij}(\rho) \partial_{v_j} w \} (t_0, v_0) \\ = -K f(\rho(t_0, v_0)) = -K f(\rho_+ \lor \alpha_+ + \varepsilon) > 0.$$

However, we see that $\partial_t w(t_0, v_0) \leq 0$ and, since $w(t_0, \cdot)$ takes the minimum value 0 at $v = v_0$, recalling (1.8) and (1.9) for $D(\rho)$, similar to [5], p. 345, p. 390, we have

$$\sum_{i,j=1}^{d} \partial_{v_i} \left\{ D_{ij}(\rho) \partial_{v_j} w \right\} (t_0, v_0) \ge 0.$$

This contradicts with (7.1) and therefore we have that $\rho(t, v) < \rho_+ \lor \alpha_+ + \varepsilon$ for all $\varepsilon > 0$ and $t \ge 0$. Thus the upper bound is shown. The proof for the lower bound is similar. \Box

8 Non-degeneracy and boundedness of diffusion matrix

This problem was studied in [26] in more general setting reversible under the Gibbs measures. Nevertheless, we give the proof, in particular that of the lower bound, since it turns out to be explicit in the setting reversible under the Bernoulli measures.

Lemma 8.1. The bound (1.9) holds for $(\theta, D(\rho)\theta)$.

Proof. The upper bound $(\theta, D(\rho)\theta) \leq c^*|\theta|^2$ is easily obtained by taking F = 0; see [26]. To show the lower bound, noting $\pi_{0,x}(\eta_x - \eta_0) = -2(\eta_x - \eta_0)$ for x : |x| = 1, we have

$$\left\langle (\eta_x - \eta_0) \pi_{0,x} \left(\sum_y \tau_y \, \theta \cdot F \right) \right\rangle_{\rho} = -2 \sum_y \langle (\eta_x - \eta_0) \tau_y \, \theta \cdot F \rangle_{\rho}$$
$$= -2 \sum_y \langle (\eta_{x+y} - \eta_y) \, \theta \cdot F \rangle_{\rho} = 0.$$

Therefore,

$$\begin{aligned} |\theta \cdot x| \langle (\eta_x - \eta_0)^2 \rangle_\rho &= \left| \left\langle (\eta_x - \eta_0) \left(\theta \cdot x(\eta_x - \eta_0) - \pi_{0,x} \left(\sum_y \tau_y \, \theta \cdot F \right) \right) \right\rangle_\rho \right| \\ &\leq \langle (\eta_x - \eta_0)^2 \rangle_\rho^{1/2} \left\langle \frac{c_{0,x}}{c_*} \left(\theta \cdot x(\eta_x - \eta_0) - \pi_{0,x} \left(\sum_y \tau_y \, \theta \cdot F \right) \right)^2 \right\rangle_\rho^{1/2}. \end{aligned}$$

Then, taking the square and summing in x : |x| = 1 in both sides, we obtain

$$(2\chi(\rho))^2 \sum_{|x|=1} |\theta \cdot x|^2 \le \frac{4\chi(\rho)}{c_*} (\theta, \widehat{c}(\rho; F)\theta),$$

for every $F \in \mathcal{F}_0^d$. This implies the lower bound $(\theta, D(\rho)\theta) \ge c_*|\theta|^2$.

Acknowledgements

The author thanks Chenlin Gu for helpful discussions, especially for his deep insights from the quantitative homogenization and its extension to the hydrodynamic limit. This work was supported in part by the International Scientists Project of BJNSF, No. IS23007.

References

- C. BERNARDIN, Regularity of the diffusion coefficient for lattice gas reversible under Bernoulli measures, Stoch. Proc. Appl., 101 (2002), 43–68.
- [2] N. CANCRINI AND F. MARTINELLI, Comparison of finite volume canonical and grand canonical Gibbs measures under a mixing condition, Markov Process. Related Fields, 6 (2000), 23–72.

- [3] P. EL KETTANI, T. FUNAKI, D. HILHORST, H. PARK AND S. SETHURAMAN, Mean curvature interface limit from Glauber+Zero-range interacting particles, Comm. Math. Phys., **394** (2022), 1173–1223.
- [4] P. EL KETTANI, T. FUNAKI, D. HILHORST, H. PARK AND S. SETHURAMAN, Singular limit of an Allen-Cahn equation with nonlinear diffusion, Tunisian J. Math., 4 (2022), 719–754.
- [5] L.C. EVANS, Partial differential equations, Graduate Studies in Mathematics, 19, American Mathematical Society, Providence, 1998. xviii+662 pp.
- [6] T. FUNAKI, Equilibrium fluctuations for lattice gas, Itô's Stochastic Calculus and Probability Theory, edited by Ikeda, Watanabe, Fukushima & Kunita, Springer, 1996, 63–72.
- [7] T. FUNAKI, Lectures on Random Interfaces, SpringerBriefs in Probability and Mathematical Statistics, 2016, xii+138 pp.
- [8] T. FUNAKI, Hydrodynamic limit for exclusion processes, Comm. Math. Statis., 6 (2018), 417–480.
- [9] T. FUNAKI, C. GU AND H. WANG, Quantitative homogenization and hydrodynamic limit of non-gradient exclusion process, arXiv:2404.12234.
- [10] T. FUNAKI, P. VAN MEURS, S. SETHURAMAN AND K. TSUNODA, Motion by mean curvature from Glauber-Kawasaki dynamics with speed change, J. Stat. Phys., 190 (2023), Article no. 45, 1–30.
- [11] T. FUNAKI, P. VAN MEURS, S. SETHURAMAN AND K. TSUNODA, Constant-speed interface flow from unbalanced Glauber-Kawasaki dynamics, Ensaios Matemáticos, 38 (2023), 223–248.
- [12] T. FUNAKI AND H. PARK, Motion of sharp interface of Allen-Cahn equation with anisotropic nonlinear diffusion, arXiv:2403.01732.
- [13] T. FUNAKI AND S. SETHURAMAN, Schauder estimate for quasilinear discrete PDEs of parabolic type, arXiv:2112.13973.
- [14] T. FUNAKI, K. UCHIYAMA AND H.-T. YAU, Hydrodynamic limit for lattice gas reversible under Bernoulli measures, Nonlinear Stochastic PDEs: Hydrodynamic Limit and Burgers' Turbulence, edited by T. Funaki & W.A. Woyczynski, IMA volume 77, Springer, 1995, 1–40.
- [15] H.O. GEORGII, Canonical Gibbs Measures, Lecture Notes in Math., 760, 1979, Springer
- [16] R.J. GLAUBER, Time-dependent statistics of the Ising model, J. Math. Phys., 4 (1963), 294–307.
- [17] M.A. KATSOULAKIS AND P.E. SOUGANIDIS, Interacting particle systems and generalized evolution of fronts, Arch. Rational Mech. Anal., 127 (1994), 133–157.

- [18] K. KAWASAKI, Diffusion constants near the critical point for time-dependent Ising models. I, Phys. Rev., 145 (1966), 224–230.
- [19] C. KIPNIS, C. LANDIM AND S. OLLA, Hydrodynamical limit for a nongradient system: the generalized symmetric exclusion process, Comm. Pure Appl. Math., 47 (1994), 1475–1545.
- [20] O.A. LADYŽENSKAJA, V.A. SOLONNIKOV AND N.N. URAL'CEVA, Linear and Quasilinear Equations of Parabolic Type, Translations Math. Monographs, 23, American Mathematical Society, 1968.
- [21] G.M. LIEBERMAN, Second Order Parabolic Differential Equations, World Scientific, 1996, 452 pp.
- [22] S.L. LU AND H.-T. YAU, Spectral gap and logarithmic Sobolev inequality for Kawasaki and Glauber dynamics, Comm. Math. Phys., 156 (1993), 399–433.
- [23] S. OLLA, S.R.S. VARADHAN AND H.-T. YAU, Hydrodynamical limit for a Hamiltonian system with weak noise, Comm. Math. Phys., 155 (1993), 523–560.
- [24] Z. QIAN AND G. XI, Parabolic equations with divergence-free drift in space $L_t^{\ell} L_x^q$, Indiana Univ. Math. J., **68** (2019), 761–797.
- [25] H. SPOHN, Large Scale Dynamics of Interacting Particles, Springer, 1991.
- [26] H. SPOHN AND H.T. YAU, Bulk diffusivity of lattice gases close to criticality, J. Stat. Phys., 79 (1995), 231–241.
- [27] D. STROOCK, Diffusion semigroups corresponding to uniformly elliptic divergence form operators, Séminaire de Probabilités, XXII, 316–347, Lecture Notes in Math., 1321, Springer, 1988.
- [28] S.R.S. VARADHAN, Nonlinear diffusion limit for a system with nearest neighbor interactions - II In: Asymptotic problems in probability theory: stochastic models and diffusions on fractals (eds. K.D. Elworthy and N. Ikeda), Longman, 1993, 75–128.
- [29] S.R.S. VARADHAN AND H.T. YAU, Diffusive limit of lattice gas with mixing conditions, Asian J. Math., 1 (1997), 623–678.
- [30] H.-T. YAU, Relative entropy and hydrodynamics of Ginzburg-Landau models, Letters Math. Phys., 22 (1991), 63–80.