

On the duality in constant-roll inflation

Yue Wang,^{1,*} Qing Gao,^{2,†} Shengqing Gao,^{1,‡} and Yungui Gong^{3,1,§}

¹*School of Physics, Huazhong University of Science and Technology, Wuhan, Hubei 430074, China*

²*School of Physical Science and Technology,*

Southwest University, Chongqing 400715, China

³*Department of Physics, School of Physical Science and Technology,*

Ningbo University, Ningbo, Zhejiang 315211, China

There is a duality in the observables n_s , r and the inflaton potential between large and small η_H for the constant-roll inflation if the slow-roll parameter ϵ_H is negligible. In general, the duality between η_H and $\bar{\eta}_H$ does not hold for the background evolution of the inflaton. For some particular solutions for the constant-roll inflation with η_H being a constant, we find that in the small field approximation, the potential takes the quadratic form and it remains the same when the parameter η_H changes to $\bar{\eta}_H = 3 - \eta_H$. If the scalar field is small and the contribution of ϵ_H is negligible, we find that there exists the logarithmic duality and the duality between large and small η_H for the primordial curvature perturbation in inflationary models with the quadratic potential.

I. INTRODUCTION

As more evidences for the existence of primordial black holes (PBHs) were provided by the observations of gravitational waves (GWs) such as the Laser Interferometer Gravitational Wave Observatory (LIGO) Scientific and Virgo Collaborations [1–6], and the Pulsar Timing Arrays (PTAs) [7–10], PBHs as dark matter attracted a lot of attention. When the density contrast of overdense regions exceeds the threshold value at the horizon reentry during radiation domination, PBHs may form through gravitational collapse [11, 12]. The large density contrast of overdense regions may be seeded from large primordial curvature perturbations at small scales generated during inflation. The large curvature perturbations are

* ywang123@hust.edu.cn

† Corresponding author. gaoqing1024@swu.edu.cn

‡ gaoshengqing@hust.edu.cn

§ gongyungui@nbu.edu.cn

also the sources of secondary gravitational waves (GWs) after the horizon reentry through the scalar-tensor mixing [13–16]. Therefore, accompanied with the production of PBHs, scalar induced gravitational waves (SIGWs) are generated [3, 13–53].

To produce large curvature perturbations at small scales during inflation, we usually consider inflationary models with an inflection point in the inflaton potential [54–57]. Near the inflection point, the slow-roll parameter $\eta_H \approx 3$ and it is called the ultra-slow-roll (USR) inflation [58, 59]. More general, the constant-roll inflation with η_H being a constant was proposed [60, 61]. If $\eta_H > 1$, then the slow-roll condition is violated in constant-roll inflation and the primordial curvature perturbations may evolve outside the horizon [59–66]. For constant-roll inflation, the inflationary potential and the background equation of motion can be solved analytically. Due to different background evolution for large and small η_H , the observational data constrained η_H to be small [67–70]. Since the slow-roll parameter ϵ_H decreases with time, so it may be negligible during inflation. By neglecting the contribution from ϵ_H , it was found that there exists a duality between η_H and $\bar{\eta}_H = 3 - \eta_H$ in the observables the scalar spectral tilt n_s and the tensor-to-scalar ratio r [71, 72]. The duality between η_H and $\bar{\eta}_H = 3 - \eta_H$ connects the constant slow-roll inflation with $|\eta_H| \ll 1$ and the USR inflation with $\eta_H \approx 3$, so the behavior of the primordial curvature perturbations in USR inflation can be understood from the usual slow-roll inflation. Recently, the logarithmic duality of the primordial curvature perturbation was found for the quadratic potential in Ref. [73]. Since the logarithmic duality applies to both the slow-roll and USR inflation, we extend the discussion on the duality to constant-roll inflation in this paper.

The paper is organized as follows. In Section II, we discuss the duality in constant-roll inflation. Motivated by the duality in constant-roll inflation, we then discuss the logarithmic duality of the primordial curvature perturbation using the δN formalism in Section III. The conclusion is drawn in Section IV.

II. THE CONSTANT-ROLL INFLATION

For the constant-roll inflation, we take the second Hubble flow slow-roll parameter

$$\eta_H = \frac{2}{H} \frac{d^2 H}{d\phi^2} = -\frac{\ddot{H}}{2H\dot{H}} = -\frac{\ddot{\phi}}{H\dot{\phi}} \quad (1)$$

to be a constant, i.e., we take $\eta_H = 3 + \alpha$ with α being a constant. If $\alpha \approx -3$, then η_H is small and the constant-roll inflation is also a slow-roll inflation; If $\alpha \approx 0$, then $\eta_H \approx 3$ and $dV/d\phi \approx 0$, the constant-roll inflation is USR inflation. For the constant-roll inflation with a constant η_H , the scalar spectral tilt is [61, 66]

$$n_s - 1 \approx 3 - |2\eta_H - 3| - \frac{2(2\eta_H^2 - 9\eta_H + 6)}{|2\eta_H - 3|} \epsilon_H, \quad (2)$$

and the tensor-to-scalar ratio is [61, 66]

$$r \approx 2^{7-|2\eta_H-3|} \left(\frac{\Gamma[3/2]}{\Gamma[|2\eta_H-3|/2]} \right)^2 \epsilon_H, \quad (3)$$

where $\epsilon_H = 2(dH/d\phi)^2/H^2 = \dot{\phi}^2/(2H^2)$ is the first Hubble flow slow-roll parameter. If the contribution of ϵ_H is negligible in Eq. (2), then the scalar spectral tilt remains unchanged with the replacement of η_H by $\bar{\eta}_H = 3 - \eta_H$. It is interesting to note that the tensor-to-scalar ratio also keeps the same under the interchange between η_H and $\bar{\eta}_H$. The behaviours of n_s and r under the swap between η_H and $\bar{\eta}_H = -\alpha$ in the constant-roll inflation are called the duality between large and small η_H in the observables n_s and r [71, 72]. The duality connects the slow-roll inflation with $|\eta_H| \ll 1$ and the USR inflation with $\eta_H \approx 3$, so it is useful for the understanding of PBH formation and SIGW generation by USR inflation from slow-roll inflation. If the contribution of ϵ_H in Eq. (2) is not negligible, then the duality in n_s does not exist, and the background evolutions are also very different [70]. Therefore, it is important to further explore the duality between η_H and $\bar{\eta}_H$.

To see when ϵ_H can be neglected, substituting $\eta_H = 3 + \alpha$ into Eq. (2), we get

$$n_s - 1 \approx -2\alpha - \frac{2(2\alpha^2 + 3\alpha - 3)}{3 + 2\alpha} \epsilon_H, \quad (4)$$

where $\alpha > -3/2$. Therefore, the condition that the contribution of ϵ_H is negligible is

$$\epsilon_H \ll \left| \frac{\alpha(3 + 2\alpha)}{2\alpha^2 + 3\alpha - 3} \right|. \quad (5)$$

For the USR inflation, $|\alpha| \ll 1$, the condition (5) means that $\epsilon_H \ll |\alpha| \ll 1$. Substituting $\bar{\eta}_H = -\alpha$ into Eq. (2), we get

$$n_s - 1 \approx -2\alpha - \frac{2(2\alpha^2 - 9\alpha + 6)}{3 + 2\alpha} \epsilon_H. \quad (6)$$

For the case with $\bar{\eta}_H$, the condition that the contribution of ϵ_H is negligible is

$$\epsilon_H \ll \left| \frac{\alpha(3 + 2\alpha)}{2\alpha^2 - 9\alpha + 6} \right|. \quad (7)$$

For slow-roll constant inflation with $|\alpha| \ll 1$, the condition (7) becomes $\epsilon_H \ll |\alpha|/2 \ll 1$. Combining the results (5) and (7), we see that the duality between the USR inflation with $\eta_H = 3 + \alpha$ and the slow-roll constant inflation with $\bar{\eta}_H = -\alpha$ holds when $\epsilon_H \ll |\alpha|/2 \ll 1$.

Now we discuss the background evolution of the constant-roll inflation with negligible ϵ_H . From the background equation

$$3H^2 = \frac{1}{2}\dot{\phi}^2 + V(\phi), \quad (8)$$

we get

$$V(\phi) = (3 - \epsilon_H)H^2, \quad (9)$$

and

$$\frac{dV}{d\phi} = (3 - \eta_H)H^2 \frac{d\phi}{dN}. \quad (10)$$

In terms of the number of e-folds N , $dN = -Hdt$, the equation of motion for the inflaton becomes

$$\frac{d^2\phi}{dN^2} + (\epsilon_H - 3)\frac{d\phi}{dN} + \frac{1}{H^2} \frac{dV}{d\phi} = 0. \quad (11)$$

Combining Eqs. (10) and (11), we get

$$\frac{d^2\phi}{dN^2} + (\epsilon_H - \eta_H)\frac{d\phi}{dN} = 0. \quad (12)$$

Eq. (12) is just the definition (1) of η_H . If ϵ_H is negligible, then Eqs. (9), (10) and (12) become

$$V(\phi) \approx 3H^2(\phi), \quad (13)$$

$$\frac{dV}{d\phi} \approx \frac{3 - \eta_H}{3}V(\phi)\frac{d\phi}{dN}, \quad (14)$$

$$\frac{d^2\phi}{dN^2} - \eta_H \frac{d\phi}{dN} \approx 0. \quad (15)$$

The solution to Eq. (15) is

$$\phi(N) \approx Ae^{\eta_H N} + B, \quad (16)$$

where A and B are integration constants. Substituting the solution (16) into Eq. (14), we get

$$\frac{dV}{V} \approx \frac{1}{3}\eta_H(3 - \eta_H)(\phi - B)d\phi. \quad (17)$$

So the potential is

$$V(\phi) \approx V_0 \exp \left[\frac{1}{3}\eta_H(3 - \eta_H) \left(\frac{1}{2}\phi^2 - B\phi \right) \right], \quad (18)$$

and $H^2 \approx V(\phi)/3$, where V_0 is an integration constant. Therefore, if ϵ_H is negligible, then the potential (18) and the Hubble parameter $H(\phi)$ are unchanged under the transformation from η_H to $\bar{\eta}_H = 3 - \eta_H$, i.e., both $H(\phi)$ and $V(\phi)$ have the duality between η_H and $\bar{\eta}_H$.

III. LOGARITHMIC DUALITY OF PRIMORDIAL CURVATURE PERTURBATION

From the definition (1) of η_H , we get

$$\frac{d^2 H}{d\phi^2} - \frac{\eta_H}{2} H = 0. \quad (19)$$

The solution to Eq. (19) is

$$H(\phi) = C_1 \exp\left(\sqrt{\frac{\eta_H}{2}} \phi\right) + C_2 \exp\left(-\sqrt{\frac{\eta_H}{2}} \phi\right), \quad (20)$$

for $\eta_H > 0$, and

$$H(\phi) = C_1 \cos\left(\sqrt{-\frac{\eta_H}{2}} \phi\right) + C_2 \sin\left(\sqrt{-\frac{\eta_H}{2}} \phi\right), \quad (21)$$

for $\eta_H < 0$, where C_1 and C_2 are integration constants. Substituting the solutions (20) and (21) into Eq. (9), we can obtain the potential $V(\phi)$. Therefore, the potential and the background evolution of the inflaton are determined by the parameter η_H only for the constant-roll inflation with the slow-roll parameter η_H being a constant [61].

For the particular solution

$$H(\phi) = M \sinh\left(\sqrt{\frac{\eta_H}{2}} \phi\right), \quad (22)$$

the potential is

$$V(\phi) = M^2 \left[3 \sinh^2\left(\sqrt{\frac{\eta_H}{2}} \phi\right) - \eta_H \cosh^2\left(\sqrt{\frac{\eta_H}{2}} \phi\right) \right]. \quad (23)$$

In the small field approximation with $|\phi| \ll 1$, we get

$$V(\phi) \approx M^2 \left[-\eta_H + \eta_H(3 - \eta_H) \frac{\phi^2}{2} \right]. \quad (24)$$

Because of the presence of the first term $-\eta_H$ in the right hand side of Eq. (24), the duality between $\eta_H < 3$ and $\bar{\eta}_H = 3 - \eta_H$ does not hold in both $H(\phi)$ and $V(\phi)$ for the solutions (22) and (23). When $\eta_H > 3$, the solution for $\bar{\eta}_H = 3 - \eta_H < 0$ is

$$H(\phi) = M \sin\left(\sqrt{-\frac{\bar{\eta}_H}{2}} \phi\right), \quad (25)$$

and the potential is

$$V(\phi) = M^2 \left[3 \sin^2 \left(\sqrt{-\frac{\bar{\eta}_H}{2}} \phi \right) + \bar{\eta}_H \cos^2 \left(\sqrt{-\frac{\bar{\eta}_H}{2}} \phi \right) \right]. \quad (26)$$

In the small field approximation with $|\phi| \ll 1$, we get

$$V(\phi) \approx M^2 \left[\bar{\eta}_H - \bar{\eta}_H(3 - \bar{\eta}_H) \frac{\phi^2}{2} \right]. \quad (27)$$

Due to the presence of the first term $\bar{\eta}_H$ in the right hand side of Eq. (27), the duality between $\eta_H > 3$ and $\bar{\eta}_H = 3 - \eta_H$ does not hold for the solutions (25) and (27) either even in the small field approximation.

For the particular solution

$$H(\phi) = M \cosh \left(\sqrt{\frac{\eta_H}{2}} \phi \right), \quad (28)$$

the potential is

$$V(\phi) = M^2 \left[3 \cosh^2 \left(\sqrt{\frac{\eta_H}{2}} \phi \right) - \eta_H \sinh^2 \left(\sqrt{\frac{\eta_H}{2}} \phi \right) \right]. \quad (29)$$

In the small field approximation with $|\phi| \ll 1$, we get

$$V(\phi) \approx M^2 \left[3 + \frac{1}{2} \eta_H (3 - \eta_H) \phi^2 \right]. \quad (30)$$

It is interesting to note that in the small field approximation with $|\phi| \ll 1$, the potential (30) keeps to be the same if we change η_H ($0 < \eta_H < 3$) to $\bar{\eta}_H = 3 - \eta_H$, i.e., there is a duality between η_H and $\bar{\eta}_H$ in the potential (30), but the duality does not exist in $H(\phi)$. If $\eta_H > 3$, then $\bar{\eta}_H = 3 - \eta_H < 0$, the solution for $\bar{\eta}_H$ is

$$H(\phi) = M \cos \left(\sqrt{-\frac{\bar{\eta}_H}{2}} \phi \right), \quad (31)$$

and the potential is

$$V(\phi) = M^2 \left[3 \cos^2 \left(\sqrt{-\frac{\bar{\eta}_H}{2}} \phi \right) + \bar{\eta}_H \sin^2 \left(\sqrt{-\frac{\bar{\eta}_H}{2}} \phi \right) \right]. \quad (32)$$

In the small field approximation with $|\phi| \ll 1$, we get

$$V(\phi) \approx M^2 \left[3 + \frac{1}{2} \bar{\eta}_H (3 - \bar{\eta}_H) \phi^2 \right]. \quad (33)$$

Therefore, in the small field approximation with $|\phi| \ll 1$, the potential (33) has the same form as (30) except that here $\bar{\eta}_H < 0$, and it seems that there exists the duality between η_H and $\bar{\eta}_H = 3 - \eta_H$ in the potential (33). When $\bar{\eta}_H < 0$, $\eta_H = 3 - \bar{\eta}_H > 0$, actually the potential (32) with $\bar{\eta}_H < 0$ should be replaced by the potential (29) with $\eta_H > 0$, so there is no duality between η_H and $\bar{\eta}_H = 3 - \eta_H$ in the potential (32) in the small field approximation. In fact, if we replace $\eta_H = 3 + \alpha$ with $\alpha > 0$ by $\bar{\eta}_H = -\alpha$, the solutions (28) and (29) should be replaced by the solutions (31) and (32), so the potential (30) is dual to the potential (33) under the interchange between $\eta_H > 3$ and $\bar{\eta}_H = 3 - \eta_H < 0$.

Since logarithmic duality of the primordial curvature perturbation was found for the quadratic potential in Ref. [73], we will discuss whether the logarithmic duality exists in the constant-roll inflation for the particular solutions (30) and (33). If $\epsilon_H \ll 1$ and $\phi \ll 1$, then substituting the potential (30) into Eq. (9), we get

$$H^2(\phi) \approx V(\phi)/3 \approx M^2. \quad (34)$$

Plugging the potential (30) into Eq. (11) and neglecting ϵ_H , we get

$$\frac{d^2\phi}{dN^2} - 3\frac{d\phi}{dN} + \eta_H(3 - \eta_H)\phi = 0. \quad (35)$$

The solution is

$$\phi(N) = C_+ e^{\lambda_+(N-N_e)} + C_- e^{\lambda_-(N-N_e)}, \quad (36)$$

where λ_+ and λ_- are

$$\begin{aligned} \lambda_+ &= \frac{3 + \sqrt{9 - 4\eta_H(3 - \eta_H)}}{2}, \\ \lambda_- &= \frac{3 - \sqrt{9 - 4\eta_H(3 - \eta_H)}}{2}, \end{aligned} \quad (37)$$

and N_e is the number of e-folds at the end of inflation. Note that both λ_+ and λ_- are invariant under the interchange between η_H and $\bar{\eta}_H$, and the inflationary model is not constant-roll inflation. Remember that for constant-roll inflation with negligible ϵ_H , the equation of motion for the scalar field satisfies Eq. (15). Combining Eqs. (35) and (15), we get

$$\frac{d\phi}{dN} = \eta_H \phi, \quad (38)$$

$$\frac{d^2\phi}{dN^2} = \eta_H^2 \phi. \quad (39)$$

So the solution (36) is not the solution to constant-roll inflation, but we can still take η_H as a parameter even though it loses the meaning of the second slow-roll parameter as defined in Eq. (1) and think that the quadratic potentials (30) and (33) are motivated from constant-roll inflation. Now back to the solution (36), to determine the coefficients C_+ and C_- , we use the velocity of the scalar field

$$\pi = -\frac{d\phi}{dN} = -C_+\lambda_+e^{\lambda_+(N-N_e)} - C_-\lambda_-e^{\lambda_-(N-N_e)}, \quad (40)$$

to get

$$C_+ = -\frac{\lambda_-\phi_e + \pi_e}{\lambda_+ - \lambda_-}, \quad C_- = \frac{\lambda_+\phi_e + \pi_e}{\lambda_+ - \lambda_-}, \quad (41)$$

where π_e is the value of $\pi(N)$ at N_e . Since λ_+ and λ_- have the duality between η_H and $\bar{\eta}_H$, so C_+ and C_- also have the duality between η_H and $\bar{\eta}_H$. If we swap λ_+ and λ_- , then C_+ and C_- are interchanged, and $\phi(N)$ and $\pi(N)$ are invariant. Therefore

$$\begin{aligned} e^{\lambda_+(N-N_e)} &= \frac{\lambda_-\phi + \pi}{\lambda_-\phi_e + \pi_e}, \\ e^{\lambda_-(N-N_e)} &= \frac{\lambda_+\phi + \pi}{\lambda_+\phi_e + \pi_e}, \end{aligned} \quad (42)$$

and

$$\begin{aligned} N - N_e &= \frac{1}{\lambda_+} \ln \left(\frac{\lambda_-\phi + \pi}{\lambda_-\phi_e + \pi_e} \right) \\ &= \frac{1}{\lambda_-} \ln \left(\frac{\lambda_+\phi + \pi}{\lambda_+\phi_e + \pi_e} \right). \end{aligned} \quad (43)$$

Using the δN formalism [74–77], we get the primordial curvature perturbation

$$\begin{aligned} \zeta &= \delta(N - N_e) \\ &= \frac{1}{\lambda_+} \ln \left(1 + \frac{\lambda_-\delta\phi + \delta\pi}{\lambda_-\phi + \pi} \right) - \frac{1}{\lambda_+} \ln \left(1 + \frac{\delta\pi_e}{\lambda_-\phi_e + \pi_e} \right) \\ &= \frac{1}{\lambda_-} \ln \left(1 + \frac{\lambda_+\delta\phi + \delta\pi}{\lambda_+\phi + \pi} \right) - \frac{1}{\lambda_-} \ln \left(1 + \frac{\delta\pi_e}{\lambda_+\phi_e + \pi_e} \right). \end{aligned} \quad (44)$$

It can be seen that under the interchange between λ_+ and λ_- , the two formulae for the primordial curvature perturbation are equivalent and the equivalence is guaranteed by taking the perturbation on Eq. (42),

$$\left(1 + \frac{\delta\pi_e}{\pi_e + \lambda_+\phi_e} \right)^{-\lambda_+} \left(1 + \frac{\delta\pi_e}{\pi_e + \lambda_-\phi_e} \right)^{\lambda_-} = \left(1 + \frac{\delta\pi + \lambda_-\delta\phi}{\pi + \lambda_-\phi} \right)^{\lambda_-} \left(1 + \frac{\delta\pi + \lambda_+\delta\phi}{\pi + \lambda_+\phi} \right)^{-\lambda_+}. \quad (45)$$

The formula (45) connects $\delta\pi_e$ with $\delta\phi$ and $\delta\pi$. In the same way, we can get the same result for the potential (33). The result is the same as that found in [73] and is called logarithmic duality. Therefore, there exists the logarithmic duality and the duality between large and small η_H for the primordial curvature perturbation in inflationary models with the potential (30) and (33).

IV. CONCLUSION

PBHs and accompanying SIGWs are usually generated in inflationary models with inflection point. Near the inflection point, the slow-roll parameter $\eta_H \approx 3$ and the inflation is also USR inflation which is a special case of the constant-roll inflation. For the constant-roll inflation, it is interesting that there is a duality between large and small η_H in the observables n_s and r if the slow-roll parameter ϵ_H is negligible. The duality between η_H and $\bar{\eta}_H = 3 - \eta_H$ connects the constant slow-roll inflation with $|\eta_H| \ll 1$ and the USR inflation with $\eta_H \approx 3$, so the behavior of primordial curvature perturbations in USR inflation can be understood from the usual slow-roll inflation. However, the duality does not hold for the background evolution of the inflaton in general. If we neglect the contribution of ϵ_H in the background evolution of the inflaton, we find that the inflaton potential has the duality between η_H and $\bar{\eta}_H$. We also derive the condition for neglecting ϵ_H . In some particular solutions of constant-roll inflation and in the small field approximation, the inflaton potential takes a quadratic form and it remains the same under the interchange of the parameter η_H and $\bar{\eta}_H$. When $0 < \eta_H < 3$, in the small field approximation, there is a duality between η_H and $\bar{\eta}_H$ in the potential (30), but there does not exist the duality in $H(\phi)$. When $\eta_H > 3$ and $\bar{\eta}_H = 3 - \eta_H < 0$, if we replace η_H by $\bar{\eta}_H$, then the potential (30) for $\eta_H > 3$ should be replaced by the potential (33) for $\bar{\eta}_H < 0$. Even in the small field approximation, both the potentials (30) and (33) take the same quadratic form and are invariant under the interchange between η_H and $\bar{\eta}_H$, there does not exist the duality between η_H and $\bar{\eta}_H$ in the potentials (30) and (33).

If we start with the quadratic potential (30) and (33) without the restriction on constant-roll inflation, then using the δN formalism we find that the primordial curvature perturbation exhibits a logarithmic duality if we neglect the contribution of ϵ_H . Since both λ_- and λ_+ are invariant under the interchange between the parameter η_H and $\bar{\eta}_H = 3 - \eta_H$, $\phi(N)$, $\pi(N)$

and the primordial curvature perturbation also exhibit the duality between the parameters η_H and $\bar{\eta}_H$.

ACKNOWLEDGMENTS

This research is supported in part by the National Key Research and Development Program of China under Grant No. 2020YFC2201504, the National Natural Science Foundation of China under Grant No. 12175184 and the Chongqing Natural Science Foundation under Grant No. CSTB2022NSCQ-MSX1324.

-
- [1] S. Bird, I. Cholis, J. B. Muñoz, Y. Ali-Haïmoud, M. Kamionkowski, E. D. Kovetz, A. Racanelli, and A. G. Riess, Did LIGO detect dark matter?, *Phys. Rev. Lett.* **116**, 201301 (2016).
 - [2] M. Sasaki, T. Suyama, T. Tanaka, and S. Yokoyama, Primordial Black Hole Scenario for the Gravitational-Wave Event GW150914, *Phys. Rev. Lett.* **117**, 061101 (2016), [Erratum: *Phys.Rev.Lett.* 121, 059901 (2018)].
 - [3] K. Inomata, M. Kawasaki, K. Mukaida, Y. Tada, and T. T. Yanagida, Inflationary primordial black holes for the LIGO gravitational wave events and pulsar timing array experiments, *Phys. Rev. D* **95**, 123510 (2017).
 - [4] V. De Luca, V. Desjacques, G. Franciolini, P. Pani, and A. Riotto, GW190521 Mass Gap Event and the Primordial Black Hole Scenario, *Phys. Rev. Lett.* **126**, 051101 (2021).
 - [5] V. De Luca, G. Franciolini, P. Pani, and A. Riotto, Bayesian Evidence for Both Astrophysical and Primordial Black Holes: Mapping the GWTC-2 Catalog to Third-Generation Detectors, *J. Cosmol. Astropart. Phys.* **05** (2021) 003.
 - [6] G. Franciolini, V. Baibhav, V. De Luca, K. K. Y. Ng, K. W. K. Wong, E. Berti, P. Pani, A. Riotto, and S. Vitale, Searching for a subpopulation of primordial black holes in LIGO-Virgo gravitational-wave data, *Phys. Rev. D* **105**, 083526 (2022).
 - [7] V. De Luca, G. Franciolini, and A. Riotto, NANOGrav Data Hints at Primordial Black Holes as Dark Matter, *Phys. Rev. Lett.* **126**, 041303 (2021).
 - [8] V. Vaskonen and H. Veermäe, Did NANOGrav see a signal from primordial black hole formation?, *Phys. Rev. Lett.* **126**, 051303 (2021).

- [9] A. Afzal *et al.* (NANOGrav), The NANOGrav 15 yr Data Set: Search for Signals from New Physics, *Astrophys. J. Lett.* **951**, L11 (2023).
- [10] J. Antoniadis *et al.* (EPTA), The second data release from the European Pulsar Timing Array: V. Implications for massive black holes, dark matter and the early Universe, [arXiv:2306.16227](https://arxiv.org/abs/2306.16227).
- [11] B. J. Carr and S. W. Hawking, Black holes in the early Universe, *Mon. Not. R. Astron. Soc.* **168**, 399 (1974).
- [12] S. Hawking, Gravitationally collapsed objects of very low mass, *Mon. Not. R. Astron. Soc.* **152**, 75 (1971).
- [13] S. Matarrese, S. Mollerach, and M. Bruni, Second order perturbations of the Einstein-de Sitter universe, *Phys. Rev. D* **58**, 043504 (1998).
- [14] S. Mollerach, D. Harari, and S. Matarrese, CMB polarization from secondary vector and tensor modes, *Phys. Rev. D* **69**, 063002 (2004).
- [15] K. N. Ananda, C. Clarkson, and D. Wands, The Cosmological gravitational wave background from primordial density perturbations, *Phys. Rev. D* **75**, 123518 (2007).
- [16] D. Baumann, P. J. Steinhardt, K. Takahashi, and K. Ichiki, Gravitational Wave Spectrum Induced by Primordial Scalar Perturbations, *Phys. Rev. D* **76**, 084019 (2007).
- [17] J. Garcia-Bellido, M. Peloso, and C. Unal, Gravitational Wave signatures of inflationary models from Primordial Black Hole Dark Matter, *J. Cosmol. Astropart. Phys.* **09** (2017) 013.
- [18] R. Saito and J. Yokoyama, Gravitational wave background as a probe of the primordial black hole abundance, *Phys. Rev. Lett.* **102**, 161101 (2009), [Erratum: *Phys.Rev.Lett.* 107, 069901 (2011)].
- [19] R. Saito and J. Yokoyama, Gravitational-Wave Constraints on the Abundance of Primordial Black Holes, *Prog. Theor. Phys.* **123**, 867 (2010), [Erratum: *Prog.Theor.Phys.* 126, 351–352 (2011)].
- [20] E. Bugaev and P. Klimai, Induced gravitational wave background and primordial black holes, *Phys. Rev. D* **81**, 023517 (2010).
- [21] E. Bugaev and P. Klimai, Constraints on the induced gravitational wave background from primordial black holes, *Phys. Rev. D* **83**, 083521 (2011).
- [22] L. Alabidi, K. Kohri, M. Sasaki, and Y. Sendouda, Observable Spectra of Induced Gravitational Waves from Inflation, *J. Cosmol. Astropart. Phys.* **09** (2012) 017.
- [23] K. Inomata, K. Kohri, T. Nakama, and T. Terada, Enhancement of Gravitational Waves

- Induced by Scalar Perturbations due to a Sudden Transition from an Early Matter Era to the Radiation Era, *Phys. Rev. D* **100**, 043532 (2019), [Erratum: *Phys.Rev.D* 108, 049901 (2023)].
- [24] M. Braglia, X. Chen, and D. K. Hazra, Probing Primordial Features with the Stochastic Gravitational Wave Background, *J. Cosmol. Astropart. Phys.* **03** (2021) 005.
- [25] Z. Yi, Primordial black holes and scalar-induced gravitational waves from the generalized Brans-Dicke theory, *J. Cosmol. Astropart. Phys.* **03** (2023) 048.
- [26] N. Orlofsky, A. Pierce, and J. D. Wells, Inflationary theory and pulsar timing investigations of primordial black holes and gravitational waves, *Phys. Rev. D* **95**, 063518 (2017).
- [27] T. Nakama, J. Silk, and M. Kamionkowski, Stochastic gravitational waves associated with the formation of primordial black holes, *Phys. Rev. D* **95**, 043511 (2017).
- [28] S.-L. Cheng, W. Lee, and K.-W. Ng, Primordial black holes and associated gravitational waves in axion monodromy inflation, *J. Cosmol. Astropart. Phys.* **07** (2018) 001.
- [29] R.-g. Cai, S. Pi, and M. Sasaki, Gravitational Waves Induced by non-Gaussian Scalar Perturbations, *Phys. Rev. Lett.* **122**, 201101 (2019).
- [30] N. Bartolo, V. De Luca, G. Franciolini, M. Peloso, D. Racco, and A. Riotto, Testing primordial black holes as dark matter with LISA, *Phys. Rev. D* **99**, 103521 (2019).
- [31] N. Bartolo, V. De Luca, G. Franciolini, A. Lewis, M. Peloso, and A. Riotto, Primordial Black Hole Dark Matter: LISA Serendipity, *Phys. Rev. Lett.* **122**, 211301 (2019).
- [32] K. Kohri and T. Terada, Semianalytic calculation of gravitational wave spectrum nonlinearly induced from primordial curvature perturbations, *Phys. Rev. D* **97**, 123532 (2018).
- [33] J. R. Espinosa, D. Racco, and A. Riotto, A Cosmological Signature of the SM Higgs Instability: Gravitational Waves, *J. Cosmol. Astropart. Phys.* **09** (2018) 012.
- [34] R.-G. Cai, S. Pi, S.-J. Wang, and X.-Y. Yang, Resonant multiple peaks in the induced gravitational waves, *J. Cosmol. Astropart. Phys.* **05** (2019) 013.
- [35] R.-G. Cai, S. Pi, S.-J. Wang, and X.-Y. Yang, Pulsar Timing Array Constraints on the Induced Gravitational Waves, *J. Cosmol. Astropart. Phys.* **10** (2019) 059.
- [36] R.-G. Cai, Z.-K. Guo, J. Liu, L. Liu, and X.-Y. Yang, Primordial black holes and gravitational waves from parametric amplification of curvature perturbations, *J. Cosmol. Astropart. Phys.* **06** (2020) 013.
- [37] R.-G. Cai, Y.-C. Ding, X.-Y. Yang, and Y.-F. Zhou, Constraints on a mixed model of dark matter particles and primordial black holes from the galactic 511 keV line,

- J. Cosmol. Astropart. Phys.* **03** (2021) 057.
- [38] G. Domènech, Induced gravitational waves in a general cosmological background, *Int. J. Mod. Phys. D* **29**, 2050028 (2020).
- [39] G. Domènech, S. Pi, and M. Sasaki, Induced gravitational waves as a probe of thermal history of the universe, *J. Cosmol. Astropart. Phys.* **08** (2020) 017.
- [40] S. Pi and M. Sasaki, Gravitational Waves Induced by Scalar Perturbations with a Lognormal Peak, *J. Cosmol. Astropart. Phys.* **09** (2020) 037.
- [41] C. Germani and T. Prokopec, On primordial black holes from an inflection point, *Phys. Dark Univ.* **18**, 6 (2017).
- [42] J. M. Ezquiaga, J. Garcia-Bellido, and E. Ruiz Morales, Primordial Black Hole production in Critical Higgs Inflation, *Phys. Lett. B* **776**, 345 (2018).
- [43] G. Ballesteros and M. Taoso, Primordial black hole dark matter from single field inflation, *Phys. Rev. D* **97**, 023501 (2018).
- [44] T.-J. Gao and Z.-K. Guo, Primordial Black Hole Production in Inflationary Models of Supergravity with a Single Chiral Superfield, *Phys. Rev. D* **98**, 063526 (2018).
- [45] I. Dalianis, S. Karydas, and E. Papantonopoulos, Generalized Non-Minimal Derivative Coupling: Application to Inflation and Primordial Black Hole Production, *J. Cosmol. Astropart. Phys.* **06** (2020) 040.
- [46] G. Ballesteros, J. Beltran Jimenez, and M. Pieroni, Black hole formation from a general quadratic action for inflationary primordial fluctuations, *J. Cosmol. Astropart. Phys.* **06** (2019) 016.
- [47] Y. Lu, Y. Gong, Z. Yi, and F. Zhang, Constraints on primordial curvature perturbations from primordial black hole dark matter and secondary gravitational waves, *J. Cosmol. Astropart. Phys.* **12** (2019) 031.
- [48] S. Passaglia, W. Hu, and H. Motohashi, Primordial black holes and local non-Gaussianity in canonical inflation, *Phys. Rev. D* **99**, 043536 (2019).
- [49] C. Fu, P. Wu, and H. Yu, Primordial Black Holes from Inflation with Nonminimal Derivative Coupling, *Phys. Rev. D* **100**, 063532 (2019).
- [50] J. Lin, Q. Gao, Y. Gong, Y. Lu, C. Zhang, and F. Zhang, Primordial black holes and secondary gravitational waves from k and G inflation, *Phys. Rev. D* **101**, 103515 (2020).
- [51] Z. Yi, Y. Gong, B. Wang, and Z.-h. Zhu, Primordial black holes and secondary gravitational

- waves from the Higgs field, *Phys. Rev. D* **103**, 063535 (2021).
- [52] Z. Yi, Q. Gao, Y. Gong, and Z.-h. Zhu, Primordial black holes and scalar-induced secondary gravitational waves from inflationary models with a noncanonical kinetic term, *Phys. Rev. D* **103**, 063534 (2021).
- [53] Z. Wang, S. Gao, Y. Gong, and Y. Wang, Primordial black holes and scalar-induced gravitational waves from the polynomial attractor model, [arXiv:2401.16069](https://arxiv.org/abs/2401.16069).
- [54] H. Di and Y. Gong, Primordial black holes and second order gravitational waves from ultra-slow-roll inflation, *J. Cosmol. Astropart. Phys.* **07** (2018) 007.
- [55] J. R. Espinosa, D. Racco, and A. Riotto, Cosmological Signature of the Standard Model Higgs Vacuum Instability: Primordial Black Holes as Dark Matter, *Phys. Rev. Lett.* **120**, 121301 (2018).
- [56] J. Garcia-Bellido and E. Ruiz Morales, Primordial black holes from single field models of inflation, *Phys. Dark Univ.* **18**, 47 (2017).
- [57] M. Sasaki, T. Suyama, T. Tanaka, and S. Yokoyama, Primordial black holes—perspectives in gravitational wave astronomy, *Classical Quantum Gravity* **35**, 063001 (2018).
- [58] N. C. Tsamis and R. P. Woodard, Improved estimates of cosmological perturbations, *Phys. Rev. D* **69**, 084005 (2004).
- [59] W. H. Kinney, Horizon crossing and inflation with large eta, *Phys. Rev. D* **72**, 023515 (2005).
- [60] J. Martin, H. Motohashi, and T. Suyama, Ultra Slow-Roll Inflation and the non-Gaussianity Consistency Relation, *Phys. Rev. D* **87**, 023514 (2013).
- [61] H. Motohashi, A. A. Starobinsky, and J. Yokoyama, Inflation with a constant rate of roll, *J. Cosmol. Astropart. Phys.* **09** (2015) 018.
- [62] S. M. Leach and A. R. Liddle, Inflationary perturbations near horizon crossing, *Phys. Rev. D* **63**, 043508 (2001).
- [63] S. M. Leach, M. Sasaki, D. Wands, and A. R. Liddle, Enhancement of superhorizon scale inflationary curvature perturbations, *Phys. Rev. D* **64**, 023512 (2001).
- [64] R. K. Jain, P. Chingangbam, and L. Sriramkumar, On the evolution of tachyonic perturbations at super-Hubble scales, *J. Cosmol. Astropart. Phys.* **10** (2007) 003.
- [65] M. H. Namjoo, H. Firouzjahi, and M. Sasaki, Violation of non-Gaussianity consistency relation in a single field inflationary model, *EPL* **101**, 39001 (2013).
- [66] Z. Yi and Y. Gong, On the constant-roll inflation, *J. Cosmol. Astropart. Phys.* **03** (2018) 052.

- [67] H. Motohashi and A. A. Starobinsky, Constant-roll inflation: confrontation with recent observational data, *EPL* **117**, 39001 (2017).
- [68] Q. Gao, The observational constraint on constant-roll inflation, *Sci. China Phys. Mech. Astron.* **61**, 070411 (2018).
- [69] J. T. Galvez Gherzi, A. Zucca, and A. V. Frolov, Observational Constraints on Constant Roll Inflation, *J. Cosmol. Astropart. Phys.* **05** (2019) 030.
- [70] Q. Gao, Y. Gong, and Z. Yi, On the constant-roll inflation with large and small η_H , *Universe* **5**, 215 (2019).
- [71] K. Tzirakis and W. H. Kinney, Inflation over the hill, *Phys. Rev. D* **75**, 123510 (2007).
- [72] M. J. P. Morse and W. H. Kinney, Large- η constant-roll inflation is never an attractor, *Phys. Rev. D* **97**, 123519 (2018).
- [73] S. Pi and M. Sasaki, Logarithmic Duality of the Curvature Perturbation, *Phys. Rev. Lett.* **131**, 011002 (2023).
- [74] M. Sasaki and E. D. Stewart, A General analytic formula for the spectral index of the density perturbations produced during inflation, *Prog. Theor. Phys.* **95**, 71 (1996).
- [75] D. Wands, K. A. Malik, D. H. Lyth, and A. R. Liddle, A New approach to the evolution of cosmological perturbations on large scales, *Phys. Rev. D* **62**, 043527 (2000).
- [76] D. H. Lyth, K. A. Malik, and M. Sasaki, A General proof of the conservation of the curvature perturbation, *J. Cosmol. Astropart. Phys.* **05** (2005) 004.
- [77] N. S. Sugiyama, E. Komatsu, and T. Futamase, δN formalism, *Phys. Rev. D* **87**, 023530 (2013).