

When Lawvere meets Peirce: an equational presentation of boolean hyperdoctrines

Filippo Bonchi

University of Pisa, Italy

Alessandro Di Giorgio 

University College London, United Kingdom

Davide Trotta

University of Pisa, Italy

Abstract

Fo-bicategories are a categorification of Peirce’s calculus of relations. Notably, their laws provide a proof system for first-order logic that is both purely equational and complete. This paper illustrates a correspondence between fo-bicategories and Lawvere’s hyperdoctrines. To streamline our proof, we introduce peircean bicategories, which offer a more succinct characterization of fo-bicategories.

2012 ACM Subject Classification Theory of computation → Logic; Theory of computation → Categorical semantics

Keywords and phrases relational algebra, hyperdoctrines, cartesian bicategories, string diagrams

1 Introduction

The first appearances of the characteristic features of first-order logic can be traced back to the works of Peirce [60] and Frege [26]. Frege was mainly motivated by the pursuit of a rigorous foundation for mathematics: his work was inspired by real analysis, bringing the concept of functions and variables into the logical realm [21]. On the other hand Peirce, inspired by the work of De Morgan [19] on relational reasoning, introduced a calculus in which operations allow the combination of relations and adhere to a set of algebraic laws. Like Boole’s algebra of classes [10], Peirce’s calculus of relations does not feature variables nor quantifiers and its sole deduction rule is substituting equals by equals.

Despite several negative results regarding axiomatizations for the entire calculus [57] and various fragments thereof [33, 69, 27, 2, 66], its lack of binder-related complexities, coupled with purely equational proofs, has rendered the calculus of relations highly influential in computer science, e.g., in the context of database theory [15], programming languages [67, 32, 44, 1, 43] and proof assistants [64, 65, 41].

In logic, the calculus played a secondary role for many years, likely because it is strictly less expressive than first-order logic [49]. This was until Tarski in [73] recognized its algebraic flavour and initiated a program of algebraizing first-order logic, including works such as [20, 31, 68]. Quoting Quine [68]:

“Logic in his adolescent phase was algebraic. There was Boole’s algebra of classes and Peirce’s algebra of relations. But in 1879 logic come of age, with Frege’s quantification theory. Here the bound variables, so characteristic of analysis rather than of algebra, became central to logic.”

Such a perspective, which regarded algebraic aspects and those concerning quantifiers as separate entities, changed with the work of Lawvere.

Thanks to the recent development of a new branch of mathematics, namely category theory, Lawvere introduced in [46, 47, 48] *hyperdoctrines* which enabled the study of logic from a pure algebraic perspective. The crucial insights of Lawvere was to show that quantifiers, as

well as many logical constructs, can be algebraically captured through the crucial notion of adjointness. Hyperdoctrines, along with many categorical structures related to logics, such as regular, Heyting, and boolean categories [37, 38], align with Frege’s functional perspective: arrows represent functions (terms), and relations are derived through specific constructions.

In the last decade, the paradigm shift towards treating data as a physical resource has motivated many computer scientists to move from traditional term-based (cartesian) syntax toward a string diagrammatic (monoidal) syntax [39, 71] (see e.g., [72, 3, 5, 9, 17, 22, 23, 29, 58, 62]). This shift in syntax enables an extension of Peirce’s calculus of relations that is as expressive as first-order logic, accompanied by an axiomatization that is purely equational and complete. The axioms are those of *first-order bicategories* [4]: see Figures 1, 2 and 3. In essence, a first-order bicategory, or fo-bicategory, encompasses a cartesian and a cocartesian bicategory [12], interacting as a linear bicategory [14], while additionally satisfying linear versions of Frobenius equations and adjointness conditions.

In this paper, we reconcile Lawvere’s understanding of logic with Peirce’s calculus of relations by illustrating a formal correspondence between boolean hyperdoctrines and first-order bicategories.

To reach such a correspondence, we found convenient to introduce the novel notion of *peircean bicategories*: these are cartesian bicategories with each homset carrying a boolean algebra where the negation behaves appropriately with *maps* – special arrows that intuitively generalize functions. Our first result (Theorem 26) establishes that peircean bicategories are equivalent to first-order bicategories.

While the definition of peircean bicategories is not purely equational, as in the case of fo-bicategories, it is notably more concise. Moreover, it allows us to reuse from [7] an adjunction between cartesian bicategories and *elementary and existential doctrines* [52, 51, 53], which are a generalisation of hyperdoctrines, corresponding to the regular (i.e., $\exists, =, \top, \wedge$) fragment of first-order logic.

Our main result (Theorem 31) reveals an adjunction between the category of first-order bicategories and the category of boolean hyperdoctrines. One can perceive this as a logical analogue of Fox’s theorem [25], which establishes an equivalence between categories with finite products and monoidal categories equipped with natural comonoids. The latter notion, like the one of fo-bicategories, is purely equational.

It is essential to note that our theorem establishes an adjunction rather than an equivalence. The discrepancy can be intuitively explained by observing that, akin to first-order logic, terms and formulas are distinct entities in hyperdoctrines. This differentiation does not exist in the calculus of relations or first-order bicategories. For instance, given two terms t_1 and t_2 , the hyperdoctrine where the formula $t_1 = t_2$ is true differs from the hyperdoctrine where t_1 and t_2 are equated as terms, a distinction not present in fo-bicategories. These issues, related to the extensionality of equality, are thoroughly analyzed in the context of doctrines in [51] and, more generally, in that of fibrations in [36].

Leveraging another result from [7], we demonstrate (Theorem 36) that the adjunction in Theorem 31 becomes an equivalence when restricted to well-behaved hyperdoctrines (i.e., those whose equality is extensional and satisfying the rule of unique choice [50]). Finally, combining this finding with a result in [50], we illustrate (Corollary 42) that *functionally complete* [12] first-order bicategories are equivalent to boolean categories [38].

Synopsis: In § 2, we provide a review of (co)cartesian bicategories, linear bicategories, and fo-bicategories. § 3 covers a recap of elementary and existential doctrines and boolean hyperdoctrines. The adjunction between cartesian bicategories and doctrines, as detailed

in [7], is presented in §4. Our original contributions commence in § 5, where we introduce peircean bicategories and establish their equivalence with fo-bicategories. This result is further used in § 6 to demonstrate the adjunction and in § 7 to establish the equivalence. § 8 elucidates the correspondence with boolean categories. All proofs are in appendix.

Terminology and Notation: All bicategories considered in this paper are just poset-enriched symmetric monoidal categories. For a bicategory \mathbf{C} , we will write \mathbf{C}^{op} for the bicategory having the same objects as \mathbf{C} but homsets $\mathbf{C}^{\text{op}}[X, Y] \stackrel{\text{def}}{=} \mathbf{C}[Y, X]$. Similarly, we will write \mathbf{C}^{co} to denote the bicategory having the same objects and arrows of \mathbf{C} but equipped with the reversed ordering \geq . The cartesian bicategories in this paper are called in [12] cartesian bicategories of relations. We refer the reader to [4, Rem. 2] for a comparison with the presentation of linear bicategories in [14].

2 From (Co)Cartesian to First-Order Bicategories

In this section we recall the notion of *first-order bicategory* from [4]. To provide a preliminary intuition, it is convenient to consider \mathbf{Rel} , the first-order bicategory of sets and relations.

It is well known that sets and relations form a symmetric monoidal category, hereafter denoted as \mathbf{Rel}° , with composition, identities, monoidal product and symmetries defined as

$$\begin{aligned} a \circ b &\stackrel{\text{def}}{=} \{(x, z) \mid \exists y \in Y. (x, y) \in a \wedge (y, z) \in b\} \subseteq X \times Z & id_X^\circ &\stackrel{\text{def}}{=} \{(x, y) \mid x = y\} \subseteq X \times X \\ a \otimes c &\stackrel{\text{def}}{=} \{((x, z), (y, v)) \mid (x, y) \in a \wedge (z, v) \in c\} \subseteq (X \times Z) \times (Y \times V) \\ \sigma_{X,Y}^\circ &\stackrel{\text{def}}{=} \{((x, y), (y', x')) \mid x = x' \wedge y = y'\} \subseteq (X \times Y) \times (Y \times X) \end{aligned} \quad (1)$$

for all sets X, Y, Z, V and relations $a \subseteq X \times Y, b \subseteq Y \times Z$ and $c \subseteq Z \times V$. As originally observed by Peirce in [61], beyond \circ there exists another form of relational composition that enjoys noteworthy algebraic properties. This different composition gives rise to another symmetric monoidal category of sets and relations, hereafter denoted by \mathbf{Rel}^\bullet and defined as follows.

$$\begin{aligned} a \bullet b &\stackrel{\text{def}}{=} \{(x, z) \mid \forall y \in Y. (x, y) \in a \vee (y, z) \in b\} \subseteq X \times Z & id_X^\bullet &\stackrel{\text{def}}{=} \{(x, y) \mid x \neq y\} \subseteq X \times X \\ a \boxtimes c &\stackrel{\text{def}}{=} \{((x, z), (y, v)) \mid (x, y) \in a \vee (z, v) \in c\} \subseteq (X \times Z) \times (Y \times V) \\ \sigma_{X,Y}^\bullet &\stackrel{\text{def}}{=} \{((x, y), (y', x')) \mid x \neq x' \vee y \neq y'\} \subseteq (X \times Y) \times (Y \times X) \end{aligned} \quad (2)$$


Note that \otimes and \boxtimes are both defined on objects as the cartesian product of sets and have as unit the singleton set $I \stackrel{\text{def}}{=} \{\star\}$. Both \mathbf{Rel}° and \mathbf{Rel}^\bullet are poset-enriched symmetric monoidal categories when taking as ordering the inclusion \subseteq and the complement \neg : $(\mathbf{Rel}^\circ)^{\text{co}} \rightarrow \mathbf{Rel}^\bullet$ is an isomorphism. As we will explain in § 2.1, the relations defined for all sets X as

$$\begin{aligned} \blacktriangleleft_X^\circ &\stackrel{\text{def}}{=} \{(x, (y, z)) \mid x = y \wedge x = z\} \subseteq X \times (X \times X) & \blacktriangleleft_X^\bullet &\stackrel{\text{def}}{=} \{(x, (y, z)) \mid x \neq y \vee x \neq z\} \subseteq X \times (X \times X) \\ \blacktriangleright_X^\circ &\stackrel{\text{def}}{=} \{((y, z), x) \mid x = y \wedge x = z\} \subseteq (X \times X) \times X & \blacktriangleright_X^\bullet &\stackrel{\text{def}}{=} \{((y, z), x) \mid x \neq y \vee x \neq z\} \subseteq (X \times X) \times X \\ !_X^\circ &\stackrel{\text{def}}{=} \{(x, \star) \mid x \in X\} \subseteq X \times I & !_X^\bullet &\stackrel{\text{def}}{=} \emptyset \subseteq X \times I \\ i_X^\circ &\stackrel{\text{def}}{=} \{(\star, x) \mid x \in X\} \subseteq I \times X & i_X^\bullet &\stackrel{\text{def}}{=} \emptyset \subseteq I \times X \end{aligned} \quad (3)$$

make \mathbf{Rel}° a cartesian bicategory, while \mathbf{Rel}^\bullet a cocartesian one.

Intuitively, a first-order bicategory \mathbf{C} consists of a cartesian bicategory \mathbf{C}° , called the “white structure”, and a cocartesian bicategory \mathbf{C}^\bullet , called the “black structure”, that interact by obeying the same laws of \mathbf{Rel}° and \mathbf{Rel}^\bullet . The name “first-order” is due to the fact that such laws provide a complete system of axioms for first-order logic.

The axioms can be conveniently given by means of a graphical representation inspired by string diagrams [39, 71]: composition is depicted as horizontal composition while



■ **Figure 1** Axioms of Cartesian bicategories

the monoidal product by vertically “stacking” diagrams. However, since there are two compositions \circ and \circ and two monoidal products \otimes and \boxtimes , to distinguish them we use different colors. All white constants have white background, mutatis mutandis for the black ones: for instance \triangleleft_X° and \triangleright_X° are drawn $x \begin{array}{|c|} \hline \triangleleft_X^\circ \\ \hline \end{array} x$ and $x \begin{array}{|c|} \hline \triangleright_X^\circ \\ \hline \end{array} x$, while for some arrows a, b, c, d of the appropriate type, $(a \otimes c) \circ (b \boxtimes d)$ is drawn as on the right of (ν_1°) in Figure 2.

2.1 (Co)Cartesian Bicategories

We commence with the notion of cartesian bicategories by Carboni and Walters [12].

► **Definition 1.** A cartesian bicategory $(\mathbf{C}, \otimes, I, \triangleleft^\circ, !^\circ, \triangleright^\circ, i^\circ)$, shorthand $(\mathbf{C}, \triangleleft^\circ, \triangleright^\circ)$, is a poset-enriched symmetric monoidal category (\mathbf{C}, \otimes, I) and, for every object X in \mathbf{C} , arrows $\triangleleft_X^\circ: X \rightarrow X \otimes X$, $!_X^\circ: X \rightarrow I$, $\triangleright_X^\circ: X \otimes X \rightarrow X$, $i_X^\circ: I \rightarrow X$ such that


1. $(\triangleleft_X^\circ, !_X^\circ)$ is a comonoid and $(\triangleright_X^\circ, i_X^\circ)$ a monoid, i.e., the equalities $(\triangleleft^\circ\text{-as})$, $(\triangleleft^\circ\text{-un})$, $(\triangleleft^\circ\text{-co})$ and $(\triangleright^\circ\text{-as})$, $(\triangleright^\circ\text{-un})$, $(\triangleright^\circ\text{-co})$ in Figure 1 hold;
2. every arrow $c: X \rightarrow Y$ is a lax comonoid homomorphism, i.e., $(\triangleleft^\circ\text{-nat})$ and $(!^\circ\text{-nat})$ hold;
3. comonoids are left adjoints to the monoids, i.e., $(\eta \triangleleft^\circ)$, $(\epsilon \triangleright^\circ)$, $(\eta !^\circ)$ and (ϵi°) hold;
4. monoids and comonoids form special Frobenius bimonoids, i.e., (F°) and (S°) hold;
5. monoids and comonoids satisfy the expected coherence conditions (see e.g. [7]).

\mathbf{C} is a cocartesian bicategory if \mathbf{C}^{co} is a cartesian bicategory. A morphism of (co)cartesian bicategories is a poset-enriched strong symmetric monoidal functor preserving monoids and comonoids. We denote by \mathbf{CIB} the category of cartesian bicategories and their morphisms.

As already mentioned, \mathbf{Rel}° with $\triangleleft_X^\circ, !_X^\circ, \triangleright_X^\circ$ and i_X° defined in (3) form a cartesian bicategory: the reader can easily check, using the definitions in (1) and (3), that all the laws in Figure 1 are satisfied. Similarly, one can observe that the opposite inequality of $(\triangleleft^\circ\text{-nat})$ holds iff the relation $c \subseteq X \times Y$ is single-valued (i.e., deterministic), while the opposite of $(!^\circ\text{-nat})$ iff c is total. In other words, c is a function iff both $(\triangleleft^\circ\text{-nat})$ and $(!^\circ\text{-nat})$ holds as equalities.

► **Definition 2.** Let $c: X \rightarrow Y$ be an arrow of a cartesian bicategory \mathbf{C} . It is a map if

$$x \begin{array}{|c|} \hline c \\ \hline \end{array} y \geq x \begin{array}{|c|} \hline \triangleleft_X^\circ \\ \hline \end{array} y \quad \text{and} \quad x \begin{array}{|c|} \hline c \\ \hline \end{array} y \geq x \begin{array}{|c|} \hline !_X^\circ \\ \hline \end{array} y. \quad (4)$$



■ **Figure 2** Axioms of closed symmetric monoidal linear bicategories

It is easy to see that maps form a monoidal subcategory of \mathbf{C} [12], hereafter denoted by $\mathbf{Map}(\mathbf{C})$. Since, by $(\blacktriangleleft^\circ\text{-nat})$, $(\blacktriangleright^\circ\text{-nat})$ and (4), comonoids are natural w.r.t. maps, Fox theorem [25] guarantees that $\mathbf{Map}(\mathbf{C})$ is a category with finite products.

In a cartesian bicategory \mathbf{C} , each homset $\mathbf{C}[X, Y]$ carries the structure of inf-semilattice, defined for all $c, d: X \rightarrow Y$ as in (5) below. Furthermore, the equation (6) defines an identity-on-objects isomorphism of cartesian bicategories $(\cdot)^\dagger: \mathbf{C} \rightarrow \mathbf{C}^{\text{op}}$.

$$c \wedge d \stackrel{\text{def}}{=} X \begin{array}{c} \boxed{c} \\ \boxed{d} \end{array} Y \quad \top \stackrel{\text{def}}{=} X \begin{array}{c} \bullet \\ \bullet \end{array} Y \quad (5) \quad c^\dagger \stackrel{\text{def}}{=} Y \begin{array}{c} \bullet \\ \bullet \end{array} \boxed{c} X \quad (6)$$

The reader can check, using (1) and (3) that in \mathbf{Rel}° , $c^\dagger: Y \rightarrow X$ is the opposite of the relation c , namely $\{(y, x) \mid (x, y) \in c\}$. It is well known that a relation c is a function iff it is left adjoint to c^\dagger . More generally in a cartesian bicategory c is a map iff it is left adjoint to c^\dagger . Summarising:

► **Proposition 3.** *Let \mathbf{C} be a cartesian bicategory and $c: X \rightarrow Y$ an arrow of \mathbf{C} . The following hold:*


1. *every homset carries the inf-semilattice structure, defined as in (5);*
2. *there is an isomorphism of cartesian bicategories $(\cdot)^\dagger: \mathbf{C} \rightarrow \mathbf{C}^{\text{op}}$, defined as in (6);*
3. *c is a map iff c is left adjoint to c^\dagger ;*
4. *$\mathbf{Map}(\mathbf{C})$ is a category with finite products; moreover, a morphism of cartesian bicategories $F: \mathbf{C} \rightarrow \mathbf{D}$ restricts to a functor $\tilde{F}: \mathbf{Map}(\mathbf{C}) \rightarrow \mathbf{Map}(\mathbf{D})$ preserving finite products.*

Hereafter, we draw $Y \begin{array}{c} \boxed{c} \\ \boxed{c} \end{array} X$ for $(X \begin{array}{c} \boxed{c} \\ \boxed{c} \end{array} Y)^\dagger$ and $X \begin{array}{c} \boxed{c} \\ \boxed{c} \end{array} Y$ for a map $c: X \rightarrow Y$.

We mentioned that \mathbf{Rel}^\bullet with $\blacktriangleleft_{X'}, \blacktriangleright_{X'}, \blacktriangleright_X$ and i_X^\bullet defined in (3) forms a cocartesian bicategory. To prove this, it is enough to observe that the complement \neg is a poset-enriched symmetric monoidal isomorphism $\neg: (\mathbf{Rel}^\circ)^{\text{co}} \rightarrow \mathbf{Rel}^\bullet$ preserving (co)monoids.

2.2 Linear Bicategories

We have seen that \mathbf{Rel}° forms a cartesian bicategory, and \mathbf{Rel}^\bullet a cocartesian bicategory. The next step consists in merging them into one entity and study their algebraic interactions. However, the coexistence of two different compositions \circ and \bullet on the same class of objects and arrows brings us out of the realm of ordinary categories. The appropriate setting is provided by *linear bicategories* [14] by Cockett, Koslowski and Seely.



■ **Figure 3** Additional axioms for fo-bicategories

► **Definition 4.** A linear bicategory $(\mathbf{C}, \circ, id^\circ, \circ, id^\bullet)$ consists of two poset-enriched categories $(\mathbf{C}, \circ, id^\circ)$ and $(\mathbf{C}, \circ, id^\bullet)$ with the same class of objects, arrows and orderings (but possibly different identities and compositions) such that \circ linearly distributes over \bullet , i.e., (δ_l) and (δ_r) in Figure 2 hold.

A symmetric monoidal linear bicategory $(\mathbf{C}, \circ, id^\circ, \circ, id^\bullet, \otimes, \sigma^\circ, \otimes, \sigma^\bullet, I)$, shortly $(\mathbf{C}, \otimes, \otimes, I)$, consists of a linear bicategory $(\mathbf{C}, \circ, id^\circ, \circ, id^\bullet)$ and two poset-enriched symmetric monoidal categories (\mathbf{C}, \otimes, I) and (\mathbf{C}, \otimes, I) s.t. \otimes and \otimes agree on objects, i.e., $X \otimes Y = X \otimes Y$, share the same unit I and

2. there are linear strengths for (\otimes, \otimes) , i.e., the inequalities (v_l°) , (v_r°) , (v_l^\bullet) and (v_r^\bullet) hold;
3. \otimes preserves id° colaxly and \otimes preserves id^\bullet laxly, i.e., (\otimes°) and (\otimes^\bullet) hold.

A morphism of symmetric monoidal linear bicategories $F: (\mathbf{C}_1, \otimes, \otimes, I) \rightarrow (\mathbf{C}_2, \otimes, \otimes, I)$ consists of two poset-enriched symmetric monoidal functors $F^\circ: (\mathbf{C}_1, \otimes, I) \rightarrow (\mathbf{C}_2, \otimes, I)$ and $F^\bullet: (\mathbf{C}_1, \otimes, I) \rightarrow (\mathbf{C}_2, \otimes, I)$ that agree on objects and arrows: $F^\circ(X) = F^\bullet(X)$ and $F^\circ(c) = F^\bullet(c)$.

All linear bicategories in this paper are symmetric monoidal. Hence, we usually omit the adjective *symmetric monoidal* and refer to them simply as linear bicategories. In linear bicategories one can define *linear adjoints*: for $a: X \rightarrow Y$ and $b: Y \rightarrow X$, a is *left linear adjoint* to b , or b is *right linear adjoint* to a , written $b \Vdash a$, if $id_X^\circ \leq a \bullet b$ and $b \circ a \leq id_Y^\bullet$.

► **Definition 5.** A linear bicategory $(\mathbf{C}, \otimes, \otimes, I)$ is said to be *closed* if every $a: X \rightarrow Y$ has both a left and a right linear adjoint and, in particular, the white symmetry σ° is both left and right linear adjoint to the black symmetry σ^\bullet ($\sigma^\bullet \Vdash \sigma^\circ \Vdash \sigma^\bullet$), i.e. $(\tau\sigma^\circ)$, $(\gamma\sigma^\circ)$, $(\tau\sigma^\bullet)$ and $(\gamma\sigma^\bullet)$ in Figure 2 hold.

Our main example is the closed linear bicategory **Rel** of sets and relations. The white structure is the symmetric monoidal category **Rel**[°] and the black structure is **Rel**[•]. Observe that the two have the same objects, arrows and ordering. The white and black monoidal products \otimes and \otimes agree on objects (they are the cartesian product of sets) and have common unit object (the singleton set I). By (1) and (2), one can easily check all the inequalities in Figure 2. Both left and right linear adjoints of a relation $c \subseteq X \times Y$ are given by $\neg c^\dagger$.

2.3 First-Order Bicategories

After (co)cartesian and linear bicategories, we can recall first-order bicategories from [4].

► **Definition 6.** A first-order bicategory \mathbf{C} consists of a closed linear bicategory $(\mathbf{C}, \otimes, \otimes, I)$, a cartesian bicategory $(\mathbf{C}, \blacktriangleleft^\circ, \blacktriangleright^\circ)$ and a cocartesian bicategory $(\mathbf{C}, \blacktriangleleft^\bullet, \blacktriangleright^\bullet)$, such that

1. the white comonoid $(\blacktriangleleft^\circ, !^\circ)$ is left and right linear adjoint to black monoid $(\blacktriangleright^\bullet, i^\bullet)$ and $(\blacktriangleright^\circ, i^\circ)$ is left and right linear adjoint to $(\blacktriangleleft^\bullet, !^\bullet)$ i.e., the 16 inequalities in the top of Figure 3 hold;
2. white and black (co)monoids satisfy the linear Frobenius laws, i.e. $(F^\bullet_\circ), (F^\circ_\bullet), (F^\bullet_\bullet), (F^\circ_\circ)$ hold.

A morphism of fo-bicategories is a morphism of linear bicategories and of (co)cartesian bicategories. We denote by \mathbf{FOB} the category of fo-bicategories and their morphisms.

We have seen that \mathbf{Rel} is a closed linear bicategory, \mathbf{Rel}° a cartesian bicategory and \mathbf{Rel}^\bullet a cocartesian bicategory. Given (3), it is easy to check the inequalities in Figure 3.

If \mathbf{C} is a fo-bicategory, then \mathbf{C}^{co} is a fo-bicategory when swapping white and black structures. Similarly, \mathbf{C}^{op} is a fo-bicategory when swapping monoids and comonoids.

In a fo-bicategory \mathbf{C} , left and right linear adjoints of an arrow c coincide and are denoted by c^\perp . The assignment $c \mapsto c^\perp$ gives rise to an identity-on-objects isomorphism of fo-bicategories $(\cdot)^\perp: \mathbf{C} \rightarrow (\mathbf{C}^{\text{co}})^{\text{op}}$. Similarly, $(\cdot)^\dagger: \mathbf{C} \rightarrow \mathbf{C}^{\text{op}}$ in (6) is also an isomorphism of fo-bicategories.

Since the diagram on the right commutes, one can define the complement as the diagonal of the square, namely $\neg(\cdot) \stackrel{\text{def}}{=} ((\cdot)^\perp)^\dagger$. Clearly $\neg: \mathbf{C} \rightarrow \mathbf{C}^{\text{co}}$ is an isomorphism of fo-bicategories. Moreover, it induces a boolean algebra on each homset of \mathbf{C} .

$$\begin{array}{ccc} \mathbf{C} & \xrightarrow{(\cdot)^\dagger} & \mathbf{C}^{\text{op}} \\ (\cdot)^\perp \downarrow & & \downarrow (\cdot)^\perp \\ (\mathbf{C}^{\text{co}})^{\text{op}} & \xrightarrow{\neg(\cdot)^\dagger} & \mathbf{C}^{\text{co}} \end{array}$$

► **Proposition 7.** *Let \mathbf{C} be a fo-bicategory. Then, every homset of \mathbf{C} is a boolean algebra.*

► **Proposition 8.** *Let $F: \mathbf{C} \rightarrow \mathbf{D}$ be a morphism of fo-bicategories. Then, $\neg F(c) = F(\neg c)$ for all arrows c , and hence F preserves the boolean structure on the homsets.*

The next properties of maps (Definition 2) plays a key role in our work.

► **Proposition 9.** *For all maps $f: X \rightarrow Y$ and arrows $c: Y \rightarrow Z$, it holds that $f \circ \neg c = \neg(f \circ c)$*

2.4 Freely Generated First-Order Bicategories

We conclude this section by giving to the reader a taste of how fo-bicategories relate to first-order theories. First, we recall from [4] the freely generated fo-bicategory \mathbf{FOB}_Σ .

Given a monoidal signature Σ , namely a set of symbols $R: n \rightarrow m$ with arity n and coarity m , \mathbf{FOB}_Σ is the fo-bicategory whose objects are natural numbers and arrows $c: n \rightarrow m$ are string diagrams generated by the following rules:

$$\begin{array}{l} \overline{\square}: 0 \rightarrow 0 \quad \overline{\text{---}}: 1 \rightarrow 1 \quad \overline{\text{X}}: 2 \rightarrow 2 \quad \overline{\boxed{R}}: n \rightarrow m \quad \overline{\boxed{c}}: n \rightarrow m, \boxed{d}: m \rightarrow o \\ \overline{\boxed{c}}: n \rightarrow m, \boxed{d}: m \rightarrow o \quad n \boxed{c} \boxed{d} o: n \rightarrow o \\ \overline{\boxed{c}}: n \rightarrow m, \boxed{d}: o \rightarrow p \\ n \boxed{c} \boxed{d} m: n + o \rightarrow m + p \\ \overline{\blacksquare}: 0 \rightarrow 0 \quad \overline{\text{---}}: 1 \rightarrow 1 \quad \overline{\text{X}}: 2 \rightarrow 2 \quad \overline{\boxed{R}}: m \rightarrow n \quad \overline{\boxed{c}}: n \rightarrow m, \boxed{d}: m \rightarrow o \\ n \boxed{c} \boxed{d} o: n \rightarrow o \\ \overline{\boxed{c}}: n \rightarrow m, \boxed{d}: o \rightarrow p \\ n \boxed{c} \boxed{d} m: n + o \rightarrow m + p \end{array}$$

More precisely, arrows are equivalence classes of string diagrams w.r.t $\lesssim \cap \gtrsim$, where \lesssim is the precongruence (w.r.t. \circ, \otimes, \circ and \otimes) generated by the axioms in Figures 1,2,3,4 (with X, Y, Z, W replaced by natural numbers, and a, b, c, d by diagrams of the appropriate type) and the axioms forcing \boxed{R} and $\boxed{\overline{R}}$ to be linear adjoints:

$$n \boxed{} \leq n \boxed{R} \boxed{R} n \quad m \boxed{\overline{R}} \boxed{R} m \leq m \boxed{} m \quad m \boxed{} m \leq m \boxed{R} \boxed{R} m \quad n \boxed{R} \boxed{R} n \leq n \boxed{} n$$

To give semantics to these diagrams we need *interpretations*, i.e. pairs $\mathcal{I} = (X, \rho)$, where X is a set and ρ is a function assigning to each $R: n \rightarrow m \in \Sigma$ a relation $\rho(R): X^n \rightarrow X^m$. Since \mathbf{FOB}_Σ is the free fo-bicategory, for any interpretation \mathcal{I} there exists a unique morphism of fo-bicategories $\mathcal{I}^\sharp: \mathbf{FOB}_\Sigma \rightarrow \mathbf{Rel}$ such that $\mathcal{I}^\sharp(1) = X$ and $\mathcal{I}^\sharp(\boxed{R}) = \rho(R) \subseteq X^n \times X^m$. Intuitively, \mathcal{I}^\sharp is defined inductively by (1), (2) and (3) with the free cases provided by \mathcal{I} .

A *diagrammatic first-order theory* is a pair $\mathbb{T} = (\Sigma, \mathbb{I})$ where Σ is a monoidal signature and \mathbb{I} is a set of *axioms*: pairs (c, d) for $c, d: n \rightarrow m$ in \mathbf{FOB}_Σ , standing for $c \leq d$. An interpretation \mathcal{I} is a *model* of \mathbb{T} if and only if, for all $(c, d) \in \mathbb{I}$, $\mathcal{I}^\sharp(c) \subseteq \mathcal{I}^\sharp(d)$. As illustrated in [4], one can generate the fo-bicategory $\mathbf{FOB}_\mathbb{T}$ and, in the spirit of Lawvere's functorial semantics [45], models of \mathbb{T} are in one-to-one correspondence with morphisms $F: \mathbf{FOB}_\mathbb{T} \rightarrow \mathbf{Rel}$.

► **Example 10.** Consider the theory $\mathbb{T} = (\Sigma, \mathbb{I})$, where $\Sigma = \{R: 1 \rightarrow 1\}$ and \mathbb{I} be as follows:


$$\{(\boxed{}, \boxed{R}), (\boxed{R} \boxed{R}, \boxed{R}), (\boxed{\overline{R}} \boxed{R}, \boxed{}), (\boxed{\overline{R}} \boxed{R}, \boxed{\overline{R}}), (\boxed{\overline{R}} \boxed{R}, \boxed{\overline{R}} \boxed{R})\}.$$

An interpretation is a set X and a relation $R \subseteq X \times X$. It is a model iff R is an order, i.e., reflexive ($id_X^\circ \subseteq R$), transitive ($R \circ R \subseteq R$), antisymmetric ($R \cap R^\circ \subseteq id^\circ$) and total ($\top \subseteq R \cup R^\circ$).

► **Remark 11.** A direct encoding of traditional first-order theories into diagrammatic ones is illustrated in [4]. Shortly, a predicate symbol P of arity n becomes a symbol $P: n \rightarrow 0 \in \Sigma$, drawn as $n \boxed{P}$, and a n -ary function symbol f becomes $f: n \rightarrow 1 \in \Sigma$, drawn as $n \boxed{f}$.

For instance, the formula $\exists x. P(x) \wedge Q(x, f(y))$ is rendered as on the right, where

$\boxed{\bullet}$ plays the role of \exists and $\boxed{\bullet \bullet}$ that of \wedge . Note that both predicate and function symbols of traditional first-order theories are regarded as symbols of the monoidal signature Σ . Function symbols are constrained to represent functions by adding to \mathbb{I} the axioms of maps, i.e., the inequalities in (4).



3 From Elementary-Existential Doctrines to Boolean Hyperdoctrines

The notion of hyperdoctrine was introduced by Lawvere in a series of seminal papers [46, 48], in order to provide an algebraic framework for first-order (intuitionistic) logic. Over the years, various generalizations and specializations of this concept have been formulated and applied across multiple domains in the fields of logic and computer science.

In this work, we employ a generalization of the notion of hyperdoctrine introduced by Maietti and Rosolini in [52, 51, 53], namely that of *elementary and existential doctrine*.

3.1 Elementary and Existential Doctrines

Elementary and existential doctrines can be thought of as a categorification of the so-called “regular fragment” of first-order intuitionistic logic, i.e. the $(\exists, =, \top, \wedge)$ -fragment.

Hereafter $\langle f, g \rangle$ denotes the pairing of f and g and Δ_X denotes $\langle id_X^\circ, id_X^\circ \rangle$.

► **Definition 12.** An elementary and existential doctrine is a functor $P: \mathbf{C}^{\text{op}} \rightarrow \text{InfSI}$ from the opposite of a category \mathbf{C} with finite products to the category of inf-semilattices such that:

- for every Y in \mathbf{C} there exists an element δ_Y in $P(Y \times Y)$, called equality predicate, such that for a morphism $\text{id}_X^\circ \times \Delta_Y: X \times Y \rightarrow X \times Y \times Y$ in \mathbf{C} and every element α in $P(X \times Y)$, the assignment

$$\exists_{\text{id}_X^\circ \times \Delta_Y}(\alpha) \stackrel{\text{def}}{=} P_{\langle \pi_1, \pi_2 \rangle}(\alpha) \wedge P_{\langle \pi_2, \pi_3 \rangle}(\delta_Y)$$

determines a left adjoint to the functor $P_{\text{id}_X^\circ \times \Delta_Y}: P(X \times Y \times Y) \rightarrow P(X \times Y)$;

- for any projection $\pi_X: X \times Y \rightarrow X$, the functor $P_{\pi_X}: P(X) \rightarrow P(X \times Y)$ has a left adjoint \exists_{π_X} , and these satisfy the Beck-Chevalley condition and Frobenius reciprocity, see [52, Sec. 2].

► **Remark 13.** In an elementary and existential doctrine, for every $f: X \rightarrow Y$ of \mathbf{C} the functor P_f has a left adjoint \exists_f that can be computed as $\exists_{\pi_Y}(P_{f \times \text{id}_Y}(\delta_Y) \wedge P_{\pi_X}(\alpha))$ for α in $P(X)$, where π_X and π_Y are the projections from $X \times Y$. These left adjoints satisfy the Frobenius reciprocity but not necessarily the Beck-Chevalley condition. See [54, Rem. 6.4].

► **Definition 14.** Let $P: \mathbf{C}^{\text{op}} \rightarrow \text{InfSI}$ and $R: \mathbf{D}^{\text{op}} \rightarrow \text{InfSI}$ be two elementary and existential doctrines. A morphism of elementary and existential doctrines is given by a pair (F, b) where

- $F: \mathbf{C} \rightarrow \mathbf{D}$ is a finite product preserving functor;
- $b: P \rightarrow F^{\text{op}} \circ R$ is a natural transformation;

satisfying the following conditions:

1. for every object X of \mathbf{C} , $b_{X \times X}(\delta_X) = \delta_{F(X) \times F(X)}$;
2. for every $\pi_X: X \times Y \rightarrow X$ of \mathbf{C} and for every α in $P(X \times Y)$, $\exists_{F(\pi_X)} b_{X \times Y}(\alpha) = b_X(\exists_{\pi_X}(\alpha))$.

$$\begin{array}{ccc} \mathbf{C}^{\text{op}} & & \\ \downarrow F^{\text{op}} & \searrow P & \\ \mathbf{D}^{\text{op}} & \xrightarrow{R} & \text{InfSI} \end{array} \quad \begin{array}{c} \downarrow b \\ \end{array}$$

We write \mathbb{EED} for the category of elementary and existential doctrines and morphisms.

► **Example 15.** The powerset functor $\mathcal{P}: \text{Set}^{\text{op}} \rightarrow \text{InfSI}$ is the archetypal example of an elementary and existential doctrine. More generally, for any regular category \mathbf{C} , the subobjects functor $\text{Sub}_{\mathbf{C}}: \mathbf{C}^{\text{op}} \rightarrow \text{InfSI}$ is an elementary and existential doctrine, see [51, 52]. This assignment extends to an inclusion of the category \mathbb{REG} of regular categories into \mathbb{EED} .

► **Example 16.** For a cartesian bicategory \mathbf{C} , the functor $\mathbf{C}[-, I]: \text{Map}(\mathbf{C})^{\text{op}} \rightarrow \text{InfSI}$ is an elementary and existential doctrine, where the actions of left adjoints is given $\exists_g(f) := f \circ g^\dagger$ [7, Thm. 20]. As we will see in §4, this assignment extends to an inclusion of \mathbf{CB} into \mathbb{EED} .

Similarly to cartesian bicategories, elementary and existential doctrines have enough structure to deal with the notion of *functional* (or single-valued) and *entire* (total) predicates.

► **Definition 17** (From [50]). Let $P: \mathbf{C}^{\text{op}} \rightarrow \text{InfSI}$ be an elementary and existential doctrine. An element $\alpha \in P(X \times Y)$ is said to be functional from X to Y if $P_{\langle \pi_1, \pi_2 \rangle}(\alpha) \wedge P_{\langle \pi_1, \pi_3 \rangle}(\alpha) \leq P_{\langle \pi_2, \pi_3 \rangle}(\delta_Y)$ in $P(X \times Y \times Y)$. Also, α is said to be entire from X to Y if $\top_X \leq \exists_{\pi_X}(\alpha)$ in $P(X)$.

► **Remark 18.** By definition, a morphism of elementary and existential doctrines preserves both \exists_{π_X} and δ_Y . Therefore it preserves functional and entire elements.

► **Example 19.** In the doctrine $\mathcal{P}: \text{Set}^{\text{op}} \rightarrow \text{InfSI}$ from Example 15, an element $\alpha \in \mathcal{P}(X \times Y)$ is functional if and only if it defines a partial function from X to Y , while it is entire if it provides a total relation from X to Y .

► **Example 20.** In the doctrine $\mathbf{C}[-, I]: \text{Map}(\mathbf{C})^{\text{op}} \rightarrow \text{InfSI}$ from Example 16, functional and entire elements are precisely maps of \mathbf{C} . A detailed proof is in Lemma 74 in Appendix E.

3.2 Boolean Hyperdoctrines

In this section we recall the notion of *boolean hyperdoctrine*, and some useful properties.

► **Definition 21** (boolean hyperdoctrine). *Let \mathbf{C} be a category with finite products. A functor $P: \mathbf{C}^{\text{op}} \rightarrow \mathbf{Bool}$ is a boolean hyperdoctrine if it is an elementary and existential doctrine.*

A morphism $(F, b): P \rightarrow R$ of boolean hyperdoctrines is a morphism of elementary and existential doctrines such that b_X is a morphism of boolean algebras for all objects X of \mathbf{C} . We denote by \mathbb{BHD} the category of boolean hyperdoctrines and their morphisms.

It is well-known that in first-order logic the universal quantifier can be derived by the existential quantifier and the negation. The same happens in boolean hyperdoctrines: for all arrows $f: X \rightarrow Y$, the functor $\forall_f(-) \stackrel{\text{def}}{=} \neg \exists_f \neg(-)$ is a right adjoint to P_f – see Appendix B.1.

► **Example 22.** The powerset functor $\mathcal{P}: \mathbf{Set}^{\text{op}} \rightarrow \mathbf{Bool}$ provides an example of boolean hyperdoctrine. This can be generalized to an arbitrary *boolean category* \mathbf{B} , namely a coherent category such that every subobject has a complement, see [38, Sec. A1.4, p. 38]. The subobjects functor on \mathbf{B} is a boolean hyperdoctrine $\text{Sub}_{\mathbf{B}}: \mathbf{B}^{\text{op}} \rightarrow \mathbf{Bool}$.

► **Example 23.** Given a standard first-order theory Th in a first-order language \mathcal{L} (for simplicity single sorted), one can consider the functor $\mathcal{L}^{\text{Th}}: \mathcal{V}^{\text{op}} \rightarrow \mathbf{Bool}$. The base category \mathcal{V} is the *syntactic* category of \mathcal{L} , i.e. the category where objects are natural numbers and morphisms are lists of terms, while the predicates of $\mathcal{L}^{\text{Th}}(n)$ are given by equivalence classes (with respect to provable reciprocal consequence \dashv) of well-formed formulae with free variables in $\{x_1, \dots, x_n\}$, and the partial order is given by the provable consequences, according to the fixed theory Th . In this case, the left adjoint to the weakening functor $\mathcal{L}_{\pi}^{\text{Th}}$ is computed by existentially quantifying the variables that are not involved in the substitution induced by the projection π . Dually, the right adjoint is computed by quantifying universally.

We conclude this section with a result that, intuitively, is the analogous of Proposition 9.

► **Lemma 24.** *Let $P: \mathbf{C}^{\text{op}} \rightarrow \mathbf{Bool}$ be a boolean hyperdoctrine and $\phi \in P(X \times Y)$ a functional and entire element from X to Y . For all $\psi \in P(Y \times Z)$, it holds that*

$$\exists \pi_{X \times Z} (P_{\pi_{X \times Y}}(\phi) \wedge P_{\pi_{Y \times Z}}(\neg \psi)) = \neg (\exists \pi_{X \times Z} (P_{\pi_{X \times Y}}(\phi) \wedge P_{\pi_{Y \times Z}}(\psi))).$$

4 An Adjunction and an Equivalence

In [7], cartesian bicategories and elementary existential doctrines are compared. The main results of [7, Thm. 28] states that there exists the following adjunction.

$$\begin{array}{ccc} \mathbf{CB} & \begin{array}{c} \xleftarrow{\text{Rel}} \\ \xrightarrow{\text{Hml}} \end{array} & \mathbb{EED} \end{array} \quad (7)$$

The embedding $\text{Hml}: \mathbf{CB} \rightarrow \mathbb{EED}$ maps a cartesian bicategory \mathbf{C} into the hom-functor $\mathbf{C}[-, I]: \text{Map}(\mathbf{C})^{\text{op}} \rightarrow \mathbf{InfSl}$ that, as explained in Example 16, is an elementary existential doctrine. A morphism of cartesian bicategories $F: \mathbf{C} \rightarrow \mathbf{D}$ is mapped to the morphism of doctrines (\tilde{F}, b^F) where $\tilde{F}: \text{Map}(\mathbf{C}) \rightarrow \text{Map}(\mathbf{D})$ is the functor F restricted to $\text{Map}(\mathbf{C})$ and $b_X^F: \mathbf{C}[X, I] \rightarrow \mathbf{D}[F(X), I]$ is defined as $b_X^F(c) \stackrel{\text{def}}{=} F(c)$ for all objects X of \mathbf{C} and arrows $c \in \mathbf{C}[X, I]$.

The functor $\text{Rel}: \mathbb{EED} \rightarrow \mathbf{CB}$ is a generalisation to elementary and existential doctrines of the construction of bicategory relations associated with a regular category (see [12, Ex. 1.4]). For $P: \mathbf{C}^{\text{op}} \rightarrow \mathbf{InfSl}$, the cartesian bicategory $\text{Rel}(P)$ is defined as follows:

- objects are those of \mathbf{C} ; for objects X, Y , the homsets $\mathbf{Rel}(P)[X, Y]$ are the posets $P(X \times Y)$;
- the identity for an object X is the equality predicate δ_X in $P(X \times X)$;
- composition of $\phi: X \rightarrow Y$ and $\psi: Y \rightarrow Z$ is given by $\exists_{\pi_{X \times Z}} (P_{\pi_{X \times Y}}(\phi) \wedge P_{\pi_{Y \times Z}}(\psi))$.

For a morphism of doctrines $(F, b): P \rightarrow Q$, the morphism of cartesian bicategories $\mathbf{Rel}(F, b): \mathbf{Rel}(P) \rightarrow \mathbf{Rel}(Q)$ is defined for all objects X as $\mathbf{Rel}(F, b)(X) \stackrel{\text{def}}{=} F(X)$ and for all arrows $\phi: X \rightarrow Y$ in $\mathbf{Rel}(P)$, i.e., elements $\phi \in P(X \times Y)$, as $\mathbf{Rel}(F, b)(\phi) \stackrel{\text{def}}{=} b_{X \times Y}(\phi)$. The reader is referred to [7] or to Appendix C for further details on the adjunction in (7).

Another result in [7, Thm. 35] shows that the adjunction in (7) restricts to an equivalence

$$\mathbf{CB} \equiv \overline{\mathbf{EED}} \quad (8)$$

where $\overline{\mathbf{EED}}$ is a full subcategory of \mathbf{EED} whose objects are particularly well-behaved doctrines. For the sake of readability, we will make clear in §7 what these doctrines are.

5 Peircean Bicategories

In this section we introduce the notion of *peircean bicategory*, and we prove that such a new notion provides an alternative presentation of fo-bicategories. The name peircean is due to the fact that, like in Peirce's algebra of relations [61], and differently from fo-bicategories, the structure of boolean algebra is taken as a primitive.

► **Definition 25.** A peircean bicategory consists of a cartesian bicategory $(\mathbf{C}, \blacktriangleleft^\circ, \blacktriangleright^\circ)$ such that

1. every homset $\mathbf{C}[X, Y]$ carries a Boolean algebra $(\mathbf{C}[X, Y], \vee, \perp, \wedge, \top, \neg)$;
2. for all maps $f: X \rightarrow Y$ and arrows $c: Y \rightarrow Z$,

$$f \circ \neg c = \neg(f \circ c). \quad (\neg M)$$

A morphism of peircean bicategories is a morphism of cartesian bicategories $F: \mathbf{C} \rightarrow \mathbf{D}$ such that $F(\neg c) = \neg F(c)$. We write \mathbf{PB} for the category of peircean bicategories and their morphisms.

By Propositions 7 and 9 every fo-bicategory is a peircean bicategory. By Proposition 8 every morphism of fo-bicategories is a morphism of peircean bicategories.

Vice versa, every peircean bicategory $(\mathbf{C}, \blacktriangleleft^\circ, \blacktriangleright^\circ)$ gives rise to a fo-bicategory. The black structure $(\mathbf{C}, \blacktriangleleft^\bullet, \blacktriangleright^\bullet)$ is defined as expected from the white one and \neg . Namely:

$$\begin{array}{llll} c \circ d \stackrel{\text{def}}{=} \neg(\neg c \circ \neg d) & id_X^\bullet \stackrel{\text{def}}{=} \neg id_X^\circ & c \otimes d \stackrel{\text{def}}{=} \neg(\neg c \otimes \neg d) & o_{X,Y}^\bullet \stackrel{\text{def}}{=} \neg o_{X,Y}^\circ \\ \blacktriangleleft_X^\bullet \stackrel{\text{def}}{=} \neg \triangleleft_X^\circ & !_X^\bullet \stackrel{\text{def}}{=} \neg !_X^\circ & \blacktriangleright_X^\bullet \stackrel{\text{def}}{=} \neg \triangleright_X^\circ & i_X^\bullet \stackrel{\text{def}}{=} \neg i_X^\circ \end{array} \quad (9)$$

With this definition, it is immediate to see that $\neg: (\mathbf{C}^{\text{co}}, \blacktriangleleft^\circ, \blacktriangleright^\circ) \rightarrow (\mathbf{C}, \blacktriangleleft^\bullet, \blacktriangleright^\bullet)$ is an isomorphism and thus to conclude that $(\mathbf{C}, \blacktriangleleft^\bullet, \blacktriangleright^\bullet)$ is a cocartesian bicategory. Proving that $(\mathbf{C}, \blacktriangleleft^\circ, \blacktriangleright^\circ)$ and $(\mathbf{C}, \blacktriangleleft^\bullet, \blacktriangleright^\bullet)$ give rise to a fo-bicategory is the main technical effort of this paper: the diagrammatic proof in Appendix D crucially exploits the boolean properties and $(\neg M)$.

► **Theorem 26.** There is an isomorphism of categories $\mathbf{FOB} \cong \mathbf{PB}$.

Note that, differently from Definition 6, Definition 25 is not purely axiomatic, since the property 2 requires f to be a map. However, the notion of a peircean bicategory is notably more succinct than that of a fo-bicategory, making it more convenient for our purposes.

6 An Equational Presentation of Boolean Hyperdoctrines

The main purpose of this section is to establish a formal link between fo-bicategories and boolean hyperdoctrines. In particular, we are going to show that the adjunction presented in (7) restricts to an adjunction between \mathbf{FOB} and \mathbf{BHD} . Theorem 26 allows us to conveniently work with peircean bicategories. We commence with the following result.

► **Proposition 27.** *Let \mathbf{C} be a peircean bicategory. Then $\mathbf{Hml}(\mathbf{C})$ is a boolean hyperdoctrine.*

Proof. By (7), $\mathbf{C}[-, I]: \mathbf{Map}(\mathbf{C})^{\text{op}} \rightarrow \mathbf{InfSl}$ is an elementary and existential doctrine and, by definition of peircean bicategories, $\mathbf{C}[X, I]$ is a boolean algebra for all objects X . To conclude that $\mathbf{C}[-, I]: \mathbf{Map}(\mathbf{C})^{\text{op}} \rightarrow \mathbf{Bool}$, one has only to show that, for all maps $f: X \rightarrow Y$, $\mathbf{C}[f, I]: \mathbf{C}[Y, I] \rightarrow \mathbf{C}[X, I]$ is a morphism of boolean algebras. Since, by (7), $\mathbf{C}[f, I]$ is a morphism of inf-semilattices, it is enough to show that it preserves negation: for all $c \in \mathbf{C}[Y, I]$

$$\begin{aligned} \mathbf{C}[f, I](\neg c) &= f \circ \neg c && \text{(Definition of } \mathbf{C}[-, I]) \\ &= \neg(f \circ c) && (\neg\mathcal{M}) \\ &= \neg\mathbf{C}[f, I](c) && \text{(Definition of } \mathbf{C}[-, I]) \end{aligned}$$

◀

The above proposition allows us to characterize peircean bicategories as follows:

► **Corollary 28.** *Let \mathbf{C} be a cartesian bicategory. Then it is a peircean bicategory if and only if $\mathbf{Hml}(\mathbf{C})$ is a boolean hyperdoctrine.*

To prove that, for any boolean hyperdoctrine P , $\mathbf{Rel}(P)$ is a peircean bicategory, we need to establish a formal correspondence between Definition 2 and Definition 17.

► **Proposition 29.** *Let $P: \mathbf{C}^{\text{op}} \rightarrow \mathbf{InfSl}$ be an elementary and existential doctrine. Then the maps of $\mathbf{Rel}(P)$ are precisely the functional and entire elements of P .*

► **Proposition 30.** *Let P be a boolean hyperdoctrine. Then $\mathbf{Rel}(P)$ is a peircean bicategory.*

Proof. By (7), $\mathbf{Rel}(P)$ is a cartesian bicategory. Since $P(X)$ is a boolean algebra for all objects X , then each hom-set $\mathbf{Rel}(P)[X, Y]$ – by definition $P(X \times Y)$ – is a boolean algebra. To conclude that $\mathbf{Rel}(P)$ is a peircean bicategory, it is enough to show that $(\neg\mathcal{M})$ holds, that is

$$\phi \circ \neg\psi = \neg(\phi \circ \psi)$$

for all maps $\phi \in \mathbf{Rel}(P)[X, Y]$ and arrows $\psi \in \mathbf{Rel}(P)[Y, Z]$. By Proposition 29, ϕ is a functional and entire element of P . Thus, one can rely on Lemma 24 to conclude that

$$\begin{aligned} \phi \circ \neg\psi &= \exists_{\pi_{X \times Z}} (P_{\pi_{X \times Y}}(\phi) \wedge P_{\pi_{Y \times Z}}(\neg\psi)) && \text{(Definition of } \mathbf{Rel}(P)) \\ &= \neg(\exists_{\pi_{X \times Z}} (P_{\pi_{X \times Y}}(\phi) \wedge P_{\pi_{Y \times Z}}(\psi))) && \text{(Lemma 24)} \\ &= \neg(\phi \circ \psi) && \text{(Definition of } \mathbf{Rel}(P)) \end{aligned}$$

◀

By Propositions 27 and 30 proving the following result amounts to a few routine checks.

► **Theorem 31.** *The adjunction in (7), restricts to the adjunction below on the left.*

$$\mathbf{PB} \begin{array}{c} \xleftarrow{\mathbf{Rel}} \\ \perp \\ \xrightarrow{\mathbf{Hml}} \end{array} \mathbf{BHD} \quad \text{Thus, by Theorem 26, there is an adjunction} \quad \mathbf{FOB} \begin{array}{c} \xleftarrow{\quad} \\ \perp \\ \xrightarrow{\quad} \end{array} \mathbf{BHD} .$$

7 Boolean Hyperdoctrines Representing First-Order Bicategories

As anticipated in §4, the adjunction in (7) becomes an equivalence for certain well-behaved doctrines. Definitions 32 and 33 state the conditions that such doctrines must satisfy.

► **Definition 32.** *An elementary and existential doctrine $P: \mathbf{C}^{\text{op}} \rightarrow \mathbf{InfSI}$ has comprehensive diagonals if for the equality predicate $\delta_X \in P(X)$ it holds that $P_{\Delta_X}(\delta_X) = \top_X$ and every arrow $f: Y \rightarrow X \times X$ such that $P_f(\delta_X) = \top_Y$ factors (uniquely) through Δ_X .*

Intuitively, a doctrine has comprehensive diagonals if its equality is *extensional*, namely if a formula $t_1 = t_2$ is true, then the terms t_1 and t_2 are syntactically equal. In the language of cartesian bicategories, for two maps t_1, t_2 , this can be stated by means of diagrams as

$$\text{if } x \begin{array}{|c|} \hline t_1 \text{---} t_2 \\ \hline \end{array} x = x \begin{array}{|c|} \hline \bullet \text{---} \bullet \\ \hline \end{array} x \text{ then } x \begin{array}{|c|} \hline t_1 \\ \hline \end{array} y = x \begin{array}{|c|} \hline t_2 \\ \hline \end{array} y. \quad (10)$$

While it is sometimes meaningful to consider syntactic doctrines (e.g. Example 23) in which the equality is not extensional, in several semantical doctrines this condition is satisfied.

► **Definition 33.** *Let $P: \mathbf{C}^{\text{op}} \rightarrow \mathbf{InfSI}$ be an elementary existential doctrine. We say that P satisfies the Rule of Unique Choice (RUC) if for every entire functional element ϕ in $P(X \times Y)$ there exists an arrow $f: X \rightarrow Y$ such that $\top_X \leq P_{\langle \text{id}_X, f \rangle}(\phi)$.*

The reader can think that a doctrine has (RUC) if for every element (intuitively formula) that is entire and functional, there exists an arrow in \mathbf{C} (intuitively a term) that represents it.

► **Example 34.** The doctrine $\mathcal{P}: \mathbf{Set}^{\text{op}} \rightarrow \mathbf{InfSI}$ has comprehensive diagonals, and it satisfies the (RUC) (since every functional and total relation can be represented by a function). More generally, every subobject doctrine $\text{Sub}_{\mathbf{C}}: \mathbf{C}^{\text{op}} \rightarrow \mathbf{InfSI}$ on a regular category, as presented in Example 15 satisfies the (RUC) and it has comprehensive diagonals, as observed in [50].

► **Example 35.** The doctrine $\mathbf{C}[-, I]: \mathbf{Map}(\mathbf{C})^{\text{op}} \rightarrow \mathbf{InfSI}$ presented in Example 16 satisfies the (RUC) and it has comprehensive diagonals, as proved in [7]. The reader can find a diagrammatic proof of (10) in Proposition 48 in Appendix A.

Hereafter –and in the equivalence in (8)– $\overline{\mathbf{IEED}}$ is the full subcategory of \mathbf{IEED} whose objects are doctrines satisfying (RUC) and with comprehensive diagonals. Similarly $\overline{\mathbf{BIHD}}$ is the full subcategory of \mathbf{BIHD} whose objects are boolean hyperdoctrines satisfying (RUC) and with comprehensive diagonals.

By means of Theorem 31, it is easy to prove that the equivalence in (8) restricts as follows.

► **Theorem 36.** $\mathbf{PB} \equiv \overline{\mathbf{BIHD}}$ and thus, by Theorem 26, $\mathbf{FOB} \equiv \overline{\mathbf{BIHD}}$.

8 Comparing Boolean Categories with First-Order Bicategories

In this section we show that boolean categories, described in Example 22, correspond to those fo-bicategories that are *functionally complete*, a property introduced in [12].

► **Definition 37.** *A cartesian bicategory $(\mathbf{C}, \blacktriangleleft, \blacktriangleright)$ is functionally complete if for every arrow $r: X \rightarrow I$ there exists a map $i: X_r \rightarrow X$, called tabulation of r , such that*

$$x_r \begin{array}{|c|} \hline i \text{---} i \\ \hline \end{array} x_r = x_r \begin{array}{|c|} \hline \\ \hline \end{array} x_r \quad \text{and} \quad x \begin{array}{|c|} \hline i \text{---} \bullet \\ \hline \end{array} = x \begin{array}{|c|} \hline r \\ \hline \end{array}.$$

► **Example 38.** Let us consider the category \mathbf{Rel}° . The tabulation of a relation $r: X \rightarrow I$ is given by the subset $X_r \subseteq X$ of those elements of X on which r is defined, together with the trivial inclusion $i: X_r \hookrightarrow X$.

The previous example emphasizes the essential intuition behind the concept of tabulation, namely, that a tabulation represents the “domain of definition of a relation”. A notion aiming to abstract the same concept has been introduced in the context of fibrations in [36] and, as particular instance, in the context of doctrines in [52], under the name of *comprehensions*.

► **Definition 39.** Let $P: \mathbf{C}^{\text{op}} \rightarrow \mathbf{InfSI}$ be an elementary and existential doctrine and α be an element of $P(X)$. A comprehension of α is an arrow $\{\alpha\}: X_\alpha \rightarrow X$ such that $P_{\{\alpha\}}(\alpha) = \top_{X_\alpha}$ and, for every $f: Y \rightarrow X$ such that $P_f(\alpha) = \top_Y$, there exists a unique arrow $g: Y \rightarrow X_\alpha$ such that $f = g \circ \{\alpha\}$. We say that P has comprehensions if every α has a comprehension. We say that P has full comprehensions if it has comprehensions and, $\alpha \leq \beta$ whenever $\{\alpha\}$ factors through $\{\beta\}$.

In the light of the previous definition, we can rephrase Definition 32, saying that an elementary doctrine has comprehensive diagonals if Δ_X is the comprehension of δ_X .

► **Example 40.** Every subobject doctrine $\text{Sub}_\mathbf{C}: \mathbf{C}^{\text{op}} \rightarrow \mathbf{InfSI}$ on a regular category, as presented in Example 15 has full comprehensions, as observed in [50]. In this case, the comprehension of a subobject is given by the subobject itself.

We prove that the notion of tabulation and that of full comprehension happen to be equivalent when we consider the doctrines associated with a cartesian bicategory.

► **Theorem 41.** A cartesian bicategory $(\mathbf{C}, \triangleleft^\circ, \triangleright^\circ)$ is functionally complete if and only if the elementary and existential doctrine $\mathbf{C}[-, I]: \mathbf{Map}(\mathbf{C})^{\text{op}} \rightarrow \mathbf{InfSI}$ has full comprehensions.

By combining Theorem 41 with Theorem 36 and Proposition 5.3 from [50], that characterizes doctrines satisfying (RUC) with comprehensive diagonals and full comprehensions (see also Appendix F), we obtain equivalences between \mathbf{FOB}_f , the full subcategory of \mathbf{FOB} of the functionally complete fo-bicategories, $\overline{\mathbf{BHD}}_c$, the full subcategory of $\overline{\mathbf{BHD}}$ of boolean doctrines with full comprehensions and \mathbf{BC} , the category of boolean categories.

► **Corollary 42.** $\mathbf{FOB}_f \equiv \overline{\mathbf{BHD}}_c \equiv \mathbf{BC}$.

9 Conclusions, Related and Future work

Theorems 31, 36 and Corollary 42 provide a solid bridge between functional and relational approaches to classical logic. The former rely on categorical structures that are usually defined by means of exactness properties; the latter on fo-bicategories which enjoy a purely equational presentation, much in the spirit of Boole’s algebra and Peirce’s calculus.

To achieve our result, we found it extremely convenient to introduce the notion of peircean bicategories that, by Theorem 26, provide a far handier characterisation of fo-bicategories.

The isomorphism between fo-bicategories and peircean bicategories might also be useful to establish a correspondence with *allegories* [27], likely the most influential approach to categorical relational algebra. Since cartesian bicategories are equivalent to unitary pretabular allegories [40], we expect that such allegories where, additionally, homsets carry boolean algebras and the negations satisfy $(\neg \mathcal{M})$ are equivalent to fo-bicategories. Despite searching the literature on allegories, we did not find analogous structures. Interestingly, the property $(\neg \mathcal{M})$ can be proven in any Peirce allegories, as shown in Proposition 4.6.1 in [59].

Boolean hyperdoctrines are used in [11] as a categorical treatment of another work of Peirce: *existential graphs* [70]. While the latter share some similarities with the graphical language of fo-bicategories there is one notable difference: negation is a primitive operator rather than a derived one, as it happens for instance also in [30] and Definition 25. In [4] and in §5, it is emphasised how this choice makes the resulting calculus less algebraic in flavour, having to deal with convoluted rules such as the one for (de)iteration or properties which are not purely equational, such as $(\neg M)$. Inspired by [11], another graphical language [56] akin to Peirce’s graphs is based on a decomposition of a hyperdoctrine into a bifibration. In this work, the categorical treatment revolves around the notion of monoidal *chiralities* [55], which are much more closer in spirit to fo-bicategories. We believe that our results might set an initial step towards a connection between fo-bicategories and chiralities.

A recent work [18] proposes a relational understanding of doctrines. However, these corresponds to the regular fragment of first-order logic, and thus it might be intriguing to understand the role of the additional black structure of first-order bicategories in this setting. Finally, it is also worth mentioning a closely related research line, exemplified by works such as [35, 28], along with the references therein. These work primarily focus on the categorical approach to classical proof theory, involving suitable variants of poset-enriched categories.

As future work we also aim to investigate how our characterizations can be extended to higher-order classical logic, which is categorically represented through the notion of *tripos* [34, 63]. Indeed, we believe that the constructions and results presented in this work, together with notion of tripos, can serve as a guide for defining a variant of fo-bicategories –hopefully, purely equational– capable of representing higher-order classical logic.

An important result in the theory of databases [13] shows that the problem of query inclusion (entailment of regular-logic formulas) is equivalent to the existence of morphisms of hypergraphs. This combinatorial characterisation found a neat algebraic understanding in [8] by means of cartesian bicategories. We hope that Theorem 26 may lead to an analogous combinatorial understanding of first-order logic.


References

- 1 Richard Bird and Oege De Moor. The algebra of programming. *NATO ASI DPD*, 152:167–203, 1996.
- 2 Benedikt Bollig, Alain Finkel, and Amrita Suresh. Bounded Reachability Problems Are Decidable in FIFO Machines. In Igor Konnov and Laura Kovács, editors, *31st International Conference on Concurrency Theory (CONCUR 2020)*, volume 171 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 49:1–49:17, Dagstuhl, Germany, 2020. Schloss Dagstuhl–Leibniz-Zentrum für Informatik. doi:10.4230/LIPIcs.CONCUR.2020.49.
- 3 Filippo Bonchi, Fabio Gadducci, Aleks Kissinger, Paweł Sobocinski, and Fabio Zanasi. String diagram rewrite theory I: rewriting with frobenius structure. *J. ACM*, 69(2):14:1–14:58, 2022. doi:10.1145/3502719.
- 4 Filippo Bonchi, Alessandro Di Giorgio, Nathan Haydon, and Paweł Sobocinski. Diagrammatic algebra of first order logic, 2024. arXiv:2401.07055.
- 5 Filippo Bonchi, Joshua Holland, Robin Piedeleu, Paweł Sobociński, and Fabio Zanasi. Diagrammatic algebra: From linear to concurrent systems. *Proceedings of the ACM on Programming Languages*, 3(POPL):25:1–25:28, January 2019. doi:10.1145/3290338.
- 6 Filippo Bonchi, Dusko Pavlovic, and Paweł Sobocinski. Functorial semantics for relational theories. *arXiv preprint arXiv:1711.08699*, 2017.
- 7 Filippo Bonchi, Alessio Santamaria, Jens Seeber, and Paweł Sobociński. On doctrines and cartesian bicategories. In *9th Conference on Algebra and Coalgebra in Computer Science (CALCO 2021)*. Schloss Dagstuhl–Leibniz-Zentrum für Informatik, 2021.

- 8 Filippo Bonchi, Jens Seeber, and Pawel Sobocinski. Graphical Conjunctive Queries. In Dan Ghica and Achim Jung, editors, *27th EACSL Annual Conference on Computer Science Logic (CSL 2018)*, volume 119 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 13:1–13:23, Dagstuhl, Germany, 2018. Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik. URL: <http://drops.dagstuhl.de/opus/volltexte/2018/9680>, doi:10.4230/LIPIcs.CSL.2018.13.
- 9 Filippo Bonchi, Pawel Sobocinski, and Fabio Zanasi. Full Abstraction for Signal Flow Graphs. In *Proceedings of the 42nd Annual ACM SIGPLAN-SIGACT Symposium on Principles of Programming Languages*, POPL '15, pages 515–526, New York, NY, USA, January 2015. Association for Computing Machinery. doi:10.1145/2676726.2676993.
- 10 George Boole. *The mathematical analysis of logic*. Philosophical Library, 1847.
- 11 Geraldine Brady and Todd Trimble. A categorical interpretation of c.s. peirce's propositional logic alpha. *Journal of Pure and Applied Algebra - J PURE APPL ALG*, 149:213–239, 06 2000. doi:10.1016/S0022-4049(98)00179-0.
- 12 A. Carboni and R. F. C. Walters. Cartesian bicategories I. *Journal of Pure and Applied Algebra*, 49(1):11–32, November 1987. doi:10.1016/0022-4049(87)90121-6.
- 13 Ashok K. Chandra and Philip M. Merlin. Optimal implementation of conjunctive queries in relational data bases. In *Proceedings of the Ninth Annual ACM Symposium on Theory of Computing*, STOC '77, pages 77–90, New York, NY, USA, May 1977. Association for Computing Machinery. doi:10.1145/800105.803397.
- 14 J. Robin B. Cockett, Jürgen Koslowski, and Robert AG Seely. Introduction to linear bicategories. *Mathematical Structures in Computer Science*, 10(2):165–203, 2000.
- 15 Edgar Frank Codd. A relational model of data for large shared data banks. *Communications of the ACM*, 26(1):64–69, 1983.
- 16 Bob Coecke and Ross Duncan. Interacting Quantum Observables. In Luca Aceto, Ivan Damgård, Leslie Ann Goldberg, Magnús M. Halldórsson, Anna Ingólfssdóttir, and Igor Walukiewicz, editors, *Automata, Languages and Programming*, Lecture Notes in Computer Science, pages 298–310, Berlin, Heidelberg, 2008. Springer. doi:10.1007/978-3-540-70583-3_25.
- 17 Bob Coecke and Ross Duncan. Interacting quantum observables: Categorical algebra and diagrammatics. *New Journal of Physics*, 13(4):043016, April 2011. doi:10.1088/1367-2630/13/4/043016.
- 18 Francesco Dagnino and Fabio Pasquali. Quotients and Extensionality in Relational Doctrines. In Marco Gaboardi and Femke van Raamsdonk, editors, *8th International Conference on Formal Structures for Computation and Deduction (FSCD 2023)*, volume 260 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 25:1–25:23, Dagstuhl, Germany, 2023. Schloss Dagstuhl – Leibniz-Zentrum für Informatik. URL: <https://drops-dev.dagstuhl.de/entities/document/10.4230/LIPIcs.FSCD.2023.25>, doi:10.4230/LIPIcs.FSCD.2023.25.
- 19 Augustus De Morgan. On the syllogism, no. iv. and on the logic of relations. 1860.
- 20 Charles J Everett and Stanislaw Ulam. Projective algebra i. *American Journal of Mathematics*, 68(1):77–88, 1946.
- 21 William Ewald. The emergence of first-order logic. 2018.
- 22 Brendan Fong, Paweł Sobociński, and Paolo Rapisarda. A categorical approach to open and interconnected dynamical systems. In *Proceedings of the 31st Annual ACM/IEEE Symposium on Logic in Computer Science*, LICS '16, pages 495–504, New York, NY, USA, July 2016. Association for Computing Machinery. doi:10.1145/2933575.2934556.
- 23 Brendan Fong and David Spivak. String diagrams for regular logic (extended abstract). In John Baez and Bob Coecke, editors, *Applied Category Theory 2019*, volume 323 of *Electronic Proceedings in Theoretical Computer Science*, pages 196–229. Open Publishing Association, Sep 2020. doi:10.4204/eptcs.323.14.
- 24 Brendan Fong and David I. Spivak. Regular and relational categories: Revisiting 'cartesian bicategories I'. *arXiv*, 2019. URL: <https://api.semanticscholar.org/CorpusID:202540886>.
- 25 T. Fox. Coalgebras and cartesian categories. *Communications in Algebra*, 4(7):665–667, 1976. doi:10.1080/00927877608822127.

- 26 GOTTLOB FREGE. *Begriffsschrift und andere Aufsätze*. Georg Olms Verlag, 1977.
- 27 Peter Freyd and Andre Scedrov. *Categories, Allegories*, volume 39 of *North-Holland Mathematical Library*. Elsevier B.V, 1990.
- 28 CARSTEN FÜHRMANN and DAVID PYM. On categorical models of classical logic and the geometry of interaction. *Mathematical Structures in Computer Science*, 17(5):957–1027, 2007. doi:10.1017/S0960129507006287.
- 29 Dan R. Ghica and Achim Jung. Categorical semantics of digital circuits. In *2016 Formal Methods in Computer-Aided Design (FMCAD)*, pages 41–48, 2016. doi:10.1109/FMCAD.2016.7886659.
- 30 Nathan Haydon and Paweł Sobociński. Compositional diagrammatic first-order logic. In *11th International Conference on the Theory and Application of Diagrams (DIAGRAMS 2020)*, 2020.
- 31 Leon Henkin. Cylindric algebras. 1971.
- 32 CAR Hoare and He Jifeng. The weakest prespecification, part i. *Fundamenta Informaticae*, 9(1):51–84, 1986.
- 33 Ian Hodkinson and Szabolcs Mikuláš. Axiomatizability of reducts of algebras of relations. *Algebra Universalis*, 43(2):127–156, August 2000. doi:10.1007/s0001200050150.
- 34 J.M.E. Hyland, P.T. Johnstone, and A.M. Pitts. Triples theory. *Math. Proc. Camb. Phil. Soc.*, 88:205–232, 1980.
- 35 Martin Hyland. Abstract interpretation of proofs: Classical propositional calculus. In Jerzy Marcinkowski and Andrzej Tarlecki, editors, *Computer Science Logic*, pages 6–21, Berlin, Heidelberg, 2004. Springer Berlin Heidelberg.
- 36 B. Jacobs. *Categorical Logic and Type Theory*, volume 141 of *Studies in Logic and the foundations of mathematics*. North Holland Publishing Company, 1999.
- 37 P.T. Johnstone. *Topos Theory*. Academic Press, 1977.
- 38 P.T. Johnstone. *Sketches of an elephant: a topos theory compendium*, volume 2 of *Studies in Logic and the foundations of mathematics*. Oxford Univ. Press, 2002.
- 39 André Joyal and Ross Street. The geometry of tensor calculus, I. *Advances in Mathematics*, 88(1):55–112, July 1991. doi:10.1016/0001-8708(91)90003-P.
- 40 Petrus Marinus Waltherus Knijnenburg and Frank Nordemann. *Two Categories of Relations*. Citeseer, 1994.
- 41 Alexander Krauss and Tobias Nipkow. Proof pearl: Regular expression equivalence and relation algebra. *Journal of Automated Reasoning*, 49(1):95–106, 2012. doi:10.1007/s10817-011-9223-4.
- 42 Stephen Lack. Composing PROPs. *Theory and Application of Categories*, 13(9):147–163, 2004. URL: <http://www.tac.mta.ca/tac/volumes/13/9/13-09abs.html>.
- 43 Ugo Dal Lago and Francesco Gavazzo. A relational theory of effects and coeffects. *Proc. ACM Program. Lang.*, 6(POPL):1–28, 2022. doi:10.1145/3498692.
- 44 Søren B Lassen. Relational reasoning about contexts. *Higher order operational techniques in semantics*, 91, 1998.
- 45 F. W. Lawvere. *Functorial Semantics of Algebraic Theories*. PhD thesis, Columbia University, New York, NY, USA, 1963.
- 46 F.W. Lawvere. Adjointness in foundations. *Dialectica*, 23:281–296, 1969.
- 47 F.W. Lawvere. Diagonal arguments and cartesian closed categories. In *Category Theory, Homology Theory and their Applications*, volume 2, pages 134–145. Springer, 1969.
- 48 F.W. Lawvere. Equality in hyperdoctrines and comprehension schema as an adjoint functor. In A. Heller, editor, *New York Symposium on Application of Categorical Algebra*, volume 2, pages 1–14. American Mathematical Society, 1970.
- 49 LEOPOLD LÖWENHEIM. Über möglichkeiten im relativkalkül. *Mathematische Annalen*, 76(4):447–470, 1915.
- 50 M.E. Maietti, F. Pasquali, and G. Rosolini. Triples, exact completions, and hilbert’s ε -operator. *Tbilisi Mathematica journal*, 10(3):141–166, 2017.
- 51 M.E. Maietti and G. Rosolini. Elementary quotient completion. *Theory App. Categ.*, 27(17):445–463, 2013.

- 52 M.E. Maietti and G. Rosolini. Quotient completion for the foundation of constructive mathematics. *Log. Univers.*, 7(3):371–402, 2013.
- 53 M.E. Maietti and G. Rosolini. Unifying exact completions. *Appl. Categ. Structures*, 23:43–52, 2013.
- 54 M.E. Maietti and D. Trotta. A characterization of generalized existential completions. *Annals of Pure and Applied Logic*, 174(4):103234, 2023.
- 55 Paul-André Melliès. Dialogue categories and chiralities. *Publications of the Research Institute for Mathematical Sciences*, 52(4):359–412, 2016.
- 56 Paul-André Melliès and Noam Zeilberger. A bifibrational reconstruction of lawvere’s presheaf hyperdoctrine. In *Proceedings of the 31st Annual ACM/IEEE Symposium on Logic in Computer Science*, pages 555–564, 2016.
- 57 Donald Monk. On representable relation algebras. *Michigan Mathematical Journal*, 11(3):207 – 210, 1964. doi:10.1307/mmj/1028999131.
- 58 Koko Muroya, Steven W. T. Cheung, and Dan R. Ghica. The geometry of computation-graph abstraction. In Anuj Dawar and Erich Grädel, editors, *Proceedings of the 33rd Annual ACM/IEEE Symposium on Logic in Computer Science, LICS 2018, Oxford, UK, July 09-12, 2018*, pages 749–758. ACM, 2018. doi:10.1145/3209108.3209127.
- 59 Jean-Pierre Olivier and Dany Serrato. Peirce allegories. identities involving transitive elements and symmetrical ones. *Journal of Pure and Applied Algebra*, 116(1-3):249–271, 1997.
- 60 Charles S. Peirce. The logic of relatives. *The Monist*, 7(2):161–217, 1897. URL: <http://www.jstor.org/stable/27897407>.
- 61 Charles Sanders Peirce. *Studies in logic. By members of the Johns Hopkins university*. Little, Brown, and Company, 1883.
- 62 Robin Piedeleu and Fabio Zanasi. A String Diagrammatic Axiomatisation of Finite-State Automata. In Stefan Kiefer and Christine Tasson, editors, *Foundations of Software Science and Computation Structures*, Lecture Notes in Computer Science, pages 469–489, Cham, 2021. Springer International Publishing. doi:10.1007/978-3-030-71995-1_24.
- 63 A.M. Pitts. Tripos theory in retrospect. *Math. Struct. in Comp. Science*, 12:265–279, 2002.
- 64 Damien Pous. Kleene algebra with tests and coq tools for while programs. In *Interactive Theorem Proving: 4th International Conference, ITP 2013, Rennes, France, July 22-26, 2013. Proceedings 4*, pages 180–196. Springer, 2013.
- 65 Damien Pous. *Automata for relation algebra and formal proofs*. PhD thesis, ENS Lyon, 2016.
- 66 Damien Pous. On the positive calculus of relations with transitive closure. In Rolf Niedermeier and Brigitte Vallée, editors, *35th Symposium on Theoretical Aspects of Computer Science, STACS 2018, February 28 to March 3, 2018, Caen, France*, volume 96 of *LIPIcs*, pages 3:1–3:16. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2018. doi:10.4230/LIPIcs.STACS.2018.3.
- 67 Vaughan R Pratt. Semantical considerations on floyd-hoare logic. In *17th Annual Symposium on Foundations of Computer Science (sfcs 1976)*, pages 109–121. IEEE, 1976.
- 68 W.V. Quine. Predicate-functor logics. In J.E. Fenstad, editor, *Proceedings of the Second Scandinavian Logic Symposium*, volume 63 of *Studies in Logic and the Foundations of Mathematics*, pages 309–315. Elsevier, 1971. URL: <https://www.sciencedirect.com/science/article/pii/S0049237X08708504>, doi:https://doi.org/10.1016/S0049-237X(08)70850-4.
- 69 Valentin N Redko. On defining relations for the algebra of regular events. *Ukrainskii Matematicheskii Zhurnal*, 16:120–126, 1964.
- 70 Don D. Roberts. *The Existential Graphs of Charles S. Peirce*. De Gruyter Mouton, 1973.
- 71 P. Selinger. A Survey of Graphical Languages for Monoidal Categories. In B. Coecke, editor, *New Structures for Physics*, volume 813 of *Lecture Notes in Physics*, pages 289–355. Springer, Berlin, Heidelberg, 2010. doi:10.1007/978-3-642-12821-9_4.
- 72 Dario Stein and Sam Staton. Probabilistic programming with exact conditions. *Journal of the ACM*, 2023.
- 73 Alfred Tarski. On the calculus of relations. *The Journal of Symbolic Logic*, 6(3):73–89, September 1941. doi:10.2307/2268577.



■ **Figure 4** Axioms of Cocartesian bicategories

A Appendix to Section 2

In this appendix we collect some useful results about (co)cartesian bicategories. Moreover, we report in diagrammatic form the axioms of cocartesian bicategories (Figure 4) and we summarise in Table 1 the properties of the isomorphism $(\cdot)^\dagger: \mathbf{C} \rightarrow \mathbf{C}^{\text{op}}$.

► **Theorem 43.** Any connected diagram $c: X^n \rightarrow X^m$ made out of $\text{id}^\circ, \sigma^\circ, \triangleleft^\circ, !^\circ, \triangleright^\circ$ and i° is equal

$$\text{to } n \left\{ \begin{array}{c} X \\ X \\ \vdots \\ X \end{array} \right\} \left[\begin{array}{c} \text{diagram} \end{array} \right] \left\{ \begin{array}{c} X \\ X \\ \vdots \\ X \end{array} \right\} m.$$

Proof. See [42, 16]. ◀

► **Remark 44.** Theorem 43 is known as the *spider theorem* and it holds in any special Frobenius algebra and thus, in particular, in a cartesian bicategory. A direct consequence of the spider theorem is that we can “rewire” parts of a diagram involving the (co)monoid structures, as long as the connectivity is preserved. This is useful in several graphical derivations for arranging diagrams into a desired shape.

► **Lemma 45.** For any $c, d: X \rightarrow Y$, $x \begin{array}{|c|} \hline c \\ \hline \end{array} y = x \begin{array}{|c|} \hline d \\ \hline \end{array} y$ if and only if $x \begin{array}{|c|} \hline c \\ \hline \end{array} \bullet = x \begin{array}{|c|} \hline d \\ \hline \end{array} \bullet$.

Proof. One direction is trivial. The other direction follows from the Frobenius axioms. ◀

► **Proposition 46.** For any $c: X \rightarrow Y$ the following inequality holds $\begin{array}{|c|} \hline c \\ \hline \end{array} \bullet \leq \begin{array}{|c|} \hline c \\ \hline \end{array} \bullet$.

Proof. See Lemma 4.3 in [6]. ◀

► **Proposition 47.** For any $c: X \rightarrow Y$ the following equality holds $\begin{array}{|c|} \hline c \\ \hline \end{array} \bullet = \begin{array}{|c|} \hline c \\ \hline \end{array} \bullet$.

Proof.

$$\begin{array}{|c|} \hline c \\ \hline \end{array} \bullet \stackrel{(6)}{=} \begin{array}{|c|} \hline c \\ \hline \end{array} \bullet \stackrel{\text{Theorem 43}}{=} \begin{array}{|c|} \hline c \\ \hline \end{array} \bullet \approx \begin{array}{|c|} \hline c \\ \hline \end{array} \bullet \stackrel{(\triangleright^\circ\text{-as})}{=} \begin{array}{|c|} \hline c \\ \hline \end{array} \bullet$$

■ **Table 1** Properties of $(\cdot)^\dagger: \mathbf{C} \rightarrow \mathbf{C}^{\text{op}}$

if $c \leq d$ then $c^\dagger \leq d^\dagger$			
$(c \circ d)^\dagger = d^\dagger \circ c^\dagger$	$(id_X^\circ)^\dagger = id_X^\circ$	$(\blacktriangleright_X^\circ)^\dagger = \blacktriangleleft_X^\circ$	$(i_X^\circ)^\dagger = !_X^\circ$
$(c \otimes d)^\dagger = c^\dagger \otimes d^\dagger$	$(\sigma_{X,Y}^\circ)^\dagger = \sigma_{Y,X}^\circ$	$(\blacktriangleleft_X^\circ)^\dagger = \blacktriangleright_X^\circ$	$(!_X^\circ)^\dagger = i_X^\circ$
$(c \wedge d)^\dagger = c^\dagger \wedge d^\dagger$		$\top^\dagger = \top$	

► **Proposition 48** (Extensional equality). *For any $t_1, t_2: X \rightarrow Y$ maps,*

$$\text{if } x \boxed{t_1 \text{---} t_2} x = x \boxed{\bullet \bullet} x \text{ then } x \boxed{t_1} Y = x \boxed{t_2} Y .$$

Proof. First observe that if $x \boxed{t_1 \text{---} t_2} x = x \boxed{\bullet \bullet} x$, then by the properties in Table 1 the following holds

$$x \boxed{t_2 \text{---} t_1} x = (x \boxed{t_1 \text{---} t_2} x)^\dagger = (x \boxed{\bullet \bullet} x)^\dagger = x \boxed{\bullet \bullet} x . \quad (11)$$

To conclude we show the two inclusions separately:

$$\begin{array}{ll}
 x \boxed{t_1} Y \leq x \boxed{\bullet \bullet \text{---} t_1} Y & (\eta!^\circ) \\
 = x \boxed{t_2 \text{---} t_1 \text{---} t_1} Y & (11) \\
 \leq x \boxed{t_2} Y & (\text{Prop. 3.3})
 \end{array}
 \quad
 \begin{array}{ll}
 x \boxed{t_2} Y \leq x \boxed{\bullet \bullet \text{---} t_2} Y & (\eta!^\circ) \\
 = x \boxed{t_1 \text{---} t_2 \text{---} t_2} Y & (\text{Hyp.}) \\
 \leq x \boxed{t_1} Y & (\text{Prop. 3.3})
 \end{array}
 \quad \blacktriangleleft$$

B Appendix to Section 3

B.1 A few properties of Boolean hyperdoctrines

In this section we recall some useful properties of boolean hyperdoctrines, and we prove a lemma which will be crucial for establishing the precise connection with peircean bicategories. All the results we are going to show here are quite natural from the perspective of first-order classical logic, and their proofs are straightforward.

First, it is well-known in first-order classical logic that the universal quantifier can be defined combining the existential quantifier with the negation. In the following lemma we provide a proof using the language of f.o. boolean hyperdoctrine of this fact.

► **Lemma 49.** *Let $P: \mathbf{C}^{\text{op}} \rightarrow \mathbf{Bool}$ be a boolean hyperdoctrine. Then for every arrow $f: A \rightarrow B$, the functor*

$$\forall_f(-) := \neg \exists_f \neg (-)$$

provides a right adjoint to P_f . Moreover, if \exists_f satisfies BCC then also \forall_f satisfies BCC.

Proof. The proof is a straightforward generalization of the ordinary proof in first-order classical logic. Indeed:

$$\alpha \leq \forall_f P_f(\alpha) = \neg \exists_f \neg P_f(\alpha)$$

holds if and only if

$$\exists_f \neg P_f(\alpha) \leq \neg \alpha$$

because P is boolean. But this holds because it is equivalent to

$$P_f(\neg\alpha) \leq P_f(\neg\alpha).$$

To prove that

$$P_f \forall_f(\beta) \leq \beta$$

we have to use again the assumption that P is boolean. In fact we have that

$$P_f \forall_f(\beta) = \neg P_f \exists_f(\neg\beta) \leq \neg\neg\beta$$

because $\neg\beta \leq P_f \exists_f(\neg\beta)$ (since $\exists_f \dashv P_f$), and using the fact that $\neg\neg\beta = \beta$ we can conclude that $P_f \forall_f(\beta) \leq \beta$. Now we prove that if \exists_f satisfies BCC then also \forall_f satisfies BCC. So let us consider a pullback

$$\begin{array}{ccc} D & \xrightarrow{f'} & C \\ g' \downarrow & \lrcorner & \downarrow g \\ A & \xrightarrow{f} & B \end{array}$$

and suppose that $P_g \exists_f(\alpha) = \exists_{f'} P_{g'}(\alpha)$ for every α in $P(A)$. From this we can deduce that $\neg P_g \exists_f(\alpha) = P_g \neg \exists_f(\alpha) = \neg \exists_{f'} P_{g'}(\alpha)$, and then that $P_g \forall_f(\neg\alpha) = \forall_{f'} P_{g'}(\neg\alpha)$ for every α element of $P(A)$. Therefore, in particular it holds for the element $\beta := \neg\alpha$, and hence (using the fact that P is boolean) we can conclude that $P_g \forall_f(\alpha) = \forall_{f'} P_{g'}(\alpha)$. ◀

► **Remark 50 (Frobenius reciprocity).** Employing the preservation of the implication \rightarrow by the functor P_f , it is straightforward to check that every first-order boolean hyperdoctrine $P: \mathbf{C}^{\text{op}} \rightarrow \mathbf{Bool}$ satisfies the so-called *Frobenius reciprocity* (FR), namely:

$$\exists_f(P_f(\alpha) \wedge \beta) = \alpha \wedge \exists_f(\beta) \text{ and } \forall_f(P_f(\alpha) \rightarrow \beta) = \alpha \rightarrow \forall_f(\beta)$$

for every morphism $f: X \rightarrow Y$, α in $P(Y)$ and β in $P(X)$. See [34, Rem. 1.3]. However, it is not guaranteed that of Beck-Chevalley conditions with respect to pullbacks along f . See [54] for more details.

► **Lemma 51.** *Let $P: \mathbf{C}^{\text{op}} \rightarrow \mathbf{Bool}$ be a boolean hyperdoctrine and $\phi \in P(X \times Y)$ a functional and entire element from X to Y . For all $\psi \in P(Y \times Z)$, it holds that*

$$\exists_{\pi_{X \times Z}}(P_{\pi_{X \times Y}}(\phi) \wedge P_{\pi_{Y \times Z}}(\neg\psi)) = \neg(\exists_{\pi_{X \times Z}}(P_{\pi_{X \times Y}}(\phi) \wedge P_{\pi_{Y \times Z}}(\psi))).$$

Proof of Lemma 24. The proof happens to be straightforward if we employ the classical arguments of natural deduction. Here we provide a completely algebraic proof. For sake of readability, here we employ the notation given by the internal language of P , writing $\exists y$ instead of the left adjoint \exists_{π_Y} and using the predicates $\phi(x, y)$ and $\psi(y, z)$ to denote the elements $\phi \in P(X \times Y)$ and $\psi \in P(Y \times Z)$. Given the well-established correspondence between a doctrine and its internal language, proving that the two previous predicates are equivalent using the logical rules of first-order classical logic is equivalent to prove that they are equal in P . Hereafter we thus prove

$$\exists y.(\phi(x, y) \wedge \neg\psi(y, z)) = \neg\exists y'.(\phi(x, y') \wedge \psi(y', z))$$

Therefore, we start by proving

$$\exists y'.(\phi(x, y') \wedge \neg\psi(y', z)) \vdash \neg\exists y.(\phi(x, y) \wedge \psi(y, z)) \quad (12)$$

The crucial point to conclude that (12) holds is to show that

$$\phi(x, y) \wedge \psi(y, z) \vdash \neg\phi(x, y') \vee \psi(y', z). \quad (13)$$

holds. In fact, from (13) we can deduce the validity of (12) because (13) implies that

$$\exists y.(\phi(x, y) \wedge \psi(y, z)) \vdash \neg\phi(x, y') \vee \psi(y', z)$$

and hence, by applying to both sides \neg , we obtain

$$\phi(x, y') \wedge \neg\psi(y', z) \vdash \neg\exists y.(\phi(x, y) \wedge \psi(y, z))$$

and hence that

$$\exists y'.(\phi(x, y') \wedge \neg\psi(y', z)) \vdash \neg\exists y.(\phi(x, y) \wedge \psi(y, z))$$

So we have to prove the validity of (13). Now since we are working with boolean algebras, we have that $\phi(x, y) \wedge \psi(y, z) = \phi(x, y) \wedge \psi(y, z) \wedge (y = y' \vee y \neq y')$. Therefore, (13) is equivalent to

$$(\phi(x, y) \wedge \psi(y, z) \wedge y = y') \vee (\phi(x, y) \wedge \psi(y, z) \wedge y \neq y') \vdash \neg\phi(x, y') \vee \psi(y', z). \quad (14)$$

To prove (14), it is enough to prove that both

$$(\phi(x, y) \wedge \psi(y, z) \wedge y = y') \vdash \neg\phi(x, y') \vee \psi(y', z) \quad (15)$$

and

$$(\phi(x, y) \wedge \psi(y, z) \wedge y \neq y') \vdash \neg\phi(x, y') \vee \psi(y', z) \quad (16)$$

hold. Now notice that (16) holds trivially, because $\phi(x, y) \wedge \psi(y, z) \wedge y = y' \vdash \psi(y', z)$. To prove (16) we have to employ the functionality of ϕ . In fact, by definition of functionality we have that

$$\phi(x, y) \wedge \phi(x, y') \vdash y = y'$$

and then, using the fact that $y = y'$ is equivalent to $(y \neq y') \rightarrow \perp$ (were $y \neq y'$ denotes $\neg(y = y')$) we can conclude that

$$\phi(x, y) \wedge y \neq y' \vdash \neg\phi(x, y') \quad (17)$$

and hence that $\phi(x, y) \wedge y \neq y' \vdash \neg\phi(x, y') \vee \psi(y', z)$. Therefore, since $\phi(x, y) \wedge \psi(y, z) \wedge y \neq y' \vdash \phi(x, y) \wedge y \neq y'$, we can conclude by transitivity that also (16) holds.

This concludes the proof that (13), and hence, (12) hold.

Now we have to prove that

$$\neg\exists y.(\phi(x, y) \wedge \psi(y, z)) \vdash \exists y'.(\phi(x, y') \wedge \neg\psi(y', z)) \quad (18)$$

First, notice that (18) is equivalent to

$$\forall y'.(\neg\phi(x, y') \vee \psi(y', z)) \vdash \exists y.(\phi(x, y) \wedge \psi(y, z)). \quad (19)$$

Now, in order to prove (19), we start by employing the assumption that ϕ is total, namely:

$$\top \vdash \exists y'.\phi(x, y'). \quad (20)$$

In fact, (20) implies that

$$\forall y'. \neg\phi(x, y') \vdash \perp$$

and since in every boolean algebra we have that $\psi(y', z) \wedge \neg\psi(y', z) \vdash \perp$, we have that

$$\forall y'. \neg\phi(x, y') = \forall y'. (\neg\phi(x, y') \vee \perp) = \forall y'. (\neg\phi(x, y') \vee (\psi(y', z) \wedge \neg\psi(y', z))) \vdash \perp \quad (21)$$

From (21), using the distributivity of \vee , and the fact that the universal quantifier of a disjunction is equivalent to the disjunction of two universal quantifiers (categorically, right adjoints preserve coproducts), we can conclude that

$$\forall y'. (\neg\phi(x, y') \vee \psi(y', z)) \wedge \forall y. (\neg\phi(x, y) \wedge \neg\psi(y, z)) \vdash \perp$$

and hence that

$$\forall y'. (\neg\phi(x, y') \vee \psi(y', z)) \vdash \neg\forall y. (\neg\phi(x, y) \wedge \neg\psi(y, z)).$$

Now, since $\neg\forall y. (\neg\phi(x, y) \wedge \neg\psi(y, z)) = \exists y. (\phi(x, y) \wedge \psi(y, z))$ we can conclude that (19), and then (18) hold. \blacktriangleleft

C Appendix to Section 4

In Section 4 we have recalled the adjunction between the category of cartesian bicategories \mathbf{CB} and that of elementary and existential doctrines \mathbf{EED} , i.e. Equation (7), from [7]. The interested reader may find all details in Sections 5, 6 and 7 of [7] however, for its convenience, we recall in this appendix some interesting facts that are omitted in the main text.

We start by recalling some details about the $\mathbf{Rel}(-)$ construction. The objects, arrows and composition of the category $\mathbf{Rel}(P)$ associated to an elementary and existential doctrine $P: \mathbf{C}^{\text{op}} \rightarrow \mathbf{InfSI}$ are described in Section 4. To see why this is a cartesian bicategory it is convenient to first illustrate the monoidal product. For $\phi: X \rightarrow Y$ and $\psi: U \rightarrow V$ arrows of $\mathbf{Rel}(P)$, namely $\phi \in P(X \times Y)$ and $\psi \in P(U \times V)$, the arrow $\phi \otimes \psi: X \otimes U \rightarrow Y \otimes V$ is defined as

$$\phi \otimes \psi \stackrel{\text{def}}{=} P_{\langle \pi_X, \pi_Y \rangle}(\phi) \wedge P_{\langle \pi_U, \pi_V \rangle}(\psi)$$

where $\langle \pi_X, \pi_Y \rangle$ and $\langle \pi_U, \pi_V \rangle$ are the projections from $X \times U \times Y \times V$ to, respectively, $X \times Y$ and $U \times V$.

The rest of the structure of cartesian bicategory is inherited from the finite products base category \mathbf{C} of P by means of the *graph functor* $\Gamma_P: \mathbf{C} \rightarrow \mathbf{Rel}(P)$. In particular, the graph functor acts as the identity on objects and mapping each arrow $f: X \rightarrow Y$ in

$$\Gamma_P(f) \stackrel{\text{def}}{=} P_{f \times \text{id}_Y}(\delta_Y) \in P(X \times Y) = \mathbf{Rel}(P)[X, Y].$$

For instance, for the doctrine $\mathcal{P}: \mathbf{Set}^{\text{op}} \rightarrow \mathbf{InfSI}$ from Example 15, the functor $\Gamma_{\mathcal{P}}: \mathbf{Set} \rightarrow \mathbf{Rel}^{\circ}$ maps every function f to its graph.

At this point, it is worth to observe that the arrow $P_{\text{id}_Y \times f}(\delta_Y)$ is right adjoint to $\Gamma_P(f)$ (see [7, Pro. 23]) and thus $\Gamma_P(f)$ is a map. Therefore, Γ_P restricts to a finite-product preserving functor $\mathbf{C} \rightarrow \mathbf{Map}(\mathbf{Rel}(P))$. One can thus define the (co)monoid structure of $\mathbf{Rel}(P)$ as

$$\begin{array}{ll} \blacktriangleleft_X^{\circ} \stackrel{\text{def}}{=} \Gamma_P(\blacktriangleleft_X^{\circ}) & \blacktriangleright_X^{\circ} \stackrel{\text{def}}{=} P_{\text{id}_{X \times X} \times \blacktriangleleft_X^{\circ}}(\delta_{X \times X}) \\ !_X^{\circ} \stackrel{\text{def}}{=} \Gamma_P(!_X^{\circ}) & !_X^{\circ} \stackrel{\text{def}}{=} P_{\text{id}_I \times !_X^{\circ}}(\delta_I) \end{array}$$

where on the right hand side of the above equation $\blacktriangleleft_X^\circ: X \rightarrow X \times X$ and $!_X^\circ: X \rightarrow I$ are copier and discard of \mathbf{C} (they exist since \mathbf{C} has finite products).

The graph functor $\Gamma_P: \mathbf{C} \rightarrow \text{Map}(\text{Rel}(P))$ is also used for defining the unit of the adjunction (7). For every elementary and existential doctrine $P: \mathbf{C}^{\text{op}} \rightarrow \text{InfSI}$, the morphism of elementary and existential doctrines $\eta_P: P \rightarrow \text{Hml}(\text{Rel}(P))$ is

$$\begin{array}{ccc}
 \mathbf{C}^{\text{op}} & \xrightarrow{P} & \text{InfSI} \\
 \downarrow \Gamma_P^{\text{op}} & \searrow \rho & \uparrow \text{Rel}(P)[- , 1] \\
 \text{Map}(\text{Rel}(P))^{\text{op}} & &
 \end{array} \quad (22)$$

where

- $\Gamma_P: \mathbf{C} \rightarrow \text{Map}(\text{Rel}(P))$ is the graph-functor;
- each component $\rho_X: P(X) \rightarrow \text{Rel}(P)[X, I] \stackrel{\text{def}}{=} P(X \times I)$ is given by the isomorphism $P(X) \cong P(X \times I)$ obtained by applying the functor P to the right unitor $X \times I \cong X$ in \mathbf{C} .

D Appendix to Section 5

In this appendix we prove that the axioms of fo-bicategories hold in peircean bicategories. In the end we show the isomorphism $\text{FOB} \equiv \text{PB}$ (Theorem 26). Before the main results, we need to prove a few useful properties of peircean bicategories.

Since most of the proofs in this appendix are diagrammatic, it is worth remarking that negation behaves graphically as a *colour switch*. Thus, for example $\neg(x \bullet \frac{X}{X}) = x \bullet \frac{X}{X}$ and $\neg(x \bullet \frac{\bullet}{\bullet}) = x \bullet \frac{\bullet}{\bullet}$. For a generic arrow \boxed{c} , we depict its negation as \boxed{c} .

Moreover, it is convenient to visualize in diagrams $(\neg M)$ on two particular cases, namely when we take as map \blacktriangleleft° or $!^\circ$:

$$\begin{array}{cc}
 \boxed{\bullet} \boxed{c} = \boxed{c} \bullet & \boxed{\bullet} \boxed{c} = \bullet \boxed{c}
 \end{array}$$

► **Lemma 52.** For all $c, d: X \rightarrow Y$, $c \vee d = \blacktriangleleft_X^\circ ; (c \otimes d) ; \blacktriangleright_Y^\circ$ and $\perp = !_X^\circ ; i_Y^\circ$, graphically rendered as follows

$$c \vee d = X \bullet \begin{array}{c} \boxed{c} \\ \boxed{d} \end{array} \bullet Y \quad \perp = X \bullet \bullet \bullet Y$$

Proof.

$ \begin{aligned} c \vee d &= \neg(\neg c \wedge \neg d) && \text{(De Morgan)} \\ &= \neg(\blacktriangleleft_X^\circ ; (\neg c \otimes \neg d) ; \blacktriangleright_Y^\circ) && (5) \\ &= \neg(\neg \blacktriangleleft_X^\circ ; \neg(c \otimes d) ; \neg \blacktriangleright_Y^\circ) && (9) \\ &= \blacktriangleleft_X^\circ ; (c \otimes d) ; \blacktriangleright_Y^\circ && (9) \end{aligned} $	$ \begin{aligned} \perp &= \neg \top && \text{(De Morgan)} \\ &= \neg(!_X^\circ ; i_Y^\circ) && (5) \\ &= \neg(\neg !_X^\circ ; \neg i_Y^\circ) && (9) \\ &= !_X^\circ ; i_Y^\circ && (9) \end{aligned} $
---	--

Given that $(\mathbf{C}, \blacktriangleleft^\bullet, \blacktriangleright^\bullet)$ is a cocartesian bicategory, there is an isomorphism $(\cdot)^\dagger: \mathbf{C} \rightarrow \mathbf{C}^{\text{op}}$ that is the identity on objects and for all arrows $c: X \rightarrow Y$, $c^\dagger: Y \rightarrow X$ is defined as follows.

$$c^\dagger \stackrel{\text{def}}{=} \begin{array}{c} X \\ \boxed{\begin{array}{c} \bullet \quad \bullet \\ \text{---} c \text{---} \\ \bullet \quad \bullet \end{array}} \\ Y \end{array} \quad (23)$$

With this definition at hand it is immediate to show that $\neg(c^\dagger) = (\neg c)^\dagger$. Moreover, we show that the two isomorphisms $(\cdot)^\dagger$ and $(\cdot)^\ddagger$ actually coincide (Lemma 57).

► **Proposition 53.** *For any $c, d: X \rightarrow Y$ the following are equivalent:*

$$1. c \leq d \quad 2. \neg d \leq \neg c \quad 3. \top \leq \neg c \vee d \quad 4. c \wedge \neg d \leq \perp$$

Proof. The four inclusions are equivalent since $\mathbf{C}[X, Y]$ is a Boolean algebra. ◀

► **Proposition 54.** *The following equality holds $x \begin{array}{c} \bullet \quad \bullet \\ \text{---} \end{array} y = x \begin{array}{c} \bullet \quad \bullet \\ \boxed{} \end{array} y$*

Proof. We prove the two inclusions separately. The \leq inclusion is trivial since $x \begin{array}{c} \bullet \quad \bullet \\ \text{---} \end{array} y$ is $\perp_{X,Y}$. For the other inclusion observe that $x \begin{array}{c} \bullet \quad \bullet \\ \text{---} \end{array} y \stackrel{(\neg M)}{=} x \begin{array}{c} \bullet \quad \bullet \\ \boxed{} \end{array} y$ and thus what is left to prove is $x \begin{array}{c} \bullet \quad \bullet \\ \boxed{} \end{array} y \leq x \begin{array}{c} \bullet \quad \bullet \\ \boxed{} \end{array} y$. We prove it by means of Lemma 45 as follows:

$$x \begin{array}{c} \bullet \quad \bullet \\ \boxed{} \end{array} y \stackrel{(\triangleright^\circ\text{-un})}{=} x \begin{array}{c} \bullet \quad \bullet \\ \boxed{} \end{array} y \approx x \begin{array}{c} \bullet \quad \bullet \\ \boxed{} \end{array} y \stackrel{(\neg M)}{=} x \begin{array}{c} \bullet \quad \bullet \\ \text{---} \end{array} y \leq x \begin{array}{c} \bullet \quad \bullet \\ \boxed{} \end{array} y$$

where the last inequality holds since the left-hand side is $\perp_{X \otimes Y, I}$. ◀

► **Lemma 55.** *The following equality holds, $(\perp_{X,Y})^\dagger = \perp_{Y,X}$.*

Proof.

$$\begin{aligned} (\perp_{X,Y})^\dagger &\stackrel{(6)}{=} \begin{array}{c} X \\ \boxed{\begin{array}{c} \bullet \quad \bullet \\ \text{---} \end{array}} \\ Y \end{array} \stackrel{\text{Proposition 54}}{=} \begin{array}{c} X \\ \boxed{\begin{array}{c} \bullet \quad \bullet \\ \text{---} \end{array}} \\ Y \end{array} \stackrel{(\neg M)}{=} \begin{array}{c} X \\ \boxed{\begin{array}{c} \bullet \quad \bullet \\ \text{---} \end{array}} \\ Y \end{array} \\ &\stackrel{(\blacktriangleleft^\circ\text{-un})}{=} \begin{array}{c} X \\ \boxed{\begin{array}{c} \bullet \quad \bullet \\ \text{---} \end{array}} \\ Y \end{array} \stackrel{(\blacktriangleright^\circ\text{-un})}{=} \begin{array}{c} X \\ \boxed{\begin{array}{c} \bullet \quad \bullet \\ \text{---} \end{array}} \\ Y \end{array} \approx \begin{array}{c} X \\ \boxed{\begin{array}{c} \bullet \quad \bullet \\ \text{---} \end{array}} \\ Y \end{array} \stackrel{(\neg M)}{=} \begin{array}{c} X \\ \boxed{\begin{array}{c} \bullet \quad \bullet \\ \text{---} \end{array}} \\ Y \end{array} \\ &\stackrel{\text{Proposition 54}}{=} \begin{array}{c} X \\ \boxed{\begin{array}{c} \bullet \quad \bullet \\ \text{---} \end{array}} \\ Y \end{array} \stackrel{\text{Lemma 52}}{=} \perp_{Y,X} \end{aligned}$$

► **Lemma 56.** *For any $c: X \rightarrow Y$, $(\neg c)^\dagger = \neg(c^\dagger)$.*

Proof. We prove the two inclusions separately. We prove \leq , on the left, by means of Proposition 53.4. For \geq , on the right, we use the properties of $(\cdot)^\dagger$ in Table 1 and the inclusion proved on the left.

$\begin{aligned} (\neg c)^\dagger \wedge \neg(c^\dagger) &= (\neg c)^\dagger \wedge c^\dagger && \text{(Definition 25.1)} \\ &= (\neg c \wedge c)^\dagger && \text{(Table 1)} \\ &= (\perp_{X,Y})^\dagger && \text{(Definition 25.1)} \\ &= \perp_{Y,X} && \text{(Lemma 55)} \end{aligned}$	$\begin{aligned} \neg(c^\dagger) &= \neg(c^\dagger)^{\dagger\dagger} && \text{(Table 1)} \\ &\leq \neg c^{\dagger\dagger\dagger} && ((\neg c)^\dagger \leq \neg(c^\dagger)) \\ &= (\neg c)^\dagger && \text{(Table 1)} \end{aligned}$
---	---

$$c^\dagger \stackrel{\text{Definition 25.1}}{=} (\neg\neg c)^\dagger \stackrel{\text{Lemma 56}}{=} \neg((\neg c)^\dagger) \stackrel{(23)}{=} (\neg\neg c)^\dagger \stackrel{\text{Definition 25.1}}{=} c^\dagger$$

Diagrammatic proof of Proposition 47. The sequence of diagrams shows the derivation of the equation $Z[c]fX = Z[f]cX$ using the following steps:

- Initial expression: $Z[c]fX$
- Step (6): $Z[c]fX \equiv$ (Diagram: Z connected to a box containing c and f , which is then connected to X).
- Proposition 47: $Z[c]fX \equiv$ (Diagram: Z connected to a box containing c and f , which is then connected to X).
- Step $(\neg M)$: $Z[c]fX \equiv$ (Diagram: Z connected to a box containing f and c , which is then connected to X).
- Lemma 57: $Z[c]fX \equiv$ (Diagram: Z connected to a box containing f and c , which is then connected to X).
- Step (23): $Z[c]fX \equiv$ (Diagram: Z connected to a box containing c and f , which is then connected to X).
- Final expression: $Z[f]cX$

The diagram shows two equations. The first equation is: a box labeled 'C' with an input line from the left and an output line to the right, followed by a dot operator (a small black circle), is equal to a dot operator followed by a box labeled 'C'. The second equation is: a dot operator followed by a box labeled 'C' is equal to a box labeled 'C' followed by a dot operator.

Figure 1 displays a collection of 10 string diagrams representing axioms and theorems in the rewriting theory of the lambda calculus. The diagrams are arranged in a grid. The first row shows the axiom of abstraction (Lemma 52) and the axiom of application (Theorem 43). The second row shows the axiom of eta-reduction (Definition 25.1) and the axiom of beta-reduction (Theorem 43). The third row shows the axiom of eta-reduction (Definition 25.1) and the axiom of beta-reduction (Theorem 43). The fourth row shows the axiom of eta-reduction (Definition 25.1) and the axiom of beta-reduction (Theorem 43). The fifth row shows the axiom of eta-reduction (Definition 25.1) and the axiom of beta-reduction (Theorem 43).

► **Lemma 60.** For any $c: X \rightarrow Y$, $\boxed{X \atop Y} \text{c} = \neg \left(\boxed{X \atop Y} \text{c} \right)$.

Proof. We prove the two inclusions separately.

We prove \leq by means of Proposition 53.4 as follows.

$$\begin{array}{c}
 \begin{array}{c} X \\ Y \end{array} \boxed{c} \wedge \neg \neg \left(\begin{array}{c} X \\ Y \end{array} \boxed{c} \right) \stackrel{\text{Definition 25.1}}{=} \begin{array}{c} X \\ Y \end{array} \boxed{c} \wedge \begin{array}{c} X \\ Y \end{array} \boxed{c} \stackrel{(5)}{=} \begin{array}{c} X \\ Y \end{array} \boxed{\begin{array}{c} c \\ c \end{array}} \\
 \stackrel{\text{Theorem 43}}{=} \begin{array}{c} X \\ Y \end{array} \boxed{\begin{array}{c} c \\ c \end{array}} \stackrel{\text{Definition 25.1}}{=} \begin{array}{c} X \\ Y \end{array} \boxed{\begin{array}{c} \bullet \\ \bullet \end{array}} \\
 \stackrel{\text{Lemma 58}}{=} \begin{array}{c} X \\ Y \end{array} \boxed{\bullet} \stackrel{(\triangleright^\circ\text{-un})}{=} \begin{array}{c} X \\ Y \end{array} \boxed{\bullet} \\
 \approx \begin{array}{c} X \\ Y \end{array} \boxed{\bullet} \stackrel{(\neg M)}{=} \begin{array}{c} X \\ Y \end{array} \boxed{\bullet} \\
 \stackrel{\text{Lemma 52}}{=} \perp_{X \otimes Y, I}
 \end{array}$$

For \geq we proceed as follows.

$$\begin{array}{c}
 \neg \left(\begin{array}{c} X \\ Y \end{array} \boxed{c} \right) = \begin{array}{c} X \\ Y \end{array} \boxed{c} \stackrel{\text{Theorem 43}}{=} \begin{array}{c} X \\ Y \end{array} \boxed{\begin{array}{c} c \\ c \end{array}} \stackrel{\text{Proposition 59}}{\leq} \begin{array}{c} X \\ Y \end{array} \boxed{\begin{array}{c} c \\ c \end{array}} \\
 \stackrel{\text{Theorem 43}}{=} \begin{array}{c} X \\ Y \end{array} \boxed{c}
 \end{array}$$

► **Lemma 61.** For any $c: X \rightarrow Y$, $\begin{array}{c} X \\ Y \end{array} \boxed{c} = \begin{array}{c} X \\ Y \end{array} \boxed{\begin{array}{c} c \\ c \end{array}}$.

Proof.

$$\begin{array}{c} X \\ Y \end{array} \boxed{c} \stackrel{\text{Definition 25.1}}{=} \neg \neg \left(\begin{array}{c} X \\ Y \end{array} \boxed{c} \right) \stackrel{\text{Lemma 60}}{=} \neg \left(\begin{array}{c} X \\ Y \end{array} \boxed{\begin{array}{c} c \\ c \end{array}} \right) = \begin{array}{c} X \\ Y \end{array} \boxed{c}$$

Now we need to prove that the axioms of fo-bicategories hold in a peircean bicategory. We do not show a proof for all of them, but only for a few representative ones. The rest are either derivable or proved in a similar manner.


► **Proposition 62.** For any $c, d, e: X \rightarrow Y$, $c \wedge (d \vee e) \leq (c \wedge d) \vee e$.

Proof.


$$\begin{aligned}
 c \wedge (d \vee e) &= (c \wedge d) \vee (c \wedge e) && \text{(Definition 25.1)} \\
 &\leq (c \wedge d) \vee (\top \wedge e) && \text{(Proposition 3)} \\
 &= (c \wedge d) \vee e && \text{(Definition 25.1)}
 \end{aligned}$$

► **Lemma 63.** For any $c: X \rightarrow I, d: Y \rightarrow I, e: Z \rightarrow I$, $c \otimes (d \otimes e) \leq (c \otimes d) \otimes e$.


Proof.



► **Lemma 64.** For any $c: X \otimes Y \rightarrow I, d: Y \otimes Z \rightarrow I, e: Z \otimes W \rightarrow I$, the following inequality holds



Proof.




◀

► **Lemma 65** (Linear distributivities). *For any $c: X \rightarrow Y, d: Y \rightarrow Z, e: Z \rightarrow W$, the following inequalities hold*

$$1. c \circ (d \circ e) \leq (c \circ d) \circ e \quad 2. (c \circ d) \circ e \leq c \circ (d \circ e)$$


Proof. We prove 1. below by means of Lemma 45. The proof for 2. is analogous.



► **Proposition 66.** *For any $c: X \rightarrow Y$, the following inequalities hold*

$$1. \begin{array}{c} X \\ \boxed{c} \\ X \end{array} \leq \begin{array}{c} X \\ \boxed{\bullet} \\ X \end{array} \quad 2. \begin{array}{c} X \\ \boxed{\bullet} \\ X \end{array} \leq \begin{array}{c} X \\ \boxed{c} \\ X \end{array}$$


Proof. We prove 1. below. The proof for 2. is analogous.




► **Lemma 67** (Linear adjoints). *For any $c: X \rightarrow Y$, the following inequalities hold*

$$1. c \circ (\neg c)^\dagger \leq id_X^\bullet \quad 2. id_Y^\circ \leq (\neg c)^\dagger \circ c \quad 3. (\neg c)^\dagger \circ c \leq id_Y^\bullet \quad 4. id_X^\circ \leq c \circ (\neg c)^\dagger$$


Proof. We prove 1. below. 2. is proved similarly, exploiting Proposition 66.2. The proofs for 3. and 4. are analogous.



► **Lemma 68** (Linear strengths I). *For any a, b, c, d properly typed, the following equalities hold*



Proof. We prove 1. by means of Proposition 53.4 below. The proof for 2. is analogous. 3. and 4. follow from 1. and 2. and Proposition 53.2.



$$\begin{aligned}
& \leq \begin{array}{c} X \\ Z \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \bullet \\ \hline \end{array} \begin{array}{c} Y \\ W \end{array} \quad (!^\circ\text{-nat}) \\
& = \begin{array}{c} X \\ Z \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \bullet \\ \hline \end{array} \begin{array}{c} Y \\ W \end{array} \quad (\text{Lemma 58}) \\
& = \perp_{X \otimes Z, Y \otimes W} \quad (\text{Lemma 52})
\end{aligned}$$

► **Proposition 69.** For any $c, d: X \rightarrow Y$, $c \wedge d \leq c \vee d$.

Proof. The following holds since $\mathbf{C}[X, Y]$ is a Boolean algebra and a \wedge -semilattice with \top :

$$\begin{aligned}
c \wedge d &= c \wedge (d \vee d) && (\text{Idempotency of } \vee) \\
&\leq (c \wedge d) \vee d && (\text{Proposition 62}) \\
&\leq (c \wedge \top) \vee d && (\top \text{ is the top element}) \\
&= c \vee d && (\top \text{ is the unit of } \wedge)
\end{aligned}$$

► **Lemma 70.** For any $c: X \rightarrow Y, d: Z \rightarrow W$, $c \otimes d \leq c \otimes d$.

Proof.

$$\begin{aligned}
& \begin{array}{c} X \\ Z \end{array} \begin{array}{|c|} \hline c \\ \hline d \\ \hline \end{array} \begin{array}{c} Y \\ W \end{array} \stackrel{(\leftarrow^\circ\text{-un})}{=} \begin{array}{c} X \\ Z \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \bullet \\ \hline \end{array} \begin{array}{c} Y \\ W \end{array} \stackrel{(5)}{=} \begin{array}{c} X \\ Z \end{array} \begin{array}{|c|} \hline c \\ \hline \bullet \\ \hline \end{array} \begin{array}{c} Y \\ W \end{array} \wedge \begin{array}{c} X \\ Z \end{array} \begin{array}{|c|} \hline \bullet \\ \hline d \\ \hline \end{array} \begin{array}{c} Y \\ W \end{array} \\
& \stackrel{\text{Proposition 69}}{\leq} \begin{array}{c} X \\ Z \end{array} \begin{array}{|c|} \hline c \\ \hline \bullet \\ \hline \end{array} \begin{array}{c} Y \\ W \end{array} \vee \begin{array}{c} X \\ Z \end{array} \begin{array}{|c|} \hline \bullet \\ \hline d \\ \hline \end{array} \begin{array}{c} Y \\ W \end{array} \stackrel{\text{Lemma 52}}{=} \begin{array}{c} X \\ Z \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \bullet \\ \hline \end{array} \begin{array}{c} Y \\ W \end{array} \\
& \stackrel{(\neg M)}{=} \begin{array}{c} X \\ Z \end{array} \begin{array}{|c|} \hline c \\ \hline d \\ \hline \end{array} \begin{array}{c} Y \\ W \end{array} \stackrel{\text{Lemma 58}}{\approx} \begin{array}{c} X \\ Z \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \bullet \\ \hline \end{array} \begin{array}{c} Y \\ W \end{array} \stackrel{(\leftarrow^\circ\text{-un})}{=} \begin{array}{c} X \\ Z \end{array} \begin{array}{|c|} \hline c \\ \hline d \\ \hline \end{array} \begin{array}{c} Y \\ W \end{array}
\end{aligned}$$

► **Corollary 71** (Linear strenghts II). For all objects X, Y , the following inequalities hold


$$id_X^\bullet \otimes id_Y^\bullet \leq id_X^\bullet \otimes id_Y^\bullet \quad id_X^\circ \otimes id_Y^\circ \leq id_X^\circ \otimes id_Y^\circ$$

Proof. Immediate by Lemma 70.


► **Proposition 72.** The following equality holds

$$\begin{array}{c} X \\ X \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \bullet \\ \hline \end{array} \begin{array}{c} X \\ X \end{array} = \begin{array}{c} X \\ X \end{array} \begin{array}{|c|} \hline \bullet \\ \hline \bullet \\ \hline \end{array} \begin{array}{c} X \\ X \end{array}$$


Proof. We prove it by means of Lemma 45 as follows.



► **Lemma 73** (Linear Frobenius). *The following equalities hold*



Proof. We prove 1. below. The proof for 2. is analogous. 3. and 4. follow from 1. and 2. and Proposition 53.2.



Proof of Theorem 26. By Propositions 7 and 9 every fo-bicategory is a peircean bicategory. By Proposition 8 every morphism of fo-bicategories is a morphism of peircean bicategories.

Now observe that a peircean bicategory \mathbf{C} is a fo-bicategory, since:

- it is a cartesian bicategory by definition;
- it is a cocartesian bicategory via the isomorphism $\neg: (\mathbf{C}^{\text{co}}, \blacktriangleleft^{\circ}, \blacktriangleright^{\circ}) \rightarrow (\mathbf{C}, \blacktriangleleft^{\bullet}, \blacktriangleright^{\bullet})$;
- it is a closed linear bicategory, since:
 - (δ_l) and (δ_r) hold by Lemma 65;
 - $(\nu_l^{\circ}), (\nu_r^{\circ}), (\nu_l^{\bullet})$ and (ν_r^{\bullet}) hold by Lemma 68 and (\otimes°) and (\otimes^{\bullet}) hold by Corollary 71;
 - every arrow $c: X \rightarrow Y$, and in particular also $\sigma^{\circ}, \sigma^{\bullet}, \blacktriangleleft^{\circ}, \blacktriangleleft^{\bullet}, \blacktriangleright^{\circ}, \blacktriangleright^{\bullet}, i^{\circ}, i^{\bullet}$, has both a left and right linear adjoint by Lemma 67;
- $(F^{\circ}), (F^{\bullet}), (F^{\circ})$ and (F^{\bullet}) hold by Lemma 73.



A morphism of peircean bicategories $F: \mathbf{C} \rightarrow \mathbf{D}$ preserves the structure defined in (9) since it preserves negation, e.g.


$$\begin{aligned}
 F(c \circ d) &= F(\neg(\neg c \circ \neg d)) & (9) \\
 &= \neg F(\neg c \circ \neg d) & (F \text{ preserves negation}) \\
 &= \neg(F(\neg c) \circ F(\neg d)) & (F \text{ is a strong symmetric monoidal functor}) \\
 &= \neg(\neg F(c) \circ \neg F(d)) & (F \text{ preserves negation}) \\
 &= F(c) \circ F(d) & (9)
 \end{aligned}$$

In particular, F is both a morphism of cartesian and cocartesian bicategories and thus a morphism of fo-bicategories. \blacktriangleleft

E Appendix to Section 6

► **Lemma 74.** Let \mathbf{C} be a cartesian bicategory, and consider the doctrine $\mathbf{C}[-, I]: \mathbf{Map}(\mathbf{C})^{\text{op}} \rightarrow \mathbf{InfSI}$. Then an element $c \in \mathbf{C}[X \times Y, I]$ is:

- functional if and only if the corresponding arrow in $\mathbf{C}[X, Y]$, i.e. , satisfies the leftmost inequality in (4);
- entire if and only if the corresponding arrow in $\mathbf{C}[X, Y]$, i.e. , satisfies the rightmost inequality in (4);

Therefore, c is both functional and entire if and only if  is a map.

Proof. By definition, an element $c \in \mathbf{C}[X \times Y, I]$ is functional from X to Y if and only if

$$\begin{array}{c} X \\ Y \\ Y \end{array} \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \begin{array}{c} \boxed{c} \\ \boxed{c} \end{array} \leq \begin{array}{c} X \\ Y \\ Y \end{array} \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \begin{array}{c} \boxed{c} \end{array} \quad (\text{functional})$$

and it is entire from X to Y if and only if

$$\begin{array}{c} X \\ \text{---} \end{array} \begin{array}{c} \boxed{c} \end{array} \leq \begin{array}{c} X \\ \text{---} \end{array} \begin{array}{c} \boxed{c} \end{array} \quad (\text{entire})$$

Now observe that (functional) holds if and only if the following holds


$$\begin{array}{c} X \\ Y \\ Y \end{array} \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \begin{array}{c} \boxed{c} \\ \boxed{c} \end{array} \leq \begin{array}{c} X \\ Y \\ Y \end{array} \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \begin{array}{c} \boxed{c} \end{array} \quad (\text{functional-alt})$$


To conclude, we need to show that (functional-alt) and (entire) are equivalent, respectively, to

$$\begin{array}{c} X \\ \text{---} \end{array} \begin{array}{c} \boxed{c} \\ \boxed{c} \end{array} \leq \begin{array}{c} X \\ \text{---} \end{array} \begin{array}{c} \boxed{c} \end{array} \quad \text{and} \quad \begin{array}{c} X \\ \text{---} \end{array} \begin{array}{c} \boxed{c} \end{array} \leq \begin{array}{c} X \\ \text{---} \end{array} \begin{array}{c} \boxed{c} \end{array}.$$

For (functional-alt) we use the fact that in any cartesian bicategory $\begin{array}{c} X \\ Y \end{array} \boxed{a} \leq \begin{array}{c} X \\ Y \end{array} \boxed{b}$ if and only if $\begin{array}{c} X \\ \text{---} \end{array} \boxed{a} \leq \begin{array}{c} X \\ \text{---} \end{array} \boxed{b}$, and thus

$$\begin{array}{c} X \\ Y \\ Y \end{array} \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \begin{array}{c} \boxed{c} \\ \boxed{c} \end{array} \leq \begin{array}{c} X \\ Y \\ Y \end{array} \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \begin{array}{c} \boxed{c} \end{array} \quad \text{iff} \quad \begin{array}{c} X \\ \text{---} \end{array} \begin{array}{c} \boxed{c} \\ \boxed{c} \end{array} \leq \begin{array}{c} X \\ \text{---} \end{array} \begin{array}{c} \boxed{c} \end{array}$$

which amounts to  using Theorem 43.

For (entire) it suffices to observe that .

Proof of Proposition 29. By Equation (7) we have that for every elementary and existential doctrine P , the unit of the adjunction $\mathbf{Rel}(-) \dashv \mathbf{Hml}(-)$ is morphism of elementary and existential doctrines $\eta_P \stackrel{\text{def}}{=} (\Gamma_P, \rho): P \rightarrow \mathbf{Hml}(\mathbf{Rel}(P))$. Moreover, each component of ρ is an iso by definition (see Appendix C for a detailed description). Therefore, ρ preserves and reflects functional and entire relations (see Remark 18), i.e. $\rho_{X \times Y}(\phi) \in \mathbf{Rel}(P)[X \times Y, I]$ is functional and entire in $\mathbf{Rel}(P)$ if and only if ϕ is functional and entire in P . On the other hand, by Lemma 74 we have that an arrow $\psi \in \mathbf{Rel}(P)[X, Y]$, i.e. an element $\psi \in P(X \times Y)$, is a map if and only if the corresponding element in $\mathbf{Rel}(P)[A \times B, I]$, i.e. $\rho_{X \times Y}(\psi)$, is functional and entire with respect to the doctrine $\mathbf{Rel}(P)[- , I]$. Therefore, we can conclude that ϕ is a map in $\mathbf{Rel}(P)$ if and only if ϕ is an entire and functional element of P . \blacktriangleleft

Proof of Theorem 31. First, we want to prove that the inclusion $\mathbf{Hml}: \mathbf{CB} \hookrightarrow \mathbf{EED}$ in (7) restricts to an inclusion of categories $\mathbf{PB} \hookrightarrow \mathbf{BHD}$. By Proposition 27, one only needs to check for morphisms in \mathbf{PB} . Given a morphism of peircean bicategories $F: \mathbf{C} \rightarrow \mathbf{D}$, $\mathbf{Hml}(F)$ is the morphism of elementary and existential doctrines (\tilde{F}, b^F) defined in Section 4. In order to conclude that it is a morphism of boolean doctrines, it is enough to show that b_X^F is a morphism of boolean algebras for all objects X . Since (\tilde{F}, b^F) is a morphism of doctrines, b_X^F is a morphism of inf-semilattices. Thus it is enough to show that b_X^F preserve negation. But this is trivial since, for all $c \in \mathbf{C}[X, I]$,

$$\begin{aligned} b_X^F(\neg c) &= F(\neg c) && \text{(Def. } b^F) \\ &= \neg F(c) && \text{(morphism of Peircean, Definition 25)} \\ &= \neg b_X^F(c) && \text{(Def. } b^F) \end{aligned}$$

Now, to prove that \mathbf{Rel} restrict to a functor $\mathbf{Rel}: \mathbf{BHD} \rightarrow \mathbf{PB}$, by Proposition 30, one only needs to check that for all morphisms of boolean hyperdoctrines $(F, b): P \rightarrow Q$, $\mathbf{Rel}(F, b): \mathbf{Rel}(P) \rightarrow \mathbf{Rel}(Q)$ is a morphism of peircean bicategories. Since by (7), $\mathbf{Rel}(F, b)$ is a morphism of cartesian bicategories, one only needs to check that it preserves the negation. But this is obvious since for all arrows $\phi \in \mathbf{Rel}(P)[X, Y]$, $\mathbf{Rel}(F, b)(\phi)$ is \neg -by definition $b_{X \times Y}(\phi)$ and $b_{X \times Y}$ is a morphism of boolean algebras.

To conclude, one only needs to check the unit and the counit of the adjunction in (7). The counit is an isomorphism of cartesian bicategories (see Equation (9) in [7]), and then it provides an isomorphism of peircean bicategories $\mathbf{C} \cong \mathbf{Rel}(\mathbf{C}[-, I])$ whenever \mathbf{C} is a peircean bicategory. The unit of the adjunction $\eta_P: P \rightarrow \mathbf{Rel}(P)[- , I]$ is the morphism of elementary and existential doctrines (Γ_P, ρ) illustrated in (22). To conclude that η_P is a morphism of boolean hyperdoctrine whenever P is a boolean hyperdoctrine, one has only to prove that ρ is a morphism of boolean algebras, but this is trivial since ρ is always an isomorphism of inf-semilattices. \blacktriangleleft

F

 Appendix to Section 7

Proof of Theorem 36. By Equation (8) we have that the Hml and Rel functors provide an equivalence between the categories $\mathbb{C}\mathbb{B}$ and $\mathbb{E}\mathbb{E}\mathbb{D}$. Now, since every peircean category is in particular a cartesian bicategory, we have that every boolean hyperdoctrine arising from a peircean bicategory satisfies (RUC) and it has comprehensive diagonals. Then, we have that the functor $\text{Hml}: \mathbb{P}\mathbb{B} \hookrightarrow \mathbb{B}\mathbb{H}\mathbb{D}$ factors through the canonical inclusion $\mathbb{B}\mathbb{H}\mathbb{D} \hookrightarrow \overline{\mathbb{B}\mathbb{H}\mathbb{D}}$:

$$\begin{array}{ccc} \mathbb{P}\mathbb{B} & \xrightarrow{\text{Hml}} & \mathbb{B}\mathbb{H}\mathbb{D} \\ & \searrow \text{Hml} & \nearrow \\ & \overline{\mathbb{B}\mathbb{H}\mathbb{D}} & \end{array}$$

By Theorem 31, we have that $\text{Hml}: \mathbb{P}\mathbb{B} \hookrightarrow \mathbb{B}\mathbb{H}\mathbb{D}$ is fully and faithful (since the counit of the adjunction is an iso), so it remains to prove that it is essentially surjective (with respect to the objects of $\overline{\mathbb{B}\mathbb{H}\mathbb{D}}$). By the equivalence presented in Equation (8), we know that every boolean hyperdoctrine (that is in particular an elementary and existential doctrine) satisfying (RUC) and having comprehensive diagonals, is isomorphic to an elementary and existential doctrine $\mathbf{C}[-, I]: \text{Map}(\mathbf{C})^{\text{op}} \rightarrow \text{InfSI}$ for some cartesian bicategory \mathbf{C} . Thus, we can conclude that $\mathbf{C}[-, I]: \text{Map}(\mathbf{C})^{\text{op}} \rightarrow \text{InfSI}$ is a boolean hyperdoctrine and, by Corollary 28, that \mathbf{C} is a peircean bicategory. This concludes the proof that $\mathbb{P}\mathbb{B} \equiv \overline{\mathbb{B}\mathbb{H}\mathbb{D}}$. ◀

G

 Appendix to Section 8

We summarize some useful properties of tabulations, which are crucial to establish the precise connection with the notion of full comprehension. We refer to [12, Lem. 3.3] for the following result:

► **Lemma 75.** *Let $i: X_r \rightarrow X$ be a tabulation of $r: X \rightarrow I$. Then*

- *for every map $f: Z \rightarrow X$ such that $f^+ \circ !_Z^{\circ} \leq r$ there exists a unique map $h: Z \rightarrow X_r$ such that $f = h \circ i$;*
- *if $f^+ \circ !_Z^{\circ} = r$, then $h^+ \circ !_Z^{\circ} = !_X^{\circ}$.*

Now we recall another useful lemma, regarding doctrines with full comprehensions. We refer to [54, Pro. 7.10] for the following result:

► **Lemma 76.** *Let $P: \mathbf{C}^{\text{op}} \rightarrow \text{InfSI}$ be a doctrine with full comprehensions. Then every comprehension is a mono for every element α of $P(X)$ we have that $\alpha = \exists_{[\alpha]}(\tau_{X_\alpha})$.*

Proof of Theorem 41. Let $(\mathbf{C}, \blacktriangleleft^{\circ}, \blacktriangleright^{\circ})$ be functionally complete, and let us consider the doctrine $\mathbf{C}[-, I]: \text{Map}(\mathbf{C})^{\text{op}} \rightarrow \text{InfSI}$ and an element $r: X \rightarrow I$ of $\mathbf{C}[X, I]$. We claim that the tabulation $i: X_r \rightarrow X$ of r is the comprehension of r .

First, notice that $\mathbf{C}[i, I](r) = \tau_{X_r}$ (i.e. $i \circ r = !_X^{\circ}$) because, by definition of functionally completeness, we have that $i \circ i^{\dagger} = \text{id}_{X_r}$ and $i^{\dagger} \circ !_X^{\circ} = r$, and hence $i \circ r = !_X^{\circ}$. Now suppose that $f: Z \rightarrow X$ is a map such that $\mathbf{C}[f, I](r) = \tau_Z$ (i.e. $f \circ r = !_Z^{\circ}$). Then, in particular, we have that $\tau_Z \leq \mathbf{C}[f, I](r)$, and we can conclude that $\exists_f(\tau_{X_r}) \leq r$ because the doctrine $\mathbf{C}[-, I]: \text{Map}(\mathbf{C})^{\text{op}} \rightarrow \text{InfSI}$ is elementary and existential, so it has left adjoints along all the arrows (see Remark 13). Now, by definition of left adjoints for this doctrine (see Example 16)

this means that $f^\dagger \circ !_Z^\circ \leq r$, so we can apply Lemma 75 and conclude that there exists a unique h such that $f = h \circ i$. This concludes the proof that if $(\mathbf{C}, \blacktriangleleft^\circ, \blacktriangleright^\circ)$ is functionally complete then $\mathbf{C}[-, I]: \mathbf{Map}(\mathbf{C})^{\text{op}} \rightarrow \mathbf{InfSI}$ has comprehensions.

Now we need to prove that comprehensions are full. So let $r, r': X \rightarrow I$ be two arrows and let i and i' their tabulation. If i factors through i' , namely $i = g \circ i'$ for some map $g: X_r \rightarrow X_{r'}$, then we have that $i^\dagger = (i')^\dagger \circ g^\dagger$. Hence, we can conclude that $r \leq r'$ because, by hypothesis, $i^\dagger \circ !_{X_r}^\circ = r$, and using the fact that $i^\dagger = (i')^\dagger \circ g^\dagger$, we can conclude that $r = (i')^\dagger \circ g^\dagger \circ !_{X_r}^\circ \leq r'$ (because $(i')^\dagger \circ !_{X_{r'}}^\circ = r'$ and $g^\dagger \circ !_{X_r}^\circ \leq !_{X_{r'}}^\circ$).

Now we show that full comprehensions implies functionally completeness. So, let us consider an arrow $r: X \rightarrow I$. We claim that the comprehension $\{r\}: X_r \rightarrow X$ is a tabulation of r . First, by Lemma 76, we have that $\exists_{\{r\}}(\top_{X_r}) = r$, namely that $\{r\}^\dagger \circ !_{X_r}^\circ = r$. Finally, we have that $\{r\} \circ \{r\}^\dagger = \text{id}_{X_r}$ because comprehensions are monomorphisms in $\mathbf{Map}(\mathbf{C})$. This concludes the proof that $\{r\}$ is the tabulation of r . \blacktriangleleft

The last part of this section is devoted to prove the final corollary, namely that

$$\mathbf{PB} \equiv \overline{\mathbf{BHD}}_c \cong \mathbf{BC}$$

To properly reach this goal, we summarize here in the language of doctrines the main results presented in this work, and some useful characterizations presented in [50]. Our result will follow by combining these results.

First, we have the following result presented in [7, Thm. 35]:

► **Theorem 77.** *Let $P: \mathbf{C}^{\text{op}} \rightarrow \mathbf{InfSI}$ be an elementary and existential doctrine. Then the following two conditions are equivalent:*

- *P has comprehensive diagonals and satisfies (RUC);*
- *$\mathbf{C} \cong \mathbf{Map}(\mathbf{Rel}(P))$ and $P \cong \mathbf{C}[-, I]$.*

In particular, $\mathbf{CB} \equiv \overline{\mathbf{EED}}$.

Combining the previous result with Theorem 41 we obtain the following corollary, where we denote by $\overline{\mathbf{EED}}_c$ the full subcategory of $\overline{\mathbf{EED}}$ given by those doctrines of $\overline{\mathbf{EED}}$ having full comprehensions, and by \mathbf{CB}_f the full subcategory of \mathbf{CB} given by functionally complete cartesian bicategories.

► **Corollary 78.** *Let $P: \mathbf{C}^{\text{op}} \rightarrow \mathbf{InfSI}$ be an elementary and existential doctrine. Then the following two conditions are equivalent:*

- *P has comprehensive diagonals, full comprehension and satisfies (RUC);*
- *$\mathbf{C} \cong \mathbf{Map}(\mathbf{Rel}(P))$, \mathbf{C} is functionally complete and $P \cong \mathbf{C}[-, I]$.*

In particular, $\mathbf{CB}_f \equiv \overline{\mathbf{EED}}_c$.

Notice that Theorem 77 can be seen as a generalization of another result regarding doctrines and regular categories presented in [50, Prop. 5.3], establishing the precise connection between regular categories and doctrines:

► **Theorem 79.** *Let $P: \mathbf{C}^{\text{op}} \rightarrow \mathbf{InfSI}$ be an elementary and existential doctrine. Then the following two conditions are equivalent:*

- *P has comprehensive diagonals, full comprehensions and satisfies (RUC);*
- *\mathbf{C} is regular and $P \cong \mathbf{Sub}_{\mathbf{C}}$.*

In particular, $\mathbb{REG} \equiv \overline{\mathbb{EED}}_c$.

Finally, we can combine the previous result with Corollary 78 obtaining as corollary the well-known equivalence (see [12] and [24]) between functionally complete bicategories and regular categories:

► **Corollary 80.** *We have the equivalences of categories $\mathbb{REG} \equiv \overline{\mathbb{EED}}_c \equiv \mathbb{CB}_f$.*

Since, by definition (see [38, Sec. A1.4, p. 38]), a category \mathbf{C} is boolean if and only if the subobjects functor on \mathbf{C} is a boolean hyperdoctrine, we have the following corollary, that is a particular instance of the previous one. Here we denote the category of boolean categories by \mathbb{BC} and that of boolean hyperdoctrines satisfying (RUC), with full comprehensions and comprehensive diagonals by \mathbb{BHD}_c :

► **Corollary 81.** *We have an equivalence of categories $\mathbb{BC} \equiv \overline{\mathbb{BHD}}_c$.*

Combining these results we obtain the proof of our final corollary:

Proof of Corollary 42. By Theorem 36 we have the equivalences $\mathbb{FOB} \equiv \overline{\mathbb{BHD}}$, so we can combine this result with Corollary 78 obtaining the equivalence $\mathbb{FOB}_f \equiv \overline{\mathbb{BHD}}_c$. Therefore, combining this equivalence with Corollary 81 we obtain

$$\mathbb{FOB}_f \equiv \overline{\mathbb{BHD}}_c \equiv \mathbb{BC}.$$

◀