

NONEXISTENCE OF MAJORANA FERMIONS IN KERR-NEWMAN TYPE SPACETIMES WITH NONTRIVIAL CHARGE

HEQUN ZHANG ² AND XIAO ZHANG ^{1,2}

ABSTRACT. In this short paper, we study Majorana fermions in terms of a modified Chandrasekhar's separation for the Dirac equation in Kerr-Newman-type spacetimes. If the electric charge or magnetic charge is nonzero, we show that nontrivial differentiable time-periodic Majorana fermions do not exist outside the event horizon in Kerr-Newman and Kerr-Newman-AdS spacetimes, and between the event horizon and the cosmological horizon in Kerr-Newman-dS spacetime.

1. INTRODUCTION

The Dirac equation in black hole spacetimes plays a significant role in the study of general relativity and quantum cosmology. In 1968, Kinnersley introduced null basis to study the Petrov type D metric [1]. In 1976, Chandrasekhar separated the Dirac equation in Kerr spacetime [2], and Page generated his method to Kerr-Newman spacetime [3]. Since then, various works have been done to discuss the Hawking radiation and the numerical solutions of Dirac fermions in various spacetime backgrounds, see e.g. [4–7]. In 2000, Finster, Kamran, Smoller and Yau applied Chandrasekhar's separation to prove nonexistence of L^2 integrable, time-periodic solutions for the Dirac equation in non-extreme Kerr-Newman spacetime. This indicates that the normalizable time-periodic Dirac fermions must either disappear into the black hole or escape to infinity [8]. In 2008, Belgiorno and Cacciatori applied the spectral properties to prove the similar results for Kerr-Newman-AdS spacetime [9]. In 2018, Wang and Zhang applied Chandrasekhar's separation to prove nonexistence of L^p integrable, time-periodic solutions with arbitrary mass and $0 < p \leq \frac{4}{3}$, or with mass greater than $q\sqrt{-\frac{\Lambda}{3}}$ and $\frac{4}{3} < p \leq \frac{4}{3-2q}$, $1 < q < \frac{3}{2}$ for the Dirac equation in non-extreme Kerr-Newman-AdS spacetime. In particular, the normalizable time-periodic Dirac fermions with mass greater than $\frac{1}{2}\sqrt{-\frac{\Lambda}{3}}$ must either disappear into the black hole or escape to infinity, where Λ is the cosmological constant [10]. For non-extreme Kerr-Newman-dS spacetime, nonexistence of L^p integrable, time-periodic Dirac fermions hold true for arbitrary mass and $p \geq 2$ [11, 12].

For the recent search for neutrinos, which is one of the most mysterious particles in the universe, we are interesting to know whether they are Majorana or Dirac fermions. The most promising way so far is through the double beta decay [13]. Various approaches were studied to distinguish the Majorana and Dirac fermions, see e.g. [14–21].

A Majorana fermion is a Dirac fermion whose antiparticle is itself. The 4-component charge conjugate operator is

$$C = \begin{pmatrix} \epsilon_{\beta\alpha} & \\ & \epsilon^{\dot{\beta}\dot{\alpha}} \end{pmatrix},$$

with the Pauli matrix σ_2 and the antisymmetric operator on spin indices, where

$$\epsilon^{\dot{\alpha}\dot{\beta}} = -\epsilon_{\alpha\beta} = i\sigma_2 = \begin{pmatrix} & 1 \\ -1 & \end{pmatrix}.$$

The charge conjugation of the Dirac fermion Ψ_{Dir} is defined by

$$\Psi^C = C\bar{\Psi}^T.$$

Therefore, one can impose the Majorana condition [22–24]

$$\Psi_{\text{Maj}} = \begin{pmatrix} \psi \\ i\sigma_2\psi^* \end{pmatrix} \quad (1.1)$$

where ψ^* is the complex conjugate of ψ .

In this short paper, we apply a modified Chandrasekhar’s separation for the Dirac equation in Kerr-Newman, Kerr-Newman-(A)dS spacetimes to prove nonexistence of differentiable time-periodic Majorana fermions outside the event horizon in Kerr-Newman and Kerr-Newman-AdS spacetimes, and between the event horizon and the cosmological horizon in Kerr-Newman-dS spacetime.

2. GEOMETRY OF KERR-NEWMAN-TYPE SPACETIMES

For convenience of discussion, we unify the Kerr-Newman, the Kerr-Newman-(A)dS metrics by taking κ as zero, real and pure imaginary respectively

$$ds_{\text{KNTYPE}}^2 = -\frac{\Delta_r}{U} \left(dt - \frac{a \sin^2 \theta}{\Xi} d\phi \right)^2 + \frac{U}{\Delta_r} dr^2 + \frac{U}{\Delta_\theta} d\theta^2 + \frac{\Delta_\theta \sin^2 \theta}{U} \left(a dt - \frac{r^2 + a^2}{\Xi} d\phi \right)^2, \quad (2.1)$$

where $\Lambda = -3\kappa^2$ is the cosmological constant, and

$$\begin{aligned} \Delta_r &= (r^2 + a^2)(1 + \kappa^2 r^2) - 2mr + Q^2 + P^2, \\ \Delta_\theta &= 1 - \kappa^2 a^2 \cos^2 \theta, \\ U &= r^2 + a^2 \cos^2 \theta, \\ \Xi &= 1 - \kappa^2 a^2 > 0. \end{aligned}$$

The metric (2.1) solves the Einstein-Maxwell field equations with the electromagnetic potential

$$A = -\frac{Qr}{U} \left(dt - \frac{a \sin^2 \theta}{\Xi} d\phi \right) - \frac{P \cos \theta}{U} \left(a dt - \frac{r^2 + a^2}{\Xi} d\phi \right), \quad (2.2)$$

where P, Q are real numbers representing the magnetic charge, the electric charge respectively.

In the following we let $0 \leq \mu, \nu \leq 3$, and $1 \leq i, j \leq 3$. On a 4-dimensional Lorentzian manifold, we choose frame $\{e_\mu\}$ such that e_0 is timelike and e_i are spacelike. Denote $\{e^\alpha\}$ the dual coframe. The Cartan structure equations are

$$de^\mu = -\omega^\mu_\nu \wedge e^\nu, \quad \omega_{\mu\nu} = g_{\mu\gamma}\omega^\gamma_\nu = -\omega_{\nu\mu}.$$

If it is spin, we use the cotangent bundle to define the Clifford multiplication, the spin connection and the Dirac operator [10, 12]. We fix the Clifford multiplications to be the following Weyl representation

$$e^0 \mapsto \begin{pmatrix} & I_{2 \times 2} \\ I_{2 \times 2} & \end{pmatrix}, \quad e^i \mapsto \begin{pmatrix} & \sigma_i \\ -\sigma_i & \end{pmatrix}, \quad (2.3)$$

where σ_i are Pauli matrices

$$\sigma_1 = \begin{pmatrix} & 1 \\ 1 & \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} & -i \\ i & \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & \\ & -1 \end{pmatrix}.$$

The spin connection is

$$\tilde{\nabla}_X \Psi = X(\Psi) - \frac{1}{4} \omega_{\mu\nu}(X) e^\mu \cdot e^\nu \cdot \Psi, \quad (2.4)$$

where X is a vector, Ψ is a spinor, $e^\alpha \cdot$ is the Clifford multiplication. The Dirac operator with electromagnetic potential A is

$$D = e^\mu \cdot \left(\tilde{\nabla}_{e_\mu} + iA(e_\mu) \right). \quad (2.5)$$

We fix our discussion in the region $\Delta_r > 0$. The coframe is

$$e^0 = \sqrt{\frac{\Delta_r}{U}} \left(dt - \frac{a \sin^2 \theta}{\Xi} d\phi \right), \quad e^1 = \sqrt{\frac{U}{\Delta_\theta}} d\theta \\ e^2 = \sqrt{\frac{\Delta_\theta}{U}} \sin \theta \left(a dt - \frac{r^2 + a^2}{\Xi} d\phi \right), \quad e^3 = \sqrt{\frac{U}{\Delta_r}} dr.$$

with dual frame

$$e_0 = \frac{r^2 + a^2}{\sqrt{U\Delta_r}} \left(\partial_t + \frac{a\Xi}{r^2 + a^2} \partial_\phi \right), \quad e_1 = \sqrt{\frac{\Delta_r}{U}} \partial_r, \\ e_2 = \sqrt{\frac{\Delta_\theta}{U}} \partial_\theta, \quad e_3 = -\frac{1}{\sqrt{U\Delta_\theta}} \left(a \sin \theta \partial_t + \frac{\Xi}{\sin \theta} \partial_\phi \right).$$

With respect to the above coframe, it gives that

$$de^0 = C_{10}^0 e^1 \wedge e^0 + C_{30}^0 e^3 \wedge e^0 + C_{12}^0 e^1 \wedge e^2, \\ de^1 = C_{31}^1 e^3 \wedge e^1, \\ de^2 = C_{30}^2 e^3 \wedge e^0 + C_{32}^2 e^3 \wedge e^2 + C_{12}^2 e^1 \wedge e^2, \\ de^3 = C_{31}^3 e^3 \wedge e^1,$$

where

$$C_{10}^0 = -\frac{a^2}{U} \sqrt{\frac{\Delta_\theta}{U}} \sin \theta \cos \theta, \quad C_{30}^0 = \partial_r \sqrt{\frac{\Delta_r}{U}}, \quad C_{12}^0 = \frac{2a}{U} \sqrt{\frac{\Delta_r}{U}} \cos \theta, \\ C_{31}^1 = \frac{r}{U} \sqrt{\frac{\Delta_r}{U}}, \quad C_{12}^2 = \frac{1}{\sin \theta} \partial_\theta \left(\sqrt{\frac{\Delta_\theta}{U}} \sin \theta \right), \quad C_{32}^2 = \frac{r}{U} \sqrt{\frac{\Delta_r}{U}},$$

$$C_{30}^2 = -\frac{2ar}{U}\sqrt{\frac{\Delta_\theta}{U}}\sin\theta, \quad C_{31}^3 = \frac{a^2}{U}\sqrt{\frac{\Delta_\theta}{U}}\sin\theta\cos\theta.$$

Thus connection 1-forms are

$$\begin{aligned} \omega_{1}^0 &= -\omega_{01} = C_{10}^0 e^0 + \frac{1}{2}C_{12}^0 e^2, & \omega_{2}^0 &= -\omega_{02} = -\frac{1}{2}C_{30}^2 e^3 - \frac{1}{2}C_{12}^0 e^1, \\ \omega_{3}^0 &= -\omega_{03} = C_{30}^0 e^0 - \frac{1}{2}C_{30}^2 e^2, & \omega_{2}^1 &= \omega_{12} = \frac{1}{2}C_{12}^0 e^0 - C_{12}^2 e^2, \\ \omega_{3}^1 &= \omega_{13} = C_{31}^3 e^3 + C_{31}^1 e^1, & \omega_{3}^2 &= \omega_{23} = \frac{1}{2}C_{30}^2 e^0 + C_{32}^2 e^2. \end{aligned} \quad (2.6)$$

Therefore we obtain spin connections

$$\begin{aligned} \tilde{\nabla}_{e_0}\Psi &= e_0\Psi - \frac{1}{2}\omega_{01}(e_0)e^0 \cdot e^1 \cdot \Psi - \frac{1}{2}\omega_{03}(e_0)e^0 \cdot e^3 \cdot \Psi \\ &\quad - \frac{1}{2}\omega_{12}(e_0)e^1 \cdot e^2 \cdot \Psi - \frac{1}{2}\omega_{23}(e_0)e^2 \cdot e^3 \cdot \Psi, \\ \tilde{\nabla}_{e_1}\Psi &= e_1\Psi - \frac{1}{2}\omega_{02}(e_1)e^0 \cdot e^2 \cdot \Psi - \frac{1}{2}\omega_{12}(e_1)e^1 \cdot e^2 \cdot \Psi, \\ \tilde{\nabla}_{e_2}\Psi &= e_2\Psi - \frac{1}{2}\omega_{01}(e_2)e^0 \cdot e^1 \cdot \Psi - \frac{1}{2}\omega_{03}(e_2)e^0 \cdot e^3 \cdot \Psi \\ &\quad - \frac{1}{2}\omega_{12}(e_2)e^1 \cdot e^2 \cdot \Psi - \frac{1}{2}\omega_{23}(e_2)e^2 \cdot e^3 \cdot \Psi, \\ \tilde{\nabla}_{e_3}\Psi &= e_3\Psi - \frac{1}{2}\omega_{02}(e_3)e^0 \cdot e^2 \cdot \Psi - \frac{1}{2}\omega_{13}(e_3)e^1 \cdot e^3 \cdot \Psi. \end{aligned}$$

They can be written as the matrix forms

$$\tilde{\nabla}_{e_\mu}\Psi = e_\mu\Psi + E_\mu \cdot \Psi, \quad E_\mu = -\frac{1}{2} \begin{pmatrix} \epsilon_\mu & 0 \\ 0 & -\epsilon_\mu^h \end{pmatrix},$$

where ϵ_μ^h is the Hermitian conjugate of ϵ_μ and

$$\begin{aligned} \epsilon_0 &= \left(C_{10}^0 - \frac{i}{2}C_{30}^2\right)\sigma_1 + \left(C_{30}^0 - \frac{i}{2}C_{12}^0\right)\sigma_3, \\ \epsilon_1 &= \left(-\frac{1}{2}C_{12}^0 + iC_{31}^1\right)\sigma_2, \\ \epsilon_2 &= \left(\frac{1}{2}C_{12}^0 - iC_{32}^2\right)\sigma_1 - \left(\frac{1}{2}C_{30}^2 - iC_{12}^2\right)\sigma_3, \\ \epsilon_3 &= \left(-\frac{1}{2}C_{30}^2 + iC_{31}^3\right)\sigma_2. \end{aligned}$$

3. TIME-PERIODIC MAJORANA FERMIONS

The Dirac equation on metric (2.1) is given by

$$(D + i\lambda)\Psi = 0, \quad (3.1)$$

where λ is certain real number, A is given by (2.2). Time-periodic Majorana fermions take the following form

$$\Psi = S^{-1}E\psi, \quad \psi = \begin{pmatrix} R_-(r)\Theta_-(\theta) \\ R_+(r)\Theta_+(\theta) \\ \bar{R}_+(r)\bar{\Theta}_+(\theta) \\ -\bar{R}_-(r)\bar{\Theta}_-(\theta) \end{pmatrix} \quad (3.2)$$

and solve the Dirac equation (3.1), where S, E are diagonal matrices

$$S = \Delta_r^{\frac{1}{4}} \text{diag} \left((r + ia \cos \theta)^{\frac{1}{2}} I_{2 \times 2}, (r - ia \cos \theta)^{\frac{1}{2}} I_{2 \times 2} \right), \\ E = \text{diag} \left(e^{-i(\omega t + (k + \frac{1}{2})\phi)} I_{2 \times 2}, e^{i(\omega t + (k + \frac{1}{2})\phi)} I_{2 \times 2} \right).$$

Now we simplify the Dirac equation. Denote $\mathcal{J} = \text{diag}(I_{2 \times 2}, -I_{2 \times 2})$. In terms of (2.4), we have

$$\begin{aligned} e^\mu \cdot e_\mu(\Psi) &= \sqrt{\frac{\Delta_r}{U}} e^3 \cdot S^{-1} E \partial_r \psi + \sqrt{\frac{\Delta_\theta}{U}} e^1 \cdot S^{-1} E \partial_\theta \psi \\ &\quad - i \frac{r^2 + a^2}{\sqrt{U} \Delta_r} \left(\omega + \frac{a\Xi}{r^2 + a^2} (k + \frac{1}{2}) \right) e^0 \cdot \mathcal{J} S^{-1} E \psi \\ &\quad + \frac{1}{2U^{\frac{3}{2}}} e^3 \cdot (S - \partial_r(U\sqrt{\Delta_r})S^{-1}) E \psi \\ &\quad + \frac{a \sin \theta}{2U^{\frac{3}{2}}} \sqrt{\frac{\Delta_\theta}{\Delta_r}} e^1 \cdot (i\mathcal{J}S + 2a \cos \theta \sqrt{\Delta_r} S^{-1}) E \psi \\ &\quad + \frac{i}{\sqrt{U} \Delta_\theta} \left(a\omega \sin \theta + \frac{\Xi}{\sin \theta} (k + \frac{1}{2}) \right) e^2 \cdot S^{-1} \mathcal{J} E \psi, \\ e^\mu \cdot E_\mu \Psi &= \frac{2\Delta_\theta - \Xi}{2\sqrt{U} \Delta_\theta} \cot \theta e^1 \cdot S^{-1} E \psi - \frac{ia}{2} \sqrt{\frac{\Delta_r \Delta_\theta}{U}} \sin \theta e^1 \cdot \mathcal{J} S^{-3} E \psi \\ &\quad + \frac{\partial_r \sqrt{\Delta_r}}{2\sqrt{U}} e^3 \cdot S^{-1} E \psi + \frac{\Delta_r}{2\sqrt{U}} e^3 \cdot S^{-3} E \psi, \\ e^\mu \cdot (iA(e_\mu)) \Psi &= - \frac{iQr}{\sqrt{U} \Delta_r} e^0 \cdot S^{-1} E \psi - \frac{iP \cot \theta}{\sqrt{U} \Delta_\theta} e^2 \cdot S^{-1} E \psi. \end{aligned}$$

Note that

$$\begin{aligned} e^\mu \mathcal{J} &= -\mathcal{J} e^\mu, \quad e^\mu E = E^{-1} e^\mu, \quad e^\mu E^{-1} = E e^\mu, \\ e^\mu S^{-1} &= \frac{1}{\sqrt{U} \Delta_r} S e^\mu, \quad e^\mu S = \sqrt{U \Delta_r} S^{-1} e^\mu. \end{aligned}$$

We obtain

$$D\Psi = \frac{1}{U\sqrt{\Delta_r}} S E^{-1} \left(\sqrt{\Delta_r} \mathcal{D}_r - \sqrt{\Delta_\theta} \mathcal{L}_\theta \right) \psi \quad (3.3)$$

with

$$\begin{aligned} \mathcal{D}_r &= e^3 \partial_r + \frac{i}{\Delta_r} \left(\omega(r^2 + a^2) + a\Xi(k + \frac{1}{2}) \right) \mathcal{J} e^0 - \frac{iQr}{\Delta_r} e^0, \\ \mathcal{L}_\theta &= -e^1 \partial_\theta + \frac{i}{\Delta_\theta} \left(\omega a \sin \theta + \frac{\Xi}{\sin \theta} (k + \frac{1}{2}) \right) \mathcal{J} e^2 \\ &\quad - \left(1 - \frac{\Xi}{2\Delta_\theta} \right) \cot \theta e^1 + \frac{iP}{\Delta_\theta} \cot \theta e^2. \end{aligned}$$

Denote $\lambda_{\omega k} = \lambda e^{-2i(\omega t + (k + \frac{1}{2})\phi)}$. Then the Dirac equation (3.1) reduces to

$$D_r \psi = L_\theta \psi, \quad \psi = \begin{pmatrix} R_-(r) \Theta_-(\theta) \\ R_+(r) \Theta_+(\theta) \\ \bar{R}_+(r) \bar{\Theta}_+(\theta) \\ -\bar{R}_-(r) \bar{\Theta}_-(\theta) \end{pmatrix} \quad (3.4)$$

where

$$D_r = \begin{pmatrix} -i\lambda_{\omega k} r & & \sqrt{\Delta_r} D_{r,00} & \\ & -i\lambda_{\omega k} r & & \sqrt{\Delta_r} D_{r,01} \\ \sqrt{\Delta_r} D_{r,11} & & -i\bar{\lambda}_{\omega k} r & \\ & \sqrt{\Delta_r} D_{r,10} & & -i\bar{\lambda}_{\omega k} r \end{pmatrix}$$

$$L_\theta = \begin{pmatrix} a\lambda_{\omega k} \cos \theta & & \sqrt{\Delta_\theta} L_{\theta,00} & \\ & a\lambda_{\omega k} \cos \theta & \sqrt{\Delta_\theta} L_{\theta,01} & \\ \sqrt{\Delta_\theta} L_{\theta,11} & & -a\bar{\lambda}_{\omega k} \cos \theta & \\ & \sqrt{\Delta_\theta} L_{\theta,10} & & -a\bar{\lambda}_{\omega k} \cos \theta \end{pmatrix}$$

and, for $l, m=0, 1$,

$$D_{r,lm} = (-1)^m \partial_r + (-1)^l \frac{i}{\Delta_r} \left(\omega(r^2 + a^2) + (k + \frac{1}{2})\Xi a \right) - \frac{iQr}{\Delta_r},$$

$$L_{\theta,lm} = -(-1)^l \partial_\theta + \frac{(-1)^{l+m}}{\Delta_\theta} \left(\omega a \sin \theta + \frac{\Xi}{\sin \theta} (k + \frac{1}{2}) \right. \\ \left. + (-1)^l P \cot \theta - (-1)^m \left(\Delta_\theta - \frac{\Xi}{2} \right) \cot \theta \right). \quad (3.5)$$

Writing down each row of (3.4), we get

$$-i\lambda_{\omega k} r R_- \Theta_- + \sqrt{\Delta_r} D_{r,00} \bar{R}_+ \bar{\Theta}_+ = a\lambda_{\omega k} \cos \theta R_- \Theta_- - \sqrt{\Delta_\theta} L_{\theta,00} \bar{R}_- \bar{\Theta}_-, \quad (3.6)$$

$$-i\lambda_{\omega k} r R_+ \Theta_+ - \sqrt{\Delta_r} D_{r,01} \bar{R}_- \bar{\Theta}_- = a\lambda_{\omega k} \cos \theta R_+ \Theta_+ + \sqrt{\Delta_\theta} L_{\theta,01} \bar{R}_+ \bar{\Theta}_+, \quad (3.7)$$

$$-i\bar{\lambda}_{\omega k} r \bar{R}_+ \bar{\Theta}_+ + \sqrt{\Delta_r} D_{r,11} R_- \Theta_- = -a\bar{\lambda}_{\omega k} \cos \theta \bar{R}_+ \bar{\Theta}_+ + \sqrt{\Delta_\theta} L_{\theta,11} R_+ \Theta_+, \quad (3.8)$$

$$i\bar{\lambda}_{\omega k} r \bar{R}_- \bar{\Theta}_- + \sqrt{\Delta_r} D_{r,10} R_+ \Theta_+ = a\bar{\lambda}_{\omega k} \cos \theta \bar{R}_- \bar{\Theta}_- + \sqrt{\Delta_\theta} L_{\theta,10} R_- \Theta_-. \quad (3.9)$$

Denote $\bar{l} = (l+1) \bmod 2$, $\bar{m} = (m+1) \bmod 2$. Note that

$$\bar{D}_{r,lm} = -D_{r,l\bar{m}}, \quad \bar{L}_{\theta,lm} = L_{\theta,lm}$$

and

$$D_{r,lm} + D_{r,l\bar{m}} = -\frac{2iQr}{\Delta_r},$$

$$L_{\theta,lm} + L_{\theta,l\bar{m}} = 2(-1)^m \frac{P \cot \theta}{\Delta_\theta}.$$

Subtracting the complex conjugation of (3.6) from (3.9), adding the complex conjugation of (3.7) with (3.8) respectively, we obtain

$$i\alpha(r) R_- \Theta_- - \beta(\theta) R_+ \Theta_+ = 0,$$

$$\beta(\theta) R_- \Theta_- + i\alpha(r) R_+ \Theta_+ = 0,$$

where

$$\alpha(r) = \frac{Qr}{\sqrt{\Delta_r}}, \quad \beta(\theta) = \frac{P \cot \theta}{\sqrt{\Delta_\theta}}.$$

Thus

$$\left(\alpha(r)^2 - \beta(\theta)^2\right) R_+ \Theta_+ = \left(\alpha(r)^2 - \beta(\theta)^2\right) R_- \Theta_- = 0.$$

Therefore, if $P \neq 0$ or $Q \neq 0$, $R_+ \Theta_+$ and $R_- \Theta_-$ could be nontrivial only in the hypersurface $\alpha(r) = \pm \beta(\theta)$. But they will not be differentiable in this case. Thus there is no differentiable time-periodic Majorana fermions in Kerr-Newman type spacetimes.

4. CONCLUSION

We conclude that there is no differentiable time-periodic Majorana fermion outside the event horizon in Kerr-Newman and Kerr-Newman-AdS spacetimes, and between the event horizon and the cosmological horizon in Kerr-Newman-dS spacetime if the electric charge or magnetic charge is nonzero.

Acknowledgement The work is supported by the National Natural Science Foundation of China 12326602.

REFERENCES

- [1] W. Kinnersley, Type D vacuum metrics. *Journal of Mathematical Physics* **10** (1969) 1195-1203.
- [2] S. Chandrasekhar, The solution of Dirac's equation in Kerr geometry. *Proceedings of the Royal Society, Series A* **349** (1976) 571-575.
- [3] D. N. Page, Dirac equation around a charged, rotating black hole. *Physical Review D*, **14** (1976) 1509-1510.
- [4] Z. Zhao, Y. X. Guei and L. Liu, On the Hawking evaporation of Dirac particles in Kerr-Newman spacetime. *Chinese Astronomy and Astrophysics*, **5** (1981) 365-372.
- [5] F. Belgiorno and M. Martellini, Quantum properties of the electron field in Kerr-Newman black hole manifolds. *Physics Letters B*, **453** (1999) 17-22.
- [6] S. K. Chakrabarti and B. Mukhopadhyay, Scattering of Dirac waves of Kerr black holes. *Monthly Notices of the Royal Astronomical Society*, **317** (2000) 979-984.
- [7] M. Angheben, M. Nadalini, L. Vanzo and S. Zerbini, Hawking radiation as tunneling for extremal and rotating black holes. *Journal of High Energy Physics*, **05** (2005) 014.
- [8] F. Finster, N. Kamran, J. Smoller and S.T. Yau, Nonexistence of time-periodic solutions of the Dirac equation in an axisymmetric black hole geometry. *Communications on Pure and Applied Mathematics*, **53** (2000) 902-929.
- [9] F. Belgiorno and S. L. Cacciatori, The Dirac equation in Kerr-Newman-AdS black hole background. *Journal of Mathematical Physics*, **51** (2010) 033517.
- [10] Y. Wang and X. Zhang, Nonexistence of time-periodic solutions of the Dirac equation in non-extreme Kerr-Newman-AdS spacetime. *Science China-mathematics*, **61** (2018) 73-82.
- [11] F. Belgiorno, S. L. Cacciatori, The absence of normalizable time-periodic solutions for the Dirac equation in the Kerr-Newman-dS black hole background. *Journal of Physics A: Mathematical and Theoretical*, 2009, **42**(13): 135207.
- [12] M. Fan, Y. Wang and X. Zhang, Nonexistence of time-periodic solutions of the Dirac equation in Kerr-Newman-(A)dS spacetime. *ArXiv:2404.13255v1 [gr-qc]* 20 Apr 2024
- [13] W. Rodejohann, Neutrino-less double Beta decay and particle physics. *International Journal of Modern Physics E*, **20** (2011) 1833-1930.

- [14] S. A. Alavi and A. Abbasnezhad, Can gravity distinguish between Dirac and Majorana neutrinos? *Gravitation and Cosmology*, **22** (2016):288-298.
- [15] T. P. Cheng and L. F. Li, $\mu \rightarrow e\gamma$ in theories with Dirac and Majorana neutrino mass terms. *Physical Review Letters*, **45** (1980) 1908-1911.
- [16] T. Garavaglia, Dirac and Majorana neutrino mass effects in neutrino-electron elastic scattering. *Physical Review D*, **29** (1984) 387.
- [17] J. Nieves and P. Pal, Minimal rephasing-invariant CP-violating parameters with Dirac and Majorana fermions. *Physical Review D*, **36** (1987) 315-317.
- [18] S. P. Rosen, Analog of the Michel parameter for neutrino-electron scattering: A test for Majorana neutrinos. *Physical Review Letters*, **48** (1982) 842.
- [19] R. E. Shrock, Electromagnetic properties and decays of Dirac and Majorana neutrinos in a general class of gauge theories. *Nuclear Physics B*, **206** (1982) 359-379.
- [20] D. Singh, N. Mobed and G. Papini. Can gravity distinguish between Dirac and Majorana neutrinos? *Physical Review Letters*, **97** (2006) 041101.
- [21] C. S. Kim, M. V. N. Murthy and D. Sahoo. Inferring the nature of active neutrinos: Dirac or Majorana? *Physical Review D*, **105** (2022) 113006.
- [22] E. Majorana, Theory of the symmetry of electrons and positrons. *Nuovo Cimento C-Colloquia and Communications in Physics*, **14** (1937) 171.
- [23] T. Prokopec and V. H. Unnithan, Majorana propagator on de Sitter space. *European Physical Journal C*, (2022) **82** (2022) (11) 1015.
- [24] C-L. Hsieh, V. Memari and M. Halilsoy. Dirac and Majorana fermions in the anti-de Sitter spacetime with tachyonic approaches. *ArXiv:2403.13810v1 [hep-th]* 4 Jan 2024.

¹ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, PR CHINA

²SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF CHINESE ACADEMY OF SCIENCES, BEIJING 100049, PR CHINA

Email address: zhanghequn@amss.ac.cn²

Email address: xzhang@amss.ac.cn^{1,2}