Study of $B_{(s)}$ meson decays to $D_0^*(2300), D_{s0}^*(2317), D_{s1}(2460)$ and $D_{s1}(2536)$ within the covariant light-front approach

You-Ya Yang, Zhi-Qing Zhang *, Hao Yang, Zhi-Jie Sun, Ming-Xuan Xie

Institute of Theoretical Physics, School of Sciences, Henan University of Technology, Zhengzhou450052, Henan, China (Dated: May 2, 2024)

Abstract

In this work, we investigate the form factors of the transitions $B_{(s)} \rightarrow D_0^*(2300), D_{s0}^*(2317), D_{s1}(2460)$ and $D_{s1}(2536)$ in the covariant light-front quark model (CLFQM), where these final states are considered as P-wave excited charmed mesons. In order to obtain the form factors for the physical transition processes, we need to extend these form factors from the space-like region to the time-like region. The q^2 -dependence for each transition form factor is also plotted. Then, combined with those form factors, the branching ratios of the two-body nonleptonic decays $B_{(s)} \rightarrow D_{(s)0}^*(2300, 2317)M, D_{s1}(2460, 2536)M$ with M being a light pseudoscalar (vector) meson or a charmed meson are calculated by considering the QCD radiative corrections to the hadronic matrix elements with the QCD factorization approach. Most of our predictions are comparable to the results given by other theoretical approaches and the present available data.

PACS numbers: 13.25.Hw, 12.38.Bx, 14.40.Nd

^{*} Electronic address: zhangzhiqing@haut.edu.cn (corresponding author)

I. INTRODUCTION

In this work, we would like to study some of excited open-charm states, such as $D_0^*(2300), D_{s0}^*(2317), D_{s1}(2460)$ and $D_{s1}(2536)$ in $B_{(s)}$ meson decays. As we know, $D_{s0}^*(2317)$ and $D_{s1}(2460)$ were first discovered by BaBar [1] and CLEO [2] in 2003, respectively. The scalar charm-strange meson $D_{s0}^*(2317)$ was observed in the invariant mass distribution of $D_s^+\pi^0$ and the axial charm-strange meson $D_{s1}(2460)$ was found in the invariant mass distribution of $D_s^*\pi^0$. $D_0^*(2300)$ as the SU(3) partner of $D_0^*(2317)$, called $D_0^*(2400)$ in the past, was discovered by Belle [3] in the three-body B decay $B^+ \to D^-\pi^+\pi^+$ in 2004. Another axial charm-strange meson with $J^P = 1^+ D_{s1}(2536)$ was first observed in the $D_s^*\gamma$ invariant mass spectrum [4] in 1987.

Previous researches suggest that the masses of the resonants $D_{s0}^*(2317)$ and $D_{s1}(2460)$ are just several tens MeV below the threshold of DK and D^*K , respectively. Furthermore, they are much lower than those given by the quark model [5]. Some abnormal properties induce that these two states are difficult to be interpreted as conventional $c\bar{s}$ mesons. Therefore, many authors consider them as the $D^{(*)}K$ molecular states [6–12], the compact tetraquark states [13–17], the chiral partners of the ground state $D_s^{(*)}$ mesons [18, 19] and the states of $c\bar{s}$ mixed with four-quark states [20, 21]. However, if considering the coupled channel effects and the fact that there are no additional states around the quark model predicted masses, these two states can be interpreted as P-wave charm-strange mesons [22–31]. As to the $D_0^*(2300)$ state, except the low mass puzzle, where the observed mass of $D_0^*(2300)$ [3, 32] is below the predictions from the quark model about 100 MeV [5, 33], there exists the SU(3) mass hierarchy puzzle, that is the masses of $D_0^*(2300)$ and $D_{s0}^*(2317)$ are almost equal to each other. These puzzles have triggered many studies on their inner structures: In Refs. [22, 34], the authors pointed out that the four-quark structure can explain the data measured by Belle [3] and BaBar [32], but not for that measured by FOCUS [35]. Another group authors solved these puzzles within the framework of unitarized chiral perturbation theory (UChPT) [36–38] and considered that there exist two states in the $D_0^*(2300)$ energy region [39], where the lighter one named as $D_0^*(2100)$ is the SU(3) partner of the $D_{s0}^*(2317)$, the heavier one is a member of a different multiplet. Certainly, other authors also explained $D_0^*(2300)$ as a mixture of two and four-quark states [21] or the bound state of $D\pi$ [40]. The axial charm-strange meson $D_{s1}(2536)$ has been confirmed and studied in the $B_{(s)}$ meson decays by BaBar [41], Belle [42] and LHCb [43]. Its measurements of mass and width are consistent with the theoretical expectations as a charm-strange meson with $J^P = 1^+$. As we know that the QCD Lagrangian is invariant under the heavy flavor and spin rotation in the heavy quark limit. For the heavy mesons, the heavy quark spin S_Q can decouple from the other degrees of freedom. Then S_Q and the total angular momentum of the light quark *j* become good quantum numbers. It is natural to label $D_{s1}(2460)$ and $D_{s1}(2536)$ by the quantum numbers L_J^j with j(L) being the total (orbital) angular momentum of the light quark, that is $P_1^{1/2}$ and $P_1^{3/2}$, denoted as $D_{s1}^{1/2}$ and $D_{s1}^{3/2}$, respectively. Since the heavy quark symmetry is not exact, the two 1⁺ states $D_{s1}(2460)$ and $D_{s1}(2536)$ can mix with each other through the following formula [44]

$$|D_{s1}(2460)\rangle = |D_{s1}^{1/2}\rangle\cos\theta_s + |D_{s1}^{3/2}\rangle\sin\theta_s, |D_{s1}(2536)\rangle = -|D_{s1}^{1/2}\rangle\sin\theta_s + |D_{s1}^{3/2}\rangle\cos\theta_s.$$
(1)

While the states $D_{s1}^{1/2}$ and $D_{s1}^{3/2}$ are expected to be a mixture of states ${}^{1}D_{s1}$ and ${}^{3}D_{s1}$ with $J^{PC} = 1^{++}$ and 1^{+-} , respectively,

$$|D_{s1}^{3/2}\rangle = \sqrt{\frac{2}{3}}|^{1}D_{s1}\rangle + \sqrt{\frac{1}{3}}|^{3}D_{s1}\rangle, |D_{s1}^{1/2}\rangle = \sqrt{\frac{1}{3}}|^{1}D_{s1}\rangle - \sqrt{\frac{2}{3}}|^{3}D_{s1}\rangle.$$
(2)

Combining Eq. (1) and Eq. (2), one can find that

$$|D_{s1}(2460)\rangle = |{}^{1}D_{s1}\rangle\sin\theta - |{}^{3}D_{s1}\rangle\cos\theta,$$

$$|D_{s1}(2536)\rangle = |{}^{1}D_{s1}\rangle\cos\theta + |{}^{3}D_{s1}\rangle\sin\theta,$$
(3)

where $\theta = \theta_s + 35.3^{\circ}$ [45].

Using the manifestly covariant of the Bethe-Salpeter (BS) approach [46, 47], Jaus, Choi, Ji and Cheng *et al.* [48–50] put forward the CLFQM around 2000. This approach provides a systematic way to explore the zero-mode effects, which are just canceled by involving the spurious contributions being proportional to the lightlike four-vector $\omega = (0, 2, 0_{\perp})$, at the same time the covariance of the matrix elements being restored [48]. Up to now the CLFQM has been used extensively to study the weak and radiative decays, as well as the features of some exotic hadrons [51–59]. In this work, we will employ the CLFQM to evaluate the $B_{(s)} \rightarrow D_0^*(2300), D_{s0}^*(2317), D_{s1}(2460), D_{s1}(2536)$ transition form factors, then the branching ratios of the relevant decays are calculated. In our calculations these hadrons are regarded as ordinary meson states. Compared with the future experimental measurements, our predictions are helpful to clarify the inner structures of these four hadrons.

The arrangement of this paper is as follows: In Section II, an introduction to the CLFQM and the expressions for the form factors of the transitions $B_{(s)} \rightarrow D_0^*(2300), D_{s0}^*(2317), D_{s1}(2460), D_{s1}(2536)$ are presented. Then the branching ratios of the $B_{(s)}$ meson decays with one of these considered P-wave excited charm sates involved are calculated under the QCD factorization approach, where the vertex corrections and the hard spectator-scattering corrections are considered. In Section III, the numerical results of the transition form factors and their q^2 -dependence are presented. Then, combined with the transition form factors, the branching ratios of the decays $B_{(s)} \rightarrow D^*_{(s)0}(2300, 2317)M, D_{s1}(2460, 2536)M$ with M being a light pseudoscalar (vector) meson or a charmed meson are calculated. In addition, detailed numerical analysis and discussion, including comparisons with the data and other model calculations, are carried out. The conclusions are presented in the final part.

II. FORMALISM

A. The Covariant Light-Front Quark Model

Under the covariant light-front quark model, the light-front coordinates of a momentum p are defined as $p = (p^-, p^+, p_\perp)$ with $p^{\pm} = p^0 \pm p_z$ and $p^2 = p^+p^- - p_\perp^2$. If the momenta of the quark and antiquark with mass $m'_1^{(\prime\prime)}$ and m_2 in the incoming (outgoing) meson are denoted as $p'_1^{(\prime\prime)}$ and p_2 , respectively, the momentum of the incoming (outgoing) meson with mass M'(M'') can be written as $P' = p'_1 + p_2(P'' = p''_1 + p_2)$. Here, we use the same notation as those in Refs. [48, 51] and M' refers to m_B for B meson decays. These momenta can be related each other through the internal variables (x_i, p'_\perp)

$$p_{1,2}^{\prime +} = x_{1,2}P^{\prime +}, \quad p_{1,2\perp}^{\prime} = x_{1,2}P_{\perp}^{\prime} \pm p_{\perp}^{\prime},$$
 (4)

with $x_1 + x_2 = 1$. Using these internal variables, we can define some quantities for the incoming meson which will be used in the following calculations:

$$M_0^{\prime 2} = (e_1^{\prime} + e_2)^2 = \frac{p_{\perp}^{\prime 2} + m_1^{\prime 2}}{x_1} + \frac{p_{\perp}^2 + m_2^2}{x_2}, \quad \widetilde{M}_0^{\prime} = \sqrt{M_0^{\prime 2} - (m_1^{\prime} - m_2)^2},$$
$$e_i^{\prime \prime} = \sqrt{m_i^{\prime \prime 2} + p_{\perp}^{\prime 2} + p_z^{\prime 2}}, \qquad p_z^{\prime} = \frac{x_2 M_0^{\prime}}{2} - \frac{m_2^2 + p_{\perp}^{\prime 2}}{2x_2 M_0^{\prime}},$$
(5)

where the kinetic invariant mass of the incoming meson M'_0 can be expressed as the energies of the quark and the antiquark $e_i^{(\prime)}$. It is similar to the case of the outgoing meson.

The form factors of the transitions $B \to D_0^{* 1}$ and $B \to {}^i D_{s1}(i = 1, 3)^{2}$ induced by the

¹ It is similar for the transition $B \to D_{s0}^*$. From now on, we will use D_0^* and D_{s0}^* to represent $D_0^*(2300)$ and $D_{s0}^*(2317)$, respectively for simplicity.

² We will use D_{s1} and D'_{s1} to represent $D_{s1}(2460)$ and $D_{s1}(2536)$, respectively. The form factors of the transitions $B \to D_{s1}$ and $B \to D'_{s1}$ can be obtained from those of the transitions $B \to {}^{1}D_{s1}$ and $B \to {}^{3}D_{s1}$ through Eq. (3).




FIG. 1: Feynman diagrams for B decay (left) and transition (right) amplitudes, where $P'^{(\prime\prime)}$ is the incoming (outgoing) meson momentum, $p'_1^{(\prime\prime)}$ is the quark momentum, p_2 is the antiquark momentum and X denotes the vector or axial vector transition vertex.

vector and aixal-vector currents are defined as

$$\langle D_0^*(P'') | A_\mu | B(P') \rangle = i \left[u_+(q^2) P_\mu + u_-(q^2) q_\mu \right], \tag{6}$$

$$\left\langle {}^{i}D_{s1}\left(P^{\prime\prime},\varepsilon\right)\left|A_{\mu}\right|B\left(P^{\prime}\right)\right\rangle = -q(q^{2})\epsilon_{\mu\nu\alpha\beta}\varepsilon^{*\nu}P^{\alpha}q^{\beta},\tag{7}$$

$$\left\langle {}^{i}D_{s1}\left(P'',\varepsilon\right)|V_{\mu}|B\left(P'\right)\right\rangle = i\left\{ l(q^{2})\varepsilon_{\mu}^{*} + \varepsilon^{*} \cdot P\left[P_{\mu}c_{+}(q^{2}) + q_{\mu}c_{-}(q^{2})\right]\right\}.$$
(8)

In calculations, the Bauer-Stech-Wirbel (BSW) [60] transition form factors are more frequently used and defined by

$$\langle D_0^*(P'') | A_\mu | B(P') \rangle = \left(P_\mu - \frac{m_B^2 - m_{D_0^*}^2}{q^2} q_\mu \right) F_1^{BD_0^*} \left(q^2 \right) + \frac{m_B^2 - m_{D_0^*}^2}{q^2} q_\mu F_0^{BD_0^*} \left(q^2 \right), \tag{9}$$

$$\langle {}^{i}D_{s1}\left(P'',\varepsilon^{\mu*}\right)|V_{\mu}|B\left(P'\right)\rangle = -i\left\{ \left(m_{B}-m_{i}_{D_{s1}}\right)\varepsilon_{\mu}^{*}V_{1}^{B\ i}D_{s1}\left(q^{2}\right) - \frac{\varepsilon^{*}\cdot P}{m_{B}-m_{i}_{D_{s1}}}P_{\mu}V_{2}^{B\ i}D_{s1}\left(q^{2}\right) - 2m_{i}_{D_{s1}}\frac{\varepsilon^{*}\cdot P}{q^{2}}q_{\mu}\left[V_{3}^{B\ i}D_{s1}\left(q^{2}\right) - V_{0}^{B\ i}D_{s1}\left(q^{2}\right)\right]\right\},$$
(10)

$$\left\langle {}^{i}D_{s1}\left(P^{\prime\prime},\varepsilon^{\mu*}\right)\left|A_{\mu}\right|B\left(P^{\prime}\right)\right\rangle = -\frac{1}{m_{B}-m_{i}D_{s1}}\epsilon_{\mu\nu\alpha\beta}\varepsilon^{*\nu}P^{\alpha}q^{\beta}A^{B^{-i}D_{s1}}\left(q^{2}\right),\tag{11}$$

where P = P' + P'', q = P' - P'', and the convention $\epsilon_{0123} = 1$ is adopted.

To smear the singularity at $q^2 = 0$ in Eq. (10), the relations $V_3^{B\ i} D_{s1}(0) = V_0^{B\ i} D_{s1}(0)$ are required, and

$$V_3^{B\ i} V_{3}^{D\ i}(q^2) = \frac{m_B - m_{i}_{D_{s1}}}{2m_{i}_{D_{s1}}} V_1^{B\ i} V_1^{D\ i}(q^2) - \frac{m_B + m_{i}_{D_{s1}}}{2m_{i}_{D_{s1}}} V_2^{B\ i} V_2^{D\ i}(q^2).$$
(12)

These two kinds of form factors are related to each other via

$$F_1^{BD_0^*}(q^2) = -u_+(q^2), F_0^{BD_0^*}(q^2) = -u_+(q^2) - \frac{q^2}{q \cdot P}u_-(q^2),$$
(13)

$$A^{B\ i}D_{s1}(q^2) = -(m_B - m_{i}D_{s1})q(q^2), V_1^{B\ i}D_{s1}(q^2) = -\frac{l(q^2)}{m_B - m_{i}D_{s1}},$$
(14)

$$V_2^{B\ i} D_{s1}(q^2) = (m_B - m_{i} D_{s1}) c_+(q^2), V_3^{B\ i} D_{s1}(q^2) - V_0^{B\ i} D_{s1}(q^2) = \frac{q^2}{2m_{i} D_{s1}} c_-(q^2).$$
(15)

The light-front wave functions (LFWFs) are needed in the form factor calculations. Although the LFWFs can be derived from solving the relativistic Schrödinger equation theoretically, it is difficult to obtained their exact solutions in many cases. Consequently, we will use the phenomenological Gaussian-type wave functions in this work,

$$\varphi' = \varphi'(x_2, p'_{\perp}) = 4 \left(\frac{\pi}{\beta'^2}\right)^{\frac{3}{4}} \sqrt{\frac{dp'_z}{dx_2}} \exp\left(-\frac{p'^2_z + p'^2_{\perp}}{2\beta'^2}\right),$$

$$\varphi'_p = \varphi'_p(x_2, p'_{\perp}) = \sqrt{\frac{2}{\beta'^2}} \varphi', \quad \frac{dp'_z}{dx_2} = \frac{e'_1 e_2}{x_1 x_2 M'_0},$$
(16)

where the parameter β' describes the momentum distribution and is approximately of order Λ_{QCD} . It can be usually determined by the decay constants through the following analytic expressions [48, 51],

$$f_{D_0^*} = \frac{N_c}{16\pi^3} \int dx_2 d^2 p'_{\perp} \frac{h'_{D_0^*}}{x_1 x_2 \left(M'^2 - M_0'^2\right)} 4\left(m'_1 x_2 - m_2 x_1\right), \tag{17}$$

$$f_{{}^{3}D_{s1}} = -\frac{N_c}{4\pi^3 M'} \int dx_2 d^2 p'_{\perp} \frac{h'_{{}^{3}D_{s1}}}{x_1 x_2 \left(M'^2 - M'_0^2\right)} \\ \times \left[x_1 M'_0^2 - m'_1 \left(m'_1 + m_2\right) - p'^2_{\perp} - \frac{m'_1 - m_2}{w'_{{}^{3}D_{s1}}} p'^2_{\perp} \right],$$
(18)

$$f_{{}^{1}D_{s1}} = \frac{N_c}{4\pi^3 M'} \int dx_2 d^2 p'_{\perp} \frac{h'_{{}^{1}D_{s1}}}{x_1 x_2 \left(M'^2 - M_0'^2\right)} \left(\frac{m'_1 - m_2}{w'_{{}^{1}D_{s1}}} p'_{\perp}^2\right), \tag{19}$$

where m'_1 and m_2 represent the constituent quarks of the states D_0^* , ${}^3D_{s1}$ and ${}^1D_{s1}$. The decay constants can be obtained through experimental measurements for the purely leptonic decays or theorical calculations. The explicit forms of h'_M are given by [51]

$$h'_{D_0^*} = \sqrt{\frac{2}{3}} h'_{3D_{s1}} = \left(M'^2 - M_0'^2\right) \sqrt{\frac{x_1 x_2}{N_c}} \frac{1}{\sqrt{2}\widetilde{M}_0'} \frac{\widetilde{M}_0'^2}{2\sqrt{3}M_0'} \varphi'_p, \tag{20}$$

$$h'_{{}^{1}D_{s1}} = \left(M^{2\prime} - M_{0}^{\prime 2}\right) \sqrt{\frac{x_{1}x_{2}}{N_{c}}} \frac{1}{\sqrt{2}\widetilde{M}_{0}^{\prime}} \varphi'_{p}.$$
(21)

B. Form factors

For the general $B \to M$ transitions with M being a scalar or axial-vector meson [51], the decay amplitude at the lowest order is

$$\mathcal{M}^{BM} = -i^3 \frac{N_c}{(2\pi)^4} \int d^4 p_1' \frac{H_B'(H_M'')}{N_1' N_1'' N_2} S^{BM}_{\mu}, \qquad (22)$$

where $N_1^{\prime(\prime\prime)} = p_1^{\prime(\prime\prime)2} - m_1^{\prime(\prime\prime)2}$, $N_2 = p_2^2 - m_2^2$ arise from the quark propagators. For our considered transitions $B \to D_0^*$ and $B \to {}^1D_{s1}$, ${}^3D_{s1}$, the traces $S_{\mu}^{BD_0^*}, S_{\mu\nu}^{B^{-1}D_{s1}}$ and $S_{\mu\nu}^{B^{-3}D_{s1}}$

can be directly obtained by using the Lorentz contraction as follows

$$S_{\mu}^{BD_{0}^{*}} = Tr\left[\left(\not p_{1}^{\prime\prime} + m_{1}^{\prime\prime}\right)\gamma_{\mu}\gamma_{5}\left(\not p_{1}^{\prime} + m_{1}^{\prime}\right)\gamma_{5}\left(-\not p_{2} + m_{2}\right)\right],$$

$$S_{\mu}^{B^{1}D_{s1}} = \left(S_{\mu}^{B^{1}D_{s1}} - S_{\mu}^{B^{1}D_{s1}}\right)$$
(23)

$$S_{\mu\nu}^{-2} = \left(S_{V}^{-1} - S_{A}^{-1} - S_{A}^{-1}\right)_{\mu\nu}$$

$$= \operatorname{Tr}\left[\left(-\frac{1}{W_{1D_{s1}}''}(p_{1}'' - p_{2})_{\nu}\right)\gamma_{5}(\not p_{1}'' + m_{1}'')(\gamma_{\mu} - \gamma_{\mu}\gamma_{5})(\not p_{1}' + m_{1}')\gamma_{5}(-\not p_{2} + m_{2})\right], \quad (24)$$

$$S_{\mu\nu}^{B^{3}D_{s1}} = \left(S_{V}^{B^{3}D_{s1}} - S_{A}^{B^{3}D_{s1}}\right)_{\mu\nu}$$

= Tr $\left[\left(\gamma_{\nu} - \frac{1}{W_{3D_{s1}}''} \left(p_{1}'' - p_{2}\right)_{\nu}\right)\gamma_{5}\left(\not p_{1}'' + m_{1}''\right)\left(\gamma_{\mu} - \gamma_{\mu}\gamma_{5}\right)\left(\not p_{1}' + m_{1}'\right)\gamma_{5}\left(-\not p_{2} + m_{2}\right)\right].$ (25)

To calculate the amplitudes for the transition form factors, we need the Feynman rules for the meson-quark-antiquark vertices $(i\Gamma'_M)$, which are listed as

$$i\Gamma'_{D_0^*} = -iH'_{D_0^*},\tag{26}$$

$$i\Gamma'_{{}^{3}D_{s1}} = -iH'_{{}^{3}D_{s1}} \left[\gamma_{\mu} + \frac{1}{W'_{{}^{3}D_{s1}}} \left(p'_{1} - p_{2} \right)_{\mu} \right] \gamma_{5}, \qquad (27)$$

$$i\Gamma'_{{}^{1}D_{s1}} = -iH'_{{}^{1}D_{s1}} \left[\frac{1}{W'_{{}^{1}D_{s1}}} \left(p'_{1} - p_{2} \right)_{\mu} \right] \gamma_{5}.$$
(28)

In practice, we employ the light-front decomposition of the Feynman loop momentum and integrate out the minus component using the contour method. Then additional spurious contributions being proportional to the light-like four-vector $\tilde{\omega} = (0, 2, \mathbf{0}_{\perp})$ will appear. While they can be eliminated by including the zero-mode contributions in a proper way. If the covariant vertex functions are not exhibit singularity during integration, the transition amplitudes will capture the singularities in the antiquark propagators. The specific rules for the p^- integration have been derived in Refs. [48, 51], and the relevant ones are summarized in Appendix A. The integration then leads to

$$N_{1}^{\prime(\prime\prime)} \rightarrow \hat{N}_{1}^{\prime(\prime\prime)} = x_{1} \left(M^{\prime(\prime\prime2} - M_{0}^{\prime(\prime\prime)2} \right), H_{B}^{\prime} \rightarrow h_{B}^{\prime}, H_{M}^{\prime\prime} \rightarrow h_{M}^{\prime\prime}, W_{M}^{\prime\prime} \rightarrow w_{M}^{\prime\prime} \quad (\text{ for the states } {}^{3}D_{s1} \text{ and } {}^{1}D_{s1}), \int \frac{d^{4}p_{1}^{\prime}}{N_{1}^{\prime}N_{1}^{\prime\prime}N_{2}} H_{B}^{\prime}H_{M}^{\prime\prime}S^{BM} \rightarrow -i\pi \int \frac{dx_{2}d^{2}p_{\perp}^{\prime}}{x_{2}\hat{N}_{1}^{\prime}\hat{N}_{1}^{\prime\prime}} h_{B}^{\prime}h_{M}^{\prime\prime}\hat{S}^{BM},$$
(29)

where

$$w_{^{\prime\prime}D_{s1}}^{\prime\prime} = \frac{\widetilde{M}_{0}^{\prime\prime2}}{m_{1}^{\prime\prime} - m_{2}}, \quad w_{^{\prime\prime}D_{s1}}^{\prime\prime} = 2, \quad M_{0}^{\prime\prime2} = \frac{p_{\perp}^{\prime\prime2} + m_{1}^{\prime\prime2}}{x_{1}} + \frac{p_{\perp}^{\prime\prime2} + m_{2}^{2}}{x_{2}}, \tag{30}$$

with $p''_{\perp} = p'_{\perp} - x_2 q_{\perp}$ and $\widetilde{M}''_0 = \sqrt{M''_0^2 - (m''_1 - m_2)^2}$. The explicit forms of h''_M have been given in Eq. (20) and Eq. (21).

Using Eq. (20)-Eq. (30) and taking the integration rules given in Refs. [48, 51], the form factors $F_1^{BD_0^*}(q^2)$, $F_0^{BD_0^*}(q^2)$ and $A^{BD_{s1}}(q^2)$, $V_0^{BD_{s1}}(q^2)$, $V_1^{BD_{s1}}(q^2)$, $V_2^{BD_{s1}}(q^2)$ can be obtained directly, which are listed in Appendix C.

C. Vertex Corrections and The Hard Spectator Function

Within the framework of QCD factorization [61], the short-distance nonfactorizable corrections including the vertex corrections and hard spectator interactions are considered. The modifications of the Wilson coefficients $a_{1,2}$ from the vertex corrections are given as

$$a_i(\mu) \to a_i(\mu) + \frac{\alpha_s(\mu)}{4\pi} C_F \frac{C_i(\mu)}{N_c} V_i(M_2), \quad i = 1, 2,$$
(31)

with M_2 being the meson emitted from the weak vertex. The vertex functions $V_{1,2}(M_2)$ are written as [61]

$$V_{1,2}(M_2) = 12 \ln \frac{m_b}{\mu} - 18 + \frac{2\sqrt{2N_c}}{f_{M_2}} \int_0^1 dx \,\Phi_{M_2}(x) \,g(x), \tag{32}$$

where f_{M_2} and $\Phi_{M_2}(x)$ are the decay constant and the twist-2 meson distribution amplitude of the meson M_2 , respectively. The hard kernel g(x) is

$$g(x) = 3\left(\frac{1-2x}{1-x}\ln x - i\pi\right) + \left[2\operatorname{Li}_{2}(x) - \ln^{2} x + \frac{2\ln x}{1-x} - (3+2i\pi)\ln x - (x\leftrightarrow 1-x)\right].$$
(33)

The modifications of the Wilson coefficients $a_{1,2}$ from the hard spectator-scattering corrections arising from a hard gluon exchange between the emitted meson and the spectator quark are written as

$$a_1(\mu) \to a_1(\mu) + \frac{C_F \pi \alpha_s C_2}{N_c^2} H_1(M_1 M_2), \quad a_2(\mu) \to a_2(\mu) + \frac{C_F \pi \alpha_s C_1}{N_c^2} H_2(M_1 M_2), \quad (34)$$

where the hard spectator functions $H_i(i = 1, 2)$ are defined as [62]

$$H_{i}(M_{1}M_{2}) = -\frac{f_{B}f_{M_{1}}}{D(M_{1}M_{2})} \int_{0}^{1} \frac{d\rho}{\rho} \Phi_{B}(\rho) \int_{0}^{1} \frac{d\xi}{\bar{\xi}} \Phi_{M_{2}}(\xi) \int_{0}^{1} \frac{d\eta}{\bar{\eta}} \left[\Phi_{M_{1}}(\eta) + r_{\chi}^{M_{1}} \frac{\bar{\xi}}{\xi} \Phi_{M_{1}}^{P}(\eta) \right], (35)$$

with $\bar{\xi} = 1 - \xi$ and $\bar{\eta} = 1 - \eta$. Φ_{M_1} and $\Phi_{M_1}^P$ are the twist-2 and twist-3 LCDAs of the meson M_1 . The definations of $D(M_1M_2)$ and $r_{\chi}^{M_1}$ can be found in Ref. [62].

III. NUMERICAL RESULTS AND DISCUSSIONS

A. Transition Form Factors

			*		
Mass(GeV)	$m_b = 4.8$	$m_c = 1.4$	$m_s = 0.37$	$m_{u,d} = 0.25$	$m_B = 5.279$
	$m_{\pi} = 0.140$	$m_K = 0.494$	$m_{\rho} = 0.775$	$m_{K^*} = 0.892$	$m_{B_s} = 5.367$
	$m_{D_0^*} = 2.343$	$m_{D_{s0}^*} = 2.317$	$m_D = 1.86966$	$m_{D_s} = 1.96835$	
	$m_{D_{s1}} = 2.460$	$m_{D_{s1}^{\prime}}=2.536$	$m_{D_s^*} = 2.1122$	$m_{D^*} = 2.010$	
Decay consta	ants(GeV)	$f_{\pi} = 0.1$	3	$f_{\rho} = 0.209 \pm 0$).002
		$f_{K} = 0.16$		$f_{K^*} = 0.217 \pm 0.005$	
		$f_B = 0.19 \pm 0.02$		$f_{B_s} = 0.231 \pm 0.015$	
		$f_D = 0.2046 \pm 0.005$		$f_{D_s} = 0.2575 \pm$	0.0046
		$f_{D^*} = 0.245 \pm 0$	$0.02^{+0.003}_{-0.002}$	$f_{D_s^*} = 0.272 \pm 0.0$	$16^{+0.003}_{-0.020}$
		$f_{D_{s1}} = 0.145$	± 0.011	$f_{D_{s1}'} = 0.032 \pm$	0.006
		$f_{D_{s1}^{3/2}} = 0.05$		$f_{D_{s1}^{1/2}} = 0.14$	45
		$f_{^3D_{s1}} = -0.121$		$f_{^{1}D_{s1}} = 0.03$	38
		$f_{D_0^*} = 0.103 \pm$	= 0.021	$f_{D_{s0}^*} = 0.067 \pm$	0.013

TABLE I: The values of the input parameters Refs. [63–67].

The input parameters, such as the masses of the initial and final mesons, the decay constants, are listed in Table I. The decay constants of the axial mesons D_{s1} and D'_{s1} can be obtained from $f_{D_{s1}^{1/2}}$ and $f_{D_{s1}^{3/2}}$ through the mixing between $D_{s1}^{1/2}$ and $D_{s1}^{3/2}$

$$f_{D_{s1}} = f_{D_{s1}^{1/2}} \cos \theta_s + f_{D_{s1}^{3/2}} \sin \theta_s,$$

$$f_{D_{s1}'} = -f_{D_{s1}^{1/2}} \sin \theta_s + f_{D_{s1}^{3/2}} \cos \theta_s.$$
(36)

Using these decay constants and the masses of the constituent quarks and mesons given in Table I, we can obtain the values of the shape parameters β' for our considered mesons, which are listed in Table II.

β'_{π}	β'_K	$\beta'_{ ho}$	β'_{K^*}	β_D'
0.317	0.37	$0.261\substack{+0.001\\-0.002}$	0.279 ± 0.004	$0.464\substack{+0.011\\-0.014}$
β_B'	β'_{B_s}	β'_{D_s}	β'_{D^*}	$\beta'_{D_s^*}$
$0.555\substack{+0.048\\-0.048}$	$0.628\substack{+0.035\\-0.034}$	$0.497\substack{+0.032\\-0.028}$	$0.409\substack{+0.021\\-0.022}$	$0.438\substack{+0.016\\-0.027}$
$\beta_{D_0^*}'$	$\beta_{D_{s0}^*}'$	${\beta'_3}_{D_{s1}}$	$\beta_{^{1}D_{s1}^{\prime}}^{\prime}$	
$0.373\substack{+0.063\\-0.059}$	$0.325\substack{+0.043\\-0.043}$	$0.342\substack{+0.030\\-0.034}$	$0.342\substack{+0.039\\-0.039}$	

TABLE II: The shape parameters β' (in units of GeV) in the Gaussian-type light-front wave functions defined in Eq. (16), and the uncertainties are from the decay constants.

It is noticed that all the computations are conducted within the $q^+ = 0$ reference frame, where the form factors can only be obtained at spacelike momentum transfers $q^2 = -q_{\perp}^2 \leq 0$. It is necessary to know the form factors in the timelike region for the physical decay processes. Here, we utilize the following double-pole approximation to parameterize the form factors in the spacelike region and then extend to the timelike region,

$$F(q^2) = \frac{F(0)}{1 - aq^2/m^2 + bq^4/m^4},$$
(37)

where *m* represents the initial meson mass and $F(q^2)$ denotes the different form factors. The values of *a* and *b* can be obtained by performing a 3-parameter fit to the form factors in the range $-15 \text{GeV}^2 \leq q^2 \leq 0$, which are collected in Tables III and V. The uncertainties arise from the decay constants of the initial $B_{(s)}$ meson and the final state mesons.

In Table III, we list the form factors of the transitions $B_{(s)} \to D_0^*, D_{s0}^*, D_{(s)}, D_{(s)}^*$. One can find that the form factors of the transitions $B_{(s)} \to D_0^*, D_{s0}^*$ are much smaller. This conclusion is also supported by other works, for example, the form factor of the transition $B \to D_0^*$ was obtained as 0.24 and 0.18 within the CLFQM [51] and the 2nd version of the Isgur-Scora-Grinstein-Wise approach (ISGW2) [66]. Furthermore, our result for the form factor of the transition $B_s \to D_{s0}^*$ is consistent with 0.20 given in the ISGW2 model [66], while smaller than 0.40 given by the QCD sum rules (QCDSR) approach [68]. As to the form factors of the transitions $B_{(s)} \to D_{(s)}, D_{(s)}^*, \pi(\rho), K(K^*)$, they have been searched by many appoaches, such as the Melikhov-Stech (MS) model [69], the relaticistic quark model (RQM) [70], the BSW model [60, 71], the Bethe-Salpeter (BS) equation [72], the QCDSR [73] and the light cone sum rules (LSCR) approach [74]. Cosidering the need for the latter branching ratio calcuations, we also give them in Table IV with other theoretical results for comparison. Obviously, our predictions are consistent well with these theoretical results.

TABLE III: Form factors of the transitions $B_{(s)} \to D_0^*, D_{s0}^*, D_{(s)}, D_{(s)}^*$ in the CLFQM. The uncertainties are from the decay constants of $B_{(s)}$ and final state mesons.

	$F_i(q^2 = 0)$	$F_i(q_{max}^2)$	a	b
$F_1^{BD_0^*}$	$0.25\substack{+0.03+0.05\\-0.02-0.05}$	$0.30\substack{+0.03+0.06\\-0.03-0.07}$	$0.70_{-0.05-0.11}^{+0.04+0.03}$	$0.65\substack{+0.08+0.03\\-0.07-0.07}$
$F_0^{BD_0^*}$	$0.25\substack{+0.03+0.05\\-0.02-0.05}$	$0.22\substack{+0.02+0.04\\-0.01-0.04}$	$-0.38\substack{+0.04+0.05\\-0.04-0.02}$	$0.21\substack{+0.07+0.08\\-0.07-0.08}$
$F_1^{B_s D_{s0}^*}$	$0.21\substack{+0.02+0.04\\-0.01-0.04}$	$0.24\substack{+0.02+0.05\\-0.01-0.05}$	$0.63\substack{+0.05+0.07\\-0.06-0.12}$	$0.78\substack{+0.08+0.01\\-0.09-0.04}$
$F_0^{B_s D_{s0}^*}$	$0.21\substack{+0.02+0.04\\-0.01-0.04}$	$0.18\substack{+0.02+0.03\\-0.01-0.03}$	$-0.43\substack{+0.01+0.01\\-0.00-0.02}$	$0.28\substack{+0.03+0.01\\-0.06-0.04}$
F_1^{BD}	$0.66\substack{+0.00+0.00\\-0.01-0.01}$	$0.81\substack{+0.00+0.00\\-0.01-0.02}$	$0.80\substack{+0.01+0.00\\-0.01-0.01}$	$0.86\substack{+0.03+0.01\\-0.03-0.01}$
F_0^{BD}	$0.66\substack{+0.00+0.00\\-0.01-0.01}$	$0.70\substack{+0.02+0.00\\-0.02-0.01}$	$0.46\substack{+0.01+0.01\\-0.00-0.00}$	$0.78\substack{+0.10+0.03\\-0.10-0.02}$
$F_1^{B_s D_s}$	$0.65\substack{+0.00+0.01\\-0.00-0.02}$	$0.79\substack{+0.00+0.01\\-0.00-0.03}$	$0.84\substack{+0.01+0.01\\-0.01-0.02}$	$1.02\substack{+0.03+0.04\\-0.04-0.05}$
$F_0^{B_s D_s}$	$0.65\substack{+0.00+0.01\\-0.00-0.02}$	$0.68\substack{+0.01+0.01\\-0.01-0.03}$	$0.50\substack{+0.01+0.01\\-0.01-0.02}$	$0.99\substack{+0.07+0.09\\-0.07-0.10}$
V^{BD^*}	$0.73\substack{+0.01+0.01\\-0.01-0.02}$	$0.89\substack{+0.01+0.01\\-0.02-0.03}$	$0.82\substack{+0.01+0.00\\-0.01-0.01}$	$0.91\substack{+0.05+0.02\\-0.05-0.02}$
$A_0^{BD^*}$	$0.67\substack{+0.00+0.02\\-0.01-0.02}$	$0.70\substack{+0.00+0.02\\-0.02-0.03}$	$0.16\substack{+0.02+0.00\\-0.03-0.01}$	$0.15\substack{+0.00+0.00\\-0.01-0.01}$
$A_1^{BD^*}$	$0.63\substack{+0.00+0.01\\-0.00-0.01}$	$0.72\substack{+0.01+0.01\\-0.00-0.01}$	$0.42\substack{+0.02+0.01\\-0.02-0.01}$	$0.22\substack{+0.03+0.02\\-0.01-0.00}$
$A_2^{BD^*}$	$0.59\substack{+0.00+0.00\\-0.01-0.01}$	$0.71\substack{+0.00+0.00\\-0.01-0.01}$	$0.75\substack{+0.01+0.01\\-0.02-0.02}$	$0.78\substack{+0.04+0.03\\-0.04-0.03}$
$V^{B_s D_s^*}$	$0.72\substack{+0.01+0.02\\-0.00-0.02}$	$0.86\substack{+0.02+0.03\\-0.00-0.02}$	$0.86\substack{+0.02+0.01\\-0.02-0.00}$	$1.10\substack{+0.05+0.04\\-0.05-0.02}$
$A_0^{B_s D_s^*}$	$0.65\substack{+0.01+0.02\\-0.00-0.03}$	$0.69\substack{+0.01+0.02\\-0.01-0.04}$	$0.23\substack{+0.02+0.00\\-0.03-0.01}$	$0.21\substack{+0.01+0.02\\-0.01-0.01}$
$A_1^{B_s D_s^*}$	$0.62\substack{+0.00+0.01\\-0.01-0.02}$	$0.72\substack{+0.00+0.00\\-0.02-0.03}$	$0.48\substack{+0.01+0.00\\-0.03-0.02}$	$0.32\substack{+0.03+0.03\\-0.02-0.01}$
$A_2^{B_s D_s^*}$	$0.57\substack{+0.00+0.00\\-0.00-0.00}$	$0.68\substack{+0.00+0.00\\-0.00-0.00}$	$0.80\substack{+0.01+0.01\\-0.02-0.02}$	$0.95\substack{+0.04+0.06\\-0.04-0.03}$

Transitions	References	$F_0(0)$	V(0)	$A_0(0)$	$A_1(0)$	$A_2(0)$
$B \rightarrow D, D^*$	This work	0.66	0.73	0.67	0.63	0.59
/	[51]	0.67	0.75	0.64	0.63	0.62
	[69]	0.67	0.76	0.69	0.66	0.62
	[60]	0.69	0.71	0.62	0.65	0.69
$B_s \to D_s, D_s^*$	This work	0.65	0.57	0.72	0.65	0.62
	[70]	0.74	0.95	0.67	0.70	0.75
	[71]	0.61	0.64	_	0.56	0.59
	[72]	0.57	0.70	0.70	0.65	0.67
	[73]	0.70	0.63	_	0.62	0.75
$B \to \pi, \rho$	This work	0.25	0.25	0.28	0.22	0.19
	[51]	0.25	0.27	0.28	0.22	0.20
	[74]	0.26	0.34	0.37	0.26	0.22
	[69]	0.29	0.31	0.29	0.26	0.24
	[60]	0.33	0.33	0.28	0.28	0.28
$B \to K, K^*$	This work	0.35	0.30	0.33	0.26	0.24
	[51]	0.35	0.31	0.31	0.26	0.24
	[74]	0.34	0.46	0.47	0.34	0.28
	[69]	0.36	0.44	0.45	0.36	0.32
	[60]	0.38	0.37	0.32	0.33	0.33

TABLE IV: Form factors of the transitions $B_{(s)} \to D_{(s)}, D^*_{(s)}, \pi(\rho), K(K^*)$ at $q^2 = 0$ together with other theoretical results.

In order to determine the physical form factors of the transitions $B_s \to D_{s1}, D'_{s1}$, we need to know the mixing angle $\theta = \theta_s + 35.3^\circ$ between ${}^1D_{s1}$ and ${}^3D_{s1}$ shown in Eq. (3). Here we take $\theta_s = 7^\circ$, which was determined from the quark potential model [66]. The results are listed in Table V, where the uncertainties are from the decay constants of B_s and the final states $({}^3D_{s1}, {}^1D_{s1})$. In Figure 2, we check the dependence of the form factors of the transitions $B_s \to D_{s1}, D'_{s1}$ on the mixing angle θ_s . We find that the form factor V_0 of the transition $B_s \to D_{s1}, D'_{s1}$ is (not) sensitive to the mixing angle. It can be used to explain why the branching ratios of the decays associated with the transition $B_s \to D_{s1}$ $(B_s \to D'_{s1})$ are (not) sensitive to the mixing angle, which will be discussed in the latter.

TABLE V: The form factors of the transitions $B_s \to D_{s1}$ and $B_s \to D'_{s1}$ in the CLFQM. The uncertainties are from the decay constants of $B_{(s)}$ and final states.

	$F_i(q^2 = 0)$	$F_i(q_{max}^2)$	a	b
$A^{B_s D_{s1}}$	$-0.08\substack{+0.01+0.02+0.00\\-0.01-0.02-0.01}$	$-0.09\substack{+0.05+0.05+0.01\\-0.06-0.07-0.03}$	$2.05_{-0.10-0.26-0.24}^{+0.13+0.24+0.26}$	$5.57_{-0.20-0.26-0.32}^{+0.25+0.34+0.36}$
$V_0^{B_s D_{s1}}$	$0.08\substack{+0.01+0.01+0.04\\-0.01-0.00-0.04}$	$0.11\substack{+0.02+0.02+0.06\\-0.01-0.00-0.07}$	$1.24\substack{+0.05+0.14+0.18\\-0.06-0.17-0.18}$	$0.74\substack{+0.02+0.16+0.14\\-0.02-0.10-0.14}$
$V_1^{B_s D_{s1}}$	$-0.17\substack{+0.02+0.03+0.02\\-0.03-0.02-0.02}$	$-0.14\substack{+0.01+0.02+0.01\\-0.03-0.02-0.02}$	$-0.52\substack{+0.06+0.02+0.05\\-0.05-0.02-0.05}$	$0.36\substack{+0.01+0.02+0.02\\-0.00-0.03-0.07}$
$V_2^{B_s D_{s1}}$	$-0.11\substack{+0.01+0.00+0.01\\-0.02-0.01-0.02}$	$-0.12\substack{+0.01+0.00+0.01\\-0.02-0.01-0.02}$	$0.25\substack{+0.06+0.00+0.06\\-0.07-0.00-0.07}$	$-0.07\substack{+0.03+0.00+0.01\\-0.04-0.01-0.03}$
$A^{B_s D'_{s1}}$	$0.20\substack{+0.01+0.02+0.01\\-0.01-0.02-0.00}$	$0.18\substack{+0.02+0.02+0.02\\-0.01-0.02-0.00}$	$-0.27\substack{+0.06+0.03+0.07\\-0.07-0.05-0.08}$	$0.11\substack{+0.02+0.01+0.02\\-0.02-0.01-0.03}$
$V_0^{B_s D_{s1}^\prime}$	$0.40\substack{+0.02+0.01+0.04\\-0.02-0.00-0.04}$	$0.42\substack{+0.02+0.01+0.05\\-0.02-0.00-0.05}$	$-0.17\substack{+0.02+0.02+0.03\\-0.04-0.04-0.04}$	$-0.02\substack{+0.01+0.00+0.00\\-0.00-0.00-0.01}$
$V_1^{B_s D'_{s1}}$	$0.58\substack{+0.01+0.02+0.02\\-0.02-0.03-0.03}$	$0.57\substack{+0.01+0.02+0.02\\-0.02-0.03-0.03}$	$-0.05\substack{+0.01+0.01+0.00\\-0.01-0.01-0.00}$	$0.02\substack{+0.00+0.01+0.01\\-0.00-0.00-0.00}$
$V_2^{B_s D_{s1}^{\prime}}$	$-0.05\substack{+0.01+0.01+0.02\\-0.00-0.00-0.01}$	$-0.05\substack{+0.01+0.01+0.03\\-0.00-0.01-0.02}$	$0.56\substack{+0.06+0.12+0.18\\-0.06-0.15-0.20}$	$2.50_{-0.20-0.85-0.98}^{+0.25+1.11+1.25}$




FIG. 2: The dependence of the form factors of the transitions $B_s \to D_{s1}$ and $B_s \to D'_{s1}$ on the mixing angle θ_s .

In Figure 3, we plot the q^2 -dependence of the form factors of the transitions $B_{(s)} \rightarrow D_0^*, D_{s0}^*, D_{s1}, D_{s1}'$. There exists the similar q^2 -dependence of the form factors $F_{0,1}(q^2)$ between the transitions $B \rightarrow D_0^*$ and $B_s \rightarrow D_{s0}^*$. It is consistent with our expectation. While it is very different in magnitude between the corresponding form factors for the transitions $B_s \rightarrow D_{s1}$ and $B_s \rightarrow D_{s1}'$.




FIG. 3: The q^2 -dependence of the form factors of the transitions $B_{(s)} \rightarrow D_0^*, D_{s0}^*, D_{s1}, D_{s1}'$.

B. Branching ratios

In addition to the parameters listed in Table I, other inputs, such as the $B_{(s)}$ meson lifetime $\tau_{B_{(s)}}$, the Wilson coefficients a_1, a_2 and the CabibboKobayashi-Maskawa (CKM) matrix elements, are listed as [64, 75]

$$\tau_{B^{\pm}} = (1.519 \pm 0.004) \times 10^{-12} s, \tau_{B_s} = (1.520 \pm 0.005) \times 10^{-12} s \tag{38}$$

$$\tau_{B_0} = (1.638 \pm 0.004) \times 10^{-12} s, a_1 = 1.018, a_2 = 0.17,$$
(39)

$$V_{cd} = 0.221 \pm 0.004, V_{cb} = (40.8 \pm 1.4) \times 10^{-3}, V_{cs} = 0.975 \pm 0.006$$
(40)

$$V_{us} = 0.2243 \pm 0.0008, V_{ud} = 0.97373 \pm 0.00031.$$
(41)

Firstly, we consider the branching ratios of the decays $B_{(s)} \to D^*_{(s)0}M$ with M being a light pseudoscalar (vector) meson P(V) or a charmed meson $(D^{(*)}, D^{(*)}_s)$, which can be calculated through the formula

$$\mathcal{B}r(B_{(s)} \to D^*_{(s)0}M) = \frac{\tau_{B_{(s)}}}{\hbar}\Gamma(B_{(s)} \to D^*_{(s)0}M),$$
(42)

where the decay width $\Gamma(B_{(s)} \to D_0^*(D_{s0}^*)M)$ for each channel is given as following

$$\Gamma\left(B^{0} \to D_{0}^{*-}P(V)\right) = \frac{\left|G_{F}V_{cb}V_{uq}^{*}a_{1}f_{P(V)}m_{B}^{2}F_{0}^{BD_{0}^{*}}(m_{P(V)}^{2})\right|^{2}}{32\pi m_{B}}\left(1-r_{D_{0}^{*}}^{2}\right),\tag{43}$$

$$\Gamma\left(B^{0} \to D_{0}^{*-}D^{(*)+}\right) = \frac{\left|G_{F}V_{cb}V_{cd}^{*}a_{1}f_{D^{(*)}}m_{B}^{2}F_{0}^{BD_{0}^{*}}(m_{D^{(*)}}^{2})\right|^{2}}{32\pi m_{B}}\left(1 - r_{D_{0}^{*}}^{2} - r_{D^{(*)}}^{2}\right), \quad (44)$$

where P(V) refers to $\pi, K(\rho, K^*)$ and the subscript q = d(s). $\Gamma\left(B^0 \to D_0^{*-} D_s^{(*)+}\right)$ can be obtained by replacing $V_{cd}, D^{(*)}$ with $V_{cs}, D_s^{(*)}$ in Eq. (44). There exists similar expressions for the cases with B^0, D_0^{*-} replaced by B_s^0, D_{s0}^{*-} in the upper decays. While for the charged Bdecays $B^+ \to \overline{D}_0^{*0} P(V)$ with P(V) being $\pi^+, K^+(\rho^+, K^{*+})$, the corresponding decay widths should be written as

$$\Gamma\left(B^{+} \to \bar{D}_{0}^{*0}\pi^{+}(K^{+})\right) = \frac{\left(G_{F}V_{cb}^{*}V_{uq}m_{B}^{2}\right)^{2}\left(1-r_{D_{0}^{*}}^{2}\right)}{32\pi m_{B}} \times \left|a_{1}f_{\pi(K)}F_{0}^{BD_{0}^{*}}(m_{\pi(K)}^{2})+a_{2}f_{D_{0}^{*}}F_{0}^{B\pi(K)}(m_{D_{0}^{*}}^{2})\right|^{2}, \quad (45)$$

$$\Gamma\left(B^{+} \to \bar{D}_{0}^{*0}\rho^{+}(K^{*+})\right) = \frac{\left(G_{F}V_{cb}^{*}V_{uq}m_{B}^{2}\right)^{2}\left(1-r_{D_{0}^{*}}^{2}\right)}{32\pi m_{B}} \times \left|a_{1}f_{\rho(K^{*})}F_{0}^{BD_{0}^{*}}(m_{\rho(K^{*})}^{2})+a_{2}f_{D_{0}^{*}}A_{0}^{B\rho(K^{*})}(m_{D_{0}^{*}}^{2})\right|^{2}. \quad (46)$$

Secondly, the decay widths of the decays $B_s \to D_{s1}^{(\prime)} M$ are defined as

$$\Gamma\left(B_s \to D_{s1}^{(\prime)}\pi(K)\right) = \frac{\left|G_F V_{cb} V_{uq}^* a_1 f_{\pi(K)} m_{B_s}^2 V_0^{B_s D_{s1}^{(\prime)}} (m_{\pi(K)}^2)\right|^2}{32\pi m_{B_s}} \left(1 - r_{D_{s1}^{(\prime)}}^2\right), \quad (47)$$

$$\Gamma\left(B_{s} \to D_{s1}^{(\prime)}D_{(s)}\right) = \frac{\left|G_{F}V_{cb}V_{cq}^{*}a_{1}f_{D_{(s)}}m_{B_{s}}^{2}V_{0}^{B_{s}D_{s1}^{(\prime)}}(m_{D_{(s)}}^{2})\right|^{2}}{32\pi m_{B_{s}}}\left(1 - r_{D_{s1}^{\prime}}^{2} - r_{D_{(s)}}^{2}\right), \quad (48)$$

$$\Gamma\left(B_{s} \to D_{s1}^{(\prime)}V\right) = \frac{\left|\vec{p}\right|}{8\pi m_{B_{s}}^{2}}\left(\left|\mathcal{A}_{L}(B_{s} \to D_{s1}^{(\prime)}V)\right|^{2} + 2\left|\mathcal{A}_{N}\left(B_{s} \to D_{s1}^{(\prime)}V\right)\right|^{2}\right)$$

$$+ 2\left|\mathcal{A}_{T}\left(B_{s} \to D_{s1}^{(\prime)}V\right)\right|^{2}\right), \quad (49)$$

where V represents a vector meson, such as $\rho, K^*, D^*_{(s)}$, and the summation of the three polarizations for the decays $B_s \to D^{(\prime)}_{s1} V$ are performed. The three-momentum \vec{p} is defined as

$$|\vec{p}| = \frac{\sqrt{\left(m_{B_s}^2 - \left(m_{D_{s1}^{(\prime)}} + m_V\right)^2\right)\left(m_{B_s}^2 - \left(m_{D_{s1}^{(\prime)}} - m_V\right)^2\right)}}{2m_{B_s}},\tag{50}$$

and the three polarization amplitudes $\mathcal{A}_L, \mathcal{A}_N$, and \mathcal{A}_T are given as

$$i\mathcal{A}_{L}(B_{s} \to D_{s1}^{(\prime)}V) = \frac{(-i)^{3}G_{F}}{\sqrt{2}}V_{cb}V_{q_{1}q_{2}}^{*}a_{1}f_{V}m_{B_{s}}^{2}\frac{1}{2r_{D_{s1}^{(\prime)}}} \\ \times \left[(1 - r_{V}^{2} - r_{D_{s1}^{(\prime)}}^{2})(1 - r_{D_{s1}^{(\prime)}})V_{1}^{B_{s}D_{s1}^{(\prime)}}(m_{V}^{2}) - \frac{\lambda(1, r_{V}^{2}, r_{D_{s1}^{\prime}}^{2})}{1 - r_{D_{s1}^{\prime}}}V_{2}^{B_{s}D_{s1}^{\prime\prime}}(m_{V}^{2}) \right], \\ i\mathcal{A}_{N}(B_{s} \to D_{s1}^{(\prime)}V) = \frac{(-i)^{3}G_{F}}{\sqrt{2}}V_{cb}V_{q_{1}q_{2}}^{*}a_{1}f_{V}m_{B_{s}}^{2}r_{V}(1 - r_{D_{s1}^{\prime\prime}})V_{1}^{B_{s}D_{s1}^{\prime\prime}}(m_{V}^{2}), \\ i\mathcal{A}_{T}(B_{s} \to D_{s1}^{\prime\prime}V) = \frac{-iG_{F}}{\sqrt{2}}V_{cb}V_{q_{1}q_{2}}^{*}a_{1}f_{V}m_{B_{s}}^{2}r_{V}\frac{\sqrt{\lambda(1, r_{V}^{2}, r_{D_{s1}^{\prime\prime}}^{2})}}{1 - r_{D_{s1}^{\prime\prime}}}A^{B_{s}D_{s1}^{\prime\prime}}(m_{V}^{2}),$$

$$(51)$$

with $q_1 = u, c$ and $q_2 = d, s, \lambda \left(1, r_V^2, r_{D_{s1}}^2\right) = \left(1 + r_V^2 - r_{D_{s1}}^2\right)^2 - 4r_V^2$. As to the decay widths of the decays $B \to D_{s1}^{(\prime)}D$ and $B \to D_{s1}^{(\prime)}D^*$, they can be obtained from Eqs. (48) and (49) with some simple replacements, respectively. Using the upper listed input parameters and the formulas given in Eq. (42)-Eq. (51), one can calculate the branching ratios for the considered decays shown in Tables VI-XIII.

TABLE VI: The CLFQM predictions for the branching ratios of the decays $B \to D_0^* \pi(K)$. The label LO, VC and HSSC mean the inclusions of the leading order, the vertex corrections and the hard spectator-scattering corrections, respectively. NLO means the inclusions of these two kinds of corrections at the same time. The upper (lower) line is corresponding to $f_{D_0^*} = 0.078(0.103)$ GeV for each decay. The first and second uncertainties are from the decay constants of B and D_0^* .

	LO	VC	HSSC	NLO	Other predictions	Refs.
$10^{-4}\times \mathcal{B}r(B^+\to \bar{D}_0^{*0}\pi^+)$	$2.98\substack{+1.07+1.72\\-0.83-1.44}$	$4.68^{+1.48+2.15}_{-1.09-1.90}$	$2.60\substack{+1.00+1.61\\-0.77-1.33}$	$4.18\substack{+1.27+2.03\\-1.02-1.77}$	8.93	PQCD [76]
					7.3	CLFQM [51]
	$5.45\substack{+1.24+2.53\\-1.09-2.36}$	$8.49\substack{+1.57+3.16\\-1.42-3.08}$	$4.71^{+1.15+2.35}_{-1.10-2.16}$	$7.59\substack{+1.48+2.98\\-1.32-2.88}$	7.7	ISGW2 [66]
					4.2	ISGW [77]
$10^{-4}\times \mathcal{B}r(B^0\to D_0^{*-}\pi^+)$	$3.68\substack{+1.03+1.82\\-0.91-1.58}$	$3.83_{-1.18-1.65}^{+0.95+1.89}$	$3.59\substack{+1.11+1.77\\-0.89-1.55}$	$3.75\substack{+1.16+1.85\\-0.93-1.61}$	4.28	PQCD [76]
	$6.69\substack{+1.32+2.66\\-1.18-2.56}$	$6.97\substack{+1.37+2.78\\-1.23-2.67}$	$6.54\substack{+1.29+2.60\\-1.15-2.51}$	$6.81\substack{+1.34+2.71\\-1.20-2.61}$	2.6	ISGW2 [66]
					4.1	ISGW [77]
$10^{-5}\times \mathcal{B}r(B^+\to \bar{D}_0^{*0}K^+)$	$2.25\substack{+0.90+1.32\\-0.68-1.11}$	$3.74\substack{+1.07+1.70\\-0.86-1.51}$	$1.94\substack{+0.76+1.23\\-0.58-1.01}$	$3.20\substack{+0.98+1.57\\-0.78-1.37}$	6.96	PQCD [76]
	$4.12\substack{+0.95+1.95\\-0.83-1.81}$	$6.78^{+1.24+2.50}_{-1.12-2.43}$	$3.57\substack{+0.88+1.81\\-0.76-1.66}$	$6.08\substack{+1.17+2.35\\-1.05-2.27}$		
$10^{-5}\times \mathcal{B}r(B^0\to D_0^{*-}K^+)$	$2.91\substack{+0.90+1.44\\-0.72-1.25}$	$3.03\substack{+0.75+1.50\\-0.94-1.30}$	$2.82\substack{+0.87+1.39\\-0.70-1.21}$	$2.94\substack{+0.91+1.45\\-0.73-1.26}$	3.57	PQCD [76]
	$5.29\substack{+1.04+2.11\\-0.93-2.03}$	$5.52\substack{+1.09+2.20\\-0.97-2.11}$	$5.13\substack{+1.01+2.04\\-1.00-1.97}$	$5.35\substack{+1.05+2.13\\-0.94-2.05}$		

From Table VI, one can find that the branching ratios of the decays $B^+ \to \bar{D}_0^{*0}\pi^+$ and $B^+ \to \bar{D}_0^{*0}K^+$ are sensitive to the vertex corrections. These two channels are contributed by two kinds of Feynman diagrams, one is associated with the $B \to D_0^*$ transition accompanied by the emission of a light pseudoscalar meson (π, K) , the other is associated with the $B \to \pi(K)$ transition accompanied by the scalar meson D_0^* emission. We can find that the former

gives the dominant contribution. The other two decay channels $B^0 \to D_0^{*-} \pi^+$ and $B^0 \to D_0^{*-} K^+$ only receive one kind of Feynman diagram contribution to the their branching ratios, which is associated with the $B \to D_0^*$ transition accompanied by a light pseudoscalar meson (π, K) emission. In any case, the contributions from the vertex corrections and the hard spectator-scatterings are destructive with each other. In view of the large uncertainty from the decay constant $f_{D_0^*}$ as mentioned before, two decay constant values are used in our calculations, which are shown in Table VI. For each decay, the upper line is the result corresponding to $f_{D_0^*} = 0.078$ GeV, and the lower is the result corresponding to $f_{D_0^*} = 0.103$ GeV. We can find that the branching ratios are sensitive to the decay constant $f_{D_0^*}$. Our predictions are consistent with other theoretical calculations, such as the PQCD approach [76], the ISGW quark model [66, 77]. Certainly, the branching ratio of the decay $B^+ \to \overline{D}_0^{*0}\pi^+$ was also calculated using the CLFQM in previous [51], which is agreement with our result.

TABLE VII: The CLFQM predictions for the branching ratios of the decays $B \to D_0^* \pi(K) \to D\pi\pi(K)$, where the upper (lower) line is corresponding to $f_{D_0^*} = 0.078(0.103)$ GeV for each decay. The first and second uncertainties are from the decay constants of B and D_0^* .

	This work	PQCD $[76]$	Data
	$2.79_{-0.68-1.18}^{+0.85+1.35}$		$6.1 \pm 0.6 \pm 0.9 \pm 1.6$ Belle [3]
$10^{-4} \times \mathcal{B}r(B^+ \to \bar{D}_0^{*0}\pi^+ \to D^-\pi^+\pi^+)$		$5.95^{+3.14}_{-2.32}$	$6.8\pm 0.3\pm 0.4\pm 2.0$ BaBar [32]
	$5.06\substack{+0.99+1.99\\-0.88-1.92}$		$5.78 \pm 0.08 \pm 0.06 \pm 0.09$ LHCb [78]
	$2.50\substack{+0.77+1.23\\-0.82-1.07}$		$0.60 \pm 0.13 \pm 0.15 \pm 0.22$ Belle [79]
$10^{-4} \times \mathcal{B}r(B^0 \to D_0^{*-} \pi^+ \to \bar{D^0} \pi^- \pi^+)$		$2.85^{+1.65}_{-1.18}$	$0.77 \pm 0.05 \pm 0.03 \pm 0.03$ LHCb [80]
	$4.54_{-0.80-1.74}^{+0.89+1.81}$		$0.80 \pm 0.05 \pm 0.08 \pm 0.04$ LHCb [80]
$10^{-5} \times \mathcal{B}r(B^+ \to \bar{D}_0^{*0}K^+ \to D^-\pi^+K^+)$	$2.13\substack{+0.65+1.05\\-0.52-0.91}$	$4.65_{-1.85}^{+2.46}$	$0.61 \pm 0.19 \pm 0.05 \pm 0.14 \pm 0.04$ LHCb [81]
	$4.05\substack{+0.78+1.57\\-0.70-1.51}$		
$10^{-5} \times \mathcal{B}r(B^0 \to D_0^{*-}K^+ \to \bar{D^0}\pi^-K^+)$	$1.96\substack{+0.61+0.97\\-0.49-0.84}$	$2.38^{+1.31}_{-0.98}$	$1.77 \pm 0.26 \pm 0.19 \pm 0.67 \pm 0.20$ LHCb [82]
	$3.57\substack{+0.70+1.42\\-0.63-1.37}$		

The branching ratio of the quasi-two-body decay $B \to D_0^* P \to D\pi P$ can be obtained from the corresponding two body decay $B \to D_0^* P$ under the narrow width approximation $\mathcal{B}r(B \to D_0^* P \to D\pi P) = \mathcal{B}r(B \to D_0^* P)\mathcal{B}r(D_0^* \to D\pi)$, where P refers to a light pseudoscalar meson (π, K) . Assuming the D_0^* state decays essentially into $D\pi$, we have $\mathcal{B}r(\bar{D}_0^{*0} \to D^-\pi^+) = \mathcal{B}r(D_0^{*-} \to \bar{D}^0\pi^-) = \frac{2}{3}$ from the Clebsch-Gordan coefficients. The branching ratios of these quasi-two-body decays are collected in Table VII, where the PQCD results and the data are also listed for comparison. One can find that if taken the bigger decay constant $f_{D_0^*} = 0.103$ GeV, the branching fraction for the decay $B^+ \to \bar{D}_0^{*0}\pi^+ \to D^-\pi^+\pi^+$ can agree with the data from Belle [3], BaBar [32] and LHCb [78] within errors. While for the decay $B^0 \to D_0^{*-} K^+ \to \overline{D}^0 \pi^- K^+$, if taken the smaller decay constant $f_{D_0^*} = 0.078$ GeV, our prediction can explain the LHCb measurement [82]. For the other two quasi-two-body decays, the predictions for their branching ratios are much larger than the present data. Similar situation exists for the comparison between the PQCD results and the measurements. Another divergence is that our results for the charged (neutral) channels by using the bigger (smaller) decay constant $f_{D_0^*} = 0.103 (0.078)$ GeV can be consistent well with the PQCD results. Further experimental and theoretical researches are needed to clarify these divergences and puzzles.

TABLE VIII: The CLFQM predictions for the branching ratios (10^{-3}) of the decays $B \rightarrow D_0^* D_{(s)}, D_0^* D_{(s)}^*, D_0^* \rho, D_0^* K^*$. The labels LO, VC, HSSC, NLO and the error sources are the same with those given in Table VI. Other theoretical predictions are also listed for comparison.

	LO	VC	HSSC	NLO	ISGW2 [66]	ISGW [77]
$\mathcal{B}r(B^+\to \bar{D}_0^{*0}D_s^+)$	$1.54_{-0.38-0.66}^{+0.47+0.76}$	$1.52\substack{+0.47+0.75\\-0.38-0.65}$	$1.52\substack{+0.47+0.75\\-0.38-0.65}$	$1.51_{-0.37-0.65}^{+0.46+0.74}$	0.80	2.7
	$2.79_{-0.49-1.07}^{+0.55+1.11}$	$2.76\substack{+0.54+1.10\\-0.49-1.06}$	$2.76\substack{+0.55+1.10\\-0.49-1.06}$	$2.74_{-0.48-1.05}^{+0.54+1.09}$		
$\mathcal{B}r(B^0\to D_0^{*-}D_s^+)$	$1.42\substack{+0.44+0.70\\-0.35-0.61}$	$1.41\substack{+0.43+0.70\\-0.35-0.61}$	$1.41\substack{+0.44+0.70\\-0.35-0.61}$	$1.40\substack{+0.43+0.69\\-0.34-0.60}$	0.73	2.6
	$2.59\substack{+0.51+1.03\\-0.46-0.99}$	$2.56\substack{+0.51+1.02\\-0.45-0.98}$	$2.53\substack{+0.50+1.01\\-0.45-0.97}$	$2.51\substack{+0.49+1.00\\-0.44-0.96}$		
$\mathcal{B}r(B^+\to \bar{D}_0^{*0}D_s^{*+})$	$1.71\substack{+0.53+0.85\\-0.42-0.74}$	$1.70\substack{+0.52+0.84\\-0.42-0.73}$	$1.73\substack{+0.53+0.85\\-0.43-0.74}$	$1.71\substack{+0.53+0.84\\-0.42-0.73}$	0.35	_
	$3.11_{-0.55-1.19}^{+0.61+1.24}$	$3.08^{+0.61+1.23}_{-0.54-1.18}$	$3.14_{-0.55-1.20}^{+0.62+1.25}$	$3.11_{-0.55-1.19}^{+0.61+1.24}$		
$\mathcal{B}r(B^0\to D_0^{*-}D_s^{*+})$	$1.59\substack{+0.49+0.78\\-0.39-0.68}$	$1.57\substack{+0.49+0.78\\-0.39-0.68}$	$1.60\substack{+0.49+0.80\\-0.39-0.69}$	$1.58\substack{+0.49+0.78\\-0.39-0.68}$	0.32	-
	$2.89\substack{+0.57+1.15\\-0.51-1.11}$	$2.86\substack{+0.56+1.14\\-0.50-1.10}$	$2.86\substack{+0.56+1.14\\-0.50-1.10}$	$2.83\substack{+0.56+1.13\\-0.50-1.09}$		
$\mathcal{B}r(B^+\to \bar{D}_0^{*0}\rho^+)$	$0.81\substack{+0.28+0.44\\-0.22-0.38}$	$0.85\substack{+0.29+0.47\\-0.22-0.40}$	$0.89\substack{+0.29+0.46\\-0.23-0.40}$	$0.94\substack{+0.30+0.48\\-0.24-0.42}$	1.30	-
	$1.47\substack{+0.32+0.65\\-0.29-0.61}$	$2.03\substack{+0.38+0.77\\-0.34-0.75}$	$1.63\substack{+0.34+0.68\\-0.30-0.65}$	$2.21\substack{+0.39+0.80\\-0.36-0.78}$		
$\mathcal{B}r(B^0\to D_0^{*-}\rho^+)$	$0.94\substack{+0.29+0.46\\-0.23-0.40}$	$0.97\substack{+0.30+0.48\\-0.24-0.42}$	$0.95\substack{+0.29+0.47\\-0.23-0.41}$	$0.99\substack{+0.30+0.49\\-0.24-0.42}$	0.64	—
	$1.70^{+0.34+0.68}_{-0.30-0.65}$	$1.77\substack{+0.35+0.71\\-0.31-0.68}$	$1.66\substack{+0.33+0.66\\-0.29-0.64}$	$1.73_{-0.31-0.66}^{+0.34+0.69}$		
$\mathcal{B}r(B^+\to \bar{D}_0^{*0}K^{*+})$	-0.012-0.021	$0.047\substack{+0.016+0.026\\-0.013-0.022}$	$0.049\substack{+0.016+0.026\\-0.013-0.022}$	$0.052\substack{+0.017+0.027\\-0.013-0.023}$	-	-
	$0.082\substack{+0.018+0.037\\-0.016-0.034}$	$0.116\substack{+0.022+0.044\\-0.020-0.043}$	$0.090\substack{+0.019+0.038\\-0.017-0.036}$	$0.126\substack{+0.022+0.045\\+0.020+0.044}$		
$\mathcal{B}r(B^0\to D_0^{*-}K^{*+})$		$0.056\substack{+0.017+0.028\\-0.014-0.024}$	$0.054\substack{+0.017+0.027\\-0.013-0.023}$	$0.056\substack{+0.017+0.028\\-0.014-0.024}$	-	—
	$0.097\substack{+0.019+0.039\\-0.017-0.037}$	$0.101\substack{+0.020+0.040\\-0.018-0.039}$	$0.095\substack{+0.019+0.038\\-0.017-0.036}$	$0.099\substack{+0.020+0.040\\-0.018-0.038}$		
$\mathcal{B}r(B^+ \to \bar{D}_0^{*0}D^+)$	$0.052\substack{+0.016+0.026\\-0.013-0.022}$	$0.051\substack{+0.016+0.025\\-0.013-0.022}$	$0.052\substack{+0.016+0.026\\-0.013-00.22}$	$0.051\substack{+0.016+0.025\\-0.013-0.022}$	-	0.114
	$0.094\substack{+0.019+0.038\\-0.017-0.036}$	$0.093\substack{+0.018+0.037\\-0.016-0.036}$	$0.094\substack{+0.019+0.037\\-0.017-0.036}$	$0.092\substack{+0.018+0.037\\-0.016-0.035}$		
$\mathcal{B}r(B^0\to D_0^{*-}D^+)$	$0.048\substack{+0.015+0.024\\-0.012-0.021}$	$0.047\substack{+0.015+0.023\\-0.012-0.020}$	$0.048\substack{+0.015+0.024\\-0.012-0.021}$	$0.047\substack{+0.015+0.023\\-0.012-0.020}$	-	0.111
	$0.087\substack{+0.017+0.035\\-0.015-0.033}$	$0.086\substack{+0.017+0.034\\-0.015-0.033}$	$0.086\substack{+0.017+0.034\\-0.015-0.033}$	$0.085\substack{+0.017+0.034\\-0.015-0.032}$		
$\mathcal{B}r(B^+\to \bar{D}_0^{*0}D^{*+})$	0.010 0.002	$0.073\substack{+0.023+0.036\\-0.018-0.031}$	$0.075\substack{+0.023+0.037\\-0.019-0.032}$	$0.074\substack{+0.023+0.037\\-0.018-0.032}$	-	-
	$0.135\substack{+0.027+0.054\\-0.024-0.052}$	$0.133\substack{+0.026+0.053\\-0.023-0.051}$	$0.137\substack{+0.027+0.054\\-0.024-0.052}$	$1.340\substack{+0.027+0.054\\-0.024-0.052}$		
$\mathcal{B}r(B^0\to D_0^{*-}D^{*+})$		$0.068\substack{+0.021+0.033\\-0.017-0.029}$	$0.070\substack{+0.022+0.034\\-0.017-0.030}$	$0.069\substack{+0.021+0.034\\-0.017-0.029}$	-	-
	$0.125\substack{+0.024+0.050\\-0.022-0.048}$	$0.123\substack{+0.024+0.049\\-0.022-0.047}$	$0.124\substack{+0.025+0.050\\-0.022-0.048}$	$0.122\substack{+0.024+0.049\\-0.022-0.047}$		

If replaced the light pseudoscalar mesons π, K in the final states with the vector meson ρ, K^* and the charmed meson $D^{(*)}, D_s^{(*)}$, the branching ratios for the conresponding decay

channels are listed in Table VIII. One can find that our predictions by using the bigger decay constant $f_{D_0^*} = 0.103$ GeV are consistent with those given in the ISGW model, it is because that the form factor of the transition $B \to D_0^*$ at maximum momentum transfer obtained in the CLFQM with $f_{D_0^*} = 0.103$ GeV is about 0.30, which is almost equal to the value calculated in the ISGW model [77], while much larger than 0.18 given in the ISGW2 model [66]. The differences between the ISGW and ISGW2 models are mainly from the q^2 -dependence of the form factor. About twteen years ago, Chua [83] studied the decay $B^+ \to \bar{D}_0^{*0} \rho^+$ in the CLFQM approach and obtained its branching fraction about 1.7×10^{-3} , which is consistent with our result by taking $f_{D_0^*} = 0.103$ GeV.

TABLE IX: The CLFQM predictions for the branching ratios (10^{-4}) of the decays $B_s \rightarrow D_{s0}^*\pi(K), D_{s0}^*\rho(K^*)$. The labels LO, VC, HSSC, NLO and the error sources are the same with those given in Table VI. Other theoretical predictions are also listed for comparison.

	$\mathcal{B}r(B_s \to D_{s0}^{*-}\pi^+)$	$\mathcal{B}r(B_s \to D_{s0}^{*-}K^+)$	$\mathcal{B}r(B_s \to D_{s0}^{*-}\rho^+)$	$\mathcal{B}r(B_s \to D_{s0}^{*-}K^{*+})$
LO	$5.06^{+0.83+2.09}_{-0.75-1.83}$	$0.40\substack{+0.07+0.17\\-0.06-0.15}$	$12.87_{-1.91-4.66}^{+2.11+5.31}$	$0.74_{-0.11-0.27}^{+0.12+0.30}$
VC	$5.27_{-0.78-1.91}^{+0.86+2.17}$	$0.42\substack{+0.07+0.17\\-0.06-0.15}$	$13.40^{+2.20+5.53}_{-1.99-4.86}$	$0.77_{-0.11-0.28}^{+0.13+0.32}$
HSSC	$4.96\substack{+0.81+2.05\\-0.74-1.80}$	$0.38\substack{+0.06+0.16\\-0.06-0.14}$	$12.59^{+2.06+5.20}_{-1.87-4.56}$	$0.72_{-0.11-0.26}^{+0.12+0.30}$
NLO	$5.17\substack{+0.85+2.13\\-0.77-1.87}$	$0.40\substack{+0.07+0.16\\-0.06-0.14}$	$13.12\substack{+2.15+5.42\\-1.95-4.76}$	$0.75\substack{+0.12+0.31\\-0.11-0.28}$
PQCD [84]	$5.49_{-1.68-0.27-0.35}^{+2.64+0.41+0.35}$	$0.51\substack{+0.06+0.01+0.01\\-0.04-0.01-0.01}$	$17.7_{-5.3-0.8-1.1}^{+8.5+1.3+1.2}$	$1.01\substack{+0.44+0.06+0.05\\-0.31-0.06-0.07}$
LSCR [67]	$5.2^{+2.5}_{-2.1}$	$0.4^{+0.2}_{-0.2}$	13^{+6}_{-5}	$0.8^{+0.4}_{-0.3}$
RQM [70]	9	0.7	22	1.2
NRQM [85]	10	0.9	27	16
ISGW2 [66]	3.3	—	8.3	—

Taking D_{s0}^* as a $c\bar{s}$ meson, we calclate the branching ratios of the decays $B_{(s)} \to D_{s0}^*M$ with M being a pseudoscalar meson (π, K, D, D_s) or a vector meson (ρ, K^*, D^*, D_s^*) in the CLFQM, which are listed in Tables IX, X and XI. From Table IX, one can find that our predictions for the decays $B_s \to D_{s0}^*\pi(K), D_{s0}^*\rho(K^*)$ are consistent well with those calculated in the LCSR [67] and the PQCD approach [84] within errors, while (much) smaller than those given by the RQM [70] and the nonrelativistic quark model (NRQM) [85]. Especially, for the pure annihilation decay $B_s \to D_{s0}^{*-}K^{*+}$, its branching fraction reaches up to 10^{-3} predicted by the NRQM [85], which seems too large compared to other theoretical results. These divergences can be clarified by the future LHCb and Super KEKB experiments. The branching ratios of the decays $B_s \to D_{s0}^{*-}\pi^+(\rho^+)$ were also calculated in the ISGW2 model [66], which are smaller than our results. It is because of the difference from the form factor of the transition $B_s \to D_{s0}^*$ and its q^2 -dependence. It is similar for the decays $B^0 \to D_0^{*-}\pi^+(\rho^+)$.

TABLE X: The CLFQM predictions for the branching ratios (10^{-4}) of the decays $B^+ \to D_{s0}^{*+} \bar{D}^{(*)0}$ and $B^0 \to D_{s0}^{*+} D^{(*)-}$. The labels LO, VC, HSSC, NLO and the error sources are the same with those given in Table VI. Other theoretical predictions and data are also listed for comparison.

	$\mathcal{B}r(B^+ \to D_{s0}^{*+}\bar{D}^0)$	$\mathcal{B}r(B^+ \to D_{s0}^{*+}\bar{D}^{*0})$	$\mathcal{B}r(B^0\to D_{s0}^{*-}D^+)$	$\mathcal{B}r(B^0\to D_{s0}^{*-}D^{*+})$
LO	$13.81_{-0.21-0.26}^{+0.04+0.19}$	$13.99_{-0.30-0.95}^{+0.12+0.87}$	$12.80^{+0.04+0.17}_{-0.19-0.24}$	$12.97_{-0.28-0.88}^{+0.11+0.80}$
VC	$13.66\substack{+0.04+0.18\\-0.21-0.26}$	$13.83\substack{+0.12+0.86\\-0.30-0.94}$	$12.66\substack{+0.04+0.17\\-0.19-0.24}$	$12.83_{-0.28-0.87}^{+0.11+0.80}$
HSSC	$13.62\substack{+0.04+0.18\\-0.21-0.25}$	$13.95\substack{+0.12+0.87\\-0.30-0.95}$	$12.62\substack{+0.04+0.17\\-0.19-0.24}$	$12.96_{-0.28-0.88}^{+0.11+0.80}$
NLO	$13.47\substack{+0.04+0.18\\-0.20-0.25}$	$13.80\substack{+0.12+0.86\\-0.30-0.94}$	$12.48^{+0.04+0.17}_{-0.19-0.23}$	$12.81\substack{+0.11+0.80\\-0.28-0.87}$
PQCD [86]	$11.2^{+4.0+0.3+0.4}_{-2.8-0.2-0.4}$	$18.3^{+7.1+2.7+0.7}_{-5.4-1.7-0.5}$	$10.5_{-3.0-0.2-0.4}^{+4.5+0.4+0.4}$	$15.9\substack{+7.0+2.4+0.6\\-4.9-1.4-0.5}$
FH [87]	10.3 ± 1.4	5.0 ± 0.7	9.6 ± 1.3	4.7 ± 0.6
TM [88]	6.77 ± 1.9	12.10 ± 3.39	6.37 ± 1.78	8.89 ± 2.49
Data [64]	$8.0^{+1.6}_{-1.3}$	$9.0^{+7.0}_{-7.0}$	$10.6^{+1.6}_{-1.6}$	$15.0^{+6.0}_{-6.0}$

In Table X, all the predictions from the different theories, including the PQCD approach [86], the factorization hypothesis (FH) [87] and the triangle mechanism (TM) [88], show that the branching ratios of the charged decay $B^+ \rightarrow D_{s0}^{*+} \bar{D}^{(*)0}$ are slightly larger than those of the corresponding neutral decays $B^0 \rightarrow D_{s0}^{*+} D^{(*)-}$. It is just contrary to the case of the data [64]. Certainly, there still exist large errors in the experimental results, especially for the branching ratios of the decays with the vector meson D^* involved. We expect more accurate measurements in the future LHCb and Super KEKB experiments. Theoretically, the decays $B^+ \to D_{s0}^{*+} \bar{D}^{(*)0}$ and $B^0 \to D_{s0}^{*+} D^{(*)-}$ have the same CKM matrix elements and Wilson coefficients for the factorizable emission amplitudes, which provide the dominant contributions to their branching ratios. Furthermore, there exist similar transition form factors for isospin symmetry among these channels. So these four decays should have similar branching ratios. From Table XI, one can find that the branching ratios for the decays $B_s \to D_{s0}^* D_s^{(*)}, D_{s0}^* D^{(*)}$ are consistent with those given in the PQCD approach [84] and the RQM [70], while much smaller than those obtained within the LCSR [67]. Further experimental and theoretical researches are needed to clarify these divergences. For the decays $B_s \to D_{s0}^{*-} D^{(*)+}$, their branching ratios are much smaller than those of other four channels mainly because of the smaller CKM matrix element V_{cd} compared with V_{cs} , that is to say there exists a suppressed factor $|V_{cd}/V_{cs}|^2 \approx 0.05$ for the decays $B_s \to D_{s0}^{*-} D^{(*)+}$ compared to other four channels.

TABLE XI: The branching ratios (10^{-3}) of the decays $B_s \to D_{s0}^* D_s^{(*)}, D_{s0}^* D^{(*)}$. The labels LO, VC, HSSC, NLO and the error sources are the same with those given in Table VI. Other theoretical predictions are also listed for comparison.

	LO	VC	HSSC	NLO	PQCD [86]	RQM [70]	LSCR [67]
$\mathcal{B}r(B_s \to D_{s0}^{*-}D_s^+)$	$1.96^{+0.32+0.81}_{-0.29-0.71}$	$1.94\substack{+0.32+0.80\\-0.29-0.70}$	$1.94_{-0.29-0.70}^{+0.32+0.80}$	$1.92^{+0.31+0.79}_{-0.28-0.69}$	$2.1\substack{+0.9+0.3+0.1\\-0.6-0.1-0.1}$	1.1	13^{+7}_{-5}
$\mathcal{B}r(B_s \to D_{s0}^{*-}D_s^{*+})$	$2.18\substack{+0.36+0.90\\-0.32-0.79}$	$2.16\substack{+0.35+0.89\\-0.32-0.78}$	$2.16\substack{+0.35+0.89\\-0.32-0.78}$	$2.14\substack{+0.35+0.88\\-0.32-0.78}$	$1.8\substack{+0.9+0.1+0.1\\-0.6-0.1-0.1}$	2.3	$6.0\substack{+2.9\\-2.4}$
$\mathcal{B}r(B_s\to D_{s0}^{*-}D^+)$	$0.066\substack{+0.011+0.027\\-0.010-0.024}$	$0.065\substack{+0.010+0.027\\-0.011-0.024}$	$0.065\substack{+0.010+0.027\\-0.011-0.024}$	$0.064\substack{+0.011+0.027\\-0.010-0.023}$	$0.065\substack{+0.034+0.006+0.002\\-0.021-0.004-0.002}$	_	$0.5^{+0.2}_{-0.2}$
$\mathcal{B}r(B_s \to D_{s0}^{*-}D^{*+})$	$0.095\substack{+0.016+0.039\\-0.014-0.034}$	$0.093\substack{+0.015+0.038\\-0.014-0.034}$	$0.094\substack{+0.015+0.039\\-0.014-0.034}$	$0.092\substack{+0.015+0.038\\-0.014-0.033}$	$0.050^{+0.026+0.004+0.002}_{-0.017-0.002-0.001}$	_	$0.2^{+0.1}_{-0.1}$
$\mathcal{B}r(B_s \to D_{s0}^{*+}D_s^-)$	$1.29\substack{+0.00+0.06\\-0.01-0.06}$	$1.28\substack{+0.00+0.06\\-0.01-0.06}$	$1.24\substack{+0.00+0.06\\-0.01-0.06}$	$1.22\substack{+0.00+0.06\\-0.01-0.06}$	$1.11\substack{+0.56+0.02+0.04\\-0.37-0.04-0.04}$	_	_
$\mathcal{B}r(B_s \to D_{s0}^{*+}D_s^{*-})$	$1.28\substack{+0.01+0.06\\-0.02-0.11}$	$1.27\substack{+0.01+0.06\\-0.01-0.06}$	$1.26\substack{+0.01+0.06\\-0.02-0.11}$	$1.24\substack{+0.01+0.06\\-0.02-0.11}$	$1.48\substack{+0.69+0.05+0.06\\-0.46-0.07-0.05}$	_	_

In the quark model the axial-vector mesons exist in two types of spectroscopic states, ${}^{3}P_{1}(J^{PC} = 1^{++})$ and ${}^{1}P_{1}(J^{PC} = 1^{+-})$. In some cases the physical particles are the mixture of these two types of states. For example, $K_{1}(1270)$ and $K_{1}(1400)$ are considered as the mixture of K_{1A} and K_{1B} for the mass difference of the strange and light quarks. Similarly, the charm-strange mesons D_{s1} and D'_{s1} are usually written as the mixture of the states ${}^{1}D_{s1}$ and ${}^{3}D_{s1}$, which are given in Eq. (3). The quark potential model determined the mixing angle $\theta_{s} \approx 7^{\circ}$ [66]. So we use $\theta_{s} = 7^{\circ}$ to calculate the branching ratios of the decays $B_{s} \rightarrow D_{s1}^{(i)}P(V)$ with P and V being the pseudoscalar mesons (π, K, D, D_{s}) and the vector mesons $(\rho, K^{*}, D^{*}, D_{s}^{*})$, respectivley, which are listed in Table XII with the results given in ISGW2 model [66, 89] for comparison. The following points can be found

- It is interesting that our predictions for the decays B_s → D'_{s1}π⁺(ρ⁺) are consistent well with the results obtained in the ISGW2 model [66], while those for the channels B_s → D_{s1}π⁺(ρ⁺) are about 5 ~ 6 times smaller than the ISGW2 calculations. Recently, the branching ratios for the decays B_s → D'_{s1}P were updated by using the ISGW2 model in Ref. [89], which strengthen the tension between these two approach predictions. Since many of these decays have large branching ratios, which lie in the range O(10⁻⁵) ~ O(10⁻³), we expect that the LHCb and Super KEKB experiments can clarify the differences between these two models in the future.
- The branching ratios of the CKM-favored decays $B_s \to D_{s1}^{(\prime)} D_s^{(*)}$ and $B_s \to D_{s1}^{(\prime)} \pi(\rho)$, which are associated with the CKM matrix elements V_{cs} and $V_{ud}(\sim 1)$, respectively, are much larger than those of the CKM-suppressed channels $B_s \to D_{s1}^{(\prime)} D^{(*)}$ and $B_s \to D_{s1}^{(\prime)} K^{(*)}$, which are associated with the CKM matrix elements V_{cd} and $V_{us} \approx 0.22$), respectively. It shows a clear hierarchical relationship for the branching raitos of these

color-favored decay modes,

$$\mathcal{B}r(B_s \to D_{s1}^{(')} D_s^{(*)}) \gg \mathcal{B}r(B_s \to D_{s1}^{(')} D^{(*)}), \quad \mathcal{B}r(B_s \to D_{s1}^{(')} \pi(\rho)) \gg \mathcal{B}r(B_s \to D_{s1}^{(')} K^{(*)}).$$
(52)

TABLE XII: The branching ratios (10^{-3}) of the decays $B_s \to D_{s1}^{(\prime)} P(V)$. The labels LO, VC, HSSC, NLO and the error sources are the same with those given in Table VI. The results given in ISGW2 model are also listed for comparison.

	LO	VC	HSSC	NLO	ISGW2 [66]	ISGW2 [89]
$\mathcal{B}r(B_s \to D_{s1}^- D_s^+)$	$0.27\substack{+0.10+0.04+0.29\\-0.08-0.03-0.19}$	$0.26\substack{+0.10+0.04+0.29\\-0.08-0.03-0.19}$	$0.32\substack{+0.11+0.04+0.31\\-0.09-0.03-0.21}$	$0.31^{+0.11+0.04+0.31}_{-0.09-0.03-0.21}$	_	1.2 ± 0.6
$\mathcal{B}r(B_s \to D_{s1}^- D_s^{*+})$	$0.42\substack{+0.15+0.01+0.30\\-0.12-0.00-0.18}$	$0.43\substack{+0.16+0.02+0.30\\-0.13-0.00-0.18}$	$0.46\substack{+0.16+0.01+0.33\\-0.13-0.01-0.21}$	$0.46\substack{+0.16+0.01+0.32\\-0.13-0.01-0.21}$	-	-
$\mathcal{B}r(B_s \to D_{s1}^- \rho^+)$	$0.20\substack{+0.07+0.02+0.18\\-0.06-0.00-0.12}$	$0.21\substack{+0.08+0.06+0.19\\-0.06-0.01-0.13}$	$0.23\substack{+0.08+0.06+0.21\\-0.06-0.04-0.14}$	$0.24\substack{+0.08+0.02+0.21\\-0.07-0.01-0.14}$	1.30	-
$\mathcal{B}r(B_s \to D_{s1}^- \pi^+)$	$0.068\substack{+0.025+0.010+0.074\\-0.020-0.007-0.049}$	$0.071^{+0.026+0.011+0.077}_{-0.021-0.007-0.051}$	$0.081\substack{+0.028+0.011+0.080\\-0.022-0.007-0.055}$	$0.084^{+0.029+0.011+0.083}_{-0.023-0.007-0.057}$	0.52	1.40 ± 0.6
$\mathcal{B}r(B_s \to D_{s1}^- K^+)$	$0.006\substack{+0.002+0.008+0.006\\-0.002-0.005-0.004}$	$0.006\substack{+0.002+0.009+0.006\\-0.002-0.006-0.004}$	$0.007^{+0.002+0.009+0.007}_{-0.002-0.006-0.004}$	$0.007^{+0.002+0.009+0.007}_{-0.002-0.006-0.005}$	-	0.11 ± 0.04
$\mathcal{B}r(B_s\to D_{s1}^-K^{*+})$	$0.012\substack{+0.004+0.009+0.011\\-0.004-0.004-0.007}$	$0.012\substack{+0.005+0.009+0.011\\-0.004-0.004-0.007}$	$0.014\substack{+0.005+0.001+0.012\\-0.004-0.005-0.008}$	$0.014\substack{+0.005+0.001+0.012\\-0.004-0.005-0.008}$		-
	$0.009\substack{+0.003+0.001+0.010\\-0.003-0.009-0.006}$	$0.009\substack{+0.003+0.001+0.009\\-0.003-0.009-0.006}$	$0.011\substack{+0.004+0.002+0.011\\-0.003-0.001-0.008}$	$0.011\substack{+0.004+0.001+0.011\\-0.003-0.009-0.007}$	-	0.055 ± 0.025
$\mathcal{B}r(B_s \to D_{s1}^- D^{*+})$	$0.018\substack{+0.007+0.001+0.013\\-0.005-0.000-0.008}$	$0.019\substack{+0.007+0.001+0.013\\-0.005-0.000-0.008}$	$0.021\substack{+0.007+0.001+0.015\\-0.006-0.000-0.010}$	$0.023\substack{+0.007+0.001+0.015\\-0.006-0.000-0.010}$	_	-
$\mathcal{B}r(B_s \to D_{s1}^{\prime -} D_s^+)$	$6.55\substack{+0.56+0.12+1.36\\-0.55-0.18-1.29}$	$6.48\substack{+0.56+0.12+1.34\\-0.55-0.17-1.28}$	$6.44\substack{+0.56+0.12+1.36\\-0.55-0.17-1.28}$	$6.38\substack{+0.56+0.12+1.34\\-0.54-0.17-1.27}$	-	10.0 ± 4
$\mathcal{B}r(B_s \to D_{s1}^{\prime -} D_s^{*+})$	$8.45\substack{+0.72+0.24+1.59\\-0.70-0.32-1.53}$	$8.48\substack{+0.72+0.25+1.58\\-0.70-0.33-1.52}$	$8.29\substack{+0.71+0.23+1.59\\-0.69-0.31-1.52}$	$8.32\substack{+0.71+0.24+1.58\\-0.70-0.32-1.52}$	-	-
$\mathcal{B}r(B_s \to D_{s1}^{\prime -} \rho^+)$	$4.49\substack{+0.39+0.10+0.90\\-0.38-0.14-0.86}$	$4.68\substack{+0.40+0.11+0.94\\-0.39-0.15-0.90}$	$4.40_{-0.38-0.13-0.86}^{+0.38+0.10+0.90}$	$4.59_{+0.39+0.14+0.89}^{+0.40+0.10+0.94}$	3.8	-
$\mathcal{B}r(B_s \to D_{s1}^{\prime -}\pi^+)$	$1.67\substack{+0.14+0.03+0.35\\-0.14-0.04-0.33}$	$1.74\substack{+0.15+0.03+0.36\\-0.15-0.05-0.34}$	$1.64\substack{+0.14+0.03+0.34\\-0.14-0.04-0.33}$	$1.70\substack{+0.15+0.03+0.36\\-0.15-0.01-0.34}$	1.5	7.2 ± 2.7
$\mathcal{B}r(B_s \to D_{s1}^{\prime-}K^+)$	$0.13\substack{+0.01+0.00+0.03\\-0.01-0.00-0.03}$	$0.14\substack{+0.01+0.00+0.03\\-0.01-0.00-0.03}$	$0.13\substack{+0.01+0.00+0.03\\-0.01-0.00-0.03}$	$0.14\substack{+0.01+0.00+0.03\\-0.01-0.00-0.03}$	-	0.52 ± 0.2
$\mathcal{B}r(B_s \to D_{s1}^{'-}K^{*+})$	$0.26\substack{+0.02+0.01+0.05\\-0.02-0.01-0.05}$	$0.27\substack{+0.02+0.01+0.05\\-0.02-0.01-0.05}$	$0.26\substack{+0.02+0.01+0.05\\-0.02-0.01-0.05}$	$0.27\substack{+0.02+0.01+0.05\\-0.02-0.01-0.05}$	-	-
$\mathcal{B}r(B_s \to D_{s1}^{\prime -}D^+)$	$0.22\substack{+0.02+0.00+0.05\\-0.02-0.01-0.04}$	$0.22\substack{+0.02+0.00+0.05\\-0.02-0.01-0.04}$	$0.22\substack{+0.02+0.00+0.05\\-0.02-0.01-0.04}$	$0.21\substack{+0.02+0.00+0.05\\-0.02-0.01-0.04}$	-	0.44 ± 0.16
$\mathcal{B}r(B_s\to D_{s1}^{'-}D^{*+})$	$0.37\substack{+0.03+0.01+0.07\\-0.03-0.01-0.07}$	$0.37\substack{+0.03+0.01+0.07\\-0.03-0.01-0.07}$	$0.36\substack{+0.03+0.01+0.07\\-0.03-0.01-0.07}$	$0.36\substack{+0.03+0.01+0.07\\-0.03-0.01-0.07}$	-	-

- Our predictions for the branching ratios of the decays $B_s \to D'_{s1}P(V)$ are at least one order larger than those of the corresponding decays $B_s \to D_{s1}P(V)$. This is because that the related form factor $V_0^{B_s D'_{s1}}$ is much larger than that of $V_0^{B_s D_{s1}}$. While it is just contrary for the decays $B \to D^{(*)}D_{s1}^{(\prime)}$, where $D_{s1}^{(\prime)}$ is at the emission position in the Feynman diagrams.
- In view of the mixing angle θ_s uncertainty, we check the dependence of the branching ratios of the decays $B_s \to D_{s1}^{(\prime)}\pi(K)$ on the mixing angle θ_s , which are shown in Figure 4. One can find that the branching ratios of the decays $B_s \to D_{s1}\pi(K)$ are very sensitive to the mixing angle, while those of the decays $B_s \to D_{s1}^{\prime}\pi(K)$ show an insensitive dependence on θ_s . Furthermore, the changing trends for the branching ratios of these two kinds of decays are just opposite.




FIG. 4: The dependence of the branching ratios of the decays $B_s \to D_{s1}^{(\prime)}\pi(K)$ on the mixing angle θ_s .

TABLE XIII: The branching ratios (10^{-3}) of the decays $B \to D^{(*)}D_{s1}^{(\prime)}$. The labels LO, VC, HSSC, NLO and the error sources are the same with those given in Table VI. Other theoretical predictions and data are also listed for comparison.

	LO	VC	HSSC	NLO	ISGW2 [66]	ISGW [91]	FH [87]	TM [88]	Data [64, 90]
$\mathcal{B}r(B^0\to D^-D^+_{s1})$	$6.41\substack{+0.02+0.09\\-0.10-0.12}$	$7.08\substack{+0.02+0.09\\-0.11-0.13}$	$6.33\substack{+0.02+0.08\\-0.10-0.12}$	$6.99\substack{+0.02+0.09\\-0.11-0.13}$	3.9	3.4	2.36 ± 0.36	1.158 ± 0.324	3.5 ± 1.1
$\mathcal{B}r(B^0\to D^{*-}D^+_{s1})$	$9.78\substack{+0.08+0.52\\-0.20-0.58}$	$10.52\substack{+0.09+0.57\\-0.22-0.63}$	$9.70\substack{+0.08+0.52\\-0.20-0.58}$	$10.44\substack{+0.08+0.56\\-0.22-0.63}$	15	—	6.85 ± 1.05	2.709 ± 0.759	9.3 ± 2.2
$\mathcal{B}r(B^+\to \bar{D}^0 D^+_{s1})$	$6.92\substack{+0.02+0.09\\-0.10-0.13}$	$7.64\substack{+0.02+0.10\\-0.12-0.14}$	$6.83\substack{+0.02+0.09\\-0.10-0.13}$	$7.54\substack{+0.02+0.10\\-0.11-0.14}$	4.3	3.5	2.54 ± 0.39	1.255 ± 0.351	$3.1^{+1.0}_{-0.9}$
$\mathcal{B}r(B^+\to \bar{D}^{*0}D^+_{s1})$	$10.55\substack{+0.09+0.56\\-0.22-0.63}$	$11.35\substack{+0.09+0.61\\-0.24-0.68}$	$10.46\substack{+0.08+0.56\\-0.22-0.62}$	$11.26\substack{+0.09+0.61\\-0.23-0.67}$	16	-	7.33 ± 1.12	3.065 ± 0.858	12.0 ± 3.0
$\mathcal{B}r(B^0\to D^-D_{s1}^{'+})$	$0.29\substack{+0.00+0.00\\-0.00-0.01}$	$0.32\substack{+0.00+0.00\\-0.00-0.01}$	$0.29\substack{+0.00+0.00\\-0.00-0.01}$	$0.32\substack{+0.00+0.00\\-0.00-0.01}$	0.28	3.3	-	-	0.39 ± 0.18^a
$\mathcal{B}r(B^0\to D^{*-}D_{s1}^{'+})$		$0.48\substack{+0.01+0.03\\-0.00-0.03}$	$0.44\substack{+0.00+0.02\\-0.01-0.03}$	$0.48\substack{+0.00+0.03\\-0.01-0.03}$	1.1	-	-	-	0.71 ± 0.28^{b}
$\mathcal{B}r(B^+\to \bar{D}^0D_{s1}^{'+})$		$0.35\substack{+0.01+0.00\\-0.00-0.01}$	$0.31\substack{+0.00+0.00\\-0.00-0.01}$	$0.34\substack{+0.00+0.00\\-0.00-0.01}$	0.31	3.4	-	-	0.35 ± 0.16^a
$\mathcal{B}r(B^+\to \bar{D}^{*0}D_{s1}^{'+})$	$0.48\substack{+0.00+0.03\\-0.01-0.03}$	$0.52\substack{+0.00+0.03\\-0.01-0.03}$	$0.48\substack{+0.00+0.03\\-0.01-0.03}$	$0.51\substack{+0.00+0.03\\-0.01-0.03}$	1.2	-	-	-	0.91 ± 0.36^{b}

^a It is obtained from the ratios $\frac{\mathcal{B}r(B \to DD_{s1})}{\mathcal{B}r(B \to DD_{s}^{*})} = 0.44 \pm 0.11$ and $\frac{\mathcal{B}r(B \to DD_{s1}^{*})}{\mathcal{B}r(B \to DD_{s}^{*})} = 0.049 \pm 0.010$ [90].

^b It is obtained from the ratios $\frac{\mathcal{B}_{T}(B \to D^{*}D_{s1})}{\mathcal{B}_{T}(B \to D^{*}D_{s}^{*})} = 0.58 \pm 0.12 \text{ and } \frac{\mathcal{B}_{T}(B \to D^{*}D_{s1})}{\mathcal{B}_{T}(B \to D^{*}D_{s}^{*})} = 0.044 \pm 0.010 \ [90].$

In Table XIII, we present our predictions for the branching ratios of the decays $B \to D^{(*)}D_{s1}^{(\prime)}$, which are associated with the $B \to D^{(*)}$ transition, accompanied by the $D_{s1}^{(\prime)}$ emission. When the emission meson is D_{s1} , our results for the decays $B^0 \to D^- D_{s1}^+$ and $B^+ \to \bar{D}^0 D_{s1}^+$ are larger than those given by the ISGW(2) [66, 91] and the FH [87] by a factor of 2 to 3. While for other two decays $B^0 \to D^{*-} D_{s1}^+$ and $B^+ \to \bar{D}^{*0} D_{s1}^+$, our predictions are consistent well with the theoretical and experimental results within errors except for those given in Ref. [88], where the triangle mechanism was used by considering D_{s1} as a molecular state. When the emission meson is D'_{s1} , the branching ratios of the decays $B^0 \to D^{(*)-} D_{s1}^{'+}$ and $B^+ \to \bar{D}^{(*)0} D_{s1}^{'+}$ are at least one order smaller than those of the decays $B^0 \to D^{(*)-} D_{s1}^{'+}$ and $B^+ \to \bar{D}^{(*)0} D_{s1}^{++}$. It is because of the much smaller decay constant $f_{D'_{s1}}$ compared to $f_{D_{s1}}$. Such character has been verified by the data shown in Table XIII.

C. SUMMARY

Firstly, we studied the form factors of the transitions $B_{(s)} \to D_0^*, D_{s0}^*, D_{s1}$ and D'_{s1} in the covariant light-front quark model (CLFQM). One can find that these form factors are (much) smaller than those of the transitions $B_{(s)} \to D_{(s)}, D^*_{(s)}$. Certainly, because of the mixing between D_{s1} and D'_{s1} , the determination of the form factors for the transitions $B_s \to D_{s1}, D'_{s1}$ are more difficult. Secondly, using the amplitudes combined via the form factors, we calculated the branching ratios of the $B_{(s)}$ nonleptonic decays with these four charmed mesons involved. Furthermore, the QCD radiative corrections to the hadronic matrix elements within the framework of QCD factorization were included. From the numerical results, we found the following points

1. The small form factors of the transitions $B_{(s)} \to D_0^*, D_{s0}^*$ are related to the small decay constants $f_{D_0^*}, f_{D_{s0}^*}$. Unfortunately, there exits large uncertainties in these two decay constants. Combined with the data, our predictions for the branching ratios of the $B_{(s)}$ decays with D_0^*, D_{s0}^* involved are helpful to probe the inner structures of these two states. Most of the decays $B_{(s)} \to D_0^* P(V), D_{s0}^* P(V)$ with P(V) being a light pseudoscalar (vector) meson or a charmed meson are not sensitive to the QCD radiative corrections including the vertex corrections and the hard spectator-scattering, except for the decays $B^+ \to \bar{D}_0^{*0}\pi^+(K^+)$, where two kinds of Feynman diagrams contribute to the branching ratios. Our predictions for the branching ratios of the quasi-twobody decays $B^+ \to \bar{D}_0^{*0}\pi^+ \to D^-\pi^+\pi^+$ and $B^0 \to D_0^{*-}K^+ \to \bar{D}^0\pi^-K^+$ can explain the data by taking appropriate decay constant value for D_0^* , while for the decays $B^+ \to \bar{D}_0^{*0}K^+ \to D^-\pi^+K^+$ and $B^0 \to D_0^{*-}\pi^+$, their branching ratios are much larger than the LHCb measurements. There exist the similar cases for the PQCD calculations compared with the data.

- 2. We checked the dependence of the branching ratios of the decays $B_s \rightarrow D_{s1}P(V), D'_{s1}P(V)$ on the mixing angle θ_s and found that the branching ratios of the decays $B_s \rightarrow D_{s1}P(V)$ are very sensitive to the mixing angle, while those of the decays $B_s \rightarrow D'_{s1}P(V)$ show an insensitive dependence on θ_s . The changing trends of the branching ratios between these two kinds of decays are just opposite. Furthermore, the branching ratios of the decays $B_s \rightarrow D'_{s1}P(V)$ are are at least one order smaller than those of the decays $B_s \rightarrow D'_{s1}P(V)$. It is because of the smaller form factor $V_0^{B_s D_{s1}}$ compared to $V_0^{B_s D'_{s1}}$. While it is just constrary for the decays $B \rightarrow D^{(*)}D^{(\prime)}_{s1}$, where the $D^{(\prime)}_{s1}$ is at the emission position in the Feynman diagrams, that is to say the branching ratios of the decays $B \rightarrow D^{(*)}D_{s1}$ are at least one order larger than those of the decays $B \rightarrow D^{(*)}D_{s1}$. It is because of the the larger decay constant $f_{D_{s1}}$ compared to $f_{D'_{s1}}$. This character has been verified by the experimental measurements.
- 3. Our predictions are helpful to clarify the different assumptions about the inner structures of these four charmed hadrons by comparing with the future data.

Acknowledgment

This work is partly supported by the National Natural Science Foundation of China under Grant No. 11347030, by the Program of Science and Technology Innovation Talents in Universities of Henan Province 14HASTIT037.

Appendix A: Some specific rules under the p^- intergration

When preforming the integraion, we need to include the zero-mode contributions. It amounts to performing the integration in a proper way in the CLFQM. Specificly we use the following rules given in Refs. [48, 51]

$$\hat{p}_{1\mu}' \doteq P_{\mu} A_1^{(1)} + q_{\mu} A_2^{(1)}, \tag{A1}$$

$$\hat{p}_{1\mu}'\hat{p}_{1\nu}' \doteq g_{\mu\nu}A_1^{(2)} + P_{\mu}P_{\nu}A_2^{(2)} + (P_{\mu}q_{\nu} + q_{\mu}P_{\nu})A_3^{(2)} + q_{\mu}q_{\nu}A_4^{(2)}, \tag{A2}$$

$$Z_2 = \hat{N}'_1 + m'^2_1 - m^2_2 + (1 - 2x_1) M'^2 + (q^2 + q \cdot P) \frac{p'_{\perp} \cdot q_{\perp}}{q^2},$$
(A3)

$$A_1^{(1)} = \frac{x_1}{2}, \quad A_2^{(1)} = A_1^{(1)} - \frac{p'_{\perp} \cdot q_{\perp}}{q^2}, \quad A_3^{(2)} = A_1^{(1)} A_2^{(1)},$$
 (A4)

$$A_4^{(2)} = \left(A_2^{(1)}\right)^2 - \frac{1}{q^2}A_1^{(2)}, \quad A_1^{(2)} = -p_{\perp}'^2 - \frac{\left(p_{\perp}' \cdot q_{\perp}\right)^2}{q^2}, \quad A_2^{(2)} = \left(A_1^{(1)}\right)^2.$$
(A5)

Appendix B: EXPRESSIONS OF $B \rightarrow D_0^*$, ${}^iD_{s1}$ FORM FACTORS

$$F_{1}^{BD_{0}^{*}}(q^{2}) = \frac{N_{c}}{16\pi^{3}} \int dx_{2}d^{2}p'_{\perp} \frac{h'_{B}h''_{D_{0}^{*}}}{x_{2}\hat{N}'_{1}\hat{N}''_{1}} \left[x_{1} \left(M'_{0}^{2} + M''_{0}^{2} \right) + x_{2}q^{2} \\ -x_{2} \left(m'_{1} + m''_{1} \right)^{2} - x_{1} \left(m'_{1} - m_{2} \right)^{2} - x_{1} \left(m''_{1} + m_{2} \right)^{2} \right], \qquad (B1)$$

$$F_{0}^{BD_{0}^{*}}(q^{2}) = F_{1}^{BD_{0}^{*}}(q^{2}) + \frac{q^{2}}{q \cdot P} \frac{N_{c}}{16\pi^{3}} \int dx_{2}d^{2}p'_{\perp} \frac{2h'_{B}h''_{D_{0}^{*}}}{x_{2}\hat{N}'_{1}\hat{N}''_{1}} \left\{ -x_{1}x_{2}M'^{2} - p'_{\perp}^{2} - m'_{1}m_{2} \\ - \left(m''_{1} + m_{2} \right) \left(x_{2}m'_{1} + x_{1}m_{2} \right) + 2\frac{q \cdot P}{q^{2}} \left(p'_{\perp}^{2} + 2\frac{\left(p'_{\perp} \cdot q_{\perp} \right)^{2}}{q^{2}} \right) + 2\frac{\left(p'_{\perp} \cdot q_{\perp} \right)^{2}}{q^{2}} \\ - \frac{p'_{\perp} \cdot q_{\perp}}{q^{2}} \left[M''^{2} - x_{2} \left(q^{2} + q \cdot P \right) - \left(x_{2} - x_{1} \right) M'^{2} + 2x_{1}M'^{2}_{0} \\ - 2 \left(m'_{1} - m_{2} \right) \left(m'_{1} - m''_{1} \right) \right] \right\}, \qquad (B2)$$

$$A^{B^{i}D_{s1}}(q^{2}) = \left(M' - M'' \right) \frac{N_{c}}{16\pi^{3}} \int dx_{2}d^{2}p'_{\perp} \frac{2h'_{B}h''_{D_{s1}}}{x_{2}\hat{N}'_{1}\hat{N}''_{1}} \left\{ x_{2}m'_{1} + x_{1}m_{2} + \left(m'_{1} + m''_{1} \right) \frac{p'_{\perp} \cdot q_{\perp}}{q^{2}} \\ + \frac{2}{w''_{r_{p}}} \left[p'_{\perp}^{2} + \frac{\left(p'_{\perp} \cdot q_{\perp} \right)^{2}}{q^{2}} \right] \right\}, \qquad (B3)$$

(B3)

$$W_{iD_{s1}}\left[\begin{array}{c} q^{2} \end{array}\right] = \frac{1}{M' - M''} \frac{N_{c}}{16\pi^{3}} \int dx_{2}d^{2}p'_{\perp} \frac{h'_{B}h''_{D_{s1}}}{x_{2}\hat{N}'_{1}\hat{N}''_{1}} \{2x_{1}\left(m_{2} - m'_{1}\right)\left(M'_{0}^{2} + M''^{2}\right) + 4x_{1}m''_{1}M'_{0}^{2} + 2x_{2}m'_{1}q \cdot P + 2m_{2}q^{2} - 2x_{1}m_{2}\left(M'^{2} + M''^{2}\right) + 2\left(m'_{1} - m_{2}\right)\left(m'_{1} - m''_{1}\right)^{2} + 8\left(m'_{1} - m_{2}\right)\right) \\ \times \left[p'_{\perp}^{2} + \frac{\left(p'_{\perp} \cdot q_{\perp}\right)^{2}}{q^{2}}\right] + 2\left(m'_{1} - m''_{1}\right)\left(q^{2} + q \cdot P\right)\frac{p'_{\perp} \cdot q_{\perp}}{q^{2}} - 4\frac{q^{2}p'_{\perp}^{2} + \left(p'_{\perp} \cdot q_{\perp}\right)^{2}}{q^{2}w''_{D_{s1}}} \\ \times \left[2x_{1}\left(M'^{2} + M'^{2}_{0}\right) - q^{2} - q \cdot P - 2\left(q^{2} + q \cdot P\right)\frac{p'_{\perp} \cdot q_{\perp}}{q^{2}} - 2\left(m'_{1} + m''_{1}\right)\left(m'_{1} - m_{2}\right)\right]\right\}, (B4)$$

$$V_{2}^{B\ iD_{s1}}(q^{2}) = \left(M' - M''\right)\frac{N_{c}}{16\pi^{3}}\int dx_{2}d^{2}p'_{\perp}\frac{2h'_{B}h''_{D_{s1}}}{x_{2}\hat{N}'_{1}\hat{N}''_{1}}\left\{(x_{1} - x_{2})\left(x_{2}m'_{1} + x_{1}m_{2}\right) - \left[2x_{1}m_{2} - m''_{1}\right] \\ + \left(x_{2} - x_{1}\right)m'_{1}\right] \times \frac{p'_{\perp} \cdot q_{\perp}}{q^{2}} - 2\frac{x_{2}q^{2} + p'_{\perp} \cdot q_{\perp}}{x_{2}q^{2}w''_{D_{s1}}}\left[p'_{\perp} \cdot p''_{\perp} + \left(x_{1}m_{2} + x_{2}m'_{1}\right)\right] \\ \times \left(x_{1}m_{2} + x_{2}m''_{1}\right)\right], \tag{B5}$$

$$V_{0}^{B^{i}D_{s1}}(q^{2}) = \frac{M' - M''}{2M''} V_{1}^{B^{i}D_{s1}}(q^{2}) - \frac{M' + M''}{2M''} V_{2}^{B^{i}D_{s1}}(q^{2}) - \frac{q^{2}}{2M''} \frac{N_{c}}{16\pi^{3}} \int dx_{2}d^{2}p'_{\perp} \frac{h'_{B}h''_{i}D_{s1}}{x_{2}\hat{N}'_{1}\hat{N}''_{1}} \\ \times \{2(2x_{1} - 3)(x_{2}m'_{1} + x_{1}m_{2}) - 8(m'_{1} - m_{2})\left[\frac{p'_{\perp}^{2}}{q^{2}} + 2\frac{(p'_{\perp} \cdot q_{\perp})^{2}}{q^{4}}\right] - [(14 - 12x_{1})m'_{1} \\ + 2m''_{1} - (8 - 12x_{1})m_{2}]\frac{p'_{\perp} \cdot q_{\perp}}{q^{2}} + \frac{4}{w'_{i}D_{s1}}([M'^{2} + M''^{2} - q^{2} + 2(m'_{1} - m_{2})(-m''_{1} + m_{2})] \\ \times \left(A_{3}^{(2)} + A_{4}^{(2)} - A_{2}^{(1)}\right) + Z_{2}\left(3A_{2}^{(1)} - 2A_{4}^{(2)} - 1\right) + \frac{1}{2}[x_{1}(q^{2} + q \cdot P) - 2M'^{2} - 2p'_{\perp} \cdot q_{\perp} \\ -2m'_{1}(-m''_{1} + m_{2}) - 2m_{2}(m'_{1} - m_{2})]\left(A_{1}^{(1)} + A_{2}^{(1)} - 1\right) \\ \times q \cdot P\left[\frac{p'_{\perp}^{2}}{q^{2}} + \frac{(p'_{\perp} \cdot q_{\perp})^{2}}{q^{4}}\right]\left(4A_{2}^{(1)} - 3\right)\right)\right\},$$
(B6)

with i = 1, 3.

- [1] B. Aubert et al. [BaBar], Phys. Rev. Lett. 90, 242001 (2003) [arXiv:hep-ex/0304021].
- [2] D. Besson *et al.* [CLEO], Phys. Rev. D 68, 032002 (2003) [Erratum: Phys. Rev. D 75, 119908 (2007)] [arXiv:hep-ex/0305100].
- [3] K. Abe, et al. [Belle], Phys. Rev. D 69 112002 (2004) [arXiv:hep-ex/0307021].
- [4] A. E. Asratian, *et al.*, Z. Phys. C **40** 483 (1988).
- [5] S. Godfrey and N. Isgur, Phys. Rev. D 32, 189 (1985).
- [6] Z. X. Xie, G. Q. Feng and X. H. Guo, Phys. Rev. D 81, 036014 (2010).
- [7] M. Cleven, H. W. Griehammer, F. K. Guo, C. Hanhart and U. G. Meiner, Eur. Phys. J. A 50, 149 (2014) [arXiv:1405.2242 [hep-ph]].
- [8] F. K. Guo, P. N. Shen, H. C. Chiang, R. G. Ping, and B. S. Zou, Phys. Lett. B 641, 278 (2006) [arXiv:hep-ph/0603072].
- [9] T. Barnes, F. E. Close and H. J. Lipkin, Phys. Rev. D 68, 054006 (2003) [arXiv:hep-ph/0305025].
- [10] E. E. Kolomeitsev and M. F. M. Lutz, Phys. Lett. B 582, 39 (2004) [arXiv:hep-ph/0307133].
- [11] J. Hofmann and M. F. M. Lutz, Nucl. Phys. A 733, 142 (2004) [arXiv:hep-ph/0308263].
- [12] C. J. Xiao, D. Y. Chen and Y. L. Ma, Phys. Rev. D 93, 094011 (2016) [arXiv:1601.06399 [hep-ph]].
- [13] L. Maiani, F. Piccinini, A. D. Polosa, and V. Riquer, Phys. Rev. D 71, 014028 (2005)
 [arXiv:hep-ph/0412098].
- [14] Z. G. Wang and S. L. Wan, Nucl. Phys. A 778, 22 (2006) [arXiv:hep-ph/0602080].
- [15] H. Y. Cheng and W. S. Hou, Phys. Lett. B 566, 193 (2003) [arXiv:hep-ph/0305038].
- [16] Y. Q. Chen and X. Q. Li, Phys. Rev. Lett. 93, 232001 (2004) [arXiv:hep-ph/0407062].
- [17] H. Kim and Y. Oh, Phys. Rev. D 72, 074012 (2005) [arXiv:hep-ph/0508251].
- [18] W. A. Bardeen, E. J. Eichten, and C. T. Hill, Phys. Rev. D 68, 054024 (2003) [arXiv:hep-ph/0305049].
- [19] M. A. Nowak, M. Rho, and I. Zahed, Acta Phys. Polon. B 35, 2377 (2004) [arXiv:hep-ph/0307102].
- [20] T. E. Browder, S. Pakvasa, and A. A. Petrov, Phys. Lett. B 578, 365 (2004) [arXiv:hep-ph/0307054].
- [21] J. Vijande, F. Fernandez, and A. Valcarce, Phys. Rev. D 73, 034002 (2006) [arXiv:hep-ph/0601143].
- [22] M. E. Bracco, A. Lozea, R. D. Matheus, F. S. Navarra and M. Nielsen, Phys. Lett. B 624,

217 (2005) [arXiv:hep-ph/0503137].

- [23] M. F. M. Lutz and M. Soyeur, Prog. Part. Nucl. Phys. 61, 155 (2008).
- [24] D. S. Hwang and D. W. Kim, Phys. Lett. B 601, 137 (2004) [arXiv:hep-ph/0408154].
- [25] X. Liu, Y. M. Yu, S. M. Zhao and X. Q. Li, Eur. Phys. J. C 47, 445 (2006) [arXiv:hep-ph/0601017].
- [26] J. Lu, X. L. Chen, W. Z. Deng and S. L. Zhu, Phys. Rev. D 73, 054012 (2006) [arXiv:hep-ph/0602167].
- [27] J. B. Liu and M. Z. Yang, JHEP 1407, 106 (2014) [arXiv:1307.4636 [hep-ph]].
- [28] Z. G. Wang, Phys. Rev. D 75, 034013 (2007) [arXiv:hep-ph/0612225].
- [29] S. Fajfer and A. P. Brdnik, Phys. Rev. D 92, 074047 (2015) [arXiv:1506.02716 [hep-ph]].
- [30] Q. T. Song, D. Y. Chen, X. Liu and T. Matsuki, Phys. Rev. D 91, 054031 (2015) [arXiv:1501.03575 [hep-ph]].
- [31] S. F. Chen, J. Liu, H. Q. Zhou, and D. Y. Chen, Eur. Phys. J. C 80, 290 (2020) [arXiv:2003.07988 [hep-ph]].
- [32] B. Aubert, et al. [BaBar], Phys. Rev. D 79 112004 (2009). [arXiv:0901.1291 [hep-ex]].
- [33] S. Godfrey and R. Kokoski, Phys. Rev. D 43 1679 (1991).
- [34] M. Nielsen, R. D. Matheus, F. S. Navarra and M. E. Bracco, Nucl. Phys. B 161 193 (2006).
- [35] J. M. Link *et al.* [FOCUS], Phys. Lett. B **586**, 11 (2004) [arXiv:hep-ex/0312060].
- [36] M. Albaladejo, P. Ferniäandez-Soler, F. K. Guo, and J. Nieves, Phys. Lett. B 767, 465 (2017) [arXiv:1610.06727 [hep-ph]].
- [37] M. L. Du, M. Albaladejo, P. Ferniäandez-Soler, F. K. Guo, C. Hanhart, U. G. Meiner, J. Nieves, and D. L. Yao, Phys. Rev. D 98, 094018 (2018) [arXiv:1712.07957 [hep-ph]].
- [38] F. K. Guo, C. Hanhart, U. G. Meiner, Q. Wang, Q. Zhao, and B. S. Zou, Rev. Mod. Phys. 90, 015004 (2018) [arXiv:1705.00141 [hep-ph]].
- [39] M. L. Du, F. K. Guo, C. Hanhart, B. Kubis and U. G. Meiner, Phys. Rev. Lett. **126** (2021), 192001 [arXiv:2012.04599 [hep-ph]].
- [40] D. Gamermann, E. Oset, D. Strottman and M. J. Vicente Vacas, Phys. Rev. D 76 (2007) 074016 [arXiv:hep-ph/0612179]
- [41] B. Aubert, et al. [BaBar], Phys. Rev. D 77 011102 (2008) [arXiv:0708.1565 [hep-ex]].
- [42] T. Aushev, et al. [Belle], Phys. Rev. D 83 051102 (2011) [arXiv:1102.0935 [hep-ex]].
- [43] R. Aaij, et al. [LHCb], Phys. Rev. D 86 112005 (2012) [arXiv:1211.1541 [hep-ex]].
- [44] V. Balagura, et al. [Belle], Phys. Rev. D 77 032001 (2008) [arXiv:0709.4184 [hep-ex]].
- [45] Z. H. Wang, Y. Zhang, T. h. Wang, Y. Jiang, Q. Li and G. L. Wang, Chin. Phys. C 42, 123101 (2018) [arXiv:1803.06822 [hep-ph]].
- [46] E. E. Salpeter and H. A. Bethe, Phys. Rev. 84 1232 (1951).

- [47] E. E. Salpeter, Phys. Rev. 87 328 (1952).
- [48] W. Jaus, Phys. Rev. D **60** 054026 (1999).
- [49] H. M. Choi and C. R. Ji, Phys. Rev. D 58 071901 (1998) [arXiv:hep-ph/9805438].
- [50] H. Y. Cheng, C. Y. Cheung, C. W. Hwang and W. M. Zhang, Phys. Rev. D 57 5598 (1998) [arXiv:hep-ph/9709412].
- [51] H. Y. Cheng, C. K. Chua and C. W. Hwang, Phys. Rev. D 69, 074025 (2004) [arXiv:hep-ph/0310359].
- [52] W. Wang, Y. L. Shen and C. D. Lu, Phys. Rev. D 79, 054012 (2009) [arXiv:0811.3748 [hep-ph]].
- [53] X. X. Wang, W. Wang and C. D. Lu, Phys. Rev. D 79, 114018 (2009) [arXiv:0901.1934 [hep-ph]].
- [54] W. Wang, Y. L. Shen and C. D. Lu, Eur. Phys. J. C 51, 841 (2007) [arXiv:0704.2493 [hep-ph]].
- [55] H. W. Ke, T. Liu and X. Q. Li, Phys. Rev. D 89, 017501 (2014) [arXiv:1307.5925 [hep-ph]].
- [56] G. Li, F. l. Shao and W. Wang, Phys. Rev. D 82, 094031 (2010) [arXiv:1008.3696 [hep-ph]].
- [57] Z. Q. Zhang, Z. J. Sun, Y. C. Zhao, Y. Y. Yang and Z. Y. Zhang, Eur. Phys. J. C 83, 477 (2023) [arXiv:2301.11107 [hep-ph]].
- [58] Z. J. Sun, S. Y. Wang, Z. Q. Zhang, Y. Y. Yang and Z. Y. Zhang, Eur. Phys. J. C 83, 945 (2023) [arXiv:2308.03114 [hep-ph]].
- [59] Z. J. Sun, Z. Q. Zhang, Y. Y. Yang and H. Yang, Eur. Phys. J. C 84, 65 (2024) [arXiv:2311.04431 [hep-ph]].
- [60] M. Wirbel, B. Stech and M. Bauer, Z. Phys. C 29, 637 (1985).
- [61] M. Beneke, G. Buchalla, M. Neubert, and C. T. Sachrajda, Phys. Rev. Lett. 83, 1914 (1999) [arXiv:hep-ph/9905312].
- [62] H. Y. Cheng, C. K. Chua and K. C. Yang, Phys. Rev. D 77, 014034 (2008) [arXiv:0705.3079 [hep-ph]].
- [63] M. Beneke and M. Neubert, Nucl. Phys. B 675, 333 (2003) [arXiv:hep-ph/0308039].
- [64] R. L. Workman *et al.* [Particle Data Group], Review of Particle Physics. PTEP **2022**, 083C01 (2022).
- [65] D. Becirevic, P. Boucaud, J. P. Leroy, V. Lubicz, G. Martinelli, F. Mescia and F. Rapuano, Phys. Rev. D 60, 074501 (1999) [arXiv:hep-lat/9811003].
- [66] H. Y. Cheng, Phys. Rev. D 68, 094005 (2003) [arXiv:hep-ph/0307168].
- [67] R. H. Li and C. D. Lu, Phys. Rev. D 80, 014005 (2009) [arXiv:0905.3259 [hep-ph]].
- [68] T. M. Aliev and M. Savci, Phys. Rev. D 73, 114010 (2006) [arXiv:hep-ph/0604002].
- [69] D. Melikhov and B. Stech, Phys. Rev. D 62, 014006 (2000) [arXiv:hep-ph/0001113].
- [70] R. N. Faustov and V. O. Galkin, Phys. Rev. D 87.034033 [arXiv:1212.3167 [hep-ph]].

- [71] G. Kramer and W. F. Palmer, Phys. Rev. D 46, 3197 (1992).
- [72] X. J. Chen, H. F. Fu, C. S. Kim and G. L. Wang, J. Phys. G 39, 045002 (2012) [arXiv:1106.3003 [hep-ph]].
- [73] P. Blasi, P. Colangelo, G. Nardulli and N. Paver, Phys. Rev. D 49, 238 (1994) [arXiv:hep-ph/9307290].
- [74] P. Ball, JHEP 09, 005 (1998) [arXiv:hep-ph/9802394].
- [75] M. Beneke, G. Buchalla, M. Neubert and C. T. Sachrajda, Nucl. Phys. B 606, 245 (2001)
 [arXiv:hep-ph/0104110].
- [76] W. F. Wang, Phys. Lett. B 788, 468 (2019) [arXiv:1809.02943 [hep-ph]].
- [77] A. C. Katoch and R. C. Verma, Int. J. Mod. Phys. A 11, 129(1996).
- [78] R. Aaij et al. [LHCb], Phys. Rev. D 94, 072001 (2016) [arXiv:1608.01289 [hep-ex]].
- [79] A. Kuzmin et al. [Belle], Phys. Rev. D 76, 012006 (2007) [arXiv:hep-ex/0611054].
- [80] R. Aaij et al. [LHCb], Phys. Rev. D 92, 032002 (2015) [arXiv:1505.01710 [hep-ex]].
- [81] R. Aaij *et al.* [LHCb], Phys. Rev. D **91**, 092002 (2015) [erratum: Phys. Rev. D **93**, 119901 (2016)] [arXiv:1503.02995 [hep-ex]].
- [82] R. Aaij et al. [LHCb], Phys. Rev. D 92, 012012 (2015) [arXiv:1505.01505 [hep-ex]].
- [83] C. K. Chua, J. Korean Phys. Soc. 45, S256 (2004) [arXiv:hep-ph/0312075].
- [84] Z. Q. Zhang, H. Guo, N. Wang and H. T. Jia, Phys. Rev. D 99, 073002 (2019) [arXiv:1903.03990 [hep-ph]].
- [85] C. Albertus, Phys. Rev. D 89, 065042 (2014) [arXiv:1401.1791 [hep-ph]].
- [86] Z. Q. Zhang, M. G. Wang, Y. C. Zhao, Z. L. Guan and N. Wang, Phys. Rev. D 103, 116030 (2021) [arXiv:2105.02688 [hep-ph]].
- [87] A. Faessler, T. Gutsche, S. Kovalenko and V. E. Lyubovitskij, Phys. Rev. D 76, 014003 (2007) [arXiv:0705.0892 [hep-ph]].
- [88] M. Z. Liu, X. Z. Ling, L. S. Geng, En-Wang and J. J. Xie, Phys. Rev. D 106, 114011 (2022) [arXiv:2209.01103 [hep-ph]].
- [89] Rohit Dhir, Eur. Phys. J. Plus 137, 324 (2022).
- [90] J. Segovia, D. R. Entem, F. Fernandez, E. Hernandez, Int. J. Mod. Phys. E [arXiv:1309.6926 [hep-ph]].
- [91] A. C. Katoch and R. C. Verma, J. Phys. G 22, 1765(1996).