

CUBIQUITOUS LATTICES AND BRANCHED COVERS BOUNDING RATIONAL BALLS

ERICA CHOI, NUR SAGLAM, JONATHAN SIMONE, KATERINA STUOPIS, AND HUGO ZHOU

ABSTRACT. In [8], Greene and Owens explore *cubiquitous* lattices as an obstruction to rational homology 3-spheres bounding rational homology 4-balls. The purpose of this article is to better understand which sublattices of $\mathbb{Z}^n$ are ubiquitous with the aim of effectively using their cubiquity obstruction. We develop a geometric obstruction (called the Wu obstruction) to cubiquity and use it as tool to completely classify which sublattices with orthogonal bases are ubiquitous. We then apply this result the double branched covers of alternating connected sums of torus links. Finally, we explore how the Wu obstruction can be used in conjunction with *contractions* to obstruct the cubiquity of infinite families of lattices.

1. INTRODUCTION

A rational homology 3-sphere is a closed 3-manifold Y satisfying $H_*(Y; \mathbb{Q}) = H_*(S^3; \mathbb{Q})$; a rational homology 4-ball is a 4-manifold X with boundary satisfying $H_*(X; \mathbb{Q}) = H_*(B^4; \mathbb{Q})$. It is a straight-forward exercise to show that the boundary of a rational homology 4-ball is a rational homology 3-sphere. The converse question—does a given rational homology 3-sphere bound a rational homology 4-ball—is a well-known problem in low-dimensional topology (c.f. Problem 4.5 in [11]). Although there are many (partial) classifications of infinite families of rational homology 3-spheres bounding rational homology 4-balls (e.g. [14], [12], [1], [2], [7], [5], [18]), the question at large is wide open.

A rather successful method of obstructing rational homology 3-spheres from bounding rational homology 4-balls comes in the form of lattice embeddings, following from Donaldson's Diagonalization Theorem [6]. A *lattice* is pair (A, Q) , where $A \cong \mathbb{Z}^n$ and Q is a nondegenerate symmetric bilinear form on A . Given two lattices (A, Q) and (B, R) , a *lattice embedding* $i : (A, Q) \rightarrow (B, R)$ is an injective linear map $i : A \rightarrow B$ satisfying $R(i(x), i(y)) = Q(x, y)$ for all $x, y \in A$.

Obstruction 1.1. *Let Y be a rational homology 3-sphere that bounds a 4-manifold X with positive definite intersection form Q_X . Suppose Y also bounds a rational homology 4-ball. Then there exists a lattice embedding*

$$i : (H_2(X), Q_X) \rightarrow (\mathbb{Z}^n, I),$$

where $n = \text{rank}(H_2(X))$ and I denotes the identity matrix.

In [9] (reinterpreted in [8]), it is shown that Heegaard Floer homology d -invariants can often be used to extract additional geometric information from the lattice embedding, which provides a more robust obstruction. This additional geometric information comes in the form of *cubiquity*. We say that a lattice is *cubiquitous* if it admits a lattice embedding into some $\mathbb{Z}^n$ such that its image Λ hits every unit cube in $\mathbb{Z}^n$; namely,

$$\Lambda \cap (x + \{0, 1\}^n) \neq \emptyset$$

for every $x \in \mathbb{Z}^n$. Moreover, we say that the sublattice Λ of $\mathbb{Z}^n$ is ubiquitous.

Obstruction 1.2. *Let Y , X , and i be as in Obstruction 1.1. If X is sharp¹, then i is ubiquitous.*

Cubiquity provides a stronger obstruction (Obstruction 1.2) than the existence of a lattice embedding (Obstruction 1.1). For this reason, this article aims to better understand the question “which sublattices of $\mathbb{Z}^n$ are ubiquitous?” We assume throughout that every sublattice Λ is *full-rank*; that is, $\text{rank}(\Lambda) = \text{rank}(\mathbb{Z}^n) = n$. It is clear that sublattices that are not full-rank cannot be ubiquitous. In [8], Greene-Owens completely classify which full-rank sublattices of $\mathbb{Z}^n$ with “large bases” are ubiquitous.

Theorem 1.3 ([8]). *Let $\Lambda \subset \mathbb{Z}^n$ be a full-rank sublattice and let B be a matrix whose columns form a basis for Λ .*

- *if $\det B > 2^n$, then Λ is not ubiquitous.*
- *if $\det B = 2^n$, then Λ is ubiquitous if and only if Λ has a Hajós basis².*

In the present article, we explore necessary conditions for a sublattice to be ubiquitous, without restriction on the size of the basis (measured via determinant). The main obstructive tool we will develop applies to a large class of sublattices having so-called *non-acute* bases; we defer the precise definition of *non-acute* to Section 2. This obstruction relies on the *Wu element* associated to the basis of a sublattice; given a sublattice with basis $S = \{v_1, \dots, v_n\}$, the Wu element of S is $W = \sum_{i=1}^n v_i$.

Proposition 1.4 (Wu Obstruction). *Let $\Lambda \subset \mathbb{Z}^n$ be a sublattice with basis $S = \{v_1, \dots, v_n\} \subset \mathbb{Z}^n$ which is non-acute. Let k_i be the unique integers satisfying $W = \sum_{i=1}^n k_i e_i$ and set $R_o := \{i : k_i \equiv 1 \pmod{2}\}$. If*

$$\sum_{i=1}^n k_i > 4n - 3|R_o|,$$

then Λ is not ubiquitous.

We use this obstruction to explore the cubiquity of full-rank sublattices with orthogonal bases, which we call *orthogonal sublattices*.

Theorem 1.5. *An orthogonal sublattice of $\mathbb{Z}^n$ is ubiquitous if and only if it admits a basis $\mathcal{B} = \{b_1, \dots, b_n\}$ such that the matrix $B = [b_1 \cdots b_n]$ is a block matrix with blocks of the form $[1]$, $[2]$, and $\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$ up to changing the basis of $\mathbb{Z}^n$.*

1.1. Application to Double Branched Covers. In [8], ubiquitous lattice embeddings are studied in relation to a large family of rational homology 3-spheres—those arising as the double branched covers over non-split alternating links in S^3 (note that the double branched cover along a split link is not a rational homology 3-sphere); we use $\Sigma(L)$ to denote the double cover of S^3 branched along L . Given such a link L , there is a well-defined positive definite lattice $\Lambda(L)$ induced by L , which is unique up to isometry (see the discussion above Theorem 1 in [8] for details). The following result can be viewed as a special case of Obstruction 1.2.

Theorem 1.6 ([8]). *Let L be a non-split alternating link. If $\Sigma(L)$ bounds a rational homology ball, then $\Lambda(L)$ is ubiquitous.*

¹We will not provide a precise definition of *sharp* in this article, as all manifolds under consideration will automatically be sharp; see, e.g. [8] for the precise definition of sharp.

²See [8] for the definition of Hajós basis.

It is further conjectured by Greene-Owens in [8] that in fact, cubiquity is a sufficient condition for $\Sigma(L)$ bounding a rational homology 4-ball.

Conjecture 1.7 ([8]). *For a non-split alternating link L , $\Sigma(L)$ bounds a rational ball $\iff \Lambda(L)$ is cubiquitous.*

In [8], Greene-Owens resolve Conjecture 1.7 for non-split alternating links with $|H_1(\Sigma(L))| = 4^n$; this follows from the second part of Theorem 1.3 (note that if $\Sigma(L)$ bounds a rational homology 4-ball, then by [4] the image of the lattice embedding given by Obstruction 1.1 has a basis whose determinant is precisely $\sqrt{|H_1(\Sigma(L))|}$). Theorem 1.5 provides us with a similar result for alternating connected sums of torus links.

Theorem 1.8. *Conjecture 1.7 holds for alternating connected sums of torus links of the same sign.*

Remark 1.9. Given a family of torus links, there are different ways to perform the connected sum, yielding non-isotopic links as a result. However, it turns out that the isomorphism type of the lattice induced by the connected sum does not depend on the way one performs the connected sum. See the discussion in Section 4.

Remark 1.10. Theorem 1.8 is already implicitly known in the literature. It follows from [13], [10], and calculations of Heegaard Floer homology d -invariants of connected sums of lens spaces. However, Theorem 1.5 provides more information. The results in the literature show that $\Sigma(L)$ —where L is an alternating connected sum of torus links of the same sign—bounds a rational homology 4-ball if and only if the lattice embedding $\Lambda(L) \rightarrow \mathbb{Z}^n$ coming from Obstruction 1.1 is cubiquitous. However, $\Lambda(L)$ can admit many lattice embeddings into $\mathbb{Z}^n$ and it is not clear which of these embeddings comes from Obstruction 1.1; hence Obstruction 1.2 cannot tell us precisely which lattice embedding is cubiquitous. Theorem 1.5, however, tells us exactly which lattice embeddings are cubiquitous.

1.2. Contractions. The notion of *contractions* (defined in Section 5) has been paramount in understanding the existence of certain kinds of sublattices (see, e.g., [14], [13], [17]), which has led a full understanding of which lens spaces bound rational homology 4-balls, for example. In particular, contractions often allow one to create infinite families of sublattices.

Lemma 1.11. *Contractions of non-acute sublattices preserve the Wu obstruction given in Proposition 1.4.*

This straight-forward calculation allows one to obstruct the cubiquity of infinite families of sublattices, which in turn, can aid one in obstructing the existence of rational homology 4-balls bounded by infinite families of rational homology 3-spheres. In Section 5, we will see how this strategy applies to sublattices with so-called *good* bases (defined by Lisca in [14]), which has implications on lens spaces bounding rational homology 4-balls.

Organization. In Section 2, we will consider a particular family of sublattices having bases called *non-acute subsets* and prove Proposition 1.4. In Section 3, we will use this theory to prove Theorem 1.5. In Section 4, we will outline the connection between orthogonal sublattices and alternating connected sums of torus links, and prove Theorem 1.8. Finally, in Section 5, we will prove Lemma 1.11 and apply it to so-called *good subsets*.

Acknowledgements. This work originated in an REU at Georgia Tech that was supported by the NSF grants 1745583, 1851843, 2244427 and the Georgia Tech College of Sciences.

2. NON-ACUTE SUBSETS

For vectors $v, w \in \mathbb{Z}^n$, we use $\langle v, w \rangle$ to denote the standard positive-definite dot product of v and w . We denote the standard orthogonal basis of $\mathbb{Z}^n$ by $\{e_1, \dots, e_n\}$.

Definition 2.1. Let $S = \{v_1, \dots, v_n\} \subset \mathbb{Z}^n$ be a subset with $\langle v_i, v_i \rangle = a_i$. We define the following notation.

$$\begin{aligned} I(S) &:= \sum_{i=1}^n (a_i - 3) \\ E_j^S &:= \{i : \langle v_i, e_j \rangle \neq 0\} \\ V_i^S &:= \{j : \langle v_i, e_j \rangle \neq 0\} \\ \mathcal{P}_i^S &:= \{j : |E_j^S| = i\} \\ p_i(S) &:= |\mathcal{P}_i^S| \end{aligned}$$

Note that the sizes of the above sets can be interpreted using the matrix $[v_1 \ v_2 \ \dots \ v_n]$; indeed, $|E_j^S|$ is the number of columns whose j -th entry is non-zero, $|V_i^S|$ is the number of rows whose i -th entry is non-zero, and $|\mathcal{P}_i^S|$ is the number of rows with exactly i non-zero entries. In some cases we will drop the superscript S from the above notation if the subset being considered is understood.

The following formula is a strengthening of Lemma 2.5 in [14].

Lemma 2.2. *Given any subset $S = \{v_1, \dots, v_n\} \subset \mathbb{Z}^n$,*

$$2p_1(S) + p_2(S) + I(S) = \sum_{j=4}^n (j-3)p_j(S) + \sum_{\substack{s,i \\ \langle v_s, e_i \rangle \neq 0}} (\langle v_s, e_i \rangle^2 - 1)$$

Proof.

$$\begin{aligned} & \sum_{j=4}^n (j-3)p_j(S) + \sum_{\substack{s,i \\ \langle v_s, e_i \rangle \neq 0}} (\langle v_s, e_i \rangle^2 - 1) \\ &= \sum_{j=4}^n j p_j(S) - 3 \sum_{j=4}^n p_j(S) + \sum_{\substack{s,i \\ \langle v_s, e_i \rangle \neq 0}} (\langle v_s, e_i \rangle^2 - 1) \\ &= \sum_{j=4}^n j p_j(S) - 3 \sum_{j=1}^n p_j(S) + 3p_1(S) + 3p_2(S) + 3p_3(S) + \sum_{\substack{s,i \\ \langle v_s, e_i \rangle \neq 0}} (\langle v_s, e_i \rangle^2 - 1) \\ &= \sum_{j=1}^n j p_j(S) - 3n + 2p_1(S) + p_2(S) + \sum_{\substack{s,i \\ \langle v_s, e_i \rangle \neq 0}} (\langle v_s, e_i \rangle^2 - 1) \\ &= \sum_{j=1}^n j p_j(S) - 3n + 2p_1(S) + p_2(S) + \sum_{\substack{s,i \\ \langle v_s, e_i \rangle \neq 0}} \langle v_s, e_i \rangle^2 - \sum_{j=1}^n j p_j(S) \end{aligned}$$

$$\begin{aligned}
&= -3n + 2p_1(S) + p_2(S) + \sum_{i=1}^n a_i \\
&= 2p_1(S) + p_2(S) + \sum_{i=1}^n (a_i - 3) \\
&= 2p_1(S) + p_2(S) + I(S)
\end{aligned}$$

□

Definition 2.3. A set $S = \{v_1, \dots, v_n\}$ is *non-acute* if $a_i \geq 1$ for all i , $\langle v_i, v_j \rangle \leq 0$ for all i, j , and

$$a_i = \langle v_i, v_i \rangle \geq - \sum_{j \neq i} \langle v_j, v_i \rangle.$$

Furthermore, S is called *mediocre* if we further impose:

$$\langle v_i, v_j \rangle = \begin{cases} 0 & |i - j| \geq 1 \\ 0, -1 & |i - j| = 1 \end{cases}$$

Finally, S is called *orthogonal*, if we impose the condition that $\langle v_i, v_j \rangle = 0$ for all $i \neq j$.

Remark 2.4. Lisca's *good subsets* are a special case of mediocre subsets. If $a_i \geq 2$ for all i , then a mediocre subset is good if it is *irreducible*; that is, the corresponding matrix cannot be written as a block matrix with more than one block.

The following technical lemma is well-known (c.f. Remark 2.1 in [14]).

Lemma 2.5. *Every mediocre subset is linearly independent.*

Definition 2.6. Given a subset $S \subset \mathbb{Z}^n$ with Wu element $W = \sum_{i=1}^n k_i e_i$, we define the following sets

$$\begin{aligned}
\mathcal{O}^S &= \{i \mid k_i = 0\} \\
R_o^S &= \{i \mid k_i \equiv 1 \pmod{2}\} \\
R_e^S &= \{i \mid k_i \equiv 0 \pmod{2}, k_i \neq 0\}
\end{aligned}$$

We will usually drop the superscript notation when the subset being considered is understood.

The following cubiquity obstruction is a generalization of Theorem 5.3 in [3].

Proposition 1.4 (Wu Obstruction). *Let $\Lambda \subset \mathbb{Z}^n$ be a sublattice with basis $S = \{v_1, \dots, v_n\} \subset \mathbb{Z}^n$ which is non-acute. Let k_i be the unique integers satisfying $W = \sum_{i=1}^n k_i e_i$ and set $R_o := \{i : k_i \equiv 1 \pmod{2}\}$. If*

$$\sum_{i=1}^n k_i > 4n - 3|R_o|,$$

then Λ is not cubiquitous.

Proof. Let $z = \frac{1}{2}W$. For a unit cube $C \subset \mathbb{Z}^n$, let $\overline{C} \subset \mathbb{R}^n$ denote the solid unit cube containing C . Let $K = \{\text{unit cubes } C \mid z \in \overline{C}\}$. We will show that $\Lambda \cap K = \emptyset$. Notice that $x \in K$ if and only if $W - x \in K$; similarly, $x \in \Lambda$ if and only if $W - x \in \Lambda$. Hence $x \in \Lambda \cap K$ if and only if $W - x \in \Lambda \cap K$.

Let $x \in \Lambda$. Then x is of the form $x = \sum_{i=1}^n x_i v_i$ and $W - x = \sum_{i=1}^n (1 - x_i) v_i$. Hence the dot product of these vectors is

$$\begin{aligned} \langle x, W - x \rangle &= \sum_{i=1}^n x_i (1 - x_i) a_i + \sum_{i < j} (x_i - 2x_i x_j + x_j) \langle v_i, v_j \rangle \\ &= \left(\sum_{i=1}^n x_i (1 - x_i) (a_i + \sum_{j \neq i} \langle v_i, v_j \rangle) \right) - \sum_{i < j} (x_i - x_j)^2 \langle v_i, v_j \rangle \\ &\leq 0. \end{aligned}$$

The last inequality follows from the fact that for non-acute subsets, $a_i \geq -\sum_{j \neq i} \langle v_j, v_i \rangle$.

Next let $x \in K$. Then x is of the form

$$x = \sum_{i \in R_o} \frac{k_i + \epsilon_i}{2} e_i + \sum_{i \in R_e} \frac{k_i + 2\epsilon_i}{2} e_i + \sum_{i \in \mathcal{O}} \epsilon_i e_i,$$

where $\epsilon_i \in \{-1, 1\}$ for all i . Hence $W - x$ is of the form

$$W - x = \sum_{i \in R_o} \frac{k_i - \epsilon_i}{2} e_i + \sum_{i \in R_e} \frac{k_i - 2\epsilon_i}{2} e_i - \sum_{i \in \mathcal{O}} \epsilon_i e_i.$$

Since $\sum_{i=1}^n k_i^2 > 4n - 3|R_o|$, we have that the dot product between these two vectors is

$$\begin{aligned} \langle x, W - x \rangle &= \sum_{i \in R_o} \frac{k_i^2 - 1}{4} + \sum_{i \in R_e} \frac{k_i^2 - 4}{4} - |\mathcal{O}| \\ &= \sum_{i=1}^n \frac{k_i^2}{4} - \frac{|R_o| + 4|R_e| + 4|\mathcal{O}|}{4} \\ &= \sum_{i=1}^n \frac{k_i^2}{4} - \frac{4n - 3|R_o|}{4} \\ &> 0. \end{aligned}$$

The result follows. $\square$

Corollary 2.7. *Let $S \subset \mathbb{Z}^n$ be an orthogonal subset with W element $W = \sum_{i=1}^n k_i e_i$. If*

$$I(S) > n - 3|R_o|,$$

then the sublattice Λ generated by S is not cubiquitous.

Proof. Since S is orthogonal, we can compute $\langle W, W \rangle$ in two simple ways. First, since $W = \sum_{i=1}^n k_i e_i$, $\langle W, W \rangle = \sum_{i=1}^n k_i^2$. On the other hand, since $W = \sum_{i=1}^n v_i$, $\langle W, W \rangle = \sum_{i=1}^n a_i$. Thus the inequality $\sum_{i=1}^n k_i^2 > 4n - 3|R_o|$ can be rewritten as $\sum_{i=1}^n a_i > 4n - 3|R_o|$, or $I(S) > n - 3|R_o|$. Now by Proposition 1.4, the result follows. $\square$

3. PROOF OF THEOREM 1.5

Let S be an orthogonal subset. Note that if there exists an integers i and s such that $V_s = \{i\}$, then by orthogonality $E_i = \{s\}$. It follows that the subset

$$\pi_i(S) := S \setminus \{v_i\} \subset \mathbb{Z}^{n-1}$$

is an orthogonal projection of S onto the subspace $\mathbb{Z}^{n-1} = \langle e_1, \dots, e_{i-1}, e_{i+1}, \dots, e_n \rangle$. Thus by performing all possible orthogonal projections of this form, we may assume that $|V_s| \geq 2$ for all s .

Lemma 3.1. *If $S = \{v_1, \dots, v_n\} \subset \mathbb{Z}^n$ is orthogonal with $|V_s| \geq 2$ for all s , then $p_1(S) = 0$.*

Proof. Assume $p_1(S) \neq 0$. Then there exist integers i and s such that $E_i = \{s\}$; hence $v_s = xe_i + f$ for some $x \in \mathbb{Z} \setminus \{0\}$ and some vector f . If $\langle f, f \rangle > 1$, then the subset

$$(S \setminus \{v_s\}) \cup \{f\} \subset \mathbb{Z}^{n-1}$$

is orthogonal; but by Lemma 2.5, this set has n linearly independent vectors in $\mathbb{Z}^{n-1}$, which is impossible. Thus $\langle f, f \rangle = 1$ and so $v_s = xe_i + e_j$ for some integer $j \neq i$. Furthermore, since S is orthogonal, $|E_j| = 1$; otherwise there would exist an integer t such that $j \in V_t$, implying that $\langle v_s, v_t \rangle \neq 0$. But the fact that $|E_j| = 1$ also leads to a contradiction. Indeed, the subset

$$S \setminus \{v_s\} \subset \mathbb{Z}^{n-2}$$

is orthogonal and hence has $n - 1$ linearly independent vectors in $\mathbb{Z}^{n-1}$ by Lemma 2.5. $\square$

Proposition 3.2. *Let $S = \{v_1, \dots, v_n\} \subset \mathbb{Z}^n$ be orthogonal with $|V_s| \geq 2$ for all s . Let i, s, t be integers such that $E_i = \{s, t\}$ for some integers i, s, t . If S is cubiquitous, then up to a change of basis, $v_s = e_i + e_j$ and $v_t = e_i - e_j$ where $E_j = \{s, t\}$.*

Proof. Let i, s, t be integers such that $E_i = \{s, t\}$. Then $v_s = xe_i + f$ and $v_t = ye_i + g$ for some integers $x, y \neq 0$ and vectors $f, g \in \mathbb{Z}^n$. First suppose $|x|, |y| = 1$. Up to negation, we may assume that $v_s = e_i + f$ and $v_t = e_i + g$. Consider the subset

$$S' = (S \setminus \{v_s, v_t\}) \cup \{f, g\} \subset \mathbb{Z}^{n-1}.$$

It is clear that S' is a mediocre subset. If $\langle f, f \rangle, \langle g, g \rangle \geq 2$, then by Lemma 2.5, S' consists of n linearly independent vectors in $\mathbb{Z}^n$, which is not possible. Thus we may assume, without loss of generality that $\langle f, f \rangle = 1$ and so $v_s = e_i + e_j$ for some integer $j \neq i$. Since $\langle v_s, v_t \rangle = 0$, we necessarily have that $j \in V_t$ and $v_t = e_i - e_j + g'$ for some vector g' . Moreover, it is clear that $E_j = \{s, t\}$. Consider the orthogonal subset

$$S'' = (S \setminus \{v_s, v_t\}) \cup \{g\} \subset \mathbb{Z}^{n-1}.$$

If $\langle g', g' \rangle > 0$, then $p_1(S'') > 0$, which contradicts Lemma 3.1. Hence $v_t = e_i - e_j$.

We call the subset S'' a *double projection* of S . Perform all possible double projections of S and, with abuse of notation, call the resulting subset S . If $p_2(S) = 0$, then there is nothing left to prove. Assume $p_2(S) \neq 0$. Then for each $i \in \mathcal{P}_2$, there exists $s(i) \in E_i$ such that $|\langle v_{s(i)}, e_i \rangle| \geq 2$. Next notice that

$$\begin{aligned} |R_o| &\geq |\{i \in \mathcal{P}_3 : k_i \text{ is odd}\}| \\ &\geq |\{i \in \mathcal{P}_3 : |\langle v_s, e_i \rangle| = 1 \text{ for all } s \in E_i\}| \\ &\geq |\mathcal{P}_3 \setminus \{i : |\langle v_s, e_i \rangle| \geq 2 \text{ for some } s \in E_i\}| \\ &\geq p_3(S) - \sum_{\substack{i \in \mathcal{P}_3 \\ s \in E_i}} (\langle v_s, e_i \rangle^2 - 1). \end{aligned}$$

Now by Lemmas 2.2 and 3.1, and the fact that $\sum_{i=2}^n p_j(S) = n$, we have that

$$\begin{aligned}
I(S) &= -p_2(S) + \sum_{j=4}^n (j-3)p_j(S) + \sum_{\substack{s,i \\ \langle v_s, e_i \rangle \neq 0}} (\langle v_s, e_i \rangle^2 - 1) \\
&\geq n - 2p_2(S) - p_3(S) + \sum_{j=4}^n (j-4)p_j(S) + \sum_{i \in \mathcal{P}_2} (\langle v_{s(i)}, e_i \rangle^2 - 1) + \sum_{\substack{i \in \mathcal{P}_3 \\ s \in E_i}} (\langle v_s, e_i \rangle^2 - 1) \\
&= n - \left(2p_2(S) - \sum_{i \in \mathcal{P}_2} (\langle v_{s(i)}, e_i \rangle^2 - 1) \right) - \left(p_3(S) - \sum_{\substack{i \in \mathcal{P}_3 \\ s \in E_i}} (\langle v_s, e_i \rangle^2 - 1) \right) \\
&\geq n + p_2(S) - |R_o| \\
&> n - 3|R_o|,
\end{aligned}$$

where the second-to-last inequality follows from the fact that $|\langle v_{s(i)}, e_i \rangle| \geq 2$ for all $i \in \mathcal{P}_2$ and last inequality follows from the fact that $p_2(S) > 0$ and $|R_o| \leq 3|R_o|$. Now by Corollary 2.7, S is not cubiqutious. The result follows. $\square$

Before proceeding to the proof of Theorem 1.5, we require one final technical lemma.

Lemma 3.3. *Let $\Lambda_1 \subset \mathbb{Z}^m$ and $\Lambda_2 \subset \mathbb{Z}^n$ be full-rank sublattices and let $\Lambda = \Lambda_1 \oplus \Lambda_2 \subset \mathbb{Z}^k$, where $k = m + n$. Then Λ is cubiqutious if and only if Λ_1 and Λ_2 are cubiqutious.*

Proof. Suppose that Λ_1 and Λ_2 are cubiqutious. Let C be a unit cube in $\mathbb{Z}^k$ and let C_1 and C_2 denote the projections of C to $\mathbb{Z}^m$ and $\mathbb{Z}^n$, respectively. Then there exists $x_1 \in C_1 \cap \Lambda_1$ and $x_2 \in C_2 \cap \Lambda_2$. Since $\Lambda = \Lambda_1 \oplus \Lambda_2$, it follows that $x = (x_1, x_2) \in C \cap \Lambda$; hence Λ is cubiqutious.

For the converse, assume that either Λ_1 or Λ_2 is not cubiqutious; without loss of generality assume the former. Then there is a unit cube $C_1 \subset \mathbb{Z}^m$ that does not intersect Λ_1 . Let $C \subset \mathbb{Z}^k$ be a unit cube that projects onto C_1 under the projection $\mathbb{Z}^k \rightarrow \mathbb{Z}^m$. Let C_2 denote the projection of C to $\mathbb{Z}^n$; then $C = C_1 \oplus C_2$. Pick a vector $x \in C$. Let $x_1 \in C_1$ and $x_2 \in C_2$ be the unique vectors such that $x = (x_1, x_2) \in C_1 \oplus C_2 = C$. Since $C_1 \cap \Lambda_1 = \emptyset$, we have that $x_1 \notin \Lambda_1$. It now follows that since $\Lambda = \Lambda_1 \oplus \Lambda_2$, $x \notin \Lambda$. Hence $C \cap \Lambda = \emptyset$ and so Λ is not cubiqutious. $\square$

Proof of Theorem 1.5. First note that if Λ is an orthogonal sublattice with basis as described in the statement of Theorem 1.5, then it is the direct sum of cubiqutious sublattices. By Lemma 3.3, Λ is cubiqutious. Now suppose $S = \{v_1, \dots, v_n\}$ is an orthogonal subset and let Λ be the lattice generated by S . Let $B = [v_1 \cdots v_n]$. If S is cubiqutious, then by Proposition 3.2, Proposition 3.2, that is, B is a block matrix with blocks of the form $[a]$ and $\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$; that is, Λ is a direct sum of sublattices with bases of the form $\{[a]\}$ and $\{[1, 1]^T, [-1, 1]^T\}$. Since $\text{Span}\{[a]\}$ not cubiqutious if $|a| \leq 3$, it follows from Lemma 3.3 that each 1×1 block must be of the form $[\pm 1]$ or $[\pm 2]$. Up to a change of basis, the result follows. $\square$

4. APPLICATION TO ALTERNATING CONNECTED SUMS OF TORUS LINKS

In this section we consider the applications of our lattice embedding results to the connected sums of torus links, and discuss some related questions. A connected sum of links




FIGURE 1. A connected sum of alternating torus links. Each integer k_i indicates the number of right-handed half twists.

is alternating if and only if each link is alternating ([15, Theorem 1]). Moreover, the only alternating torus links are $T_{2,k}$ for some integer k (see for example, [16, Theorem 7.3.2]), whose double branched cover is $\Sigma(T_{2,k}) = L(k, 1)$. We will assume throughout that each torus knot is negative; that is it is of the form $T(2, k)$, where $k < 1$.

In general, the connected sum of two links L_1 and L_2 depends on the components of each link that are summed together. However, the double branched cover of both of these connected sums are diffeomorphic to $\Sigma(L_1) \# \Sigma(L_2)$. Let L be a connected sum of the torus links $T(2, k_1), \dots, T(2, k_n)$. Note that if we connect sum another alternating torus link $T(2, k_{n+1})$ to L , then the choice of component of $T(2, k_{n+1})$ does not matter; each choice will yield isotopic links (if we do not order the link components of $T(2, k_{n+1})$). Let K and K' be components of L and consider the connected sums L_K and $L_{K'}$ obtained by connect summing $T(2, k_{n+1})$ to K and K' , respectively. We will see that lattices induced by L_K and $L_{K'}$ are isometric; hence we may focus on the lattices induced by the connected sums of the form given in Figure 1.

Proposition 4.1. *Let B_K and $B_{K'}$ denote the black checkerboard surfaces for L_K and $L_{K'}$, respectively. Then $\Sigma(B_K)$ is diffeomorphic to $\Sigma(B_{K'})$.*

Proof. By convention, B_K and $B_{K'}$ are bounded. Let Γ_B and $\Gamma_{B'}$ denote the respective Tait graphs of B and B' . Let v_a and v_b denote the vertices of the Tait graph for L corresponding to K and K' . Let $-a$ and $-b$ denote the weights of these vertices. Consider the Tait graphs of $T(2, k_{n+1})$ and L shown in Figure 2a. The Tait graphs Γ_B and $\Gamma_{B'}$ are then the graphs shown in Figure 2b. To produce handlebody diagrams for $\Sigma(B)$ and $\Sigma(B')$, we must first delete a vertex from each graph. Deleting the vertices corresponding to v_a in both graphs, yields the graphs in Figure 2c. It is now easy to see that the obvious handlebody diagram for $\Sigma(B')$ can be obtained from the obvious handlebody diagram of $\Sigma(B)$ by sliding the 2-handle corresponding to v_b over the 2-handle corresponding to the isolated vertex with weight $-k_{n+1}$. The result follows. $\square$

Corollary 4.2. *Let L be a connected sum of negative torus links $T(2, k_1), \dots, T(2, k_n)$. Then $\Sigma(L)$ bounds a rational ball if and only if $k_i \in \{-1, -2, -4\}$ for all i and there is an even number of parameters equal to -2 .*

Proof. By Proposition 4.1, the lattice embedding associated to L does not depend on the way one performs the connected sum. Consider the specific connected sum diagram depicted in Figure 1. Denote by W the unbounded checkerboard surface of L . Note that if we delete the




FIGURE 2

vertex associated to the unbounded region in the orresponding Tait graph Γ_W , the resulting diagram consists of isolated vertices with weight $-k_i$ for $i = 1, \dots, n$. It follows that $\Lambda(L)$ has matrix representation $\text{diag}\{-k_1, \dots, -k_n\}$.

If $\Sigma(L)$ bounds a rational homology 4-ball, then by Theorem 1.5 and Theorem 1.6 we have that $k_i \in \{-1, -2, -4\}$ for all i and there is an even number of parameters equal to -2 . On the other hand, to show that if the parameters are as stated, then $\Sigma(L)$ bounds a rational homology 4-ball, one can show that the the link shown in Figure 1 bounds a smooth, properly embedded surface in B^4 made up of a disk and a Mobius band; the double branched cover of such a link necessarily bounds a rational homology 4-ball. See [13] for details. $\square$

Proof of Theorem 1.8. The “ $\implies$ ” direction follows from Theorem 1.6. For the “ $\impliedby$ ” direction, let L be an alternating connected sum of n negative torus links. By the discussion above, these torus links are of the form $T(2, k_1), \dots, T(2, k_n)$ for integers $k_i \geq 1$. The result now follows from Corollary 4.2. $\square$

5. A NOTE ON CUBIQUITY AND CONTRACTIONS

The notion of *contractions* has been used in the literature ([14], [13], [17]) to help understand the behavior of particular subsets, which often belong to infinite families. We will see in this section, that contractions preserve the Wu obstruction (Proposition 1.4). This, in turn, allows us to obstruct cubiquity for a large class of subsets.

Definition 5.1. Suppose $S = \{v_1, \dots, v_n\} \subset \mathbb{Z}^n$, $n \geq 3$, such that there exist integers i, s, t , and u such that $E_i = \{s, t, u\}$, where $\langle v_s, v_t \rangle = -1$, $\langle v_s, e_i \rangle = -\langle v_t, e_i \rangle = \pm 1$, $|\langle v_u, e_i \rangle| = 1$, and $a_u \geq 3$. Then the subset $S' \subset \mathbb{Z}^{n-1} = \langle e_1, \dots, e_{i-1}, e_{i+1}, \dots, e_n \rangle$ defined by

$$S' = (S \setminus \{v_s, v_t, v_u\}) \cup \{v_s + v_t, v_u - \langle v_u, e_i \rangle e_i\},$$

is called a *contraction* of S .

Lemma 1.11. *Contractions of non-acute sublattices preserve the Wu obstruction given in Proposition 1.4.*

Proof. Let S be a subset with Wu element $W = \sum_{\alpha=1}^n k_{\alpha} e_{\alpha}$. Let i, s, t, u , and S' be as in Definition 5.1. Let $W' = \sum_{\alpha=1, \alpha \neq i}^n k'_{\alpha} e_{\alpha}$ denote the Wu element of S' . It is easy to see that $k'_i = k_i$ for all $i \neq \alpha$ and hence $W = W' \pm e_i$. It follows that $\sum_{\alpha=1}^n k_{\alpha} > 4n - 3|R_o^S|$ if and only if $\sum_{\alpha=1, \alpha \neq i}^n k'_{\alpha} > 4n - 3|R_o^{S'}|$. In particular, the Wu obstruction given by Proposition 1.4 is preserved by contractions. $\square$

We will now restrict our attention to good subsets as defined by Lisca in [14]. It is worth mentioning that Lisca defines contractions differently than the definition presented here; however these definitions coincide for the subsets considered.

Proposition 5.2. *If S is a good subset with $p_1(S) \neq 0$ then S is cubiquitous if and only if $I(S) \leq 0$.*

Proof. By Lemma 3.2 in [14], any good subset with $p_1(S) > 0$ can be contracted to a length 3 subset with $p_1(S) > 0$. It can be checked by hand that the only such subsets are of the form

$$S' = \{e_1 + e_2, -e_2 + x e_3, e_2 - e_1\},$$

where $x \neq 0$. The Wu element of this subset is $W = x e_3 + e_2$. If $I(S) \leq 0$, then $|x| \geq 3$; by Proposition 1.4, S' is not cubiquitous. Hence S is not cubiquitous. If $I(S) > 0$, then we have either $|x| = 1$ or $|x| = 2$. It is known (see [14]) that in both cases, S corresponds to a lens space that bounds a rational homology 4-ball. It follows that S is cubiquitous. $\square$

Remark 5.3. The proof strategy of Proposition 5.2 can be applied to good subsets with $p_1(S) = 0$, but a general argument is not as clear. In the $p_1(S) \neq 0$ case, good subsets can be contracted to length 3 subsets; this is not the case when $p_1(S) = 0$. In fact, it is unknown in general how small the length of a subset with $p_1 = 0$ can be made via contractions.

REFERENCES

- [1] P. Aceto and M. Golla. “Dehn surgeries and rational homology balls”. In: *Algebr. Geom. Topol.* 17.1 (2017), pp. 487–527. ISSN: 1472-2747. DOI: [10.2140/agt.2017.17.487](https://doi-org.proxy01.its.virginia.edu/10.2140/agt.2017.17.487). URL: <https://doi-org.proxy01.its.virginia.edu/10.2140/agt.2017.17.487>.
- [2] S. Akbulut and K. Larson. “Brieskorn spheres bounding rational balls”. In: *Proc. Amer. Math. Soc.* 146.4 (2018), pp. 1817–1824. ISSN: 0002-9939. DOI: [10.1090/proc/13828](https://doi-org.proxy01.its.virginia.edu/10.1090/proc/13828). URL: <https://doi-org.proxy01.its.virginia.edu/10.1090/proc/13828>.
- [3] V. Brejevs and J. Simone. “On slice quasi-alternating 3-braid closures”. In: *arXiv e-prints* (). arXiv: [2301.06573](https://arxiv.org/abs/2301.06573) [math.GT].
- [4] A. J. Casson and C. M. Gordon. “Cobordism of classical knots”. In: *À la recherche de la topologie perdue*. Vol. 62. Progr. Math. With an appendix by P. M. Gilmer. Birkhäuser Boston, Boston, MA, 1986, pp. 181–199.
- [5] A. Casson and J. Harer. “Some homology lens spaces which bound rational homology balls”. In: *Pacific Journal of Mathematics* 96(1) (1981), pp. 23–36.
- [6] S. K. Donaldson. “The orientation of Yang-Mills moduli spaces and 4-manifold topology”. In: *J. Differential Geom.* 26.3 (1987), pp. 397–428. ISSN: 0022-040X. URL: <http://projecteuclid.org/euclid.jdg/1214441485>.

- [7] H. C. Fickle. “Knots, $\mathbf{Z}$ -homology 3-spheres and contractible 4-manifolds”. In: *Houston J. Math.* 10.4 (1984), pp. 467–493. ISSN: 0362-1588.
- [8] J. Greene and B. Owens. “Alternating links, rational balls, and cube tilings”. In: *arXiv e-prints* (). arXiv: [2212.06248](https://arxiv.org/abs/2212.06248) [[math.GT](https://arxiv.org/archive/math)].
- [9] J. Greene and S. Jabuka. “The slice-ribbon conjecture for 3-stranded pretzel knots”. In: *Amer. J. Math.* 133.3 (2011), pp. 555–580. ISSN: 0002-9327. DOI: [10.1353/ajm.2011.0022](https://doi.org/10.1353/ajm.2011.0022). URL: <https://doi.org/10.1353/ajm.2011.0022>.
- [10] J. E. Greene. “A note on applications of the d -invariant and Donaldson’s theorem”. In: *J. Knot Theory Ramifications* 26.2 (2017), pp. 1740006, 8. ISSN: 0218-2165,1793-6527. DOI: [10.1142/S0218216517400065](https://doi.org/10.1142/S0218216517400065). URL: <https://doi.org/10.1142/S0218216517400065>.
- [11] “Problems in low-dimensional topology”. In: *Geometric topology (Athens, GA, 1993)*. Ed. by R. Kirby. Vol. 2. AMS/IP Stud. Adv. Math. Amer. Math. Soc., Providence, RI, 1997, pp. 35–473.
- [12] A. G. Lecuona. “On the slice-ribbon conjecture for Montesinos knots”. In: *Trans. Amer. Math. Soc.* 364.1 (2012), pp. 233–285. ISSN: 0002-9947. DOI: [10.1090/S0002-9947-2011-05385-7](https://doi.org/10.1090/S0002-9947-2011-05385-7). URL: <https://doi-org.proxy01.its.virginia.edu/10.1090/S0002-9947-2011-05385-7>.
- [13] P. Lisca. “Sums of lens spaces bounding rational balls”. In: *Algebr. Geom. Topol.* 7 (2007), pp. 2141–2164. ISSN: 1472-2747. DOI: [10.2140/agt.2007.7.2141](https://doi.org/10.2140/agt.2007.7.2141). URL: <https://doi.org/10.2140/agt.2007.7.2141>.
- [14] P. Lisca. “Lens spaces, rational balls and the ribbon conjecture”. In: *Geom. Topol.* 11 (2007), pp. 429–472. ISSN: 1465-3060. DOI: [10.2140/gt.2007.11.429](https://doi.org/10.2140/gt.2007.11.429). URL: <https://doi.org/10.2140/gt.2007.11.429>.
- [15] W. Menasco. “Closed incompressible surfaces in alternating knot and link complements”. In: *Topology* 23.1 (1984), pp. 37–44. ISSN: 0040-9383. DOI: [10.1016/0040-9383\(84\)90023-5](https://doi.org/10.1016/0040-9383(84)90023-5). URL: [https://doi.org/10.1016/0040-9383\(84\)90023-5](https://doi.org/10.1016/0040-9383(84)90023-5).
- [16] K. Murasugi. *Knot theory and its applications*. Translated from the 1993 Japanese original by Bohdan Kurpita. Birkhäuser Boston Inc. Boston MA, 1996, pp. viii+341. ISBN: 0-8176-3817-2.
- [17] J. Simone. “Classification of torus bundles that bound rational homology circles”. In: *Algebr. Geom. Topol.* 23.6 (2023), pp. 2449–2518. ISSN: 1472-2747,1472-2739. DOI: [10.2140/agt.2023.23.2449](https://doi.org/10.2140/agt.2023.23.2449). URL: <https://doi.org/10.2140/agt.2023.23.2449>.
- [18] J. Simone. “Using rational homology circles to construct rational homology balls”. In: *Topology Appl.* 291 (2021), Paper No. 107626, 16. ISSN: 0166-8641,1879-3207. DOI: [10.1016/j.topol.2021.107626](https://doi.org/10.1016/j.topol.2021.107626). URL: <https://doi.org/10.1016/j.topol.2021.107626>.