

# FREE GROUP OF HAMEL BIJECTIONS OF BIG SIZE

TOMASZ NATKANIEC

**ABSTRACT.** A  $f: \mathbb{R} \rightarrow \mathbb{R}$  is called *Hamel function* if its graph is a Hamel basis of the linear space  $\mathbb{R}^2$  over rationals. We construct, assuming CH, a free group of the size  $2^c$  contained in the class of all Hamel functions, with the identity function included. This answers, consistently, Question 1 from [1].

## 1. INTRODUCTION

We consider  $\mathbb{R}^2$  as a linear space over the field of rationals. The following notions were introduced by Plotka in his dissertation [3], see also [4]. A function  $f: A \rightarrow \mathbb{R}$ , where  $A \subset \mathbb{R}$ , is called *partially linearly independent* ( $f \in \text{PLIF}$ ) if its graph is a linearly independent subset of  $\mathbb{R}^2$ . If  $A = \mathbb{R}$  then we say that  $f$  is linearly independent ( $f \in \text{LIF}$ ). A function  $f \in \text{LIF}$  is called *Hamel function* ( $f \in \text{HF}$ ) if its graph is a basis of the space  $\mathbb{R}^2$ . Those notions were studied also by other authors. In particular, Matusik and Natkaniec in 2010 paper [2] constructed a bijection  $f: \mathbb{R} \rightarrow \mathbb{R}$  which is a Hamel function. Authors of the recent paper [1] construct a free group of continuum many generators within the family of all Hamel auto-bijections of  $\mathbb{R}$  and they ask if it is possible to find such a free group with  $2^c$  many generators, see [1, Question 1]. In this note we show that if  $2^{<c} = c$  then this query has affirmative answer. Recall that the assumption  $2^{<c} = c$  is independent with ZFC, it follows e.g. from CH, and its negation is a consequence of equalities  $2^{\omega_1} = 2^{\omega_2}$  and  $c = \omega_2$ . We are unable to prove an analogous result in ZFC without additional set-theoretic assumptions.

We use standard notations. In particular,  $|X|$  denotes the cardinality of the set  $X$  and  $Y^X$  denotes the class of all functions from  $X$  to  $Y$ . An ordinal number  $\alpha$  is identified with the set of all ordinals less than  $\alpha$ . An ordinal  $\kappa$  is a cardinal number if it is not equivalent to any ordinal  $\xi < \kappa$ . A function is identified with its graph. For a cardinal  $\kappa$ ,  $2^\kappa$  denotes the set of all functions from  $\kappa$  (i.e., the set of all ordinals less than  $\kappa$ ) to 2 (where  $2 = \{0, 1\}$ ). The symbol  $2^{<c}$  denotes the binary tree of the high  $c$ , i.e. the set of all functions  $f: \xi \rightarrow 2$ ,  $\xi < c$ , ordered by the inclusion.

We will modify the proof of [1, Theorem 1]. For a set  $S$  of injections defined on subsets of  $\mathbb{R}$  let  $G(S)$  denote the group under composition generated by  $S$ , i.e., the family of all compositions  $s_k^{n_k} \circ \dots \circ s_0^{n_0}$ , where  $s_0, \dots, s_k \in S$ ,  $n_0, \dots, n_k \in \mathbb{Z} \setminus \{0\}$ . Note that  $|G(S)| \leq c$  whenever  $|S| \leq c$ . We say that  $G(S)$  is a *free group* if each element of  $G$  has a unique representation as a product  $s_0^{n_0} \circ \dots \circ s_k^{n_k}$ , where

---

*Date:* 09.05.2023.

2010 *Mathematics Subject Classification.* Primary: 20B99; Secondary: 03E35, 15A03, 26A18, 46A35.

*Key words and phrases.* Hamel bases, Hamel functions, free group.

$s_0, \dots, s_k \in S$ ,  $n_0, \dots, n_k \in \mathbb{Z} \setminus \{0\}$  and  $s_i \neq s_{i+1}$  for  $i = 0, \dots, k-1$ . Then the set  $S$  will be called the set of *free generators* of  $G(S)$ .

For  $A \subset \mathbb{R}^i$ ,  $i = 1, 2$ , the symbol  $\text{LIN}(A)$  denotes the linear subspace of  $\mathbb{R}^i$  over  $\mathbb{Q}$  spanned by  $A$ . We will use the following facts.

**Lemma 1.** [2, Fact 2.2]. *Let  $f \in \text{LIF}$ . If  $\langle 0, x \rangle \in \text{LIN}(f)$  then  $f \in \text{HF}$ .*

**Lemma 2.** *Assume that  $A, B, C$  are subsets of  $\mathbb{R}$  such that*

- (1)  $|A| = \mathfrak{c}$  and  $|C| < \mathfrak{c}$ ;
- (2)  $A \cap B = \emptyset$  and  $A \cup B$  is linearly independent over  $\mathbb{Q}$ .

*Then  $A \setminus \text{LIN}(C \cup B) \neq \emptyset$ .*

*Proof.* We may assume that  $C$  is infinite. Let  $|C| = \kappa$ , list  $\text{LIN}(C) = \{c_\xi : \xi < \kappa\}$ . Then  $\text{LIN}(C \cup B) = \bigcup_{\xi < \kappa} (c_\xi + \text{LIN}(B))$ . Suppose that  $A \subset \text{LIN}(C \cup B)$ . Then there are different  $x_1, x_2 \in A$  and  $\xi < \kappa$  such that  $x_1, x_2 \in c_\xi + \text{LIN}(B)$ . Hence  $0 \neq x_1 - x_2 \in \text{LIN}(A) \cap \text{LIN}(B)$ , contrary to (2).  $\square$

**Theorem 3.** *Assume that  $2^{<\mathfrak{c}} = \mathfrak{c}$ . Then there exists a family  $\{f_s : s \in 2^{\mathfrak{c}}\}$  of Hamel bijections which are free generators of a group under composition contained in the class of Hamel bijections plus the identity.*

*Proof.* Let  $\mathbb{R} = \{x_\alpha : \alpha < \mathfrak{c}\}$  with  $x_0 = 0$ . Let  $\{H_{\kappa, \alpha, i} : \kappa, \alpha < \mathfrak{c}, i < 3\}$  be the family of pairwise disjoint subsets of  $\mathbb{R}$ , each of size  $\mathfrak{c}$  with the union  $\bigcup_{\kappa, \alpha < \mathfrak{c}, i < 3} H_{\kappa, \alpha, i}$  being linearly independent. Let

$$W_\kappa = \text{LIN} \left( \{x_\xi : \xi < \kappa\} \cup \bigcup_{\xi < \kappa, \alpha < \mathfrak{c}, i < 3} H_{\xi, \alpha, i} \right).$$

We will construct, by induction, a family of real-valued functions  $\{f_s : s \in 2^{<\mathfrak{c}}\}$  defined on subsets of  $\mathbb{R}$ , such that for each  $\kappa < \mathfrak{c}$  the family  $S_\kappa := \{f_s : s \in 2^\kappa\}$  satisfies the following conditions

- (I):  $f_{s|_\lambda} \subset f_s$  for each  $s \in 2^\kappa$  and  $\lambda < \kappa$ ;
- (II):  $f_s$  is a 1-1 map from a subset of the set  $W_\kappa$  into  $W_\kappa$  for each  $s \in 2^\kappa$ ;
- (III):  $G(S_\kappa) \subset \text{PLIF}$ ;
- (IV):  $\langle 0, x_\kappa \rangle \in \text{LIN}(f)$  for every  $f \in G(S_{\kappa+1})$ ;
- (V):  $\text{dom}(f_s \setminus f_{s|_\kappa}) \subset \text{LIN} \left( \{x_\kappa\} \cup \bigcup_{\alpha < \mathfrak{c}, i < 3} H_{\kappa, \alpha, i} \right)$  for  $s \in 2^{\kappa+1}$ ;
- (VI):  $x_\kappa \in \text{dom}(f_s) \cap \text{rng}(f_s)$  for all  $s \in 2^{\kappa+1}$ ;

Then, for  $s \in 2^{\mathfrak{c}}$ , let  $f_s = \bigcup_{\kappa < \mathfrak{c}} f_{s|_\kappa}$  and let  $S = \{f_s : s \in 2^{\mathfrak{c}}\}$ . Note that conditions (I), (II), and (VI) imply that  $f_s$  is an autobijection on  $\mathbb{R}$  for every  $s \in 2^{\mathfrak{c}}$ . Statements (III) and (IV) give that  $G(S) \subset \text{HF}$ . Moreover, the condition  $G(S) \subset \text{LIF}$  yields also that  $S$  generates a free group – see [1]. Indeed, suppose that there is a function  $g \in G(S)$  with two different representations

$$f_{s_k}^{n_k} \circ \dots \circ f_{s_0}^{n_0} = f_{t_k}^{m_k} \circ \dots \circ f_{t_0}^{m_0}.$$

Then

$$f_{s_k}^{n_k} \circ \dots \circ f_{s_0}^{n_0} \circ f_{t_0}^{-m_0} \circ \dots \circ f_{t_k}^{-m_k} = \text{id}_{\mathbb{R}}$$

where the left side written in reduced form remains non-trivial, so it belongs to the group  $G(S)$  and consequently to  $\text{LIF}$ , while  $\text{id}_{\mathbb{R}} \notin \text{LIF}$ . In particular, we get  $f_s \neq f_{s'}$  for  $s \neq s'$ . Thus  $|S| = 2^{\mathfrak{c}}$  and therefore also  $|G(S)| = 2^{\mathfrak{c}}$ .

So, fix  $\lambda < \mathfrak{c}$  and assume that for all  $\xi < \lambda$  we have functions  $f_t$ ,  $t \in 2^\xi$  satisfying the conditions (I)–(VI). If  $\lambda$  is a limit ordinal then set

$$f_s = \bigcup_{\xi < \lambda} f_{s|\xi}$$

for each  $s \in 2^\lambda$ . Evidently,  $f_s$ 's satisfy conditions (I)–(VI). So we can assume that  $\lambda = \kappa + 1$  and  $f_s$  are defined for all  $s \in 2^\kappa$ .

**Step I.** The main problem is to extend all functions  $f_s$ ,  $s \in 2^\kappa$ , so as to keep the conditions (III) and (IV). Since the group  $G(S_\kappa)$  has cardinality  $\leq \mathfrak{c}$ , we can write

$$G(S_\kappa) = \{g_\alpha : \alpha < \mathfrak{c}\}.$$

Let  $g_\alpha = f_{s_k}^{n_k} \circ \dots \circ f_{s_0}^{n_0}$  where  $f_{s_i} \in S_\kappa$ ,  $n_i \in \mathbb{Z} \setminus \{0\}$  for  $i \leq k$ , and  $s_i \neq s_{i+1}$  for  $i = 0, \dots, k-1$ . For every  $s \in 2^\kappa$  and  $\alpha < \mathfrak{c}$  we will choose, inductively on  $\alpha$ , the function  $f_s^{(\alpha)}$  such that

- $f_s^{(0)} = f_s$ ;
- if  $\beta < \alpha$  then  $f_s^{(\beta)} \subset f_s^{(\alpha)}$ ;
- $f_s^{(\alpha)}$  is one-to-one;
- $\langle 0, x_\kappa \rangle \in \text{LIN}(g_\alpha^{(\alpha)})$ , where  $g_\alpha^{(\alpha)} = (f_{s_k}^{(\alpha)})^{n_k} \circ \dots \circ (f_{s_0}^{(\alpha)})^{n_0}$ , i.e.,  $g_\alpha^{(\alpha)}$  is derived from  $g_\alpha$  by substituting the component functions  $f_{s_i}$  with  $f_{s_i}^{(\alpha)}$ ;
- $G(S_\kappa^{(\alpha)}) \subset \text{PLIF}$ , where  $S_\kappa^{(\alpha)} = \{f_s^{(\alpha)} : s \in 2^\kappa\}$ .

Fix  $\alpha < \mathfrak{c}$ . Let  $f'_s = \bigcup_{\xi < \alpha} f_s^{(\xi)}$  for  $s \in 2^\kappa$  and let  $g'_\alpha = (f'_{s_k})^{n_k} \circ \dots \circ (f'_{s_0})^{n_0}$ . Let  $k_\alpha = \sum_{i=0}^k |n_i|$ . We will consider two cases.

**Case 1.** If  $\langle 0, x_\kappa \rangle \in \text{LIN}(g'_\alpha)$  then set  $f_s^{(\alpha)} = f'_s$  for all  $s \in 2^\kappa$ .

**Case 2.** Otherwise we choose finite sequences  $\langle z_0^\alpha, \dots, z_{k_\alpha}^\alpha \rangle$ ,  $\langle r_0^\alpha, \dots, r_{k_\alpha}^\alpha \rangle$  such that

- (1)  $z_i^\alpha \in H_{\kappa, \alpha, 0} \setminus \text{LIN}(\{z_j^\alpha : j < i\} \cup \{x_\kappa\} \cup \bigcup_{\beta < \alpha, i < 3} H_{\kappa, \beta, i} \cup \bigcup_{\xi < \kappa} W_\xi)$  for  $i = 0, \dots, k_\alpha - 1$ ;
- (2)  $r_0^\alpha = -z_0^\alpha$ ;
- (3)  $r_i^\alpha \in H_{\kappa, \alpha, 1} \setminus \text{LIN}(\{r_j^\alpha : j < i\} \cup H_{\kappa, \alpha, 0} \cup \{x_\kappa\} \cup \bigcup_{\beta < \alpha, i < 3} H_{\kappa, \beta, i} \cup \bigcup_{\xi < \kappa} W_\xi)$  for  $i = 1, \dots, k_\alpha$ ;
- (4)  $z_{k_\alpha}^\alpha = x_\kappa - r_{k_\alpha}^\alpha$ .

Such a choice is possible by Lemma 2 as

$$\begin{aligned} & \text{LIN} \left( \{z_j^\alpha : j < i\} \cup \{x_\kappa\} \cup \bigcup_{\beta < \alpha, i < 3} H_{\kappa, \beta, i} \cup \bigcup_{\xi < \kappa} W_\xi \right) = \\ & \text{LIN} \left( \{z_j^\alpha : j < i\} \cup \{x_\xi : \xi \leq \kappa\} \cup \bigcup_{\beta < \alpha, i < 3} H_{\kappa, \beta, i} \cup \bigcup_{\xi < \kappa, \beta < \mathfrak{c}, i < 3} H_{\xi, \beta, i} \right) \end{aligned}$$

and the set

$$\text{LIN} \left( \{r_j^\alpha : j < i\} \cup H_{\kappa, \alpha, 0} \cup \{x_\kappa\} \cup \bigcup_{\beta < \alpha, i < 3} H_{\kappa, \beta, i} \cup \bigcup_{\xi < \kappa} W_\xi \right)$$

can be written in a similar way. Our goal is to extend a partial functions  $f'_s$ ,  $s \in 2^\kappa$ , to  $f_s^{(\alpha)}$  to get the conditions  $g_\alpha^{(\alpha)}(z_0^\alpha) = z_{k_\alpha}^\alpha$ , and  $g_\alpha^{(\alpha)}(r_0^\alpha) = r_{k_\alpha}^\alpha$ . The

idea of this construction is exactly the same as in [1]. We wander, for each  $\alpha < \mathfrak{c}$ , according to the structure of  $g_\alpha$  from  $z_0^\alpha$  to  $z_{k_\alpha}^\alpha$  via  $z$ 's and from  $r_0^\alpha$  to  $r_{k_\alpha}^\alpha$  via  $r$ 's. Then  $\langle z_0^\alpha, z_{k_\alpha}^\alpha \rangle, \langle r_0^\alpha, r_{k_\alpha}^\alpha \rangle \in g_\alpha^{(\alpha)}$ , and since  $r_0^\alpha = -z_0^\alpha$ ,  $r_{k_\alpha}^\alpha = x_\kappa - z_{k_\alpha}^\alpha$ , so  $\langle 0, x_\kappa \rangle \in \text{LIN}(g_\alpha^{(\alpha)})$ . So, if  $s \neq s_i$  for all  $i \leq k$  then  $f_s^{(\alpha)} = f'_s$ . Otherwise define an increasing sequence

- $p_0 = 0$ ;
- $p_j = \sum_{i=0}^{j-1} |n_i|$  for  $0 < j < k$ ;
- $p_k = k_\alpha$ .

Then for  $j < k$  we extend  $f'_{s_j}$  in the following way.

- If  $n_j < 0$  then

$$f_{s_j}^{(\alpha)} = f'_{s_j} \cup \left\{ \langle z_{p_j+1}^\alpha, z_{p_j}^\alpha \rangle, \dots, \langle z_{p_{j+1}}^\alpha, z_{p_{j+1}-1}^\alpha \rangle, \langle r_{p_j+1}^\alpha, r_{p_j}^\alpha \rangle, \dots, \langle r_{p_{j+1}}^\alpha, r_{p_{j+1}-1}^\alpha \rangle \right\}.$$

- If  $n_j > 0$  then

$$f_{s_j}^{(\alpha)} = f'_{s_j} \cup \left\{ \langle z_{p_j}^\alpha, z_{p_{j+1}}^\alpha \rangle, \dots, \langle z_{p_{j+1}-1}^\alpha, z_{p_{j+1}}^\alpha \rangle, \langle r_{p_j}^\alpha, r_{p_{j+1}}^\alpha \rangle, \dots, \langle r_{p_{j+1}-1}^\alpha, r_{p_{j+1}}^\alpha \rangle \right\}.$$

To prove the inclusion  $G(S_\kappa^{(\alpha)}) \subset \text{PLIF}$  fix  $\beta < \mathfrak{c}$  and assume that

- $g'_\beta$  is derived from  $g_\beta$  by substituting the component functions  $f_{s_i}$  with  $\bigcup_{\xi < \alpha} f_{s_i}^{(\xi)}$ , and
- $g_\beta^{(\alpha)}$  is derived from  $g_\beta$  by substituting the functions  $f_{s_i}$  with  $f_{s_i}^{(\alpha)}$  (hence  $g'_\beta \subset g_\beta^{(\alpha)}$ ).

Suppose that

$$\sum_{i=0}^m q_i \langle v_i, y_i \rangle = 0$$

for some  $q_0, \dots, q_m \in \mathbb{Q}$  and  $\langle v_0, y_0 \rangle, \dots, \langle v_m, y_m \rangle \in g_\beta^{(\alpha)}$  with  $v_i \neq v_j$  for  $i \neq j$ . Then  $\sum_{i=0}^m q_i v_i = 0$ . Let  $N_0 := \{i \leq m : v_i \in \text{dom}(g'_\beta)\}$  and  $N_1 := \{i \leq m : v_i \notin \text{dom}(g'_\beta)\}$ . If  $N_1 = \emptyset$  then  $q_1 = \dots q_m = 0$  since  $g'_\beta \in \text{PLIF}$ . So assume that  $N_1 \neq \emptyset$ . Observe that then  $g_\beta$  is “a part” of  $g_\alpha$  in the sense that if  $g_\alpha = f_{s_k}^{n_k} \circ \dots \circ f_{s_0}^{n_0}$  then there are  $k' \leq k$  and  $i_0 \geq 0$  such that

$$g_\beta = f_{s_{k'}}^{m_{k'}} \circ \dots \circ f_{s_{i_0}}^{m_{i_0}}$$

with  $m_i = n_i$  for  $i \in \{i_0 + 1, \dots, k' - 1\}$  and  $m_{i_0} \in [0, n_{i_0}]$ ,  $m_{k'} \in [0, n_{k'}]$ .

We have two cases to consider.

- First, assume that  $\{r_0^\alpha, z_0^\alpha\} \not\subset \{v_i : i \leq m\}$ . Then the set  $\{v_i : i \in N_1\}$  is linearly independent and

$$\text{LIN}(\{v_i : i \in N_1\}) \cap \text{LIN}(\{v_i : i \in N_0\}) = \{0\},$$

so  $q_i = 0$  for all  $i \in N_1$ . Hence

$$\sum_{i \in N_0} q_i \langle v_i, y_i \rangle = \langle 0, 0 \rangle$$

and since  $g'_\beta \in \text{PLIF}$ , so  $q_i = 0$  for  $i \in N_0$ .

- (ii) Otherwise,  $\{r_0^\alpha, z_0^\alpha\} \subset \{v_i : i \leq m\}$ . Then  $i_0 = 0$  and  $m_{i_0} = n_{i_0}$ . Let  $r_0^\alpha = v_j$ ,  $z_0^\alpha = v_k$ . Then

$$0 = \sum_{i \leq m} q_i y_i = \sum_{i \in N_0} q_i g'_\beta(v_i) + \sum_{i \in N_1} q_i g_\beta^{(\alpha)}(v_i).$$

Now we have the following subcases.

- If  $g_\beta^{(\alpha)}(\{v_j, v_k\}) \cap \{r_{k_\alpha}^\alpha, z_{k_\alpha}^\alpha\} = \emptyset$  then the set  $\{g_\beta^{(\alpha)}(v_i) : i \in N_1\}$  is linearly independent and  $\text{LIN}(\{g_\beta^{(\alpha)}(v_i) : i \in N_1\}) \cap \text{LIN}(\text{rng}(g'_\beta)) = \{0\}$ . Hence  $q_i = 0$  for  $i \in N_1$  and since  $g'_\beta \in \text{PLIF}$ ,  $q_i = 0$  for  $i \in N_0$ .
- Otherwise  $k' = k$  and  $m_{k'} = n_k$  and therefore  $g_\beta = g_\alpha$ .

**Step II.** At this step we will expand each  $f_s'' := \bigcup_{\alpha < \mathfrak{c}} f_s^{(\alpha)}$ ,  $s \in 2^\kappa$ , to two functions  $f_{s \smallfrown 0}$  and  $f_{s \smallfrown 1}$  so as to provide the condition (VI). Since  $|2^\lambda| \leq \mathfrak{c}$ , we can write  $2^\lambda = \{s_\alpha : \alpha < \mathfrak{c}\}$ . We will work inductively on  $\alpha$ . Note that

$$V_\alpha := \bigcup_{t \in 2^\kappa} (\text{dom}(f'_t) \cup \text{rng}(f'_t)) \subset \text{LIN} \left( \{x_\kappa\} \cup \bigcup_{\xi < \kappa} V_\xi \cup \bigcup_{\alpha < \mathfrak{c}, i < 2} H_{\kappa, \alpha, i} \right).$$

Fix  $\alpha < \mathfrak{c}$  and set  $t = s_\alpha \smallfrown \kappa$ . If  $x_\kappa \in \text{dom}(f''_t)$  then  $f''_{s_\alpha} = f''_t$ . Otherwise choose

$$y_\alpha \in H_{\kappa, \alpha, 2} \setminus \text{LIN}(\{y_\beta, z_\beta : \beta < \alpha\} \cup V_\alpha).$$

Such a choice is possible by Lemma 2. Let  $f'''_t := f''_t \cup \langle x_\kappa, y_\alpha \rangle$ . Now, if  $x_\kappa \in \text{rng}(f'''_t)$  then  $f_{s_\alpha} = f'''_t$ . Otherwise, again use Lemma 2 and choose

$$z_\alpha \in H_{\kappa, \alpha, 2} \setminus \text{LIN}(\{x_\alpha\} \cup \{y_\beta, z_\beta : \beta < \alpha\} \cup V_\alpha)$$

and put  $f_{s_\alpha} = f'''_t \cup \langle z_\alpha, x_\kappa \rangle$ .

□

## REFERENCES

- [1] M. Lichman, M. Pawlikowski, S. Smolarek, J. Swaczyna, *Free group of Hamel functions*, Banach Cent. Publ., 125 (2023), 91–98.
- [2] G. Matusik, T. Natkaniec, *Algebraic properties of Hamel functions*, Acta Math. Hung. 126 (3), (2010), 209–229.
- [3] K. Plotka, *Set-theoretic and algebraic properties of certain families of real functions*, Ph.D. Thesis, West Virginia University 2001.
- [4] K. Plotka, *On functions whose graph is a Hamel basis*, Proc. Amer. Math. Soc., 131 (2003), 1031–1041.

INSTITUTE OF MATHEMATICS, FACULTY OF MATHEMATICS, PHYSICS AND INFORMATICS, UNIVERSITY OF GDAŃSK, 80-308 GDAŃSK, POLAND  
Email address: Tomasz.Natkaniec@ug.edu.pl