

CHARACTERISTIC DETERMINANTS FOR A SECOND ORDER DIFFERENCE EQUATION ON THE HALF-LINE ARISING IN HYDRODYNAMICS

YURI LATUSHKIN AND SHIBI VASUDEVAN

To Yuriy Karlovich on his seventy fifth birthday

ABSTRACT. We study the point spectrum of a second order difference operator with complex potential on the half-line via Fredholm determinants of the corresponding Birman-Schwinger operator pencils, the Evans and the Jost functions. An application is given to instability of a generalization of the Kolmogorov flow for the Euler equation of ideal fluid on the two dimensional torus.

1. INTRODUCTION AND MAIN RESULTS

In this paper we continue the work began in [39] and study the eigenvalues of the following boundary value problem for the second order asymptotically autonomous difference equation on the half-line $\mathbb{Z}_+ = \{0, 1, \dots\}$,

$$z_{n-1} - z_{n+1} + (b_n c_n) z_n = \lambda z_n, \quad n \geq 0, \quad (1)$$

$$z_{-1} = 0, \quad (2)$$

where $\lambda \in \mathbb{C}$ is the spectral parameter,

$$(b_n)_{n \geq 0} \in \ell^2(\mathbb{Z}_+; \mathbb{C}) \text{ and } (c_n)_{n \geq 0} \in \ell^2(\mathbb{Z}_+; \mathbb{C}) \quad (3)$$

are two given complex valued sequences that, in general, may depend on λ holomorphically. We are seeking to characterize the values of the spectral parameter such that (1)–(2) has a nontrivial solution $\mathbf{z} = (z_n)_{n \geq 0} \in \ell^2(\mathbb{Z}_+; \mathbb{C})$. This question is important in stability issues for special steady state solutions of the two-dimensional Euler equation, the so called *generalized Kolmogorov flow*, and our main application is a result on its instability. Specifically, in the current paper we define (and prove that they are being equal) four functions of the spectral parameter whose zeros are the eigenvalues of (1)–(2). We call the functions *characteristic determinants*. Our overall strategy is as in [39] where the full line case has been considered, but the treatment of the important half-line case is technically more challenging; in particular, in the current paper we have to prove from the outset for the half-line case analogs of some results obtained in [12] for the line case and essentially used in [39]. By reflection, analogous results hold for the equations on the negative half-line.

Date: May 3, 2024.

Key words and phrases. 2D Euler equations, unidirectional flows, continued fractions, Jost function, Fredholm determinants, Evans function, unstable eigenvalues, half-line equations.

Y.L was supported by the NSF grant DMS-2106157, and would like to thank the Courant Institute of Mathematical Sciences at NYU and especially Prof. Lai-Sang Young for their hospitality.

An important particular case of (1)–(2) is the following eigenvalue problem for a difference equation arising in stability analysis of the generalized Kolmogorov flow of the Euler equation of ideal fluids on 2D torus, as seen below and considered in [14] and [39],

$$z_{n-1} - z_{n+1} = \lambda z_n / \rho_n, \quad n \geq 0, \quad z_{-1} = 0, \quad (4)$$

where $(\rho_n)_{n \geq 0}$ is a given sequence satisfying the following conditions:

$$\rho_0 < 0, \rho_n \in (0, 1), n \geq 1, \text{ and } \rho_n = 1 + O(1/n^2) \text{ as } |n| \rightarrow \infty. \quad (5)$$

The problem (4) is reduced to (1)–(2) by setting

$$b_n = -\lambda \sqrt{1 - \rho_n} / \rho_n \text{ and } c_n = \sqrt{1 - \rho_n}. \quad (6)$$

For (4) we use continued fractions to define yet another, fifth function, also proven to be equal to the previous four characteristic determinants, whose zeros are the eigenvalues. This result, in turn, yields instability of the generalized Kolmogorov flow.

It is convenient to re-write (1)–(2) as

$$(S - S^{-1} + \text{diag}_{n \in \mathbb{Z}_+} \{b_n c_n\}) \mathbf{z} = \lambda \mathbf{z}, \quad (7)$$

where we denote by $S : (z_n)_{n \geq 0} \mapsto (0, z_0, z_1, \dots)$ the right shift operator on $\ell^2(\mathbb{Z}_+; \mathbb{C})$ and by $S^{-1} = S^* : \mathbf{z} \mapsto (z_1, z_2, \dots)$ the left shift operator such that $S^{-1}S = I_{\ell^2(\mathbb{Z}_+; \mathbb{C})}$, the identity operator, while $\text{ran}(SS^{-1}) = (\text{span}\{(1, 0, \dots)\})^\perp$. By the spectral mapping theorem $\text{Spec}(S - S^{-1}) = [-2i, 2i]$ since $\text{Spec}(S) = \{\lambda \in \mathbb{C} : |\lambda| \leq 1\}$, and so throughout the paper we assume that $\lambda \notin [-2i, 2i]$ thus looking for the *isolated eigenvalues* of (1)–(2).

Letting $y_n = \begin{bmatrix} z_n \\ z_{n-1} \end{bmatrix} \in \mathbb{C}^2$, the problem (1)–(2) is equivalent to the following boundary value problem for the first order (2×2) -system of difference equations,

$$y_{n+1} = A_n^\times y_n, \quad n \geq 0, \quad (8)$$

$$y_0 \in \text{ran}(Q_+). \quad (9)$$

Here and throughout the paper we use the following notations,

$$A_n^\times = A + B_n C_n := \begin{bmatrix} b_n c_n - \lambda & 1 \\ 1 & 0 \end{bmatrix} \in \mathbb{C}^{2 \times 2}, \quad n \geq 0. \quad (10)$$

$$A = A(\lambda) = \begin{bmatrix} -\lambda & 1 \\ 1 & 0 \end{bmatrix}, Q_+ = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, Q_- = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}, B_n = b_n Q_+, C_n = c_n Q_{11}$$

The eigenvalues of the matrix $A(\lambda)$ solve the quadratic equation $\mu^2 + \lambda\mu - 1 = 0$, and so our standing assumption $\lambda \notin [-2i, 2i]$ is equivalent to the fact that the eigenvalues of $A(\lambda)$ are off the unit circle. We let $\mu_+ = \mu_+(\lambda)$ and $\mu_- = \mu_-(\lambda)$ denote the roots of the equation $\mu^2 + \lambda\mu - 1 = 0$ satisfying the inequalities

$$|\mu_+(\lambda)| < 1 < |\mu_-(\lambda)| \quad (12)$$

and denote by $P_\pm = P_\pm(\lambda)$ the spectral projections for $A(\lambda)$ in $\mathbb{C}^2$ such that $\text{Spec}(A(\lambda)|_{\text{ran } P_\pm}) = \{\mu_\pm(\lambda)\}$. Finally, we let $R_+ = R_+(\lambda)$ denote the projection in $\mathbb{C}^2$ onto $\text{ran } P_+$ parallel to $\text{ran } Q_+$, and set $R_- = I_{2 \times 2} - R_+$.

The choice of the projection R_+ is important for the half-line case. Indeed, the constant coefficient difference equation $y_{n+1} = A y_n$ has exponential dichotomy on $\mathbb{Z}_+$ with the dichotomy projection P_+ whose range is the uniquely determined subspace of the initial values of the bounded solutions to the equation. Unlike the full

line case, the exponential dichotomy on $\mathbb{Z}_+$ is not unique, but the only requirement on the dichotomy projection is that its range must be equal to the subspace $\text{ran } P_+$. The choice of R_+ whose kernel is $\text{ran } Q_+$ as the dichotomy projection will allow us to satisfy the boundary condition in (2), see, e.g., formula (34) below.

We now proceed to define our first characteristic determinant associated with (1)–(2). We consider the following Birman-Schwinger-type pencil of operators acting in $\ell^2(\mathbb{Z}_+; \mathbb{C})$,

$$K_\lambda^+ = -\text{diag}_{n \in \mathbb{Z}_+} \{c_n\} (S - S^{-1} - \lambda)^{-1} \text{diag}_{n \in \mathbb{Z}_+} \{b_n\}, \quad (13)$$

analogous to K_λ studied in [39] for the full line. By (3), the operator K_λ^+ is of trace class and so we may define our first characteristic determinant $\det(I - K_\lambda^+)$.

Next, we re-write (8) as $(S^{-1} - \text{diag}_{n \in \mathbb{Z}_+} \{A_n^\times\})\mathbf{y} = 0$ for $\mathbf{y} = (y_n)_{n \in \mathbb{Z}_+}$ where we continue to denote by S^{-1} the shift acting in the space of vector valued sequences. We stress that the operator $S^{-1} - \text{diag}_{n \in \mathbb{Z}_+} \{A_n^\times\}$ in (8)–(9) acts from the subspace

$$\ell_{\text{ran}(Q_+)}^2(\mathbb{Z}_+; \mathbb{C}^2) := \{(y_n)_{n \geq 0} \in \ell^2(\mathbb{Z}_+; \mathbb{C}^2) : y_0 \in \text{ran}(Q_+)\} \quad (14)$$

into $\ell^2(\mathbb{Z}_+; \mathbb{C}^2)$ and refer to [6] for results on equivalence of invertibility of the operator and dichotomy of the difference equation (8) on the half-line. We introduce the Birman-Schwinger-type pencil of operators acting in $\ell^2(\mathbb{Z}_+; \mathbb{C}^2)$,

$$\mathcal{T}_\lambda^+ = \text{diag}_{n \in \mathbb{Z}_+} \{C_n\} (S^{-1} - \text{diag}_{n \in \mathbb{Z}_+} \{A_n\})^{-1} \text{diag}_{n \in \mathbb{Z}_+} \{B_n\}, \quad (15)$$

analogous to $\mathcal{T}_\lambda$ studied in [39] for the full line case. The operator $\mathcal{T}_\lambda^+$ is of trace class by (3) and we define our second characteristic determinant $\det(I - \mathcal{T}_\lambda^+)$.

We now consider the (2×2) -matrix valued Jost solution $\mathbf{Y}^+ = \mathbf{Y}^+(\lambda) = (Y_n^+)_{n \geq 0}$, $Y_n^+ \in \mathbb{C}^{2 \times 2}$, cf. [12, 23, 39], defined as the solution to the difference equation (8) (without the boundary condition) satisfying

$$\|(\mu_+)^{-n} (Y_n^+ - A^n R_+)\|_{\mathbb{C}^{2 \times 2}} \rightarrow 0 \text{ as } n \rightarrow +\infty \text{ and } Y_0^+ = Y_0^+ R_+. \quad (16)$$

As we will prove below, this solution is unique. Also, $Y_n^+ = Y_n^+ R_+$ for all $n \geq 0$. Using the projection $R_- = R_-(\lambda)$ onto $\text{ran } Q_+$ parallel to $\text{ran } P_+(\lambda)$, analogously to $\mathcal{E}(\lambda)$ in [39] for the full line case, we introduce our third characteristic determinant, the *Evans function*, by the formula

$$\mathcal{E}^+(\lambda) = \det(Y_0^+(\lambda) + R_-(\lambda)). \quad (17)$$

We call a solution $\mathbf{z}^+ = (z_n^+)_{n \geq -1}$ of the second order difference equation (1) (without the boundary condition) the *Jost solution*, cf. [9], provided

$$(\mu_+)^{-n} z_n^+ - 1 \rightarrow 0 \text{ as } n \rightarrow +\infty, \quad (18)$$

and denote by $\mathbf{z}^r = (z_n^r)_{n \geq -1}$ the *regular* solution of the boundary value problem (1)–(2) satisfying the additional boundary condition $z_0^r = 1$ (which is of course unique but not necessarily bounded at infinity). As in the case of the discrete Schrödinger equations, the Jost solution $\mathbf{z}^+ = \mathbf{z}^+(\lambda)$ is unique. Introducing the notation $\mathcal{W}(\mathbf{u}, \mathbf{v})_n = (-1)^{n-1} (u_{n-1} v_n - u_n v_{n-1})$, $n \geq 0$, for the Wronskian of any two sequences $\mathbf{u} = (u_n)_{n \geq -1}$ and $\mathbf{v} = (v_n)_{n \geq -1}$, we note that the Wronskian of the solutions to (1) is n -independent, and that $\mathcal{W}(\mathbf{z}^+, \mathbf{z}^r)_0 = z_{-1}^+(\lambda)$. We define our fourth characteristic determinant, the *Jost function*, analogously to $\mathcal{F}$ from [39] for the full line case,

$$\mathcal{F}^+(\lambda) = \mu_+(\lambda) \mathcal{W}(\mathbf{z}^+(\lambda), \mathbf{z}^r(\lambda))_0, \quad (19)$$

where the scaling factor $\mu_+(\lambda)$ is chosen such that $\mathcal{F}^+(\lambda) = 1$ provided $b_n c_n = 0$ for $n \geq 0$ when $z_n^+ = \mu_+^n$ for $n \geq -1$.

Our fifth characteristic determinant will be defined only for the difference equation (4) assuming (5) and uses continued fractions; we introduce the notation

$$g_+(\lambda) = \frac{1}{\frac{\lambda}{\rho_1} + \frac{1}{\frac{\lambda}{\rho_2} + \dots}}. \quad (20)$$

The continued fraction converges for $\operatorname{Re}(\lambda) > 0$ and $|\arg(\lambda)| \leq \pi/2 - \delta$ for any $\delta \in (0, \pi/2)$ by the classical Van Vleck Theorem [32, Theorem 4.29] and we may now define our fifth function of interest by the formula, cf. $\mathcal{G}$ from [39],

$$\mathcal{G}^+(\lambda) = \mu_+(\lambda) z_0^+(\lambda) (g_+(\lambda) + \lambda/\rho_0), \quad (21)$$

where $z_0^+(\lambda)$ is the 0-th entry of the Jost solution, $\mu_+(\lambda)$ is the eigenvalue of $A(\lambda)$ inside of the unit disk, and $g_+(\lambda)$ is defined in (20), and we assume that $\operatorname{Re}(\lambda) > 0$.

We are ready to formulate the main results of this paper.

Theorem 1.1. *Assume $\lambda \notin [-2i, 2i]$ and that the sequences in (3) depend on λ holomorphically. The functions introduced in (13), (15), (17) and (19) are holomorphic in λ and equal,*

$$\det(I - K_\lambda^+) = \det(I - \mathcal{T}_\lambda^+) = \mathcal{E}^+(\lambda) = \mathcal{F}^+(\lambda). \quad (22)$$

As a result, λ is a simple discrete eigenvalue of (1)–(2) if and only if λ is a zero of each of the functions in (22).

Corollary 1.2. *Assume $\operatorname{Re}(\lambda) > 0$ and (5), and consider the difference equation (4). Define (b_n) and (c_n) by (6) and use the sequences to construct all four characteristic determinants in (22). Then $\mathcal{G}^+(\lambda)$ defined in (21) via the continued fraction is equal to the functions in (22).*

We refer to [39] for a detailed discussion of the literature related to the results, and mention here only the following. Regarding the Birman-Schwinger operator pencils see [2, 8]. The papers most relevant to the current setup are [12, 23] and [38]. We are not aware of any literature on the Birman-Schwinger type pencils specific for the first order systems (1) on the half-line. For the full line case they have been studied in [12] which is a companion paper to [23] dealing with differential equations where, among many other things, the first equality in (22) was proved for the case of the Schrödinger differential operators. We are not aware of results of this sort for the difference equations of type (1) on the half-line. This equation is of course a particular case of eigenvalue problems for three diagonal (Jacobi) matrices, cf. [29, 31] and also [7, 18, 19, 30, 35, 45]. The dichotomies on the half-line were studied in great detail, see [4, 5, 6] and the references therein. Regarding the Evans function we refer to [34, 44] where one can find many other sources. The most relevant papers are again [12, 23], where, in particular, the second equality in (22) has been proved for quite general first order systems of difference and differential equations but on the full line, and so the results in the current paper for the half-line case seem to be new. The Evans function is a relatively new topic in the study of the 2D-Euler-related difference equations as in (4); we are aware of only [15, 39]. The Jost solutions are classical [9] and so is the equality of $\det(I - K_\lambda^+)$ and $\mathcal{F}^+$ in (22) for the Schrödinger case, see [33, 43] and generalizations in [25, 26]; however,

it is quite possible that the use of the Jost solutions as well as the equality of $\mathcal{F}^+$ and $\mathcal{E}^+$ and $\det(I - K_\lambda^+)$ in the context of (1), see [39], appear to be new for the half-line case. Regarding continued fractions in this context see [14, 21, 32, 42, 46].

We now briefly discuss connections to the Euler equation again referring to [14, 39] for more details and concentrating on just one particularly important case of stability of the so-called unidirectional, or generalized Kolmogorov, flow for the Euler equations of ideal fluid on the two dimensional torus $\mathbb{T}^2$,

$$\partial_t \Omega + U \cdot \nabla \Omega = 0, \operatorname{div} U = 0, \Omega = \operatorname{curl} U, \mathbf{x} = (x_1, x_2) \in \mathbb{T}^2, \quad (23)$$

where the two-dimensional vector $U = U(\mathbf{x})$ is the velocity and the scalar Ω is the vorticity of the fluid. The unidirectional (or generalized Kolmogorov) flow is the steady state solution to the Euler equations on $\mathbb{T}^2$ of the form

$$\Omega^0(\mathbf{x}) = \alpha e^{i\mathbf{p} \cdot \mathbf{x}}/2 + \alpha e^{-i\mathbf{p} \cdot \mathbf{x}}/2 = \alpha \cos(\mathbf{p} \cdot \mathbf{x}), \quad (24)$$

where $\mathbf{p} \in \mathbb{Z}^2 \setminus \{0\}$ is a given vector and $\alpha \in \mathbb{R}$. In particular, $\mathbf{p} = (m, 0) \in \mathbb{Z}^2$ for $m \in \mathbb{N}$ corresponds to the classical Kolmogorov flow. It is thus a classical problem to study (linear) stability of the flow given by (24). To this end, using Fourier series $\Omega(\mathbf{x}) = \sum_{\mathbf{k} \in \mathbb{Z}^2 \setminus \{0\}} \omega_{\mathbf{k}} e^{i\mathbf{k} \cdot \mathbf{x}}$ for vorticity, we rewrite (23) as a system of nonlinear equations for $\omega_{\mathbf{k}}$, $\mathbf{k} \in \mathbb{Z}^2$ as in [14, 36, 39, 40]. Linearizing this system about the unidirectional flow, one obtains the following operator in $\ell^2(\mathbb{Z}^2; \mathbb{C})$,

$$L : (\omega_{\mathbf{k}})_{\mathbf{k} \in \mathbb{Z}^2} \mapsto (\alpha\beta(\mathbf{p}, \mathbf{k} - \mathbf{p})\omega_{\mathbf{k} - \mathbf{p}} - \alpha\beta(\mathbf{p}, \mathbf{k} + \mathbf{p})\omega_{\mathbf{k} + \mathbf{p}})_{\mathbf{k} \in \mathbb{Z}^2}, \quad (25)$$

where the coefficients $\beta(\mathbf{p}, \mathbf{q})$ for $\mathbf{p} = (p_1, p_2)$, $\mathbf{q} = (q_1, q_2) \in \mathbb{Z}^2$ are defined as

$$\beta(\mathbf{p}, \mathbf{q}) = \frac{1}{2}(\|\mathbf{q}\|^{-2} - \|\mathbf{p}\|^{-2})(\mathbf{p} \wedge \mathbf{q}), \quad \text{with } \mathbf{p} \wedge \mathbf{q} := \det \begin{bmatrix} p_1 & q_1 \\ p_2 & q_2 \end{bmatrix}, \quad (26)$$

for $\mathbf{p} \neq 0$, $\mathbf{q} \neq 0$, and $\beta(\mathbf{p}, \mathbf{q}) = 0$ otherwise. The flow (24) is called (linearly) unstable if the operator L has nonimaginary spectrum. We refer to [37, 41] for results on stability and instability for the 2D Euler equations and related models.

To study the spectrum of L , (see the discussion in [14, pp. 2054-2057]; see also [36, 39]) we decompose this operator into a sum of operators $L_{\mathbf{q}}$, $\mathbf{q} \in \mathbb{Z}^2$, acting in the space $\ell^2(\mathbb{Z}; \mathbb{C})$, by “slicing” the grid $\mathbb{Z}^2$ along lines parallel to $\mathbf{p}$ such that $\operatorname{Spec}(L) = \cup_{\mathbf{q}} \operatorname{Spec}(L_{\mathbf{q}})$, where

$$L_{\mathbf{q}} : (w_n) \mapsto (\alpha\beta(\mathbf{p}, \mathbf{q} + (n-1)\mathbf{p})w_{n-1} - \alpha\beta(\mathbf{p}, \mathbf{q} + (n+1)\mathbf{p})w_{n+1}), \quad n \in \mathbb{Z}, \quad (27)$$

and for $\mathbf{k} = \mathbf{q} + n\mathbf{p}$ from (25) we denote $w_n = \omega_{\mathbf{q} + n\mathbf{p}}$. In other words, $L_{\mathbf{q}} = \alpha(S - S^*) \operatorname{diag}_{n \in \mathbb{Z}} \{\beta(\mathbf{p}, \mathbf{q} + n\mathbf{p})\}$. By (26), if $\mathbf{q} \parallel \mathbf{p}$ then $L_{\mathbf{q}} = 0$ and thus we assume throughout that $\mathbf{q} \nparallel \mathbf{p}$. Moreover, since $L_{\mathbf{q}}$ contains a scalar multiple $\alpha \in \mathbb{R}$, with no loss of generality we may rescale this operator or, equivalently, will assume the normalization condition $\alpha(\mathbf{q} \wedge \mathbf{p})\|\mathbf{p}\|^{-2}/2 = 1$.

The operators $L_{\mathbf{q}}$ are classified based on the location of the line $B_{\mathbf{q}} = \{\mathbf{q} + n\mathbf{p} : n \in \mathbb{Z}\}$ through the point $\mathbf{q} \in \mathbb{Z}^2$ relative to the disc of radius $\|\mathbf{p}\|$ centered at zero, see, again, the discussion in [14, pp. 2054-2057]; see also [36, 39]. Spectral properties of the operators drastically depend on the location. For instance, if none of the vectors $\mathbf{q} + n\mathbf{p}$, $n \in \mathbb{Z}$, is located inside the open disc then $L_{\mathbf{q}}$ has no unstable eigenvalues [36, 40]. In the current paper we consider only the case when $\mathbf{p}$ and $\mathbf{q}$ are such that

$$\|\mathbf{q}\| < \|\mathbf{p}\|, \quad \|\mathbf{q} - \mathbf{p}\| = \|\mathbf{p}\| \quad \text{and} \quad \|\mathbf{q} + n\mathbf{p}\| > \|\mathbf{p}\| \quad \text{for all } n \in \mathbb{Z} \setminus \{-1, 0\}; \quad (28)$$

a typical example of this is $\mathbf{p} = (3, 1)$, $\mathbf{q} = (2, -2)$. This assumption corresponds to the case I_- described in [14], and we refer to this paper for a discussion regarding other possible cases. The case I_+ when -1 in (28) is replaced by $+1$ can be treated similarly to I_- while the case I_0 when $\|\mathbf{q} + n\mathbf{p}\| > \|\mathbf{p}\|$ for all $n \neq 0$ has been considered in [39]. Geometrically, the case I_- and I_+ occur when one of the points in $B_{\mathbf{q}}$ is inside of the open disc of radius $\|\mathbf{p}\|$ and one of the points in $B_{\mathbf{q}}$ is on the boundary of the disk while in case I_0 one point of $B_{\mathbf{q}}$ is inside of the open disk while all others are outside of the closed disk. An interesting open question is to describe the spectrum of $L_{\mathbf{q}}$ in the case II when two points of $B_{\mathbf{q}}$ are located inside of the open disk of radius $\|\mathbf{p}\|$ centered at zero.

Assuming that we are in the case I_- , that is, that (28) holds, we introduce the sequence $\rho_n = 1 - \|\mathbf{p}\|^2 \|\mathbf{q} + n\mathbf{p}\|^{-2}$, $n \in \mathbb{Z}$, such that the operator $L_{\mathbf{q}}$ in (27) reads $L_{\mathbf{q}} = (S - S^*) \text{diag}_{n \in \mathbb{Z}} \{\rho_n\}$, see (26). The eigenvalue equation for $L_{\mathbf{q}}$ is

$$\rho_{n-1}w_{n-1} - \rho_{n+1}w_{n+1} = \lambda w_n, \quad n \in \mathbb{Z}, \quad \text{with } (w_n)_{n \in \mathbb{Z}} \in \ell^2(\mathbb{Z}; \mathbb{C}). \quad (29)$$

By (28) we have $\rho_{-1} = 0$, $\rho_0 < 0$, and $\rho_n \in (0, 1)$ for all $n \in \mathbb{Z} \setminus \{0, -1\}$. In particular, the sequence $(\rho_n)_{n \geq 0}$ satisfies (5). Now Theorem 1.1 and Corollary 1.2 can be applied because the solution of (29) must be supported on $\mathbb{Z}_+$ as seen in Lemma 2.5 given in the next section. Theorem 1.1 implies the following.

Corollary 1.3. *Assume that a given $\mathbf{p} \in \mathbb{Z}^2$ is such that there exists a $\mathbf{q} \in \mathbb{Z}^2$ satisfying (28). Then the eigenvalues of $L_{\mathbf{q}}$ with positive real parts are in one-to-one correspondence with zeros of each of the five functions in (22) and (21). Moreover, the operator $L_{\mathbf{q}}$ from (27), and thus L from (25), has a positive eigenvalue. As a result, the unidirectional flow (24) is linearly unstable.*

The instability of the unidirectional flow in the current setting has been established in [14, Theorem 2.9]. Nevertheless, the first assertion in Corollary 1.3 is an improvement of the part of [14, Theorem 2.9] where the correspondence was established between only the *positive* roots of the function $g_+(\lambda) + \lambda/\rho_0$ and the *positive* eigenvalues of $L_{\mathbf{q}}$ but under the additional assumption that the respective eigensequences satisfy some special property [14, Property 2.8]. By applying Theorem 1.1 in the proof of Corollary 1.3 we were able to show that this assumption is redundant.

We conclude this section with references on the literature on stability of unidirectional flows. This topic is quite classical and well-studied, and we refer to [1, 3, 20, 21, 22, 42]. The setup used herein was also used in many papers [16, 17, 36, 40, 46], but the closest to the current work is [14, 39]. We mention also [38] regarding connections to the Birman-Schwinger operators. Finally, connections between the Evans function and the linearization of the 2D Euler equation has been studied in a recent important paper [15].

2. PROOFS

We begin with several general comments regarding the objects introduced in the previous section. For a start, it is sometimes convenient to diagonalize the matrix A from (11). To this end we introduce matrices

$$W = \begin{bmatrix} \mu_+ & \mu_- \\ 1 & 1 \end{bmatrix}, \quad W^{-1} = (\mu_+ - \mu_-)^{-1} \begin{bmatrix} 1 & -\mu_- \\ -1 & \mu_+ \end{bmatrix}, \quad \tilde{A} = \text{diag}\{\mu_+, \mu_-\}, \quad (30)$$

so that one has, cf. (11),

$$W^{-1}AW = \tilde{A}, \quad W^{-1}P_{\pm}W = Q_{\pm}. \quad (31)$$

We will use below the explicit formulas

$$R_+ = \begin{bmatrix} 0 & \mu_+ \\ 0 & 1 \end{bmatrix}, \quad R_- = \begin{bmatrix} 1 & -\mu_+ \\ 0 & 0 \end{bmatrix}, \quad P_+ = (\mu_+ - \mu_-)^{-1} \begin{bmatrix} \mu_+ & 1 \\ 1 & -\mu_- \end{bmatrix}, \quad A^{-1} = \begin{bmatrix} 0 & 1 \\ 1 & \lambda \end{bmatrix}. \quad (32)$$

The operator $S - S^{-1} - \lambda$ is invertible in $\ell^2(\mathbb{Z}_+; \mathbb{C})$ if and only if $\lambda \notin [-2i, 2i]$ and in this case the inverse operator $(S - S^{-1} - \lambda)^{-1}$ is given by the infinite matrix $[a_{nk}]_{n,k=0}^{\infty}$ whose entries are defined by the formula

$$a_{nk} = (\mu_+ - \mu_-)^{-1} \times \begin{cases} (\mu_-)^{n-k} \left(1 - \left(\frac{\mu_+}{\mu_-}\right)^{n+1}\right) & \text{for } n \leq k, \\ (\mu_+)^{n-k} \left(1 - \left(\frac{\mu_+}{\mu_-}\right)^{k+1}\right) & \text{for } n > k. \end{cases} \quad (33)$$

One can check this either directly by multiplying the respective infinite matrices and using the equation $\mu^2 + \lambda\mu - 1 = 0$, or by using [27, Theorem 1.1] as follows: we factorize $S - S^{-1} - \lambda = \mu_- (I + \mu_+ S^{-1})(I + \mu_-^{-1} S)$ and use the Neumann series to obtain the inverse of each factor.

From now on we abbreviate $A = \text{diag}_{n \in \mathbb{Z}_+} \{A\}$. The operator $S^{-1} - A$ acting from $\ell_{\text{ran } Q_+}^2(\mathbb{Z}_+; \mathbb{C}^2)$ into $\ell^2(\mathbb{Z}_+; \mathbb{C}^2)$, cf. (14) and [6], is invertible if and only if $\lambda \notin [-2i, 2i]$ and in this case the inverse operator $(S^{-1} - A)^{-1}$ from $\ell^2(\mathbb{Z}_+; \mathbb{C}^2)$ onto $\ell_{\text{ran } Q_+}^2(\mathbb{Z}_+; \mathbb{C}^2)$ for $\mathbf{u} = (u_n)_{n \geq 0} \in \ell^2(\mathbb{Z}_+; \mathbb{C}^2)$ is given by the formula

$$((S^{-1} - A)^{-1} \mathbf{u})_n = - \sum_{k=n}^{+\infty} A^n R_- A^{-(k+1)} u_k + \sum_{k=0}^{n-1} A^n R_+ A^{-(k+1)} u_k, \quad n \geq 0; \quad (34)$$

here and below we always set $\sum_{k=0}^{-1} = 0$, and so the RHS of (34) for $n = 0$ is a vector from $\text{ran } R_- = \text{ran } Q_+$ as required in (14). Formula (34) can be either taken from [6] or checked directly by multiplying $S^{-1} - A$ and the operator in (34) and taking into account that $\text{ran } R_+ = \text{ran } P_+$ is the set of the initial data for the *bounded* on $\mathbb{Z}_+$ solutions of the difference equation $y_{n+1} = Ay_n$, the projection P_+ is a dichotomy projection for this equation on $\mathbb{Z}_+$, and therefore the projection R_+ onto $\text{ran } P_+$ parallel to $\text{ran } Q_+$ is also a dichotomy projection since $\text{ran } P_+ \oplus \text{ran } Q_+ = \mathbb{C}^2$, cf. [4, 5, 6] and also [10], [11] or [13, Chapter 4] for discussions of dichotomies on the half-line.

Formula (34) shows that the operator $\mathcal{T}_{\lambda}^+$ in (15) is an operator with semi-separable kernel, that is, $\mathcal{T}_{\lambda}^+$ can be written as an infinite matrix $\mathcal{T}_{\lambda}^+ = [T_{nk}]_{n,k=0}^{+\infty}$ so that $(\mathcal{T}_{\lambda}^+ \mathbf{u})_n = \sum_{k=0}^{+\infty} T_{nk} u_k$, where

$$T_{nk} = \begin{cases} -C_n A^n R_- A^{-(k+1)} B_k & \text{for } 0 \leq n \leq k, \\ -C_n A^n R_+ A^{-(k+1)} B_k & \text{for } 0 \leq k \leq n. \end{cases} \quad (35)$$

We refer to [28, Chapter IX] and [24] for a discussion of the operators with semi-separable kernels; the results therein are used below in the proof of the second equality in (22).

The matrix-valued Jost solution $\mathbf{Y}^+ = (Y_n^+)_{n \geq 0}$ of (8) (with no boundary condition) is obtained as a solution to the following Volterra equation,

$$Y_n^+ - A^n R_+ = - \sum_{k=n}^{+\infty} A^{n-(k+1)} B_k C_k Y_k^+, \quad (36)$$

defined first for $n \geq N$ with N large enough and then extended to all of $\mathbb{Z}_+$ as a solution to the difference equation (8) via $Y_n^+ := (A_n^\times)^{-1} Y_{n+1}^+$ for $n = 0, 1, \dots, N-1$. The sequence $(Y_n^+)_{n \geq 0}$ thus defined will satisfy the Volterra equation (36) as the following inductive step shows: Suppose we know that (36) holds for $n = N$ and that $B_{N-1} C_{N-1} Y_{N-1}^+ = Y_N^+ - A Y_{N-1}^+$. We then use this and (36) with $n = N$ in the following calculation yielding (36) for $n = N-1$,

$$\begin{aligned} A^{N-1} R_+ &= \sum_{k=N-1}^{+\infty} A^{(N-1)-(k+1)} B_k C_k Y_k^+ \\ &= A^{-1} (A^N R_+ - \sum_{k=N}^{+\infty} A^{N-(k+1)} B_k C_k Y_k^+ - Y_N^+) + Y_{N-1}^+ = Y_{N-1}^+. \end{aligned}$$

Lemma 2.1. *There is a large enough N such that equation (36) for $n \geq N$ has a unique solution thus yielding the matrix-valued Jost solution $\mathbf{Y}^+ = (Y_n^+)_{n \geq 0}$ of (8) satisfying (16).*

Proof. We recall that $|\mu_+| < 1$ and introduce the space

$$\ell_N^\infty := \{\mathbf{u} = (u_n)_{n \geq N} : \|\mathbf{u}\|_{\ell_N^\infty} := \sup \{|\mu_+|^{-n} \|u_n\|_{\mathbb{C}^2} : n \geq N\} < \infty\}$$

of exponentially decaying at infinity $\mathbb{C}^2$ -valued sequences on $[N, +\infty) \cap \mathbb{Z}_+$. Let T_N denote the operator on ℓ_N^∞ that appears in the RHS of (36),

$$(T_N \mathbf{u})_n = - \sum_{k=n}^{+\infty} A^{n-(k+1)} B_k C_k u_k, \quad \mathbf{u} = (u_n)_{n \geq N} \in \ell_N^\infty. \quad (37)$$

Denoting $q_N = \sum_{n=N}^{+\infty} \|B_k C_k\|_{\mathbb{C}^2 \times 2}$, we have $q_N \rightarrow 0$ as $N \rightarrow \infty$ by (3) and (11). Using (31) we have¹ $\|A^{n-(k+1)}\|_{\mathbb{C}^2 \times 2} \lesssim |\mu_+|^{n-(k+1)}$ for $k \geq n$. Thus, for $n \geq N$,

$$|\mu_+|^{-n} \|u_n\|_{\mathbb{C}^2} \lesssim \sum_{k=n}^{+\infty} |\mu_+|^{-n} |\mu_+|^{n-(k+1)} \|B_k C_k\|_{\mathbb{C}^2 \times 2} \|u_k\|_{\mathbb{C}^2} \lesssim q_N \|\mathbf{u}\|_{\ell_N^\infty} \quad (38)$$

and the norm of T_N in ℓ_N^∞ is dominated by q_N and therefore is less than, say, $1/2$ provided N is large enough. Then $(Y_n^+)_{n \geq N} = (I - T_N)^{-1} (A^n R_+)_{n \geq N}$ is the unique solution of (36) whose columns are in ℓ_N^∞ and so for $n \geq N$ we infer

$$\begin{aligned} |\mu_+|^{-n} \|Y_n^+ - A^n R_+\|_{\mathbb{C}^2 \times 2} &\leq \|(Y_n^+)_{n \geq N} - (A^n R_+)_{n \geq N}\|_{\ell_N^\infty} = \|T_N (Y_n^+)_{n \geq N}\|_{\ell_N^\infty} \\ &\lesssim q_N \|(Y_n^+)_{n \geq N}\|_{\ell_N^\infty} = q_N \|(I - T_N)^{-1} (A^n R_+)_{n \geq N}\|_{\ell_N^\infty} \\ &\lesssim q_N \|(A^n R_+)_{n \geq N}\|_{\ell_N^\infty} \lesssim q_N \end{aligned}$$

because $\|(A^n R_+)_{n \geq N}\|_{\ell_N^\infty} \lesssim 1$ since $\text{ran } R_+ = \text{ran } P_+$. This yields the first assertion in (16) while the second follows by multiplying (36) by R_+ from the right and using the uniqueness of the solution $(Y_n^+)_{n \geq N} = (Y_n^+ R_+)_{n \geq N}$. $\square$

¹We write $a \lesssim b$ if $a \leq cb$ for a constant c independent of any parameters contained in a and b .

Remark 2.2. Assertion $Y_n^+ = Y_n^+ R_+$, cf. (16), and formula (30) for R_+ show that the first column of the matrix Y_n^+ is zero. Since $(Y_n^+)_{n \geq 0}$ solves the difference equation (8), we conclude that there is a sequence $(z_n)_{n \geq -1}$ that solves the difference equation (1) (with no boundary condition) such that

$$Y_n^+ = \begin{bmatrix} 0 & z_n \\ 0 & z_{n-1} \end{bmatrix}, \quad n \geq 0. \quad (39)$$

Next, we proceed to discuss the Jost solution $\mathbf{z}^+ = (z_n^+)_{n \geq -1}$ obtained as a solution to the following scalar Volterra equation,

$$z_n^+ - (\mu_+)^n = -(\mu_+ - \mu_-)^{-1} \sum_{k=n}^{+\infty} b_k c_k ((\mu_+)^{n-k} - (\mu_-)^{n-k}) z_k^+, \quad (40)$$

at first for $n \geq N$ with N sufficiently large. A computation using $(\mu_+)^2 + \lambda \mu_+ - 1 = 0$ shows that z_n^+ from (40) satisfy (1), see [29] for a similar computation. Therefore, the solution $(z_n^+)_{n \geq N}$ to (40) can be propagated backward as solution to (1), and thus z_n^+ can be defined for all $n \geq -1$. We now record properties of the Jost solution.

Remark 2.3. The Jost solution $\mathbf{z}^+(\lambda)$ is unique and holomorphic in λ . This follows by a standard argument, cf. [12, 23] and the references therein, presented in [39, Remark 2.2] and based on passing in (40) to the new unknowns z_n^+/μ_+^n and proving that the RHS of the resulting equation is a strict contraction in the space of exponentially decaying at infinity sequences, see the analogous proof of Lemma 2.1. As in the lemma, this yields the property (18).

We stress that the Jost solution $\mathbf{z}^+ = (z_n^+(\lambda))_{n \geq -1}$ is defined for $n \geq -1$ as the solution of the difference equation (1), with no boundary condition (2). Theorem 1.1 shows, in particular, that $z_{-1}^+(\lambda) = 0$ if and only if λ is an eigenvalue (which is not surprising as in this case $\mathbf{z}^+(\lambda)$ is the exponentially decaying solution to (1) that also satisfies the boundary condition (2)).

Remark 2.4. Let us consider the difference equation (4) which is a particular case of (1) with b_n, c_n as in (6) and ρ_n satisfying (5). Then the proof of [39, Lemma 2.3] applies and shows that the Jost solution satisfies $z_n^+ \neq 0$ for all $n \geq 0$.

Proof of Theorem 1.1. The three equalities in (22) are proved as follows.

1. We prove $\det(I - K_\lambda^+) = \det(I - \mathcal{T}_\lambda^+)$. We claim that K_λ^+ from (13) is the $(1, 1)$ -block of the operator $\mathcal{T}_\lambda^+$ from (15) in the decomposition

$$\ell^2(\mathbb{Z}_+; \mathbb{C}^2) = \text{ran}(\text{diag}\{Q_+\}) \oplus \text{ran}(\text{diag}\{Q_-\}),$$

with $\text{diag} = \text{diag}_{\mathbb{Z}_+}$, for the projections $Q_\pm$ from (11), that is, that

$$\text{diag}\{C_n\}(S^{-1} - A)^{-1} \text{diag}\{B_n\} = \begin{bmatrix} -\text{diag}\{c_n\}(S - S^{-1} - \lambda)^{-1} \text{diag}\{b_n\} & 0 \\ 0 & 0 \end{bmatrix}. \quad (41)$$

Clearly, this implies the required equality.

To begin the proof of the claim, we fix $\mathbf{u} = (u_n)_{n \geq 0} \in \ell^2(\mathbb{Z}_+; \mathbb{C}^2)$ and denote $v_n = (\text{diag}_{n \in \mathbb{Z}_+}\{Q_+\}(S^{-1} - A) \text{diag}_{n \in \mathbb{Z}_+}\{Q_+\}\mathbf{u})_n$. Formulas (11) and (32) show that $AR_+ = \mu_+ R_+$ and $R_- A^{-1} = -\mu_+ R_-$, and thus (34) and (31) imply that v_n

is equal to

$$-\sum_{k=n}^{+\infty} (-\mu_+)^{k+1} Q_+ W \tilde{A}^n W^{-1} R_- Q_+ u_k + \sum_{k=0}^{n-1} \mu_+^n Q_+ R_+ W \tilde{A}^{-(k+1)} W^{-1} Q_+ u_k. \quad (42)$$

We now use formulas (30) and (32) to compute the matrices $Q_+ W$, $W^{-1} R_- Q_+$, $Q_+ R_+ W$, $W^{-1} Q_+$. Plugging this into (42) and using that $\mu_{\pm}$ solve the equation $\mu^2 + \lambda\mu - 1 = 0$, after a tedious computation we conclude that the first component of the vector $v_n = Q_+ v_n \in \mathbb{C}^2$ is given by the formula

$$(\mu_+ - \mu_-)^{-1} \left(\sum_{k=n}^{+\infty} (\mu_-)^{n-k} \left(1 - \left(\frac{\mu_+}{\mu_-}\right)^{n+1}\right) z_k + \sum_{k=0}^{n-1} (\mu_+)^{n-k} \left(1 - \left(\frac{\mu_+}{\mu_-}\right)^{k+1}\right) z_k \right).$$

We now use (33) to recognize that the last expression is $((S - S^{-1} - \lambda)^{-1} \mathbf{z})_n$ where z_n denote the first component of the vector $u_n \in \mathbb{C}^2$, $n \geq 0$. Multiplying by c_n and b_n and recalling (11) yields the required claim (41).

2. We prove $\det(I - \mathcal{T}_{\lambda}^+) = \mathcal{E}^+(\lambda)$ in three steps. The first step is to show that it suffices to prove the equality only for finitely supported $(B_n)_{n \geq 0}$ and $(C_n)_{n \geq 0}$. This is fairly standard and follows, say, as the *claim* in the proof of [12, Theorem 4.6] or the proof of [24, Theorem 4.3]: Indeed, replace B_k, C_k in (37) by $\mathbf{1}_M(k) B_k, \mathbf{1}_M(k) C_k$ where $\mathbf{1}_M$ is the characteristic function of $[0, M]$ equals to one on the segment and to zero outside. As in the proof of Lemma 2.1, we denote the respective operator by $T_N^{(M)}$ and the respective matrix-valued Jost solution by $\mathbf{Y}^{+,M}$. Choosing a large N and even larger M the proof of Lemma 2.1 yields $\|T_N - T_N^{(M)}\|_{\mathcal{B}(\ell_N^{\infty})} \lesssim q_{M+1} \rightarrow 0$ as $M \rightarrow \infty$ for the operator norm on ℓ_N^{∞} . Then $\|Y_0^+ - Y_0^{+,M}\|_{\mathbb{C}^{2 \times 2}} \rightarrow 0$ as in the proof of Lemma 2.1 and therefore $\mathcal{E}^{+,M}(\lambda) \rightarrow \mathcal{E}^+(\lambda)$ as $M \rightarrow \infty$ for the Evans function $\mathcal{E}^{+,M}$ obtained by replacing Y_0^+ in (17) by $Y_0^{+,M}$. Analogously, let $\mathcal{T}_{\lambda}^{+,M}$ denote the operator in (15) obtained by replacing B_k, C_k by $\mathbf{1}_M(k) B_k, \mathbf{1}_M(k) C_k$. Then (3) yields convergence of $\mathcal{T}_{\lambda}^{+,M}$ to $\mathcal{T}_{\lambda}^+$ in the trace class norm and so $\det(I - \mathcal{T}_{\lambda}^{+,M}) \rightarrow \det(I - \mathcal{T}_{\lambda}^+)$ as $M \rightarrow \infty$ completing the first step in the proof. From now on we thus assume that sequences $(B_n)_{n \geq 0}, (C_n)_{n \geq 0}$ are finitely supported.

The second step in the proof is a reduction of the infinite dimensional determinant $\det(I - \mathcal{T}_{\lambda}^+)$ to a finite dimensional one. It follows a well established path, see [23, 24, 26, 28]. Multiplying the operator in the RHS of (34) by $\text{diag}_{n \in \mathbb{Z}_+} \{C_n\}$ and $\text{diag}_{k \in \mathbb{Z}_+} \{B_k\}$ and adding and subtracting $\sum_{k=n}^{+\infty} C_n A^n R_- A^{-(k+1)} B_k u_k$ to the result we arrive at the identity

$$\mathcal{T}_{\lambda}^+ = H_+ + H_2 H_3, \text{ where } H_+ = H_0 + H_1, \quad (43)$$

and we introduce notations

$$\begin{aligned} H_0 &= -\text{diag}_{n \in \mathbb{Z}_+} \{C_n A^{-1} B_n\}, \quad (H_1 \mathbf{u})_n = -\sum_{k=n+1}^{+\infty} C_n A^{n-(k+1)} B_k u_k \\ (H_2 y)_n &= C_n A^n R_+ y, \quad H_3 \mathbf{u} = R_+ \sum_{k=0}^{+\infty} A^{-(k+1)} B_k u_k \end{aligned} \quad (44)$$

for $y \in \mathbb{C}^2$ and $\mathbf{u} = (u_n)_{n \geq 0} \in \ell^2(\mathbb{Z}_+; \mathbb{C}^2)$. Here, the operators H_0 and H_1 in $\ell^2(\mathbb{Z}_+; \mathbb{C}^2)$ are of trace class by (3) while the operators H_2 and H_3 are of rank one as H_2 acts from $\text{ran } R_+ = \text{ran } P_+ \subset \mathbb{C}^2$ into $\ell^2(\mathbb{Z}_+; \mathbb{C}^2)$ while H_3 acts from

$\ell^2(\mathbb{Z}_+; \mathbb{C}^2)$ into $\text{ran } R_+ = \text{ran } P_+ \subset \mathbb{C}^2$. The diagonal operator $I - H_0$ is invertible with $\det(I - H_0)$ being equal to 1 by (10) because

$$\prod_{n=0}^{+\infty} \det(I + C_n A^{-1} B_n) = \prod_{n=0}^{+\infty} \det(I + A^{-1} B_n C_n) = \prod_{n=0}^{+\infty} \det(A^{-1}) \det(A + B_n C_n).$$

The operator H_1 is block-lower-triangular, and thus $I - H_+$ is invertible with $\det(I - H_+) = 1$. Writing $I - \mathcal{T}_\lambda^+ = (I - H_+)(I - (I - H_+)^{-1} H_2 H_3)$ and changing the order of factors in the determinant, we arrive at the identity

$$\det(I_{\ell^2(\mathbb{Z}_+; \mathbb{C}^2)} - \mathcal{T}_\lambda^+) = \det_{\mathbb{C}^{2 \times 2}} (I_{2 \times 2} - H_3(I - H_+)^{-1} H_2) \quad (46)$$

that reduces the computation of the infinite dimensional determinant for $I - \mathcal{T}_\lambda^+$ to the finite dimensional determinant for the operator $H_3(I - H_+)^{-1} H_2$ acting in $\text{ran } R_+$ thus completing the second step in the proof.

The third step relates the RHS of (46) and the solution $(Y_n^+)_{n \geq 0}$ to the Volterra equation (36). Indeed, for any $y \in \mathbb{C}^2$ we let $u_n = C_n Y_n^+ y$ and $\mathbf{u} = (u_n)_{n \geq 0}$. Multiplying (36) from the left by C_n and applying the resulting matrices to y yields

$$u_n = C_n A^n R_+ y - \sum_{k=n}^{+\infty} C_n A^{n-(k+1)} B_k u_k = (H_2 y)_n + (H_+ \mathbf{u})_n, \quad n \in \mathbb{Z}_+,$$

or $\mathbf{u} = (I - H_+)^{-1} H_2 y$. Multiplying by H_3 from the left, by the definition of H_3 ,

$$H_3(I - H_+)^{-1} H_2 y = H_3 \mathbf{u} = R_+ \sum_{k=0}^{+\infty} A^{-(k+1)} B_k u_k.$$

Since y is arbitrary, the respective matrices are equal, that is,

$$H_3(I - H_+)^{-1} H_2 = R_+ \sum_{k=0}^{+\infty} A^{-(k+1)} B_k C_k Y_k^+.$$

Since $Y_n^+ = Y_n^+ R_+$ by (16), the last identity and formula (46) yield

$$\det(I - \mathcal{T}_\lambda^+) = \det_{\mathbb{C}^{2 \times 2}} (I_{2 \times 2} - R_+ \sum_{k=0}^{+\infty} A^{-(k+1)} B_k C_k Y_k^+ R_+). \quad (47)$$

Writing matrices in the block form using the direct sum decomposition $\mathbb{C}^2 = \text{ran } R_+ \oplus \text{ran } R_-$ gives the equality of the determinants of the following two matrices,

$$\begin{aligned} I_{2 \times 2} - \sum_{k=0}^{+\infty} A^{-(k+1)} B_k C_k Y_k^+ R_+ &= \begin{bmatrix} R_+ - R_+ \sum_{k=0}^{+\infty} A^{-(k+1)} B_k C_k Y_k^+ R_+ & 0 \\ -R_- \sum_{k=0}^{+\infty} A^{-(k+1)} B_k C_k Y_k^+ R_+ & R_- \end{bmatrix}, \\ I_{2 \times 2} - R_+ \sum_{k=0}^{+\infty} A^{-(k+1)} B_k C_k Y_k^+ R_+ &= \begin{bmatrix} R_+ - R_+ \sum_{k=0}^{+\infty} A^{-(k+1)} B_k C_k Y_k^+ R_+ & 0 \\ 0 & R_- \end{bmatrix}. \end{aligned}$$

This, (36) with $n = 0$, and (47) show that $\det(I - \mathcal{T}_\lambda^+)$ is equal to the determinant of the matrix

$$I_{2 \times 2} - \sum_{k=0}^{+\infty} A^{-(k+1)} B_k C_k Y_k^+ R_+ = R_+ - \sum_{k=0}^{+\infty} A^{-(k+1)} B_k C_k Y_k^+ + R_- = Y_0^+ + R_-,$$

completing the proof of the equality $\det(I - \mathcal{T}_\lambda^+) = \mathcal{E}^+(\lambda)$ by (17).

3. We prove $\mathcal{E}^+(\lambda) = \mathcal{F}^+(\lambda)$. Using the Jost solution $(z_n^+)_{n \geq -1}$ and the regular solution satisfying $z_0^+ = 1$, we recall that $\mathcal{W}(\mathbf{z}^+, \mathbf{z}^r)_0 = z_{-1}^+$ in (19). By Remark 2.2 the solution $(Y_n^+)_{n \geq 0}$ is of the form (39) with some $(z_n)_{n \geq -1}$. We claim that the solution $(z_n)_{n \geq -1}$ in formula (39) for Y_n^+ satisfies $z_n = \mu_+ z_n^+$ where $(z_n^+)_{n \geq -1}$ is the Jost solution from (40). Assuming the claim, we use formulas (17), (39), (32), and (19) to obtain the desired result,

$$\mathcal{E}^+(\lambda) = \det(Y_0^+ + R_-) = \det \begin{bmatrix} 1 & -\mu_+ + \mu_+ z_0^+ \\ 0 & \mu_+ z_{-1}^+ \end{bmatrix} = \mu_+(\lambda) z_{-1}^+(\lambda) = \mathcal{F}^+(\lambda).$$

It remains to prove the claim $z_n = \mu_+ z_n^+$ in formula (39). Using (30) we multiply equation (36) by W^{-1} from the left and W from the right passing to the new unknowns $\tilde{Y}_n^+$ and using (30) and $(z_n)_{n \geq -1}$ from (39),

$$\tilde{Y}_n^+ = W^{-1} Y_n^+ W = (\mu_+ - \mu_-)^{-1} \begin{bmatrix} z_n - \mu_- z_{n-1} & z_n - \mu_- z_{n-1} \\ -(z_n - \mu_+ z_{n-1}) & -(z_n - \mu_+ z_{n-1}) \end{bmatrix}.$$

Noting that $A^n R_+ = \mu_+^n R_+$ by (11) and (32), and explicitly computing all matrices that appear in the equation for $\tilde{Y}_n^+$, we arrive at the following equations,

$$\begin{aligned} (\mu_+ - \mu_-)^{-1} (z_n - \mu_- z_{n-1}) &= \mu_+^n - (\mu_+ - \mu_-)^{-1} \sum_{k=n}^{+\infty} b_k c_k \mu_+^{n-(k+1)} z_k, \\ (\mu_+ - \mu_-)^{-1} (-z_n + \mu_+ z_{n-1}) &= -(\mu_+ - \mu_-)^{-1} \sum_{k=n}^{+\infty} b_k c_k (-\mu_-^{n-(k+1)}) z_k. \end{aligned}$$

Multiplying the first equation by μ_+ and the second by μ_- and adding the results,

$$z_n = \mu_+^{n+1} - (\mu_+ - \mu_-)^{-1} \sum_{k=n}^{+\infty} b_k c_k (\mu_+^{n-k} - \mu_-^{n-k}) z_k.$$

Divided by μ_+ , this is equation (40) for z_n/μ_+ and the uniqueness of the solution of the equation yields the claim thus completing the proof of (22) in the theorem.

4. To prove the last assertion in the theorem, we rely on the Birman-Schwinger principle saying that $\lambda \in \text{Spec}(S - S^{-1} + \text{diag}_{n \in \mathbb{Z}_+} \{b_n c_n\})$ if and only if λ is a zero of $\det(I - K_\lambda^+)$ with the same multiplicity which follows from the fact that the operator $S - S^{-1} + \text{diag}_{n \in \mathbb{Z}_+} \{b_n c_n\} - \lambda$ can be written as the product

$$(S - S^{-1} - \lambda)(I_{\ell^2(\mathbb{Z}_+; \mathbb{C})} - (S - S^{-1} - \lambda)^{-1} \text{diag}_{n \in \mathbb{Z}_+} \{b_n\} \times \text{diag}_{n \in \mathbb{Z}_+} \{c_n\})$$

and the standard property $\det(I - K_1 \times K_2) = \det(I - K_2 \times K_1)$ of the operator determinants. That the eigenvalues λ are simple follows from the fact that $\mathbf{z}$ is an eigensequence if and only if $\mathbf{z}$ is proportional to $\mathbf{z}^+(\lambda)$ with λ satisfying $z_{-1}^+(\lambda) = 0$ (in other words, the dichotomy projection for (8) has rank one). $\square$

Proof of Corollary 1.2. We follow [39] and show that $\mathcal{F}^+(\lambda) = \mathcal{G}^+(\lambda)$ for $\mathcal{F}^+$ from (19) and $\mathcal{G}^+$ from (21). Here, we assume that $\text{Re}(\lambda) > 0$. The function $\mathcal{F}^+$ is holomorphic in λ by Remark 2.3. The convergent continued fraction in (20) is holomorphic in λ by the classical Stieltjes-Vitali Theorem [32, Theorem 4.30]. Thus, it is enough to show the desired equality only for $\lambda > 0$ which we assume from now on. We recall that $z_n^+ \neq 0$ for $n \geq 0$ by Remark 2.4, introduce $v_n^\pm = z_{n-1}^\pm / z_n^\pm$ and re-write equation (4) for $\mathbf{z}^\pm$ as $v_n^+ = \frac{\lambda}{\rho_n} + \frac{1}{v_{n+1}^+}$ for $n \geq 0$. The last formula iterated forward produces the continued fraction (20). Because the

continued fraction converges and $\lambda > 0$, an argument from [14, pp.2063] involving monotonicity of the sequences formed by the odd and even truncated continued fractions yields $v_0^+ = g_+(\lambda) + \lambda/\rho_0$. Therefore,

$$\mathcal{W}(\mathbf{z}^+, \mathbf{z}^r)_0 = z_{-1}^+ = z_0^+ v_0^+ = z_0^+ (g_+(\lambda) + \lambda/\rho_0),$$

and the desired equality of $\mathcal{F}^+$ and $\mathcal{G}^+$ follows by (19) and (21). $\square$

Lemma 2.5. *Assume (28) and $\lambda > 0$. Then the solutions of (29) on $\mathbb{Z}$ are in one-to-one correspondence with the $\ell^2(\mathbb{Z}_+; \mathbb{C})$ -solutions to the eigenvalue problem (4) given by $z_n = \rho_n w_n$ for $n \geq -1$ and $w_n = 0$ for $n \leq -2$.*

Proof. Suppose that $(w_n)_{n \in \mathbb{Z}}$ solves (29) and let $z_n = \rho_n w_n$ for all $n \in \mathbb{Z}$. Then $z_{-1} = 0$ since $\rho_{-1} = 0$ and the sequence $(z_n)_{n \leq -1}$ solves the equation

$$z_{n-1} - z_{n+1} = \lambda z_n / \rho_n \text{ for } n \leq -2 \text{ and } z_{-1} = 0. \quad (48)$$

If $z_{-2} \neq 0$ then the sequence $(z_n)_{n \leq -2}$ due to $\lambda > 0$ exponentially grows as $n \rightarrow -\infty$ as shown in the proof of [39, Lemma 2.3] contradicting $(w_n)_{n \in \mathbb{Z}} \in \ell^2(\mathbb{Z}; \mathbb{C})$ in (29). So one has $z_{-2} = 0$, $z_{-1} = 0$ and, as a result, $z_n = 0$ for all $n \leq -2$. This, in turn, implies that $w_n = 0$ for all $n \leq -2$.

Conversely, suppose that $(z_n)_{n \geq 0} \in \ell_2(\mathbb{Z}_+)$, $z_{-1} = 0$ solves (4) and let $w_n = z_n / \rho_n$ for $n \geq 0$, $w_{-1} = -z_0 / \lambda$, and $w_n = 0$ for $n \leq -2$. Then $(w_n)_{n \in \mathbb{Z}}$ so defined solves (29). $\square$

Proof of Corollary 1.3. We follow the proof of [39, Corollary 1.3]. Due to Lemma 2.5, Theorem 1.1 yields the first assertion in the corollary. To finish the proof it suffices to show that there is a positive root of the function $\mathcal{G}^+$ defined in (21), equivalently, that for some $\lambda > 0$ and g_+ defined in (20) one has $-\lambda/\rho_0 = g_+(\lambda)$. We recall that $\rho_0 < 0$ and $g_+(\lambda) > 0$ for $\lambda > 0$ because condition (5) holds. The proof is completed by using the relations $\lim_{\lambda \rightarrow 0^+} g_+(\lambda) = 1$ and $\lim_{\lambda \rightarrow +\infty} g_+(\lambda) = 0$ established in [14, Lemma 2.10(4)]. $\square$

REFERENCES

- [1] M. Beck and C. E. Wayne, *Metastability and rapid convergence to quasi-stationary bar states for the two-dimensional Navier-Stokes equations*. Proc. Royal Soc. Edinburgh Sect. A - Math. **143** (2013), 905–927.
- [2] J. Behrndt, A. F. M. ter Elst, and F. Gesztesy, *The generalized Birman-Schwinger principle*. Trans. Amer. Math. Soc. **375** (2022), no. 2, 799–845.
- [3] L. Belenkaya, S. Friedlander and V. Yudovich, *The unstable spectrum of oscillating shear flows*. SIAM J. App. Math. **59** (5) (1999), 1701–1715.
- [4] A. Ben-Artzi and I. Gohberg, *Dichotomy of systems and invertibility of linear ordinary differential operators*, Oper. Theory Adv. Appl. **56** (1992), 90–119.
- [5] A. Ben-Artzi, I. Gohberg, and M. A. Kaashoek, *Invertibility and dichotomy of differential operators on a half line*, J. Dynam. Diff. Eqns. **5** (1993), 1–36.
- [6] A. Ben-Artzi, I. Gohberg and M. A. Kaashoek, *Invertibility and dichotomy of singular difference equations*, Operator Theory: Advances and Applications **48** (1990), 157–184.
- [7] S. Bögli and F. Stampach, *On Lieb-Thirring inequalities for one-dimensional non-self-adjoint Jacobi and Schrödinger operators*. J. Spectr. Theory **11** (2021) 1391–1413.
- [8] A. Carey, F. Gesztesy, G. Levitina, R. Nichols, F. Sukochev, D. Zanin, *The limiting absorption principle for massless Dirac operators, properties of spectral shift functions, and an application to the Witten index of non-Fredholm operators*. Mem. European Math. Soc. **4**. European Math. Soc., Berlin, 2023, viii+213 pp.
- [9] K. Chadan and P. C. Sabatier, *Inverse Problems in Quantum Scattering Theory*, 2nd ed., Springer, New York, 1989.

- [10] C. Chicone and Y. Latushkin, *Evolution Semigroups in Dynamical Systems and Differential Equations*, Math. Surv. Monogr. **70**, AMS, Providence, 1999.
- [11] W. A. Coppel, *Dichotomies in stability theory*, Lect. Notes Math., vol. 629, Springer-Verlag, New York, 1978.
- [12] D. Cramer and Y. Latushkin, *Fredholm determinants and the Evans function for difference equations*. Perspectives in operator theory, pp. 111–135, Banach Center Publ., **75**, Polish Acad. Sci. Inst. Math., Warsaw, 2007.
- [13] J. Daleckiĭ and M. Krein, *Stability of differential equations in Banach space*, Amer. Math. Soc., Providence, RI, 1974.
- [14] H. Dullin, Y. Latushkin, R. Marangell, S. Vasudevan and J. Worthington, *Instability of the unidirectional flows for the 2D α -Euler equations*. Comm. Pure. Appl. Analysis., **19** (4) (2020), 2051–2079.
- [15] H. Dullin and R. Marangell, *An Evans function for the linearised 2D Euler equations using Hill's determinant*. Phys. D, **457** (2024), Paper No. 133954, 14 pp.
- [16] H. R. Dullin, R. Marangell and J. Worthington, *Instability of equilibria for the 2D Euler equations on the torus*. SIAM J. Appl. Math. **76** (4) (2016), 1446–1470.
- [17] H. R. Dullin and J. Worthington, *Stability results for idealized shear flows on a rectangular periodic domain*. J. Math. Fluid Mech. **20** (2) (2018), 473–484.
- [18] I. Egorova and L. Golinskii, *On the location of the discrete spectrum for complex Jacobi matrices*. Proc. Amer. Math. Soc. **133** (2005) 3635–3641.
- [19] R. L. Frank, A. Laptev and T. Weidl, *Schrödinger Operators: Eigenvalues and Lieb-Thirring Inequalities*, vol. **200** of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2023.
- [20] A. L. Frenkel and X. Zhang, *Large-scale instability of generalized oscillating Kolmogorov flows*. SIAM Journal on Applied Mathematics. **58** (2) (1998), 540–564.
- [21] S. Friedlander and L. Howard, *Instability in parallel flows revisited*. Studies Appl. Math. **101** (1) (1998), 1–21.
- [22] S. Friedlander, M. Vishik and V. Yudovich, *Unstable eigenvalues associated with inviscid fluid flows*. J. Math. Fluid. Mech. **2** (4) (2000), 365–380.
- [23] F. Gesztesy, Y. Latushkin and K. Makarov, *Evans functions, Jost functions, and Fredholm determinants*. Arch. Rational Mech. Anal. **186** (2007), 361–421.
- [24] F. Gesztesy and K. Makarov, *(Modified) Fredholm determinants for operators with matrix-valued semi-separable integral kernels revisited*. Integr. Equ. Oper. Theory **48** (2004), 561–602.
- [25] F. Gesztesy, M. Mitrea and M. Zinchenko, *Variations on a theme of Jost and Pais*. J. Funct. Anal. **253** (2007), no. 2, 399–448.
- [26] F. Gesztesy and R. Nichols, *A Jost-Pais-type reduction of (modified) Fredholm determinants for semi-separable operators in infinite dimensions*. Recent advances in inverse scattering, Schur analysis and stochastic processes, 287–314, Oper. Theory Adv. Appl. **244**, Linear Oper. Linear Syst., Birkhäuser/Springer, Cham, 2015.
- [27] I. C. Gohberg and I. A. Feldman, *Convolution Equations and Projection Methods for Their Solution*, Transl. Math. Monogr. **41**, AMS, Providence, RI, 1974.
- [28] I. Gohberg, S. Goldberg and M. A. Kaashoek, *Classes of Linear Operators. Vol. I*. Birkhauser Verlag, Basel, 1990.
- [29] L. Golinskii, *Volterra-type discrete integral equations and spectra of non-self-adjoint Jacobi operators*. Integr. Equ. Oper. Theory. **93** (2021).
- [30] L. Golinskii, *Perturbation determinants and discrete spectra of semi-infinite non-self-adjoint Jacobi operators*. J. Spectr. Theory **12** (2022) 835–856.
- [31] J. Janas, and S. Naboko, *On the point spectrum of some Jacobi matrices*. J. Operator Theory **40** (1998), no. 1, 113–132.
- [32] W.B. Jones and W.J. Thron, *Continued Fractions: Analytic Theory and Applications*. Cambridge University Press, 1984.
- [33] R. Jost and A. Pais, *On the scattering of a particle by a static potential*, Phys. Rev. **82**, 840–851 (1951).
- [34] T. Kapitula and K. Promislow, *Spectral and Dynamical Stability of Nonlinear Waves*, Appl. Math. Sci. **185**, Springer, New York, 2013.

- [35] D. Krejcirik, A. Laptev and F. Stampach, *Spectral enclosures and stability for non-self-adjoint discrete Schrödinger operators on the half-line*. Bull. Lond. Math. Soc. **54** (2022), 2379–2403.
- [36] Y. Latushkin, Y. C. Li and M. Stanislavova, *The spectrum of a linearized 2D Euler operator*. Studies Appl. Math. **112** (2004), 259–270.
- [37] Y. Latushkin and S. Vasudevan, *Stability criteria for the 2D α -Euler equations*. J. Math. Anal. Appl., **472** (2) (2019), 1631–1659.
- [38] Y. Latushkin and S. Vasudevan, *Eigenvalues of the linearized 2D Euler equations via Birman-Schwinger and Lin's operators*. J. Math. Fluid. Mech. **20** (4) (2018), 1667–1680.
- [39] Y. Latushkin and S. Vasudevan, *Fredholm determinants, continued fractions, Jost and Evans functions for a Jacobi matrix associated with the 2D-Euler equations*, In: S. Rogosin (ed.) *Analysis without borders: dedicated to Ilya Spitkovsky in occasion of his 70th anniversary*, Operator Theory Adv. Appl. **297** Springer Nature, Switzerland (2024), to appear.
- [40] Y. (Charles) Li, *On 2D Euler equations. I. On the energy-Casimir stabilities and the spectra for linearized 2D Euler equations*, J. Math. Phys. **41** (2000), 728–758.
- [41] Zhiwu Lin, *Some stability and instability criteria for ideal plane flows*. Comm. Math. Phys. **246** (2004), 87–112.
- [42] L. D. Meshalkin and Ia. G. Sinai, *Investigation of the stability of a stationary solution of a system of equations for the plane movement of an incompressible viscous liquid*. J. Appl. Math. Mech. **25** (1961), 1700–1705.
- [43] R. G. Newton, *Relation between the three-dimensional Fredholm determinant and the Jost function*, J. Math. Phys. **13** (1972), 880–883.
- [44] B. Sandstede, *Stability of travelling waves*, in *Handbook of dynamical systems*, Vol. 2, B. Hasselblatt and A. Katok (eds.), North-Holland, Elsevier, Amsterdam, 2002, pp. 983–1055.
- [45] F. Stampach and P. Stovicek, *The characteristic function for Jacobi matrices with applications*. Linear Algebra Appl. **438** (2013) 4130–4155.
- [46] S. Vasudevan, *Instability of Unidirectional Flows for the 2D Navier–Stokes Equations and Related α -Models*. Journal of Math. Fluid Mechanics, **23** (2021), 1 – 29.

UNIVERSITY OF MISSOURI, COLUMBIA, MO 65211, USA

Email address: latushkiny@missouri.edu

KREA UNIVERSITY, SRI CITY, ANDHRA PRADESH, 517646, INDIA

Email address: shibi.vasudevan@krea.edu.in