

CHARACTERIZING LIPSCHITZ IMAGES OF INJECTIVE METRIC SPACES

TARAS BANAKH, JUDYTA BĄK, JOANNA GARBULIŃSKA-WĘGRZYN,
MAGDALENA NOWAK, MICHAŁ POPŁAWSKI

ABSTRACT. A metric space X is *injective* if every non-expanding map $f : B \rightarrow X$ defined on a subspace B of a metric space A can be extended to a non-expanding map $\bar{f} : A \rightarrow X$. We prove that a metric space X is a Lipschitz image of an injective metric space if and only if X is *Lipschitz connected* in the sense that for every points $x, y \in X$, there exists a Lipschitz map $f : [0, 1] \rightarrow X$ such that $f(0) = x$ and $f(1) = y$. In this case the metric space X carries a well-defined intrinsic metric. A metric space X is a Lipschitz image of a compact injective metric space if and only if X is compact, Lipschitz connected and its intrinsic metric is totally bounded. A metric space X is a Lipschitz image of a separable injective metric space if and only if X is a Lipschitz image of the Urysohn universal metric space if and only if X is analytic, Lipschitz connected and its intrinsic metric is separable.

1. INTRODUCTION

By the classical Hahn–Mazurkiewicz Theorem [11, 16], a Hausdorff space X is a continuous image of the unit segment $[0, 1]$ if and only if X is a Peano continuum, i.e., X is compact, metrizable, connected and locally connected. A metric counterpart of the Hahn–Mazurkiewicz Theorem is due to Ważewski [19] and Eilenberg, Harrold [7] who proved that a metric space X is the image of the closed interval $[0, 1]$ under a Lipschitz map if and only if X has finite length. The unit interval and the real line are the most important examples of injective metric spaces. So, it is natural to ask which metric spaces are Lipschitz images of injective metric spaces. In this paper we address this problem and characterize metric spaces which are Lipschitz images of (separable or totally bounded) injective metric spaces.

For a metric space X , its metric will be denoted by d_X . For a map $f : X \rightarrow Y$, its *Lipschitz constant* is defined as a (finite or infinite) number

$$\text{Lip}(f) := \sup_{\substack{x, x' \in X \\ x \neq x'}} \frac{d_Y(f(x), f(x'))}{d_X(x, x')} \in [0, \infty].$$

In this definition we assume that $\sup \emptyset = 0$.

A map $f : X \rightarrow Y$ between metric spaces X, Y is called

- *Lipschitz* if $\text{Lip}(f) < \infty$;
- *L-Lipschitz* for a real constant L , if $\text{Lip}(f) \leq L$;
- *non-expanding* if it is 1-Lipschitz.

A metric space X is *Lipschitz connected* if for every points $x, y \in X$ there exists a Lipschitz map $f : [0, 1] \rightarrow X$ such that $f(0) = x$ and $f(1) = y$. Such a map f is called a *Lipschitz path* connecting the points x, y . It is clear that every Lipschitz connected metric space is a path connected topological space. Every Lipschitz connected metric space X carries the *intrinsic metric* $d_{IX} : X \times X \rightarrow \mathbb{R}$ defined by

$$d_{IX}(x, y) := \inf\{\text{Lip}(f) : f \in X^{[0,1]} \wedge f(0) = x \wedge f(1) = y\}.$$

for any points $x, y \in X$. In the above formula, $X^{[0,1]}$ stands for the set of all functions from $[0, 1]$ to X .

Date: May 6, 2024.

2010 Mathematics Subject Classification. Primary: 54E35, 54E40; Secondary: 51F30, 54C55, 54E45, 54E50, 54F15.

Key words and phrases. Injective metric space, Lipschitz map, Lipschitz image, Urysohn metric space, Lipschitz connected metric space.

A metric space X is called *injective* if every 1-Lipschitz map $f : B \rightarrow X$ defined on a subspace B of a metric space A extends to a 1-Lipschitz map $\tilde{f} : A \rightarrow X$.

Injective metric spaces form a well-known and well-studied class of complete metric spaces, [4], [10, p. 47]. Such spaces are known under two alternative names: hyperconvex metric spaces [1, 9, 14] and 1-Lipschitz Absolute Retracts [5, Ch.I.1.]. The following (known) characterization of injective metric spaces can be deduced from [1, 5].

Theorem 1.1. *For a metric space X , the following conditions are equivalent:*

- (1) X is an injective metric space;
- (2) X is 1-Lipschitz Absolute Retract, which means that for every metric space Y containing X , there exists a 1-Lipschitz map $f : Y \rightarrow X$ such that $f(x) = x$ for all $x \in X$;
- (3) X is hyperconvex in the sense that for any indexed family of closed balls $(B[x_i, r_i])_{i \in I}$ in X with $d_X(x_i, x_j) \leq r_i + r_j$ for all $i, j \in I$, the intersection $\bigcap_{i \in I} B[x_i, r_i]$ is not empty.
- (4) X is metrically convex (for every $a, b \in X$ and $t \in [0, 1]$ there exists $x \in X$ such that $d_X(a, x) = t \cdot d_X(a, b)$ and $d_X(x, b) = (1 - t) \cdot d_X(a, b)$) and has a binary intersection property (every collection of mutually intersecting closed balls in X has a common point).
- (5) for every metric space Y containing X and for every metric space Z , every Lipschitz map $X \rightarrow Z$ has an extension $Y \rightarrow Z$ with the same Lipschitz constant.

Proof. The implications (5) $\Leftrightarrow$ (1) $\Leftrightarrow$ (2) are proved in [5, Proposition 1.2]; (3) $\Rightarrow$ (2) in [1, Theorem 8], and (2) $\Leftrightarrow$ (4) in [5, Proposition 1.4]. To see that (4) $\Rightarrow$ (3), take any indexed family $(B[x_i, r_i])_{i \in I}$ of closed balls in X such that $d_X(x_i, x_j) \leq r_i + r_j$ for every $i, j \in I$. The metric convexity of X ensures that for any distinct indices $i, j \in I$, the closed balls $B[x_i, r_i]$ and $B[x_j, r_j]$ have a common point. The binary intersection property guarantees that $\bigcap_{i \in I} B[x_i, r_i] \neq \emptyset$. $\square$

Another important source of injective metric spaces are complete $\mathbb{R}$ -trees, see [6, 14]. A metric space X is called an $\mathbb{R}$ -tree if for every points $x, y \in X$ the *metric segment*

$$[x, y] := \{z \in X : d_X(x, y) = d_X(x, z) + d_X(z, y)\}$$

is isometric to the interval $[0, d_X(x, y)]$ on the real line, and for every points $x, y, z \in X$ with $[x, y] \cap [y, z] = \{y\}$ we have $d_X(x, z) = d_X(x, y) + d_X(y, z)$. By [14], every complete $\mathbb{R}$ -tree is an injective metric space. Nevertheless, classes of $\mathbb{R}$ -trees and injective metric spaces are distinct. For instance $(0, 1)$ (with the euclidean metric) is $\mathbb{R}$ -tree, which is not injective, since injective metric spaces are complete. On the other hand, $[0, 1]^2$ with the maximum metric is an injective metric space, which is not an $\mathbb{R}$ -tree (the metric segment $[(0, 0), (1, 0)]$ is not isometric to $[0, 1]$).

A metric space Y is called a *Lipschitz image* (resp. *1-Lipschitz image*) of a metric space X if there exists a surjective Lipschitz map $f : X \rightarrow Y$ (with $\text{Lip}(f) \leq 1$).

One can easily reformulate Hahn-Mazurkiewicz theorem [11, 16] to the following version: a Hausdorff space X is a continuous image of the unit segment $[0, 1]$ if and only if X is a continuous image of some compact injective metric space Y . However, the Ważewski result in [19] implies that the class of Lipschitz images of the unit interval $[0, 1]$ is strictly smaller than the class of Lipschitz images of compact injective metric spaces: a counterexample is the injective metric space $[0, 1]^2$ which is not the Lipschitz image of the unit segment $[0, 1]$.

The following three characterizations proved in Sections 5, 6, 7 are the main results of this paper.

Theorem 1.2. *For a metric space X , the following conditions are equivalent:*

- 1) X is a Lipschitz image of an injective metric space;
- 2) X is a Lipschitz image of a complete $\mathbb{R}$ -tree;
- 3) X is Lipschitz connected.

Theorem 1.3. *For a metric space X , the following conditions are equivalent:*

- 1) X is a Lipschitz image of a totally bounded injective metric space;
- 2) X is a Lipschitz image of a compact $\mathbb{R}$ -tree;
- 3) X is compact, Lipschitz connected, and its intrinsic metric is totally bounded.

Theorem 1.4. *For a metric space X , the following conditions are equivalent:*

- 1) X is a Lipschitz image of a separable injective metric space;
- 2) X is a Lipschitz image of a separable complete $\mathbb{R}$ -tree;
- 3) X is a 1-Lipschitz image of the Urysohn universal space;
- 4) X is analytic, Lipschitz connected, and its intrinsic metric is separable.

Let us recall that a topological space X is *analytic* if it is a continuous image of a separable complete metric space. The *Urysohn universal metric space* is a complete separable metric space $\mathbb{U}$ such that any isometric embedding of a subset A of a finite metric space B admits an extension to an isometric embedding of B in $\mathbb{U}$. It is well-known [18] that a Urysohn universal metric space exists and is unique up to an isometry.

2. $\mathbb{R}$ -CONVEX METRIC SPACES AND $\mathbb{R}$ -TREES

Definition 2.1. A metric space X is called *$\mathbb{R}$ -convex* if for every points $x, z \in X$, the metric segment

$$[x, z] := \{y \in X : d_X(x, y) + d_X(y, z) = d_X(x, z)\}$$

is isometric to the segment $[0, d_X(x, z)]$ of the real line.

Proposition 2.2. *For a metric space X , the following conditions are equivalent:*

- 1) X is $\mathbb{R}$ -convex;
- 2) for every points $x, y \in X$, there exists a unique set $I \subseteq X$ such that $x, y \in I$ and I is isometric to the segment $[0, d_X(x, y)]$ on the real line;
- 3) for every points $x, y \in X$ there exists a unique 1-Lipschitz map $f : [0, d_X(x, y)] \rightarrow X$ such that $f(0) = x$ and $f(d_X(x, y)) = y$.

Proof. The implications (1) $\Rightarrow$ (2) $\Rightarrow$ (3) are obvious. To prove that (3) $\Rightarrow$ (1), assume that a metric space X satisfies the condition (3). Given any points $x, y \in X$, we need to show that the metric segment $[x, y]$ is isometric to the interval $[0, d_X(x, y)]$ in the real line. By the condition (3), there exists a unique 1-Lipschitz map $f : [0, d_X(x, y)] \rightarrow X$ such that $f(0) = x$ and $f(d_X(x, y)) = y$. The triangle inequality implies that the image $I := f([0, d_X(x, y)])$ is a subset of the metric segment $[x, y]$. Assuming that $[x, y] \neq I$, we can find a point $z \in [x, y] \setminus I$. Then $d_X(x, y) = d_X(x, z) + d_X(z, y)$. By condition (3), there exist 1-Lipschitz maps $\alpha : [0, d_X(x, z)] \rightarrow X$ and $\beta : [0, d_X(z, y)] \rightarrow X$ such that $\alpha(0) = x$, $\alpha(d_X(x, z)) = z = \beta(0)$ and $\beta(d_X(z, y)) = y$. The 1-Lipschitz maps α, β compose a 1-Lipschitz map $\gamma : [0, d_X(x, y)] \rightarrow X$ defined by

$$\gamma(t) = \begin{cases} \alpha(t) & \text{if } t \in [0, d_X(x, z)]; \\ \beta(t - d_X(x, z)) & \text{if } t \in [d_X(x, z), d_X(x, y)]. \end{cases}$$

Since $\gamma(0) = x = f(0)$, $\gamma(d_X(x, y)) = y = f(d_X(x, y))$, and $\gamma \neq f$, we obtain a contradiction with the uniqueness of the 1-Lipschitz path f . This contradiction shows that $[x, y] = I$. The triangle inequality implies that the 1-Lipschitz map $f : [0, d_X(x, y)] \rightarrow [x, y]$ is an isometry, witnessing that the metric space X is $\mathbb{R}$ -convex. Indeed, for every u, z such that $0 \leq u \leq z \leq d_X(x, y)$ we have $|u - z| \geq d_X(f(u), f(z)) \geq d_X(x, y) - d_X(x, f(u)) - d_X(f(z), y) \geq d_X(x, y) - |0 - u| - |z - d_X(x, y)| = |u - z|$. $\square$

Proposition 2.3. *If a metric space X is $\mathbb{R}$ -convex, then for every points $x, y, z \in X$, the intersection $[x, y] \cap [x, z]$ is equal to the metric segment $[x, u]$ for a unique point $u \in [x, y] \cap [x, z]$.*

Proof. Since the metric space X is $\mathbb{R}$ -convex, there exist isometries $\alpha : [0, d_X(x, y)] \rightarrow [x, y]$ and $\beta : [0, d_X(x, z)] \rightarrow [x, z]$ such that $\alpha(0) = x = \beta(0)$, $\alpha(d_X(x, y)) = y$, and $\beta(d_X(x, z)) = z$. By the compactness of $[x, z]$, the closed set $\alpha^{-1}[x, z]$ in $[0, d_X(x, y)]$ has the largest element s . Since α is an isometry, $s = d_X(\alpha(0), \alpha(s)) = d_X(x, \alpha(s))$. Let $u = \alpha(s)$ and $t = \beta^{-1}(u)$. Since β is an isometry, $t = d_X(\beta(0), \beta(t)) = d_X(x, u) = s$. Observe that $\alpha' := \alpha|_{[0, s]} : [0, s] \rightarrow X$ and $\beta' := \beta|_{[0, t]} : [0, t] \rightarrow X$ are two 1-Lipschitz maps from the interval $[0, s] = [0, t]$ to X such that $\alpha'(0) = x = \beta'(0)$

and $\alpha'(s) = u = \beta'(t)$. Proposition 2.2 ensures that $\alpha' = \beta'$ and hence $[x, y] \cap [x, z] = \alpha[[0, s]] = [x, u]$. The uniqueness of the point u implies from the uniqueness of the end-points of the segment $[x, u] = [x, y] \cap [x, z]$. $\square$

We observe that a metric space X is $\mathbb{R}$ -tree if X is an $\mathbb{R}$ -convex metric space such that for every points $x, y, z \in X$ with $[x, y] \cap [y, z] = \{y\}$, we have $d_X(x, z) = d_X(x, y) + d_X(y, z)$.

The following important fact can be found in [14].

Proposition 2.4. *Every complete $\mathbb{R}$ -tree is an injective metric space.*

3. LIPSCHITZ CONNECTED METRIC SPACES

In this section we study Lipschitz connected and L -Lipschitz connected metric spaces in more details.

Definition 3.1. Let L be a real number. A metric space X is called *L -Lipschitz connected* if for every points $x, y \in X$, there exists an L -Lipschitz path $f : [0, d_X(x, y)] \rightarrow X$ such that $f(0) = x$ and $f(d_X(x, y)) = y$.

If an L -Lipschitz connected metric space contains more than one point, then $L \geq 1$. Therefore, the 1-Lipschitz connectedness is the strongest notion among the L -Lipschitz connectedness properties.

Example 3.2. Every injective metric space X is 1-Lipschitz connected.

Proof. Given any point $x, y \in X$, consider the doubleton $\{0, d_X(x, y)\}$ in the real line and the 1-Lipschitz map $f : \{0, d_X(x, y)\} \rightarrow X$ defined by $f(0) = x$ and $f(d_X(x, y)) = y$. By the injectivity of X , this 1-Lipschitz map admits a 1-Lipschitz extension $\bar{f} : [0, d_X(x, y)] \rightarrow X$, which is a 1-Lipschitz path witnessing that X is 1-Lipschitz connected. $\square$

As we have mentioned in the introduction, every Lipschitz connected metric space X carries the intrinsic metric $d_{IX} : X \times X \rightarrow \mathbb{R}$ defined by

$$\begin{aligned} d_{IX}(x, y) &= \inf\{\text{Lip}(f) : f \in X^{[0,1]} \wedge f(0) = x \wedge f(1) = y\} \\ &= \inf\{C \in [0, \infty) : \exists f \in X^{[0,C]} (f(0) = x \wedge f(C) = y \wedge \text{Lip}(f) \leq 1)\} \end{aligned}$$

for any points $x, y \in X$. The second equality in the definition of the intrinsic metric holds because a function $f \in X^{[0,1]}$ is L -Lipschitz if and only if the function $f_L : [0, L] \rightarrow X$, $f_L : t \mapsto f(t/L)$, is 1-Lipschitz.

The definition of the intrinsic metric implies that $d_X \leq d_{IX}$ for every Lipschitz connected metric space X . Also any 1-Lipschitz map $f : X \rightarrow Y$ between Lipschitz connected metric spaces remains 1-Lipschitz with respect to the intrinsic metrics on X and Y , which means that

$$d_{IY}(f(x), f(x')) \leq d_{IX}(x, x')$$

for every points $x, x' \in X$.

The following lemma describes an interplay between the completions of a Lipschitz connected metric space with respect to the intrinsic metric and the original metric.

Lemma 3.3. *Let X be a Lipschitz connected metric space, $\overline{X}$ be the completion of X , $\overline{X}_I$ be the completion of the metric space $X_I := (X, d_{IX})$, and $\overline{\text{id}} : \overline{X}_I \rightarrow \overline{X}$ be the unique continuous extension of the identity map $\text{id} : X_I \rightarrow X$. Then $\overline{\text{id}}^{-1}[X] = X_I$. In other words, every Cauchy sequence $(x_n)_{n \in \mathbb{N}}$ in (X, d_{IX}) , which is convergent to a point x in (X, d_X) is convergent to x in (X, d_{IX}) , too.*

Proof. Given any point $x \in \overline{X}_I$ with $\overline{\text{id}}(x) \in X$, we need to prove that $x \in X_I$. Since $x \in \overline{X}_I$, there exists a sequence $(x_n)_{n \in \mathbb{N}}$ in X_I such that $d_{IX}(x, x_n) < \frac{1}{2^{n+2}}$ for all $n \in \mathbb{N}$. For every $n \in \mathbb{N}$ we have

$$d_{IX}(x_n, x_{n+1}) \leq d_{IX}(x_n, x) + d_{IX}(x, x_{n+1}) < \frac{1}{2^{n+2}} + \frac{1}{2^{n+3}} < \frac{1}{2^{n+1}}.$$

So, we can find a 1-Lipschitz path $\gamma_n : [\frac{1}{2^{n+1}}, \frac{1}{2^n}] \rightarrow X$ such that $\gamma_n(\frac{1}{2^{n+1}}) = x_{n+1}$ and $\gamma_n(\frac{1}{2^n}) = x_n$. The paths γ_n , $n \in \mathbb{N}$, compose a 1-Lipschitz path $\gamma : [0, 1] \rightarrow X$ defined by $\gamma(0) = \overline{\text{id}}(x)$ and $\gamma|_{[\frac{1}{2^{n+1}}, \frac{1}{2^n}]} = \gamma_n$ for all $n \in \mathbb{N}$. The definition of γ guarantees that

$$\lim_{n \rightarrow \infty} d_{IX}(x, x_n) = 0 = \lim_{n \rightarrow \infty} d_{IX}(x_n, \text{id}^{-1}(\overline{\text{id}}(x)))$$

and hence $x = \text{id}^{-1}(\overline{\text{id}}(x)) \in \text{id}^{-1}[X] = X_I$. $\square$

A metric d on a set X will be called *compact* (resp. *complete*, *separable*, *analytic*) if so is the metric space (X, d) .

Lemma 3.4. *Let (X, d_X) be a Lipschitz connected metric space.*

- (1) *If the metric d_X is complete, then so is the intrinsic metric d_{IX} .*
- (2) *The metric d_{IX} is analytic if and only if d_{IX} is separable and the metric d_X is analytic.*
- (3) *If the metric d_{IX} is separable (resp. totally bounded, compact), then so is the metric d_X .*

Proof. (1) Suppose that the metric d_X is complete and take a Cauchy sequence $(x_n)_{n \in \mathbb{N}}$ in the metric space (X, d_{IX}) . Since $d_X \leq d_{IX}$, the sequence $(x_n)_{n \in \mathbb{N}}$ remains Cauchy in the metric space (X, d_X) . Since (X, d_X) is complete, this sequence converges to some point $x \in X$ in the metric space (X, d_X) . Lemma 3.3 guarantees that $\lim_{n \rightarrow \infty} x_n = x$ in (X, d_{IX}) too, witnessing that the intrinsic metric d_{IX} is complete.

(2) If the metric d_{IX} is analytic, then d_{IX} is separable and d_X is analytic, because of the continuity of the identity map $(X, d_{IX}) \rightarrow (X, d_X)$.

Next, assume that the metric d_{IX} is separable and d_X is analytic. Let $\overline{X}$ be the completion of the metric space (X, d_X) and $\overline{X}_I$ be the completion of the metric space $X_I := (X, d_{IX})$. Let $\bar{I} : \overline{X}_I \rightarrow \overline{X}$ be the continuous extension of the (1-Lipschitz) identity map $I : X_I \rightarrow X$. By Lemma 3.3, $X_I = \bar{I}^{-1}[X]$. Since the spaces $\overline{X}_I$ and $\overline{X}$ are Polish, the preimage $X_I = \bar{I}^{-1}[X]$ is an analytic subspace of $\overline{X}_I$, by [13, 14.4].

The statement (3) follows from the 1-Lipschitz property of the identity map $(X, d_{IX}) \rightarrow (X, d_X)$. $\square$

The following simple example shows that the implications of Lemma 3.4 cannot be reversed in a general case.

Example 3.5. Let L be any line in the complex plane $\mathbb{C}$ and $o \in \mathbb{C} \setminus L$ be any point. For every uncountable compact zero-dimensional set $K \subseteq L$, the cone

$$X := \bigcup_{x \in K} [o, x]$$

over K with vertex o is a Lipschitz connected compact (by [8, 3.2.11]) metric space with respect to the metric d_X inherited from the complex plane. Thus (X, d_X) is also separable and totally bounded. Since $d_{IX}(x, y) \geq 2 \cdot \text{dist}(o, L)$ for any distinct points $x, y \in K$, the metric space (X, d_{IX}) is not separable (and hence not totally bounded and not compact).

If the compact space K has no isolated points, then for every countable dense set $D \subseteq K$, the Lipschitz connected metric subspace $Y = \bigcup_{x \in D} [o, x]$ of X is not complete (by the Baire Theorem) whereas the intrinsic metric d_{IY} is σ -compact and complete.

Example 3.6. The metric subspace $X = \{(x, \sin(\frac{1}{x})) : x > 0\}$ of the Euclidean plane is Lipschitz connected and not complete whereas the intrinsic metric on X is complete.

4. THE $\mathbb{R}$ -TREES GENERATED BY WEIGHTED TREES

In this section we describe a general construction of a (complete) $\mathbb{R}$ -tree, determined by a weighted tree.

Definition 4.1. A *forest* is a partially ordered set $(T, \preceq)$ such that for every $t \in T$ the set

$$\downarrow t := \{x \in T : x \preceq t\}$$

is finite and linearly ordered.

For any elements $s, t \in T$, the largest element of the linearly ordered set $\downarrow s \cap \downarrow t$ is called the *infimum* of s and t and is denoted by $s \wedge t$.

For an element t of a forest T , the set $\text{succ}(t) := \min\{s \in T : t \prec s\}$ is called the set of *immediate successors* of t in the forest.

A subset B of a forest T is called a *branch* of T if B is a maximal linearly ordered subset of T . The *boundary* ∂T of a forest T is the set of all infinite branches in T .

For a forest T by $\min T$ denote the set of minimal elements of T . For a nonempty forest T , the set $\min T$ is not empty. A forest T is a *tree* if the set $\min T$ is a singleton, consisting of the smallest element of T , called the *root* of the tree T and denoted by r_T .

Observe that for every element r of a forest T , the set

$$\uparrow r := \{x \in T : r \preceq x\}$$

is a tree with root r .

Definition 4.2. A *weighted tree* is a pair (T, w) consisting of a tree T and a function $w : T \rightarrow [0, +\infty)$ assigning to every vertex $t \in T$ its *weight* $w_t = w(t)$ so that $w_t = 0$ if and only if t is the root of T .

Every weighted tree (T, w) determines an $\mathbb{R}$ -tree

$$T_w = (\{r_T\} \times \{0\}) \cup \bigcup_{t \in T} \{t\} \times (0, w_t],$$

endowed with the metric

$$d_w((t, t'), (s, s')) := \begin{cases} |t' - s'| & \text{if } s = t; \\ t' - w_t + s' - w_s + \sum_{x \in \downarrow s \Delta \downarrow t} w_x & \text{otherwise,} \end{cases}$$

where $A \Delta B$ stands for the symmetric difference $(A \setminus B) \cup (B \setminus A)$ of the sets A, B . It is easy to check that the metric space (T_w, d_w) is indeed an $\mathbb{R}$ -tree. The $\mathbb{R}$ -tree T_w will be called the *$\mathbb{R}$ -tree of the weighted tree (T, w)* .

The $\mathbb{R}$ -tree $T_w \subset T \times \mathbb{R}$ carries a natural lexicographic order

$$(s, s') \leq_L (t, t') \iff s \prec t \vee (s = t \wedge s' \leq t').$$

So, for every element $t'' \in T_w$ we can consider its lower and upper sets

$$\downarrow t'' := \{s'' \in T_w : s'' \leq_L t''\} \quad \text{and} \quad \uparrow t'' := \{s'' \in T_w : t'' \leq_L s''\}$$

in the partially ordered set $(T_w, \leq_L)$. Since T is a tree, the lower set $\downarrow t''$ of every element $t'' \in T_w$ is linearly ordered. For two elements $a'', b'' \in T_w$, let $a'' \wedge b''$ denote the largest element of the linearly ordered set $\downarrow a'' \cap \downarrow b''$ in the partially ordered set $(T_w, \leq_L)$.

Now we shall characterize weighted trees whose $\mathbb{R}$ -trees are totally bounded or separable.

Definition 4.3. A weighted tree (T, w) is defined to be *precompact* if for every $\varepsilon > 0$ there exists a finite set $F \subseteq T$ such that for every finite linearly ordered set $L \subseteq T \setminus F$ we have $\sum_{x \in L} w_x < \varepsilon$.


Proposition 4.4. A weighted tree (T, w) is precompact if and only if the metric space T_w is totally bounded.

Proof. To prove the “if” part, assume that the metric space T_w is totally bounded. Then for every $\varepsilon > 0$, there exists a finite set $F'' \subseteq T_w$ such that for every $x'' \in T_w$, there exists $y'' \in F''$ with $d_w(x'', y'') < \varepsilon$. Since F'' is finite, there exists a finite set $F \subseteq T$ such that $F'' \subseteq \bigcup \{\downarrow(t, w_t) : t \in F\} \subseteq T_w$. Replacing F by the finite set $\downarrow F := \bigcup_{t \in F} \downarrow t$, we can assume that $F = \downarrow F$. We claim that $\sum_{x \in L} w_x < \varepsilon$ for every finite linearly ordered set $L \subseteq T \setminus F$. Since L is finite and linearly ordered, there exists an element $u \in L$ such that $L \subseteq \downarrow u$. By the choice of the set F'' , there exists a point

$y'' \in F''$ such that $d_w((u, w_u), y'') < \varepsilon$. Write the element y'' as (y, y') for some $y \in T$ and $y' \in [0, w_y]$. The choice of the set F ensures that $y \in F$ and hence $L \subseteq \downarrow u \setminus F \subseteq \downarrow u \setminus \downarrow y$. Then

$$\sum_{x \in L} w_x \leq \sum_{x \in \downarrow u \setminus \downarrow y} w_x \leq d_w((u, w_u), y'') < \varepsilon,$$

witnessing that the weighted tree (T, w) is precompact.



To prove the “only if” part, assume that the weighted tree (T, w) is precompact. To prove that the metric space T_w is totally bounded, fix any $\varepsilon > 0$. By the precompactness of the weighted tree (T, w) , there exists a nonempty finite set $F \subseteq T$ such that $\sum_{x \in L} w_x < \varepsilon$ for every finite linearly ordered set $L \subseteq T \setminus F$. Replacing F with the finite set $\downarrow F := \bigcup_{t \in F} \downarrow t$, we can assume that $F = \downarrow F$. Since the set F is finite, the set

$$K := \bigcup \{ \downarrow(t, w_t) : t \in F \} = (\{r_T\} \times \{0\}) \cup \bigcup_{t \in F} (\{t\} \times (0, w_t])$$

is compact and hence totally bounded in the metric space T_w . Then there exists a finite set $F'' \subseteq K$ such that for every $x'' \in K$ there exists $y'' \in F''$ with $d_w(x'', y'') < \varepsilon$. Replacing F'' by the set $F'' \cup \{(t, w_t) : t \in F\}$, we can assume that $\{(t, w_t) : t \in F\} \subseteq F''$. We claim that for every $x'' \in T_w$, there exists $y'' \in F''$ such that $d_w(x'', y'') < \varepsilon$.

Indeed, if $x'' \in K$, then the existence of a point $y'' \in F''$ with $d_w(x'', y'') < \varepsilon$ follows from the choice of the set F'' . So, we assume that $x'' \notin K$. In this case, $x'' = (x, x')$ for some $x \in T \setminus F$ and $x' \in (0, w_x]$. Since $r_T \in \downarrow F = F$, there exists a point $y \in F \cap \downarrow x$ such that $F \cap \downarrow x = \downarrow y$. The choice of the set F'' ensures that it contains the point $y'' := (y, w_y)$. The choice of F guarantees that $\sum_{t \in \downarrow x \setminus \downarrow y} w_t = \sum_{t \in \downarrow x \setminus F} w_t < \varepsilon$ and hence $d_w(x'', y'') = x' - w_x + \sum_{t \in \downarrow x \setminus \downarrow y} w_t \leq \sum_{t \in \downarrow x \setminus \downarrow y} w_t < \varepsilon$, by the definition of the metric d_w . This shows that the metric space T_w is totally bounded. $\square$

Proposition 4.5. *For a weighted tree (T, w) , the $\mathbb{R}$ -tree T_w is separable if and only if the tree T is countable.*

Proof. If the tree T is countable, then $T_w \cap (T \times \mathbb{Q})$ is a countable dense set in the $\mathbb{R}$ -tree, witnessing that the metric space T_w is separable.

Next, assume that the tree T is uncountable. The definition of metric d_w ensures that for every node $t \in T$, the set $\{t\} \times (0, w_t)$ is open in the metric space T_w . Therefore, the metric space T_w contains the uncountable family of pairwise disjoint open sets $\{\{t\} \times (0, w_t) : t \in T\}$ and hence T_w cannot be separable, according to [8, Corollary 4.1.16]. $\square$

Let (T, w) be a weighted tree. In the set ∂T of all infinite branches of T , consider the subset

$$\partial T_w := \{L \in \partial T : |L| = \infty, \sum_{t \in L} w_t < \infty\}.$$

On the set $\bar{T}_w := T_w \cup \partial T_w$ we define a metric $\bar{d}_w$, that extends the metric d_w on T_w as follows:

$$\bar{d}_w(x'', y'') := \begin{cases} |x' - y'| & \text{if } x'' = (x, x'), y'' = (y, y') \in T_w \text{ and } x = y; \\ x' - w_x + y' - w_y + \sum_{t \in \downarrow x \Delta \downarrow y} w_t & \text{if } x'' = (x, x'), y'' = (y, y') \in T_w \text{ and } x \neq y; \\ x' - w_x + \sum_{t \in \downarrow x \Delta y''} w_t & \text{if } x'' = (x, x') \in T_w, y'' \in \partial T_w \text{ and } x \notin y''; \\ w_x - x' + \sum_{t \in y'' \setminus \downarrow x} w_t & \text{if } x'' = (x, x') \in T_w, y'' \in \partial T_w \text{ and } x \in y''; \\ \sum_{t \in x'' \Delta y''} w_t & \text{if } x'', y'' \in \partial T_w. \end{cases}$$

Proposition 4.6. *For every weighted tree (T, w) , the metric space $\bar{T}_w$ is a completion of the metric space T_w .*

Proof. The definition of the metric on $\bar{T}_w$ implies that the metric space T_w is a dense subspace of the metric space $\bar{T}_w$. To prove that $\bar{T}_w$ is a completion of T_w , it suffices to show that every Cauchy sequence $(s''_n)_{n \in \omega}$ in the metric space T_w has a limit in the space $\bar{T}_w$. If for some element t of T_w , the set $\{n \in \omega : s''_n = t\}$ is infinite, then t is the limit of the Cauchy sequence $(s''_n)_{n \in \omega}$ and we are done. So, we can assume that all elements of the Cauchy sequence $(s''_n)_{n \in \omega}$ are pairwise distinct and moreover all of them are distinct from the root $(r_T, 0)$ of the $\mathbb{R}$ -tree T_w . For every $n \in \omega$, write the element s''_n of T_w as a pair (s_n, s'_n) where $s_n \in T$ and $s'_n \in (0, w_{s_n}]$. Let $\check{s}_n := \max(\downarrow s_n \setminus \{s_n\})$ be the immediate predecessor of s_n in the tree T .

In the tree T , consider the set

$$\Lambda := \{t \in T : |\{n \in \omega : t \preceq \check{s}_n\}| = \omega\}$$

and observe that it contains the root r_T of the tree T and hence is not empty. Two cases are possible.


1. First assume that the set Λ has no maximal elements. In this case Λ contains an increasing sequence $(\lambda_n)_{n \in \omega}$. For this sequence the set $b := \bigcup_{n \in \omega} \downarrow \lambda_n$ is an infinite branch of the tree T such that for every $t \in b$ the set $\{n \in \omega : t \preceq \check{s}_n\}$ is infinite. Write the branch b as $b = \{b_n\}_{n \in \omega}$, where b_n is a unique element of b such that $|\downarrow b_n| = n + 1$. We claim that $b \in \partial T_w$. Since the sequence $(s''_n)_{n \in \omega}$ is Cauchy, for every $\varepsilon > 0$, there exists $m \in \mathbb{N}$ such that $d_w(s''_n, s''_m) < \varepsilon$ for every $n \geq m$.

Since the sequence $(b_n)_{n \in \omega}$ is unbounded in the tree T , there exists a number $i \in \omega$ such that $b_i \not\preceq \check{s}_m$. Given any number $j \geq i$, find a number $n \geq m$ such that $b_j \preceq \check{s}_n$. Observe that $b_i, b_j, \check{s}_n$ and $\check{s}_n \wedge \check{s}_m$ are elements of the linearly ordered set $\downarrow \check{s}_n$. Assuming that $\check{s}_m \wedge \check{s}_n \not\preceq b_i$, we conclude that $b_i \preceq \check{s}_m \wedge \check{s}_n \preceq \check{s}_m$, which contradicts the choice of the number i . This contradiction shows that $\check{s}_m \wedge \check{s}_n \preceq b_i$.

In this case, the definition of the metric d_w ensures that

$$\sum_{k=i+1}^j w_{b_k} \leq -w_{s_n} - w_{s_m} + \sum_{x \in \downarrow s_n \Delta \downarrow s_m} w_x \leq d_w(s''_m, s''_n) < \varepsilon$$

and hence $\sum_{k=i+1}^{\infty} w_{b_k} \leq \varepsilon < \infty$, witnessing that $b \in \partial T_w$.




Since the sequence $(s''_n)_{n \in \omega}$ is Cauchy and $\varepsilon > 0$ was arbitrary, the equality $b = \lim_{n \rightarrow \infty} s''_n$ will follow as soon as we check that $\bar{d}_w(b, s''_n) < 2\varepsilon$ for infinitely many numbers n . Indeed, for every number $n \geq m$ in the infinite set $\{n \in \omega : b_i \preceq s_n\}$ we have $s_n \wedge s_m \preceq b_i$ and

$$\begin{aligned} \bar{d}_w(s''_n, b) &= s'_n - w_{s_n} + \sum_{x \in \downarrow s_n \triangle b} w_x = s'_n - w_{s_n} + \sum_{x \in \downarrow s_n \setminus b} w_x + \sum_{x \in b \setminus \downarrow s_n} w_x \\ &\leq s'_n - w_{s_n} + \sum_{x \in \downarrow s_n \setminus \downarrow b_i} w_x + \sum_{k=i+1}^{\infty} w_{b_k} \\ &\leq s'_n - w_{s_n} + \sum_{x \in \downarrow s_n \setminus \downarrow s_m} w_x + \sum_{k=i+1}^{\infty} w_{b_k} \leq d_w(s''_n, s''_m) + \varepsilon < 2\varepsilon. \end{aligned}$$

2. Next, assume that the set Λ has a maximal element λ . Consider the infinite set $I := \{n \in \omega : \lambda \preceq s_n\}$. If for some $s \in T$, the set $J := \{n \in I : s = s_n\}$ is infinite, then the maximality of λ ensures that λ is the immediate predecessor of s in the tree T . Then the subsequence $(s''_n)_{n \in J}$ of the sequence $(s''_n)_{n \in \omega}$ is contained in the compact set $\{(\lambda, w_\lambda)\} \cup (\{s\} \times (0, w_s])$ and hence the Cauchy sequence $(s''_n)_{n \in \omega}$ converges to some points of this compact set. So, assume that for every $s \in T$, the set $\{n \in I : s_n = s\}$ is finite. In this case we shall show that the Cauchy sequence $(s''_n)_{n \in \omega}$ converges to the element $\lambda'' := (\lambda, w_\lambda)$ of the $\mathbb{R}$ -tree T_w . Since the sequence $(s''_n)_{n \in \omega}$ is Cauchy and consists of pairwise distinct elements, it suffices to show that for every $\varepsilon > 0$ there exists $n \in \omega$ such that $d_w(\lambda'', s''_n) < \varepsilon$. Since the sequence $(s''_n)_{n \in \omega}$ is Cauchy, there exists $m \in \omega$ such that $d_w(s''_n, s''_k) < \varepsilon$ for all $n, k \geq m$. We can assume that $\lambda \preceq s_m \prec s_m$. Thus the element λ has an immediate successor $\hat{\lambda}$ in the tree T such that $\lambda \prec \hat{\lambda} \preceq s_m$. The maximality of λ ensures that $\hat{\lambda} \notin \Lambda$ and hence the set $\{n \in \omega : \hat{\lambda} \preceq s_n\}$ is finite. Then we can choose a number $n \in I$ such that $n \geq m$ and $\hat{\lambda} \not\preceq s_n$. Then $s_n \wedge s_m = s_n = \lambda$ and

$$d_w(\lambda'', s''_n) \leq s'_n - w_{s_n} + s'_m - w_{s_m} + \sum_{x \in \downarrow s_n \triangle \downarrow s_m} w_x = d_w(s''_n, s''_m) < \varepsilon.$$



□

Propositions 2.4, 4.4, 4.5, 4.6 and the definition of the metric $\bar{d}_w$ imply the following corollary describing the property of the completion $\bar{T}_w$ of the $\mathbb{R}$ -tree T_w of a weighted tree (T, w) .

Corollary 4.7. *For every weighted tree (T, w) ,*

- (1) *the complete metric space $\bar{T}_w$ is an $\mathbb{R}$ -tree and hence is an injective metric space;*
- (2) *the space $\bar{T}_w$ is compact if and only if the weighted tree (T, w) is precompact;*
- (3) *the space $\bar{T}_w$ is separable if and only if the tree T is countable.*

5. PROOF OF THEOREM 1.2

To prove Theorem 1.2, we need to demonstrate the equivalence of the following conditions for every metric space X :

- (1) X is a Lipschitz image of an injective metric space;
- (2) X is a Lipschitz image of some complete $\mathbb{R}$ -tree;
- (3) X is Lipschitz connected.

The implication (2) $\Rightarrow$ (1) follows from Proposition 2.4 (saying that every complete $\mathbb{R}$ -tree is an injective metric space), and the implication (1) $\Rightarrow$ (3) follows from the Lipschitz connectedness of injective metric spaces, proved in Example 3.2.

To prove that (3) $\Rightarrow$ (2), assume that a metric space X is Lipschitz connected. Fix any point $x_0 \in X$ and for every point $a \in X$ find a real number w_a and a 1-Lipschitz path $\gamma_a : [0, w_a] \rightarrow X$ such that $\gamma_a(0) = x_0$ and $\gamma_a(w_a) = a$. If $a = x_0$, then will assume that $w_a = 0$.

Consider the tree $T := X$ endowed with the partial order

$$\preceq := \{(x, y) \in X \times X : x = y \vee x = x_0\}$$

The function $w : X \rightarrow [0, \infty)$, $w : a \mapsto w_a$, turns T into a weighted tree. The weighted tree (T, w) generates the $\mathbb{R}$ -tree T_w . Since the tree T contains no infinite branches, the $\mathbb{R}$ -tree T_w coincides with its completion $\bar{T}_w = T_w \cup \partial T_w$ and hence T_w is a complete $\mathbb{R}$ -tree.

The 1-Lipschitz maps γ_a , $a \in X$, compose a surjective 1-Lipschitz map $\gamma : T_w \rightarrow X$ such that $\gamma((r_T, 0)) = x_0$ and $\gamma((a, t)) = \gamma_a(t)$ for $(a, t) \in \bigcup_{a \in X} \{a\} \times (0, w_a]$. Indeed for any $(a, t), (b, s) \in T_w$ if $a = b$

$$d_X(\gamma((a, t)), \gamma((b, s))) = d_X(\gamma_a(t), \gamma_a(s)) \leq |t - s|,$$

because γ_a is a 1-Lipschitz map. If $a \neq b$

$$\begin{aligned} d_X(\gamma((a, t)), \gamma((b, s))) &= d_X(\gamma_a(t), \gamma_b(s)) \leq d_X(\gamma_a(t), x_0) + d_X(x_0, \gamma_b(s)) = \\ &= d_X(\gamma_a(t), \gamma_a(0)) + d_X(\gamma_b(0), \gamma_b(s)) \leq t + s, \end{aligned}$$

because γ_a, γ_b are 1-Lipschitz maps. The 1-Lipschitz map γ witnesses that X is a 1-Lipschitz image of the complete $\mathbb{R}$ -tree T_w .

6. PROOF OF THEOREM 1.3

Theorem 1.3 will follow as soon as we check the equivalence of the following conditions for every metric space X :

- (1) X is a Lipschitz image of some compact $\mathbb{R}$ -tree;
- (2) X is a Lipschitz image of a totally bounded injective metric space;
- (3) X is a Lipschitz image of a compact 1-Lipschitz connected metric space;
- (4) X is a Lipschitz image of an L -Lipschitz connected compact metric space for some $L \geq 1$;
- (5) X is compact, Lipschitz connected, and the intrinsic metric d_{IX} is totally bounded.

The implication (1) $\Rightarrow$ (2) follows from Proposition 2.4 (saying that every complete $\mathbb{R}$ -tree is an injective metric space), and the total boundedness of compact metric spaces.

The implication (2) $\Rightarrow$ (3) follows from the completeness and 1-Lipschitz connectedness of injective metric spaces, and the implication (3) $\Rightarrow$ (4) is obvious.

(4) $\Rightarrow$ (5) Assume that a metric space (X, d_X) is the image of some L -Lipschitz connected compact metric space (Y, d_Y) under a Lipschitz map $f : Y \rightarrow X$. Then X is compact and Lipschitz connected, so the intrinsic metric d_{IX} on X is well-defined. Scaling the metric of the metric space Y , we can assume that the Lipschitz map f is 1-Lipschitz. In this case the definition of the intrinsic metric implies that the function f remains 1-Lipschitz with respect to the intrinsic metrics on the metric spaces X and Y . The L -Lipschitz connectedness of the metric space Y implies that $d_Y \leq d_{IY} \leq L \cdot d_Y$, which implies that the identity map $(Y, d_Y) \rightarrow (Y, d_{IY})$ is Lipschitz and hence the metric space (Y, d_{IY}) is totally bounded and so is its 1-Lipschitz image (X, d_{IX}) .

(5) $\Rightarrow$ (1) Assume that the metric space (X, d_X) is compact, Lipschitz connected, and its intrinsic metric d_{IX} is totally bounded. Let $D := \sup\{d_{IX}(x, y) : x, y \in X\}$ be the diameter of the (totally) bounded metric space (X, d_{IX}) .

Since the metric space (X, d_{IX}) is totally bounded, there exists a sequence $(\mathcal{F}_n)_{n \in \omega}$ of finite partitions of X such that $\mathcal{F}_0 = \{X\}$ and for every $n \in \omega$, every set $F \in \mathcal{F}_n$ has d_{IX} -diameter $\sup\{d_{IX}(x, y) : x, y \in F\} < \frac{D+1}{2^n}$. Replacing each partition $\mathcal{F}_n$ with $n \in \mathbb{N}$ by the partition $\mathcal{F}_n \wedge \mathcal{F}_{n-1} := \{A \cap B : A \in \mathcal{F}_{n-1}, B \in \mathcal{F}_n, A \cap B \neq \emptyset\}$, we can assume that every set $F \in \mathcal{F}_n$ is contained in a unique set $\check{F} \in \mathcal{F}_{n-1}$. For every $n \in \omega$ and $F \in \mathcal{F}_n$, choose a point $c_F \in F$.

Consider the tree $T := \bigcup_{n \in \omega} (\{n\} \times \mathcal{F}_n)$ endowed with the partial order $\preceq$, defined by $(n, F) \preceq (m, E)$ iff $n \leq m$ and $E \subseteq F$. Next, endow the tree T with the weight function $w : T \rightarrow [0, \infty)$ defined by

$$w(n, F) = w_{(n, F)} = \begin{cases} \frac{D+1}{2^{n-1}} & \text{if } n > 0; \\ 0 & \text{if } n = 0. \end{cases}$$




FIGURE 1. The weighted tree (T, w) .

Let T_w be the $\mathbb{R}$ -tree of the weighted tree (T, w) . Since the families $\mathcal{F}_n$ are finite and $\sum_{t \in b} w_t = 2(D+1)$ for every $b \in \partial T$, the weighted tree (T, w) is precompact. By Proposition 4.4, the metric space T_w is totally bounded and hence its completion $\overline{T}_w$ is a compact $\mathbb{R}$ -tree.

It remains to construct a 1-Lipschitz surjective map $\bar{\gamma} : \overline{T}_w \rightarrow X$. For every $n \in \mathbb{N}$ and $F \in \mathcal{F}_n$, choose a 1-Lipschitz function $\gamma_{n, F} : [0, w_{(n, F)}] \rightarrow X$ such that $\gamma_{(n, F)}(0) = c_{\check{F}}$ and $\gamma_{(n, F)}(w_{(n, F)}) = c_F$. The function $\gamma_{(n, F)}$ exists because $\{c_{\check{F}}, c_F\} \subseteq \check{F} \cup F = \check{F}$ and the set $\check{F} \in \mathcal{F}_{n-1}$ has d_{IX} -diameter $< \frac{D+1}{2^{n-1}} = w_{(n, F)}$. Next, consider the function $\gamma : T_w \rightarrow X$ defined by $\gamma((0, X), 0) = c_X$ and $\gamma((n, F), t) = \gamma_{n, F}(t)$ for every $n \in \mathbb{N}$, $F \in \mathcal{F}_n$ and $t \in (0, w_{(n, F)})$. The definition of the metric d_w on the $\mathbb{R}$ -tree T_w ensures that the map $\gamma : T_w \rightarrow X$ is 1-Lipschitz. Observe that the image $\gamma[T_w]$ is dense in X (because it contains the dense subset $\{c_F : n \in \mathbb{N}, F \in \mathcal{F}_n\}$ of X).

Since the metric space X is complete, the 1-Lipschitz map $\gamma : T_w \rightarrow X$ admits a unique 1-Lipschitz extension $\bar{\gamma} : \overline{T}_w \rightarrow X$. The total boundedness of the $\mathbb{R}$ -tree T_w implies the compactness of its completion $\overline{T}_w$. Then the image $\bar{\gamma}[\overline{T}_w]$ is a compact dense subset of X , witnessing that the 1-Lipschitz map $\bar{\gamma} : \overline{T}_w \rightarrow X$ is surjective and hence X is a 1-Lipschitz image of the compact $\mathbb{R}$ -tree $\overline{T}_w$.

Remark 6.1. Although compactness of (X, d_X) follows from any of the conditions (1)–(5), the compactness of (X, d_X) is not sufficient to infer these conditions, which is showed in Example 3.5.

7. PROOF OF THEOREM 1.4

Theorem 1.4 will follow as soon as we check the equivalence of the following conditions for every metric space X :

- (1) X is a Lipschitz image of some separable complete $\mathbb{R}$ -tree;
- (2) X is a Lipschitz image of a separable injective metric space;
- (3) X is a 1-Lipschitz image of the Urysohn universal metric space;
- (4) X is a 1-Lipschitz image of a 1-Lipschitz connected separable complete metric space;
- (5) X is a Lipschitz image of an L -Lipschitz connected analytic metric space for some $L \geq 1$;
- (6) X is analytic, Lipschitz connected, and the intrinsic metric d_{IX} is separable.

The implication (1) $\Rightarrow$ (2) follows from Proposition 2.4 (saying that every complete $\mathbb{R}$ -tree is an injective metric space).

To prove that (2) $\Rightarrow$ (3), assume that the metric space X is the image of a separable injective space Y under a Lipschitz map $f : Y \rightarrow X$. Scaling the metric of the metric space Y , we can assume that the Lipschitz map f is 1-Lipschitz. By [18], the Urysohn universal metric space $\mathbb{U}$ contains an isometric copy of any separable metric space. So, there exists an isometric embedding $e : Y \rightarrow \mathbb{U}$. By the injectivity of Y , the 1-Lipschitz map $e^{-1} : e[Y] \rightarrow Y$ defined on the subset $e[Y]$ of $\mathbb{U}$ admits a 1-Lipschitz extension $g : \mathbb{U} \rightarrow Y$. Then $f \circ g : \mathbb{U} \rightarrow X$ is a surjective 1-Lipschitz map, witnessing that X is a 1-Lipschitz image of the Urysohn universal metric space $\mathbb{U}$.

To see the implication (3) $\Rightarrow$ (4) it suffices to check that the Urysohn universal metric space $\mathbb{U}$ is 1-Lipschitz connected. Given any points $x, y \in \mathbb{U}$, consider the function $f : \{0, d(x, y)\} \rightarrow \{x, y\} \subseteq \mathbb{U}$ such that $f(0) = x$ and $f(d(x, y)) = y$. By [3, 12] (see also [2, 15, 17]), the isometric embedding $f : \{0, d(x, y)\} \rightarrow \mathbb{U}$ extends to a 1-Lipschitz map $f : [0, d(x, y)] \rightarrow \mathbb{U}$, witnessing that $\mathbb{U}$ is 1-Lipschitz connected.

The implication (4) $\Rightarrow$ (5) is trivial.

(5) $\Rightarrow$ (6) Assume that for some $L \geq 1$, a metric space (X, d_X) is the image of some L -Lipschitz connected analytic metric space (Y, d_Y) under a Lipschitz map $f : Y \rightarrow X$. Then X is analytic and Lipschitz connected, so the intrinsic metric d_{IX} on X is well-defined. Scaling the metric of the metric space Y , we can assume that the Lipschitz map f is 1-Lipschitz. In this case the definition of the intrinsic metric implies that the function f remains 1-Lipschitz with respect to the intrinsic metrics on the metric spaces X and Y . The L -Lipschitz connectedness of the metric space Y implies that $d_Y \leq d_{IY} \leq L \cdot d_Y$, which implies that the identity map $(Y, d_Y) \rightarrow (Y, d_{IY})$ is Lipschitz and hence the metric space (Y, d_{IY}) is separable and so is its 1-Lipschitz image (X, d_{IX}) .

(6) $\Rightarrow$ (1) Assume that the metric space (X, d_X) is analytic, Lipschitz connected, and its intrinsic metric d_{IX} is separable. By Proposition 3.4, the metric space $X_I := (X, d_{IX})$ is analytic and hence it is the image of a separable complete metric space (Z, d_Z) under a surjective continuous map $f : Z \rightarrow X_I$. Consider the metric $\hat{d}_Z$ on Z , defined by the formula $\hat{d}_Z(z, z') := \max\{d_Z(z, z'), d_{IX}(f(z), f(z'))\}$. The continuity of the function f ensures that the metric $\hat{d}_Z$ is topologically equivalent to the metric d_Z . Since $d_Z \leq \hat{d}_Z$, the completeness of the metric d_Z implies the completeness of the metric $\hat{d}_Z$. Replacing the metric d_Z by the metric $\hat{d}_Z$, we can assume that the function $f : Z \rightarrow X_I$ is 1-Lipschitz.

Since the metric space Z is separable, there exists a sequence $(\mathcal{F}_n)_{n \in \omega}$ of countable partitions of Z such that $\mathcal{F}_0 = \{Z\}$ and for every $n \in \mathbb{N}$, every set $F \in \mathcal{F}_n$ has diameter $\sup\{d_Z(x, y) : x, y \in F\} < \frac{1}{2^n}$. Replacing each partition $\mathcal{F}_n$ by the partition $\mathcal{F}_n \wedge \mathcal{F}_{n-1} := \{A \cap B : A \in \mathcal{F}_{n-1}, B \in \mathcal{F}_n, A \cap B \neq \emptyset\}$, we can assume that every set $F \in \mathcal{F}_n$ is contained in a unique set $\tilde{F} \in \mathcal{F}_{n-1}$. For every $n \in \omega$ and $F \in \mathcal{F}_n$, choose a point $c_F \in F$.

Consider the countable tree $T := \bigcup_{n \in \omega} (\{n\} \times \mathcal{F}_n)$ endowed with the partial order $\preceq$, defined by $(n, F) \preceq (m, E)$ iff $n \leq m$ and $F \subseteq E$. Next, endow the tree T with the weight function $w : T \rightarrow [0, \infty)$

defined by $w_{(0,Z)} = 0$ and $w_{(n,F)} = d_{IX}(f(c_{\tilde{F}}), f(c_F)) + \frac{1}{2^n}$ for every $n \in \mathbb{N}$ and $F \in \mathcal{F}_n$ (in this formula $\tilde{F}$ denotes the unique set in $\mathcal{F}_{n-1}$ that contains the set F).

Let T_w be the $\mathbb{R}$ -tree of the weighted tree (T, w) and $\overline{T}_w = T_w \cup \partial T_w$ be its completion. Proposition 4.5 ensures that the $\mathbb{R}$ -tree T_w is separable and hence its completion $\overline{T}_w$ is a separable complete $\mathbb{R}$ -tree.

It remains to construct a 1-Lipschitz surjective map $\overline{\gamma} : \overline{T}_w \rightarrow X$. For every $n \in \mathbb{N}$ and $F \in \mathcal{F}_n$, choose a 1-Lipschitz function $\gamma_{n,F} : [0, w_{(n,F)}] \rightarrow X$ such that $\gamma_{n,F}(0) = f(c_{\tilde{F}})$ and $\gamma_{n,F}(w_{(n,F)}) = f(c_F)$. The function $\gamma_{n,F}$ exists because $d_{IX}(f(c_{\tilde{F}}), f(c_F)) < w_{(n,F)}$. Next, consider the function $\gamma : T_w \rightarrow X$ defined by $\gamma((0, X), 0) = f(c_Z)$ and $\gamma((n, F), t) = \gamma_{n,F}(t)$ for every $n \in \mathbb{N}$, $F \in \mathcal{F}_n$ and $t \in (0, w_{(n,F)})$. The definition of the metric d_w on the $\mathbb{R}$ -tree T_w ensures that the map $\gamma : T_w \rightarrow X$ is 1-Lipschitz.

Consider the set ∂T of all infinite branches of the tree T and observe that every branch $b \in \partial T$ is equal to the set $\{(n, F_n) : n \in \omega\}$ for some decreasing sequence $(F_n)_{n \in \omega} \in \prod_{n \in \omega} \mathcal{F}_n$. Observe that for every $n \geq 2$, the set $\{c_{\tilde{F}_n}, c_{F_n}\}$ is a subset of the set $\tilde{F}_n \in \mathcal{F}_{n-1}$ and hence $d_Z(c_{F_{n-1}}, c_{F_n}) = d_Z(c_{\tilde{F}_n}, c_{F_n}) < \frac{1}{2^{n-1}}$. Then the sequence $(c_{F_n})_{n \in \omega}$ is Cauchy in the complete metric space Z and has a limit $\lambda(b) := \lim_{n \rightarrow \infty} c_{F_n} \in Z$, which coincides with the unique point of the intersection $\bigcap_{n \in \omega} \overline{F_n}$. The correspondence $\lambda : \partial T \mapsto \lambda(b)$, determines a function $\lambda : \partial T \rightarrow Z$. Observe that for every point $z \in Z$ there exists a unique branch $b = \{(n, F_n)\}_{n \in \omega} \in \partial T$ such that $z \in \bigcap_{n \in \omega} F_n = \bigcap_{n \in \omega} \overline{F_n} = \{\lambda(b)\}$, witnessing that the function $\lambda : \partial T \rightarrow Z$ is surjective. Taking into account that diameters of the sets in the partition $\mathcal{F}_n$ tend to zero as n tends to infinity, one can show that the map $\lambda : \partial T \rightarrow Z$ is continuous with respect to the topology on ∂T inherited from the Tychonoff product $\prod_{n \in \omega} (\{n\} \times \mathcal{F}_n)$ of discrete spaces $\{n\} \times \mathcal{F}_n$, $n \in \omega$.

Observe that for every branch $b = \{(n, F_n)\}_{n \in \omega}$ of the tree T and every $n \in \mathbb{N}$ we have $w_{(n, F_n)} = d_{IX}(f(c_{F_{n-1}}), f(c_{F_n})) + \frac{1}{2^n} \leq d_Z(c_{F_{n-1}}, c_{F_n}) + \frac{1}{2^n} < \frac{1}{2^{n-1}} + \frac{1}{2^n}$ and hence $\sum_{t \in b} w_t = \sum_{n \in \omega} w_{(n, F_n)} < \infty$, witnessing that $\partial T_w = \partial T$.

Since the map $\gamma : T_w \rightarrow X$ is 1-Lipschitz, there exists a unique 1-Lipschitz extension $\overline{\gamma} : \overline{T}_w \rightarrow \overline{X}$ to the completion of the space X . We claim that $\overline{\gamma}[\overline{T}_w] = X$.

The inclusion $\overline{\gamma}[\overline{T}_w] \subseteq X$ will follow as soon as we check that $\overline{\gamma}[\partial T] \subseteq X$. Fix any element $b \in \partial T$ and find a unique decreasing sequence of sets $(F_n)_{n \in \omega} \in \prod_{n \in \omega} \mathcal{F}_n$ such that $b = \{(n, F_n)\}_{n \in \omega}$. Then $\lambda(b) = \lim_{n \rightarrow \infty} c_{F_n}$. The definition of the metric on the $\mathbb{R}$ -tree $\overline{T}_w$ ensures that

$$b = \lim_{n \rightarrow \infty} ((n, F_n), w_{(n, F_n)})$$

in $\overline{T}_w$ and hence $\overline{\gamma}(b) = \lim_{n \rightarrow \infty} \gamma((n, F_n), w_{(n, F_n)}) = \lim_{n \rightarrow \infty} \gamma_{n, F}(w_{(n, F_n)}) = \lim_{n \rightarrow \infty} f(c_{F_n}) = f(\lim_{n \rightarrow \infty} c_{F_n}) = f(\lambda(b)) \in f[Z] = X$.

Next, we show that $X \subseteq \overline{\gamma}[\overline{T}_w]$. Given any point $x \in X$, choose any point $z \in f^{-1}(x) \subseteq Z$ and find a unique branch $b = \{(n, F_n)\}_{n \in \omega} \in \partial T$ such that $z \in \bigcap_{n \in \omega} F_n$. Then $z = \lambda(b)$ and $x = f(\lambda(b)) = \overline{\gamma}(b)$.

Therefore, $\overline{\gamma} : \overline{T}_w \rightarrow X$ is a 1-Lipschitz surjective map witnessing that X is a 1-Lipschitz image of the separable complete $\mathbb{R}$ -tree $\overline{T}_w$.

Remark 7.1. Although analyticity and separability of (X, d_X) follows from each of the conditions (1)–(6), the analyticity of (X, d_X) is not sufficient to infer the equivalent conditions of Theorem 1.4, which is showed in Example 3.5.

ACKNOWLEDGEMENT

This work was co-financed by the Minister of Science (Poland) under the ‘‘Regional Excellence Initiative’’ program (project no.: RID/SP/0015/2024/01).

REFERENCES

- [1] N. Aronszajn, P. Panitchpakdi, *Extension of uniformly continuous transformations and hyperconvex metric spaces*, Pacific J. Math. 6:3 (1956), 405–439.
- [2] T. Banakh, F. Stobin, *Embedding topological fractals in universal spaces*, J. Fractal Geom. 2 (2015), no. 4, pp. 377–388.
- [3] S. A. Bogatyĭ, *Compact homogeneity of Uryson’s universal metric space* (Russian), Russ. Math. Surv. 55, 332, 2000.
- [4] B. H. Bowditch, *Median and injective metric spaces*, Mathematical Proceedings of the Cambridge Philosophical Society 168:1 (2020) 43–55.
- [5] Y. Beniamini, J. Lindenstrauss, *Geometric Nonlinear Functional Analysis*, American Mathematical Society Colloquium Publications, v. 48, 2000.
- [6] I. Chiswell, *Introduction to Λ -trees*, World Scientific Publishing Co. Pte. Ltd., 2001.
- [7] S. Eilenberg, O.G. Harrold, Jr. *Continua of finite linear measure. I*, Amer. J. Math. 65 (1943), 137–146.
- [8] R. Engelking, *General topology*, Helderman Verlag Berlin, Sigma series in pure mathematics, vol. 6, 1989.
- [9] M. Espinola, M.A. Khamsi, *Introduction to hyperconvex spaces* In: Kirk, W.A., Sims, B. (eds) Handbook of Metric Fixed Point Theory. Springer, Dordrecht. DOI:10.1007/978-94-017-1748-9_13.
- [10] K. Goebel and W. A. Kirk, *Topics in Metric Fixed Point Theory*, Cambridge Univ. Press, Cambridge, 1990.
- [11] H. Hahn, *Mengentheoretische Charakterisierung der stetigen Kurven*, Sitzungsber. Akad. Wiss. Wien 123 (1914), 2433–2489.
- [12] G. E. Huhunaišvili, *On a property of Uryson’s universal metric space* (Russian), Soviet Math. Dokl., 101 (1955), 607–610.
- [13] A. Kechris, *Classical Descriptive Set Theory*, Springer, 1995.
- [14] W. Kirk, *Hyperconvexity of $\mathbb{R}$ -trees*, Fund. Math. 156 (1998), 67–72.
- [15] W. Kubiś, M. Rubin, *Extension theorems and reconstruction theorems for the Urysohn universal space*, Czechoslovak Math. J. 60:1 (2010), 1–29.
- [16] S. Mazurkiewicz, *Sur les lignes de Jordan*, Fund. Math. 1 (1920), 166–209.
- [17] J. Melleray, *On the geometry of Urysohn’s universal metric space*, Topology Appl. 154:2 (2007), 384–403.
- [18] P. S. Urysohn, *Sur un espace métrique universel*, Bull. Sci. Math 51 (1927) 43–64 and 74–96.
- [19] T. Ważewski, *Kontinua prostowalne w związku z funkcjami i odwzorowaniami absolutnie ciągłymi*, Annales de la Société Polonaise de Mathématique, supplement, vol. 3 (1937), pp. 9–49.

T. BANAKH (ORCID 0000-0001-6710-4611): IVAN FRANKO NATIONAL UNIVERSITY OF LVIV (UKRAINE), AND MATHEMATICS DEPARTMENT, JAN KOCHANOWSKI UNIVERSITY IN KIELCE, UNIWERSYTECKA 7, 25-406 KIELCE, POLAND

Email address: t.o.banakh@gmail.com

J. BĄK: (ORCID 0000-0001-8027-7226): MATHEMATICS DEPARTMENT, JAN KOCHANOWSKI UNIVERSITY IN KIELCE, UNIWERSYTECKA 7, 25-406 KIELCE, POLAND

Email address: judyta.bak@ujk.edu.pl

J. GARBULIŃSKA-WĘGRZYN (ORCID 0000-0001-7217-2002): MATHEMATICS DEPARTMENT, JAN KOCHANOWSKI UNIVERSITY IN KIELCE, UNIWERSYTECKA 7, 25-406 KIELCE, POLAND

Email address: jgarbulinska@ujk.edu.pl

M. NOWAK (ORCID 0000-0003-1915-0001): MATHEMATICS DEPARTMENT, JAN KOCHANOWSKI UNIVERSITY IN KIELCE, UNIWERSYTECKA 7, 25-406 KIELCE, POLAND

Email address: magdalena.nowak805@gmail.com

M. POPLAWSKI (ORCID 0000-0002-2725-9675): MATHEMATICS DEPARTMENT, JAN KOCHANOWSKI UNIVERSITY IN KIELCE, UNIWERSYTECKA 7, 25-406 KIELCE, POLAND

Email address: michal.poplawski.m@gmail.com

