

# SUBORDINATION INVOLVING GAUSS HYPERGEOMETRIC FUNCTION

ANISH KUMAR AND SOURAV DAS

**ABSTRACT.** The primary objective of this work is to obtain some sufficient conditions so that normalized Gauss hypergeometric function satisfies exponential starlikeness and convexity in the unit disk. Moreover, conditions on parameter of this function has been derived for being Janowski convexity and starlikeness. Furthermore, some sufficient conditions are also obtained so that Gauss hypergeometric function posses lemniscate starlikeness and convexity. Results established in this work are presumably new and their significance is illustrated by several consequences, graphical representations and examples.

## 1. INTRODUCTION AND MOTIVATION

In the last few years, generalized hypergeometric functions have sparked great interest among researchers. Mainly used by L. de. Branges [4] in 1985 into the proof of Melin conjecture which prompted into the Bieberbach conjecture. A few year ago to this proof only some literature deals with the univalent function theory for different kind of Gauss hypergeometric function, confluent hypergeometric function, other generalized hypergeometric function like Bessel function, Struve function, Lommel function using various methods. In [1, 2, 12–17], researchers derived several require conditions on the parameters for these special functions lies on geometric characteristics such as convexity, starlikeness, close-to-convexity and uniformly convex. In this article, various sufficient conditions have been obtained on the involved parameters in Gauss hypergeometric function  $F$  which satisfies geometric properties such as exponential starlikeness and convexity, Janowski starlikeness and convexity, Lemniscate starlikeness and convexity.

Let  $\mathcal{A}$  denote the normalized collection of analytic functions, which satisfy the condition  $F'(0) - 1 = F(0) = 0$ . A function  $F \in \mathcal{A}$  which is univalent in the unit disk such that  $U = \frac{zF'}{F}$  or  $U = 1 + \frac{zF''}{F'}$  lies in the domain  $|\log U| < 1$  of the right half plane connected with the exponential function. Subordination play an important character in defining these geometric properties. If  $g$  and  $h$  are two analytic function in  $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$ , then  $g$  is subordinate to  $h$  if there exists a Schwartz function  $l$  in  $\mathbb{D}$  such that  $g = hol$  in  $\mathbb{D}$ . It can be denoted as  $g \prec h$  if  $g(0) = h(0)$  and  $g(D) \subseteq h(D)$ . The Class  $\mathcal{P}_e$  contains analytic function  $p$  in  $\mathbb{D}$  with  $p(z) \prec e^z$  and  $p(0) = 1$  for every  $z \in \mathbb{D}$ . A function  $F \in \mathcal{A}$  is called exponential convex (or starlike) if  $1 + \frac{zF''(z)}{F'(z)}$  or  $\frac{zF'(z)}{F(z)}$  belong to  $\mathcal{P}_e$ . It is denoted as  $\mathcal{K}_e$  and  $\mathcal{K}_e^*$ . Ma and Minda [6] introduced particular cases of subclass of starlike  $S^*$  and convex function  $\mathcal{K}^*$ . Mathematical charecterization of  $S^*$  and  $\mathcal{K}^*$  is defined as follows:

$$\begin{aligned}\mathcal{S}^* &= \left\{ F \in \mathcal{A} : \Re \left( \frac{zF'(z)}{F(z)} \right) > 0 \quad \text{for all } z \in \mathbb{D} \right\}, \\ \mathcal{K} &= \left\{ F \in \mathcal{A} : \Re \left( 1 + \frac{zF''(z)}{F'(z)} \right) > 0 \quad \forall z \in \mathbb{D} \right\}.\end{aligned}$$

A function  $F \in \mathcal{A}$  is lemniscate convex if  $1 + \frac{zF''(z)}{F'(z)}$  lie in bounded region by right half of lemniscate of Bernouli  $\{k : |k^2 - 1| = 1\}$ . In terms of subordination lemniscate starlike and lemniscate convexity is defined as  $\frac{zF'(z)}{F(z)} \prec \sqrt{1+z}$  and  $1 + \frac{zF''(z)}{F'(z)} \prec \sqrt{1+z}$  respectively. For  $-1 \leq D < C \leq 1$ , suppose that  $P[C, D]$  be the class containing normalized analytic function

---

2020 *Mathematics Subject Classification.* 30D15; 30C45; 30H10.

*Key words and phrases.* Fox-Wright functions, Analytic functions, Univalent functions, Convex functions, Starlike functions, Hardy spaces.

$p(z) = 1 + c_1(z) + \dots$  in  $\mathbb{D}$  satisfying  $p(z) \prec \frac{1+Cz}{1+Dz}$ . For  $0 \leq \beta < 1$ , then  $P[1 - 2\beta, -1]$  is the class of function  $1 + c_1(z) + \dots$  holds  $\Re(p(z)) > \beta$  in  $\mathbb{D}$ . The class  $S^*[C, D]$  of Janowski Starlike function [5] if  $F \in \mathcal{A}$  satisfy  $\frac{zf'(z)}{F(z)} \in P[C, D]$ .

Assume  $w \neq 0, -1, -2, \dots$  then function

$$F(u, v; w; z) = \sum_{n=0}^{\infty} \frac{(u)_n(v)_n}{(w)_n} \frac{z^n}{n!}$$

is said to be Gauss hypergeometric function is analytic in  $\mathbb{D}$  and satisfy the differential equations  $z(1-z)F'' + [w - (u+v+1)z]F' - uvF = 0$ .  $F$  holds some recurrence relation [8] as:

$$\begin{aligned} F(u, v; w; z) &= F(v, u; w; z), \\ wF'(u, v; w; z) &= uvF(u+1, v+1; w+1; z), \\ F(u, v; w; z) &= (1-z)^{w-u-v}F(w-u, w-v; w; z). \end{aligned}$$

The paper is systemized as follows. In Section 2, we have mentioned some useful lemmas, which will be helpful to derive our key results. Section 3 discusses some sufficient conditions so that the Gauss hypergeometric function possesses such as exponential starlikeness and convexity. Moreover, some consequences, important remarks and graphical representations have been shown in this section. In Section 4, we have considered normalized Gauss hypergeometric function for which Janowski starlikeness and convexity have been discussed. Furthermore some sufficient conditions are established in Section 5, so that this function belong lemniscate starlikeness and convexity. Consequence and remarks related to this have also been shown in this section.

## 2. USEFUL LEMMAS

Some Lemmas have been recollected in the below section, which will be useful to show our main results.

**Lemma 1.** [10] Let  $\Omega$  be a subset of  $\mathbb{C}$  and the function  $\psi : \mathbb{C}^3 \times \mathbb{D} \rightarrow \mathbb{C}$  holds the condition  $\psi(r, s, t; z) \notin \Omega$  whenever:  $s = me^{i\theta}e^{e^{i\theta}}, r = e^{e^{i\theta}}, \Re((s+t)e^{-i\theta}e^{-e^{i\theta}}) \geq 0, \theta \in [0, 2\pi], z \in \mathbb{D}$  and  $1 \leq m$ . If  $q$  is an analytic function in  $\mathbb{D}$  with  $q(0) = 1$  and  $\psi(q(z), zq'(z), z^2q''(z); z) \in \Omega$  for  $z \in \mathbb{D}$ , then  $q \in \mathcal{P}_e$ .

It can be noted that the admissibility condition  $\psi(r, s, t; z) \notin \Omega$  is verified for all:  $s = me^{i\theta}e^{e^{i\theta}}, r = e^{e^{i\theta}}, \Re((s+t)e^{-i\theta}e^{-e^{i\theta}}) \geq 0, \theta \in [0, 2\pi], z \in \mathbb{D}$  and  $1 \leq m$ .

Furthermore for  $\psi : \mathbb{C}^2 \times \mathbb{D} \rightarrow \mathbb{C}$  the admissibility condition convert into  $\psi(e^{e^{i\theta}}, me^{i\theta}e^{e^{i\theta}}) \notin \Omega$ , where  $z \in \mathbb{D}, \theta \in [0, 2\pi)$  and  $m \geq 1$ .

**Lemma 2.** [8, 9] Suppose that  $\Omega \subset \mathbb{C}$  and  $\psi : \mathbb{C}^3 \times \mathbb{D} \rightarrow \mathbb{C}$  hold  $\psi(i\rho, \sigma, \mu + iv; z) \notin \Omega$  when  $z \in \mathbb{D}$ ,  $\rho$  real  $\sigma \leq -\frac{1+\rho^2}{2}$  and  $\sigma + \mu \leq 0$ . If  $q$  is analytic in  $\mathbb{D}$  with  $q(0) = 1$  and  $\psi(q(z), zq'(z), z^2q''(z); z) \in \Omega$  for  $z \in \mathbb{D}$ , then  $\Re(q(z)) > 0$  in  $\mathbb{D}$ .

In this case  $\psi : \mathbb{C}^2 \times \mathbb{D} \rightarrow \mathbb{C}$ , then the condition Lemma 2 convert into  $\psi(i\rho, \sigma; z) \notin \Omega$  for  $\rho$  real and  $\sigma \leq -\frac{1+\rho^2}{2}$ .

**Lemma 3.** [7] Assume  $q$  in normalized class of analytic function with  $q(z) \neq 1$ . Let  $\Omega \in \mathbb{C}$  and  $\psi : \mathbb{C}^3 \times \mathbb{D} \rightarrow \mathbb{C}$  hold  $\psi(i\rho, \sigma, \mu + iv; z) \notin \Omega$  when  $z \in \mathbb{D}, r = \sqrt{2 \cos 2\theta}e^{i\theta}, s = \frac{me^{3i\theta}}{2\sqrt{2 \cos 2\theta}}, \Re((s+t)e^{-3i\theta}) \geq \frac{3m^2}{8\sqrt{2 \cos 2\theta}}, m \geq n \geq 1$  and  $-\frac{\pi}{4} < \theta < \frac{\pi}{4}$ . If  $(q(z), zq'(z), z^2q''(z); z) \in \mathbb{D}$  for  $z \in \mathbb{D}$  and  $\psi(q(z), zq'(z), z^2q''(z); z) \in \Omega, z \in \mathbb{D}$  then  $q(z) \prec \sqrt{1+z}$ .

In this case  $\psi : \mathbb{C}^2 \times \mathbb{D} \rightarrow \mathbb{C}$ , the condition in Lemma 3, reduces to  $\psi(r, s; z)$  whenever  $r = \sqrt{2 \cos 2\theta}e^{i\theta}, s = \frac{me^{3i\theta}}{2\sqrt{2 \cos 2\theta}}, m \geq n \geq 1$  and  $-\frac{\pi}{4} < \theta < \frac{\pi}{4}$  and  $z \in \mathbb{D}$ .

### 3. EXPONENTIAL STARLIKENESS AND CONVEXITY OF GAUSS HYPERGEOMETRIC FUNCTION

In this section some of main results related exponential starlikeness and convexity has been discussed.

**Theorem 1.** Assume that the parameters  $u, v \in \mathbb{R}$ ,  $w \neq 0, -1, -2, \dots$  be constrained and  $u, v$  and  $w$  also fulfill the condition:  $\frac{1}{e}(\Re(w - 1 - u - v)) - |(u + v - 1 + (u - 1)(v - 1))e| \geq 0$ , then  $F(u, v; w; z) \in \mathcal{P}_e$ .

*Proof.* Let us assume a function  $q: \mathbb{D} \rightarrow \mathbb{C}$

$$q(z) = \frac{w}{uv} F'(u, v; w; z).$$

Now, we claim that  $\frac{w}{uv} F'(u, v; w; z) \in \mathcal{P}_e$ . Since  $F(u, v; w; z)$  satisfy the differential equation  $z(1-z)F''(u, v; w; z) + [w - (u+v+1)z]F'(u, v; w; z) - uvF(u, v; w; z) = 0$ . Consequently,

$$\begin{aligned} & z(1-z)F'''(z) + [w + 1 - (u + v + 3)z]F'' - (u + v + 1 + uv)F'(z) \\ &= z(1-z)\frac{uv}{w}q''(z) + \frac{uv}{w}[w + 1 - (u + v + 3)z]q'(z) - \frac{uv}{w}(u + v + 1 + uv)q(z) = 0. \end{aligned} \quad (3.1)$$

Now, multiplying both sides by  $z$  in (3.1), we have

$$(1-z)z^2q''(z) + [w + 1 - (u + v + 3)z]zq'(z) - (u + v + 1 + uv)zq(z) = 0.$$

Further, we define another function  $\psi: \mathbb{C}^3 \times \mathbb{D} \rightarrow \mathbb{C}$  by :

$$\psi(r, s, t; z) = (1-z)t + [w + 1 - (u + v + 3)z]s - (u + v + uv + 1)zr = 0, \quad (3.2)$$

and suppose  $\Omega = 0$ , then differential equation (3.2) can be rewritten as:

$$\psi(q(z), zq'(z), z^2q''(z); z) = \Omega.$$

To show that  $q(z) \prec e^z$ , we have to prove  $\psi(r, s, t; z) \neq \Omega$ , whenever  $s = me^{i\theta}e^{e^{i\theta}}, r = e^{e^{i\theta}}$ ,  $\Re((s+t)e^{-i\theta}e^{-e^{i\theta}}) \geq 0$ ,  $\theta \in [0, 2\pi]$ ,  $z \in \mathbb{D}$  and  $1 \leq m$  taking into account Lemma 1 and using triangle inequality. We can observe that,

$$\begin{aligned} |\psi(r, s, t; z)| &= |(1-z)(t+s) + [w - (u+v+2)z]s - (u+v+uv+1)zr| \\ &\geq |e^{e^{i\theta}}e^{i\theta}| |(1-z)(t+s)e^{-e^{i\theta}e^{-i\theta}} + [w - (u+v+2)z]m| - |(u+v+uv+1)ze^{e^{i\theta}}| \\ &\geq e^{\cos\theta} ((1-x)\Re((s+t)e^{-i\theta}e^{-e^{i\theta}}) + \Re(w - (u+v+2)z)m) - |(u+v+uv+1)e| \\ &\geq \frac{1}{e}(\Re(w - (u+v+2)z)) - |(u+v+uv+1)e|. \end{aligned} \quad (3.3)$$

Since,  $\Re(z) < 1$  and  $\frac{1}{e}(\Re(w - 2 - u - v)) - |(u + v + uv + 1)e| \geq 0$ , (3.3) yields

$$|\psi(r, s, t; z)| > \frac{1}{e}(\Re(w - 2 - u - v)) - |(u + v + uv + 1)e| \geq 0. \quad (3.4)$$



This shows that  $|\psi(r, s, t; z)| > 0$ , hence  $q(z) \prec e^z$  for all  $z \in \mathbb{D}$ . Therefore we have proved that  $\frac{w}{uv} F'(u, v; w; z) \in \mathcal{P}_e$  under the given hypothesis. Using relation  $\frac{w}{uv} F'(u, v; w; z) = F(w+1, u+1; v+1; z)$ , one can get  $F(u, v; w; z) \in \mathcal{P}_e$  under the requirement  $\frac{1}{e}(\Re(w - 1 - u - v)) - |(u + v - 1 + (u - 1)(v - 1))e| \geq 0$ . This completes the proof of the theorem.  $\square$

**Example 1.**  $F(-2, 1, 5)$  and  $F(-2, 1, 3) \in \mathcal{P}_e$

In the below theorem sufficient condition on parameter  $u, v$  and  $w$  have been established such that Gauss hypergeometric function  $F(u, v; w; z)$  is exponential convex.

**Theorem 2.** Suppose that the parameters  $u, v \in \mathbb{R}$  and  $w \neq 0, -1, -2, -3, \dots$  be constrained and  $u, v; w$  also hold the condition :

$$\Re\left(\frac{w-2}{e}\right) - |uv| - |u+v||e| - |1-w| > 0, \quad (3.5)$$

FIGURE 1. For  $u = -2, v = 1, w = 5$ FIGURE 2. For  $u = -2, v = 1, w = 3$ 

then the function  $\Delta(u, v; w; z) \in \mathcal{K}_e$ .

*Proof.* Let  $\Delta(u, v; w; z) = (F(u, v; w; z) - 1) \frac{w}{uv}$  and define a function  $p : \mathbb{D} \rightarrow \mathbb{C}$  as:

$$p(z) = 1 + \frac{z\Delta''(u, v; w; z)}{\Delta'(u, v; w; z)} = 1 + \frac{zF''(u, v; w; z)}{F'(u, v; w; z)}. \quad (3.6)$$

Assume that  $F'(u, v; w; z) \neq 0$  and since  $F(u, v; w; z)$  satisfy the differential equation  $z(1 - z)F''(u, v; w; z) + [w - (u + v + 1)z]F'(u, v; w; z) - uvF(u, v; w; z) = 0$ .

Differentiation both sides, we get

$$\begin{aligned} &z(1 - z)F'''(u, v; w; z) + (1 - 2z)F''(u, v; w; z) + [w - (u + v + 1)z]F''(u, v; w; z) \\ &- (u + v + 1 + uv)F'(u, v; w; z) = 0. \end{aligned} \quad (3.7)$$

Dividing both sides by  $F'(u, v; w; z)$  and multiplying by  $z$  in (3.7), we have

$$\begin{aligned} z^2(1-z) \frac{F'''(u, v; w; z)}{F'(u, v; w; z)} + (1-2z) \frac{zF''(u, v; w; z)}{F'(u, v; w; z)} + [w - (u+v+1)z] \frac{zF''(u, v; w; z)}{F'(u, v; w; z)} \\ - (u+v+1+uv)z = 0. \end{aligned} \quad (3.8)$$

By derivative of the equation (3.6)  $(p(z)-1)F'(u, v; w; z) = zF''(u, v; w; z)$ , we obtain

$$\begin{aligned} (p(z)-1)F''(u, v; w; z) + p'(z)F'(u, v; w; z) &= zF'''(u, v; w; z) + F''(u, v; w; z) \\ \text{and } (q(z)-1) \frac{zF''(u, v; w; z)}{F'(u, v; w; z)} + zp'(z) &= z^2 \frac{F'''(u, v; w; z)}{F'(u, v; w; z)} + \frac{zF''(u, v; w; z)}{F'(u, v; w; z)}. \end{aligned}$$

So that,

$$(q(z)-1)^2 - (q(z)-1) + zq'(z) = z^2 \frac{F'''(u, v; w; z)}{F'(u, v; w; z)}.$$

Now, substituting all the values in (3.8),  $q(z)$  satisfies the first order differential equation

$$\begin{aligned} (1-z)((q(z)-1)^2 - (q(z)-1) + zq'(z)) + (1-2z)(q(z)-1) + [w - (u+v+1)z](q(z)-1) \\ - (u+v+1+uv)z \\ = (1-z)(q^2(z) - 3q(z) + 2 + zq'(z)) + (1-2z)(q(z)-1) + [w - (u+v+1)z](q(z)-1) \\ - (u+v+1+uv)z \\ = (1-z)zq'(z) + (1-z)q^2(z) + (w-2-(u+v)z)q(z) + (1-w-uvz) = 0. \end{aligned} \quad (3.9)$$

The above equation is also satisfies for  $z = 0$ . Further we define a function  $\psi : \mathbb{C}^2 \times \mathbb{D} \rightarrow \mathbb{C}$  by :

$$\psi(r, s; z) = (1-z)s + (1-z)r^2 + (w-2-(u+v)z)r + (1-w-uvz) = 0$$

and suppose  $\Omega = 0$ , then differential equation (3.9) can be rewritten as:

$$\psi(q(z), zq'(z), z) = \Omega.$$

Now, we can observe that for  $s = me^{i\theta}e^{i\theta}, r = e^{i\theta}, \theta \in [0, 2\pi], z \in \mathbb{D}$  and  $m \geq 1$ , we have

$$\begin{aligned} |\psi(r, s; z)| &= |(s+r^2) + (w-2)r + (1-w) - (s+r^2)z - (u+v)zr - uvz| \\ &\geq |(w-2)r + (1-z)(s+r^2)| - |z||uv| - |u+v||rz| - |1-w| \\ &\geq \Re((w-2)r) + \Re(1-z)(s+r^2) - |uv| - |u+v||e| - |1-w| \\ &\geq \Re\left(\frac{w-2}{e}\right) + (1-x)\Re(s+r^2) - |uv| - |u+v||v| - |1-w| \\ &\geq \Re\left(\frac{w-2}{e}\right) - |uv| - |u+v||e| - |1-w|, \quad \theta \in [0, 2\pi]. \end{aligned} \quad (3.10)$$

Now, by using (3.10) and above conditions

$$|\psi(r, s, z)| \geq \Re\left(\frac{w-2}{e}\right) - |uv| - |u+v||e| - |1-w| > 0.$$

whenever given hypothesis (3.5) holds,  $\psi(r, s; z) > 0$ . Therefore, by Lemma 1,  $\Delta(u, v; w; z) = (F(u, v; w; z) - 1) \frac{w}{uv} \in \mathcal{K}_e$ .  $\square$

Applying the duality theorem between two classes  $\mathcal{P}_e$  and  $\mathcal{K}_e$ , which state as :  $F \in \mathcal{K}_e$  if and only if  $zF' \in \mathcal{P}_e$ . we have the property  $wF'(u, v; w; z) = uvF(u+1, v+1; w+1; z)$ , sufficient conditions for  $zF(u, v; w; z)$  is exponential starlike can be obtained.

**Corollary 1.** Let the parameters  $u, v \in \mathbb{R}$  and  $w \neq 0, -1, -2, -3, \dots$  be constrained and  $u, v$  and  $w$  satisfy the condition :

$\Re\left(\frac{w-3}{e}\right) - |(u-1)(v-1)| - |u+v-2||e| - |2-w| > 0$ , then the function  $zF(u, v; w; z) \in \mathcal{P}_e^*$  or  $zF(u, v; w; z)$  is exponential starlike function.

In the next section, we have obtained sufficient conditions such that Gauss hypergeometric function  $\phi(u, v; w; z)$  belong to Janowski convexity. Further we will show that  $z\phi(u, v; w; z)$  belong to Janowski starlikeness  $S^*[C, D]$ .

#### 4. JANOWSKI STARLIKENESS AND CONVEXITY OF GAUSS HYPERGEOMETRIC FUNCTION

**Theorem 3.** *Let parameters  $u, v$  and  $w \neq 0, -1, -2, \dots$  be constrained such that it satisfy the conditions for  $-1 \leq D < C \leq 1$ ,*

$$1 + C - D + f(1 + D) - \left| 1 + C - D + (c + d + 2)(1 + D) + \frac{(u + v + uv + 1)(D + 1)^2}{C - D} \right| > 0, \quad (4.1)$$

further

$$\begin{aligned} & (2 + \frac{w(1 + D)}{2} + \frac{w(D - 1)}{2} - \left( (D - C) - (u + v + 2)D + \frac{(u + v + uv + 1)(1 - D^2)}{D - C} \right)^2) \\ & 2 \left( \left( (D - C) - (u + v + 2)D - \frac{(u + v + uv + 1)(1 - D^2)}{C - D} \right)^2 \right) + \\ & \left( \frac{(u + v + uv + 1)}{2(C - D)}(1 - D)^2 - 1 - \frac{(u + v + 2)(1 + D)}{2} - \frac{(u + v + uv + 1)(D + 1)^2}{2(D - C)} + \right. \\ & \left. \left( -1 - \frac{[(u + v + 2)]}{2}(D - 1) \right) \right)^2 \geq 0, \end{aligned} \quad (4.2)$$

whenever,

$$\begin{aligned} & \left( \frac{(u + v + uv + 1)}{2(C - D)}(1 - D)^2 - 1 - \frac{(u + v + 2)(1 + D)}{2} - \frac{(u + v + uv + 1)(D + 1)^2}{2(C - D)} \right) \\ & + \left( -1 - \frac{[(u + v + 2)]}{2}(D - 1) \right) \\ & \geq \left( (D - C) - (u + v + 2)D - \frac{(u + v + uv + 1)(1 - D^2)}{C - D} \right)^2. \end{aligned} \quad (4.3)$$

another inequality

$$\begin{aligned} & 2 \left( \left( (D - C) - (u + v + 2)D - \frac{(u + v + uv + 1)(1 - D^2)}{C - D} \right)^2 \right) + \\ & \left( \frac{(u + v + uv + 1)}{2(D - C)}(1 - D)^2 - 1 - \frac{(u + v + 2)(1 + D)}{2} - \frac{(u + v + uv + 1)(D + 1)^2}{2(C - D)} \right) \\ & + \left( -1 - \frac{[(u + v + 2)]}{2}(D - 1) \right) \geq 0, \end{aligned} \quad (4.4)$$

whenever

$$\begin{aligned} & \left( \frac{(u + v + uv + 1)}{2(C - D)}(1 - D)^2 - 1 - \frac{(u + v + 2)(1 + D)}{2} - \frac{(u + v + uv + 1)(D + 1)^2}{2(D - C)} \right) \\ & + \left( -1 - \frac{[(u + v + 2)]}{2}(D - 1) \right) \\ & \leq \left( (D - C) - (u + v + 2)D - \frac{(u + v + uv + 1)(1 - D^2)}{C - D} \right)^2. \end{aligned} \quad (4.5)$$

If  $0 \notin \phi'((D))$  and  $0 \notin \phi''((D))$ , then

$$1 + z \frac{\phi''(z)}{\phi'(z)} \prec \frac{1 + Cz}{1 - Dz}.$$

*Proof.* We define a function  $q : \mathcal{D} \rightarrow \mathbb{C}$

$$q(z) = \frac{(C - D)\phi'(z) + (1 - D)z\phi''(z)}{(C - D)\phi'(z) - (1 + D)z\phi''(z)},$$

Then

$$\frac{z\phi''(z)}{\phi'(z)} = \frac{(C - D)(q(z) - 1)}{(q(z) + 1) + D(q(z) - 1)}.$$

Taking logarithmic derivative both sides,

$$\begin{aligned} \frac{1}{z} + \frac{\phi'''(z)}{\phi''(z)} - \frac{\phi''(z)}{\phi'(z)} &= \frac{q'(z)}{q(z) - 1} - \frac{q'(z)(D + 1)}{(q(z) + 1) + D(q(z) - 1)} \\ &= 1 + \frac{z\phi'''(z)}{\phi''(z)} - \frac{z\phi''(z)}{\phi'(z)} = \frac{zq'(z)}{q(z) - 1} - \frac{zq'(z)(D + 1)}{(q(z) + 1) + D(q(z) - 1)}. \end{aligned} \quad (4.6)$$

From (4.6), we get

$$\begin{aligned} \frac{z\phi'''(z)}{\phi''(z)} &= \frac{2zq'(z)}{(q(z) - 1)((q(z) + 1) + D(q(z) - 1))} - 1 + \frac{z\phi''(z)}{\phi'(z)} \text{ or,} \\ \frac{z\phi'''(z)}{\phi''(z)} \frac{z\phi''(z)}{\phi'(z)} &= \frac{2(C - D)(q(z) - 1)zq'(z)}{(q(z) - 1)((q(z) + 1) + D(q(z) - 1))^2} - \frac{(C - D)(q(z) - 1)}{(q(z) + 1) + D(q(z) - 1)} \\ &+ \frac{(C - D)^2(q(z) - 1)^2}{((q(z) + 1) + D(q(z) - 1))^2}. \end{aligned}$$

Differentiating both sides the differential equation  $z(1-z)\phi''(u, v; w; z) + [w - (u+v+1)z]\phi'(u, v; w; z) - uv\phi(u, v; w; z) = 0$  and followed by dividing by  $\phi'(z)$  and multiplying  $z$ , we have

$$\begin{aligned}
 & (1-2z)\phi''(z) + z(1-z)\phi'''(z) + [w - (u+v+1)z]\phi''(z) - (u+v+1)\phi'(z) - uv\phi'(z) \\
 &= z(1-z)\frac{z\phi'''(z)}{\phi'(z)} + [w + 1 - (u+v+3)z]\frac{z\phi''(z)}{\phi'(z)} - (u+v+uv+1)z \\
 &= (1-z)\frac{z\phi'''(z)}{\phi''(z)}\frac{z\phi''(z)}{\phi'(z)} + [w + 1 - (u+v+3)z]\frac{z\phi''(z)}{\phi'(z)} - (u+v+uv+1)z \\
 &= (1-z)\left(\frac{2(C-D)(q(z)-1)zq'(z)}{(q(z)-1)((q(z)+1)+B(q(z)-1))^2} - \frac{(D-C)(q(z)-1)}{(q(z)+1)+D(q(z)-1)}\right. \\
 &\quad \left.+ \frac{(C-D)^2(q(z)-1)^2}{((q(z)+1)+D(q(z)-1))^2}\right) + [1+w-(w+u+3)z]\frac{(C-D)(q(z)-1)}{(q(z)+1)+D(q(z)-1)} \\
 &\quad - (u+v+uv+1)z \\
 &= \frac{2(C-D)(1-z)zq'(z)}{(q(z)+1)+D(q(z)-1))^2} + \frac{(C-D)^2(1-z)(q(z)-1)^2}{(q(z)+1)+D(Q(z)-1))^2} - (u+v+uv+1)z \\
 &\quad + [w - (u+v+2)z]\frac{(C-D)(q(z)-1)}{(q(z)+1)+D(q(z)-1)} \\
 &= (1-z)zq'(z) + \frac{(C-D)}{2}(1-z)(q(z)^2 + 1 - 2q(z)) + \frac{w - (u+v+2)z}{2}((q(z))^2 - 1) \\
 &\quad + D((q(z))^2 + 1 - 2q(z)) - \frac{(u+v+uv+1)z}{2(C-D)}(q(z)^2(D+1)^2 + (1-D)^2 + 2q(z)(1-D^2)) \\
 &= (1-z)zq'(z) \\
 &\quad + \left(\frac{C-D}{2}(1-z) + \frac{[w - (u+v+2)z]}{2}(1+D) - \frac{(u+v+uv+1)}{2(C-D)}z(1+D)^2\right)q(z)^2 \\
 &\quad + \left[(C-D)(z-1) - [w - (u+v+2)z]D - \frac{u+v+uv+1}{C-D}(1-D^2)z\right]q(z) \\
 &\quad + \frac{C-D}{2}(1-z) + \frac{[w - (u+v+2)z]}{2}(D-1) - \frac{(u+v+uv+1)}{2(C-D)}z(1-D)^2 = 0. \tag{4.7}
 \end{aligned}$$

Let us consider a function  $\psi(q(z), zq'(z); z) = F_1zq'(z) + F_2(q(z))^2 + F_3q(z) + F_4$ , where  $F_1 = 1-z$ ,

$$\begin{aligned}
 F_2 &= \left(\frac{C-D}{2}(1-z) + \frac{[w - (u+v+2)z]}{2}(1+D) - \frac{(u+v+uv+1)}{2(C-D)}z(1+D)^2\right), \\
 F_3 &= \left[(C-D)(z-1) - [w - (u+v+2)z]D - \frac{u+v+uv+1}{C-D}(1-D^2)z\right], \\
 F_4 &= \frac{C-D}{2}(1-z) + \frac{[w - (u+v+2)z]}{2}(D-1) - \frac{(u+v+uv+1)}{2(C-D)}z(1-D)^2.
 \end{aligned}$$

With the help of (4.7),  $\Omega = 0$  imply that  $\psi(q(z), zq'(z); z) \in \Omega$ . Assume that  $G_1 = \Re(F_1) = 1-x$ ,  $G_2 =$

$$\begin{aligned}
 \Re(F_2) &= \frac{C-D}{2}(1-x) + \frac{w(1+D)}{2} - \frac{(u+v+2)x(1+D)}{2} - \frac{(u+v+uv+1)x(D+1)^2}{2(C-D)}, \\
 G_3 &= \Re(iF_3) = (C-D)(-y) + (u+v+2)(-y)D + \frac{(u+v+uv+1)(1-D^2)}{C-D}y, \\
 G_4 &= \frac{C-D}{2}(1-x) + \frac{[w - (u+v+2)x]}{2}(D-1) - \frac{(u+v+uv+1)}{2(C-D)}x(1-D)^2.
 \end{aligned}$$

For  $\sigma \leq -\frac{(1+\rho^2)}{2}$ ,  $\rho \in \mathbb{R}$ .

$$\begin{aligned}
\Re\psi(i\rho, \sigma; z) &= G_1\sigma + G_2(i\rho)^2 + G_3\rho + G_4 \\
&= G_1\sigma - G_2(\rho)^2 + G_3\rho + G_4 \\
&\leq G_1\left(\frac{-(1+\rho^2)}{2}\right) - G_2\rho^2 + G_3\rho + G_4 \\
&\leq \frac{-G_1}{2} - \frac{G_1\rho^2}{2} - G_2\rho^2 + G_3\rho + G_4 \\
&= \frac{-G_1 - G_1\rho^2 - 2G_2\rho^2 + 2G_3\rho + 2G_4}{2} \\
&= \frac{-1}{2}[(G_1 + 2G_2)\rho^2 - 2G_3\rho + G_1 - 2G_4] = Q(\rho).
\end{aligned}$$

By given hypothesis (4.1)  $(G_1 + 2G_2) > 0$ ,  $Q(\rho)$  has maximum value at  $\rho = \frac{G_3}{G_1 + 2G_2}$ . Now finding value of  $Q(\rho)$  at  $\rho = \frac{G_3}{G_1 + 2G_2}$ , for all  $\rho, |x|, |y| < 1$ , we have

$$\begin{aligned}
Q(\rho) &= \frac{-1}{2}[(G_1 + 2G_2)\left(\frac{G_3}{G_1 + 2G_2}\right)^2 - 2G_3\left(\frac{G_3}{G_1 + 2G_2}\right) - 2G_4 + G_1] < 0 \\
&= \frac{-1}{2}\left[\frac{-(G_3)^2}{G_1 + 2G_2} - 2G_4 + G_1\right] \leq 0 \\
&= \frac{G_3^2}{G_1 + 2G_2} \leq G_1 - 2G_4 \\
&= G_3^2 \leq (G_1 + 2G_2)(G_1 - 2G_4).
\end{aligned}$$

For  $|x| < 1, |y| < 1$  and  $y^2 < 1 - x^2$  for above inequality, we have to prove

$$\begin{aligned}
&((C - D)(-y) + (u + v + 2)(-y)D + \frac{(u + v + uv + 1)(1 - D^2)}{C - D}y)^2 \\
&\leq (1 - x + 2(\frac{C - D}{2}(1 - x) + \frac{w(1 + D)}{2} - \frac{(u + v + 2)x(1 + D)}{2}) \\
&\quad - \frac{(u + v + uv + 1)x(D + 1)^2}{2(C - D)})(1 - x - 2(\frac{C - D}{2}(1 - x) \\
&\quad + \frac{[w - (u + v + 2)x]}{2}(D - 1) - \frac{(u + v + uv + 1)}{2(C - D)}x(1 - D)^2)) \\
&\implies ((D - C) - (u + v + 2)D + \frac{(u + v + uv + 1)(1 - D^2)}{C - D})^2(1 - x^2) \\
&\leq (1 - x + 2(\frac{C - D}{2}(1 - x) + \frac{w(1 + D)}{2} - \frac{(u + v + 2)x(1 + D)}{2}) \\
&\quad - \frac{(u + v + uv + 1)x(D + 1)^2}{2(C - D)})(1 - x - 2(\frac{C - D}{2}(1 - x) \\
&\quad + \frac{[w - (u + v + 2)x]}{2}(D - 1) - \frac{(u + v + uv + 1)}{2(C - D)}x(1 - D)^2)).
\end{aligned}$$

The above inequality yields

$H(x) = h_1(C, D, O)x^2 + h_2(C, D, O)x + h_3(C, D, O) \geq 0$ , where

$$\begin{aligned} h_1(C, D, O) &= \left( (D - C) - (u + v + 2)D - \frac{(u + v + uv + 1)(1 - D^2)}{D - C} \right)^2, \\ h_2(C, D, O) &= \left( \frac{(u + v + uv + 1)}{2(C - D)}(1 - D)^2 - 1 - \frac{(u + v + 2)(1 + D)}{2} - \frac{(u + v + uv + 1)(D + 1)^2}{2(C - D)} \right) \\ &\quad + \left( -1 - \frac{[(u + v + 2)]}{2}(D - 1) \right), \\ h_3(C, D, O) &= 2 + \frac{w(1 + D)}{2} + \frac{w(D - 1)}{2} \\ &\quad - \left( (D - C) - (u + v + 2)D + \frac{(u + v + uv + 1)(1 - D^2)}{C - D} \right)^2. \end{aligned}$$

To obtain (4.1), in first case (4.3) holds. Then  $|\frac{-h_2}{2h_1}| = |\rho| \leq 1$  and therefore  $H'(x) = 0$  at  $x = \rho \in (-1, 1)$ . Since  $H''(x) > 0$ , then minimum value at  $x = \frac{-h_2}{2h_1}$  and from (4.2) it satisfies that  $H(x) \geq H(\rho) = h_3 - \frac{h_2^2}{2h_1} \geq 0$ . Now, let (4.5) be hold. In this case  $|\rho| \geq 1$  and  $H'(x) = 2h_1(x) + h_2 \leq 2h_1 + h_2 \leq 0$  equivalent to (4.4),  $H(x)$  is decreasing. Hence  $H(x) \geq H(1) = h_1 + h_2 + h_3 \geq 0$ . By help of Lemma 2,  $\Re(p(z)) > 0$ , it is equivalent to

$$q(z) = \frac{(C - D)\phi'(z) + (1 - D)z\phi''(z)}{(C - D)\phi'(z) - (1 + D)z\phi''(z)} \prec \frac{1 + z}{1 - z}.$$

Using definition of subordination,  $\exists$  a self-map  $l(z)$  such that  $l(0) = 0$  and

$$q(z) = \frac{(C - D)\phi'(z) + (1 - D)z\phi''(z)}{(C - D)\phi'(z) - (1 + D)z\phi''(z)} = \frac{1 + l(z)}{1 - l(z)}.$$

After simplifying above equation, we get

$$1 + z \frac{\phi''(z)}{\phi'(z)} = \frac{1 + Cl(z)}{1 - Dl(z)}$$

and therefore

$$1 + z \frac{\phi''(z)}{\phi'(z)} \prec \frac{1 + Cz}{1 - Dz}.$$

Let us consider the relation, we have

$$\frac{z(z\phi(u, v; w; z))'}{z\phi(u, v; w; z)} = 1 + z \frac{\phi''(u - 1, v - 1; w - 1; z)}{\phi'(u - 1, v - 1; w - 1; z)}.$$

Simultaneously with the Theorem 3, yields the following theorem as  $z\phi(u, v; w; z) \in S^*[C, D]$ .  $\square$

In next section, we have obtained sufficient condition such that Gauss hypergeometric function belong to lemniscate starlike and convex.

## 5. LEMNISCATE CONVEXITY AND STARLIKENESS OF GAUSS HYPERGEOMETRIC FUNCTION

**Theorem 4.** *If the parameters  $u, v \in \mathbb{R}$ ,  $w \neq 0, -1, -2 \dots$  and  $\Re(1 - w) - |uv| - |u + v||\sqrt{2}| - |(w - 2)\sqrt{2}| > 0$ , then  $F(z)$  is lemniscate convex in  $\mathbb{D}$ .*

*Proof.* Let us consider a function  $p : \mathbb{D} \rightarrow \mathbb{C}$

$$q(z) = 1 + \frac{zF''(z)}{F'(z)}.$$

It's clear that  $F(z)$  is univalent in  $\mathbb{D}$  and  $F'(z) \neq 0$  for all  $z \in \mathbb{D}$ . Therefore the function  $q(z)$  is analytic in  $\mathbb{D}$  and  $q(0) = 1$ . Since  $F(z)$  satisfy the differential equation  $z(1-z)F''(u, v; w; z) + [w - (u+v+1)z]F'(u, v; w; z) - uvF(u, v; w; z) = 0$ ,  $q(z)$  satisfy the below equation by proof of the Theorem 2

$$z(1-z)q'(z) + (w-2-(u+v)z)q(z) + (1-z)q^2(z) + 1-w-uvz = 0.$$

Let  $\Omega = 0$  then  $\psi(q(z), zq'(z); z) \in \Omega$  for all  $z \in \mathbb{D}$ . Suppose that a function  $\psi$  defined as:  $\psi : \mathbb{C}^2 \times \mathbb{D} \rightarrow \mathbb{C}$  such that

$$\psi(r, s; z) = (1-z)s + (w-2-(u+v)z)r + (1-z)r^2 + 1-w-uvz = 0.$$

For  $r = \sqrt{2 \cos 2\theta} e^{i\theta}$ ,  $s = \frac{me^{3i\theta}}{2\sqrt{2 \cos 2\theta}}$ ,  $m \geq n \geq 1$  for  $\frac{-\pi}{4} < \theta < \frac{\pi}{4}$  by Lemma 3, we have

$$\begin{aligned} |\psi(r, s; z)| &= |(s+r^2) + (w-2-(u+v)z)r - z(s+r^2) + 1-w-uvz| \\ &\geq |(1-w) + (1-z)(s+r^2)| - |z||uv| - |u+v||rz| - |(w-2)r| \\ &\geq \Re(1-w) + \Re(1-z)(s+r^2) - |uv| - |u+v||\sqrt{2}| - |w-2|r| \\ &\geq \Re(1-w) + (1-x)\Re(s+r^2) - |uv| - |u+v||\sqrt{2}| - |(w-2)\sqrt{2}| \\ &\geq \Re(1-w) - |uv| - |u+v||\sqrt{2}| - |(w-2)\sqrt{2}|. \end{aligned}$$

$$\implies F(z) \prec \sqrt{1+z}.$$

Under the given hypothesis proof of this theorem readily follow.  $\square$

Applying the result, which state as:  $F$  is lemniscate convex imply that  $zF'$  is lemniscate starlike. we have the property  $wF'(u, v; w; z) = uvF(u+1, v+1; w+1; z)$ , sufficient conditions for  $zF(u, v; w; z)$  is lemniscate starlike can be obtained.

**Corollary 2.** *Let the parameters  $u, v \in \mathbb{R}$  and  $w \neq 0, -1, -2, -3, \dots$  be constrained and  $u, v$  and  $w$  satisfy the condition :*

*$\Re(2-w) - |(u-1)(v-1)| - |u+v-2||\sqrt{2}| - |(w-2)\sqrt{2}| > 0$ , then the function  $zF(u, v; w; z)$  is lemniscate starlike function.*

## 6. CONCLUSION

The Investigation successfully established new and significant conditions for the normalized Gauss hypergeometric function, determining its geometric properties concerning exponential starlikeness, convexity, Janowski convexity, and starlikeness based on specific parameter values. The findings also extended to include lemniscate starlikeness and convexity using subordination. The implications of these results were highlighted through various consequences, graphical representations, and illustrative examples, underscoring the importance and novelty of these established conditions. These contributions add valuable insights to the understanding of the Gauss hypergeometric function's behavior within the unit disk, enriching the field of mathematical analysis.

## REFERENCES

- [1] R. M. Ali, S. R. Mondal, V. Ravichandran, On the Janowski convexity and starlikeness of the confluent hypergeometric function. *Bull. Belg. Math. Soc. Simon Stevin* 22(2), 227–250 (2015).
- [2] A. Baricz, N. Yȧgmur, Geometric properties of some Lommel and Struve functions, *Ramanujan J.* 42(2), 325–346 (2017).
- [3] N. Bohra, V. Ravichandran, On confluent hypergeometric functions and generalized Bessel functions. *Anal. Math.* 43(4), 533–545 (2017).
- [4] L. de Branges, A proof of the Bieberbach conjecture, *Acta Math.* 154 (1985), no. 1-2, 137–152.
- [5] W. Janowski, Some extremal problems for certain families of analytic functions. I. *Ann. Polon. Math.* 28 (1973), 297–326.
- [6] W. C. Ma and D. Minda, A unified treatment of some special classes of univalent functions, in *Proceedings of the Conference on Complex Analysis (Tianjin, 1992)*, 157–169, *Conf. Proc. Lecture Notes Anal.*, I, Int. Press, Cambridge, MA.

- [7] V. Madaan, A. Kumar and V. Ravichandran, Starlikeness associated with lemniscate of Bernoulli, *Filomat* 33 (2019), no. 7, 1937–1955.
- [8] S. S. Miller and P. T. Mocanu, Differential subordinations and inequalities in the complex plane, *J. Differential Equations* 67 (1987), no. 2, 199–211.
- [9] S. S. Miller and P. T. Mocanu, Differential subordinations, Monographs and Textbooks in Pure and Applied Mathematics, 225, Marcel Dekker, Inc., New York, 2000.
- [10] A. Naz, S. Nagpal and V. Ravichandran, Star-likeness associated with the exponential function, *Turkish J. Math.* 43 (2019), no. 3, 1353–1371.
- [11] A. Naz, S. Nagpal and V. Ravichandran, Exponential starlikeness and convexity of confluent hypergeometric, Lommel, and Struve functions, *Mediterranean Journal of Mathematics* 17, 6 (2020), 204.
- [12] S. Noreen, M. Raza, E. Deniz, S. Kazimōglu, On the Janowski class of generalized Struve functions. *Afr. Mat.* 30(1–2), 23–35 (2019).
- [13] H. Orhan, N. Yagmur, Geometric properties of generalized Struve functions. *An. S̄tiint̄. Univ. Al. I. Cuza Iāsi. Mat. (N.S.)* 63(2), 229–244 (2017).
- [14] S. Owa, H. M. Srivastava, Univalent and starlike generalized hypergeometric functions. *Can. J. Math.* 39(5), 1057–1077 (1987).
- [15] Y. Sim, O. Kwon, N. E. Cho, Geometric properties of Lommel functions of the first kind. *Symmetry* 10, 455 (2018).
- [16] N. Yāgmur, Hardy space of Lommel functions. *Bull. Korean Math. Soc.* 52(3), 1035–1046 (2015).
- [17] N. Yāgmur, H. Orhan, Starlikeness and convexity of generalized Struve functions. *Abstr. Appl. Anal.* 2013, 6 (2013).

**Anish Kumar**

DEPARTMENT OF MATHEMATICS, NATIONAL INSTITUTE OF TECHNOLOGY JAMSHEDPUR,  
JAMSHEDPUR 831014, JHARKHAND, INDIA

*Email address:* ak8107690@gmail.com

**Sourav Das**

DEPARTMENT OF MATHEMATICS, NATIONAL INSTITUTE OF TECHNOLOGY JAMSHEDPUR,  
JAMSHEDPUR 831014, JHARKHAND, INDIA

*Email address:* souravdasmath@gmail.com, souravdas.math@nitjsr.ac.in