Integrating Molecular Dynamics Simulations and Experimental Data for **Azeotrope Predictions in Binary Mixtures**

Xiaoyu Wang^{1, a)} and Bingqing Cheng^{2, 1, b)}

(Dated: 6 May 2024)

An azeotrope is a constant boiling point mixture, and its behavior is important for fluid separation processes. Predicting azeotropes from atomistic simulations is difficult, due to the complexities and convergence problems in Monte Carlo and free-energy perturbation techniques. Here, we present a methodology for predicting the azeotropes of binary mixtures, which computes the compositional dependence of chemical potentials from molecular dynamics simulations using the S0 method, and employs experimental boiling point and vaporization enthalpy data. Using this methodology, we reproduce the azeotropes or the lack of in five case studies, including ethanol/water, ethanol/isooctane, methanol/water, hydrazine/water, and acetone/chloroform mixtures. We find that it is crucial to use the experimental boiling point and vaporization enthalpy for reliable azeotrope predictions, as empirical force fields are not accurate enough for these quantities. Finally, we use regular solution models to rationalize the azeotropes, and reveal that they tend to form when the mixture components have similar boiling points and strong interactions.

I. INTRODUCTION

An azeotrope is a special liquid mixture of two or more substances; it maintains identical compositions in both the liquid and the vapor phases upon vaporization. Studies reveal that about half of binary mixtures exhibit azeotropic behaviors^{1,2} and their characteristics impact the chemical industry extensively.³ For example, azeotropes play a crucial role in the purification process, such as alcohol distillation. Azeotropes have also been widely exploited as flammable azeotropic fuels to improve vehicle engine performance^{4,5} by altering boiling points and compositions, as well as sustainable multipurpose solutions in vapor phase cleaning processes.^{6,7}

Thermodynamically, one can determine the azeotrope in a binary mixture by satisfying two conditions:

$$x_i^{\mathcal{L}} = x_i^{\mathcal{V}},\tag{1}$$

$$x_{i}^{L} = x_{i}^{V}, \tag{1}$$

$$\mu_{i}^{L}(T, P, x_{i}^{L}) = \mu_{i}^{V}(T, P, x_{i}^{V}), \tag{2}$$

 $x_i^{\rm V}$ and $x_i^{\rm L}$ are the mole fraction of each component, i, in the vapor and the liquid phase, respectively, and Eq. 1 states that they must be identical. Simultaneously, Eq.2 outlines that an azeotrope forms when the chemical potentials of each component i in both the vapor and liquid phases, denoted as $\mu_i^{\rm V}(T,P,x_i^{\rm V})$ and $\mu_i^{\rm L}(T,P,x_i^{\rm L})$ respectively, are the same.

Predicting the formation of azeotropes remains a theoretical challenge, as one needs to accurately characterize the vaporization and the free energies of mixing of liquids. Calculating μ_i of the mixture involves both energetic and entropic contributions, and the latter often necessitates complicated simulation setups. Secondly, modeling the coexistence of vapor and liquid poses difficulties for the applied empirical force fields in atomistic simulations. To accurately capture vapor-liquid coexistence, it is often neccesary to refine force field parameters using experimental measurements from the vapor-liquid equilibria (VLE) conditions.^{8,9}

Gibbs ensemble Monte Carlo (GEMC) method¹⁰ is commonly used to compute the excess chemical potential μ_i^{Ex} of fluids. GEMC introduces two separate simulation boxes to simultaneously represent the liquid and vapor phases, and perform Monte Carlo exchanges of particles and volumes to determine the equilibrium densities and compositions of the coexisting phases. GEMC is popular for simulating the azeotrope in multiple systems such as ethanol/water, 11 methanol/n-hexane, 12 ethanol/n-hexane, 12 methanol/acetonitrile, 13 1-pentanol/nhexane, ¹⁴ 1-propanol/acetonitrile, ¹⁴ ethyl acetate/ethanol, ¹⁵ methanol/ethyl acetate, ¹⁵ and ethanol/benzene. ¹⁶ However, GEMC can require substantial computational resources to achieve statistical convergence. 17–20 The convergence requirements become particularly demanding for dense liquids or large molecules for which particle insertion is difficult, and near critical points, where phase distinction proves problematic. Similarly, the free energy perturbation (FEP) method²¹ can simulate chemical potentials through particle insertion based on molecular dynamic (MD) simulations. Yet this approach also struggles with the same convergence issues as GEMC.

One can also estimate μ_i^{Ex} using approximative thermodynamic models, which are based on simplified expressions of molecular interactions. In these models, the activity coefficient, γ_i , is derived from interaction parameters and utilized to compute μ_i^{Ex} as $\mu_i^{\text{Ex}} = k_{\text{B}}T \ln \gamma_i$. The expression of γ_i varies among different thermodynamic models, such as the structuralgroup-contribution-based model UNIQUAC²² and advanced machine-learning-based models such as SPT-NRTL,²³ artificial neural network,²⁴ and random forest models.²⁴ The broader use of these models, however, is restricted by limited experimental data on mixture densities and VLE compositions, which are crucial for model calibration. More recently, the COnductor-like Screening MOdel for Real Solvents (COSMO-RS)²⁵ in conjunction with Density Functional Theory (DFT) computations is developed as a general predictive

¹⁾Institute of Science and Technology Austria, Am Campus 1, 3400 Klosterneuburg, Austria

²⁾Department of Chemistry, University of California, Berkeley, CA 94720, USA

a) Electronic mail: Xiaoyu. Wang@ist.ac.at

b) Electronic mail: bingqingcheng@berkeley.edu

method, which computes γ_i from the probability distribution of the surface charge of molecules.

In this paper, we first introduce a methodology that streamlines the prediction of azeotropes in binary mixtures in Sec II. This approach utilizes the S0 method²⁶ to compute the compositional dependence of chemical potentials from MD simulations. Combined with the experimental boiling point T_b and vaporization enthalpy $\Delta_{vap}H$, we derive the vapor-liquid equilibrium of the mixture and subsequently locate the azeotrope. In Sec IV, the application of this approach is demonstrated through the simulations of five case studies, including two minimum boiling mixtures (ethanol/water and ethanol/isooctane), one non-azeotropic mixture (methanol/water), and two maximum boiling mixtures (hydrazine/water and acetone/chloroform). Our findings reveal the necessity of using experimental boiling points and vaporization enthalpy for accurate azeotrope predictions since the empirical force fields are not accurate enough for these quantities. In Sec V, we utilize the regular solution model to discuss the thermodynamic prerequisites for the formation of azeotropes in binary mixtures.

II. METHODS

The procedure to predict the azeotrope is captured in Fig. 1. First, a reference state is established using the boiling points of pure substances under fixed pressure ($P_0 = 1$ bar). This step sets the baseline of chemical potentials, as the chemical potential of component i in the vapor phase $\mu_{i, \text{pure}}^{\text{V}}(T_{\text{b}}, P_0)$ is equal to that in the liquid phase $\mu_{i, \text{pure}}^{\text{L}}(T_{\text{b}}, P_0)$. The relevant T_{b} and $\Delta_{\text{vap}}H$ can be determined either from VLE MD simulations or thermodynamical databases such as ChemSpider, ²⁷ NIST, ²⁸ PubChem, ²⁹ and Reaxys. ³⁰ Further information is elaborated in Fig. 1(a) and Sec II A.

The second step (Fig. 1(b)) involves calculating the chemical potential of each component in the vapor and liquid mixtures across the composition-temperature (x-T) parameter space, utilizing the S0 method. ²⁶ Specific descriptions of chemical potentials in the liquid $\mu_i^{\rm L}(T,P,x_i^{\rm L})$ and vapor $\mu_i^{\rm V}(T,P,x_i^{\rm V})$ are discussed in Secs II B and II C, respectively.

We then proceed to establish the vapor-liquid equilibrium of the mixture based on $\mu_i^L(T,P,x_i^L) = \mu_i^V(T,P,x_i^V)$, and create the phase diagram of the mixture. Finally, we evaluate the presence of azeotropes and locate the minimum/maximum azeotropic temperature and the corresponding composition, as demonstrated in Sec II D.

A. Vapor-liquid coexistence of pure substances

One can either use experimental measurements of the boiling point T_b and the enthalpy of vaporization $\Delta_{\rm vap}H$ for pure substances, or compute these from simulations. The Clausius–Clapeyron relation can be used to describe the VLE con-

ditions for pure substances:

$$\frac{\mathrm{d}P}{\mathrm{d}T} = \frac{\left\langle H^{\mathrm{V}} \right\rangle_{T,P} - \left\langle H^{\mathrm{L}} \right\rangle_{T,P}}{T(V^{\mathrm{V}} - V^{\mathrm{L}})} = \frac{\Delta_{\mathrm{vap}}H}{T\Delta V},\tag{3}$$

where $\Delta V = V^{\rm V} - V^{\rm L}$ is the molar volume difference between the vapor and liquid phases at VLE conditions. Eq. 3 can be further simplified based on the following three assumptions: First, $\Delta_{\rm vap}H = \left\langle H^{\rm V}\right\rangle_{T,P} - \left\langle H^{\rm L}\right\rangle_{T,P}$ can be considered to be constant within the investigated temperature range close to the boiling point $T_{\rm b}$ at the constant pressure $P_0 = 1$ bar. Second, ΔV is dominated by the vapor volume $V^{\rm V}$ that is typically 3 orders of magnitude larger than that of liquid. Third, $V^{\rm V}$ can be computed from the equation of state for an ideal gas $PV^{\rm V} = nk_{\rm B}T$ at low pressures, and then Eq. 3 is thus simplified as:

$$\frac{\mathrm{d}\ln P}{\mathrm{d}1/T} = -\Delta_{\mathrm{vap}}H.\tag{4}$$

Consequently, $\Delta_{\text{vap}}H$ can be calculated by evaluating the slope on the $\ln P$ -T diagram, as presented in Fig. 1(a). Through MD simulations of the vapor-liquid interfacial systems, we derive the equilibrium pressures of the coexisting phases at varying temperatures, as detailed in Sec III A. We then compute the boiling temperature $T_{\rm b}$ by setting P=1 bar in Eq. 4.

Moreover, regarding the chemical potentials of pure substances at temperatures other than the boiling point, along an isobaric path, the thermodynamic integration (TI) expression of $\Delta\mu_{L,\text{pure}}^{L\to V}(T)$ is specified as follows:

$$\Delta\mu_{i, \text{ pure}}^{L \to V}(T) = -k_B T \int_{T_b}^{T} \frac{\Delta_{\text{vap}} H_i}{N k_B T^2} dT, \tag{5}$$

where $\varDelta\mu_{i,\;\mathrm{pure}}^{\mathrm{L}\to\mathrm{V}}(T)$ is the free energy change needed to transform N molecules from the liquid into the vapor state at a given temperature T.

B. Chemical potentials of the liquid

As the azeotrope prediction in this work is performed at a constant pressure $P_0 = 1$ bar, we will omit the notion of P in subsequent discussion. Here we base μ_i^L of component i in the solution on its chemical potential under its boiling conditions, where $\mu_{i, \text{pure}}^L(T_b) = \mu_{i, \text{pure}}^V(T_b)$. This baseline simplifies computations by setting a consistent reference point for both the vapor and the liquid phases, representing a thermodynamic cycle construction between the vapor and the liquid through an initial coexistence point. Accordingly, the chemical potential of component i in the liquid mixture can be described as:

$$\mu_i^{\mathsf{L}}(T, x_i) = \mu_{i, \text{ pure}}^{\mathsf{L}}(T_{\mathsf{b}}) + \Delta \mu_{i, \text{ pure}}^{\mathsf{L} \to \mathsf{V}}(T)$$

$$+ k_{\mathsf{B}} T \ln x_i^{\mathsf{L}} + k_{\mathsf{B}} T \ln \gamma_i.$$

$$(6)$$




FIG. 1 A schematic representation of the simulation workflow for identifying an azeotrope. The different components are symbolized by blue circles and orange diamonds. (a) T_b and $\Delta_{vap}H$ are derived, either from MD simulations or existing databases such as ChemSpider, ²⁷ NIST, ²⁸ PubChem, ²⁹ and Reaxys. ³⁰ From vapor-liquid coexistence MD simulations under NVT ensembles, one can extract the vapor pressure at different temperatures to fit the Clausius-Clapeyron relation. The vaporization enthalpy $\Delta_{vap}H$ is derived from the gradient of $d \ln P/d1/T$, and the boiling point T_b is the T at $P_0 = 1$ bar. (b) The chemical potentials (μ_i^V, μ_i^L) of each component in the vapor and liquid mixtures are computed using the S0 method. ²⁶ As the chemical potentials depend on temperature T and composition x, one performs sets of simulations for different compositions at different isotherm conditions.

Combined with Eq. 5, Eq. 6 becomes:

$$\mu_i^{L}(T, x_i) = \mu_{i, \text{ pure}}^{V}(T_b) + \Delta_{\text{vap}} H_i(\frac{T}{T_b} - 1) + k_B T \ln(x_i^{L} \gamma_i).$$
 (7)

We then apply the newly established S0 method²⁶ to compute γ_i . This method utilizes the thermodynamic relationship between the fluctuations in particle number and the derivatives of the chemical potentials with respect to the molar fraction.^{31,32} It relies on the static structure factors S(k) computed from equilibrium MD simulations at different mixture fractions, effectively overcoming the finite-size effect. For binary systems, the derivatives of the chemical potentials $\partial \mu_i$ with respect to $\ln x_i$ is:

$$\left(\frac{\partial \mu_i}{\partial \ln x_i}\right)_{T,P} = \frac{k_B T}{x_j S_{ii}^0 + x_i S_{ii}^0 - 2\sqrt{x_i x_j} S_{ij}^0},\tag{8}$$

where i and j denote different components respectively, while S_{ii}^0 , S_{jj}^0 , S_{ij}^0 are the static structure factors for pairs of particles i-i, j-j, and i-j in small-k limit.

Specifically, we conduct equilibrium MD simulations in the isothermal–isobaric ensemble (*NPT*) at different mole fractions and then perform numerical integration, i.e.

$$\mu_{i}^{L}(T, x_{i}^{L}) = \mu_{i, \text{ pure}}^{L}(T_{b}) + \Delta_{\text{vap}}H_{i}(\frac{T}{T_{b}} - 1) + k_{B}T \int_{0}^{\ln x_{i}} d\ln(x_{i}) \frac{k_{B}T}{x_{j}S_{ii}^{0} + x_{i}S_{jj}^{0} - 2\sqrt{x_{i}x_{j}}S_{ij}^{0}}.$$
(9)

C. Chemical potentials of the vapor

We follow the same procedure outlined in Sec IIB to calculate the chemical potentials of the component in the vapor phase. At ambient pressure, the ideal gas assumption can offer a useful approximation: the ideal gas behaviors result in ideal mixing in the vapor mixture with no excess chemical potential, i.e. $\mu_i^{\text{V,Ex}} = 0$. Moreover, the partial pressure P_i of the component i is equivalent to $x_i^{\text{V}}P$ according to the Dalton law for ideal gases in the vapor mixture. Hence μ_i^{V} is simplified to:

$$\mu_i^{V}(T, y_i) = \mu_{i, \text{ pure}}^{V}(T_b) + k_B T \ln \frac{P_i}{P_0}$$

$$= \mu_{i, \text{ pure}}^{V}(T_b) + k_B T \ln x_i^{V}.$$
(10)

D. Vapor-liquid equilibrium and azeotrope of the binary system

The vapor-liquid coexistence curve of a mixture is established when each component shares the same chemical potentials in the vapor and the liquid, denoted as $\mu_i^V(T, x_i^V) = \mu_i^L(T, x_i^L)$. Specifically, the vapor-liquid equilibria of mixtures are defined by combining Eqs. 7 and 10:

$$\Delta_{\text{vap}} H_i(\frac{T}{T_b} - 1) + k_B T \ln(x_i^L \gamma_i) = k_B T \ln x_i^V. \tag{11}$$

By solving Eq. 11, one obtains a set of x_i^{V} - x_i^{L} values at variable temperatures, which give the vapor and liquid compositions at VLE respectively. The azeotrope is then identified by determining the temperature at which $x_i^{\text{V}} = x_i^{\text{L}}$ according to the

definition of an azeotrope stated in Eq. 2. We also estimate the statistical errors in the computed values via random sampling using the Monte Carlo method.

III. SIMULATION DETAILS

All MD runs are initiated using empirical force fields within GROMACS (V2022.03).³³ Specifically, the Coulomb and 12–6 Lennard-Jones (LJ) terms for non-bonded interactions are applied with a truncation of 1.4 nm. Particle Mesh Ewald (PME) summations are used for the electrostatic interactions with a Fourier space grid of 0.16 nm. A fourth-order interpolating function is applied with a relative tolerance of 10⁻⁴ for the PME solver. With the united-atom (UA) models, we employ the Lorentz-Berthelot mixing rule^{34,35} for pairwise interactions between different atoms. In contrast, for the OPLS models,³⁶ we use the geometric-mean mixing rule instead. Additionally, the LINCS algorithm³⁷ is utilized to restrict the lengths of all bonds in the UA models, as well as H-bonds in all-atom (AA) models. This algorithm allows for longer integration time steps such as 2 fs.

A. Boiling point and vaporization enthalpy

MD simulations of pure substances for vapor-liquid interfacial systems are performed in the *NVT* ensemble to determine the boiling temperature (T_b) and its enthalpy of vaporization $(\Delta_{\text{vap}}H)$ as per Eq. 4. To create a vapor-liquid coexistent system, an orthorhombic cell that contains 1000 molecules is built. This simulation cell features an elongated z-axis $(L_z > L_x = L_y)$, leading to adjacent liquid and vapor slabs with interfaces spanning parallel to the xy plane, as demonstrated in Fig. 1(a).

We prevent large vapor pressure and surface tension oscillations due to finite-size effects by verifying the dimensions of simulaiton cell (L_x, L_y, L_z) against two standards, thereby ensuring an appropriate vapor phase. First, it has been reported that the side length of the interfacial area (L_x, L_y) has to be at least ten times the diameter of molecules or the radius of gyration D_i , $^{38-41}$ i.e., $L_x = L_y > 10D_i$. Meanwhile, the elongated axis L_z is at least 3 times the side length of the crossing section L_x .

Regarding the production run, a timestep of 2 fs is used with a \sim 2 ns equilibration run in the *NVT* ensemble using the stochastic velocity rescaling thermostat with a time constant of 0.4 ps in GROMACS.⁴² For accurate simulations of nonhomogeneous systems, especially with larger molecules such as isooctane, we set a cutoff of 2.3 nm for LJ and electrostatic interactions. The long-range dispersion correction for pressure is disabled for the interfacial system in GROMACS.

To estimate the vapor pressure P_{vap} , we use the diagonal element of the pressure tensor along the elongated z-direction, denoted as P_{zz} . By fitting the Clausius–Clapeyron relation in Eq 4, T_{b} and $\Delta_{\text{vap}}H$ of the pure substance at 1 bar are determined using the least-square method.

B. Activity coefficients

MD simulations are performed on binary mixtures with varying mole fractions at 1 bar using GROMACS. The temperature range covers the boiling points of the pure components. The target of these simulations is to establish a functional relationship between activity coefficients $\gamma(x,T)$ and mixture compositions as well as temperature. Therefore, we could explore x-T parameter space to solve Eq. 11 through linear interpolation.

Each simulation box contains 10^4 molecules in total with different fractions of two components to avoid the system size effect. After energy minimization of the initial configuration, the equilibration run is carried out in the *NPT* ensemble for ~ 0.5 ns with a time step of 1 fs. Production runs take place following this for another 1.0 ns.

In each snapshot taken every 2000 MD steps, the atoms close to the geometrical centers of the molecules are taken as the positions of the molecules, which are then used to calculate S(k) via Fourier transformation. The time-averaged S(k) of all pairs of components i-i, j-j, and i-j, along with the associated uncertainty, are determined based on the correlation analysis in the time series. These S(k) values are then fitted to the Ornstein–Zernike form with a maximum cutoff $k^2 = 0.01$ A to obtain the values for S^0 . Finally, the obtained structure factors are utilized to calculate the chemical potentials and the activity coefficients of each species as expressed in Eq. 8.

IV. APPLICATIONS

TABLE I Simulation and experimental results of boiling temperature T_b (K) and molar enthalpy of vaporization $\Delta_{\text{vap}}H$ (kJ/mol) at 1 bar.

Species	Force field	$T_{\rm b,sim}$	$T_{\rm b,exp}$	$\Delta_{\mathrm{vap}}H_{\mathrm{sim}}$	$\Delta_{\mathrm{vap}}H_{\mathrm{exp}}$
Water	TIP4P-Ew	397.7 ± 1.3	373.15 ⁸	51.4 ± 1.0	40.6°
	SPC/E	397.6 ± 1.7		47.6 ± 3.0	
Ethanol	TraPPE-UA	344.7 ± 1.2		39.6 ± 2.1	. – – – -
	OPLS-AA	334.9 ± 1.0	351.5 ⁴³	38.0 ± 1.6	42.3^{43}
	KBUA	390.6 ± 1.9		37.5 ± 1.5	
Isooctane	TraPPE-UA	361.6 ± 2.7	372.444	31.0 ± 3.3	
Methanol	TraPPE-UA	334.5 ± 1.3	337.8 ⁴³	35.9 ± 0.8	37.6^{43}
Hydrazine	OPLS	425.4 ± 1.1	386.95^{45}	48.6 ± 2.1	44.5^{46}
Chloroform	AA	342.2 ± 1.5	334.3 ⁴⁷	28.9 ± 1.0	31.3^{47}
Acetone	AA	315.0 ± 1.3	329.344	28.1 ± 1.7	29.1 ⁴⁴

A. Ethanol/water mixtures

We examine three force field parameterization strategies for ethanol/water: the TraPPE-UA model⁴⁸ of ethanol in TIP4P-Ew water, ⁴⁹ OPLS-AA model of ethanol⁵⁰ in TIP4P-Ew water, and KB-based UA model (KBUA) of ethanol⁵¹ in SPC/E

water.⁵² The simulated boiling temperature and enthalpy of vaporization of each model are summarized in Table I, with corresponding activity coefficients of each component presented in Fig. 2.

As listed in Table I, VLE simulations reveal that both SPC/E and TIP4P-Ew water have similar boiling points, 7% higher than the observed experimental values.⁸ These findings are consistent with prior research that documented similar boiling points, as $T_b = 392 \pm 2$ K⁴⁹ and $T_b = 394$ K⁸ for TIP4P-Ew water and $T_b = 396$ K⁵³ for SPC/E water. Consequently, the higher boiling temperature for TIP4P-Ew water can be traced back to inherent deficiencies of the force field. For ethanol, the TraPPE-UA model accurately predicts the boiling temperature and enthalpy of vaporization that align closely with experimental data, with a discrepancy of less than 5%, while the boiling points estimated from the OPLS-AA and KUBA models diverge significantly from experimental results with a difference of ~ 10 %.

The activity coefficients of water and ethanol from different force fields are calculated with the S0 method, as shown in Fig. 2. For comparison, the activity coefficients in the UNI-QUAC model are also calculated based on experimental data,⁵⁴ using the Phasepy⁵⁵ python package. As depicted in Fig. 2, the combination of TraPPE-UA ethanol and TIP4P-Ew water best fits the experimental fitting, so we use them for the subsequent determination of the azeotrope.

The simulated activity coefficients γ , boiling temperature $T_{\rm b}$, and enthalpy of vaporization $\Delta_{\rm vap}H$ based on TraPPE-UA ethanol and TIP4P-Ew water are used to derive the chemical potentials of water and ethanol in relation to the mole fractions in both the liquid and vapor phases, as presented in Fig. 3. To locate the vapor-liquid coexistence curve and azeotrope of the ethanol/water mixture, we follow the process outlined in Sect IID. In essence, we first deduce the temperature range that allows for vapor-liquid coexistence, including discerning whether the solution has a minimum- or maximum-boiling azeotrope, which is finalized by solving Equation 11. For the ethanol/water mixture specifically, it indicates a boiling point that is less than that of any pure component, essentially a minimum-boiling azeotrope. Furthermore, this finding is confirmed by Fig. 2(b), which displays that the chemical potentials of both water and ethanol exceed the values of an ideal solution. In other words, the interactions between water and ethanol destabilize the mixture and result in a minimumboiling azeotrope.

We further deduce the vapor and liquid compositions at various temperatures within the temperature range where both phases exist. In detail, the equilibrium phase compositions for both liquid and vapor can be detailed in simultaneous Eqs. 12 and Fig. 3:

$$\begin{cases} \Delta_{\text{vap}} H_{\text{eth}} \left(\frac{T}{T_{b,\text{eth}}} - 1 \right) + k_{\text{B}} T \ln(x_{\text{eth}}^{\text{L}} \gamma_{\text{eth}}) = k_{\text{B}} T \ln x_{\text{eth}}^{\text{V}} \\ \Delta_{\text{vap}} H_{\text{water}} \left(\frac{T}{T_{b,\text{water}}} - 1 \right) + k_{\text{B}} T \ln(x_{\text{water}}^{\text{L}} \gamma_{\text{water}}) = k_{\text{B}} T \ln x_{\text{water}}^{\text{V}} \\ x_{\text{eth}}^{\text{L}} + x_{\text{water}}^{\text{L}} = 1 \\ x_{\text{eth}}^{\text{V}} + x_{\text{water}}^{\text{V}} = 1 \end{cases}$$

$$(12)$$

After determining vapor-liquid equilibria, we can produce the phase diagram of the ethanol/water mixture at 1 bar; see Fig. 4(a). Fig. 4(a) also compares with the experimental results⁵⁴ for reference. The computed azeotropic composition $x_{\rm eth} = 99.0$ mol% (blue pentagon in Fig. 4(a)) diverges significantly from experimental data due to discrepancies in $T_{\rm b}$ and $\Delta_{\rm vap}H$ between the experimental and simulated results. Alternatively, we recalculate the phase diagram with experimental $T_{\rm b}$ and $\Delta_{\rm vap}H$ values with computed γ . The corrected diagram (red lines in Fig. 4(a)) predicts an azeotropic temperature of 350.8 \pm 0.7 K and an azeotropic composition of 89.3 \pm 0.1 mol%. These predictions are in good agreement with experimental azeotrope that has a temperature of 351.55 K for a mixture containing 89 mol% ethanol and 11 mol% water.⁵⁴




FIG. 2 Activity coefficients of ethanol/water at 355 K calculated using different force fields. Blue and red symbols denote the activity coefficients γ of ethanol and water with respect to the mole fraction of ethanol $x_{\rm eth}$, respectively. Three different symbols denote different selections of force field parameters: 1) open circles (o) for OPLS-AA ethanol in TIP4P-Ew water; 50 2) solid stars (\star) for TraPPE-UA ethanol in TIP4P-Ew water; and 3) crosses (\times) for KBFF ethanol 51 in SPC/E water. 52 The solid curves represent the fitted values of the UNIQUAC model with Phasepy, 55 using experimental data. 54

B. Ethanol/isooctane mixture

The ethanol/isooctane mixture is representative of common fuel mixtures. ⁵⁶ We employ the TraPPE-UA parameterization for the ethanol/isooctane system. The corresponding values of $T_{\rm b}$ and $\Delta_{\rm vap}H$ at 1 bar are listed in Table I.

The original phase diagram of the ethanol/isooctane mixture at 1 bar, derived directly from the calculated quantities, is shown by the blue lines in Fig. 4(b). In addition, Fig. 4(b) presents a corrected phase diagram (red lines) incorporating experimental T_b and $\Delta_{\rm vap}H$, alongside experimental data for comparison.⁵⁷ The comparison between original data (blue lines) and updated data (red lines) in Fig. 4(b) addresses the




FIG. 3 The chemical potential of (a) water and (b) ethanol relative to the mixture composition in the liquid (solid lines) and vapor (dashed lines) at 352 K, using the TIP4P-Ew water and TraPPE-UA ethanol models. Black dots show equilibrium compositions for both liquid and vapor, with corresponding chemical potentials represented by vertical dashed lines. The grey lines indicate the chemical potential of ideal mixing where $\gamma_i = 1$.

influence of $T_{\rm b}$ and $\Delta_{\rm vap}H$ on phase diagram predictions and the challenge of accurately modeling vapor-liquid equilibria with the presumed potential energy surface. Based on the corrected phase diagram, the azeotropic temperature of the ethanol/isooctane mixture is predicted to be 344.9 ± 0.1 K, with an azeotropic composition of $x_{\rm eth} = 63 \pm 1$ mol%. This result aligns closely with a prior experiment that documented an azeotropic temperature of 344.42 K with a mixture of 64.5 mol% ethanol under a pressure of 1.01 bar. 57,58

C. Methanol/water mixture

We perform a negative test using the methanol/water mixture, which does not exhibit any azeotropic behavior when heated to boiling. We utilize the TraPPE-UA model for methanol and the TIP4P-Ew model for water to initiate MD simulations. The simulated boiling point and vaporization enthalpy of methanol are displayed in Table I, which agrees well with experimental results. 43

The derived phase diagram is presented in Fig. 4(c), where no azeotrope is identified as expected. Once updating our calculations with experimental T_b and $\Delta_{\text{vap}}H$, the final phase (red lines in Fig. 4(c)) matches well with previous laboratory measurements at 1.01 bar (grey lines in Fig. 4(c)).⁵⁹

D. Hydrazine/water mixture

Regarding the maximum boiling solutions, a benchmark examination is performed on the hydrazine/water mixture at 1 bar. The parameters from the OPLS-AA/1.14*CM1A force field are used for hydrazine. The charge distribution within N₂H₄ is generated using the LigParGen online server. 60,61 This model of N₂H₄ results in lower vapor pressures and leads to a higher boiling temperature of 425.4 \pm 1.1 K at 1 bar, approximately 38 degrees higher than the experimental value, despite an acceptable $\varDelta_{\rm vap}H$ simulated.

Even if the empirical parameterization of the H₂O and

 N_2H_4 force fields causes a significant deviation in the calculated phase diagram, the occurrence of the maximum-boiling azeotropic phenomenon is successfully predicted, as illustrated in Fig. 4(d). Furthermore, the corrected phase diagram, depending on accurate values of T_b and $\Delta_{\rm vap}H$, is presented as red lines in Fig. 4(d). Accordingly, the updated azeotropic temperature of the hydrazine/water mixture is 393.6 ± 0.3 K, with an azeotropic composition of 62.5 ± 0.5 mol% hydrazine. By comparing our predicted azeotropic composition at 1 bar with experimental data indicating $X_{N_2H_4} = 55$ mol% at 0.93 bar and $X_{N_2H_4} = 58$ mol% at 1.03 bar, 62 we notice that our estimate falls within a 5% difference range. On the other hand, the computed azeotropic temperature aligns well with the experimental value of 393.65 K at 1.03 bar.

E. Acetone/chloroform mixture

The acetone/chloroform mixture represents a classic negative azeotrope that has been extensively studied. 63–67 The force field parameters applied are adapted from previous literature.9 The simulated values of T_b and $\Delta_{vap}H$ for acetone and chloroform satisfactorily match experimental data, differing by less than 5%, as detailed in Table I. However, as depicted by the blue lines in Fig. 5(e), the azeotrope at 1 bar cannot be accurately identified based only on the simulated quantities. Only by substituting experimentally determined values for $T_{\rm b}$ and $\Delta_{\text{vap}}H$ can the prediction of azeotrope be achieved, marked by red lines in Fig. 5(e). Moreover, for the acetone/chloroform mixture, the estimated azeotropic temperature is 335.0 ± 0.3 K, with an azeotropic composition of $x_{\text{act}} = 26.7 \pm 2.1 \text{ mol}\%$. These results are in good agreement with previous experimental observations, showing a temperature of 337.6 K and an acetone composition of 40 mol% at a pressure of 1.01 bar.⁶⁸

V. REGULAR SOLUTION MODEL AND AZEOTROPE FORMATION

To understand the driving force underlying the formation of azeotropes and to rationalize the associated thermodynamic conditions, we apply a regular solution model. ^{70,71} In this framework, the Gibbs free energy during mixing $\Delta_{\rm mix}G(x)$ is expressed as:

$$\Delta_{\text{mix}}G(x) = \Delta_{\text{mix}}G^{\text{Id}}(x) + \Delta_{\text{mix}}G^{\text{Ex}}(x)$$
$$= Nk_{\text{B}}T\sum_{i}x_{i}\ln x_{i} + \Delta_{\text{mix}}H^{\text{Ex}}, \qquad (13)$$

where the Gibbs free energy of ideal mixing $\Delta_{\rm mix}G^{\rm Id}(x)$ is only contributed by mixing entropy, and all the non-ideal factor $\Delta_{\rm mix}G^{\rm Ex}(x)$ of a regular solution is referred to as the excess enthalpy of mixing $\Delta_{\rm mix}H^{\rm Ex}$. Specifically, for a binary mixture, one polynomial expression of the excess Gibbs free energy of mixing 72 is noted as:

$$\Delta_{\text{mix}}H^{\text{Ex}} = \Delta_{\text{mix}}G^{\text{Ex}} = x_i x_j \left[\omega_0 + \omega_1 \left(x_i - x_j \right) + \ldots \right]. \tag{14}$$




FIG. 4 The phase diagrams for the vapor-liquid equilibrium of (a) ethanol/water, (b) ethanol/isooctane, (c) methanol/water, (d) hydrazine/water, and (e) acetone/chloroform, at 1 bar. Liquid and vapor compositions are displayed by solid and dashed lines respectively. Colors indicate different results: blue for direct computations with empirical force fields (S0+MD), red for updates using activity coefficients from the S0 method and experimental T_b and $\Delta_{vap}H$ (S0+EXP), and grey for previous experimental measurements at 1 bar (EXP). Moreover, azeotropes predicted using just MD simulations (S0+MD) are represented by blue pentagons, updated azeotropes using experimental boiling points T_b and vaporization enthalpies $\Delta_{vap}H$ (S0+EXP) by red stars, and experimental measurements by grey squares.

It should be noted that a regular solution strictly involves only ω_0 , while solutions described with higher order terms $\omega_{k,k>0} \neq 0$ are defined as sub-regular solutions. As a result, $\Delta_{\text{mix}}G(x)$ is described as:

$$\Delta_{\text{mix}}G(x) = Nk_{\text{B}}T \sum_{i} x_{i} \ln x_{i} + x_{i}x_{j} \left[\omega_{0} + \omega_{1} \left(x_{i} - x_{j} \right) + \ldots \right]. \tag{15}$$

Here we fit our simulation results with the regular solution model in Eq. 15, as shown in Fig. 5. The regular solution model aligns well with $\Delta_{\rm mix}G(x)$ of two maximum-boiling solutions (hydrazine/water in Fig. 5(c) and acetone/chloroform in Fig. 5(d)), as the higher order ω terms $\omega_{k>0}$ are negligible. This is also evidenced by the near-symmetry of $\Delta_{\rm mix}G(x)$ with respect to the mixture composition x. Furthermore, the negative sign of ω_0 for maximum-boiling solutions demonstrates that the interactions between different molecules stabilize the system compared to the ideal mixing. On the other hand, the asymmetry in $\Delta_{\rm mix}G(x)$ requires higher order terms $\omega_{j,j\neq 0}$ to fit the model for the two minimum-boiling solutions (ethanol/water in Fig. 5(a) and ethanol/isooctane Fig. 5(b)). The positive sign of ω_0 for minimum-boiling solutions in-

dicates the interactions between different molecules are no stronger than those between the same molecules. In other words, a positive ω_0 represents a minimum-boiling azeotrope when $\Delta_{\rm mix}G(x)>\Delta_{\rm mix}G^{\rm Id}(x)$, while a negative ω_0 implies a maximum-boiling azeotrope when $\Delta_{\rm mix}G(x)<\Delta_{\rm mix}G^{\rm Id}(x)$.

We continue to investigate the compositional and temperature prerequisites for binary mixtures to form azeotropes. The effect of the temperature on γ_i is expressed by:

$$\frac{\partial \ln \left(\gamma_{i}\right)}{\partial T} = -\frac{\bar{H}_{i}^{\rm Ex}}{k_{\rm B}T^{2}},\tag{16}$$

where \bar{H}_i^{Ex} refers to the partial excess enthalpy of component i, which is linked to the excess enthalpy H^{Ex} of mixing via:⁷³

$$\bar{H}_i^{\text{Ex}} = H^{\text{Ex}} + (1 - x_i) \frac{\partial H^{\text{Ex}}}{\partial x_i}.$$
 (17)

Combining Eqs. 14, 16 and 17, the activity coefficients γ can be derived from the Redlich–Kister relation:⁷⁴

$$\mu_i^{\text{Ex}} = k_{\text{B}} T \ln \gamma_i = x_j^2 (\omega_0 + \omega_1 (3x_i - x_j) + \ldots).$$
 (18)




FIG. 5 Fitting results for both the regular and sub-regular solution models across five mixtures at specific temperatures: (a) ethanol/water at 350 K, (b) ethanol/isooctane at 344 K, (c) methanol/water at 340 K, (d) hydrazine/water at 391 K, and (e) acetone/chloroform at 338 K. Blue squares denote simulation outcomes, with dashed teal and solid red lines representing regular (ω_0) and sub-regular ($\omega_{k,k>0}$) solution models, respectively.

Substituting Eq. 18 into Eq. 11 gives the compositional criteria for azeotrope formation:

$$\omega_0 \frac{x_{\text{aze},i}^2}{A_{\text{van}}H_i} - \omega_0 \frac{T_{\text{b},i}}{T_{\text{b},i}} \frac{x_{\text{aze},j}^2}{A_{\text{van}}H_i} + \frac{T_{\text{b},i}}{T_{\text{b},i}} - 1 = 0.$$
 (19)

Furthermore, the azeotropic temperature can also be determined by taking Eq. 1 into account:

$$\frac{\omega_0}{T_{\text{aze}}} \left(\frac{x_{\text{aze},j}}{\Delta_{\text{vap}} H_i} - \frac{x_{\text{aze},i}}{\Delta_{\text{vap}} H_j} \right) = \frac{1}{T_{\text{b},j}} - \frac{1}{T_{\text{b},i}}.$$
 (20)

Consequently, the formation of an azeotrope occurs provided Eqs. 19 and 20 yield a solution.

To illustrate the predictive power of the regular solution model, we conducted two hypothetical tests on mixtures of methanol/water and ethanol/water. Fig. 6 illustrates the conditions needed for Eqs. 19 and 20 to result in an azeotrope. We examine azeotrope formation by varying differences in boiling temperatures of mixture components, as well as the value of ω_0 ; meanwhile, the respective experimental values of the vaporization enthalpy are fixed accordingly. As depicted in Fig. 6, the closer the boiling temperatures, the higher the likelihood of azeotrope formation. In addition, since ω_0 naturally measures how far a solution deviates from an ideal mixing,

larger ω_0 values facilitate the formation of azeotropes. For instance, the methanol/water mixture in Fig. 6(a) does not exhibit an azeotrope due to the large difference in the boiling temperatures ($\frac{T_{b,\text{water}}}{T_{b,\text{met}}} = 1.10$) and a small value of ω_0 ($\omega_0 = 2.19$). Conversely, the ethanol/water mixture in Fig. 6(b) displays a minimum-boiling azeotrope because the boiling temperatures of water and ethanol are closer ($\frac{T_{b,\text{water}}}{T_{b,\text{eth}}} = 1.06$) and the mixture has a larger ω_0 value ($\omega_0 = 4.19$).

The analyses based on the regular solution model thus reveal that azeotropes tend to form when the mixture's components have similar boiling points, indicated by $\frac{T_{b,A}}{T_{b,B}}$ approaching 1.0, and when intermolecular interactions between different components are strong, indicated by larger absolute values of ω . Specifically, a positive or negative ω_0 indicates a minimum-boiling or maximum-boiling azeotrope, respectively.

VI. CONCLUSIONS

In summary, we have developed a workflow that facilitates straightforward and reliable predictions of the azeotrope in binary solutions, as illustrated in Fig. 1. This approach determines the compositional dependence of chemical potentials from MD simulations using the S0 method. It also in-




FIG. 6 Two hypothetical tests illustrating azeotrope formation conditions in (a) methanol/water and (b) ethanol/water. We alter the boiling temperature differences between the binary mixture components (y-axis) and the ω_0 value (x-axis), maintaining the experimental vaporization enthalpy $\Delta_{\text{vap}}H$. The colorbar⁶⁹ shows the difference between the computed azeotropic temperature and predicted boiling temperature from ideal mixing, with blue and red zones indicating minimum-boiling and maximum-boiling azeotropes, respectively. The blank mid-section shows no azeotrope formation. (a) In methanol/water, the black circle refers to methanol/water's corresponding conditions ($\frac{T_{b,\text{water}}}{T_{b,\text{met}}} = 1.10$ and $\omega_0 = 2.19$) at 1 bar, situated at the no-azeotrope zone, showing no azeotrope. (b) In ethanol/water, the black circle denotes ethanol/water's corresponding conditions ($\frac{T_{b,\text{water}}}{T_{b,\text{eth}}} = 1.06$ and $\omega_0 = 4.19$) at 1 bar, located in the minimum-boiling azeotrope zone, implying an azeotrope presence.

volves using experimental data on boiling temperature and vaporization enthalpy data to establish the reference state for chemical potentials. Compared to the state-of-the-art methods, 16,21,23,24,75,76 we streamline the procedure of computing the excess chemical potential of each component in the mixture. Applying this approach, we reproduce the formation of azeotropes or the lack of in five test cases, including ethanol/water, ethanol/isooctane, methanol/water, hydrazine/water, and acetone/chloroform mixtures. We find that it is crucial to use the experimental boiling point and vaporization enthalpy for reliable azeotrope predictions, as empirical force fields are not accurate enough for these quantities.

We also explore the conditions necessary for the formation of azeotropes with the regular solution model. We find that azeotropes typically form when the mixture components have comparable boiling points and strong interactions.

We expect that our workflow for predicting azeotropes holds considerable potential for various technologically important systems. First, it can contribute to the understanding of azeotrope formation at the atomistic scale through MD simulations. Second, it may support the development of extraction and purification techniques in the chemical industry. Furthermore, this approach can provide information on the properties of flammable azeotropic fuels and the combustion chemistry of fuel blends.

Acknowledgments BC thanks Alessandro Laio, who introduced the phenomenon of azeotrope and suggested to use of the S0 method to compute it. BC and XW thank Felix Wodaczek for insightful comments and suggestions on the manuscript.

BC and XW acknowledge resources provided by the Cambridge Tier-2 system operated by the University of Cambridge Research Computing Service funded by EPSRC Tier-2 capital grant EP/P020259/1.

Data availability statement All simulation setups, analysis scripts, and raw data in the study are available in the SI repository https://github.com/Xiaoyu-Wang-Stone/Azeotrope_S0.

¹S. Widagdo and W. D. Seider, "Journal review. azeotropic distillation," AIChE Journal 42, 96–130 (1996).

²L. H. Horsley and W. S. Tamplin, *Azeotropic data—III* (ACS Publications, 1973).

³W. L. Luyben and I.-L. Chien, *Design and control of distillation systems for separating azeotropes* (John Wiley & Sons, 2011).

⁴S. A. Shirazi, B. Abdollahipoor, B. Windom, K. F. Reardon, and T. D. Foust, "Effects of blending C3-C4 alcohols on motor gasoline properties and performance of spark ignition engines: A review," Fuel Processing Technology **197**, 106194 (2020).

⁵A. Loyte, J. Suryawanshi, G. Bhiogade, Y. Devarajan, and G. Subbiah, "Recent developments in utilizing hydrous ethanol for diverse engine technologies," Chemical Engineering and Processing-Process Intensification 177, 108985 (2022).

⁶B. Kanegsberg and E. Kanegsberg, Handbook for critical cleaning: cleaning agents and systems (CRC press, 2011).

⁷D. L. Williams, "Development of azeotropic blends to replace TCE and nPB in vapor degreasing operations," Sam Houston State University Huntsville United States (2016).

⁸C. Vega, J. Abascal, and I. Nezbeda, "Vapor-liquid equilibria from the triple point up to the critical point for the new generation of TIP4P-like models: TIP4P/Ew, TIP4P/2005, and TIP4P/ice," The Journal of chemical physics 125 (2006)

⁹G. Kamath, G. Georgiev, and J. J. Potoff, "Molecular modeling of phase

- behavior and microstructure of acetone- chloroform- methanol binary mixtures," The Journal of Physical Chemistry B **109**, 19463–19473 (2005).
- ¹⁰ A. Z. Panagiotopoulos, "Direct determination of phase coexistence properties of fluids by monte carlo simulation in a new ensemble," Molecular Physics 61, 813–826 (1987).
- ¹¹ M. Lísal, W. R. Smith, and I. Nezbeda, "Accurate vapour-liquid equilibrium calculations for complex systems using the reaction gibbs ensemble monte carlo simulation method," Fluid Phase Equilibria 181, 127–146 (2001).
- ¹²B. Chen, J. J. Potoff, and J. I. Siepmann, "Monte carlo calculations for alcohols and their mixtures with alkanes. transferable potentials for phase equilibria. 5. united-atom description of primary, secondary, and tertiary alcohols," The Journal of Physical Chemistry B 105, 3093–3104 (2001).
- ¹³ A. K. Sum, S. I. Sandler, R. Bukowski, and K. Szalewicz, "Prediction of the phase behavior of acetonitrile and methanol with ab initio pair potentials. i. pure components," The Journal of chemical physics 116, 7627–7636 (2002).
- ¹⁴C. D. Wick, J. M. Stubbs, N. Rai, and J. I. Siepmann, "Transferable potentials for phase equilibria. 7. primary, secondary, and tertiary amines, nitroalkanes and nitrobenzene, nitriles, amides, pyridine, and pyrimidine," The Journal of Physical Chemistry B 109, 18974–18982 (2005).
- ¹⁵H. Li, J. Zhang, D. Li, X. Li, and X. Gao, "Monte carlo simulations of vapour–liquid phase equilibrium and microstructure for the system containing azeotropes," Molecular Simulation 43, 1125–1133 (2017).
- ¹⁶D. Li, Z. Gao, N. K. Vasudevan, H. Li, X. Gao, X. Li, and L. Xi, "Molecular mechanism for azeotrope formation in ethanol/benzene binary mixtures through gibbs ensemble monte carlo simulation," The Journal of Physical Chemistry B 124, 3371–3386 (2020).
- ¹⁷ A. L. Ferguson, P. G. Debenedetti, and A. Z. Panagiotopoulos, "Solubility and molecular conformations of n-alkane chains in water," The Journal of Physical Chemistry B 113, 6405–6414 (2009).
- ¹⁸M. Khawaja, A. Sutton, and A. Mostofi, "Molecular simulation of gas solubility in nitrile butadiene rubber," The Journal of Physical Chemistry B 121, 287–297 (2017).
- ¹⁹T. R. Rogers and F. Wang, "Comparing alchemical free energy estimates to experimental values based on the ben-naim formula: How much agreement can we expect?" The Journal of Physical Chemistry B 124, 840–847 (2020).
- ²⁰D. Frenkel and B. Smit, Understanding molecular simulation: from algorithms to applications (Elsevier, 2023).
- ²¹M. Li, Y. Hu, L. Lv, T. Fang, L. Hao, S. Li, Y. Wu, X. Dong, and H. Liu, "Implementation of solvation free energy framework to predict the vapor-liquid equilibrium behaviors for the water–hydrazine and ethanol–water azeotropic systems," Chemical Engineering Science 275, 118751 (2023).
- ²²D. S. Abrams and J. M. Prausnitz, "Statistical thermodynamics of liquid mixtures: a new expression for the excess gibbs energy of partly or completely miscible systems," AIChE journal 21, 116–128 (1975).
- ²³B. Winter, C. Winter, T. Esper, J. Schilling, and A. Bardow, "Spt-nrtl: A physics-guided machine learning model to predict thermodynamically consistent activity coefficients," Fluid Phase Equilibria 568, 113731 (2023).
- ²⁴G. Sun, Z. Zhao, S. Sun, Y. Ma, H. Li, and X. Gao, "Vapor-liquid phase equilibria behavior prediction of binary mixtures using machine learning," Chemical Engineering Science 282, 119358 (2023).
- ²⁵ A. Klamt and F. Eckert, "Cosmo-rs: a novel and efficient method for the a priori prediction of thermophysical data of liquids," Fluid Phase Equilibria 172, 43–72 (2000).
- ²⁶B. Cheng, "Computing chemical potentials of solutions from structure factors," The Journal of Chemical Physics 157 (2022).
- ²⁷"Chemspider," http://www.chemspider.com, accessed: 2024-04-29.
- 28"National Institute of Standards and Technology (NIST)," https://www.nist.gov/, accessed: 2024-04-29.
- ²⁹"PubChem," https://pubchem.ncbi.nlm.nih.gov/, accessed: 2024-04-29.
- 30"Reaxys," https://www.reaxys.com/, accessed: 2024-04-29.
- ³¹J. G. Kirkwood and F. P. Buff, "The statistical mechanical theory of solutions. i," The Journal of chemical physics **19**, 774–777 (1951).
- ³²A. Ben-Naim, *Molecular theory of solutions* (OUP Oxford, 2006).
- ³³M. J. Abraham, T. Murtola, R. Schulz, S. Páll, J. C. Smith, B. Hess, and E. Lindahl, "Gromacs: High performance molecular simulations through multi-level parallelism from laptops to supercomputers," SoftwareX 1, 19– 25 (2015).
- ³⁴H. A. Lorentz, "Ueber die anwendung des satzes vom virial in der kinetischen theorie der gase," Annalen der physik 248, 127–136 (1881).

- ³⁵D. Berthelot, "Sur le mélange des gaz," Compt. Rendus **126**, 15 (1898).
- ³⁶J. Pranata, S. G. Wierschke, and W. L. Jorgensen, "Opls potential functions for nucleotide bases, relative association constants of hydrogen-bonded base pairs in chloroform," Journal of the American Chemical Society 113, 2810– 2819 (1991).
- ³⁷B. Hess, H. Bekker, H. J. Berendsen, and J. G. Fraaije, "Lincs: A linear constraint solver for molecular simulations," Journal of computational chemistry 18, 1463–1472 (1997).
- ³⁸M. González-Melchor, P. Orea, J. López-Lemus, F. Bresme, and J. Alejandre, "Stress anisotropy induced by periodic boundary conditions," The Journal of Chemical Physics 122 (2005).
- ³⁹P. Orea, J. López-Lemus, and J. Alejandre, "Oscillatory surface tension due to finite-size effects," The Journal of chemical physics 123 (2005).
- ⁴⁰J. Janeček, "Effect of the interfacial area on the equilibrium properties of lennard-jones fluid," The Journal of chemical physics 131 (2009).
- ⁴¹E. A. Muller, ^A. Ervik, and A. Mejía, "A guide to computing interfacial properties of fluids from molecular simulations [article v1. 0]," Living Journal of Computational Molecular Science 2, 21385–21385 (2020).
- ⁴²G. Bussi, D. Donadio, and M. Parrinello, "Canonical sampling through velocity rescaling," The Journal of chemical physics 126 (2007).
- ⁴³P. Linstorm, "Nist chemistry webbook, nist standard reference database number 69," J. Phys. Chem. Ref. Data, Monograph 9, 1–1951 (1998).
- ⁴⁴ V. Majer and V. Svoboda, Enthalpies of vaporization of organic compounds: a critical review and data compilation (Journal of the American Chemical Society, 1986).
- ⁴⁵T. Tipton, D. A. Stone, K. KuBulat, and W. B. Person, "Experimental and theoretical studies of the infrared spectra of hydrazines: N2H4, N2H3D, N2H2D2, N2HD3, and N2D4," The Journal of Physical Chemistry 93, 2917–2927 (1989).
- ⁴⁶D. Scott, G. Oliver, M. E. Gross, W. Hubbard, and H. M. Huffman, "Hydrazine: Heat capacity, heats of fusion and vaporization, vapor pressure, entropy and thermodynamic functions," Journal of the American Chemical Society 71, 2293–2297 (1949).
- ⁴⁷J. A. Manion, "Evaluated enthalpies of formation of the stable closed shell c1 and c2 chlorinated hydrocarbons," Journal of Physical and Chemical Reference Data 31, 123–172 (2002).
- ⁴⁸B. L. Eggimann, A. J. Sunnarborg, H. D. Stern, A. P. Bliss, and J. I. Siepmann, "An online parameter and property database for the trappe force field," Molecular Simulation 40, 101–105 (2014).
- ⁴⁹ H. W. Horn, W. C. Swope, J. W. Pitera, J. D. Madura, T. J. Dick, G. L. Hura, and T. Head-Gordon, "Development of an improved four-site water model for biomolecular simulations: TIP4P-Ew," The Journal of chemical physics 120, 9665–9678 (2004).
- ⁵⁰W. L. Jorgensen, "Optimized intermolecular potential functions for liquid alcohols," The Journal of Physical Chemistry 90, 1276–1284 (1986).
- ⁵¹E. A. Ploetz, S. Karunaweera, N. Bentenitis, F. Chen, S. Dai, M. B. Gee, Y. Jiao, M. Kang, N. L. Kariyawasam, N. Naleem, *et al.*, "Kirkwood–Buff-derived force field for peptides and proteins: philosophy and development of KBFF20," Journal of Chemical Theory and Computation 17, 2964–2990 (2021)
- ⁵²H. J. Berendsen, J. R. Grigera, and T. P. Straatsma, "The missing term in effective pair potentials," Journal of Physical Chemistry 91, 6269–6271 (1987).
- ⁵³M. Fugel and V. C. Weiss, "A corresponding-states analysis of the liquid-vapor equilibrium properties of common water models," The Journal of Chemical Physics **146** (2017).
- ⁵⁴R. M. Rieder and A. R. Thompson, "Vapor-liquid equilibria measured by a gillespie still-ethyl alcohol-water system," Industrial & Engineering Chemistry 41, 2905–2908 (1949).
- ⁵⁵G. Chaparro and A. Mejía, "Phasepy: A python based framework for fluid phase equilibria and interfacial properties computation," Journal of Computational Chemistry 41, 2504–2526 (2020).
- ⁵⁶R. A. Stein, J. E. Anderson, and T. J. Wallington, "An overview of the effects of ethanol-gasoline blends on si engine performance, fuel efficiency, and emissions," SAE International Journal of Engines 6, 470–487 (2013).
- ⁵⁷H.-C. Ku and C.-H. Tu, "Isobaric vapor-liquid equilibria for mixtures of acetone, ethanol, and 2, 2, 4-trimethylpentane at 101.3 kpa," Fluid phase equilibria 231, 99–108 (2005).
- ⁵⁸T. Hiaki, K. Takahashi, T. Tsuji, M. Hongo, and K. Kojima, "Vapor-liquid equilibria of ethanol with 2, 2, 4-trimethylpentane or octane at 101.3 kpa,"

- Journal of Chemical and Engineering Data 39, 720-722 (1994).
- ⁵⁹J. Gmehling, "Vapor-liquid equilibrium data collection," DECHEMA Chem. Ser. 1, 618 (1981).
- ⁶⁰W. L. Jorgensen and J. Tirado-Rives, "Potential energy functions for atomic-level simulations of water and organic and biomolecular systems," Proceedings of the National Academy of Sciences 102, 6665–6670 (2005).
- ⁶¹L. S. Dodda, I. Cabeza de Vaca, J. Tirado-Rives, and W. L. Jorgensen, "Lig-pargen web server: an automatic opls-aa parameter generator for organic ligands," Nucleic acids research 45, W331–W336 (2017).
- ⁶²J. G. Burtle, "Vapor pressure-composition measurements on aqueous hydrazine solutions," Industrial & Engineering Chemistry 44, 1675–1676 (1952).
- ⁶³A. Campbell and E. Kartzmark, "The energy of hydrogen bonding in the system: acetone–chloroform," Canadian Journal of Chemistry 38, 652–655 (1960).
- ⁶⁴A. Capparelli, H. Hertz, B. Kwatra, and R. Tutsch, "On the nature of association in the system chloroform—acetone. a nuclear magnetic relaxation study," Zeitschrift für Physikalische Chemie 103, 279–294 (1976).
- ⁶⁵A. Apelblat, A. Tamir, and M. Wagner, "Thermodynamics of acetone—chloroform mixtures," Fluid Phase Equilibria 4, 229–255 (1980).
- ⁶⁶V. A. Durov and I. Y. Shilov, "Molecular structure and physicochemical properties of acetone–chloroform mixtures," Journal of the Chemical Society, Faraday Transactions **92**, 3559–3563 (1996).
- ⁶⁷ J.-P. Ricard, "Process for the production of anhydrous hydrazine and device for its implementation," (2005), uS Patent 6,849,161.

- ⁶⁸Q. Gao, Z. Zhao, P. Jia, and X. Zhang, "Effect of ionic liquids on the isobaric vapor-liquid equilibrium behavior of acetone-chloroform," Applied Sciences 8, 1519 (2018).
- ⁶⁹F. Crameri, G. E. Shephard, and P. J. Heron, "The misuse of colour in science communication," Nature communications 11, 5444 (2020).
- ⁷⁰E. A. Guggenheim, "The statistical mechanics of regular solutions," Proceedings of the Royal Society of London. Series A-Mathematical and Physical Sciences 148, 304–312 (1935).
- ⁷¹J. Ganguly, *Thermodynamics in earth and planetary sciences* (Springer, 2008).
- ⁷²E. Guggenheim, "The theoretical basis of raoult's law," Transactions of the Faraday Society 33, 151–156 (1937).
- ⁷³O. Redlich and A. Kister, "On the thermodynamics of non-electrolyte solutions and its technical applications iii. systems with associated components," The Journal of Chemical Physics 15, 849–855 (1947).
- ⁷⁴L. Darken, "Application of the gibbs-duhem equation to ternary and multicomponent systems," Journal of the American Chemical Society **72**, 2909– 2914 (1950)
- ⁷⁵M. Boussaha, F. Amireche, R. Sahki, B. F. Belaribi, I. Mokbel, and J. Jose, "Isothermal vapor-liquid equilibria for binary liquid mixtures of ethylmer-captan and propylmercaptan with some common solvents," Fluid Phase Equilibria 569, 113763 (2023).
- ⁷⁶L. Zhang, Y. Yang, K. Yin, and Y. Liu, "A review of GEMC method and its improved algorithms," Acta Geochimica, 1–26 (2023).