DRAFT VERSION APRIL 12, 2024 Published: April 24, 2024: Research Notes of the AAS, Volume 8, Number 4, 110 Citation: Zhou, Ai-Ying: 2024, Res. Notes AAS, 8, 110 Publisher web: https://iopscience.iop.org/article/10.3847/2515-5172/ad414e

# Validating Gaia DR3 Pulsating Variable Classifications with *TESS*: Building Reliable $\delta$ Scuti and $\gamma$ Doradus Stars Catalogs (In Progress)<sup>\*</sup>

Ai-Ying Zhou<sup>1, 2</sup>

<sup>1</sup>National Astronomical Observatories, Chinese Academy of Sciences, A20 Datun Road, Chaoyang District, Beijing 100101, P.R. China <sup>2</sup>Key Laboratory of Radio Astronomy and Technology (CAS)

# ABSTRACT

Gaia DR3 revealed 748,058 pulsating variable stars of mixed DSCT|GDOR|SXPHE types. This project undertakes a comprehensive examination to validate and distinguish these stars using *TESS* data. Aiming for reliable catalogs of bona fide  $\delta$  Scuti and  $\gamma$  Doradus stars, I have validated 1715  $\delta$  Scuti stars, 1403  $\gamma$  Doradus stars, and identified 260 eclipsing binaries, one RR Lyrae star, and 460+ rotating variables from an initial sample of 16,690 objects. Notably, 15 of the newfound eclipsing binaries harbor pulsating  $\gamma$  Doradus components.

Keywords: Pulsating variable stars(1307) —  $\delta$  Scuti variable stars(370) —  $\gamma$  Doradus variable stars(2101) — Eclipsing binary stars(444)

# INTRODUCTION

Gaia Data Release 3 released 12.4 million variables, including 748,058 pulsating stars of mixed DSCT |GDOR|SXPHE types among the variability classification results of all classifiers – 9,976,881 objects (Gaia Collaboration et al. 2023)

With its extensive catalog of variable stars, *Gaia* DR3 presents a transformative opportunity to explore the properties of  $\delta$  Scuti and  $\gamma$  Doradus stars. However, independent verification using dedicated observations remains crucial, as the catalog includes known variables, eclipsing binaries, and non-variables (evident from a trial screening of the leading 16,690 objects, ~2.2% of total).

NASA's Transiting Exoplanet Survey Satellite (*TESS*, Ricker et al. 2015), provides exquisite photometry, making it ideal to identify stellar variability. This project leverages the synergy of *Gaia* catalogs and *TESS* high-precision light curves for a rigorous verification campaign of these 'Gaia variables'. My approach combines visual inspection of light curves, periodograms, and stellar parameters evaluation for stars location on the Hertzsprung-Russell diagram to validate variability and identify genuine pulsators. This process improves *Gaia* variable catalogs by removing false positives, yielding reliable catalogs of confirmed DSCT and GDOR.

#### SAMPLE AND DATA

This project works on two sources:

- 748,058 pulsating variable stars of mixed 'DSCT|GDOR|SXPHE' types from *Gaia* DR3 (Gaia Collaboration et al. 2023);
- 8,882 'DSCT\_SXPHE' and 3,018 stars of short time-scale from the *Gaia* DR2 Variability Results of 363,969 records (Gaia Collaboration et al. 2018).

Candidates were chosen according to the following selection criteria: effective temperature 5,500 to 12,000 K, brighter than TESS magnitude 16<sup>m</sup>, and in 16<sup>m</sup>, -19<sup>m</sup>, with 'score of the best class' higher than 0.3.

Corresponding author: Ai-Ying Zhou aiying@nao.cas.cn

1

<sup>\*</sup> Dedicated to my wife Jingyun Zhang

<sup>&</sup>lt;sup>1</sup> https://cdsarc.cds.unistra.fr/viz-bin/cat/I/358

For each star, I downloaded the shortest-exposure light curve from the latest observations (Sectors 1-71), prioritizing SPOC, TESS-SPOC, QLP, and TASOC products available at Mikulski Archive for Space Telescopes (MAST<sup>2</sup>):

- 20-second and 2-minute cadence observations (MAST Team 2021a,b),
- MIT Quick-Look Pipeline (Huang 2020),
- TESS Light Curves From Full Frame Images (TESS-SPOC, Caldwell et al. 2020),
- TESS Data For Asteroseismology Lightcurves (TASOC, Handberg, Rasmus et al. 2019).

Astronomical parameters are based on Simbad, *TESS* Input Catalog (TIC v8.2, Paegert et al. 2021) and *Gaia* DR2/DR3 (Gaia Collaboration et al. 2023, 2016). Spectral types were empirically derived using effective temperatures if unavailable.

### METHODOLOGY AND IDENTIFICATION

A star's variability type is determined on following factors: morphological characteristics and periodograms of their light curves, astrophysical parameters and their H-R diagram location.

I first check each candidate against the most relative catalogues: 70 552 DSCT and 8 080 GDOR (Zhou 2024); 123 841 and 84 206 *TESS* variables (Balona 2022; Fetherolf et al. 2023); and the new variables identified by Zhou (2023a,b,c). Then I check against Simbad. Candidates already identified as known variables in existing catalogs and those lacking corresponding *TESS* data are excluded. Each candidate underwent a two-step confirmation process: visually inspect the downloaded light curves, retrieve stellar parameters and evaluate their locations on H-R diagram.

A custom Python program (adapted from Zhou 2023a) facilitated the interactive screening, which performed various tasks: Online data retrieval from Simbad, TIC, MAST, and *Gaia*; Data processing and visualization; Periodogram and noise-level computation; Significant peaks identification; Cross-matching *Gaia* DR3 identifiers with TIC and Simbad main identifiers.

#### RESULTS

An initial cross-match of 16,690 pulsating stars from *Gaia* DR3 against Simbad and known catalogs revealed: 1454 pulsating variables ('PulsV', 8.7%, e.g. NP Del, HD 83628); 143 eclipsing binaries ('EclBin', 0.86%, e.g. FI Boo, V452 Cam); 346 generic variables ('Variable', 2.1%); 1108 known  $\delta$  Sct (6.6%, e.g. GG UMa, V1209 Her); 68 known  $\gamma$  Dor (0.41%, e.g. LS UMa, LU UMa). However, some Simbad 'PulsV' classifications potentially relied on *Gaia* DR3.

TESS light curves screened 7,360 objects out of 16,690 (remaining lacked TESS data or were not suitable), validated following Gaia DR3 discoveries: 1715  $\delta$  Sct, 1403  $\gamma$  Dor, one RR Lyr star (=TYC 2553-889-1), 460+ rotating variables, 260+ eclipsing binaries (including 15 with pulsating GDOR components), and others. Additionally, 760+ known variables, 2600+ unclassified variables, and 1050+ non-variables were identified at the precision of TESS photometry. Figure 1 shows ten of the TESS-validated 'Gaia variables' that actually do not belong to DSCT|GDOR|SXPHE types. Current validated variables as "Version 1" is available at Zenodo DOI:10.5281/zenodo.10023631 and this website too.

#### DISCUSSION

The preliminary results of initial examination of 16,690 objects from *Gaia* DR3 pulsating variables of DSCT|GDOR|SXPHE types using *TESS* data, emphasizes the importance of cross-validation with complementary data sources for robust scientific results. While automated algorithms are essential for classifying massive datasets, critical evaluation is crucial to avoid limitations.

Assuming current validation rates hold for the remaining sample, I would expect to validate over 85,000  $\delta$  Scuti and 66,000  $\gamma$  Doradus stars, exceeding a success rate of 20% for the whole sample. The number of stars that can be identified and confirmed ultimately depends on *TESS* data coverage. This is an ongoing work, I shall provide updates as the validation progresses.

<sup>&</sup>lt;sup>2</sup> https://mast.stsci.edu/portal/Mashup/Clients/Mast/Portal.html



Figure 1. Example *TESS* light curves of ten newly identified variables not DSCT|GDOR|SXPHE types. Top left: a new RRab star; Top right: EA+flare; 2nd and 3rd rows: four new heartbeat stars; 4th row left: a new rotating ellipsoidal variable star; Right, a candidate planetary transit; 5th rows: two new eclipsing binaries with pulsating  $\gamma$  Dor primary components.

I am indebted to my wife Jingyun Zhang for her unwavering support throughout my research. This work includes data collected with *TESS* mission, I acknowledge the use of *TESS* data, which are derived from pipelines at *TESS* Science Processing Operations Center. *TESS* High Level Science Products produced by the Quick-Look Pipeline at *TESS* Science Office at MIT, which are publicly available from the Mikulski Archive for Space Telescopes data archive at the Space Telescope Science Institute. Funding for *TESS* mission is provided by NASA Explorer Program. STScI is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5–26555. This research has made use of SIMBAD, operated at CDS, Strasbourg, France; data from the European Space Agency mission *Gaia*, processed by the *Gaia* Data Processing and Analysis Consortium. Funding for the DPAC has been provided by national institutions, in particular the institutions participating in the *Gaia* Multilateral Agreement.

Software: Astropy (Astropy Collaboration et al. 2013, 2018, 2022); Astroquery (Ginsburg et al. 2019); Lightkurve (Lightkurve Collaboration et al. 2018).

### REFERENCES

- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, A&A, 558, A33
- Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, AJ, 156, 123
- Astropy Collaboration, Price-Whelan, A. M., Lim, P. L., et al. 2022, ApJ, 935, 167
- Balona, L. A. 2022, arXiv e-prints, arXiv:2212.10776.
- Caldwell, D. A., Jenkins, J. M., & Ting, E. B. 2020, TESS Light Curves From Full Frame Images ("TESS-SPOC"), STScI/MAST, doi: 10.17909/T9-WPZ1-8S54
- Fetherolf, T., Pepper, J., Simpson, E., et al. 2023, ApJS, 268, 4
- Gaia Collaboration, Prusti, T., et al. 2016, A&A, 595, A1
- Gaia Collaboration, Brown, et al. 2018, A&A, 616, A1
- Gaia Collaboration, Vallenari, A., et al. 2023, A&A, 674, A1
- Ginsburg, A., Sipőcz, B. M., et al. 2019, AJ, 157, 98
- Handberg, Rasmus, Lund, Mikkel, Huber, Daniel, &

Buzasi, Derek. 2019, TESS Data For Asteroseismology Lightcurves ("TASOC"), STScI/MAST, doi: 10.17909/T9-4SMN-DX89

- Huang, C. X. 2020, TESS Lightcurves From The MIT Quick-Look Pipeline ("QLP"), STScI/MAST, doi: 10.17909/T9-R086-E880
- Lightkurve Collaboration, Cardoso, J. V. d. M., Hedges, C., et al. 2018, Lightkurve: Kepler and TESS time series analysis in Python, Astrophysics Source Code Library. http://ascl.net/1812.013
- MAST Team, 2021a, TESS "Fast" Light Curves All Sectors, STScI/MAST, doi: 10.17909/T9-ST5G-3177
- —. 2021b, TESS Light Curves All Sectors, STScI/MAST, doi: 10.17909/T9-NMC8-F686
- Paegert, M., Stassun, K. G., et al. 2021, arXiv e-prints, arXiv:2108.04778.
- Ricker, G. R., Winn, J. N., Vanderspek, R., et al. 2015, Journal of Astronomical Telescopes, Instruments, and Systems, 1, 014003, doi: 10.1117/1.JATIS.1.1.014003
- Zhou, A.-Y. 2023a, Research Notes of the AAS, 7, 210, doi: 10.3847/2515-5172/acffc2
- —. 2023b, Research Notes of the AAS, 7, 227, doi: 10.3847/2515-5172/ad06b9
- —. 2023c, Research Notes of the AAS, 7, 262, doi: 10.3847/2515-5172/ad12a3
- 2024, NewA, 105, 102081,
  doi: 10.1016/j.newast.2023.102081