

ON MANOLESCU'S κ -INVARIANTS OF RATIONAL HOMOLOGY 3-SPHERES

MASAAKI UE

ABSTRACT. We give an estimate for Manolescu's κ -invariant of a rational homology 3-sphere Y by the data of a spin 4-orbifold bounded by Y . By an appropriate choice of a 4-orbifold, sometimes we can restrict and determine the value of κ . we give such examples in case of rational homology 3-spheres obtained by Dehn surgeries along knots in S^3 .

Let Y be a rational homology 3-sphere with spin structure $\mathfrak{s}$. Then there exists a compact smooth 4-manifold with spin structure $(X, \mathfrak{s}_X)$ that satisfies $\partial(X, \mathfrak{s}_X) = (Y, \mathfrak{s})$. Manolescu's κ -invariant $\kappa(Y, \mathfrak{s})$ (which is simply denoted by $\kappa(Y)$ if Y is an integral homology 3-sphere), defined via a Seiberg-Witten Floer homotopy type of $(Y, \mathfrak{s})$ [10], [11], gives constraints on the intersection form of X [12]. In case of almost rational homology 3-spheres, Dai-Sasahira-Stoffregen [3] determines their Seiberg-Witten Floer homotopy types and the values of κ -invariants. But it is still difficult to determine $\kappa(Y, \mathfrak{s})$ in general. In this paper we give an estimate of $\kappa(Y, \mathfrak{s})$ by the data of a compact spin 4-orbifold X bounded by Y . If $b_2(X)$ is small enough, the value of $\kappa(Y, \mathfrak{s})$ is restricted and sometimes determined. In §2 we apply it to Dehn surgeries along knots in S^3 . For example, let $K_{1/n}$ be a $(1/n)$ -surgery along a knot K . Then $\kappa(K_{1/n}) = 0$ for any knot K if n is even and positive (Theorem 5), while if n is odd and positive, every integer is realized as the value of $\kappa(K_{1/n})$ for a certain knot K (an explicit example is given in Proposition 3).

1. AN ESTIMATE OF $\kappa(Y, \mathfrak{s})$ BY A SPIN 4-ORBIFOLD BOUNDED BY Y

Let $(Y, \mathfrak{s})$ be a rational homology 3-sphere with spin structure. Then Y bounds a compact connected 4-orbifold X with spin structure $\mathfrak{s}_X$, which induces $\mathfrak{s}$ on Y .

Theorem 1. *Let $(Y, \mathfrak{s})$ be a rational homology 3-sphere with spin structure, and $(X, \mathfrak{s}_X)$ be a compact connected spin 4-orbifold bounded by $(Y, \mathfrak{s})$. We assume that the set of the singular points of X consists of isolated points only, which is denoted by $\{x_i \mid i = 1, \dots, n\}$. Then a regular neighborhood of x_i is a cone cS_i over some spherical 3-manifold S_i . Let $\mathfrak{s}_i$ be the spin structure on S_i induced by $\mathfrak{s}_X$, and $k^\pm = b_2^\pm(X)$. Then $\kappa(Y, \mathfrak{s})$ has the following properties.*

(1)

$$\kappa(Y, \mathfrak{s}) \equiv -\frac{k^+ - k^-}{8} - \sum_{i=1}^n \bar{\mu}(S_i, \mathfrak{s}_i) \pmod{2\mathbf{Z}}.$$

(2)

$$\begin{aligned}
-k^+ - \frac{k^+ - k^-}{8} - \epsilon^+ - \sum_{i=1}^n \bar{\mu}(S_i, \mathfrak{s}_i) &\leq \kappa(Y, \mathfrak{s}) \\
&\leq k^- - \frac{k^+ - k^-}{8} + \epsilon^- - \sum_{i=1}^n \bar{\mu}(S_i, \mathfrak{s}_i),
\end{aligned}$$

where

$$\epsilon^- = \begin{cases} 0 & (k^- \equiv 0 \pmod{2}) \\ 1 & (k^- \equiv 1 \pmod{2}) \end{cases}, \quad \epsilon^+ = \begin{cases} 0 & (k^+ = 0) \\ -1 & (k^+ \equiv 1 \pmod{2}) \\ -2 & (k^+ \equiv 0 \pmod{2}, k > 0) \end{cases}$$

and $\bar{\mu}(S_i, \mathfrak{s}_i)$ is the Neumann-Siebenmann invariant of $(S_i, \mathfrak{s}_i)$ [13], [17].

The proof of the following corollary is straightforward from Theorem 1.

Corollary 1. *Let $\kappa_0 = -\frac{k^+ - k^-}{8} - \sum_{i=1}^n \bar{\mu}(S_i, \mathfrak{s}_i)$. Then*

$$\kappa(Y, \mathfrak{s}) = \begin{cases} \kappa_0 & \text{if } k^+ \leq 2 \text{ and } k^- = 0, \\ \kappa_0 \text{ or } \kappa_0 - 2 & \text{if } k^+ = 3, 4 \text{ and } k^- = 0, \\ \kappa_0 \text{ or } \kappa_0 + 2 & \text{if } k^+ \leq 2 \text{ and } k^- = 1, 2. \end{cases}$$

The proof of Theorem 1 is based on the following inequalities by Manolescu. (In [20] we give the outline of the proof in case of a rational homology 3-sphere.)

Theorem 2. [12]

Let $(Y_i, \mathfrak{s}_i)$ ($i = 0, 1$) be rational homology 3-spheres with spin structures and $(W, \mathfrak{s}_W)$ be a spin cobordism with $b_1(W) = 0$ from $(Y_0, \mathfrak{s}_0)$ to $(Y_1, \mathfrak{s}_1)$. Then

$$-b_2^-(W) + \frac{\sigma(W)}{8} - 1 \leq \kappa(Y_0, \mathfrak{s}_0) - \kappa(Y_1, \mathfrak{s}_1) \leq b_2^+(W) + \frac{\sigma(W)}{8} + 1.$$

The first term of the above inequality can be replaced by $-b_2^-(W) + \frac{\sigma(W)}{8}$ if $b_2^-(W)$ is even, and the last term can be replaced by $b_2^+(W) + \frac{\sigma(W)}{8}$ if $b_2^+(W)$ is even.

Furthermore if $(Y_0, \mathfrak{s}_0)$ is Floer K_G split¹, then the second inequality is improved as follows.

$$\begin{aligned}
&\kappa(Y_0, \mathfrak{s}_0) - \kappa(Y_1, \mathfrak{s}_1) \\
&\leq \begin{cases} b_2^+(W) + \frac{\sigma(W)}{8} - 1 & \text{if } b_2^+(W) \text{ is odd,} \\ b_2^+(W) + \frac{\sigma(W)}{8} - 2 & \text{if } b_2^+(W) \text{ is even and } b_2^+(W) > 0. \end{cases}
\end{aligned}$$

¹See [12] for the definition of this terminology.

Proof of Theorem 1.

Let cS_i ($i = 1, \dots, n$) be the cones over the spherical 3-manifolds S_i , which are the regular neighborhoods of the singular points of X . If $n \geq 2$, S_i 's can be connected by 1-handles in $X \setminus \text{Int} \cup_{i=1}^n cS_i$ to form a spin cobordism $W \subset X \setminus \text{Int} \cup_{i=1}^n cS_i$ from $\cup_{i=1}^n S_i$ to the connected sums $S_0 = \sharp_{i=1}^n S_i$ of S_i 's. Then $X_0 = X \setminus \text{Int}(\cup_{i=1}^n cS_i \cup W)$ is a spin cobordism from S_0 to Y . Let $\mathfrak{s}_0, \mathfrak{s}_W, \mathfrak{s}_{X_0}$ be the spin structures on S_0, W , and X_0 induced by $\mathfrak{s}_X$ respectively.

Since S_i ($i \geq 0$) has a metric g_i of positive scalar curvature, $\kappa(S_i, \mathfrak{s}_i)$ is related to the correction term $n(S_i, \mathfrak{s}_i, g_i)$ of the Seiberg-Witten-Floer homotopy type of $(S_i, \mathfrak{s}_i)$ with respect to the metric g_i by the following proposition.

Proposition 1. [12] *Let $(Y, \mathfrak{s})$ be a rational homology 3-sphere with spin structure that admits a metric g of positive scalar curvature. Then $\kappa(Y, \mathfrak{s}) = -n(Y, \mathfrak{s}, g)$ and $(Y, \mathfrak{s})$ is Floer K_G split.*

If $i \geq 1$, $\kappa(S_i, \mathfrak{s}_i)$ is also related to the Fukumoto-Furuta invariant $w(S_i, cS_i, \mathfrak{s}_{cS_i})$, where $\mathfrak{s}_{cS_i}$ is the restriction of $\mathfrak{s}_X$ to cS_i .

Definition 1. Let $(X, \mathfrak{s}_X)$ be a compact spin 4-orbifold with spin structure $\mathfrak{s}_X$ bounded by a rational homology 3-sphere $(Y, \mathfrak{s})$ with spin structure. Choose a compact spin 4-manifold $(X', \mathfrak{s}_{X'})$ with $\partial(X', \mathfrak{s}_{X'}) = (Y, \mathfrak{s})$ and put $(Z, \mathfrak{s}_Z) = (X \cup (-X'), \mathfrak{s}_X \cup \mathfrak{s}_{X'})$. Then the Fukumoto-Furuta invariant $w(Y, X, \mathfrak{s}_X)$ is defined to be

$$w(Y, X, \mathfrak{s}_X) = -\text{ind}_{\mathbb{C}} \mathcal{D}_Z(\mathfrak{s}_Z) + \frac{\sigma(X')}{8},$$

where $\mathcal{D}_Z(\mathfrak{s}_Z)$ is the Dirac operator of $(Z, \mathfrak{s}_Z)$. The value of $w(Y, X, \mathfrak{s}_X)$ does not depend on the choice of X' . (The sign of w is opposite to the original one in [4].)

Since $\dim \ker \mathcal{D}_{S_i}(\mathfrak{s}_i) = 0$, $n(S_i, \mathfrak{s}_i, g_i)$ ($i \geq 1$) is represented by the eta invariants of the Dirac and the signature operators on S_i as follows:

$$n(S_i, \mathfrak{s}_i, g_i) = -\frac{1}{8}(4\eta^{\text{Dir}}(S_i, \mathfrak{s}_i, g_i) + \eta^{\text{sign}}(S_i, g_i)).$$

Here we assume that the Clifford multiplication c of the volume form vol_{S_i} on S_i satisfies $c(\text{vol}_{S_i}) = -1$ as in [20]. If we choose c satisfying $c(\text{vol}_{S_i}) = 1$, $\eta^{\text{Dir}}(S_i, \mathfrak{s}_i, g_i)$ should be replaced with $-\eta^{\text{Dir}}(S_i, \mathfrak{s}_i, g_i)$ ([20], Remark 4).

The right hand side of the above formula coincides with $w(S_i, cS_i, \mathfrak{s}_{cS_i})$, which is the same as $\bar{\mu}(S_i, \mathfrak{s}_i)$ ([20]), and hence

$$(*) \quad \kappa(S_i, \mathfrak{s}_i) = -n(S_i, \mathfrak{s}_i, g_i) = -\bar{\mu}(S_i, \mathfrak{s}_i).$$

By considering the spin cobordism W from $\cup_{i=1}^n S_i$ to S_0 , we also have the relation $n(S_0, \mathfrak{s}_0, g_0) = \sum_{i=1}^n n(S_i, \mathfrak{s}_i, g_i)$ ([20]). It follows that

$$(**) \quad \kappa(S_0, \mathfrak{s}_0) = -n(S_0, \mathfrak{s}_0, g_0) = -\sum_{i=1}^n n(S_i, \mathfrak{s}_i, g_i) = -\sum_{i=1}^n \bar{\mu}(S_i, \mathfrak{s}_i).$$

Then applying the inequality in Theorem 2 (note that $(S_0, \mathfrak{s}_0)$ is Floer K_G split) to the spin cobordism X_0 and $(**)$, we obtain the inequality in Theorem 1 (2). We note that for a rational homology 3-sphere with spin structure $(Y, \mathfrak{s})$, the mod $2\mathbf{Z}$ reduction of $\bar{\mu}(Y, \mathfrak{s})$ is the Rokhlin invariant $\mu(Y, \mathfrak{s})$ of $(Y, \mathfrak{s})$. We also have $\kappa(Y, \mathfrak{s}) \equiv -n(Y, \mathfrak{s}, g) \equiv -\mu(Y, \mathfrak{s}) \pmod{2\mathbf{Z}}$ for some metric g . Thus by the additivity of signature, if we choose a spin 4-manifold $(W_i, \mathfrak{s}_{W_i})$ with $\partial(W_i, \mathfrak{s}_{W_i}) = (S_i, \mathfrak{s}_i)$, we have

$$\kappa(Y, \mathfrak{s}) \equiv -\bar{\mu}(Y, \mathfrak{s}) \equiv -\frac{1}{8}(\sigma(X_0) + \sum_{i=1}^n \sigma(W_i)) \equiv -\frac{1}{8}\sigma(X_0) - \sum_{i=1}^n \bar{\mu}(S_i, \mathfrak{s}_i) \pmod{2\mathbf{Z}}.$$

This proves Theorem 1 (1). $\square$

On the other hand, $w(Y, X, \mathfrak{s}_X)$ for (Y, X) in Theorem 1 is represented via the eta invariants of S_i 's and $(*)$ as follows ([20]).

$$w(Y, X, \mathfrak{s}_X) = \frac{\sigma(X)}{8} + \sum_{i=1}^n \bar{\mu}(S_i, \mathfrak{s}_i).$$

Furthermore $w(Y, X, \mathfrak{s}_X) \equiv \mu(Y, \mathfrak{s}) \pmod{2\mathbf{Z}}$ since $\dim \text{ind}_{\mathbf{C}} \mathcal{D}_Z(\mathfrak{s}_Z)$ is even, where Z is a closed spin orbifold in the definition of $w(Y, X, \mathfrak{s}_X)$. The following theorem is deduced from these facts and Theorem 1.

Theorem 3. *Let $(Y, \mathfrak{s})$ be a rational homology 3-sphere with spin structure and $(X, \mathfrak{s}_X)$ be a spin 4-orbifold bounded by $(Y, \mathfrak{s})$ as in Theorem 1. Then*

$$\begin{aligned} \kappa(Y, \mathfrak{s}) + w(Y, X, \mathfrak{s}_X) &\equiv 0 \pmod{2\mathbf{Z}}, \\ -k^+ - \epsilon^+ &\leq \kappa(Y, \mathfrak{s}) + w(Y, X, \mathfrak{s}_X) \leq k^- + \epsilon^-. \end{aligned}$$

If Y is a Seifert rational homology 3-sphere, we can choose X so that $w(Y, X, \mathfrak{s}_X) = \bar{\mu}(Y, \mathfrak{s})$ and $k^\pm$ are small, which restricts the value of $\kappa(Y, \mathfrak{s}) + \bar{\mu}(Y, \mathfrak{s})$ ([20]).

We also note that the κ -invariant of an almost rational (AR) homology 3-sphere is determined by Dai-Sasahira-Stoffregen [3].

Theorem 4. [3] *Let $(Y, \mathfrak{s})$ be an AR homology 3-sphere with spin structure. Then either one of the followings holds.*

- (1) *If $-\bar{\mu}(Y, \mathfrak{s}) = \delta(Y, \mathfrak{s})$, then $\kappa(Y, \mathfrak{s}) = -\bar{\mu}(Y, \mathfrak{s})$.*
- (2) *If $-\bar{\mu}(Y, \mathfrak{s}) < \delta(Y, \mathfrak{s})$, then $\kappa(Y, \mathfrak{s}) = -\bar{\mu}(Y, \mathfrak{s}) + 2$.*

Here $\delta(Y, \mathfrak{s})$ is the monopole Frøyshov invariant of $(Y, \mathfrak{s})$. Note that we have always $-\bar{\mu}(Y, \mathfrak{s}) \leq \delta(Y, \mathfrak{s})$ [1].

Remark 1. In [6], [8], the Frøyshov invariant is defined as $h(Y, \mathfrak{s}) = -\delta(Y, \mathfrak{s})$ so that $d(Y, \mathfrak{s}) = -2h(Y, \mathfrak{s})$, where $d(Y, \mathfrak{s})$ is the Ozsváth-Szabó's correction term [15].

For an AR homology 3-sphere Y , the correction term $\underline{d}(Y, \mathfrak{s})$ of the involutive Floer homology is equal to $-2\bar{\mu}(Y, \mathfrak{s})$ [2]. If Y is also an L -space, then the correction terms $\underline{d}(Y, \mathfrak{s})$, $d(Y, \mathfrak{s})$, $\bar{d}(Y, \mathfrak{s})$ are all the same ([7]). Since $\delta(Y, \mathfrak{s}) = d(Y, \mathfrak{s})/2$, we have $-\bar{\mu}(Y, \mathfrak{s}) = \delta(Y, \mathfrak{s}) = \kappa(Y, \mathfrak{s})$ by the above theorem. (*) is also deduced from this fact because a spherical 3-manifold is rational and also an L -space ([14]).

2. κ -INVARIANTS OF DEHN SURGERIES ALONG KNOTS IN S^3

We apply Theorem 1 to a rational homology 3-sphere obtained by Dehn surgery along a knot K in S^3 . We denote by $K_{p/q}$ the (p/q) -surgery along K . The set of spin structures on the complement $X_K := S^3 \setminus \text{Int}N(K)$ of the tubular neighborhood $N(K)$ of K is identified with $\text{Hom}_{\mathbf{Z}_2}(H_1(X_K, \mathbf{Z}_2), \mathbf{Z}_2) \cong \mathbf{Z}_2$. Hence there are 2 spin structures $\mathfrak{s}_0$ and $\mathfrak{s}_1$ on X_K . Here $\mathfrak{s}_0$ is the one that extends to the unique spin structure on S^3 , and $\mathfrak{s}_1$ is the one that does not extend to S^3 . The spin structure on X_K corresponding to $c \in \text{Hom}_{\mathbf{Z}_2}(H_1(X_K, \mathbf{Z}_2), \mathbf{Z}_2)$ extends to $K_{p/q}$ if and only if $pc(\mu) \equiv pq \pmod{2}$, where μ is the meridian of K . It follows that only $\mathfrak{s}_0$ (resp. $\mathfrak{s}_1$) extends to the spin structure on $K_{p/q}$ if p is odd and q is even (resp. both p and q are odd), while both $\mathfrak{s}_0$ and $\mathfrak{s}_1$ extend to $K_{p/q}$ if p is even and q is odd. We denote the extended spin structures on $K_{p/q}$ by the same symbols as those on X_K .

Theorem 5. *Let K be any knot in S^3 , and p and q are positive integers such that $\gcd(p, q) = 1$, $p + q \equiv 1 \pmod{2}$. Then*

$$\begin{aligned}\kappa(K_{p/q}, \mathfrak{s}_0) &= \kappa(L(p, -q), \mathfrak{s}_0), \\ \kappa(K_{-p/q}, \mathfrak{s}_0) &= \kappa(L(p, q), \mathfrak{s}_0) \text{ or } \kappa(L(p, q), \mathfrak{s}_0) + 2.\end{aligned}$$

Here $\mathfrak{s}_0$ on $L(p, \pm q)$ is the spin structure $\mathfrak{s}_0$ on the $\mp p/q$ -surgery along the unknot, which is $L(p, \pm q)$. We note that $\kappa(L(p, \pm q), \mathfrak{s}_0) = -\bar{\mu}(L(p, \pm q), \mathfrak{s}_0)$. In particular, we have


$$\kappa(K_{1/2n}, \mathfrak{s}_0) = \begin{cases} 0 & \text{if } n > 0, \\ 0 \text{ or } 2 & \text{if } n < 0. \end{cases}$$

Proof. First suppose that $p > q > 1$. Since $p + q \equiv 1 \pmod{2}$, we have the continued fraction expansion of p/q of the following form.

$$(\dagger) \quad \frac{p}{q} = [\alpha_1, \alpha_2, \dots, \alpha_n] = \alpha_1 - \frac{1}{\alpha_2 - \frac{1}{\dots - \frac{1}{\alpha_n}}}$$

where $n \geq 2$ and $\alpha_i \equiv 0 \pmod{2}$ and $|\alpha_i| \geq 2$ for all α_i . Then $K_{p/q}$ bounds a 4-manifold X represented by the framed link $\mathcal{L}$ in Figure 1.

Since every α_i is even, X has a spin structure $\mathfrak{s}_X$ whose restriction to $K_{p/q}$ is $\mathfrak{s}_0$ (see Remark 2 (1) below). The sublink $\mathcal{L}_0$ of $\mathcal{L}$ consisting of the components of framing α_i

FIGURE 1. The framed link $\mathcal{L}$

($i \geq 2$) represents a plumbing X_Γ whose boundary is a lens space $L(p', -q')$, where $p'/q' = [\alpha_2, \dots, \alpha_n]$. We can push X_Γ into $\text{Int}X$ and attach the cone $cL(p', -q')$ over $L(p', -q')$ to the complement $X_0 = X \setminus \text{Int}X_\Gamma$ along $L(p', -q')$ to form a 4-orbifold $\widehat{X}$. The spin structure $\mathfrak{s}_X$ over X_0 can be extended to $cL(p', -q')$ and form the orbifold spin structure $\mathfrak{s}_{\widehat{X}}$ on $\widehat{X}$ (This procedure is similar to those in [16], [19]). The intersection matrices of X and X_Γ are congruent to the diagonal matrices whose diagonal entries are $\{[\alpha_i, \alpha_{i-1}, \dots, \alpha_n] \mid i \geq 1\}$ and $\{[\alpha_i, \alpha_{i-1}, \dots, \alpha_n] \mid i \geq 2\}$ respectively. Hence $\sigma(X_0) = \sigma(X) - \sigma(X_\Gamma)$ is the sign of $[\alpha_1, \dots, \alpha_n] = p/q$, which is 1. Since $b_2(X_0) = b_2(X) - b_2(X_\Gamma) = 1$, we have $b_2^+(\widehat{X}) = b_2^+(X_0) = 1$ and $b_2^-(\widehat{X}) = b_2^-(X_0) = 0$ and hence by Corollary 1,

$$\kappa(K_{p/q}, \mathfrak{s}_0) = -\frac{1}{8} - \bar{\mu}(L(p', -q'), \mathfrak{s}'),$$

where $\mathfrak{s}'$ is the restriction of $\mathfrak{s}_X$ to $L(p', -q')$. Since $\text{sgn}[\alpha_i, \dots, \alpha_n] = \text{sgn} \alpha_i$ and $\text{sgn} \alpha_1 = 1$ and the characteristic sublink of $\mathcal{L}_0$ associated with $\mathfrak{s}'$ is empty, we have $\bar{\mu}(L(p', -q'), \mathfrak{s}') = \frac{1}{8}\sigma(X_\Gamma)$ and

$$\kappa(K_{p/q}, \mathfrak{s}_0) = -\frac{1}{8}(1 + \sigma(X_\Gamma)) = -\frac{1}{8} \sum_{i=1}^n \text{sgn} \alpha_i,$$

which is the same as $-\bar{\mu}(L(p, -q), \mathfrak{s}_0) = \kappa(L(p, -q), \mathfrak{s}_0)$. In fact, $\kappa(K_{p/q}, \mathfrak{s}_0)$ does not depend on K , and hence is the same as the κ -invariant of the p/q -surgery along the unknot, which is $L(p, -q)$. In case of $K_{-p/q}$, we form a spin 4-manifold X' with $\partial X' = K_{-p/q}$ represented by a framed link $\mathcal{L}'$, which is obtained from $\mathcal{L}$ by replacing the framings α_i with $-\alpha_i$. There is a plumbing $-X_\Gamma$ corresponding to the sublink of $\mathcal{L}'$, which consists of the components of framing $-\alpha_i$ ($i \geq 2$) with $\partial(-X_\Gamma) = L(p', q')$ in X' . Then by replacing it with the cone over $L(p', q')$, we also obtain the spin 4-orbifold $\widehat{X}'$. In this case $b_2^+(\widehat{X}') = 0$ and $b_2^-(\widehat{X}') = 1$ and hence by Corollary 1,

$$\kappa(K_{-p/q}, \mathfrak{s}_0) = \frac{1}{8} - \bar{\mu}(L(p', q'), \mathfrak{s}') \text{ or } 2 + \frac{1}{8} - \bar{\mu}(L(p', q'), \mathfrak{s}'),$$

where $\mathfrak{s}'$ is the restriction of the spin structure on X' to $L(p', q')$. Here we have

$$\frac{1}{8} - \bar{\mu}(L(p', q'), \mathfrak{s}') = \frac{1}{8} \sum_{i=1}^n \text{sgn } \alpha_i = -\frac{1}{8} \bar{\mu}(L(p, q), \mathfrak{s}_0) = \kappa(L(p, q), \mathfrak{s}_0).$$

If p is even and $q = 1$, K_p bounds a spin 4-manifold X represented by a framed link K with framing p whose spin structure $\mathfrak{s}_X$ induces $\mathfrak{s}_0$ on K_p . In this case we apply Manolescu inequality in Theorem 2 (by putting $Y_0 = S^3$) directly to $(X, \mathfrak{s}_X)$. Then $\kappa(K_p, \mathfrak{s}_0) = -1/8 = -\bar{\mu}(L(p, -1), \mathfrak{s}_0) = \kappa(L(p, -1), \mathfrak{s}_0)$, and $\kappa(K_{-p}, \mathfrak{s}_0)$ is either $\kappa(L(p, 1), \mathfrak{s}_0)$ or $\kappa(L(p, 1), \mathfrak{s}_0) + 2$ since $\kappa(Y_{\pm p}, \mathfrak{s}_0) \equiv -\mu(Y_{\pm p}, \mathfrak{s}_0) \equiv \mp 1/8 \pmod{2\mathbf{Z}}$. Thus we prove Theorem 5 in the first case.

Next suppose that $q > p > 0$. Then there is a continued fraction expansion of q/p of the form

$$\frac{q}{p} = [\beta_2, \dots, \beta_n] \quad n \geq 2, \quad |\beta_i| \geq 2, \quad \beta_i \equiv 0 \pmod{2} \text{ for all } i \geq 2.$$

In this case $K_{p/q}$ bounds a spin 4-manifold X' represented by a framed link $\mathcal{L}'$, which is obtained from the above $\mathcal{L}$ by replacing the framing α_1 with 0, and α_i with $-\beta_i$ for $i \geq 2$. The spin structure $\mathfrak{s}_{X'}$ on X' induces $\mathfrak{s}_0$ on $K_{p/q}$. Then we have a plumbing X'_Γ in X' corresponding to the sublink, which consists of the components of framing $-\beta_i$ ($i \geq 2$). The boundary of X'_Γ is a lens space $L(p', q')$, where $p'/q' = [\beta_2, \dots, \beta_n]$. Again we can push X'_Γ into $\text{Int} X'$ and replace it with the cone $cL(p', q')$ over $L(p', q')$ to form a spin 4-orbifold $\widehat{X}'$. The intersection matrices of X' and X'_Γ are congruent to the diagonal matrices whose diagonal entries are $\{1/[\beta_2, \dots, \beta_n], -[\beta_2, \dots, \beta_n], \dots, -\beta_n\}$ and $\{-[\beta_2, \dots, \beta_n], \dots, -\beta_n\}$ respectively. Hence $\sigma(\widehat{X}')$ is the sign of $1/[\beta_2, \dots, \beta_n] = p/q$ and hence 1. It follows that $b_2^+(\widehat{X}') = 1$ and $b_2^-(\widehat{X}') = 0$. Thus again by Corollary 1 we have

$$\begin{aligned} \kappa(K_{p/q}, \mathfrak{s}_0) &= -\frac{1}{8} - \bar{\mu}(L(p', q'), \mathfrak{s}') = -\frac{1}{8} (1 - \sum_{i=2}^n \text{sgn}[\beta_i, \dots, \beta_n]) \\ &= -\frac{1}{8} (\text{sgn}(1/[\beta_2, \dots, \beta_n]) + \sum_{i=2}^n \text{sgn}[-\beta_i, \dots, -\beta_n]) \quad (\mathfrak{s}' = \mathfrak{s}_{X'}|_{L(p', q')}), \end{aligned}$$

which is the same as $-\bar{\mu}(L(p, -q), \mathfrak{s}_0) = \kappa(L(p, -q), \mathfrak{s}_0)$. In case of $K_{-p/q}$, it bounds a spin 4-manifold X'' represented by a framed link $\mathcal{L}''$, which is obtained from $\mathcal{L}'$ by replacing the framing $-\beta_i$ with β_i ($i \geq 2$). The spin structure $\mathfrak{s}_{X''}$ also induces $\mathfrak{s}_0$ on $K_{-p/q}$. Then we obtain the spin 4-orbifold $\widehat{X}''$ by deleting the plumbing $-X'_\Gamma$ with $\partial(-X'_\Gamma) = L(p', -q')$ from X'' and attaching the cone $cL(p', -q')$ over $L(p', -q')$. In this case $b_2^+(\widehat{X}'') = 0$ and $b_2^-(\widehat{X}'') = 1$ and hence by Corollary 1 we have

$$\kappa(K_{-p/q}, \mathfrak{s}_0) = \frac{1}{8} - \bar{\mu}(L(p', -q'), \mathfrak{s}') \text{ or } 2 + \frac{1}{8} - \bar{\mu}(L(p', -q'), \mathfrak{s}'),$$

where $\frac{1}{8} - \bar{\mu}(L(p', -q'), \mathfrak{s}') = -\bar{\mu}(L(p, q), \mathfrak{s}_0) = \kappa(L(p, q), \mathfrak{s}_0)$. Thus we obtain the required result in the second case. $\square$

On the other hand, the value of κ may depend on the choice of K in case of a p/q surgery with p and q odd. In fact, examples below show the following proposition.

Proposition 2. *Every integer is realized as the value of the κ -invariant of a $(+1)$ surgery along some torus knot.*

Remark 2. (1) Let L_i be the component of $\mathcal{L}$ with framing α_i ($L_1 = K$), and μ_i be a meridian of L_i . Then L_i and μ_i are oriented so that we have the following relations in $H_1(K_{p/q}, \mathbf{Z})$.

$$\begin{pmatrix} -\alpha_i & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} \mu_i \\ \mu_{i-1} \end{pmatrix} = \begin{pmatrix} \mu_{i+1} \\ \mu_i \end{pmatrix} \quad (1 \leq i \leq n),$$

where (μ_1, μ_0) corresponds to the pair (μ, λ) of the meridian and the preferred longitude of $K \subset S^3$, and (μ_{n+1}, μ_n) corresponds to the pair of the meridian and the longitude of the solid torus attached to X_K by the (p/q) surgery along K (and hence $\mu_0 = \mu_{n+1} = 0$ in $H_1(K_{p/q}, \mathbf{Z})$). If we put

$$\begin{pmatrix} \tilde{p} & \tilde{q} \\ \tilde{r} & \tilde{s} \end{pmatrix} = \begin{pmatrix} -\alpha_n & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} -\alpha_{n-1} & -1 \\ 1 & 0 \end{pmatrix} \cdots \begin{pmatrix} -\alpha_1 & -1 \\ 1 & 0 \end{pmatrix},$$

then $(\tilde{p}, \tilde{q}) = \pm(p, q)$. We note that the set of spin structures on $K_{p/q}$ is also identified with the set of $\tilde{c} \in \text{Hom}(H_1(S^3 \setminus \cup_{i=1}^n L_i, \mathbf{Z}), \mathbf{Z}_2)$ satisfying

$$\sum_{j=1}^n \text{lk}(L_i, L_j) \tilde{c}(\mu_j) \equiv \text{lk}(L_i, L_i) = \alpha_i \pmod{2} \quad (1 \leq i \leq n).$$

It follows that $\tilde{c} \equiv 0$ (corresponding to the spin structure on $K_{p/q}$ induced by $\mathfrak{s}_X$), if and only if $c(\mu) \equiv 0$ (corresponding to $\mathfrak{s}_0$) since α_i is even. On the other hand, if p is even and q is odd, then n is odd in the continued fraction $(\dagger)$. Hence $\tilde{c} \in \text{Hom}(H_1(S^3 \setminus \cup_{i=1}^n L_i, \mathbf{Z}), \mathbf{Z}_2)$ satisfying

$$\tilde{c}(\mu_i) \equiv 1 \pmod{2} \quad \text{if } i \text{ is odd, } \tilde{c}(\mu_i) \equiv 0 \pmod{2} \quad \text{if } i \text{ is even}$$

corresponds to $\mathfrak{s}_1$, whose associated characteristic sublink of $\mathcal{L}$ is $\cup_{i \text{ odd}} L_i$.

- (2) $K_{1/2m}$ bounds a compact spin 4-manifold X represented by a framed link consisting of K and its meridian with framings 0 and $-2m$ respectively. A direct application of Theorem 2 (by putting $Y_0 = S^3$) to X shows that $\kappa(K_{1/2m}, \mathfrak{s}_0) = 0$ or 2. Theorem 5 slightly improves the estimate of κ when $m > 0$.
- (3) If we reverse the orientation of $K_{p/q}$, we obtain $\overline{K}_{-p/q}$, where $\overline{K}$ is the mirror image of K . In case of $K_{p/q}$ with $p > 0$, $q > 0$ and $p + q \equiv 1 \pmod{2}$, the computation of the above theorem shows that

$$\kappa(\overline{K}_{-p/q}, \mathfrak{s}_0) = -\kappa(K_{p/q}, \mathfrak{s}_0) \text{ or } -\kappa(K_{p/q}, \mathfrak{s}_0) + 2.$$




FIGURE 2. Γ corresponding to $\Sigma(2, 4k + 1, 8k + 3)$

We can see that both cases actually occur (examples below).

- (4) Let $(Y, \mathfrak{s})$ be a rational homology 3-sphere Y with spin structure $\mathfrak{s}$, which bounds a compact 4-manifold X represented by a framed link $\mathcal{L}$. Suppose that there is a sublink $\mathcal{L}_0$ of $\mathcal{L}$ which forms a plumbing X_Γ whose boundary is a spherical 3-manifold S , and contains the characteristic sublink of $\mathcal{L}$ corresponding to $\mathfrak{s}$. Then by replacing X_Γ with the cone over S we obtain a spin 4-orbifold with smaller Betti numbers, from which we could restrict the value of $\kappa(Y, \mathfrak{s})$. In case of $(K_{p/q}, \mathfrak{s}_1)$ with $p + q \equiv 1 \pmod{2}$ or $(K_{p/q}, \mathfrak{s}_1)$ with $p \equiv q \equiv 1 \pmod{2}$, $K_{p/q}$ bounds a compact 4-manifold X represented by a framed link $\mathcal{L}$ similar to Figure 1, which is obtained by the continued fraction expansion of p/q . But in either case, we cannot obtain an orbifold by collapsing the characteristic sublink $\mathcal{L}_0$ of $\mathcal{L}$ corresponding to $\mathfrak{s}_1$ if K is nontrivial, because $\mathcal{L}_0$ contains a component K .

Some examples.

Let $T_{p,q}$ be the right-handed (p, q) torus knot. Then the $(1/n)$ -surgery along $T_{p,q}$ is diffeomorphic to a Brieskorn homology 3-sphere up to orientation as follows:


$$\begin{aligned} (T_{p,-q})_{-1/n} &\cong \Sigma(p, q, pqn - 1), & (T_{p,q})_{-1/n} &\cong \Sigma(p, q, pqn + 1), \\ (T_{p,q})_{1/n} &\cong -\Sigma(p, q, pqn - 1), & (T_{p,-q})_{1/n} &\cong -\Sigma(p, q, pqn + 1) \end{aligned}$$

for $p, q, n > 0$. Then by Theorem 5, $\kappa(-\Sigma(p, q, 2pqm - 1)) = \kappa(-\Sigma(p, q, 2pqm + 1)) = 0$, while $\kappa(\Sigma(p, q, 2pqm - 1))$ and $\kappa(\Sigma(p, q, 2pqm + 1))$ are either 0 or 2 if $m > 0$, and both cases occur. For example, $(T_{2,3})_{-1/2} \cong \Sigma(2, 3, 13)$ bounds a contractible manifold, and hence $\kappa(\Sigma(2, 3, 13)) = 0$. On the other hand $(T_{2,-3})_{-1/2m} \cong \Sigma(2, 3, 12m - 1)$ and $\kappa(\Sigma(2, 3, 12m - 1)) = 2$ [12]. Another example is $(T_{2,-7})_{-1/2k} \cong \Sigma(2, 7, 28k - 1)$. In this case $\bar{\mu} = -\beta = 0$ and $\delta = 2$ ([18]), and hence $\kappa(\Sigma(2, 7, 28k - 1)) = 2$ by Theorem 4.

On the other hand, $\kappa(K_{1/n})$ for n odd may vary according to the choice of K . For example, $(T_{2,-4k-1})_1 \cong -\Sigma(2, 4k + 1, 8k + 3)$, where $\Sigma(2, 4k + 1, 8k + 3)$ has the Seifert invariants of the form

$$\{(2, 1), (4k + 1, 2k), (8k + 3, -8k - 2)\}$$

(we follow the sign convention in [9], which is opposite to those in [5], [19]), and bounds a spin plumbing $X(\Gamma)$ corresponding to the weighted graph Γ in Figure 2.

FIGURE 3. $K(4c, 2d + 1)$ and $\mathcal{L}$

Hence we have $\bar{\mu}(\Sigma(2, 4k + 1, 8k + 3)) = \frac{1}{8}\sigma(X(\Gamma)) = k$. (We note that $\sigma(X(\Gamma))$ is the sum of the sign of $\deg(\partial X(\Gamma))$ (which is -1) and the signs of the weights of the vertices of Γ other than the central one in Figure 2.) In [20] we claim that if a Seifert rational homology 3-sphere Y contains a singular fiber of even multiplicity and has positive degree, then $\kappa(Y, \mathfrak{s}) = -\bar{\mu}(Y, \mathfrak{s})$. It follows that

$$\kappa(-\Sigma(2, 4k + 1, 8k + 3)) = -\bar{\mu}(-\Sigma(2, 4k + 1, 8k + 3)) = \bar{\mu}(\Sigma(2, 4k + 1, 8k + 3)) = k.$$

Likewise $(T_{2,4k+1})_1 \cong -\Sigma(2, 4k + 1, 8k + 1)$, where $\Sigma(2, 4k + 1, 8k + 1)$ has the Seifert invariants of the form

$$\{(2, -1), (4k + 1, -2k), (8k + 1, 8k)\}$$

and hence

$$\kappa(-\Sigma(2, 4k + 1, 8k + 1)) = \bar{\mu}(\Sigma(2, 4k + 1, 8k + 1)) = -k,$$

by computation similar to the above case. This proves Proposition 2.

As another example, we consider the knot $K(4c, 2d + 1)$ with $c \neq 0$ in Figure 3. Here $\boxed{\ell}$ means ℓ times right-handed half twists if $\ell > 0$, and $|\ell|$ times left-handed half twists if $\ell < 0$.

Proposition 3. *Let $Y = K(4c, 2d + 1)_{1/n}$ with $c \neq 0$ and n odd. Then for the unique spin structure $\mathfrak{s}$ on Y , we have*

$$\kappa(Y, \mathfrak{s}) = \begin{cases} -c & \text{if } n > 0, \\ -c \text{ or } -c + 2 & \text{if } n < 0. \end{cases}$$

We note that $\kappa(Y, \mathfrak{s}) = 0$ if $c = 0$ since $K(0, 2d + 1)$ is the unknot and hence $Y = S^3$. To prove Proposition 3, we consider a 4-manifold X represented by a framed link $\mathcal{L}$ in Figure 3, which is bounded by Y . We denote the components of $\mathcal{L}$ by L_1, L_2, L_3 and L_4 from left to right. Let x_i be the basis of $H_2(X, \mathbf{Z})$ represented by a union of the core of the 2-handle attached along L_i and the cone over L_i in the 4-ball. Then the characteristic sublink of $\mathcal{L}$ associated with $\mathfrak{s}$ is L_3 , and the Poincare

dual of $x_3 \bmod 2$ is the obstruction to extending $\mathfrak{s}$ to the spin structure on X . Then the 4-manifolds $\tilde{X}$ and $\tilde{X}'$ represented by L_1 and $L_3 \cup L_4$ have boundaries $L(2c, -1)$ and $L(p, q)$ respectively, where $p/q = [8c, n]$ (since L_3 is also the unknot). Hence by pushing $\tilde{X}$ and $\tilde{X}'$ separately into $\text{Int}X$ and replacing them with the cones over $L(2c, -1)$ and $L(p, q)$, we obtain a 4-orbifold $\hat{X}$ with spin structure $\hat{\mathfrak{s}}$. Let $\tilde{\mathfrak{s}}$ and $\tilde{\mathfrak{s}}'$ be the spin structures on $L(2c, -1)$ and $L(p, q)$ induced from $\hat{\mathfrak{s}}$ respectively. Then

$$\begin{aligned}\bar{\mu}(L(2c, -1), \tilde{\mathfrak{s}}) &= \frac{1}{8}(\sigma(\tilde{X})) = \frac{1}{8} \text{sgn } c, \\ \bar{\mu}(L(p, q), \tilde{\mathfrak{s}}') &= \frac{1}{8}(\sigma(\tilde{X}') - x_3^2) = \frac{1}{8}(-\text{sgn}[8c, n] - \text{sgn } n + 8c),\end{aligned}$$

since $\tilde{\mathfrak{s}}$ extends to the spin structure on $\tilde{X}$, while the characteristic sublink of $L_3 \cup L_4$ associated with $\tilde{\mathfrak{s}}'$ is also L_3 . The intersection matrix of X is congruent to the diagonal matrix whose diagonal entries are $\{1/4n, 4/[8c, n], -[8c, n], -n\}$ and hence $\sigma(X) = 0$. It follows that $b_2(\hat{X}) = 1$ and $\sigma(\hat{X}) = \sigma(X) - \sigma(\tilde{X}) - \sigma(\tilde{X}') = \text{sgn } n$ since $\text{sgn}[8c, n] = \text{sgn } c$. Hence $b_2^+(\hat{X}) = 1$ and $b_2^-(\hat{X}) = 0$ if $n > 0$, and $b_2^+(\hat{X}) = 0$ and $b_2^-(\hat{X}) = 1$ if $n < 0$. Thus we deduce from Corollary 1 that

$$\kappa(Y, \mathfrak{s}) = -\frac{\sigma(\hat{X})}{8} - (\bar{\mu}(L(2c, -1), \tilde{\mathfrak{s}}) + \bar{\mu}(L(p, q), \tilde{\mathfrak{s}}')) = -\frac{1}{8}(1 - \text{sgn } n + 8c) = -c$$

if $n > 0$, and $\kappa(Y, \mathfrak{s}) = -c$ or $-c + 2$ if $n < 0$.

We note that $K(4c, -1) = T_{2, 4c+1}$ if $c > 0$, and hence if $n > 0$ and n is odd,

$$\kappa(K(4c, -1)_{1/n}) = \kappa(-\Sigma(2, 4c+1, (8c+2)n-1)) = \bar{\mu}(\Sigma(2, 4c+1, (8c+2)n-1)),$$

which is $-c$ by computation similar to the above case.

On the other hand, if we consider $K(2c, 2d)$ (the double twist knot with c times full twists and d times full twists) instead, $K(2c, 2d)_{1/n}$ is diffeomorphic to the Dehn surgery along the Borromean rings with framings $-1/c$, $-1/d$, and $1/n$ respectively, and also diffeomorphic to $K(2c, -2n)_{-1/d}$ and $K(2d, -2n)_{-1/c}$ by the symmetry of the Borromean rings. Hence if c or d is even, $\kappa(K(2c, 2d)_{1/n}) = 0$ or 2 , and if $c < 0$ or $d < 0$ in addition, it is 0 by Theorem 5 for every nonzero integer n .

Proposition 3 can be generalized to the case of the $\pm p/q$ -surgery along $K := K(4c, 2d+1)$ ($c \neq 0$) as follows.

Proposition 4. *Let $K = K(4c, 2d+1)$ with $c \neq 0$ and $\mathfrak{s}_1$ be the spin structure on $K_{\pm p/q}$ defined as in the first paragraph of this section.*

- (1) *Suppose that $p \equiv 0 \pmod{2}$, $q \equiv 1 \pmod{2}$, and $p > q > 0$. Then there exist $\beta_i \in \mathbf{Z}$ ($1 \leq i \leq n$) such that $p/q = [\beta_1, \dots, \beta_n]$, $\beta_i \equiv 0 \pmod{2}$, and*

$|\beta_i| \geq 2$ (note that n is odd in this case). Let

$$\begin{aligned}\kappa_0 &= -\frac{1}{8}(1 + \sum_{i=2}^n \operatorname{sgn} \beta_i - \sum_{i \text{ odd}} \beta_i) - c, \\ \kappa'_0 &= \frac{1}{8}(1 + \sum_{i=2}^n \operatorname{sgn} \beta_i - \sum_{i \text{ odd}} \beta_i) - c.\end{aligned}$$

Then

$$\begin{aligned}\kappa(K_{p/q}, \mathfrak{s}_1) &= \begin{cases} \kappa_0 & (\text{if } c > \frac{p}{8q} \text{ or } c < 0), \\ \kappa_0 \text{ or } \kappa_0 + 2 & (\text{if } 0 < c < \frac{p}{8q}), \end{cases} \\ \kappa(K_{-p/q}, \mathfrak{s}_1) &= \begin{cases} \kappa'_0 & (\text{if } -\frac{p}{8q} < c < 0), \\ \kappa'_0 \text{ or } \kappa'_0 + 2 & (\text{if } c < -\frac{p}{8q} < 0 \text{ or } c > 0). \end{cases}\end{aligned}$$

- (2) Suppose that $q > p > 0$, $p \equiv 0 \pmod{2}$ and $q \equiv 1 \pmod{2}$. Then there exist $\gamma_i \in \mathbf{Z}$ ($2 \leq i \leq n$) such that $q/p = [\gamma_2, \dots, \gamma_n]$, $|\gamma_i| \geq 2$, $\gamma_i \equiv 0 \pmod{2}$ (n is also odd). Let

$$\begin{aligned}\kappa_1 &= -\frac{1}{8}(1 - \sum_{i=2}^n \operatorname{sgn} \gamma_i + \sum_{i \text{ odd}} \gamma_i) - c, \\ \kappa'_1 &= \frac{1}{8}(1 - \sum_{i=2}^n \operatorname{sgn} \gamma_i + \sum_{i \text{ odd}} \gamma_i) - c.\end{aligned}$$

Then

$$\begin{aligned}\kappa(K_{p/q}, \mathfrak{s}_1) &= \kappa_1 \\ \kappa(K_{-p/q}, \mathfrak{s}_1) &= \kappa'_1 \text{ or } \kappa'_1 + 2\end{aligned}$$

- (3) Suppose that $p > 0$, $q > 0$, $p \equiv q \equiv 1 \pmod{2}$. Then there exist $\delta_i \in \mathbf{Z}$ ($1 \leq i \leq n$) such that $(p+q)/p = [\delta_1, \dots, \delta_n]$, $|\delta_i| \geq 2$, $\delta_i \equiv 0 \pmod{2}$. Let

$$\begin{aligned}\kappa_2 &= -\frac{1}{8}(1 - \sum_{i=1}^n \operatorname{sgn} \delta_i) - c, \\ \kappa'_2 &= \frac{1}{8}(1 - \sum_{i=1}^n \operatorname{sgn} \delta_i) - c.\end{aligned}$$

Then

$$\begin{aligned}\kappa(K_{p/q}, \mathfrak{s}_1) &= \begin{cases} \kappa_2 & (\text{if } c < 0 \text{ or } \frac{p}{8q} < c), \\ \kappa_2 \text{ or } \kappa_2 + 2 & (\text{if } 0 < c < \frac{p}{8q}), \end{cases} \\ \kappa(K_{-p/q}, \mathfrak{s}_1) &= \begin{cases} \kappa'_2 & (\text{if } -\frac{p}{8q} < c < 0), \\ \kappa'_2 \text{ or } \kappa'_2 + 2 & (\text{if } c > 0 \text{ or } c < -\frac{p}{8q}). \end{cases}\end{aligned}$$

Proof. Suppose that $p > q > 0$, $p \equiv 0 \pmod{2}$, and $q \equiv 1 \pmod{2}$. Let $\mathcal{L}_{(1)}$ be a framed link obtained from $\mathcal{L}$ in the proof of Proposition 3 by replacing L_4 with $\cup_{i=1}^n L'_i$, where

- $L'_1 = L_4$ and L'_i is a meridian of L'_{i-1} ($i \geq 2$),
- the framings of L'_i are $-8c + \beta_1$ ($i = 1$), and β_i ($i \geq 2$) respectively.

Then $K_{p/q}$ bounds a 4-manifold $X_{(1)}$ represented by $\mathcal{L}_{(1)}$ and the characteristic sublink of $\mathcal{L}_{(1)}$ associated with $\mathfrak{s}_1$ is $\cup_{i \text{ odd}} L'_i$. Furthermore $\cup_{i=1}^n L'_i$ represents a plumbing $X_{\Gamma_{(1)}}$ with $\partial X_{\Gamma_{(1)}} = L(p - 8cq, -q)$. The intersection matrix of $X_{(1)}$ is congruent to the diagonal matrix whose diagonal entries are

$$\{p/(4q), -4q/(p - 8cq), (p - 8cq)/q, [\beta_i, \dots, \beta_n] \ (i \geq 2)\}.$$

Furthermore the intersection matrices of the 4-manifold X_{L_1} corresponding to L_1 and $X_{\Gamma_{(1)}}$ are congruent to the diagonal matrices whose diagonal entries are $\{2c\}$ and $\{(p - 8cq)/q, [\beta_i, \dots, \beta_n] \ (i \geq 2)\}$ respectively. Then by pushing $X_{\Gamma_{(1)}}$ and X_{L_1} into $X_{(1)}$ and replacing them with the cones over $L(p - 8cq, -q)$ and $L(2c, -1)$ we obtain a spin orbifold $\widehat{X}_{(1)}$ bounded by $K_{p/q}$ whose spin structure $\widehat{\mathfrak{s}}$ induces $\mathfrak{s}_1$ on $K_{p/q}$. Let $\mathfrak{s}_{(1)}$ and $\mathfrak{s}_{L_1}$ be the spin structures on $L(p - 8cq, q)$ and $L(2c, -1)$ induced by $\widehat{\mathfrak{s}}$ respectively. It follows that $\sigma(\widehat{X}_{(1)}) = \sigma(X_{(1)}) - \sigma(X_{\Gamma_{(1)}}) - \sigma(X_{L_1}) = 1 - \text{sgn}(p - 8cq) - \text{sgn } c$ and hence

$$\begin{aligned} b_2^+(\widehat{X}_{(1)}) &= 1, \quad b_2^-(\widehat{X}_{(1)}) = 0 \text{ if } c < 0 \text{ or } p/q < 8c, \\ b_2^+(\widehat{X}_{(1)}) &= 0, \quad b_2^-(\widehat{X}_{(1)}) = 1 \text{ if } 0 < 8c < p/q. \end{aligned}$$

Furthermore we have

$$\bar{\mu}(L(p - 8cq, q), \mathfrak{s}_{(1)}) = \frac{1}{8}(\text{sgn}(p - 8cq) + \sum_{i=2}^n \text{sgn } \beta_i - \sum_{i \text{ odd}} \beta_i) + c,$$

and $\bar{\mu}(L(2c, -1)) = \frac{1}{8} \text{sgn } c$ since $\text{sgn}[\beta_i, \dots, \beta_n] = \text{sgn } \beta_i$. Let $\mathcal{L}'_{(1)}$ be a framed link obtained from $\mathcal{L}_{(1)}$ by replacing the framings of L'_i with $-8c - \beta_1$ ($i = 1$) and $-\beta_i$ ($i \geq 2$) respectively. Then $\mathcal{L}'_{(1)}$ represents a 4-manifold $X'_{(1)}$ bounded by $K_{-p/q}$ and the characteristic sublink of $\mathcal{L}'_{(1)}$ is also $\cup_{i \text{ odd}} L'_i$. In this case $\cup_{i=1}^n L'_i$ represents a plumbing $X_{\Gamma'_{(1)}}$ with $\partial X_{\Gamma'_{(1)}} = L(p + 8cq, q)$. We obtain a spin orbifold $\widehat{X}'_{(1)}$ with $\partial \widehat{X}'_{(1)} = K_{-p/q}$ by replacing $X_{\Gamma'_{(1)}}$ and X_{L_1} with the cones over their boundaries, whose spin structure $\widehat{\mathfrak{s}}'$ induces s_1 on $K_{-p/q}$. By computation similar to the above case, we have $\sigma(\widehat{X}'_{(1)}) = \text{sgn}(p + 8cq) - 1 - \text{sgn } c$ and hence

$$\begin{aligned} b_2^+(\widehat{X}'_{(1)}) &= 1, \quad b_2^-(\widehat{X}'_{(1)}) = 0 \text{ if } -p/q < 8c < 0, \\ b_2^-(\widehat{X}'_{(1)}) &= 0, \quad b_2^+(\widehat{X}'_{(1)}) = 1 \text{ if } c > 0 \text{ or } 8c < -p/q. \end{aligned}$$

Furthermore we have

$$\bar{\mu}(L(p + 8cq, q), \mathfrak{s}_{(1)'}) = \frac{1}{8}(-\operatorname{sgn}(p + 8cq) - \sum_{i=2}^n \operatorname{sgn} \beta_i + \sum_{i \text{ odd}} \beta_i) + c,$$

where $\mathfrak{s}_{(1)'}$ is the spin structure induced from $\widehat{\mathfrak{s}}'$.

Next suppose that $q > p > 0$, $p \equiv 0 \pmod{2}$ and $q \equiv 1 \pmod{2}$. Let $\mathcal{L}_{(2)}$ be a framed link obtained from $\mathcal{L}$ by replacing L_4 with $\cup_{i=1}^n L'_i$, where

- $L'_1 = L_4$, L'_i is a meridian of L'_{i-1} ($i \geq 2$),
- the framings of L'_i are $-8c$ ($i = 1$), $-\gamma_i$ ($i \geq 2$) respectively.

Then $\mathcal{L}_{(2)}$ represents a 4-manifold $X_{(2)}$ bounded by $K_{p/q}$ and the characteristic sublink of $\mathcal{L}_{(2)}$ associated with $\mathfrak{s}_1$ is $\cup_{i \text{ odd}} L'_i$. Furthermore $\cup_{i=1}^n L'_i$ represents a plumbing $X_{\Gamma_{(2)}}$ with $\partial X_{\Gamma_{(2)}} = L(p - 8cq, -q)$. We can also form a spin orbifold $\widehat{X}_{(2)}$ from $X_{(2)}$ by replacing $X_{\Gamma_{(2)}}$ and X_{L_1} with the cones over their boundaries, whose spin structure $\widehat{\mathfrak{s}}_{(2)}$ induces $\mathfrak{s}_1$ on $K_{p/q}$. The intersection matrix of $X_{(2)}$ is congruent to the diagonal matrix whose diagonal entries are

$$\{p/(4q), -4q/(p - 8cq), (p - 8cq)/q, -[\gamma_i, \dots, \gamma_n] \ (i \geq 2)\}$$

and we have $\sigma(\widehat{X}_{(2)}) = 1 - \operatorname{sgn}(p - 8cq) - \operatorname{sgn} c$ by computation similar to the above case. Hence we have $b_2^+(\widehat{X}_{(2)}) = 1$, $b_2^-(\widehat{X}_{(2)}) = 0$ since $0 < p/q < 1$ and

$$\bar{\mu}(L((p - 8cq, -q), \mathfrak{s}_{(2)})) = \frac{1}{8}(\operatorname{sgn}(p - 8cq) - \sum_{i=2}^n \operatorname{sgn} \gamma_i + \sum_{i \text{ odd}} \gamma_i) + c$$

and $\bar{\mu}(L(2c, -1), \mathfrak{s}_{L_1}) = \operatorname{sgn} c$, where $\mathfrak{s}_{(2)}$ and $\mathfrak{s}_{L_1}$ are the spin structures induced by $\widehat{\mathfrak{s}}_{(2)}$. On the other hand, $K_{-p/q}$ bounds a 4-manifold $X'_{(2)}$ represented by a framed link $\mathcal{L}'_{(2)}$, which is obtained from $\mathcal{L}_{(2)}$ by replacing the framings of L'_i with γ_i ($i \geq 2$). The characteristic sublink of $\mathcal{L}'_{(2)}$ associated with $\mathfrak{s}_1$ is also $\cup_{i \text{ odd}} L'_i$ and $\cup_{i=1}^n L'_i$ forms a plumbing $X_{\Gamma'_{(2)}}$ with $\partial X_{\Gamma'_{(2)}} = L(p + 8cq, q)$. We also form a spin 4-orbifold $\widehat{X}'_{(2)}$ from $X'_{(2)}$ by replacing $X_{\Gamma'_{(2)}}$ and X_{L_1} with the cones over their boundaries, whose spin structure $\widehat{\mathfrak{s}}_{(2)'}$ induces $\mathfrak{s}_1$ on $K_{-p/q}$. Then computation similar to the above case shows that $\sigma(\widehat{X}'_{(2)}) = \operatorname{sgn}(p + 8cq) - 1 - \operatorname{sgn} c$, and $b_2^+(\widehat{X}'_{(2)}) = 0$, $b_2^-(\widehat{X}'_{(2)}) = 1$. Furthermore we have

$$\bar{\mu}(L(p + 8cq, q), \mathfrak{s}'_{(2)}) = \frac{1}{8}(-\operatorname{sgn}(p + 8cq) + \sum_{i=2}^n \operatorname{sgn} \gamma_i - \sum_{i \text{ odd}} \gamma_i) + c$$

where $\mathfrak{s}'_{(2)}$ is induced from $\widehat{\mathfrak{s}}_{(2)}'$.

Finally suppose that $p > 0$, $q > 0$, and $p \equiv q \equiv 1 \pmod{2}$. In this case we consider a framed link $\mathcal{L}_{(2)}$ obtained from $\mathcal{L}$ by replacing L_4 with $\cup_{i=1}^{n+2} L'_i$, where

- $L'_1 = L_4$, L'_i is a meridian of L'_{i-1} ($i \geq 2$),

- the framings of L'_i are $-8c-1$ ($i=1$), -1 ($i=2$), $-\delta_{i-2}$ ($i \geq 3$) respectively.

Then $\mathcal{L}_{(2)}$ represents a 4-manifold $X_{(2)}$ bounded by $K_{p/q}$, and $\cup_{i=1}^{n+2} L'_i$ represents a plumbing $X_{\Gamma_{(2)}}$ bounded by $L(p-8cq, -q)$. The characteristic sublink of $\mathcal{L}_{(2)}$ associated with $\mathfrak{s}_1$ is L'_1 . The intersection matrices of $X_{(2)}$ and $X_{\Gamma_{(2)}}$ are congruent to the diagonal matrices whose diagonal entries are

$$\begin{aligned} &\{p/(4q), -4q/(p-8cq), (p-8cq)/q, -q/(p+q), -[\delta_i, \dots, \delta_n] \ (i \geq 1)\} \\ &\{(p-8cq)/q, -q/(p+q), -[\delta_i, \dots, \delta_n] \ (i \geq 1)\}. \end{aligned}$$

It follows that if we form a spin 4-orbifold $\widehat{X}_{(2)}$ from $X_{(2)}$ by replacing $X_{\Gamma_{(2)}}$ and X_{L_1} with the cones over their boundaries, whose spin structure $\widehat{\mathfrak{s}}_{(2)}$ induces $\mathfrak{s}_1$ on $K_{p/q}$, we have $\sigma(\widehat{X}_{(2)}) = 1 - \text{sgn}(p-8cq) - \text{sgn } c$ and hence

$$\begin{aligned} b_2^+(\widehat{X}_{(2)}) &= 1, \quad b_2^-(\widehat{X}_{(2)}) = 0 \text{ if } c < 0 \text{ or } 8c > p/q, \\ b_2^+(\widehat{X}_{(2)}) &= 0, \quad b_2^-(\widehat{X}_{(2)}) = 1 \text{ if } 0 < 8c < p/q. \end{aligned}$$

Furthermore we have

$$\bar{\mu}(L(p-8cq, -q), \mathfrak{s}_{(2)}) = \frac{1}{8}(\text{sgn}(p-8cq) - \sum_{i=1} \text{sgn } \delta_i) + c,$$

where $\mathfrak{s}_{(2)}$ is induced by $\widehat{\mathfrak{s}}_{(2)}$. On the other hand $K_{-p/q}$ bounds a 4-manifold $X'_{(2)}$ represented by a framed link $\mathcal{L}'_{(2)}$, which is obtained from $\mathcal{L}_{(2)}$ by replacing the framings of L'_i with $-8c+1$ ($i=1$), $+1$ ($i=2$), δ_{i-2} ($i \geq 3$) respectively. In this case $\cup_{i=1}^{n+2} L'_i$ forms a plumbing $X_{\Gamma'_{(2)}}$ bounded by $L(p+8cq, q)$, and the characteristic sublink of $\mathcal{L}'_{(2)}$ is also L'_1 . If we form a spin orbifold $\widehat{X}'_{(2)}$ from $X'_{(2)}$ by replacing $X_{\Gamma'_{(2)}}$ and X_{L_1} with the cones over their boundaries, whose spin structure $\widehat{\mathfrak{s}}'_{(2)}$ induces $\mathfrak{s}_1$ on $K_{-p/q}$ as before, computation similar to the above case shows that $\sigma(\widehat{X}'_{(2)}) = \text{sgn}(p+8cq) - 1 - \text{sgn } c$,

$$\begin{aligned} b_2^+(\widehat{X}'_{(2)}) &= 1, \quad b_2^-(\widehat{X}'_{(2)}) = 0 \text{ if } -p/q < 8c < 0, \\ b_2^+(\widehat{X}'_{(2)}) &= 0, \quad b_2^-(\widehat{X}'_{(2)}) = 1 \text{ if } c > 0 \text{ or } 8c < -p/q, \end{aligned}$$

and also

$$\bar{\mu}(L(p+8cq, q), \mathfrak{s}'_{(2)}) = \frac{1}{8}(-\text{sgn}(p+8cq) + \sum_{i=1}^n \text{sgn } \delta_i) + c,$$

where $\mathfrak{s}'_{(2)}$ is induced from $\widehat{\mathfrak{s}}_{(2)}'$.

We obtain the required results by applying Corollary 1 to the above computation. $\square$

REFERENCES

- [1] I. Dai, On the $\text{Pin}(2)$ -equivariant monopole Floer homology of plumbed 3-manifolds, *Mich. Math. J.* **67** (2018), 423–447.
- [2] I. Dai, C. Manolescu, Involutive Heegaard Floer homology and plumbed three-manifolds, *J. Inst. Math. Jussieu* **18** (2019), 1115–1155.
- [3] I. Dai, H. Sasahira, M. Stoffregen, Lattice homology and Seiberg-Witten-Floer spectra, *arXiv:2309.01253*, (2023)
- [4] Y. Fukumoto, M. Furuta, Homology 3-spheres bounding acyclic 4-manifolds, *Math. Res. Lett.* **7** (2000), 757–766
- [5] Y. Fukumoto, M. Furuta, M. Ue, W invariants and the Neumann-Siebenmann invariants for Seifert homology 3-spheres, *Topol. and its appl.* **116** (2001), 333–369
- [6] K. A. Frøyshov, Monopole Floer homology for rational homology 3-manifolds, *Duke Math. J.* **155** (2010), 519–576
- [7] K. Hendricks, C. Manolescu, Involutive Heegaard Floer Homology, *Duke Math. J.* **166** (7) (2017), 1211–1299.
- [8] P. Kronheimer, T. Mrowka, Monopoles and Three-Manifolds, *New Mathematical Monograph* **10** Cambridge Univ. Press, (2007)
- [9] T. Mrowka, P. Ozsváth, and B. Yu, Seiberg-Witten monopoles on Seifert fibered spaces, *Comm. Anal. Geom.*, **5** (1997), No. 4, 685–791
- [10] C. Manolescu, Seiberg-Witten Floer stable homotopy type of three manifolds with $b_1 = 0$, *Geom. Topol.* **7** (2003), 889–932.
- [11] C. Manolescu, $\text{Pin}(2)$ -equivariant Seiberg-Witten Floer homology and the triangulation conjecture, *J. Amer. Math. Soc.* **29** (2016), 147–176.
- [12] C. Manolescu, On the intersection forms of spin 4-manifolds with boundary, *Math. Ann.* **359** (2014), 695–728.
- [13] W. D. Neumann, An invariant of plumbed homology spheres, in *Topology Symposium, Siegen 1979* (Proc. Sympos., Univ. Siegen), Springer, Berlin Vol. **788** *Lectures Notes in Math.*, pp.125–144, (1980)
- [14] A. Némethi, On the Ozsváth-Szabó invariant of negative definite plumbed 3-manifolds, *Geom. Topol.* **9** (2005), 991–1042
- [15] P. Ozsváth, Z. Szabó, Absolutely graded Floer homologies and intersection forms for four manifolds, *Adv. in Math.*, **173** (2003), 179–261
- [16] N. Saveliev, Fukumoto-Furuta invariants of plumbed homology 3-spheres, *Pacific J. Math.* **205** (2002), 465–490
- [17] L. Siebenmann, On vanishing of the Rohlin invariant and nonfinitely amphicheiral homology 3-spheres, in *Topology Symposium, Siegen 1979* (proc. Sympos., Univ. Siegen, 1979) , Springer, Berlin, Vol. **788** *lecture Notes in Math.*, pp. 172–222, (1980)
- [18] M. Stoffregen, $\text{Pin}(2)$ -equivariant Seiberg-Witten Floer homology of Seifert fibrations, *Compos. Math.* **156** (2) (2020), 199–250.
- [19] M. Ue, The Neumann-Siebenmann invariant and Seifert surgery, *Math.Z.* **250** (2005), 475–493
- [20] M. Ue, Constraints on intersection forms of spin 4-manifolds bounded by Seifert rational homology 3-spheres in terms of $\bar{\mu}$ and κ invariants, *arXiv:2206.05412*, (2022)

KYOTO UNIVERSITY, KYOTO, 606-8502, JAPAN

Email address: ue.masaaki.87m@st.kyoto-u.ac.jp