1. INTRODUCTION

ON FINDING BIFURCATIONS FOR NON-VARIATIONAL ELLIPTIC SYSTEMS BY THE EXTENDED QUOTIENTS

METHOD

YAVDAT IL'YASOV

ABSTRACT. We develop a novel method for finding bifurcations for nonlinear systems of equations based on directly finding bifurcations through saddle points of extended quotients. The method is applied to find the saddle-node bifurcation point for elliptic equations with the nonlinearity of the general convex-concave type. The main result justifies the variational formula for the detection of the maximum saddle-node type bifurcation point of stable positive solutions. As a consequence, a precise threshold value separating the interval

This paper develops a method of detecting bifurcation introduced in [20, 25], which provides a direct way of finding bifurcations by identifying saddle points of the corresponding extended Rayleigh quotient. We develop the method by finding saddle-node bifurcation point for the following system of equations:

(1.1)
$$\begin{cases} -\Delta u_i = a_i(x)u_i^q + \lambda g_i(x, u), & x \in \Omega, \\ u_i \ge 0, & x \in \Omega, \\ u_i|_{\partial\Omega} = 0, & i = 1, \dots, m. \end{cases}$$

of the existence of stable positive solutions is established.

Here $q_i \in (0,1), i = 1, \ldots, m, \Omega$ is a bounded connected domain in \mathbb{R}^d with $\partial \Omega \in C^2, d \geq 1, \lambda \in \mathbb{R}, u := (u_1, \ldots, u_m)$. For $i = 1, \ldots, m, a_i \in L^{\infty}(\Omega), a_i > 0$ in $\Omega, a_i(\cdot)$ is a Hölder continuous function in $\Omega, g_i(\cdot, u)$ is measurable function in $\Omega, \forall u \in \mathbb{R}^m$, and $g_i(x, \cdot) \in C^1(\mathbb{R}^m, \mathbb{R})$. Furthermore,

 (g_1) : $\exists c_0, c_1 > 0$ and $\exists \gamma \in (1, +\infty)$ such that

$$0 \le g_i(x, u) \le c_0 |u| + c_2 |u|^{\gamma}, \ x \in \Omega, \ u \in \mathbb{R}^m_+, \ i = 1, ..., m;$$

 (g_2) : $\exists c_2, c_3 > 0$ and $\exists \gamma_0 \in (1, \gamma)$ such that

$$(g_{i,u_i}(x,u)u_i^2 - g_i(x,u)u_i) \ge c_2 |u|^{\gamma_0+1} + c_3 |u|^{\gamma+1}, \ x \in \Omega, \ u \in \mathbb{R}^m_+.$$

Throughout this paper the summation convention is in place: we sum over any index that appears twice. A particular example of functions g_i , i = 1, ..., m that meets condition $(g_1) - (g_2)$ is as follows: $g_i = b_i(x)u_i + b(x)\sum_{j=1}^m |u_j|^{\gamma-1}u_j$, i = 1, ..., m with $\gamma_0 = \gamma > m$ and $b, b_i \in L^{\infty}(\Omega)$.

Hereafter, we denote $\mathcal{W} := (\check{W}_2^1(\Omega) \cap L^{\gamma})^m$, $F_i(u,\lambda) := -\Delta u_i - a_i |u_i|^{q-1} u_i - \lambda g_i(x,u)$, $i = 1, \ldots m$, $F(u,\lambda) = (F_1(u,\lambda), \ldots, F_m(u,\lambda))^T$, $u \in \mathcal{W}$, $\lambda \in \mathbb{R}$. $(\mathcal{W})^*$

²⁰²⁰ Mathematics Subject Classification. 35J60; 35J96; 35R35; 53C45.

means the dual space of \mathcal{W} . A point $(u, \lambda) \in \mathcal{W} \times \mathbb{R}$ is called a weak solution of (1.1) if $F(u, \lambda) = 0$ holds true in $(\mathcal{W})^*$.

Define $S := \{u \in C^1(\overline{\Omega}) \mid \exists c_u > 0, u > c_u \operatorname{dist}(x, \partial\Omega) \text{ in } \Omega, u|_{\partial\Omega} = 0\}.$ We show below that the map $F(\cdot, \lambda) : \mathcal{W} \to \mathcal{W}^*$ is Fréchet differentiable on S^m , $\forall \lambda \in \mathbb{R}$ (see Proposition 2.1 below). A solution $u_{\lambda} \in S^m$ of (1.1) is said to be *stable* if $\lambda_1(F_u(u_{\lambda}, \lambda)) \geq 0$, and *asymptotically stable* if $\lambda_1(F_u(u_{\lambda}, \lambda)) > 0$, cf. [8, 11]. Hereafter, $\lambda_1(F_u(u, \lambda))$, for $u \in S^m$ denotes the first eigenvalue of the operator $F_u(u, \lambda)(\cdot)$.

Introduce

$$\mathcal{W}_s := \{ u \in \mathcal{W} \cap S^m : \lambda_1(F_u(u,\tau)) \ge 0, \ \tau = \mathcal{R}(u,u) \},\$$

We call a solution $(\hat{u}, \hat{\lambda}) \in \mathcal{W}_s \times \mathbb{R}$ of (1.1) the saddle-node bifurcation point in \mathcal{W}_s (or, equivalently, fold, turning point) (cf. [26, 27]) if the following is fulfilled: (i) the nullspace $N(F_u(\hat{u}, \hat{\lambda}))$ of the Fréchet derivative $F_u(\hat{u}, \hat{\lambda})$ is not empty; (ii) there exists $\varepsilon > 0$ and a neighborhood $U_1 \subset \mathcal{W}$ of \hat{u} such that for any $\lambda \in (\hat{\lambda}, \hat{\lambda} + \varepsilon)$ equation (1.1) has no solutions in $\mathcal{W}_s \cap U$; (iii) for each $\lambda \in (\hat{\lambda} - \varepsilon, \hat{\lambda})$, the equation has precisely two distinct solutions in $\mathcal{W}_s \cap U$. This definition corresponds to the solution's curve turning back at the bifurcation value $\hat{\lambda}$. The solution's curve turning forward is defined similar. In the case only (i)-(ii) are satisfied, we call $(\hat{u}, \hat{\lambda})$ the saddle-node type bifurcation point of (1.1) in \mathcal{W}_s . A saddle-node type bifurcation $(\hat{u}, \hat{\lambda})$ of (1.1) in \mathcal{W}_s .

A model example for (1.1) in the scalar case is the Ambrosetti–Brezis–Cerami problem [2] with concave–convex nonlinearity

(1.2)
$$-\Delta u = |u|^{q-1}u + \lambda |u|^{\gamma-1}u, \quad u|_{\partial\Omega} = 0,$$

where $0 < q < 1 < \gamma$. It is why the nonlinearity in (1.1) can be considered to be of the convex-concave type. From [2] it follows that there exists an extremal value $\lambda^* > 0$ such that for any $\lambda \in (0, \lambda^*]$, (1.2) has a stable positive solution u_{λ} , while for $\lambda > \lambda^*$, (1.2) does not admit weak positive solutions. According to [2], the solution u_{λ} for $\lambda \in (0, \lambda^*)$ is obtained by super-subsolution methods, while u_{λ^*} is shown to exist as a limit point of (u_{λ}) . Unfortunately, this method is not easily adaptable to systems of equations like (1.1). Indeed, the super-sub solution method for a system of equations differs considerably from that which is used for a scalar equation.

In general cases, system (1.1) is not a variational or Hamiltonian. It should be noted that in contrast to the extensive literature concerning the existence of solutions for variational and Hamiltonian systems (see the survey [15]), relatively little research is devoted to nonvariational and non-Hamiltonian systems of equations (see, e.g., [1, 5, 7, 10, 34, 35] and references therein).

The finding of bifurcations of solutions to equations poses a more complex challenge, requiring a comprehensive approach that considers both the finding solutions themselves and the analysis of the structure of the family of solutions. This problem is still quite challenging even when dealing with scalar equations. The complete answer to the question on the existence of the saddle-node bifurcation point and the exact shape of the positive solution curves for instance of the scalar equation (1.2) was obtained only in radially symmetric solutions [28, 30, 33]. An additional obstacle encountered when studying equations (1.1) and (1.2) is the presence of singular derivatives of the right-hand sides. Specifically, standard methods (see [8,9,26,27]) are not readily applicable for verifying that the solution's curve turning back at $(\hat{u}, \hat{\lambda})$, due to the difficulties in testing conditions for the second derivative of $F(u, \lambda)$.

The existence of positive solutions and multiplicity results were studied only in some special cases of system (1.1) in the variational form (see, e.g., [18, 19, 25, 35] and references therein). A recent study [25] answered the question of whether positive solutions of system (1.1) in the variational form have a saddle-node type bifurcation point. However, in the general cases of system (1.1), to the best of my knowledge, no studies have been conducted on the existence of non-negative solutions and saddle-node bifurcation points.

Let us state our main results. Observe that by the definition the saddle-node type bifurcation point $(u, \lambda) \in \mathcal{W}_s \times \mathbb{R}$ of (1.1) should satisfy the system of equations

(1.3)
$$\begin{cases} F(u,\lambda) = 0, \\ F_u(u,\lambda)(v) = 0, \end{cases}$$

with some $v \in N(F_u(u, v))$. To analyze this system, following [25] we introduce the extended Rayleigh quotient (extended quotient for short) associated with (1.1)

$$\mathcal{R}(u,v) := \frac{\int (\nabla u_i, \nabla v_i) - \int a_i u_i^q v_i}{\int g_i(x, u) v_i}, \quad u \in \mathcal{W}_s, v \in \Sigma(u).$$

Here $\Sigma(u) := \{ v \in \mathcal{W} : \int g_i(x, u) v_i \neq 0 \}$ for $u \in \mathcal{W}_s$. Observe,

$$\begin{cases} \lambda = \mathcal{R}(u, v), \\ \mathcal{R}_v(u, v) = 0, \\ \mathcal{R}_u(u, v) = 0 \end{cases} \Leftrightarrow \begin{cases} F(u, \lambda) = 0, \\ F_u(u, \lambda)(v) = 0, \end{cases}$$

that is, the set of saddle-node type bifurcation points of (1.1) contains in the set of critical points of $\mathcal{R}(u, v)$ on $\mathcal{W}_s \times \mathcal{W}$.

In our approach, the following minimax formula plays a major role (cf. [25])

(1.4)
$$\lambda_s^* := \sup_{u \in \mathcal{W}_s} \inf_{v \in \Sigma(u)} \mathcal{R}(u, v)$$

The main result of the work is as follows

Theorem 1.1. Assume $(g_1) - (g_2)$, $q_i \in (0, 1)$, i = 1, ..., m.

- (1°) Then $0 \leq \lambda_s^* < +\infty$.
 - (a): For $\lambda = \lambda_s^*$, there exists a weak positive solution $u_s^* \in (C^{1,\alpha}(\overline{\Omega}))^m \cap \mathcal{W}$ of system (1.1).
 - (b): For any $\lambda > \lambda_s^*$, system (1.1) has no stable weak positive solutions.

(2°) Assume in addition that $\gamma < 2^*$, and $q_i < (2^* - 2)/2 \equiv 2/(d - 2)$ if d > 2. Then $0 < \lambda_s^* < +\infty$, (u_s^*, λ_s^*) is a maximal saddle-node type bifurcation point of (1.1) in \mathcal{W}_s . Moreover, u_s^* is a stable solution of (1.1).

Here $2^* = 2d/(d-2)$ if $d \ge 3$, and $2^* = +\infty$ if d = 1, 2.

Remark 1.1. Statement (1°) can be supplemented as follows. There exists $\lambda \in [0, \lambda_s^*]$ such that (1.1) has a stable positive weak solution $u_{\lambda} \in (C^{1,\alpha}(\overline{\Omega}))^m \cap \mathcal{W}$. Indeed, we will see below that (1.1) has a stable positive weak solution at least for $\lambda = 0$. The following can also be considered in conjunction with the value (1.4)

(1.5)
$$\lambda_{as}^* := \sup_{u \in \mathcal{W}_{as}} \inf_{v \in \Sigma(u)} \mathcal{R}(u, v)$$

4

Here $\mathcal{W}_{as} := \{u \in \mathcal{W} \cap S^m : \lambda_1(F_u(u,\tau)) > 0, \tau = \mathcal{R}(u,u)\}$. It easily see that $\lambda_{as}^* \leq \lambda_s^*$. For (1.5), it can be obtained a result similar to Theorem 1.1. In particular, we have the following

Theorem 1.2. Assume $(g_1) - (g_2)$, $q_i \in (0, 1)$, i = 1, ..., m, and $q_i < (2^* - 2)/2 \equiv 2/(d-2)$ if d > 2. Then $0 < \lambda_{as}^* < +\infty$, and

- (a): For $\lambda = \lambda_{as}^*$, system (1.1) has a stable weak positive solution $u_{as}^* \in (C^{1,\alpha}(\overline{\Omega}))^m \cap \mathcal{W}$. Furthermore, $(u_{as}^*, \lambda_{as}^*)$ is a maximal saddle-node type bifurcation point of (1.1) in \mathcal{W}_{as} .
- (b) : For any $\lambda > \lambda_{as}^*$, (1.1) has no asymptotically stable weak positive solutions.
- (c): There exists a sequence of asymptotically stable weak positive solutions $u_{\lambda_n} \in (C^{1,\alpha}(\overline{\Omega}))^m \cap \mathcal{W}$ of (1.1) with $\lambda = \lambda_n > 0$, $n = 1, \ldots$ such that $u_{\lambda_n} \to u_{as}^*$ in \mathcal{W} and $\lambda_n \to \lambda_{as}^*$ as $n \to +\infty$.

Remark 1.2. It is natural to expect that $u_s^* = u_{as}^*$, $\lambda_s^* = \lambda_{as}^*$ and (u_s^*, λ_s^*) is indeed a saddle-node bifurcation point of (1.1) in W_s . It should be noted that assertions (1°), (b) and (2) of Theorems 1.1 and 1.2 do not necessarily mean that (1.1) has no any positive solutions for $\lambda > \lambda_s^*$. Furthermore, such a behavior is possible if (1.1) has an S-shaped bifurcation curve (see [4, 6, 17]).

Remark 1.3. We believe that the variational formula (1.4) has the potential provide a useful tool in further analyzing saddle-node bifurcation points and constructing numerical methods for finding them (cf. [21-24, 31]).

The rest of the paper is organised as follows. Section 2 presents preliminaries. In Section 3, we prove Theorems 1.1, 1.2. In Appendix, we present a proof of a version of Ekeland's principal for smooth functional.

2. Preliminaries

We use the standard notation $L^p := L^p(\Omega)$ for the Lebesgue spaces, $1 \le p \le +\infty$, and denote by $\|\cdot\|_p$ the associated norm. By $\mathring{W}_2^1 := \mathring{W}_2^1(\Omega)$ we denote the standard Sobolev space, endowed with the norm $\|u\|_{1,2} = (\int |\nabla u|^2)^{1/2}$. Hereafter, we denote $W := \mathring{W}_2^1(\Omega) \cap L^\gamma, d(x) := \operatorname{dist}(x, \partial \Omega).$

For $\delta > 0$, define $S(\delta) := \{ u \in S \mid u(x) > \delta d(x) \text{ in } \Omega \}$, S(0) := S. Clearly, $S(\delta), \forall \delta \ge 0$ is an open subset in $C^1 := C^1(\overline{\Omega})$. Let Y be a topological space such that $S^m(\delta) \subset Y$. The set $S^m(\delta)$ endowed with topology of Y we denote by $S_Y^m(\delta)$. $\mathcal{L}(\mathcal{W}, \mathcal{W}^*)$ denotes the Banach space of bounded linear operators from \mathcal{W} into \mathcal{W}^* .

Proposition 2.1. Let $\lambda \in \mathbb{R}$, $0 < q_i < 1$, i = 1, ..., m. Then $F_i(u, \lambda) : \mathcal{W} \to \mathcal{W}^*$ is Fréchet differentiable at any $u \in S^m$, and $F_i(\cdot, \lambda) \in C^1(S^m_{C^1}, \mathcal{L}(\mathcal{W}, \mathcal{W}^*))$. Furthermore, if in addition $q_i \leq (\bar{\gamma} - 2)/2$, where $\bar{\gamma} = \max\{\gamma, 2^*\}$, then $F_{i,u}(\cdot, \lambda) \in C(S_{\mathcal{W}}(\delta), \mathcal{L}(\mathcal{W}, \mathcal{W}^*))$, $\forall \delta > 0$.

Proof. We develop an approach proposed in [2]. We verify the assertion only for the map $Q(u) := a_i u^q$, $u \in S$ since for the remaining terms in F_i the statement is trivial. Using the inequality $u(x) \ge c(u)d(x)$ in Ω and Hölder's inequalities we derive

$$\begin{split} |\int a_{i}u^{q-1}\phi\psi\,dx| = & |\int a_{i}u^{q}\left(\frac{\phi}{u}\right)\psi\,dx| \leq \\ & \frac{\|a_{i}\|_{\infty}}{c(u)}\|u\|_{p}^{q}\cdot\|\frac{\phi}{d(\cdot)}\|_{2}\cdot\|\psi\|_{\gamma}, \ \ u\in S, \ \forall\phi,\psi\in C_{0}^{\infty}(\Omega), \end{split}$$

where $p = 2q\gamma/(\gamma - 2)$. By the Hardy inequality, $\|\phi/d(\cdot)\|_2 \leq C \|\phi\|_{1,2}, \forall \phi \in C_0^{\infty}(\Omega)$, and thus, using the Sobolev inequalities we derive

(2.1)
$$|\int a_i u^{q-1} \phi \psi \, dx| \le \frac{C ||a_i||_{\infty}}{c(u)} ||u||_p^q ||\phi||_{1,2} ||\psi||_{1,2}, \ u \in S, \ \forall \phi, \psi \in C_0^{\infty}(\Omega),$$

where $C \in (0, +\infty)$ does not depend on u, ϕ, ψ . This implies that $Q(\cdot) : W \to W^*$ is Fréchet differentiable at any $u \in S$. In the same manner we can see that $Q_u(\cdot) \in C(S_{C^1}, \mathcal{L}(S_{C^1}, W^*))$, and thus, $F_i(\cdot, \lambda) \in C^1(S_{C^1}^m; \mathcal{L}(W, W^*))$.

Let us prove the second part. For simplicity we assume that $\bar{\gamma} = \gamma$. Suppose $u_k \to u$ in L^{γ} as $k \to +\infty$. This means that there exist $\bar{u} \in L^{\gamma}$ and a subsequence (which is denoted again by (u_k)) such that $|u_k|, |u| \leq \bar{u}$ in Ω . Hence, $u_k^{q-1}d(x) \leq \bar{u}^q$ in Ω , $k = 1, \ldots$, and therefore, Lebesgue's dominated convergence theorem yields $u_k^{q-1}d(x) \to u^{q-1}d(x)$ in $L^{\gamma/q}$ as $k \to +\infty$. Notice that $q \leq (\gamma - 2)/2$ implies $p \leq \gamma$. Similar to (2.1) we have

$$\left|\int a_{i}(u_{k}^{q-1}-u^{q-1})\phi\psi\,dx\right| \leq \frac{C}{c(u)}\|(u_{k}^{q-1}-u^{q-1})d(\cdot)\|_{p}^{q}\|\phi\|_{1,2}\|\psi\|_{1,2}, \quad \forall \phi, \psi \in W,$$

for some $C < +\infty$ which does not depend on $u, u_k \in S$, $\phi, \psi \in (W)^*$. Hence, using the Sobolev inequalities we derive $Q_u(\cdot) \in C(S_W(\delta), \mathcal{L}(W, W^*))$, and thus, $F_{i,u}(\cdot, \lambda) \in C(S_W(\delta), \mathcal{L}(W, W^*)), \forall \delta > 0.$

Proposition 2.2. If $u \in W$ is a weak non-negative solution to (1.1), then $u_i \in C^{1,\alpha}(\overline{\Omega})$ for any $\alpha \in (0,1)$, and $u_i > 0$ in Ω , $i = 1, \ldots, m$.

Proof. Note that equality (1.1) implies that $-\Delta u_i \geq 0$, $i = 1, \ldots, m$, and thus, by the maximum principals for the elliptic problems, $u_i > 0$ in Ω , $i = 1, \ldots, m$. The standard bootstrap argument and Sobolev's embedding theorem entail that $u_i \in L^{\infty}(\Omega), i = 1, \ldots, m$. This by the regularity results for elliptic problems from [29] implies that $u_i \in C^{1,\alpha}(\overline{\Omega})$ for any $\alpha \in (0, 1), i = 1, \ldots, m$.

Let i = 1, ..., m. By Brezis-Oswald's result [3] there exists a unique solution $w_i \in \mathring{W}_2^1(\Omega) \cap C^{1,\alpha}(\overline{\Omega}), \alpha \in (0,1)$ of

(2.3)
$$\begin{cases} -\Delta w = a_i w^q \text{ in } \Omega \\ w|_{\partial\Omega} = 0. \end{cases}$$

By the assumption $a_i(\cdot)$ is a Hölder continuous function in Ω , and hence, by the Schauder estimates (see, e.g., [16]), $w_i \in C^2(\Omega)$. Furthermore, the strong maximum principals for the elliptic problems imply that $\min_{x'\in\partial\Omega} \frac{\partial w_i(x')}{\partial \nu(x')} > 0$, where $\nu(x')$ denotes the interior unit normal at $x' \in \partial\Omega$, see, e.g., Lemma 3.4 in [16]. Thus, $w_i \in S$.

Moreover, $\overline{w} := (w_1, \ldots, w_m)$ is a stable solution of (2.3) with $\lambda = 0$, i.e., $\overline{w} \in \mathcal{W}_s$. Indeed, from Proposition 2.1 it follows that $F(\overline{w}, 0) \in C^1(S_{C^1}^m; \mathcal{L}(\mathcal{W}, \mathcal{W}^*))$, and therefore, $\lambda_1(F_u(\overline{w}, 0))$ is well defined. It is not hard to show that \overline{w} is

a minimizer of $E(v) := \left(\frac{1}{2}\sum_{i=1}^{m}\int |\nabla v_i|^2 - \frac{1}{q+1}\int a_i|v_i|^{q+1}\right)$ on $(\mathring{W}_2^1)^m$, that is, $E(\overline{w}) = \inf_{v \in (\mathring{W}_2^1)^m} E(v)$. In particular, this means

(2.4)
$$\lambda_1(F_u(\overline{w},0)) \ge 0.$$

Lemma 2.1. Suppose $u^0 \in \mathcal{W}_s$ such that

$$-\infty < \lambda_0 := \inf_{v \in \Sigma(u^0)} \mathcal{R}(u^0, v) < +\infty.$$

Then u^0 is a weak solution of (1.1) for $\lambda = \lambda_0$.

Proof. Let $v^k \in \Sigma(u^0)$, $k = 1, \ldots$, such that

$$\lambda_k \equiv \mathcal{R}(u^0, v^k) \to \inf_{v \in \Sigma(u^0)} \mathcal{R}(u^0, v) \equiv \lambda_0 \text{ as } k \to +\infty.$$

Since $\mathcal{R}(u, v) = \mathcal{R}(u, sv), \forall s \in \mathbb{R} \setminus 0, \forall v \in \Sigma(u), \forall u \in \mathcal{W}_s$, we may assume that

(2.5)
$$\int g_i(x, u^0) v_i^k = 1, \quad k = 1, \dots$$

Calculate

$$\begin{split} \mathcal{R}_{v}(u^{0},v^{k})(\xi) &= \frac{\int (\nabla u_{i}^{0},\nabla\xi_{i}) - \int a_{i}(u_{i}^{0})^{q}\xi_{i} - \mathcal{R}(u^{0},v^{k}) \int g_{i}(x,u^{0})\xi_{i}}{\int g_{i}(x,u^{0})v_{i}^{k}},\\ \mathcal{R}_{vv}(u^{0},v^{k})(\xi,\zeta) &= \\ &- \frac{\left(\int (\nabla u_{i}^{0},\nabla\xi_{i}) - \int a_{i}(u_{i}^{0})^{q}\xi_{i} - \lambda_{k} \int g_{i}(x,u^{0})\xi_{i}\right) \cdot \int g_{i}(x,u^{0})\zeta_{i}}{(\int g_{i}(x,u^{0})v_{i}^{k})^{2}} - \\ &\frac{\left(\int (\nabla u_{i}^{0},\nabla\zeta_{i}) - \int a_{i}(u_{i}^{0})^{q}\zeta_{i} - \lambda_{k} \int g_{i}(x,u^{0})\zeta_{i}\right) \cdot \int g_{i}(x,u^{0})\xi_{i}}{(\int g_{i}(x,u^{0})v_{i}^{k})^{2}}, \ \zeta, \xi \in \mathcal{W}. \end{split}$$

Let $\phi \in \mathcal{W}$, $\|\phi\|_{\mathcal{W}} = 1$. Using (2.5) and the Hölder and Sobolev inequalities one can see that

(2.6)
$$\left| \int (g_i(x, u^0)(v_i^k + \tau \phi_i)) \right| = |1 + \tau \int g_i(x, u^0) \phi_i| \ge 1 - a_0 |\tau|$$

where $a_0 \in (0, \infty)$ does not depend on ϕ and $k = 1, \ldots$. Hence $v^k + \tau \phi \in \Sigma(u^0)$ for any $k = 1, \ldots$ and τ such that $|\tau| < \tau_0 := 1/a_0$.

By (2.6), (g_1) we have

$$\begin{aligned} \|\mathcal{R}_{vv}(u^{0}, v^{k} + \tau\phi)\|_{(\mathcal{W}\times\mathcal{W})^{*}} &= \frac{2}{|\int g_{i}(x, u^{0})(v_{i}^{k} + \tau\phi_{i})|^{2}} \times \\ \sup_{\xi,\zeta\in\mathcal{W}} \frac{|\left(\int (\nabla u_{i}^{0}, \nabla\xi_{i}) - \int a_{i}(u_{i}^{0})^{q}\xi_{i} - \lambda_{k}\int g_{i}(x, u^{0})\xi_{i}\right) \cdot \int a_{i}(u_{i}^{0})^{q}\zeta_{i}|}{\|\xi_{i}\|_{\mathcal{W}}} \leq \\ \frac{2}{(1 - a_{0}|\tau|)^{2}} \left(\sum_{i=1}^{m} \|-\Delta u_{i}^{0} - a_{i}(u_{i}^{0})^{q} - \lambda_{k}g_{i}(x, u^{0})\|_{\mathcal{W}^{*}}\right) \|u^{0}\|_{\mathcal{W}^{*}}}{\leq \frac{C_{0}}{(1 - a_{0}|\tau|)^{2}}} \end{aligned}$$

where $C_0 \in (0, \infty)$ does not depend on $k = 1, \ldots$. We thus may apply Theorem 4.1 to the functional $G(v) := \mathcal{R}(u^0, v)$ defined in the open domain $V := \Sigma(u^0) \subset \mathcal{W}$.

Indeed, it is easily seen that $G \in C^2(\Sigma(u^0))$, and (2.7) implies (4.2), while by (2.6) there holds (4.3). Thus, we have

$$\epsilon_k := \|\mathcal{R}_v(u^0, v^k)\|_{\mathcal{W}*} \to 0 \quad \text{as} \quad k \to +\infty,$$

which by (2.5) yields:

$$\left|\int (\nabla u_i^0, \nabla \xi) - \int a_i(u_i^0)^q \xi - \int \lambda_k g_i(x, u^0) \xi\right| \le \epsilon_k \|\xi\|_{\mathcal{W}}, \quad \forall \xi \in W.$$

 $i = 1, \ldots, m$. Now passing to the limit as $k \to +\infty$ we obtain (1.1).

3. Proof of main results

Proof of Theorem 1.1

Let us prove (1^o) . Let $\overline{w} := (w_1, \ldots, w_m)$, where $w_i, i = 1, \ldots, m$ is a solution of (2.3). By the above $\overline{w} \in \mathcal{W}_s$, and thus, we have

$$\lambda_s^* := \sup_{u \in \mathcal{W}_s} \inf_{v \in \Sigma(u)} \mathcal{R}(u, v) \ge \inf_{v \in \Sigma(\overline{w})} \frac{\int (\nabla w_i, \nabla v_i) - \int a_i w_i^q v_i}{\int g_i(x, \overline{w}) v_i} = 0$$

Since $0 \leq \lambda_s^* \leq +\infty$, there exists a maximizing sequence $u^n \in \mathcal{W}_s$, $n = 1, \ldots$, such that

$$\lambda_n := \lambda(u^n) := \inf_{v \in \Sigma(u^n)} \mathcal{R}(u^n, v) \to \lambda_s^* \text{ as } n \to +\infty.$$

By Lemma 2.1,

(3.1)
$$\begin{cases} -\Delta u_i^n = a_i (u_i^n)^q + \lambda_n g_i(x, u^n), & x \in \Omega, \\ u_i^n |_{\partial \Omega} = 0, & i = 1, \dots, m. \end{cases}$$

Testing (3.1) by u_i , i = 1, ..., m and integrating by parts we derive

(3.2)
$$\|u^n\|_{1,2}^2 = \int a_i |u_i^n|^{q_i+1} + \lambda_n \int g_i(x, u^n) u_i^n, \quad n = 1, \dots.$$

Since $u^n \in \mathcal{W}_s$, $n = 1, \dots,$ (3.3)

$$\lambda_1(F(u^n,\lambda_n)) := \inf_{\phi \in \mathcal{W}} \frac{\int |\nabla \phi|^2 - q_i \int a_i(u_i^n)^{q_i-1} |\phi_i|^2 - \lambda_n \int g_{i,u_j}(u^n) \phi_j \phi_i}{\int |\phi|^2} \ge 0.$$

Hence

(3.4)
$$||u^n||_{1,2}^2 \ge q_i \int a_i |u^n|^{q_i+1} + \lambda_n \int g_{i,u_i}(u^n)(u_i^n)^2 \quad n = 1, \dots$$

Subtraction (3.2) from (3.4), and using (g_2) we obtain

$$(1-q_i)\int a_i |u^n|^{q_i+1} \ge \lambda_n \int \left(g_{i,u_i}(u^n)(u_i^n)^2 - g_i(x,u^n)u_i^n\right) \ge \lambda_n (c_2 ||u^n||_{\gamma_0}^{\gamma_0} + c_3 ||u^n||_{\gamma}^{\gamma}), \quad n = 1, \dots.$$

Applying Hölder's inequality we derive

(3.5)
$$C_1 \ge \lambda_n (c_2 \| u^n \|_{\gamma_0}^{\gamma_0 - q} + c_3 \| u^n \|_{\gamma}^{\gamma - q}), \quad n = 1, \dots$$

Let us show that there exists $C_2 \in (0, +\infty)$ which does not depends on $n = 1, \ldots$, such that

(3.6)
$$||u^n||_{\gamma_0}, ||u^n||_{\gamma} \ge C_2, \quad n = 1, \dots, .$$

To this end we need the following assertion that is derived by the same method as Lemma 3.3 in [2]

Lemma 3.1. Assume that f(t) is a function such that $t^{-1}f(t)$ is decreasing for t > 0, $a \in L^{\infty}$, a > 0 in Ω . Let v and w satisfy: u > 0, w > 0 in Ω , v = w = 0 on $\partial\Omega$, and

$$-\Delta w \le a f(w), \quad -\Delta u \ge a f(u), \quad in \quad \Omega.$$

Then $u \geq w$.

By the assumption $g_i(x, u) \ge 0, x \in \Omega, i = 1, ..., m, u \in \mathbb{R}^m_+$, and therefore,

$$-\Delta u_i^n \ge a_i(u_i^n)^q$$
 in Ω , $i = 1, \ldots, m$, $n = 1, \ldots$

Hence, (2.3) and Lemma 3.1 yield

(3.7)
$$u_i^n \ge w_i, \quad i = 1, \dots, m, \quad n = 1 \dots,$$

and as a result, we get (3.6). Clearly, (3.5), (3.6) imply that $\lambda_s^* < +\infty$, and $\|u^n\|_{\gamma} \leq C_2, n = 1, \ldots$ This by (3.2), (g_1) implies that $\|u^n\|_{1,2} \leq C_3, n = 1, \ldots$, where $C_2, C_3 \in (0, +\infty)$ do not depend on $n = 1, \ldots$ Thus, (u^n) is bounded in \mathcal{W} , and therefore, by the Banach–Alaoglu and the Sobolev theorems there exists a subsequence (again denoted by (u^n)) such that

(3.8)
$$u^n \to u_s^*$$
 weakly in \mathcal{W} ,

(3.9)
$$u^n \to u^*_s$$
 strongly in $(L^r)^m$, $1 \le r < 2^*$

as $n \to +\infty$ for some $u_s^* \in \mathcal{W}$. From (3.7) it follows that $u_{s,i}^* \ge w_i > 0, i = 1, \ldots, m, n = 1 \ldots$

Passing to the limit in (3.1) as $n \to +\infty$ we obtain

(3.10)
$$-\Delta u_{s,i}^* = a_i (u_i^*)^q + \lambda_s^* g_i(x, u_s^*), \quad x \in \Omega, \quad i = 1, \dots, m.$$

Using Proposition 2.2 we conclude that $u_s^* \in C^{1,\alpha}(\overline{\Omega})$ for any $\alpha \in (0,1)$ and $u_i^* > 0$ in Ω , $i = 1, \ldots, m$. Thus, we obtain (1^o) , (a).

To show (1°), (b), suppose conversely that for $\lambda > \lambda_s^*$ there exists a stable weak non-negative solution u_{λ} of (1.1). Then by Proposition 2.2, $u_{\lambda} \in \mathcal{W}_s$, and consequently, (1.4) yields $\inf_{v \in \Sigma(u_{\lambda})} \mathcal{R}(u_{\lambda}, v) < \lambda$. Hence, there exists $v \in \Sigma(u_{\lambda})$ such that $\mathcal{R}(u_{\lambda}, v) < \lambda$. Assume that $\int u_{\lambda,i}^q v_i > 0$. Then

$$\int (\nabla u_{\lambda,i}, \nabla v_i) - \int a_i u_{\lambda,i}^q v_i - \lambda \int g_i(x, u_\lambda) v_i < 0$$

which contradicts (1.1), and as a result we get (1°) , (b).

Let us prove (2°). For simplicity we assume that d > 2. Using $\gamma < 2^*$, it is not hard to show from (3.8),(3.9), (3.10), and $u_s^* \neq 0$ that

(3.11)
$$u^n \to u_s^*$$
 strongly in \mathcal{W} as $n \to +\infty$.

Clearly, by the maximum principle $w_i \in S(\delta)$, $i = 1, \ldots, m$ with some $\delta > 0$. Hence, (3.7) imply that $u_n \in S^m(\delta)$, $n = 1, \ldots$. The assumption $q_i < (2^* - 2)/2 \equiv 2/(d-2)$ for d > 2 and $\gamma < 2^*$ implies by Proposition 2.1 that $F_u(u, \lambda) \in C(S_W^m(\delta); \mathcal{L}(W, W^*))$. Hence,

(3.12)
$$\langle F_u(u_n,\lambda_n)(\phi),\psi\rangle \to \langle F_u(u_s^*,\lambda_s^*)(\phi),\psi\rangle \text{ as } n\to+\infty, \quad \forall \phi,\psi\in\mathcal{W},$$

and consequently, $\lambda_1(F_u(u_n, \lambda_n)) \to \lambda_1(F_u(u_s^*, \lambda_s^*)) \ge 0$ as $n \to +\infty$. Thus we get that $u_s^* \in \mathcal{W}_s$.

Let us show that $\lambda_1(F_u(u_s^*, \lambda_s^*)) = 0$. Suppose, contrary to our claim, that $\lambda_1(F_u(u_s^*, \lambda_s^*)) > 0$. Then $F_u(u_s^*, \lambda_s^*)(\cdot) : \mathcal{W} \to \mathcal{W}^*$ is nonsingular linear operator. From Proposition 2.1 we have $F(\cdot, \lambda) \in C^1(S_{C_1}^m; \mathcal{L}(\mathcal{W}, \mathcal{W}^*))$. Hence, by the Implicit Functional Theorem (see, e.g., [12]) there is a neighbourhood $V \times U \subset \mathbb{R} \times S_{C_1}^m$ of (λ_s^*, u_s^*) and a mapping $V \ni \lambda \mapsto u_\lambda \in U$ such that $u_\lambda|_{\lambda=\lambda_s^*} = u_s^*$ and $F(u_\lambda, \lambda) = 0$, $\forall \lambda \in V$. Furthermore, the map $u_{(\cdot)} : V \to U$ is continuous. Since $\lambda_1(F_u(u_s^*, \lambda_s^*)) > 0$, there exists a neighbourhood $V_1 \subset V$ of λ_s^* such that $\lambda_1(F_u(u_\lambda, \lambda)) > 0$ for every $\lambda \in V_1$. However, this contradicts assertion (1^o) of the theorem. Thus, $\lambda_1(F_u(u_s^*, \lambda_s^*)) = 0$, and (u_s^*, λ_s^*) is a saddle-node type bifurcation point of (1.1) in \mathcal{W}_s . Since (1^o) , (b), (u_s^*, λ_s^*) is a maximal saddle-node type bifurcation point of (1.1) in \mathcal{W}_s .

Finally, let us show that $0 < \lambda_s^*$. Suppose the converse $\lambda_s^* = 0$. Then by the above $0 = \lambda_1(F_u(u_s^*, \lambda_s^*)) = \lambda_1(-\Delta - q|u_s^*|^{q-1})$. However, this is clearly impossible.

Proof of Theorem 1.2 is similar to the proof of Theorem 1.1. We only need to show assertion (c) of Theorem. Indeed, by the construction there are sequences λ_n , $u_{as}^n \in \mathcal{W}_{as}$, $n = 1, \ldots$ such that $F_u(u_{as}^n, \lambda_n) = 0$, $n = 1, \ldots, \lambda_n \to \lambda_{as}^*$, and $u_{as}^n \to u_{as}^*$ strongly in \mathcal{W} as $n \to +\infty$. Moreover, $u_{as}^* \in \overline{\mathcal{W}}_{as}$, and $\lambda_1(F_u(u_{as}^*, \lambda_{as}^*)) = 0$. Thus, $u_{as}^* \in \overline{\mathcal{W}}_{as} \setminus \mathcal{W}_{as}$. On the other hand, $u_{as}^n \in \mathcal{W}_{as}$, $n = 1, \ldots$ Hence, $u_{as}^n \neq u_{as}^*$, $n = 1, \ldots$, and we thus obtain the proof of (c).

4. Appendix A

Let X be a Banach space and $V \subset X$ be an open domain. Denote $B_r := \{ \phi \in X : \|\phi\|_X \leq r \}, r > 0$. Assume that $G : V \to \mathbb{R}, G \in C^2(V)$. Consider

(4.1)
$$\hat{G} = \inf_{v \in V} G(v).$$

Theorem 4.1. Assume that $|\hat{G}| < +\infty$. Suppose that there exist $\tau_0, a_0, C_0 \in (0, +\infty)$, and a minimizing sequence $(v_k) \subset V$ of (4.1) such that

(4.2)
$$\|G_{vv}(v_k + \tau\phi)\|_{(X \times X)^*} < \frac{C_0}{(1 - |\tau|a_0)^2} < +\infty,$$

(4.3)
$$v_k + \tau \phi \in V, \quad \forall \tau \in (-\tau_0, \tau_0), \quad \forall \phi \in B_1, \quad \forall k = 1, \dots$$

Then

$$\|G_v(v_k)\|_{X^*} := \sup_{\xi \in X \setminus 0} \frac{|G_v(v_k)(\xi)|}{\|\xi\|_X} \to 0 \quad as \quad k \to +\infty.$$

Proof. Suppose, contrary to our claim, that there exists $\alpha > 0$ such that

$$||G_v(v_k)||_{X^*} > \alpha, \quad \forall k = 1, \dots$$

This means that for every k = 1, ...,there exists $\phi_k \in V, \|\phi_k\|_X = 1$ such that $|G_v(v_k)(\phi_k)| > \alpha$. By the Taylor expansion

$$G(v_{k} + \tau\phi_{k}) = G(v_{k}) + \tau G_{v}(v_{k})(\phi_{k}) + \frac{\tau^{2}}{2}G_{vv}(v_{k} + \theta_{k}\tau\phi_{k})(\phi_{k},\phi_{k}),$$

for sufficiently small $|\tau|$, and some $\theta_k \in (0, 1)$, $k = 1, \ldots$ Suppose, for definiteness, that $G_v(v_k)(\phi_k) > \alpha$. Then for $\tau \in (-\tau_0, 0)$, by (4.2)

$$G(v_k + \tau \phi_k) \le G(v_k) + \tau \alpha + \frac{\tau^2}{2} \frac{C_0}{(1 + \tau a_0)^2}, \quad k = 1, \dots$$

It is easily seen that there exists $\tau_1 \in (0, \tau_0)$ such that

$$\kappa(\tau) := \tau \left(\alpha + \frac{\tau}{2} \frac{C_0}{(1 + \tau a_0)^2} \right) < 0, \quad \forall \tau \in (-\tau_1, 0).$$

Since (v_k) is a minimizing sequence, for any $\varepsilon > 0$ there exists $k(\varepsilon)$ such that

$$G(v_k) < \hat{G} + \varepsilon, \quad \forall k > k(\varepsilon).$$

Take $\tau \in (-\tau_1, 0)$ and $\varepsilon_0 = -\kappa(\tau)/2$. Then by the above

$$G(v_k + \tau \phi_k) < \hat{G} + \varepsilon_0 + \kappa(\tau) = \hat{G} + \kappa(\tau)/2 < \hat{G}, \quad \forall k > k(\varepsilon_0),$$

and thus, in view of (4.3) we get a contradiction.

References

- C.O. Alves, D.G. De Figueiredo, Nonvariational elliptic systems via Galerkin methods. Function Spaces, Differential Operators and Nonlinear Analysis: The Hans Triebel Anniversary Volume, (2003) 47-57.
- [2] A. Ambrosetti, H. Brezis, G. Cerami, Combined effects of concave and convex nonlinearities in some elliptic problems, J. Funct. Anal. 122(2), (1994) 519–543.
- [3] H. Brezis and L. Oswald, Remarks on sublinear elliptic equations. Nonlinear Analysis: Theory, Methods & Applications, 10(1), (1986) 55-64.
- [4] K. J. Brown, M. M. A. Ibrahim, R. Shivaji, S-shaped bifurcation curves. Nonlinear Analysis: Theory, Methods & Applications, 5(5), (1981) 475-486.
- [5] L. Caffarelli, S. Patrizi, V. Quitalo, On a long range segregation model. Journal of the European Mathematical Society (EMS Publishing). Dec 1;19(12) (2017).
- [6] M.L.Carvalho, Y. Il'yasov, C.A. Santos, Existence of S-shaped type bifurcation curve with dual cusp catastrophe via variational methods. Journal of Differential Equations, 334, (2022) 256-279.
- [7] M. Clapp, A. Szulkin, Non-variational weakly coupled elliptic systems, Analysis and Mathematical Physics, 12(2), (2022): 57.
- [8] M. G. Crandall, P. H. Rabinowitz, Bifurcation, perturbation of simple eigenvalues, and linearized stability. Arch. Rat. Mech. Anal. 52(2), (1973) 161–180.
- M. G. Crandall, P. H. Rabinowitz, Some continuation and variational methods for positive solutions of nonlinear elliptic eigenvalue problems, Arch. Ration. Mech. Anal. 58, (1975) 207–218.
- [10] E.C.M. Crooks, E.N. Dancer, Highly nonlinear large-competition limits of elliptic systems. Nonlinear Analysis: Theory, Methods & Applications, 73(5), (2010) 1447-1457.
- [11] L. Dupaigne, Stable Solutions of Elliptic Partial Differential Equations, Chapman & Hall/CRC Monographs and Surveys in Pure and Applied Mathematics 143, Boca Raton, FL, 2011.
- [12] J. Dieudonné, Foundations of Modern Analysis. Academic Press, New York, 1964.
- [13] I. Ekeland, On the variational principle. J. Math. Anal. Appl. 47(2), (1974) 324–353.
- [14] M. Escobedo, T. Cazenave, and F. Dickstein, A semilinear heat equation with concaveconvex nonlinearity, Rend. Mat. Appl. (7) 19 (1999) 211–242.
- [15] D. De Figueiredo, Semilinear elliptic systems: Existence, multiplicity, symmetry of solutions, Handbook of Differential Equations: Stationary Partial Differential Equations, 5 (2008) 1–48.
- [16] D. Gilbarg, N.S. Trudinger, Elliptic partial differential equations of second order. Berlin: Springer, 1977.
- [17] R. Gilmore, Catastrophe theory for scientists and engineers. Courier Corporation, 1993.
- [18] X. He, M. Squassina, and W. Zou, The Nehari manifold for fractional systems involving critical nonlinearities. Communications on Pure and Applied Analysis, 15(4), (2016) 1285-1308.
- [19] T.S. Hsu, H.L. Lin, Multiple positive solutions for a critical elliptic system with concave—convex nonlinearities. Proceedings of the Royal Society of Edinburgh Section A: Mathematics, 139(6), (2009) 1163-1177.
- [20] Y.S. Il'yasov, Bifurcation calculus by the extended functional method, Funct. Anal. & Appl. 41 (1), (2007) 18–30.

11

- [21] Y. Il'yasov, A duality principle corresponding to the parabolic equations, Physica D, 237(5), (2008) 692–698.
- [22] Y.S. Il'yasov, A.E. Ivanov, Finding bifurcations for solutions of nonlinear equations by quadratic programming methods, Comp. Math. Math. Phys. 53(3), (2013) 350–364.
- [23] Y.S. Il'yasov, A.E. Ivanov, Computation of maximal turning points to nonlinear equations by nonsmooth optimization, Opt. Meth. and Softw. 31(1), (2016) 1–23.
- [24] Y. Ilyasov, Finding Saddle-Node Bifurcations via a Nonlinear Generalized Collatz-Wielandt Formula, International Journal of Bifurcation and Chaos 31, no. 01 (2021): 2150008.
- [25] Y. Il'yasov, A finding of the maximal saddle-node bifurcation for systems of differential equations. Journal of Differential Equations 378, (2024) 610-625.
- [26] H.B. Keller, Numerical solution of bifurcation and nonlinear eigenvalue problems, in: Application of bifurcation theory, ed. 2. Rabinowitz (Academic Press, New York), 1977, 359–384.
- [27] H. Kielhöfer, Bifurcation theory: An introduction with applications to PDEs. Vol. 156, Springer Science & Business Media, 2006.
- [28] P. Korman, On uniqueness of positive solutions for a class of semilinear equations. Discrete and Continuous Dynamical Systems, 8(4), (2002) 865-872.
- [29] O. Ladyzhenskaya, N. Uraltseva, Linear and quasilinear elliptic equations, Acad. Press, New York 20, 1968.
- [30] T. Ouyang, and J. Shi, Exact multiplicity of positive solutions for a class of semilinear problem, II. Journal of Differential Equations, 158(1), (1999) 94-151.
- [31] P.D. Salazar ,Y. Ilyasov, L.F. Alberto, E.C. Costa, M.B. Salles, Saddle-node bifurcations of power systems in the context of variational theory and nonsmooth optimization, IEEE Access. 16, 2020; 8:110986-93.
- [32] R. Seydel, Practical Bifurcation and Stability Analysis, Interdisciplinary Applied Mathematics Vol. 5, Springer, 2010.
- [33] M.Tang, Exact multiplicity for semilinear elliptic Dirichlet problems involving concave and convex nonlinearities. Proceedings of the Royal Society of Edinburgh Section A: Mathematics, 133(3), (2003) 705-717.
- [34] N. N. Ural'tseva, Boundary-value problems for quasi-linear elliptic equations and systems with principal part of divergence type, Dokl. Akad. Nauk SSSR, 147:2 (1962), 313–316.
- [35] T.F. Wu, The Nehari manifold for a semilinear elliptic system involving sign-changing weight functions. Nonl. Anal.: Th. Meth. & Appl., 68(6), (2008) 1733-1745.

INSTITUTE OF MATHEMATICS OF UFRC RAS, 112, CHERNYSHEVSKY STR., 450008 UFA, RUSSIA *Email address:* ilyasov02@gmail.com