

ULAM'S METHOD FOR COMPUTING STATIONARY DENSITIES OF INVARIANT MEASURES FOR PIECEWISE CONVEX MAPS WITH COUNTABLY INFINITE NUMBER OF BRANCHES

MD SHAFIQUUL ISLAM, PAWEŁ GÓRA, AND A H M MAHBUBUR RAHMAN

ABSTRACT. Let $\tau : I = [0, 1] \rightarrow [0, 1]$ be a piecewise convex map with countably infinite number of branches. In [Góra et al.(2022)], the existence of absolutely continuous invariant measure (ACIM) μ for τ and the exactness of the system (τ, μ) has been proven. In this paper, we develop an Ulam method for approximation of f^* , the density of ACIM μ . We construct a sequence $\{\tau_n\}_{n=1}^\infty$ of maps $\tau_n : I \rightarrow I$ s. t. τ_n has a finite number of branches and the sequence τ_n converges to τ almost uniformly. Using supremum norms and Lasota-Yorke type inequalities, we prove the existence of ACIMs μ_n for τ_n with the densities f_n . For a fixed n , we apply Ulam's method with k subintervals to τ_n and compute approximations $f_{n,k}$ of f_n . We prove that $f_{n,k} \rightarrow f^*$ as $n \rightarrow \infty, k \rightarrow \infty$, both a.e. and in L^1 .

We provide examples of piecewise convex maps τ with countably infinite number of branches, their approximations τ_n with finite number of branches and for increasing values of parameter k show the errors $\|f^* - f_{n,k}\|_1$.

1. INTRODUCTION

The existence and properties of absolutely continuous invariant measures (ACIMs) of deterministic dynamical systems reflect their long time behaviour and play an important role in understanding their chaotic nature [Boyarsky & Góra(1997), Lasota & Mackey(1994), Lasota & Yorke(1973)]. Let $\mathcal{B}$ be the Lebesgue σ -algebra of subsets of $I = [0, 1]$ and let λ be the normalized Lebesgue measure on I . Let $\tau : I \rightarrow I$ be a non-singular measurable transformation. A measure μ on $\mathcal{B}$ is τ -invariant if $\mu(\tau^{-1}(A)) = \mu(A)$ for all $A \in \mathcal{B}$. The Frobenius-Perron (F-P) operator $P_\tau : L^1(I, \mathcal{B}, \lambda) \rightarrow L^1(I, \mathcal{B}, \lambda)$ of τ plays an important role for the existence, approximations and properties of ACIMs. The F-P operator P_τ is defined by

$$(1.1) \quad \int_A P_\tau f \, d\lambda = \int_{\tau^{-1}(A)} f \, d\lambda, \quad \forall A \in \mathcal{B}, \forall f \in L^1.$$

It can be shown that for τ with countable number of monotonic branches, the F-P operator P_τ has the following representation [Boyarsky & Góra(1997)] :

$$(1.2) \quad P_\tau f(x) = \sum_{i \geq 1} \frac{f(\tau_i^{-1}(x))}{|\tau'(\tau_i^{-1}(x))|} \chi_{I_i}(x) = \sum_{z \in \{\tau^{-1}(x)\}} \frac{f(z)}{|\tau'(z)|},$$

Key words and phrases. Stationary densities; Absolutely continuous invariant measures; Ulam's method; Convex maps with countably infinite number of branches; Frobenius-Perron (F-P) Operator.

where τ_i^{-1} are inverse branches of $\tau_i = \tau|_{I_i}$, $i \geq 1$, and $\{I_i\}_{i \geq 1}$ are the subintervals of the monotonicity partition. The F-P operator P_τ has a number of useful properties including: (i) Linearity: $P_\tau : L^1 \rightarrow L^1$ is a linear operator; (ii) Positivity: If $f \in L^1$ and $f \geq 0$, then $P_\tau f \geq 0$; (iii) Contractivity: $P_\tau : L^1 \rightarrow L^1$ is a contraction, i.e., $\|P_\tau f\|_1 \leq \|f\|_1$ for any $f \in L^1$. Moreover, $P_\tau : L^1 \rightarrow L^1$ is continuous with respect to norm topology; (iv) P_τ preserves integrals, i.e., $\int_0^1 P_\tau f d\lambda = \int_0^1 f d\lambda$; (v) If $\tau_1, \tau_2 : I \rightarrow I$ are non-singular transformations, then $P_{\tau_1 \circ \tau_2} f = P_{\tau_1} \circ P_{\tau_2} f$. In particular, $P_{\tau^n} f = P_\tau^n f$; (vi) Operator P_τ is conjugated to the Koopman operator $Kg = g \circ \tau$ on L^∞ , i.e.,

$$\int_I g \cdot P_\tau f d\lambda = \int_I g \circ \tau \cdot f d\lambda, f \in L^1, g \in L^\infty;$$

(vii) P_τ has a fixed point f^* if and only if the measure $\mu = f^* \cdot \lambda$ defined by $\mu(A) = \int_A f^* d\lambda$ is τ -invariant. From the property (vii) above one can see that the fixed point of P_τ is the density f^* of the ACIM μ of τ (see [Boyarsky & Góra(1997)]).

In [Lasota & Yorke(1973)], Lasota and Yorke proved the existence of ACIMs for piecewise expanding maps. In this paper, we consider transformations that are not necessarily expanding, i.e., their derivatives may be smaller than 1, but they possess another property which makes them very special, namely piecewise convexity. In [Lasota & Yorke(1982)], the authors studied the existence of ACIMs and the exactness for piecewise convex transformations with a finite number of branches and with a strong repeller. The authors in [Lasota & Yorke(1982)] considered the following properties as key properties for the proof of the existence of ACIMs (i) the F-P operator P_τ maps non-increasing functions to non-increasing functions; (ii) If $f : [0, 1] \rightarrow \mathbb{R}^+$ is a non-increasing function, then

$$\|P_\tau f\|_\infty \leq A \|f\|_\infty + B \|f\|_1,$$

where $A < 1$ and $B > 0$ are some constants. In [Rahman(2019)], the author studied ACIMs for piecewise convex maps with infinite number of branches, where 1 is the limit point of partition points. The literature on piecewise convex maps with a finite number of branches is quite rich, see for example [Bose et al.(2002)], [Ding(1996)], [Inoue(1992)], [Inoue(2023)], [Jabłoński & Malczak(1984)], [Jabłoński & Malczak(1983)].

In [Góra et al.(2022)], we have studied ACIMs of maps in two classes $\mathcal{T}_{pc}^\infty(I)$, $\mathcal{T}_{pc}^{\infty,0}(I)$ of piecewise convex maps $\tau : I \rightarrow I$ with countably infinite number of branches. Any τ in the first class $\mathcal{T}_{pc}^\infty(I)$ is a piecewise convex map $\tau : I \rightarrow I$ with a countable number of branches and arbitrary countable number of limit points of partition points separated from 0. Let $\{0 = a_0, a_1, a_2, \dots, a_n, \dots\}$ be a countable partition of I such that $a_0 < a_1$ and all $a_2, a_3, \dots \in [a_1, 1]$. We do not assume the sequence $\{a_2, \dots, a_n, \dots\}$ to be increasing or decreasing. For any $i \in \{0, 1, 2, \dots\}$, let $n(i)$ be the index such that the interval $[a_i, a_{n(i)}]$ does not contain any other points of the partition. If a_k is the limit point of decreasing subsequence of a_n 's, the $n(k)$ is not defined. For such points the interval $[a_k, a_k]$ is not considered an element of the partition. We allow any countable number of the limit points of the points of the partition. We assume that the union of intervals $[a_i, a_{n(i)}]$ covers the whole interval I except of a countable number of points. At any such point x we assume $\tau(x) = 0$ and $\tau'(x) = +\infty$. These points do not contribute to the images of the P_τ operator.

Any τ in the second class $\mathcal{T}_{pc}^{\infty,0}(I)$ is a piecewise convex map $\tau : I \rightarrow I$ with countably infinite number of branches such that there exists a countable partition

$\{0 = a_0 < \cdots < a_{0,-n} < a_{0,-(n-1)} < \cdots < a_{0,-2} < a_{0,-1} = a_1, a_2, a_3, \dots, a_n, \dots\}$ of I with $\lim_{n \rightarrow \infty} a_{0,-n} = 0$. It is proved in [Góra et al.(2022)] that any $\tau \in \mathcal{T}_{pc}^\infty(I) \cup \mathcal{T}_{pc}^{\infty,0}(I)$ has the unique stationary density f^* of the ACIM μ^* . In this paper, we apply Ulam's method for approximation of f^* . Fixed points of the F-P operator P_τ of $\tau \in \mathcal{T}_{pc}^\infty(I) \cup \mathcal{T}_{pc}^{\infty,0}(I)$ are the stationary densities of τ . The F-P operator P_τ is an infinite dimensional operator. Except some simple cases (where τ is piecewise linear Markov and the Frobenius-Perron operator has a matrix representation), it is not easy to obtain an analytical solution of the F-P equation $P_\tau f = f$.

A method for numerical computations of invariant measures of dynamical systems was suggested by Ulam [Ulam(1960)]. For piecewise expanding deterministic transformations [Lasota & Yorke(1973)], T-Y Li [Li(1976)] first proved the convergence of Ulam's approximation. Since then, Ulam's method has been applied to one and higher dimensional expanding deterministic transformations (see, for example, [Boyarsky & Lou(1991), Bose & Murray(2001), Boltt & Santitissadeekorn(2013), Ding & Zhou(1996)]). Ulam's method is one of the most used and the best understood methods among numerical methods for the approximation of ACIMs for deterministic maps [Li(1976), Ulam(1960)]. However, approximate invariant densities via Ulam's methods are piecewise constant on subintervals of the partition of the state space. In [Miller(1994)], Miller proved the convergence of Ulam's method for piecewise convex transformations with a finite number of branches and a strong repeller. J. Ding [Ding(1996)] developed and presented piecewise linear and piecewise quadratic Markov finite approximation methods for such piecewise convex transformations.

In Section 2, we introduce notations and review the existence of stationary densities of ACIMs for $\tau \in \mathcal{T}_{pc}^\infty(I) \cup \mathcal{T}_{pc}^{\infty,0}(I)$. In Section 3, we construct a sequence $\{\tau_n\}_{n=1}^\infty$ of maps $\tau_n : I \rightarrow I$ s. t. τ_n has a finite number of branches and the sequence τ_n converges to τ almost uniformly. Using supremum norms and Lasota-Yorke type inequalities, we prove the existence of densities f_n of ACIMs μ_n for τ_n . Moreover, we prove that f_n converges to f^* . In Section 3.2 we apply Ulam's method with k subintervals to τ_n , compute an approximation $f_{n,k}$ of f_n and prove that $f_{n,k} \rightarrow f_n$ as $k \rightarrow \infty$. In Theorem 5 we prove that $f_{n,k} \rightarrow f^*$, as $n, k \rightarrow \infty$, both a.e. and in L^1 . Theoretically, we could apply Ulam's method directly to the map τ . Practically, such method would be impossible since the calculation of the Ulam's matrix $M_\tau^{(k)}$ would require solving of infinitely many equations. In Section 4, we present numerical examples.

2. ACIMs FOR PIECEWISE CONVEX MAPS IN CLASSES $\mathcal{T}_{pc}^\infty(I)$ AND $\mathcal{T}_{pc}^{\infty,0}(I)$

In this section, we review results on the existence of stationary densities of ACIMs of piecewise convex maps with countably infinite number of branches. We closely follow [Góra et al.(2022)].

2.1. ACIMs for piecewise convex maps in $\mathcal{T}_{pc}^\infty(I)$. For the maps in $\mathcal{T}_{pc}^\infty(I)$ the limit points of the partition points are separated from 0.

We say that $\tau \in \mathcal{T}_{pc}^\infty(I)$ if

- (1) $\tau_0 = \tau|_{[0,a_1]}$ is continuous and convex;
- $\tau_i = \tau|_{[a_i, a_{n(i)})}$ is continuous and convex, $i = 1, 2, \dots$;

$$(2) \quad \tau(a_i) = 0, \tau'(a_i) > 0, i = 1, 2, \dots;$$

$$(3) \quad \tau(0) = 0, \tau'(0) = \alpha_1 > 1;$$

$$(4) \quad \sum_{i=1}^{\infty} \frac{1}{\tau'(a_i)} < \infty.$$

For $\tau \in \mathcal{T}_{pc}^{\infty}(I)$, $f \in L^1(I)$, $f \geq 0$ the F-P operator P_{τ} can be represented as

$$(2.1) \quad P_{\tau}f(x) = \frac{f(\tau_0^{-1}(x))}{\tau'(\tau_0^{-1}(x))} \chi_{\tau[0, a_1)}(x) + \sum_{i=1}^{\infty} \frac{f(\tau_i^{-1}(x))}{\tau'(\tau_i^{-1}(x))} \chi_{\tau[a_i, a_{n(i)})}(x).$$

The following results are proved in [Góra et al.(2022)].

Lemma 1. *Let $\tau \in \mathcal{T}_{pc}^{\infty}(I)$, $f \in L^1(I)$, $f \geq 0$, f be non-increasing. Then*

$$(1) \quad P_{\tau}(f) \in L^1(I).$$

$$(2) \quad P_{\tau}(f) \geq 0.$$

$$(3) \quad P_{\tau}(f) \text{ is non-increasing.}$$

$$(4) \quad \|P_{\tau}(f)\|_{\infty} \leq C \|f\|_{\infty}, \text{ where } C = \left(\frac{1}{\alpha_1} + \sum_{i=1}^{\infty} \frac{1}{\tau'(a_i)} \right).$$

Proposition 1. *If $f \geq 0$ and f is non-increasing, then $f(x) \leq \frac{1}{x} \lambda(f)$, for $x \in [0, 1]$, where*

$$\lambda(f) = \int_0^1 f(x) d\lambda(x).$$

Lemma 2. *If $f : I \rightarrow \mathbb{R}^+$ is non-increasing and $\tau \in \mathcal{T}_{pc}^{\infty}(I)$, then*

$$(2.2) \quad \|P_{\tau}(f)\|_{\infty} \leq \frac{1}{\alpha_1} \|f\|_{\infty} + D \|f\|_1,$$

$$\text{where } D = \left(\sum_{i=1}^{\infty} \frac{1}{a_i} \frac{1}{\tau'(a_i)} \right).$$

Theorem 1. *Let $\tau \in \mathcal{T}_{pc}^{\infty}(I)$. Then τ admits an ACIM $\mu = f^* \cdot \lambda$ with non-increasing density function f^* . The ACIM μ is unique and the system (τ, μ) is exact.*

2.2. ACIMs for piecewise convex maps in $\mathcal{T}_{pc}^{\infty,0}(I)$. For the maps in $\mathcal{T}_{pc}^{\infty,0}(I)$ the point 0 is the limit point of partition points.

We say that $\tau \in \mathcal{T}_{pc}^{\infty,0}(I)$ if

$$(1) \quad \begin{aligned} \tau_{-j} = \tau|_{[a_0, -(j+1), a_0, -j)} & \text{ is continuous and convex, } j = 1, 2, \dots; \\ \tau_i = \tau|_{[a_i, a_{n(i)})} & \text{ is continuous and convex, } i = 1, 2, \dots; \end{aligned}$$

$$(2) \quad \tau(a_0, -j) = 0, \tau'(a_0, -j) > 0, j = 1, 2, \dots;$$

$$\tau(a_i) = 0, \tau'(a_i) > 0, i = 1, 2, \dots;$$

$$(3) \quad \sum_{i=1}^{\infty} \frac{1}{\tau'(a_i)} < \infty;$$

$$(4) \quad D_1 = \sum_{j=1}^{\infty} \frac{1}{\tau'(a_0, -j)} < 1.$$

Remark 1. *The Condition 3 and the Condition 4 can be replaced by the following condition:*

$$(3^+) \quad \sum_{j=1}^{\infty} \frac{1}{\tau'(a_{0,-j})} + \sum_{i=1}^{\infty} \frac{1}{\tau'(a_i)} < \infty.$$

If (3^+) is satisfied, then we can find an integer $J \geq 1$ such that

$$\sum_{j=J}^{\infty} \frac{1}{\tau'(a_{0,-j})} < 1,$$

and after proper renaming of partition points, the Condition 3 and the Condition 4 are satisfied.

The following results are proved in [Góra et al.(2022)].

Lemma 3. *Let $\tau \in \mathcal{T}_{pc}^{\infty,0}(I)$, $f \in L^1(I)$, $f \geq 0$, f be non-increasing. Then*

- (1) $P_{\tau}(f) \in L^1(I)$.
- (2) $P_{\tau}(f) \geq 0$.
- (3) $P_{\tau}(f)$ is non-increasing.
- (4) $\|P_{\tau}(f)\|_{\infty} \leq C \|f\|_{\infty}$, where $C = \left(D_1 + \sum_{i=1}^{\infty} \frac{1}{\tau'(a_i)}\right)$.

Lemma 4. *If $f : [0, 1] \rightarrow \mathbb{R}^+$ is non-increasing and $\tau \in \mathcal{T}_{pc}^{\infty,0}(I)$, then*

$$(2.3) \quad \|P_{\tau}(f)\|_{\infty} \leq D_1 \|f\|_{\infty} + D \|f\|_1,$$

where $D = \sum_{i=1}^{\infty} \frac{1}{a_i \tau'(a_i)}$.

Theorem 2. *Let $\tau \in \mathcal{T}_{pc}^{\infty,0}(I)$. Then τ admits an ACIM $\mu = f^* \cdot \lambda$ with non-increasing density function f^* . The ACIM μ is unique and the system (τ, μ) is exact.*

3. ULAM'S METHOD FOR PIECEWISE CONVEX MAPS WITH COUNTABLE NUMBER OF BRANCHES

In this section our main goal is to develop Ulam's method for stationary densities of ACIMs for piecewise convex maps with countably infinite number of branches. We do this in the following two steps.

3.1. Approximation of maps with infinite number of branches. In this section we approximate piecewise convex maps with countably infinite number of branches by piecewise convex maps with a finite number of branches.

We concentrate on a case of a map $\tau \in \mathcal{T}_{pc}^{\infty,0}(I)$. On the example of point 0, we show how to perform an approximation of τ with a point of a partition that is a limit point of other points of the partition. The method can be applied to maps in $\mathcal{T}_{pc}^{\infty}(I)$, that have points with this property without substantial changes.

Let $\tau \in \mathcal{T}_{pc}^{\infty,0}(I)$. We assume that there are no other limit points of the partition points. For simplicity we rename the partition points. The new partition points are a_n , $n = 0, 1, 2, \dots$, with $a_0 = 1$, $a_{n+1} < a_n$, $n \geq 0$ and $\lim_{n \rightarrow \infty} a_n = 0$. Then, the assumptions (3) and (4) of Section 2.2 are restated as:

(3') There exists an $N \geq 1$ such that

$$(3.1) \quad D_1 = \sum_{n=N+1}^{\infty} \frac{1}{\tau'(a_n)} < 1.$$

Then, the Lemma 3 and the Lemma 4 of Section 2.2 hold with changed constants

$$(3.2) \quad \begin{aligned} C &= D_1 + \sum_{n=1}^N \frac{1}{\tau'(a_n)} = \sum_{n=1}^{\infty} \frac{1}{\tau'(a_n)}, \\ D &= \sum_{n=1}^N \frac{1}{a_n \tau'(a_n)}. \end{aligned}$$

For $n \geq N$, we construct a sequence $\{\tau_n\}_{n=N}^{\infty}$ of maps $\tau_n : I \rightarrow I$ s.t. τ_n has a finite number of branches and the sequence τ_n converges to τ almost uniformly. Using supremum norms and Lasota-Yorke type inequalities, we prove the existence of stationary densities f_n of ACIMs μ_n for τ_n . We approximate $\tau : I \rightarrow I$ with the following sequence of maps $\tau_n : I \rightarrow I$, $n \geq N$, with a finite number of branches:

$$(3.3) \quad \tau_n(x) = \begin{cases} x/a_n, & 0 \leq x < a_n; \\ \tau(x), & a_n \leq x \leq 1. \end{cases}$$

In the following, we show that for each $n \geq 0$, the map τ_n has an ACIM. It can be easily shown that each τ_n is a piecewise convex maps with the finite partition $\{1 = a_0, a_1, a_2, \dots, a_n, a_{n+1} = 0\}$ and τ_n satisfies following conditions:

- (1) $\tau_{n_j} = \tau_n|_{(a_j, a_{j-1}]}$ is continuous and convex, $j = 1, 2, \dots, n+1$;
- (2) $\tau_n(a_j) = 0, \tau'_n(a_j) > 0, j = 1, \dots, n+1$;
- (3) $\tau_n(0) = 0, \tau'_n(0) = \frac{1}{a_n} > 1$.
- (4) $\sum_{j=1}^{n+1} \frac{1}{\tau'_n(a_j)} < \infty$.

Let $f \in L^1(I), f \geq 0$. Then, the F-P operator P_{τ_n} is defined as

$$(3.4) \quad P_{\tau_n} f(x) = \frac{f(\tau_a^{-1}(x))}{1/a_n} + \sum_{j=1}^n \frac{f(\tau_{n_j}^{-1}(x))}{\tau'_n(\tau_{n_j}^{-1}(x))} \chi_{\tau_n(a_j, a_{j-1}]}(x),$$

where $\tau_a = x/a_n$ on the interval $[0, a_n)$.

Lemma 5. *Let $f \in L^1(I), f \geq 0, f$ be a non-increasing function. Then*

- (1) $P_{\tau_n}(f) \in L^1(I)$ for each $n = 1, 2, \dots$.
- (2) $P_{\tau_n}(f) \geq 0$ for each $n = 1, 2, \dots$.
- (3) $P_{\tau_n}(f)$ is non-increasing for each $n = 1, 2, \dots$.
- (4) $\|P_{\tau_n}(f)\|_{\infty} \leq C_n \|f\|_{\infty} \leq (1+C) \|f\|_{\infty}$ for each $n = 1, 2, \dots$ where $C_n = \left(a_n + \sum_{j=1}^n \frac{1}{\tau'_n(a_j)}\right)$.

Proof. (1) Holds for the F-P operator of any map.

- (2) $P_{\tau_n} f(x)$ is shown in formula (3.4) and each summand is clearly non-negative.
- (3) It can be easily shown that each summand in formula (3.4) is non-increasing.
- (4) We have

$$\begin{aligned}
P_{\tau_n} f(x) &= \frac{f(\tau_n^{-1}(x))}{1/a_n} + \sum_{j=1}^n \frac{f(\tau_{n_j}^{-1}(x))}{\tau'_n(\tau_{n_j}^{-1}(x))} \chi_{\tau_n(a_j, a_{j-1}]}(x) \\
&\leq a_n \|f\|_\infty + \sum_{j=1}^n \frac{\|f\|_\infty}{\tau'_n(\tau_{n_j}^{-1}(x))} \leq a_n \|f\|_\infty + \sum_{j=1}^n \frac{\|f\|_\infty}{\tau'_n(a_j)} \\
&= \left(a_n + \sum_{j=1}^n \frac{1}{\tau'_n(a_j)} \right) \|f\|_\infty \leq (1+C) \|f\|_\infty
\end{aligned}$$

□

Lemma 6. *If $f : [0, 1] \rightarrow \mathbb{R}^+$ is non-increasing, then for each $n \geq N$,*

$$(3.5) \quad \|P_{\tau_n}(f)\|_\infty \leq (a_n + D_1) \|f\|_\infty + D \|f\|_1,$$

Proof. Since f is non-increasing, $f(0) \geq \|f\|_\infty$, and by Lemma 5, $P_{\tau_n} f(0) = \|P_{\tau_n} f\|_\infty$. Now,

$$\begin{aligned}
P_{\tau_n} f(0) &= \frac{f(\tau_n^{-1}(0))}{1/a_n} + \sum_{j=1}^n \frac{f(\tau_{n_j}^{-1}(0))}{\tau'_n(\tau_{n_j}^{-1}(0))} \chi_{\tau_n(a_j, a_{j-1}]}(0) \\
&= \frac{1}{\tau'_n(0)} f(0) + \sum_{j=1}^N \frac{f(\tau_{n_j}^{-1}(0))}{\tau'_n(\tau_{n_j}^{-1}(0))} + \sum_{j=N+1}^n \frac{f(\tau_{n_j}^{-1}(0))}{\tau'_n(\tau_{n_j}^{-1}(0))} \\
&\leq a_n f(0) + \sum_{j=1}^N \frac{f(a_{n_j})}{\tau'_n(a_{n_j})} + \sum_{j=N+1}^n \frac{f(a_{n_j})}{\tau'_n(a_{n_j})} \\
&\leq (a_n + D_1) f(0) + \sum_{j=1}^N \frac{\lambda(f)}{a_j} \frac{1}{\tau'_n(a_j)} \\
&\leq (a_n + D_1) \|f\|_\infty + D \|f\|_1.
\end{aligned}$$

□

Theorem 3. *For each $n \in \mathbb{N}$, τ_n admits an ACIM $\mu_n = f_n^* \cdot \lambda$ with non-increasing density function f_n^* .*

Proof. Our proof works for all $n \geq n_0$, where we have $a_{n_0} + D_1 < 1$ and gives a uniform estimate. For the $n < n_0$ the claim follows from Subsection 2.1.

Let $f \in L^1$ be non-negative and non-increasing. Consider the sequence $\{P_{\tau_n}^k f\}_{k=0}^\infty$. Since every $P_{\tau_n}^k f$ is non-negative and non-increasing, by part (3) of Lemma 5 we can apply Lemma 6 iteratively and obtain

$$\|P_{\tau_n}^k f\|_\infty \leq (a_{n_0} + D_1)^k \|f\|_\infty + \frac{D}{1 - (a_{n_0} + D_1)} \|f\|_1.$$

So the sequence $\{P_{\tau_n}^k f\}_{k=0}^\infty$ is uniformly bounded and weakly compact. By Yosida-Kakutani Theorem [Yosida & Kakutani(1941)], $\frac{1}{k} \sum_{j=1}^k P_{\tau_n}^j f$ converges in L^1 to a P_{τ_n} invariant function f_n^* . It is non-increasing since it is the limit of non-increasing functions. □

Definition 1. Let τ and τ_n , $n = 1, 2, \dots$ be maps from I into itself. We say that τ_n converges to τ almost uniformly if for any $\epsilon > 0$, there exists a measurable set $A_\epsilon \subset [0, 1]$, $\lambda(A_\epsilon) > 1 - \epsilon$, such that $\tau_n \rightarrow \tau$ uniformly on A_ϵ .

Lemma 7. Let τ be a piecewise convex map with countably many branches and the sequence τ_n , $n = 1, 2, \dots$ be defined by formula (3.3). Then, τ_n converges to τ almost uniformly.

Proof. Let $\epsilon > 0$. Choose the decreasing partition $\{1 = a_0, a_1, a_2, \dots, a_n, a_{n+1} = 0\}$ of $[0, 1]$ for τ_n such that $a_n < \epsilon$. Let $A_\epsilon = (a_n, 1)$. Then, $\lambda(A_\epsilon) = 1 - a_n > 1 - \epsilon$ and we have $\tau_n = \tau$ on A_ϵ . $\square$

3.2. Ulam's method. In this section, first, we describe Ulam's method for a finite dimensional approximation $P_{n,k}$ of the Perron-Frobenius operator P_{τ_n} of τ_n . Ulam's method computes $f_{n,k}$ on a partition of k subintervals of the state space as an approximation of the actual stationary density function f_n of τ_n , $n \geq 1$. Moreover, we show that $f_{n,k}$ converges to f_n as $k \rightarrow \infty$. We follow [Li(1976)], [Miller(1994)] and [Ding(1996)]. In general, the Ulam's method is widely studied and used, see for example [Bose & Murray(2001)], [Boltt & Santitissadeekorn(2013)], [Ding & Zhou(1996)], [Froyland(1999)], [Góra & Boyarsky(1989)] and the references within.

Let τ_n be an approximation of $\tau \in \mathcal{T}_{pc}^\infty(I)$. Then, by the Theorem 3, τ_n has an ACIM μ_n with a stationary density function f_n . Now, we describe Ulam's method for approximating f_n . Let k be a positive integer. Let $\mathcal{P}^{(k)} = \{J_1, J_2, \dots, J_k\}$ be a partition of the interval $[0, 1]$ into k equal subintervals. Now, construct the matrix

$$\mathbb{M}_n^{(k)} = \left(\frac{\lambda(\tau_n^{-1}(J_j) \cap J_i)}{\lambda(J_i)} \right)_{1 \leq i, j \leq k}.$$

Let $L^{(k)} \subset L^1([0, 1], \lambda)$ be a subspace of L^1 consisting of functions which are constant on elements of the partition $\mathcal{P}^{(k)}$. We will represent functions in $L^{(k)}$ as vectors: vector $f = [f_1, f_2, \dots, f_k]$ corresponds to the function $f = \sum_{i=1}^k f_i \chi_{J_i}$. Let $Q^{(k)}$ be the isometric projection of L^1 onto $L^{(k)}$:

$$(3.6) \quad Q^{(k)}(f) = \sum_{i=1}^k \left(\frac{1}{\lambda(J_i)} \int_{J_i} f d\lambda \right) \chi_{J_i} = \left[\frac{1}{\lambda(J_1)} \int_{J_1} f d\lambda, \dots, \frac{1}{\lambda(J_k)} \int_{J_k} f d\lambda \right]$$

Let $f = [f_1, f_2, \dots, f_k] \in L^{(k)}$. We define the operator $P_{\tau_n}^{(k)} : L^{(k)} \rightarrow L^{(k)}$ by

$$(3.7) \quad P_{\tau_n}^{(k)} f = [f_1, f_2, \dots, f_k] \cdot \left(\mathbb{M}_n^{(k)} \right)^T,$$

which is a finite dimensional approximation to the operator P_{τ_n} . A^T denotes the transpose of the matrix A .

The following Lemma will be used several times in the sequel.

Lemma 8. Let $\{g_n\}_{n=1,2,\dots}$ be a sequence of non-increasing functions uniformly bounded in L^∞ . If $g_n \rightarrow h$, as $n \rightarrow \infty$, weakly in L^1 , then $g_n \rightarrow h$, as $n \rightarrow \infty$, in L^1 and almost everywhere (a. e.).

Proof. Since g_n 's are non-increasing and uniformly bounded in L^∞ , they are also of uniformly bounded variation. By Helly's Theorem [Rudin(1976)] there is a subsequence g_{n_k} convergent a.e. to some function h_1 . Since $g_{n_k} \rightarrow h_1$ weakly in L^1

we have $h_1 = h$. Considering all possible subsequences we prove that $g_n \rightarrow h$ a.e. Since g_n converge to h a.e. and they are uniformly bounded in L^∞ , the convergence is also in L^1 (e.g. by Lebesgue Dominated Convergence theorem). $\square$

Since each map τ_n is exact [Lasota & Yorke(1982)], by Proposition 1.2 of [Hunt & Miller(1992)] we obtain that the invariant densities $f_{n,k}$ of $P_{\tau_n}^{(k)}$ are unique.

Lemma 9. *The invariant density $f_{n,k}$ of $P_{\tau_n}^{(k)}$ is non-increasing for any $n, k > 1$.*

Proof. Let $f \in L^{(k)}$. Since $P_{\tau_n}^{(k)} f = Q^{(k)} P_{\tau_n} f$, and both operators P_{τ_n} and $Q^{(k)}$ transform non-increasing functions into non-increasing functions, the operators $P_{\tau_n}^{(k)}$ also have this property. Let $f = 1$ be a constant function understood as $[1, 1, \dots, 1] \in L^{(k)}$. It is non-increasing. Thus, all the functions $(P_{\tau_n}^{(k)})^m f$, $m = 1, 2, \dots$, are non-increasing. Similarly as the estimate (3.5) was obtained, they can be shown to be uniformly bounded in L^∞ and thus weakly compact in L^1 . Then, Yosida-Kakutani theorem [Yosida & Kakutani(1941)] shows that the sequence $1/s \sum_{m=1}^s (P_{\tau_n}^{(k)})^m f$ converges in L^1 to the invariant density $f_{n,k}$. By Lemma 8 the convergence is also a.e. and $f_{n,k}$ is non-increasing. $\square$

Using Ulam's method and corresponding convergence analysis described in [Li(1976), Miller(1994), Ding(1996)], the following theorem can be proved.

Theorem 4. *Let $\tau \in \mathcal{T}_{pc}^{\infty,0}(I)$ be a piecewise convex map with countably many branches. Let $\{\tau_n\}_{n=1}^\infty$ be the approximating sequence of piecewise convex maps with a finite number of branches where τ_n are defined in Equation (3.3) in Subsection 3.1. If $f_{n,k}$ is a normalized fixed point of $P_{\tau_n}^{(k)}$, $k = 1, 2, \dots$, defined in (3.7), then the sequence $\{f_{n,k}\}_{k=1}^\infty$ is weakly pre-compact in L^1 . Any limit point f_n^* of the sequence $\{f_{n,k}\}_{k=1}^\infty$ is a fixed point of P_{τ_n} .*

Proof. Let $P_{\tau_n}^{(k)}$ be the Ulam's approximation of the F-P operator P_{τ_n} of τ_n . Let $Q^{(k)}$ be the isometric projection defined in (3.6). It can be shown that (see: (4), page 3 [Miller(1994)]; def. (2.1), page 5 [Li(1976)]):

$$(3.8) \quad P_{\tau_n}^{(k)} Q^{(k)} f = Q^{(k)} P_{\tau_n} Q^{(k)} f.$$

By the definition of $Q^{(k)}$ it is obvious that

$$(3.9) \quad \|Q^{(k)} f\|_\infty \leq \|f\|_\infty.$$

By equations (2.3), (3.8) and (3.9) $P_{\tau_n}^{(k)}$ satisfies the following Lasota-Yorke type inequality

$$(3.10) \quad \|P_{\tau_n}^{(k)} f\|_\infty \leq (a_n + D_1) \|f\|_\infty + D \|f\|_1,$$

which implies the following uniform estimate

$$(3.11) \quad \|f_{n,k}\|_\infty \leq \frac{D}{1 - (a_n + D_1)}.$$

Now, the proof continues exactly as in [Miller(1994)]. $\square$

Theorem 5. *Let $\tau \in \mathcal{T}_{pc}^{\infty,0}(I)$ be a piecewise convex map with countably many branches. As described at the beginning of Subsection 3.1, let $\{\tau_n\}_{n=1}^\infty$ be the approximating sequence of piecewise convex maps with finite numbers of branches. Let*

$P_{\tau_n}^{(k)}$, $k = 1, 2, \dots$ be the sequence of Ulam's operators approximating the operators P_{τ_n} . Let $f_{n,k}$ be the normalized (in L^1) fixed point of $P_{\tau_n}^{(k)}$. Then, the family $\{f_{n,k}\}_{n=1,2,\dots,k=1,2,\dots}$ is weakly compact in L^1 and uniformly bounded in L^∞ . If f_{n_j,k_j} , $j = 1, 2, \dots$ is a weakly convergent subsequence, then it converges in L^1 (and almost everywhere) to a function f which is a fixed point of P_τ , $P_\tau f = f$.

Proof. The stationary densities $\{f_n\}_{n \geq 1}$ of $\{\tau_n\}_{n \geq 1}$ are uniformly bounded in L^∞ by Theorem 3. Moreover, the densities $f_{n,k}$, the piecewise constant approximations of f_n 's are also uniformly bounded in L^∞ by formula (3.11). Thus, the set $\{f_{n,k}\}_{k=1}^\infty$ is weakly compact in L^1 . Assume that $\{f_{n,k}\}_{k=1}^\infty$ has weakly convergent subsequence $\{f_{n_j,k_j}\}$ with limit f . Since the functions $\{f_{n_j,k_j}\}$ are decreasing and uniformly bounded, by Lemma 8 they converge to f a.e. and in L^1 .

It remains to show that f is a fixed point of P_τ , $P_\tau f = f$. We will show that the measures $f d\lambda$ and $(P_\tau f) d\lambda$ are equal. It is enough to show that for any $g \in C(I)$, we have $\int g(f - P_\tau f) d\lambda = 0$. To simplify the notation we assume that the whole sequence $f_{n,k}$ converges to f . We have,

$$\begin{aligned} \left| \int g(f - P_\tau f) d\lambda \right| &\leq \left| \int g(f - f_{n,k}) d\lambda \right| + \left| \int g(f_{n,k} - P_{\tau_n}^{(k)} f_{n,k}) d\lambda \right| \\ &+ \left| \int g(P_{\tau_n}^{(k)} f_{n,k} - P_{\tau_n} f_{n,k}) d\lambda \right| + \left| \int g(P_{\tau_n} f_{n,k} - P_{\tau_n} f) d\lambda \right| + \left| \int g(P_{\tau_n} f - P_\tau f) d\lambda \right|. \end{aligned}$$

Since $f_{n,k} \rightarrow f$ in L^1 , the first and the fourth term go to 0 as $n, k \rightarrow \infty$. Since $f_{n,k}$ are the fixed points of $P_{\tau_n}^{(k)}$, the second term is 0. We have

$$\int g \left[P_{\tau_n}^{(k)} f_{n,k} - P_{\tau_n} f_{n,k} \right] d\lambda = \int g \left[Q^{(k)}(P_{\tau_n} f_{n,k}) - P_{\tau_n} f_{n,k} \right] d\lambda = \int \left[Q^{(k)} g - g \right] P_{\tau_n} f_{n,k} d\lambda,$$

by the self adjointness of $Q^{(k)}$. Since $Q^{(k)} g \rightarrow g$ in L^1 and the densities $\{f_{n,k}\}$ are uniformly bounded, $\int g \left[P_{\tau_n}^{(k)} f_{n,k} - P_{\tau_n} f_{n,k} \right] d\lambda$ converges to 0 as $k \rightarrow +\infty$. By the property (vi) of the F-P operator we have

$$\left| \int g(P_{\tau_n} f - P_\tau f) d\lambda \right| = \left| \int (g \circ \tau_n - g \circ \tau) f d\lambda \right| = \left| \int_{[0, a_n]} (g \circ \tau_n - g \circ \tau) f d\lambda \right| \leq a_n \cdot 2M \cdot \|f\|_1,$$

where $M = \sup |g|$. This shows that $\left| \int g(P_{\tau_n} f - P_\tau f) d\lambda \right|$ converges to 0 as $n \rightarrow +\infty$. $\square$

4. EXAMPLES

Example 1. Consider the piecewise expanding and piecewise linear map $T : [0, 1] \rightarrow [0, 1]$ with countable number of branches defined as

$$(4.1) \quad T(x) = i(i+1) \left(x - \frac{1}{i+1} \right) \quad \text{on} \quad \left[\frac{1}{i+1}, \frac{1}{i} \right], i = 1, 2, \dots$$

It is easy to show that the Lebesgue measure is invariant under T and $f = \mathbf{1}$ is the invariant density of T . Now, consider the conjugation $h : [0, 1] \rightarrow [0, 1]$ defined by $h(x) = 1 - (1 - x)^2$. We construct the piecewise convex map $\tau : [0, 1] \rightarrow [0, 1]$ with countable number of branches defined by $\tau = h^{-1} \circ T \circ h$. See Figure 1 for a graph of τ . The piecewise convex map τ is topologically conjugate to the piecewise linear and piecewise expanding map T via the conjugation h . Therefore, the stationary




FIGURE 1. The graph of the piecewise convex map τ with countably infinite number of branches for Example 1.

density g of τ is given by $g = f \circ h \times |h'|$. Now, $h(x) = 1 - (1 - x)^2$. Hence, $g(x) = f(h(x)) \times |h'(x)| = |2(1 - x)|$.

The map τ is defined on the partition

$$\left\{ \left[1 - \sqrt{\frac{i}{i+1}}, 1 - \sqrt{\frac{i-1}{i}} \right] \right\}_{i=1}^{\infty},$$

i.e., the partition points are

$$\left\{ \dots, 1 - \sqrt{\frac{4}{5}}, 1 - \sqrt{\frac{3}{4}}, 1 - \sqrt{\frac{2}{3}}, 1 - \sqrt{\frac{1}{2}}, 1 \right\}.$$

We have

$$\tau(x) = 1 - \sqrt{1 - i^2 + i(i+1)(1-x)^2} \text{ for } x \in \left[1 - \sqrt{\frac{i}{i+1}}, 1 - \sqrt{\frac{i-1}{i}} \right], \quad i = 1, 2, \dots$$

The first few, starting from 1 to the left, branches of τ are

$$\begin{aligned} \tau_1(x) &= 1 - \sqrt{2}(1-x) && \text{for } 1 - \sqrt{1/2} \leq x \leq 1; \\ \tau_2(x) &= 1 - \sqrt{6(1-x)^2 - 3} && \text{for } 1 - \sqrt{2/3} \leq x < 1 - \sqrt{1/2}; \\ \tau_3(x) &= 1 - \sqrt{12(1-x)^2 - 8} && \text{for } 1 - \sqrt{3/4} \leq x < 1 - \sqrt{2/3}; \\ \tau_4(x) &= 1 - \sqrt{20(1-x)^2 - 15} && \text{for } 1 - \sqrt{4/5} \leq x < 1 - \sqrt{3/4}. \end{aligned}$$

Now, consider the following sequence $\{\tau_n\}_{n \geq 0}$ of piecewise convex maps $\tau_n : [0, 1] \rightarrow [0, 1]$ with a finite number of branches:

$$\tau_n(x) = \begin{cases} \frac{1}{1-\sqrt{\frac{n}{n+1}}}x, & 0 \leq x < 1 - \sqrt{\frac{n}{n+1}}; \\ \tau(x), & 0 \leq x < 1 - \sqrt{\frac{n}{n+1}} \leq x \leq 1. \end{cases}$$

See Figure 2 for a graph of τ_n with $n = 5$. The sequence $\{\tau_n\}_{n \geq 0}$ of piecewise convex




FIGURE 2. The graph of the piecewise convex map τ_n with a finite number of branches ($n = 8$).

map $\tau_n : [0, 1] \rightarrow [0, 1]$ with finite number of branches converges almost uniformly to τ with the countable number of branches.

In Figure 3, we present the graph of the actual density f^* (in red) of the piecewise convex map τ with countable number of branches and graphs of the approximate stationary densities $f_{n,k}$ (in blue) via Ulam's method for maps τ_n with a finite number of branches. Numerical computations are performed for a number of cases. In the following table, we present the L^1 norm error $\|f^* - f_{n,k}\|_1$. Note that for each n , τ_n is a map with a finite number of branches which approximates the piecewise convex map τ in Figure 1 with countable number of branches.

n	k	$\ f^* - f_{n,k}\ _1$
5	100	0.2195470623
5	1000	0.2195243505
6	100	0.1943541673
6	1000	0.1943541673
7	1000	0.1742407352
10	1000	0.133493040
12	1000	0.11551923

The above table shows that as we increase n , the L^1 norm error $\|f^* - f_{n,k}\|_1$ gets smaller. For the fixed n , the increasing of k is not effective. For example, the




FIGURE 3. The graph of the actual invariant density f^* of the piecewise convex map τ with infinite number of branches (in red) and the graph of the approximating density $f_{n,k}$ (in blue): $n = 5, k = 1000$ on the left and $n = 10, k = 1000$ on the right hand side.

errors for $k = 100$ and $k = 1000$ are almost the same. The main method to lower the error is to increase the number of branches of τ_n .

By Theorem 5, L^1 norm error can be made arbitrarily small by making n and k large enough.

Example 2. Consider the piecewise convex map $\tau : [0, 1] \rightarrow [0, 1]$ with countable number of branches defined as

$$(4.2) \quad \tau(x) = \frac{1}{\frac{2i+1}{i(i+1)} - x} - i \quad \text{on} \quad \left[\frac{1}{i+1}, \frac{1}{i} \right], \quad i = 1, 2, \dots$$

See Figure 4. The map τ is defined on the countable partition of $[0, 1]$, with partition points $a_0 = 1, a_1 = \frac{1}{2}, a_2 = \frac{1}{3}, \dots, a_n = \frac{1}{n+1}, \dots$.




FIGURE 4. The graph of the piecewise convex map with countable number of branches for Example 2.

It is shown in [Góra et al.(2022)] that $\tau \in \mathcal{T}_{pc}^{\infty,0}(I)$ and hence by Theorem 2, τ has an ACIM. The actual density of τ is not known. The sequence $\{\tau_n\}_{n \geq 0}$ of piecewise convex maps $\tau_n : [0, 1] \rightarrow [0, 1]$ with finite number of branches, see Figure 5 for graphs of τ_n with $n = 8$ and $n = 10$, converges almost uniformly to τ , where

$$\tau_n(x) = \begin{cases} (n+1)x, & 0 \leq x < \frac{1}{n+1}; \\ \tau(x), & \frac{1}{n+1} \leq x \leq 1. \end{cases}$$

In Figure 6, we present the graph of the Ulam's approximation $f_{n,k}, n = 10, k = 1000$ of the actual invariant density of $f_n, n = 10$, of the piecewise convex map $\tau_n, n = 10$, with finite number of branches. By Theorem 5 it is also an approximation of the invariant density f^* of the map τ . The same Figure 6 shows also the approximation $f_{n,k}, n = 10, k = 500$, as the densities $f_{10,1000}$ and $f_{10,500}$ are indistinguishable at this scale. We have $\|f_{10,1000} - f_{10,500}\|_1 \sim 0.00055$. In Figure 7 we show the enlargement of both graphs on a small subinterval. The scales on x, y axes are different.




FIGURE 5. The graphs of the piecewise convex maps τ_n with finite number of branches $n = 8$ and $n = 10$.

Similarly as in Example 1, the effect of increasing k (after some threshold) is negligible. The improvement of the approximation is achieved by increasing the parameter n . The density $f_{11,1000}$ is indistinguishable from $f_{10,1000}$ at the paper illustrations scale with $\|f_{10,1000} - f_{11,1000}\|_1 \sim 0.00035$. The error $\|f_{11,1000} - f_{12,1000}\|_1 \sim 0.00029$.




FIGURE 6. The graph of the Ulam's approximation $f_{n,k,n} = 10, k = 1000$.

Acknowledgment: The authors are very grateful to anonymous reviewers for very detailed and useful comments and suggestions and also for stating a new problem related with their note. The research was supported by the NSERC Discovery Grant (DG) of the first author and second author. The third author is grateful to Concordia University and University of Prince Edward Island for financial support via NSERC DG grants of the first author and the second author respectively.

REFERENCES

- [Boltt & Santitissadeekorn(2013)] Boltt, E. M. & Santitissadeekorn, N. [2013], "Applied and computational measurable dynamics," *Math. Model. Comput.*, **18**, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA.
- [Bose et al.(2002)] Bose, C., Maume-Deschamps, V., Schmitt, B. & Shinand, S. S. [2002], "Invariant measures for piecewise convex transformations of an interval," *Studia Mathematica*, **152**, 263–297.
- [Bose & Murray(2001)] Bose, C. & Murray, R. [2001], "The exact rate of approximation in Ulam's method," *Discr. Contin. Dyn. Syst.* **7**, 219–235.
- [Boyarsky & Góra(1997)] Boyarsky, A. & Góra, P. [1997], *Laws of chaos. Invariant measures and dynamical systems in one dimension*, Probability and its Applications. Birkhäuser Boston, Inc., Boston, MA.
- [Boyarsky & Lou(1991)] Boyarsky, A. & Lou, Y.-S. [1991], "Approximating measures invariant under higher-dimensional chaotic transformations," *J. Approx. Theory* **65**, no. 2, 231–244.
- [Ding(1996)] Ding, J. [1996], "Computing Invariant Measures of Piecewise Convex Transformations," *Journal of Statistical Physics*, **83** (3 – 4), 623 – 635.
- [Ding & Zhou(1996)] Ding, J. & Zhou, A. [1996], "Finite approximations of Frobenius-Perron operators. A solution of Ulam's conjecture to multi-dimensional transformations," *Phys. D* **92**, no. 1-2, 61–68.




FIGURE 7. Enlargement of the approximating densities $f_{10,1000}$ (in red) and $f_{10,500}$ (in blue) on $[0, 0.02]$.

- [Froyland(1999)] Froyland, G. [1999], “Ulam’s method for random interval maps,” *Nonlinearity* **12**, 1029–1052.
- [Góra & Boyarsky(1989)] Góra, P. & Boyarsky, A. [1989], “Approximating the invariant densities of transformations with infinitely many pieces on the interval,” *Proc. Amer. Math. Soc.* **105**, no. 4, 922–928.
- [Góra et al.(2022)] Góra, P., Islam, M. S. & Rahman, AHM. M. [2022], “Absolutely continuous invariant measures for piecewise convex maps with countable number of branches,” *Dyn. of Contin. Disc. and Impul. Systems Series A* (under review, submitted on Nov. 10, 2022).
- [Hunt & Miller(1992)] Hunt, F. Y. & Miller, W. M. [1992], “On the approximation of invariant measures,” *J. Statist. Phys.* **66**, no.1–2, 535–548.
- [Inoue(1992)] Inoue, T. [1992], “Asymptotic stability of densities for piecewise convex maps,” *Ann. Polon. Math.* **57**, no. 1, 83–90.
- [Inoue(2023)] Inoue, T. [2023], “Invariant measures for random piecewise convex maps,” <https://arxiv.org/abs/2303.10873>.
- [Jabłoński & Malczak(1984)] Jabłoński, M. & Malczak, J. [1984], “A central limit theorem for piecewise convex transformations of the unit interval,” *Colloq. Math.* **49**, no. 1, 59–65.
- [Jabłoński & Malczak(1983)] Jabłoński, M. & Malczak, J. [1983], “The rate of the convergence of iterates of the Frobenius-Perron operator for piecewise convex transformations,” *Bull. Polish Acad. Sci. Math.* **31**, no. 5-8, 249–254.
- [Lasota & Yorke(1973)] Lasota, A. & Yorke, J.A. [1973], “On the existence of invariant measures for piecewise monotonic transformations,” *Trans. Amer. Math. Soc.* **186** 481 – 488.
- [Lasota & Mackey(1994)] Lasota A. & Mackey, M. C. [1994], *Chaos, fractals, and noise, Stochastic aspects of dynamics*, Applied Mathematical Sciences 97, Springer-Verlag, New York.
- [Lasota & Yorke(1982)] Lasota, A. & Yorke, J. A. [1982], “Exact dynamical systems and the Frobenius-Perron operator,” *Trans. Amer. Math. Soc.* **273** (1), pp. 375–384.
- [Li(1976)] Li, T-Y. [1976], “Finite approximation for the Frobenius-Perron operator: A solution to Ulam’s conjecture,” *J. Approx. Theory.* **17**, 177 – 186.

- [Miller(1994)] Miller, W. M. [1994], “Stability and approximation of invariant measures for a class of non-expanding transformations,” *Nonlinear Anal.* **23** (8), 1013 – 1025.
- [Rahman(2019)] Rahman, M. H. [2019], “Absolutely Continuous Invariant Measures for Piecewise Convex Maps of Interval with Infinite Number of Branches,” Master’s Thesis, Concordia University, Canada.
- [Rudin(1976)] Rudin, Walter [1976], *Principles of mathematical analysis*, Third edition, Internat. Ser. Pure Appl. Math., McGraw-Hill Book Co., New York-Auckland-Düsseldorf.
- [Ulam(1960)] Ulam, S. M. [1960], “A collection of Mathematical problems Interscience Tracts in pure and Applied Math.,” **8**, Interscience, New York, USA.
- [Yosida & Kakutani(1941)] Yosida, K. & Kakutani, S. [1941], “Operator-Theoretical Treatment of Markoff’s Process and Mean Ergodic Theorem,” *Annals of Mathematics*, Second Series, Vol. **42**, No. 1 , pp. 188–228.

(Md Shafiqul Islam) SCHOOL OF MATHEMATICAL AND COMPUTATIONAL SCIENCES, UNIVERSITY OF PRINCE EDWARD ISLAND, 550 UNIVERSITY AVE, CHARLOTTETOWN, PE, C1A 4P3, CANADA
Email address, Md Shafiqul Islam: sislam@upei.ca

(P. Góra) DEPARTMENT OF MATHEMATICS AND STATISTICS, CONCORDIA UNIVERSITY, 1455 DE MAISONNEUVE BLVD. WEST, MONTREAL, QUEBEC H3G 1M8, CANADA
Email address, P. Góra: pawel.gora@concordia.ca

(A H M Mahbubur Rahman) DEPARTMENT OF MATHEMATICS AND STATISTICS,, CONCORDIA UNIVERSITY, 1455 DE MAISONNEUVE BLVD. W., MONTREAL, QUEBEC, H3G 1M8, CANADA
Email address, A H M Mahbubur Rahman: mahbubur.rahman@concordia.ca