

CANONICAL DATA-RECONSTRUCTIONS VIA KERNELS, HILBERT SPACE-VALUED GAUSSIAN PROCESSES, AND QUANTUM STATES

PALLE E.T. JORGENSEN AND JAMES TIAN

ABSTRACT. We offer new results and new directions in the study of operator-valued kernels and their factorizations. Our approach provides both more explicit realizations and new results, as well as new applications. These include: (i) an explicit covariance analysis for Hilbert space-valued Gaussian processes, (ii) optimization results for quantum gates (from quantum information), (iii) new results for positive operator-valued measures (POVMs), and (iv) a new approach/result in inverse problems for quantum measurements.

CONTENTS

1. Introduction	1
2. Factorization of $\mathcal{L}(H)$ -valued p.d. kernels	2
3. Relation to classical dilation theory	4
4. H -valued Gaussian processes	6
5. Completeness, POVMs, quantum states, and quantum information	8
5.1. An inverse problem	10
References	12

1. INTRODUCTION

We present a new approach to inverse problems, combining classical and quantum. Applications include new covariance analyses for Hilbert space-valued Gaussian processes, and to a variety of dilation constructions. We further present new results for optimization with quantum gates; a new approach/result in inverse problems for quantum measurements, and to inference with positive operator-valued measures (POVMs).

With our starting point being that of operator-valued positive definite kernels, our present paper then proceeds to offer a new two-fold framework (i) and (ii) for the following problems: one is that of Hilbert space and representations, and the other, new classes of Gaussian processes. In more detail: We present: (i) a new approach to the interplay between a variety of systems which allow for realizations in specific and concrete Hilbert spaces H , on the one hand, and which at the same time also admit extensions, typically called dilations, in “larger” Hilbert spaces, say L . (By “dilation,” we mean that the two systems are connected via an isometry

2020 *Mathematics Subject Classification.* 46E22, 46N50, 47A20, 47B32, 47N50, 60G15, 81P15.

Key words and phrases. Positive definite functions, kernels, Gaussian processes, covariance, dilation, POVMs, PVMs, measurement, quantum states.

V from H to L , which intertwines the respective systems/representations.) And for (ii), we make use of these constructions in our analysis of Hilbert space-valued Gaussian processes. The constructions we present in (i) are more explicit than those in earlier related analyses, such as Gelfand Naimark Segal (GNS), from states to representations; Stinespring, from completely positive maps to representations; or Halmos, from powers contraction to unitary power dilations; and finally from positive operator-valued measures (POVMs) to projection valued measures. See, e.g., [DM20, PTD23, Ver22, BJ97, BJ99, BJKW00, DJ12, DJ14, DHJ15, DJ15]. Our approach serves multiple purposes: one is to unify theories, and the other is to make these constructions explicit.

Our new results include (i) an explicit covariance analysis for Hilbert space-valued Gaussian processes (Section 4), (ii) optimization results for quantum gates (Section 5), (iii) new results for positive operator-valued measures (POVMs) (Sections 2–5), and (iv) a new approach/result in inverse problems for quantum measurements (Section 5.1).

Notation. Throughout the paper, we use the physics convention that inner products are linear in the second variable. Let $|a\rangle\langle b|$ denote Dirac’s rank-1 operator, $c \mapsto a \langle b, c \rangle$. $\mathcal{L}(H)$ denotes the algebra of all bounded linear operators in a Hilbert space H .

For a positive definite (p.d.) kernel $K : S \times S \rightarrow \mathbb{C}$, let H_K be the corresponding reproducing kernel Hilbert space (RKHS). H_K is the Hilbert completion of

$$\text{span}_{\mathbb{C}} \{K_y(\cdot) := K(\cdot, y) \mid y \in S\} \quad (1.1)$$

with respect to the inner product

$$\left\langle \sum_i c_i K(\cdot, x_i), \sum_j d_j K(\cdot, y_j) \right\rangle_{H_K} := \sum_i \sum_j \bar{c}_i d_j K(x_i, y_j). \quad (1.2)$$

The following reproducing property holds:

$$\varphi(x) = \langle K(\cdot, x), \varphi \rangle_{H_K}, \forall x \in S, \forall \varphi \in H_K. \quad (1.3)$$

Any scalar-valued kernel K as above is associated with a zero-mean Gaussian process, where K is the covariance:

$$K(s, t) = \mathbb{E} [\overline{W_s} W_t], \quad (1.4)$$

and $W_s \sim N(0, K(s, s))$.

An $\mathcal{L}(H)$ -valued kernel $K : S \times S \rightarrow \mathcal{L}(H)$ is p.d. if

$$\sum_{i=1}^n \langle a_i, K(s_i, s_j) a_j \rangle_H \geq 0 \quad (1.5)$$

for all $(a_i)_1^n$ in H , and all $n \in \mathbb{N}$.

For the theory of reproducing kernels, the literature is extensive, both pure and recent applications, we refer to the following, including current, [AS56, Aro48, AJ21, PR16, AFMP94, LYA23, CY17, Alp91, AD93, AL95, AB97, AD06, AJ15].

2. FACTORIZATION OF $\mathcal{L}(H)$ -VALUED P.D. KERNELS

In this section, we present our universal construction in a general framework. This will then, in adapted forms, go into a set of diverse applications. The latter are addressed in subsequent sections.

In our analysis of kernels and factorizations we want to highlight the following two distinctions: First is the distinction between the traditional results which only yield existence of factorizations involving abstract and unstructured completions, as opposed to arising as a universal and concrete solution, realized directly from the given operator valued kernel as an explicitly given space of functions. The second is the distinction between universal, as opposed to case-by-case analyses. In both senses, our results below fall in the category of universal factorizations.

Theorem 2.1 (Universal Factorization). *Let $K : S \times S \rightarrow \mathcal{L}(H)$ be a p.d. kernel. Then there exists a RKHS $H_{\tilde{K}}$, and a family of operators $V_s : H \rightarrow H_{\tilde{K}}$, $s \in S$, such that*

$$H_{\tilde{K}} = \overline{\text{span}} \{V_s a : a \in H, s \in S\} \quad (2.1)$$

and

$$K(s, t) = V_s^* V_t. \quad (2.2)$$

Conversely, if there is a Hilbert space L and operators $V_s : H \rightarrow L$, $s \in S$, such that

$$L = \overline{\text{span}} \{V_s a : a \in H, s \in S\} \quad (2.3)$$

and (2.2) holds, then $L \simeq H_{\tilde{K}}$.

Proof. First, assume that K is of the form (2.2). For all $(a_i)_{i=1}^n$ in H , and $(s_i)_{i=1}^n$ in S ,

$$\begin{aligned} \sum_{i,j} \langle a_i, K(s_i, s_j) a_j \rangle_H &= \sum_{i,j} \langle a_i, V_{s_i}^* V_{s_j} a_j \rangle_H \\ &= \sum_{i,j} \langle V_{s_i} a_i, V_{s_j} a_j \rangle_L = \left\| \sum_i V_{s_i} a_i \right\|_L^2 \geq 0. \end{aligned}$$

Thus, K is p.d.

Now, given a p.d. kernel $K : S \times S \rightarrow \mathcal{L}(H)$, let

$$\tilde{K} : (S \times H) \times (S \times H) \rightarrow \mathbb{C}$$

be the scalar-valued kernel defined by

$$\tilde{K}((s, a), (t, b)) := \langle a, K(s, t) b \rangle_H.$$

Then $\tilde{K}$ is p.d. and let $H_{\tilde{K}}$ be the associated RKHS.

Set $V_s : H \rightarrow H_{\tilde{K}}$ by

$$V_s a = \tilde{K}(\cdot, (s, a)) = \tilde{K}(\cdot, (s, a)) : (S \times H) \rightarrow \mathbb{C}, \quad \forall a \in H.$$

Since

$$\left\langle V_s a, \tilde{K}(\cdot, (t, b)) \right\rangle_{H_{\tilde{K}}} = \langle a, K(s, t) b \rangle_H, \quad \forall a, b \in H, \forall s, t \in S,$$

it follows that

$$V_s^* \tilde{K}(\cdot, (t, b)) = K(s, t) b.$$

One concludes that

$$K(s, t) = V_s^* V_t.$$

Finally, suppose there is a Hilbert space L and family of operators $V_s : H \rightarrow L$, $s \in S$, such that (2.2)–(2.3) hold. Then the mapping

$$\tilde{K}(\cdot, (s, a)) \mapsto V_s a$$

extends (by linearity and density) to a unique isometric isomorphism from $H_{\tilde{K}}$ onto L , and so any such space L is isomorphic to the RKHS $H_{\tilde{K}}$. $\square$

Corollary 2.2. *Any function $F \in H_{\tilde{K}}$ satisfies that*

$$F(s, a) = \langle a, V_s^* F \rangle_{H_{\tilde{K}}}, \quad (s, a) \in S \times H. \quad (2.4)$$

In particular, F is conjugate linear in the second variable.

Proof. This follows from the reproducing property (see (1.3)) of $H_{\tilde{K}}$:

$$F(s, a) = \left\langle \tilde{K}(\cdot, (s, a)), F \right\rangle_{H_{\tilde{K}}} = \langle V_s a, F \rangle_{H_{\tilde{K}}} = \langle a, V_s^* F \rangle_H.$$

$\square$

Corollary 2.3. *Let (φ_i) be any orthonormal basis (ONB) in $H_{\tilde{K}}$. The following operator identity holds:*

$$K(s, t) = \sum_i |V_s^* \varphi_i\rangle \langle V_t^* \varphi_i|. \quad (2.5)$$

Proof. Using the identity in $H_{\tilde{K}}$,

$$I_{H_{\tilde{K}}} = \sum_i |\varphi_i\rangle \langle \varphi_i|,$$

we get that

$$K(s, t) = V_s^* V_t = V_s^* I_{H_{\tilde{K}}} V_t = \sum_i |V_s^* \varphi_i\rangle \langle V_t^* \varphi_i|.$$

$\square$

3. RELATION TO CLASSICAL DILATION THEORY

The notion of dilation theory dates back to the fundamental Hilbert space axioms, including to the study of such basic notions in operator theory as contractive operators in Hilbert space H , of states defined as positive linear functionals (on algebras of operators), and of the notion and properties of Stinespring-completely positive operator-valued maps. While the early advances were motivated primarily by pure operator theory, and by representation theory, recent studies have served to link much more directly to modern quantum information, see also Section 5 below. The cast of characters in the general context was (and is) perhaps focused on such areas of dilation theory as: dilations for (i) subnormal operators, (ii) for operator valued measures, (iii) for contraction operators, (iv) for operator spaces, and their role as extensions in dilation theory, and (v) for commuting sets of operators, and semigroups of completely positive maps. Early applications include systems theory, prediction theory, and more recently quantum information. The literature is vast, and here we call attention to [Arv10], and the papers cited there. We now turn to the role our above results (Section 2) play as tools in both classical and modern dilation theory; as well as in sharpening the conclusions, and in presenting new directions.

Indeed, the setting in Theorem 2.1 is quite general. It includes and extends well known constructions in classical dilation theory. Notably, passing from K to a scalar-valued p.d. kernel $\tilde{K}$ offers a function-based approach to dilation theory, in contrast with traditional abstract spaces of equivalence classes. It also allows for direct evaluation and manipulation of functions, due to the RKHS structure of the

dilation space, as well as the associated H -valued Gaussian processes (Section 4), making it especially relevant for practical applications.

For instance, consider a completely positive map φ from a C^* -algebra $\mathfrak{A}$ to the algebra of bounded operators on a Hilbert space H . The Stinespring Dilation Theorem [Sti55] states that there exists a Hilbert space L (potentially larger than H), a representation $\pi : \mathfrak{A} \rightarrow \mathcal{L}(L)$, and a bounded operator $V : H \rightarrow L$, such that

$$\varphi(A) = V^* \pi(A) V, \quad A \in \mathfrak{A}. \quad (3.1)$$

Moreover, if φ is unital, then V is isometric and H can be identified as a subspace of L . The space L is minimal if the span of $\{\pi(A) V a : a \in H, A \in \mathcal{L}(H)\}$ is dense in L , in which case L is unique up to unitary equivalence.

A key step in the proof of Stinespring's theorem is to introduce a sesquilinear form on the algebraic tensor product $\mathfrak{A} \otimes H$,

$$\left\langle \sum_i A_i \otimes a_i, \sum_j B_j \otimes b_j \right\rangle_L := \sum_{i,j} \langle a_i, \varphi(A^* B) b_j \rangle_H,$$

and then setting L to be the corresponding Hilbert completion, modulo elements with zero norm, i.e., elements such that $\|\sum A_i \otimes a_i\|_L = 0$. The space L is then an abstract Hilbert space of equivalence classes.

To compare with Theorem 2.1, one may consider the $\mathcal{L}(H)$ -valued p.d. kernel

$$K(A, B) = \varphi(A^* B)$$

defined on $\mathfrak{A} \times \mathfrak{A}$, and define the scalar-valued p.d. kernel $\tilde{K} : (\mathfrak{A} \times H) \times (\mathfrak{A} \times H) \rightarrow \mathbb{C}$ by

$$\tilde{K}((A, a), (B, b)) = \langle a, \varphi(A^* B) b \rangle_H.$$

Corollary 3.1. *The space L is then the RKHS $H_{\tilde{K}}$, consisting of functions F on $\mathfrak{A} \times H$, with the reproducing property:*

$$F(A, a) = \left\langle \tilde{K}(\cdot, (A, a)), F \right\rangle_{H_{\tilde{K}}}.$$

Note that $H_{\tilde{K}}$ is a function space on $\mathfrak{A} \times H$, as opposed to an abstract space of equivalence classes.

The operator V , and the representation π , are also explicit:

Lemma 3.2. *Set $V : H \rightarrow H_{\tilde{K}}$ by*

$$V a = \tilde{K}(\cdot, (I, a)).$$

The adjoint $V^ : H_{\tilde{K}} \rightarrow H$ is given by*

$$V^* \tilde{K}(\cdot, (B, b)) = \varphi(B) b.$$

Define $\pi : \mathfrak{A} \rightarrow \mathcal{L}(H_{\tilde{K}})$ as

$$\pi(A) \tilde{K}(\cdot, (B, b)) = \tilde{K}(\cdot, (AB, b)).$$

Then (3.1) holds. Indeed, since, by definition,

$$\text{span} \{K(\cdot, (A, a)) : A \in \mathfrak{A}, a \in H\}$$

is dense in $H_{\tilde{K}}$, $(\mathfrak{A}, \pi, V, H_{\tilde{K}})$ forms a minimal Stinespring dilation of φ .

Proof. We verify the adjoint V^* is as stated. Since the span of

$$\left\{ \tilde{K}(\cdot, (A, a)) : A \in \mathfrak{A}, a \in H \right\}$$

is dense in $H_{\tilde{K}}$, it is enough to check that

$$\begin{aligned} \left\langle Va, \tilde{K}(\cdot, (B, b)) \right\rangle_{H_{\tilde{K}}} &= \left\langle \tilde{K}(\cdot, (I, a)), \tilde{K}(\cdot, (B, b)) \right\rangle_{H_{\tilde{K}}} \\ &= \langle a, \varphi(B)b \rangle_H \end{aligned}$$

which holds for all $a, b \in H$, and all $B \in \mathcal{L}(H)$. This means $V^* \tilde{K}(\cdot, (B, b)) = \varphi(B)b$, for all $B \in \mathfrak{A}$, and all $b \in H$.

Then,

$$\begin{aligned} V^* \pi(A) Va &= V^* \pi(A) \tilde{K}(\cdot, (I, a)) \\ &= V^* \tilde{K}(\cdot, (A, a)) = \varphi(A)a \end{aligned}$$

which is (3.1).

Additionally, if φ is unital, then

$$\begin{aligned} \|Va\|_{H_{\tilde{K}}}^2 &= \left\langle \tilde{K}(\cdot, (I, a)), \tilde{K}(\cdot, (I, a)) \right\rangle_{H_{\tilde{K}}} \\ &= \langle a, K(I, I)a \rangle_H = \langle a, \varphi(I)a \rangle_H = \langle a, a \rangle_H = \|a\|_H^2, \end{aligned}$$

so that V is isometric. $\square$

4. H -VALUED GAUSSIAN PROCESSES

Since the foundational work of Ito, the stochastic analysis of Gaussian processes has relied heavily on Hilbert space and associated operators. For reference, see works such as [Sch52, AJ15, AJ21] along with the early literature cited therein. In this discussion, we highlight two distinct scenarios: (i) scalar-valued Gaussian processes indexed by a Hilbert space, often through an Ito-isometry; and (ii) Gaussian processes that take values in a Hilbert space themselves, see e.g., [Lop23, Gue22, KM20, FS20, Kat19, JT23]. Our current focus is on the latter. This setting offers the advantage of enabling more flexible representations of covariance structures, which are increasingly relevant in the analysis of large data sets. We now proceed with the construction details.

Let S be a set, and H a Hilbert space. We say $\{W_s\}_{s \in S}$ is an H -valued Gaussian process if, for all $a, b \in H$, and all $s, t \in S$,

$$\mathbb{E}[\langle a, W_s \rangle_H \langle W_t, b \rangle_H] = C_{a,b}(s, t), \quad (4.1)$$

where

$$\langle W_s, a \rangle_H \sim N(0, C_{a,a}(s, s)), \quad (4.2)$$

i.e., mean zero Gaussian, with variance $C_{a,a}(s, s)$.

Theorem 4.1. *Every operator-valued p.d. kernel $K : S \times S \rightarrow \mathcal{L}(H)$ is associated with an H -valued Gaussian process $\{W_s\}_{s \in S}$, such that*

$$\mathbb{E}[\langle a, W_s \rangle_H \langle W_t, b \rangle_H] = \langle a, K(s, t)b \rangle_H.$$

for all $s, t \in S$, and all $a, b \in H$.

Conversely, any H -valued Gaussian process $\{W_s\}_{s \in S}$ is obtained from such a p.d. kernel.

Proof. Suppose $\{W_s\}_{s \in S}$ is an H -valued (mean zero) Gaussian process realized in a probability space $(\Omega, \mathbb{P})$, with expectation $\mathbb{E}(\cdot) = \int_{\Omega} (\cdot) d\mathbb{P}$.

Fix $s, t \in S$. For all $a, b \in H$,

$$(a, b) \longmapsto \mathbb{E} [\langle a, W_s \rangle_H \langle W_t, b \rangle_H]$$

is a bounded sesquilinear form on H , and so

$$\mathbb{E} [\langle a, W_s \rangle_H \langle W_t, b \rangle_H] = \langle a, K(s, t) b \rangle_H$$

for some operator $K(s, t) \in \mathcal{L}(H)$. Also,

$$K : S \times S \rightarrow \mathcal{L}(H)$$

is positive definite, since

$$\begin{aligned} \sum_{i,j=1}^n \langle a_i, K(s_i, s_j) a_j \rangle_H &= \sum_{i,j=1}^n \mathbb{E} [\langle a_i, W_{s_i} \rangle_H \langle W_{s_j}, a_j \rangle_H] \\ &= \mathbb{E} \left[\left| \sum_{i=1}^n \langle W_{s_i}, a_i \rangle_H \right|^2 \right] \geq 0 \end{aligned}$$

for all $a_1, \dots, a_n \in H$ and $s_1, \dots, s_n \in S$.

Conversely, given $K : S \times S \rightarrow \mathcal{L}(H)$ p.d., introduce the scalar-valued p.d. kernel $\tilde{K} : (S \times H) \times (S \times H) \rightarrow \mathbb{C}$,

$$\tilde{K}((s, a), (t, b)) = \langle a, K(s, t) b \rangle_H.$$

Let $H_{\tilde{K}}$ be the RKHS of $\tilde{K}$, and let

$$\{W_{(s,a)} : (s, a) \in S \times H\}$$

be the associated Gaussian process (see (1.4), and also Theorem 4.3 for a realization). That is,

$$\tilde{K}((s, a), (t, b)) = \langle a, K(s, t) b \rangle_H = \mathbb{E} [\overline{W_{(s,a)}} W_{(t,b)}].$$

Let $V_s : H \rightarrow H_{\tilde{K}}$ be given by $V_s a = \tilde{K}(\cdot, (s, a))$. Using the isometry

$$H_{\tilde{K}} \ni V_s a = \tilde{K}(\cdot, (s, a)) \longrightarrow W_{(s,a)} \in L^2(\mathbb{P})$$

it follows that $H \ni a \rightarrow W_{(s,a)}$ is linear. The assertion follows. $\square$

Definition 4.2. Consider the H -valued Gaussian process $W_s : \Omega \rightarrow H$,

$$W_t = \sum_i (V_t^* \varphi_i) Z_i, \quad (4.3)$$

where (φ_i) is an ONB in $H_{\tilde{K}}$. Following standard conventions, here $\{Z_i\}$ refers to a choice of an independent identically distributed (i.i.d.) system of standard scalar Gaussians $N(0, 1)$ random variables, and with an index matching the choice of ONB.

Theorem 4.3. *We have*

$$\mathbb{E} [\langle a, W_s \rangle_H \langle W_t, b \rangle_H] = \langle a, K(s, t) b \rangle_H. \quad (4.4)$$

Proof. Note that $\mathbb{E}[Z_i Z_j] = \delta_{i,j}$. From this, we get

$$\begin{aligned}
\text{LHS}_{(4.4)} &= \mathbb{E}[\langle a, W_s \rangle_H \langle W_t, b \rangle_H] \\
&= \sum_i \sum_j \langle a, V_s^* \varphi_i \rangle \langle V_t^* \varphi_j, b \rangle \mathbb{E}[Z_i Z_j] \\
&= \sum_i \langle V_s a, \varphi_i \rangle_{H_{\bar{K}}} \langle \varphi_i, V_t b \rangle_{H_{\bar{K}}} \\
&= \langle V_s a, V_t b \rangle_{H_{\bar{K}}} = \langle a, V_s^* V_t b \rangle_H = \langle a, K(s, t) b \rangle_H.
\end{aligned}$$

Also see (2.5). $\square$

Remark 4.4. The Gaussian process $\{W_t\}_{t \in S}$ in (4.3) is well defined and possesses the stated properties. This is an application of the central limit theorem to the choice $\{Z_i\}$ of i.i.d. $N(0, 1)$ Gaussians on the right-side of (4.3).

Varying the choices of ONBs (φ_i) and i.i.d. $N(0, 1)$ Gaussian random variables (Z_i) will result in different Gaussian processes $\{W_t\}_{t \in S}$, but all will adhere to the covariance condition specified in (4.4). It is also important to note that if (φ_i) , (Z_i) are fixed, the resulting Gaussian process $\{W_t\}_{t \in S}$ is uniquely determined by its first two moments.

5. COMPLETENESS, POVMS, QUANTUM STATES, AND QUANTUM INFORMATION

In this section we apply our construction to the case of positive operator-valued measures (POVMs). While POVMs have long played an important role in operator theory, they have recently found applications in quantum information [ANW17, EHD⁺13, CDP⁺13, KLSW12, KSW08, KW01, Wer01]. In more detail: For a given Hilbert space H , and a measure space Ω , a positive operator-valued measure (POVM) on Ω is a sigma-additive measure whose values are positive semi-definite operators on H . Thus the notion of POVM is a generalization of projection-valued measures (PVMs). In quantum information, the latter are called projection-measurements. (Here by a projection, we mean selfadjoint projection P in H .) Note that therefore quantum measurements realized as POVMs are generalizations of quantum measurement described by PVMs. By analogy, a POVM is to a PVM what a mixed state is to a pure state. Recall that in quantum theory, mixed states specify those quantum states occurring in subsystems of larger systems. Hence the study of purification of quantum states. The POVMs describe the effect on a subsystem of a projection measurement performed on a larger system. In other words, POVMs are the most general kind of measurement in quantum mechanics, and they in the field of quantum information. Here quantum information refers the information of states of quantum systems, thus leading to a variety of quantum information processing techniques. The literature is extensive. For current references we here mention only a sample, [LK24, QY23, DKM24, HKLP24, WNGB24, FLZG24, Hol22, RMC21, HCW21] and the papers cited there. Quantum information refers to both the technical definition of von Neumann entropy as well as to the general computational devices. Recently, quantum computing has become especially active research area, serving as a key part of modern computation and communication.

A main difference between classical probability, and its quantum counterpart is played by the role of entangled quantum states, see e.g., [LKKK24, QLY⁺24, FMBS24]. The latter in turn have their origin with the pioneers. But in the past

decade, they have come to play a crucial role in modern quantum computing. (We make a note on their origin: Erwin Schrödinger, in studying the Einstein-Podolsky-Rosen (EPR) state of two particles influencing each other instantly at a distance, introduced the concept of entangled (quantum) states. He used the German term “verschränkung”, which was subsequently translated into English as entanglement.)

In quantum information theory, a measurement channel is a POVM P , defined on a measurable space $(\Omega, \mathcal{B}_\Omega)$, acting in a Hilbert space H . The set of quantum states (density operators) $\mathcal{S}_1^1(H)$ consists of positive trace class operators ρ with $\text{tr}(\rho) = 1$. A measurement channel P is said to be information-complete if

$$\rho \mapsto \text{tr}(\rho P(\cdot))$$

from $\mathcal{S}_1^1(H)$ to the set of Borel probability measures $\mathcal{M}_1(\Omega)$ on Ω , is injective. In other words, measurements uniquely determine states. In this context, an $\mathcal{L}(H)$ -valued p.d. kernel on $S \times S$ may be naturally viewed as a generalized quantum measurement channel.

In what follows, let $\mathcal{K}(S, \mathcal{L}(H))$ be the set of all $\mathcal{L}(H)$ -valued p.d. kernels on $S \times S$. In particular, $\mathcal{K}(S, \mathbb{C})$ is the set of all scalar-valued p.d. kernels. $\mathcal{S}_p(H)$ denotes the p th Schatten-class operators, with norm

$$\|T\|_p = \left(\sum_n s_n(T)^p \right)^{1/p} < \infty,$$

where $s_n(T)$ are the singular numbers of T . Here, $\mathcal{S}_1(H)$ is the trace class, and $\mathcal{S}_2(H)$ the class of Hilbert-Schmidt operators. $\mathcal{S}_1^+(H)$ is the cone of positive trace class operators, and $\mathcal{S}_1^1(H) = \{T \in \mathcal{S}_1^+(H) : \text{tr}(T) = 1\}$ consists of quantum states.

Lemma 5.1. *Fix $K \in \mathcal{K}(S, \mathcal{L}(H))$. For all $\rho \in \mathcal{S}_1^+(H)$, the kernel $c : S \times S \rightarrow \mathbb{C}$,*

$$c(s, t) := \text{tr}(\rho K(s, t))$$

is positive definite, that is, $c \in \mathcal{K}(S, \mathbb{C})$.

Proof. Write

$$\rho = \sum \lambda_i |\varphi_i\rangle\langle\varphi_i|, \quad \lambda_i > 0,$$

where (φ_i) is an ONB in H . Then,

$$(s, t) \mapsto \text{tr}(\rho K(s, t)) = \sum \lambda_i \langle \varphi_i, K(s, t) \varphi_i \rangle_H$$

is a sum of scalar valued p.d. kernels, and so it is in $\mathcal{K}(S, \mathbb{C})$. $\square$

Theorem 5.2. *Fix $K \in \mathcal{K}(S, \mathcal{L}(H))$. Consider the map $\Psi : \mathcal{S}_1^+(H) \rightarrow \mathcal{K}(S, \mathbb{C})$, where*

$$\Psi(\rho)(s, t) = \text{tr}(\rho K(s, t)), \quad (s, t) \in S \times S.$$

Then Ψ is 1-1 if and only if the double commutant $\{K(s, t)\}''$ is weak dense in $\mathcal{L}(H)$.*

Proof. This follows from von Neumann’s double commutant theorem. Recall that $\mathcal{S}_1^+(H)$ is the space of normal states, and it is the predual of $\mathcal{L}(H)$. Therefore, the injectivity of Ψ in question is equivalent to the von Neumann algebra generated by $\{K(s, t) : s, t \in S\}$ being weak* dense in $\mathcal{L}(H)$. $\square$

Definition 5.3. An $\mathcal{L}(H)$ -valued p.d. kernel K is said to be complete if the map

$$\mathcal{S}_1^+(H) \ni \rho \mapsto \text{tr}(\rho K(\cdot, \cdot)) \in \mathcal{K}(S, \mathbb{C})$$

is 1-1. That is, the scalar-valued kernel $(s, t) \mapsto \text{tr}(\rho K(s, t))$ on $S \times S$ uniquely determines $K \in \mathcal{K}(S, \mathcal{L}(H))$.

Corollary 5.4. Fix $K \in \mathcal{K}(S, \mathcal{L}(H))$. Let $H_{\tilde{K}}$ be the RKHS of the corresponding scalar-valued kernel $\tilde{K}$, defined on $(S \times H) \times (S \times H)$. Let $\{W_t(\cdot)\}_{t \in S}$ be the associated H -valued Gaussian process. Assume that K is complete. Then the system of vectors

$$\left\{ V_t a = \tilde{K}(\cdot, (t, a)) \right\}_{t \in S} \subset H_{\tilde{K}}$$

determines each $a \in H$ uniquely. Similarly, the scalar valued Gaussian process

$$\{\langle W_t, a \rangle_H\}_{t \in S} \subset L^2(\Omega, \mathbb{P})$$

determines each $a \in H$ uniquely.

Proof. Let $a \in H$. Apply Theorem 5.2 to $\rho = |a\rangle\langle a| \in \mathcal{S}_1^+(H)$, and use the fact that

$$\langle V_s a, V_t a \rangle_{H_{\tilde{K}}} = \mathbb{E}[\langle a, W_s \rangle_H \langle W_t, a \rangle_H] = \langle a, K(s, t) a \rangle_H = \text{tr}(\rho K(s, t)).$$

□

5.1. An inverse problem. In the theory of quantum measurement, a pertinent question is how to recover the state of a system given measurement data. More precisely, for generalized channels, one may consider the following:

Question 5.5. Given a generalized channel $K \in \mathcal{K}(S, \mathcal{L}(H))$, and measurement data $c(s, t) = \text{tr}(\rho K(s, t))$, how to recover the quantum state $\rho \in \mathcal{S}_1^1(H)$?

Alternatively, given K , $(s_i)_{i=1}^N \subset S$ and $(c_{i,j})_{i,j=1}^N \subset \mathbb{C}$, consider the following optimization problem:

$$\min \left\{ \sum_{i,j=1}^N |\text{tr}(\rho K(s_i, s_j)) - c_{i,j}|^2 : \rho \in \mathcal{S}_1^1(H) \right\}. \quad (5.1)$$

Since every state $\rho \in \mathcal{S}_1^+(H)$ is a product of Hilbert-Schmidt operators ($\rho = (\sqrt{\rho})^2$), we will first extend $\mathcal{L}(H)$ -valued p.d. kernels to the case where $H = \mathcal{S}_2$.

In fact, every $\mathcal{L}(H)$ -valued p.d. kernel is also an $\mathcal{L}(\mathcal{S}_2)$ -valued kernel. Here, $\mathcal{S}_2 = \mathcal{S}_2(H)$ is the class of Hilbert-Schmidt operators in H , which is a Hilbert space with inner product $\langle A, B \rangle_2 = \text{tr}(A^* B)$. More precisely, one has:

Proposition 5.6. $\mathcal{K}(S, \mathcal{L}(H)) \subset \mathcal{K}(S, \mathcal{L}(\mathcal{S}_2))$.

Proof. Let $K \in \mathcal{K}(S, \mathcal{L}(H))$. Note that $K \in \mathcal{K}(S, \mathcal{L}(\mathcal{S}_2))$ if and only if

$$\sum_{i,j} \langle A_i, K(s_i, s_j) A_j \rangle_2 = \sum_{i,j} \text{tr}(A_i^* K(s_i, s_j) A_j) \geq 0, \quad \forall (A_i)_{i=1}^N \subset \mathcal{S}_2.$$

It suffices to verify that

$$\sum_{i,j} \langle \varphi, A_i^* K(s_i, s_j) A_j \varphi \rangle_H \geq 0, \quad \forall \varphi \in H.$$

This follows from the p.d. assumption of K , since

$$\begin{aligned} \sum_{i,j} \langle \varphi, A_i^* K(s_i, s_j) A_j \varphi \rangle_H &= \sum_{i,j} \langle A_i \varphi, K(s_i, s_j) A_j \varphi \rangle_H \\ &= \sum_{i,j} \langle h_i, K(s_i, s_j) h_j \rangle_H \geq 0, \quad h_i := A_i \varphi. \end{aligned}$$

□

Remark. Thus, there is an associated scalar-valued p.d. kernel $\tilde{K} : (S \times \mathcal{S}_2) \times (S \times \mathcal{S}_2) \rightarrow \mathbb{C}$, with

$$\tilde{K}((s, A), (t, B)) = \langle A, K(s, t) B \rangle_2.$$

And, an $\mathcal{S}_2$ -valued Gaussian process W satisfying that

$$\mathbb{E}[\langle A, W_s \rangle_2 \langle W_t, B \rangle_2] = \langle A, K(s, t) B \rangle_2$$

for all $s, t \in S$, and $A, B \in \mathcal{S}_2$.

Below, we analyze a simple case when $\dim H < \infty$, which reduces (5.1) to a constraint kernel approximation (least-squares) problem.

Assume H is finite-dimensional. Let

$$K(s, t) = Q(s \cap t), \quad s, t \in \mathcal{B}_\Omega, \quad (5.2)$$

where $Q : (\Omega, \mathcal{B}_\Omega) \rightarrow \mathcal{L}(H)$ is a POVM over a measurable space $(\Omega, \mathcal{B}_\Omega)$. Recall that $Q(\Omega) = I_H$. Introduce $\tilde{K}$ and its RKHS $H_{\tilde{K}}$ as follows:

$$\begin{aligned} \tilde{K}((s, A), (t, B)) &= \left\langle \tilde{K}_{(s,A)}, \tilde{K}_{(t,B)} \right\rangle_{H_{\tilde{K}}} \\ &= \langle A, Q(s \cap t) B \rangle_2 = \text{tr}(A^* Q(s \cap t) B), \end{aligned}$$

for all Borel sets $s, t \in \mathcal{B}_\Omega$, and all $A, B \in \mathcal{S}_2(H)$ (i.e., all $A, B \in \mathcal{L}(H)$, since $\dim H < \infty$).

Set $V : \mathcal{S}_2(H) \rightarrow H_{\tilde{K}}$ by

$$VA = \tilde{K}(\cdot, (\Omega, A)).$$

This is an isometric embedding of $\mathcal{S}_2(H)$ into $H_{\tilde{K}}$, since

$$\|VA\|_{H_{\tilde{K}}}^2 = \left\langle \tilde{K}(\cdot, (\Omega, A)), \tilde{K}(\cdot, (\Omega, A)) \right\rangle_{H_{\tilde{K}}} = \langle A, Q(\Omega) A \rangle_2 = \|A\|_2^2.$$

Lemma 5.7. *Given $(c_i)_{i=1}^N$ in $\mathbb{R}_+$, and $(s_i)_{i=1}^N$ in $\mathcal{B}_\Omega$, we have*

$$\sum_i |\text{tr}(AQ(s_i)) - c_i|^2 = \sum_i \left| \left\langle \tilde{K}_{(s_i, I_H)}, \tilde{K}_{(\Omega, A)} \right\rangle_{H_{\tilde{K}}} - c_i \right|^2.$$

Proof. Note that

$$\begin{aligned} \text{tr}(AQ(s)) &= \text{tr}(I_H^* Q(s) A) \\ &= \langle I_H, Q(s) A \rangle_2 = \left\langle \tilde{K}_{(s, I_H)}, \tilde{K}_{(\Omega, A)} \right\rangle_{H_{\tilde{K}}}. \end{aligned}$$

□

Corollary 5.8. *In the current setting, (5.1) is converted to the following constraint kernel approximation problem:*

$$\begin{aligned} \min \left\{ \sum_{i=1}^N \left| \left\langle \tilde{K}_{(s_i, I_H)}, \tilde{K}_{(\Omega, A)} \right\rangle_{H_{\tilde{K}}} - c_i \right|^2 : A \in \mathcal{S}_1^1(H) \right\} \\ \Updownarrow \\ \min \left\{ \sum_{i=1}^N |F(s_i, I_H) - c_i|^2 : F \in V(\mathcal{S}_1^1(H)) \subsetneq H_{\tilde{K}} \right\}. \end{aligned} \quad (5.3)$$

Proof. Recall that $VA(\cdot) = \tilde{K}(\cdot, (\Omega, A))$, so that

$$\left\langle \tilde{K}_{(s_i, I_H)}, \tilde{K}_{(\Omega, A)} \right\rangle_{H_{\tilde{K}}} = (VA)(s_i, I_H)$$

by the reproducing property of $H_{\tilde{K}}$. Then F in (5.3) is of the form $F = VA$, $A \in \mathcal{S}_1^1(H)$. $\square$

Remark 5.9. Kernel approximation methods are widely used in statistical learning for tasks such as regression, classification, and density estimation. They are particularly useful in settings where the underlying functional relationships are complex and non-linear, and where a balance between fitting the training data and maintaining generalizability to new data is crucial. The standard formulation is as follows:

$$\min \left\{ \sum_{i=1}^N |f(x_i) - y_i|^2 + \lambda \|f\|_{H_K}^2 : f \in H_K \right\} \quad (5.4)$$

where H_K is the RKHS of a scalar-valued kernel K on some set $S \times S$, $(y_i) \subset \mathbb{R}$, and λ is a regularization parameter.

A key difference between (5.3) and (5.4) is that, in (5.3), F is restricted to a convex subset, rather than being defined over the entire space $H_{\tilde{K}}$. A standard method to solve (5.3) is to use projected gradient descent.

Remark 5.10. For a given POVM Q as in (5.2), the construction of $H_{\tilde{K}}$ and V in the above example can be used to show that Q has a PVM dilation in $H_{\tilde{K}}$. The verification is similar to that of the Stinespring dilation theorem in Section 3 (also see [JT24, Section 2]).

Acknowledgement. For numerous enlightening discussions, we are grateful to Prof Wayne Polyzou, and to members of the Univ of Iowa weekly Math Physics seminar. We also benefited from discussions with our other colleagues, including Profs Daniel Alpay, Sergii Bezuglyi, Myung-Sin Song, Ilwoo Cho, and Shrey Sanadhya.

REFERENCES

- [AB97] D. Alpay and V. Bolotnikov, *On tangential interpolation in reproducing kernel Hilbert modules and applications*, Topics in interpolation theory (Leipzig, 1994), Oper. Theory Adv. Appl., vol. 95, Birkhäuser, Basel, 1997, pp. 37–68. MR 1473250
- [AD93] Daniel Alpay and Harry Dym, *On a new class of structured reproducing kernel spaces*, J. Funct. Anal. **111** (1993), no. 1, 1–28. MR 1200633
- [AD06] D. Alpay and C. Dubi, *Some remarks on the smoothing problem in a reproducing kernel Hilbert space*, J. Anal. Appl. **4** (2006), no. 2, 119–132. MR 2223568
- [AFMP94] Gregory T. Adams, John Froelich, Paul J. McGuire, and Vern I. Paulsen, *Analytic reproducing kernels and factorization*, Indiana Univ. Math. J. **43** (1994), no. 3, 839–856. MR 1305949

- [AJ15] Daniel Alpay and Palle Jorgensen, *Spectral theory for Gaussian processes: reproducing kernels, boundaries, and L^2 -wavelet generators with fractional scales*, Numer. Funct. Anal. Optim. **36** (2015), no. 10, 1239–1285. MR 3402823
- [AJ21] Daniel Alpay and Palle E. T. Jorgensen, *New characterizations of reproducing kernel Hilbert spaces and applications to metric geometry*, Opuscula Math. **41** (2021), no. 3, 283–300. MR 4302453
- [AL95] Daniel Alpay and Juliette Leblond, *Traces of Hardy functions and reproducing kernel Hilbert spaces*, Arch. Math. (Basel) **64** (1995), no. 6, 490–499. MR 1329822
- [Alp91] Daniel Alpay, *Some reproducing kernel spaces of continuous functions*, J. Math. Anal. Appl. **160** (1991), no. 2, 424–433. MR 1126127
- [ANW17] Pablo Arrighi, Vincent Nesme, and Reinhard F. Werner, *Bounds on the speedup in quantum signaling*, Phys. Rev. A **95** (2017), no. 1, 012331, 11. MR 3681182
- [Aro48] N. Aronszajn, *Reproducing and pseudo-reproducing kernels and their application to the partial differential equations of physics*, Harvard University, Graduate School of Engineering, Cambridge, MA, 1948, Studies in partial differential equations. Technical report 5, preliminary note. MR 31663
- [Arv10] William Arveson, *Dilation theory yesterday and today*, A glimpse at Hilbert space operators, Oper. Theory Adv. Appl., vol. 207, Birkhäuser Verlag, Basel, 2010, pp. 99–123. MR 2743416
- [AS56] N. Aronszajn and K. T. Smith, *Functional spaces and functional completion*, Ann. Inst. Fourier (Grenoble) **6** (1955/56), 125–185. MR 80878
- [BJ97] Ola Bratteli and Palle E. T. Jorgensen, *Isometries, shifts, Cuntz algebras and multiresolution wavelet analysis of scale N* , Integral Equations Operator Theory **28** (1997), no. 4, 382–443. MR 1465320
- [BJ99] ———, *Iterated function systems and permutation representations of the Cuntz algebra*, Mem. Amer. Math. Soc. **139** (1999), no. 663, x+89. MR 1469149
- [BJKW00] O. Bratteli, P. E. T. Jorgensen, A. Kishimoto, and R. F. Werner, *Pure states on $\mathcal{O}_d$* , J. Operator Theory **43** (2000), no. 1, 97–143. MR 1740897
- [CDP⁺13] Giulio Chiribella, Giacomo Mauro D’Ariano, Paolo Perinotti, Dirk Schlingemann, and Reinhard Werner, *A short impossibility proof of quantum bit commitment*, Phys. Lett. A **377** (2013), no. 15, 1076–1087. MR 3038250
- [CLPL19] Wen-Sheng Chen, Jingmin Liu, Binbin Pan, and Yugao Li, *Block kernel nonnegative matrix factorization for face recognition*, Int. J. Wavelets Multiresolut. Inf. Process. **17** (2019), no. 1, 1850059, 28. MR 3911880
- [CY17] Rupak Chatterjee and Ting Yu, *Generalized coherent states, reproducing kernels, and quantum support vector machines*, Quantum Inf. Comput. **17** (2017), no. 15-16, 1292–1306. MR 3700848
- [DHJ15] Dorin Ervin Dutkay, John Haussermann, and Palle E. T. Jorgensen, *Atomic representations of Cuntz algebras*, J. Math. Anal. Appl. **421** (2015), no. 1, 215–243. MR 3250475
- [DJ12] Dorin Ervin Dutkay and Palle E. T. Jorgensen, *Spectral measures and Cuntz algebras*, Math. Comp. **81** (2012), no. 280, 2275–2301. MR 2945156
- [DJ14] ———, *Monic representations of the Cuntz algebra and Markov measures*, J. Funct. Anal. **267** (2014), no. 4, 1011–1034. MR 3217056
- [DJ15] ———, *Representations of Cuntz algebras associated to quasi-stationary Markov measures*, Ergodic Theory Dynam. Systems **35** (2015), no. 7, 2080–2093. MR 3394108
- [DKM24] Ali Dadkhah, Mohsen Kian, and Mohammad Sal Moslehian, *Decomposition of tracial positive maps and applications in quantum information*, Anal. Math. Phys. **14** (2024), no. 3, Paper No. 48. MR 4736299
- [DM20] Nicolò Drago and Valter Moretti, *The notion of observable and the moment problem for $*$ -algebras and their GNS representations*, Lett. Math. Phys. **110** (2020), no. 7, 1711–1758. MR 4114386
- [EHD⁺13] Tobias Eberle, Vitus Händchen, Jörg Duhme, Torsten Franz, Fabian Furrer, Roman Schnabel, and Reinhard F. Werner, *Gaussian entanglement for quantum key distribution from a single-mode squeezing source*, New J. Phys. **15** (2013), no. May, 053049, 14. MR 3081872

- [FLZG24] Lingxuan Feng, Shunlong Luo, Yan Zhao, and Zhihua Guo, *Equioverlapping measurements as extensions of symmetric informationally complete positive operator valued measures*, Phys. Rev. A **109** (2024), no. 1, Paper No. 012218, 10. MR 4705975
- [FMBS24] Samuel Fedida, Anupam Mazumdar, Sougato Bose, and Alessio Serafini, *Entanglement entropy in scalar quantum electrodynamics*, Phys. Rev. D **109** (2024), no. 6, Paper No. 065028. MR 4735667
- [FS20] Martin Forde and Benjamin Smith, *The conditional law of the Bacry-Muzy and Riemann-Liouville log correlated Gaussian fields and their GMC, via Gaussian Hilbert and fractional Sobolev spaces*, Statist. Probab. Lett. **161** (2020), 108732, 11. MR 4073554
- [Gue22] J. C. Guella, *On Gaussian kernels on Hilbert spaces and kernels on hyperbolic spaces*, J. Approx. Theory **279** (2022), Paper No. 105765, 36. MR 4414825
- [HCW21] Shan Huang, Zeng-Bing Chen, and Shengjun Wu, *Entropic uncertainty relations for general symmetric informationally complete positive operator-valued measures and mutually unbiased measurements*, Phys. Rev. A **103** (2021), no. 4, Paper No. 042205, 9. MR 4256782
- [HKLP24] Teiko Heinosaari, Oskari Kerppo, Leevi Leppäjärvi, and Martin Plávala, *Simple information-processing tasks with unbounded quantum advantage*, Phys. Rev. A **109** (2024), no. 3, Paper No. 032627. MR 4732043
- [Hol22] A. S. Holevo, *On optimization problem for positive operator-valued measures*, Lobachevskii J. Math. **43** (2022), no. 7, 1646–1650. MR 4509531
- [JT23]alle E. T. Jorgensen and James Tian, *Operator-valued Gaussian processes and their covariance kernels*, Infinite Dimensional Analysis, Quantum Probability and Related Topics **0** (2023), no. 0, 2350020.
- [JT24]alle E. T. Jorgensen and James Tian, *Hilbert space valued Gaussian processes, their kernels, factorizations, and covariance structure*, arXiv: 2404.14685 (2024).
- [JYZ21] Hao Jiang, Ming Yi, and Shihua Zhang, *A kernel non-negative matrix factorization framework for single cell clustering*, Appl. Math. Model. **90** (2021), 875–888. MR 4160499
- [Kat19] Victor Katsnelson, *On the completeness of Gaussians in a Hilbert functional space*, Complex Anal. Oper. Theory **13** (2019), no. 3, 637–658. MR 3940383
- [KLSW12] J. Kiukas, P. Lahti, J. Schultz, and R. F. Werner, *Characterization of informational completeness for covariant phase space observables*, J. Math. Phys. **53** (2012), no. 10, 102103, 11. MR 3050559
- [KM20] Vít Kubelka and Bohdan Maslowski, *Filtering of Gaussian processes in Hilbert spaces*, Stoch. Dyn. **20** (2020), no. 3, 2050020, 22. MR 4101087
- [KSW08] Dennis Kretschmann, Dirk Schlingemann, and Reinhard F. Werner, *A continuity theorem for Stinespring’s dilation*, J. Funct. Anal. **255** (2008), no. 8, 1889–1904. MR 2462580
- [KW01] M. Keyl and R. F. Werner, *Estimating the spectrum of a density operator*, Phys. Rev. A (3) **64** (2001), no. 5, 052311, 5. MR 1878924
- [LK24] A. Lebedev and A. Khrennikov, *Quantum-like modeling of the order effect in decision making: POVM viewpoint on the Wang-Busemeyer QQ-equality*, Infinite dimensional analysis, quantum probability and related topics, QP-PQ: Quantum Probab. White Noise Anal., vol. 32, World Sci. Publ., Hackensack, NJ, [2024] ©2024, pp. 123–128. MR 4678079
- [LKKK24] Su-Yong Lee, Jihwan Kim, Zaeill Kim, and Duk Y. Kim, *Asymmetric entanglement for quantum target sensing*, Phys. Rev. A **109** (2024), no. 4, Paper No. 042429. MR 4740579
- [Lop23] Oleh Lopushansky, *Bernstein-Jackson inequalities on Gaussian Hilbert spaces*, J. Fourier Anal. Appl. **29** (2023), no. 5, Paper No. 58, 21. MR 4641110
- [LYA23] Binglin Li, Changwon Yoon, and Jeongyoun Ahn, *Reproducing kernels and new approaches in compositional data analysis*, J. Mach. Learn. Res. **24** (2023), Paper No. [327], 34. MR 4690276
- [OCL24] Eoin O’Connor, Steve Campbell, and Gabriel T. Landi, *Fisher information rates in sequentially measured quantum systems*, New J. Phys. **26** (2024), no. March, Paper No. 033048, 15. MR 4730881

- [PPPR20] Satish K. Pandey, Vern I. Paulsen, Jitendra Prakash, and Mizanur Rahaman, *Entanglement breaking rank and the existence of SIC POVMs*, J. Math. Phys. **61** (2020), no. 4, 042203, 14. MR 4085703
- [PR16] Vern I. Paulsen and Mrinal Raghupathi, *An introduction to the theory of reproducing kernel Hilbert spaces*, Cambridge Studies in Advanced Mathematics, vol. 152, Cambridge University Press, Cambridge, 2016. MR 3526117
- [PTD23] J. E. Pascoe and Ryan Tully-Doyle, *Induced Stinespring factorization and the Wittstock support theorem*, Results Math. **78** (2023), no. 4, Paper No. 135, 15. MR 4581177
- [PTG15] Andrés Esteban Páez-Torres and Fabio A. González, *Online kernel matrix factorization*, Progress in pattern recognition, image analysis, computer vision, and applications, Lecture Notes in Comput. Sci., vol. 9423, Springer, Cham, 2015, pp. 651–658. MR 3487535
- [QL17] Likuan Qin and Vadim Linetsky, *Long-term factorization of affine pricing kernels*, Math. Financ. Econ. **11** (2017), no. 4, 479–498. MR 3709384
- [QLY⁺24] Ji Qi, Ke Li, Zhe Yang, Rui-Yang Yuan, and Bao-Cang Ren, *Error identification entanglement purification for stationary system using high-dimensional entanglement*, Phys. Rev. A **109** (2024), no. 4, Paper No. 042423. MR 4740573
- [QY23] Minglong Qin and Penghui Yao, *Decidability of fully quantum nonlocal games with noisy maximally entangled states*, 50th International Colloquium on Automata, Languages, and Programming, LIPIcs. Leibniz Int. Proc. Inform., vol. 261, Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern, 2023, pp. Art. No. 97, 20. MR 4612967
- [RMC21] Benjamin Robinson, Bill Moran, and Doug Cochran, *Positive operator-valued measures and densely defined operator-valued frames*, Rocky Mountain J. Math. **51** (2021), no. 1, 265–272. MR 4280112
- [Sch52] Laurent Schwartz, *Théorie des noyaux*, Proceedings of the International Congress of Mathematicians, Cambridge, Mass., 1950, vol. 1, Amer. Math. Soc., Providence, RI, 1952, pp. 220–230. MR 45307
- [Sch64] ———, *Sous-espaces hilbertiens d’espaces vectoriels topologiques et noyaux associés (noyaux reproduisants)*, J. Analyse Math. **13** (1964), 115–256. MR 179587
- [Shi05] Serguei Shimorin, *Commutant lifting and factorization of reproducing kernels*, J. Funct. Anal. **224** (2005), no. 1, 134–159. MR 2139107
- [Sti55] W. Forrest Stinespring, *Positive functions on C^* -algebras*, Proc. Amer. Math. Soc. **6** (1955), 211–216. MR 69403
- [Ver22] Dominic Verdon, *A covariant Stinespring theorem*, J. Math. Phys. **63** (2022), no. 9, Paper No. 091705, 49. MR 4482713
- [Wer01] R. F. Werner, *All teleportation and dense coding schemes*, vol. 34, 2001, Quantum information and computation, pp. 7081–7094. MR 1863141
- [WNGB24] Elisabeth Wagner, Ramil Nigmatullin, Alexei Gilchrist, and Gavin K. Brennen, *Information flow in non-unitary quantum cellular automata*, SciPost Phys. **16** (2024), no. 1, Paper No. 014, 41. MR 4710381

(PALLE E.T. JORGENSEN) DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF IOWA, IOWA CITY, IA 52242-1419, U.S.A.

Email address: `palle-jorgensen@uiowa.edu`

(JAMES F. TIAN) MATHEMATICAL REVIEWS, 416 4TH STREET ANN ARBOR, MI 48103-4816, U.S.A.

Email address: `jft@ams.org`