

ASYMPTOTIC PROFILES OF GROUND STATE SOLUTIONS FOR CHOQUARD EQUATIONS WITH A GENERAL LOCAL PERTURBATION

SHIWANG MA

*School of Mathematical Sciences and LPMC, Nankai University
Tianjin 300071, China*

VITALY MOROZ

*Department of Mathematics, Swansea University
Fabian Way, Swansea SA1 8EN, Wales, UK*

ABSTRACT. In this paper, we study the asymptotic behavior of ground state solutions for the nonlinear Choquard equation with a general local perturbation

$$-\Delta u + \varepsilon u = (I_\alpha * |u|^p)|u|^{p-2}u + g(u), \quad \text{in } \mathbb{R}^N, \quad (P_\varepsilon)$$

where $N \geq 3$ is an integer, $p = \frac{N+\alpha}{N}$, or $\frac{N+\alpha}{N-2}$, I_α is the Riesz potential and $\varepsilon > 0$ is a parameter. Under some mild conditions on $g(u)$, we show that as $\varepsilon \rightarrow \infty$, after a *suitable rescaling* the ground state solutions of (P_ε) converge to a particular solution of some limit equations, and establish a sharp asymptotic characterisation of such a rescaling, which depend in a non-trivial way on the asymptotic behavior of the function $g(s)$ at infinity and the space dimension N . Based on this study, we also present some results on the existence and asymptotic behaviors of positive normalized solutions of (P_ε) with the normalization constraint $\int_{\mathbb{R}^N} |u|^2 = a^2$. Particularly, we obtain the asymptotic behavior of positive normalized solutions of such a problem as $a \rightarrow 0$ and $a \rightarrow \infty$.

E-mail addresses: shiwangm@nankai.edu.cn, v.moroz@swansea.ac.uk.

2010 Mathematics Subject Classification. Primary 35J60, 35Q55; Secondary 35B25, 35B40, 35R09, 35J91.

Key words and phrases. Nonlinear Choquard equation; groundstate solution; normalized solution; concentration compactness; asymptotic behavior.

1. INTRODUCTION

We study standing-wave solutions of the nonlinear Schrödinger equation with attractive combined nonlinearity

$$i\psi_t = \Delta\psi + (I_\alpha * |\psi|^p)|\psi|^{p-2}\psi + f(|\psi|)\psi \quad \text{in } \mathbb{R}^N \times \mathbb{R}, \quad (1.1)$$

where $N \geq 3$ is an integer, $\psi : \mathbb{R}^N \times \mathbb{R} \rightarrow \mathbb{C}$, $p \in [\frac{N+\alpha}{N}, \frac{N+\alpha}{N-2}]$, and I_α is the Riesz potential defined for every $x \in \mathbb{R}^N \setminus \{0\}$ by

$$I_\alpha(x) = \frac{A_\alpha(N)}{|x|^{N-\alpha}}, \quad A_\alpha(N) = \frac{\Gamma(\frac{N-\alpha}{2})}{\Gamma(\frac{\alpha}{2})\pi^{N/2}2^\alpha},$$

where Γ denotes the Gamma function, f is a continuous function to be specified later.

A theory of NLS with combined powers nonlinearity was developed by Tao, Visan and Zhang [33] and attracted a lot attention during the past decade (cf. [1, 2, 7, 8, 13, 15, 16, 17, 18, 22, 28, 38] and further references therein). The nonlocal convolution term in (1.1) represents the Newtonian gravitational attraction between bosonic particles. The local term takes into account the short-range self-interaction between bosons. For more background of Choquard equation, we refer the readers to [30] and references therein.

A standing-wave solutions of (1.1) with a frequency $\varepsilon > 0$ is a finite energy solution in the form

$$\psi(t, x) = e^{-i\varepsilon t}u(x).$$

Put $g(u) = f(|u|)u$, then this ansatz yields the equation for u in the form

$$-\Delta u + \varepsilon u = (I_\alpha * |u|^p)|u|^{p-2}u + g(u), \quad \text{in } \mathbb{R}^N. \quad (P_\varepsilon)$$

A solution $u_\varepsilon \in H^1(\mathbb{R}^N)$ of (P_ε) is a critical point of the Action functional defined by

$$I_\varepsilon(u) = \frac{1}{2} \int_{\mathbb{R}^N} |\nabla u|^2 + \frac{\varepsilon}{2} \int_{\mathbb{R}^N} |u|^2 - \frac{1}{2p} \int_{\mathbb{R}^N} (I_\alpha * |u|^p)|u|^p - \int_{\mathbb{R}^N} G(u), \quad (1.2)$$

where $G(u) = \int_0^u g(s)ds$. The existence and properties of ground state $u_\varepsilon \in H^1(\mathbb{R}^N)$ to (P_ε) can be obtained by using the argument as in [15, 16]. In the present paper, we are interested in the limit asymptotic profile of the groundstates u_ε of the problem (P_ε) , and in the asymptotic behaviors of different norms of u_ε , as $\varepsilon \rightarrow \infty$.

Many authors consider the following Schrödinger equation with two focusing exponents p and q :

$$-\Delta u + \varepsilon u = |u|^{p-2}u + |u|^{q-2}u, \quad \text{in } \mathbb{R}^N, \quad (1.3)$$

where $N \geq 3$, $2 < q < p \leq 2^* := \frac{2N}{N-2}$ and $\varepsilon > 0$ is a parameter. It is well-known [4, 24, 40] that when $p = 2^*$ and $q \in (2, 2^*)$ if $N \geq 4$, or $q \in (4, 6)$ if $N = 3$, the equation (1.3) admits a ground state solution u_ε for every $\varepsilon > 0$. Recently, T. Akahori et al. [3], and Wei and Wu [37] proved that (1.3) admits a ground state for all $N \geq 3$ and $\varepsilon > 0$ small, while, for $N = 3$ and $q \in (2, 4]$, (1.3) has no ground state for large $\varepsilon > 0$. When $p = 2^*$ and $q \in (2, 2^*)$, T. Akahori et al. [2] proved that for small $\varepsilon > 0$ the ground state is unique and as $\varepsilon \rightarrow 0$, the unique ground state u_ε tends to the unique positive solution of the equation $-\Delta u + u = u^{q-1}$. After a suitable rescaling, T. Akahori et al. [1] established a uniform decay estimate for the ground state u_ε , and then proved the uniqueness and nondegeneracy in $H_r^1(\mathbb{R}^N)$ of ground states u_ε for $N \geq 5$ and large $\varepsilon > 0$, and show that for $N \geq 3$, $q \in (2, 2^*)$ if $N \geq 4$, or $q \in (4, 6)$ if $N = 3$, as $\varepsilon \rightarrow \infty$, u_ε tends to a particular solution of the critical Emden–Fowler equation.

More recently, by using a global bifurcation argument, Jeanjean, Zhang and Zhong [12] also study the asymptotic behavior of solutions as $\varepsilon \rightarrow 0$ and $\varepsilon \rightarrow \infty$ for the equation (1.3) with a general *subcritical* nonlinearity and discuss the connection to the existence, non-existence and multiplicity of prescribed mass positive solutions to (1.3) with the associated L^2 constraint condition $\int_{\mathbb{R}^N} |u|^2 = a^2$. But the precise asymptotic profiles of positive solutions are not addressed there. For other related papers, we refer the reader to [26] and the references therein.

Li, Ma and Zhang [15] and Li and Ma [16] considered (P_ε) with $g(u) = |u|^{q-2}u$, where $q \in (2, 2^*]$, and established the existence, regularity and symmetry of the ground state solutions of (P_ε) . Especially, for any $\varepsilon > 0$, the equation (P_ε) admits a ground state solution if $p = \frac{N+\alpha}{N}$ and $q \in (2, 2 + \frac{4}{N})$, or $p = \frac{N+\alpha}{N-2}$ and $q \in (2, 2^*)$ for $N \geq 4$, $q \in (4, 6)$ for $N = 3$. More recently, in [27], the authors consider again the equation (P_ε) with $g(u) = |u|^{q-2}u$, and establish the precisely asymptotic behaviors of positive ground states as $\varepsilon \rightarrow 0$ and $\varepsilon \rightarrow \infty$. However, to the best of our knowledge, so far nothing has been achieved as for the precise asymptotic behaviors of ground states of (P_ε) with a general perturbation $g(u)$.

In [28], the second author and Muratov first study the asymptotic properties of ground states for a class of scalar field equations with a defocusing large exponent p and a focusing smaller exponent q . More precisely, the following equation

$$-\Delta u + \varepsilon u = |u|^{p-2}u - |u|^{q-2}u, \quad \text{in } \mathbb{R}^N, \quad (1.4)$$

is discussed, where $N \geq 3$, $q > p > 2$. Later, in [14], M. Lewin and S. Nodari prove a general result about the uniqueness and non-degeneracy of positive radial solutions to the above equation, and then the non-degeneracy of the unique solution u_ε allows them to derive its behavior in the two limits $\varepsilon \rightarrow 0$ and $\varepsilon \rightarrow \varepsilon_*$, where ε_* is a threshold of existence. Amongst other things, a precise asymptotic expression of $M(\varepsilon) = \|u_\varepsilon\|_2^2$ is obtained. This gives the uniqueness of energy minimizers at fixed mass in certain regimes.

In [22], Zeng Liu and the second author extend the results in [28] to a class of Choquard type equation

$$-\Delta u + \varepsilon u = (I_\alpha * |u|^p)|u|^{p-2}u - |u|^{q-2}u, \quad \text{in } \mathbb{R}^N, \quad (1.5)$$

where $N \geq 3$ is an integer. Under some assumptions on the exponents p and q , the limit profiles of the ground states are discussed in the two cases $\varepsilon \rightarrow 0$ and $\varepsilon \rightarrow \infty$. But the precisely asymptotic behaviors of ground states was not studied in [22].

Alternatively, one can search for solutions to (P_ε) having prescribed mass, and in this case $\varepsilon \in \mathbb{R}$ is part of the unknown. That is, for a fixed $a > 0$, search for $u \in H^1(\mathbb{R}^N)$ and $\lambda \in \mathbb{R}$ satisfying

$$\begin{cases} -\Delta u = \lambda u + (I_\alpha * |u|^p)|u|^{p-2}u + g(u), & \text{in } \mathbb{R}^N, \\ u \in H^1(\mathbb{R}^N), \quad \int_{\mathbb{R}^N} |u|^2 = a^2. \end{cases} \quad (1.6)$$

The solution of (1.6) is usually denoted by a pair (u, λ) and called a normalized solution.

This approach seems to be particularly meaningful from the physical point of view, and often offers a good insight of the dynamical properties of the standing-wave solutions for (1.1), such as stability or instability.

It is standard to check that the Energy functional

$$E(u) = \frac{1}{2} \int_{\mathbb{R}^N} |\nabla u|^2 - \frac{1}{2p} \int_{\mathbb{R}^N} (I_\alpha * |u|^p)|u|^p - \int_{\mathbb{R}^N} G(u) \quad (1.7)$$

is of class C^1 and that a critical point of E restricted to the (mass) constraint

$$S(c) = \{u \in H^1(\mathbb{R}^N) : \|u\|_2^2 = a^2\} \quad (1.8)$$

gives a solution to (1.6). Here $\lambda := -\varepsilon$ arises as a Lagrange multiplier. We refer the readers to [17, 18, 38] and the references therein.

For quite a long time the paper [10] was the only one dealing with existence of normalized solutions in cases when the energy is unbounded from below on the L^2 -constraint. More recently, however, problems of this type received much attention. In particular, many authors have made intensive studies to the following nonlinear Schrödinger equation with combined powers nonlinearity

$$\begin{cases} -\Delta u = \lambda u + |u|^{p-2}u + \mu|u|^{q-2}u, & \text{in } \mathbb{R}^N, \\ u \in H^1(\mathbb{R}^N), \quad \|u\|_2^2 = a^2, \end{cases} \quad (1.9)$$

where $N \geq 3$, $2 < q < p \leq 2^*$, $\mu > 0$ is a new parameter and $\lambda \in \mathbb{R}$ appears as a Lagrange multiplier. We refer the readers to [11, 12, 31, 32, 36, 37] and references therein for the existence and multiplicity of normalized solutions to the problem (1.9). In [36], Wei and Wu study the existence and asymptotic behaviors of normalized solutions for (1.9) with $p = 2^*$, and obtain some precise asymptotic behaviors of ground states and mountain-pass solutions as $\mu \rightarrow 0$ and μ goes to its upper bound. It is worth mentioned that some ODE technique plays a crucial role in the studies in [36]. We refer the readers to [11, 12, 31, 32, 37] for the asymptotic behaviors of normalized solutions when the parameter μ varies in its range.

Roughly speaking, the parameter μ in (1.9) plays a role of changing some thresholds for the existence. But it does not change the qualitative properties of solutions. In a sense, changing the parameter μ is equivalent to changing the mass $a > 0$. More precisely, it follows from the reduction given in [37] that finding a normalized solution of (1.9) with $p = 2^*$ and mass constrained condition $\int_{\mathbb{R}^N} |u|^2 = a^2$ is equivalent to finding a normalized solution of the problem

$$\begin{cases} -\Delta u = \lambda u + |u|^{2^*-2}u + |u|^{q-2}u, & \text{in } \mathbb{R}^N, \\ u \in H^1(\mathbb{R}^N), \quad \int_{\mathbb{R}^N} |u|^2 = a^2 \mu^{\frac{4}{2N-q(N-2)}}. \end{cases} \quad (1.10)$$

In particular, sending $\mu \rightarrow 0$ (resp. $\mu \rightarrow \infty$) is equivalent to sending the mass $a \rightarrow 0$ (resp. $a \rightarrow \infty$).

A very complete theory has been established as to the existence and multiplicity of normalized solutions to (1.9). However, to the best of our knowledge, little has been done for the Schrödinger equation with combined general nonlinearity such as

$$\begin{cases} -\Delta u = \lambda u + |u|^{2^*-2}u + g(u), & \text{in } \mathbb{R}^N, \\ u \in H^1(\mathbb{R}^N), \quad \|u\|_2^2 = a^2. \end{cases} \quad (1.11)$$

The main reason is that the so called fibreing map is complicated and the technique to handle the Schrödinger equation with combined powers nonlinearity does not work any more in this case. To investigate the existence and asymptotic behavior of normalized solutions to (1.11), it seems that some new approach must be employed.

As for the problem (1.6) with $g(u) = \mu|u|^{q-2}u$, by adopting the technique used in [11, 12, 31, 32], some similar existence results on normalized solutions are already obtained in [17, 18, 19, 20, 38, 39] in the whole possible range of parameters, but the precisely asymptotic behavior of the normalized solutions are not addressed there.

The purpose of this paper is two-fold. Firstly, by using some new technique, we give some precise asymptotic behaviors of ground state solutions of (P_ε) under some mild subcritical growth conditions on the function $g(u)$. Secondly, as an application of our main results, we present some existence results of positive normalized solutions to (1.6) and their precise asymptotic behaviors. To explore the existence and precise asymptotic behaviors of normalized solutions of (1.6), it seems that the classical method developed recently in [11, 12, 31, 32] is hard to be adopted since the associated fibreing map is more complicated in our setting. Besides, due to the appearance of the nonlocal term, the ODE technique does not work well any more, this prevent us from adopting the technique used in [36] to derivate the precise asymptotic behaviors of normalized solutions.

Organization of the paper. In Section 2, we state the main results in this paper. In Section 3, we give some preliminary results which are needed in the proof of our main results. Sections 4–5 are devoted to the proofs of Theorems 2.1–2.2, respectively. Finally, in the last section, we present some existence results of positive normalized solutions and their precise asymptotic behaviors.

Basic notations. Throughout this paper, we assume $N \geq 3$. $C_c^\infty(\mathbb{R}^N)$ denotes the space of the functions infinitely differentiable with compact support in \mathbb{R}^N . $L^p(\mathbb{R}^N)$ with $1 \leq p < \infty$ denotes the Lebesgue space with the norms $\|u\|_p = (\int_{\mathbb{R}^N} |u|^p)^{1/p}$. $H^1(\mathbb{R}^N)$ is the usual Sobolev space with norm $\|u\|_{H^1(\mathbb{R}^N)} = (\int_{\mathbb{R}^N} |\nabla u|^2 + |u|^2)^{1/2}$. $D^{1,2}(\mathbb{R}^N) = \{u \in L^{2^*}(\mathbb{R}^N) : |\nabla u| \in L^2(\mathbb{R}^N)\}$. $H_r^1(\mathbb{R}^N) = \{u \in H^1(\mathbb{R}^N) : u \text{ is radially symmetric}\}$. B_r denotes the ball in \mathbb{R}^N with radius $r > 0$ and centered at the origin, $|B_r|$ and B_r^c denote its Lebesgue measure and its complement in \mathbb{R}^N , respectively. As usual, C, c , etc., denote generic positive constants. For any large $\epsilon > 0$ and two nonnegative functions $f(\epsilon)$ and $g(\epsilon)$, we write

- (1) $f(\epsilon) \lesssim g(\epsilon)$ or $g(\epsilon) \gtrsim f(\epsilon)$ if there exists a positive constant C independent of ϵ such that $f(\epsilon) \leq Cg(\epsilon)$.
- (2) $f(\epsilon) \sim g(\epsilon)$ if $f(\epsilon) \lesssim g(\epsilon)$ and $f(\epsilon) \gtrsim g(\epsilon)$.

If $|f(\epsilon)| \lesssim |g(\epsilon)|$, we write $f(\epsilon) = O(g(\epsilon))$. We also denote by $\Theta = \Theta(\epsilon)$ a generic positive function satisfying $C_1\epsilon \leq \Theta(\epsilon) \leq C_2\epsilon$ for some positive numbers $C_1, C_2 > 0$, which are independent of ϵ . Finally, if $\lim f(\epsilon)/g(\epsilon) = 1$ as $\epsilon \rightarrow \infty$, then we write $f(\epsilon) \simeq g(\epsilon)$ as $\epsilon \rightarrow \infty$.

2. MAIN RESULTS

In the present paper, we study the asymptotic behavior of positive groundstates of (P_ε) involving a critical exponent in the sense of Hardy-Littlewood-Sobolev inequality. More precisely, if $p = \frac{N+\alpha}{N}$, or $\frac{N+\alpha}{N-2}$, we show that as $\varepsilon \rightarrow \infty$, after a *suitable rescaling* the ground state solutions u_ε of (P_ε) converge in $H^1(\mathbb{R}^N)$ to a particular solution of some limit equations. We also establish a sharp asymptotic characteritaton of such a rescaling, and the precise asymptotic behaviors of $u_\varepsilon(0)$, $\|\nabla u_\varepsilon\|_2^2$, $\|u_\varepsilon\|_2^2$ and the least energy of the groundstates.

To state our main results, we make the following assumptions:

- (H1) $g \in C([0, \infty), [0, \infty))$ satisfies $g(s) = o(s)$ as $s \rightarrow 0$;
- (H2) there exists $q \in (2, 2^*)$ such that $\lim_{s \rightarrow +\infty} g(s)s^{1-q} = A > 0$.

When $p = \frac{N+\alpha}{N}$, $q \in (2, 2 + \frac{4\alpha}{N(2+\alpha)})$ and $\varepsilon \rightarrow \infty$, we are going to show that after a suitable rescaling the limit equation for (P_ε) is given by the critical Hardy-Littlewood-Sobolev equation

$$U = (I_\alpha * |U|^{\frac{N+\alpha}{N}}) U^{\frac{\alpha}{N}} \quad \text{in } \mathbb{R}^N. \quad (2.1)$$

It is well-known that the radial ground states of (2.1) are given by the function

$$U_1(x) := \left(\frac{A_0}{1 + |x|^2} \right)^{\frac{N}{2}} \quad (2.2)$$

with a suitable constant $A_0 > 0$, and the family of its rescalings

$$U_\rho(x) := \rho^{-\frac{N}{2}} U_1(x/\rho), \quad \rho > 0. \quad (2.3)$$

If $p = \frac{N+\alpha}{N}$, $q \in (2 + \frac{4\alpha}{N(2+\alpha)}, 2 + \frac{4}{N})$ and $\varepsilon \rightarrow \infty$, then it turns out that after some suitable rescaling the limit equation for (P_ε) is given by

$$-\Delta W + W = AW^{q-1}. \quad (2.4)$$

For convenience, we set

$$S_1 := \inf_{u \in H^1(\mathbb{R}^N) \setminus \{0\}} \frac{\int_{\mathbb{R}^N} |u|^2}{\left(\int_{\mathbb{R}^N} (I_\alpha * |u|^{\frac{N+\alpha}{N}} |u|^{\frac{N+\alpha}{N}})^{\frac{N}{N+\alpha}} \right)}, \quad (2.5)$$

and

$$S_q := \inf_{u \in H^1(\mathbb{R}^N) \setminus \{0\}} \frac{\int_{\mathbb{R}^N} |\nabla u|^2 + |u|^2}{\left(\int_{\mathbb{R}^N} |u|^q \right)^{\frac{2}{q}}}. \quad (2.6)$$

Theorem 2.1. *Assume (H1), (H2) hold, $p = \frac{N+\alpha}{N}$ and $q \in (2, 2 + \frac{4}{N})$, then for large $\varepsilon > 0$, the problem (P_ε) admits a positive ground state $u_\varepsilon \in H^1(\mathbb{R}^N)$, which is radially symmetric and radially nonincreasing. Moreover, if $q \in (2, 2 + \frac{4\alpha}{N(2+\alpha)})$ and $\varepsilon \rightarrow \infty$, then there exists $\zeta_\varepsilon \in (0, +\infty)$ verifying*

$$\zeta_\varepsilon \sim \varepsilon^{-\frac{N(q-2)}{\alpha[4-N(q-2)]}}$$

such that as $\varepsilon \rightarrow \infty$, the rescaled family of ground states $\tilde{w}_\varepsilon(x) = \varepsilon^{-\frac{N}{2\alpha}} \zeta_\varepsilon^{\frac{N}{2}} u_\varepsilon(\zeta_\varepsilon x)$ converges in $H^1(\mathbb{R}^N)$ to the extremal function U_{ρ_0} with

$$\rho_0 = \left(\frac{2q \int_{\mathbb{R}^N} |\nabla U_1|^2}{NA(q-2) \int_{\mathbb{R}^N} |U_1|^q} \right)^{\frac{2}{4-N(q-2)}}. \quad (2.7)$$

If $q = 2 + \frac{4\alpha}{N(2+\alpha)}$ and $\varepsilon \rightarrow \infty$, then up to a subsequence, the rescaled family of ground states $v_\varepsilon(x) = \varepsilon^{-\frac{1}{q-2}} u_\varepsilon(\varepsilon^{-\frac{1}{2}} x)$ converges in $H^1(\mathbb{R}^N)$ to a positive solution of the equation

$$-\Delta v + v = (I_\alpha * |v|^{\frac{N+\alpha}{N}}) |v|^{\frac{N+\alpha}{N}-2} v + A |v|^{\frac{4\alpha}{N(2+\alpha)}} v.$$

If $q \in (2 + \frac{4\alpha}{N(2+\alpha)}, 2 + \frac{4}{N})$ and $\varepsilon \rightarrow \infty$, then the rescaled family of ground states $v_\varepsilon(x) = \varepsilon^{-\frac{1}{q-2}} u_\varepsilon(\varepsilon^{-\frac{1}{2}} x)$ converges in $H^1(\mathbb{R}^N)$ to the unique positive solution W of the equation (2.4).

Furthermore, as $\varepsilon \rightarrow \infty$, the least energy m_ε of the ground state satisfies

$$m_\varepsilon = \begin{cases} \varepsilon^{\frac{N+\alpha}{\alpha}} \left[\frac{\alpha}{2(N+\alpha)} S_1^{\frac{N+\alpha}{\alpha}} - \Theta(\varepsilon^{-\frac{4\alpha-N(2+\alpha)(q-2)}{\alpha[4-N(q-2)]}}) \right], & \text{if } q \in (2, 2 + \frac{4\alpha}{N(2+\alpha)}), \\ \varepsilon^{\frac{2N-q(N-2)}{2(q-2)}} \left[\frac{q-2}{2q} A^{-\frac{2}{q-2}} S_q^{\frac{q}{q-2}} + o_\varepsilon(1) \right], & \text{if } q \in (2 + \frac{4\alpha}{N(2+\alpha)}, 2 + \frac{4}{N}). \end{cases}$$

Corollary 2.1. Assume (H1), (H2) hold, $p = \frac{N+\alpha}{N}$ and $q \in (2, 2 + \frac{4}{N})$. Let u_ε be the ground state solution of (P_ε) , then as $\varepsilon \rightarrow \infty$

$$\begin{aligned} u_\varepsilon(0) &\sim \begin{cases} \varepsilon^{\frac{2N}{\alpha[4-N(q-2)]}}, & \text{if } q \in (2, 2 + \frac{4\alpha}{N(2+\alpha)}), \\ \varepsilon^{\frac{1}{q-2}}, & \text{if } q \in [2 + \frac{4\alpha}{N(2+\alpha)}, 2 + \frac{4}{N}), \end{cases} \\ \|u_\varepsilon\|_2^2 &\sim \begin{cases} \varepsilon^{\frac{N}{\alpha}}, & \text{if } q \in (2, 2 + \frac{4\alpha}{N(2+\alpha)}), \\ \varepsilon^{\frac{4-N(q-2)}{2(q-2)}}, & \text{if } q \in [2 + \frac{4\alpha}{N(2+\alpha)}, 2 + \frac{4}{N}), \end{cases} \\ \|\nabla u_\varepsilon\|_2^2 \sim \|u_\varepsilon\|_q^q &\sim \begin{cases} \varepsilon^{\frac{N[2N-q(N-2)]}{\alpha[4-N(q-2)]}}, & \text{if } q \in (2, 2 + \frac{4\alpha}{N(2+\alpha)}), \\ \varepsilon^{\frac{2N-q(N-2)}{2(q-2)}}, & \text{if } q \in [2 + \frac{4\alpha}{N(2+\alpha)}, 2 + \frac{4}{N}), \end{cases} \\ \int_{\mathbb{R}^N} (I_\alpha * |u_\varepsilon|^{\frac{N+\alpha}{N}}) |u_\varepsilon|^{\frac{N+\alpha}{N}} &\sim \begin{cases} \varepsilon^{\frac{N+\alpha}{\alpha}}, & \text{if } q \in (2, 2 + \frac{4\alpha}{N(2+\alpha)}), \\ \varepsilon^{\frac{(N+\alpha)[4-N(q-2)]}{2N(q-2)}}, & \text{if } q \in [2 + \frac{4\alpha}{N(2+\alpha)}, 2 + \frac{4}{N}). \end{cases} \end{aligned}$$

Remark 2.1. Corollary 2.1 is a direct consequence of Theorem 2.1. Surprisingly, in Theorem 2.1, if $q \in (2, 2 + \frac{4\alpha}{N(2+\alpha)})$, then as $\varepsilon \rightarrow \infty$, the rescaled family of ground states \tilde{w}_ε converges to the extremal function U_{ρ_0} in $H^1(\mathbb{R}^N)$ other than in $L^2(\mathbb{R}^N)$. Besides, from the main body of the paper, the following more precise estimates are valid:

If $q \in (2, 2 + \frac{4\alpha}{N(2+\alpha)})$ and $\varepsilon \rightarrow \infty$, then the ground state solutions satisfy

$$\begin{aligned} \|u_\varepsilon\|_2^2 &= \varepsilon^{\frac{N}{\alpha}} \left[S_1^{\frac{N+\alpha}{\alpha}} + O(\varepsilon^{-\frac{4\alpha-N(2+\alpha)(q-2)}{\alpha[4-N(q-2)]}}) \right], \\ \|\nabla u_\varepsilon\|_2^2 &= \frac{NA(q-2)}{2q} \|u_\varepsilon\|_q^q + o(\varepsilon^{\frac{N[2N-q(N-2)]}{\alpha[4-N(q-2)]}}) \sim \varepsilon^{\frac{N[2N-q(N-2)]}{\alpha[4-N(q-2)]}}, \\ \int_{\mathbb{R}^N} (I_\alpha * |u_\varepsilon|^{\frac{N+\alpha}{N}}) |u_\varepsilon|^{\frac{N+\alpha}{N}} &= \varepsilon^{\frac{N+\alpha}{\alpha}} \left[S_1^{\frac{N+\alpha}{\alpha}} + O(\varepsilon^{-\frac{4\alpha-N(2+\alpha)(q-2)}{\alpha[4-N(q-2)]}}) \right]. \end{aligned}$$

If $q \in (2 + \frac{4\alpha}{N(2+\alpha)}, 2 + \frac{4}{N})$ and $\varepsilon \rightarrow \infty$, then the ground state solutions satisfy

$$\begin{aligned} \|u_\varepsilon\|_2^2 &= \varepsilon^{\frac{4-N(q-2)}{2(q-2)}} \left[\frac{2N-q(N-2)}{2q} A^{-\frac{2}{q-2}} S_q^{\frac{q}{q-2}} + o_\varepsilon(1) \right], \\ \|\nabla u_\varepsilon\|_2^2 &= \varepsilon^{\frac{2N-q(N-2)}{2(q-2)}} \left[\frac{N(q-2)}{2q} A^{-\frac{2}{q-2}} S_q^{\frac{q}{q-2}} + o_\varepsilon(1) \right], \\ \|u_\varepsilon\|_q^q &= \varepsilon^{\frac{2N-q(N-2)}{2(q-2)}} \left[A^{-\frac{q}{q-2}} S_q^{\frac{q}{q-2}} + o_\varepsilon(1) \right]. \end{aligned}$$

When $p = \frac{N+\alpha}{N-2}$ and $\varepsilon \rightarrow \infty$, we are going to show that after a suitable rescaling the limit equation for (P_ε) is given by the critical Choquard equation

$$-\Delta V = (I_\alpha * |V|^{\frac{N+\alpha}{N-2}}) V^{\frac{2+\alpha}{N-2}} \quad \text{in } \mathbb{R}^N. \quad (2.8)$$

It is well-known that the radial ground states of (2.8) are given by the Talenti function

$$V_1(x) := [N(N-2)]^{\frac{N-2}{4}} \left(\frac{1}{1+|x|^2} \right)^{\frac{N-2}{2}} \quad (2.9)$$

and the family of its rescalings

$$V_\rho(x) := \rho^{-\frac{N-2}{2}} V_1(x/\rho), \quad \rho > 0. \quad (2.10)$$

For convenience, we set

$$S_\alpha := \inf_{v \in D^{1,2}(\mathbb{R}^N) \setminus \{0\}} \frac{\int_{\mathbb{R}^N} |\nabla v|^2}{\left(\int_{\mathbb{R}^N} (I_\alpha * |v|^{\frac{N+\alpha}{N-2}}) |v|^{\frac{N+\alpha}{N-2}} \right)^{\frac{N-2}{N+\alpha}}}. \quad (2.11)$$

Theorem 2.2. *Assume (H1) and (H2) hold, $p = \frac{N+\alpha}{N-2}$, $q \in (2, 2^*)$ if $N \geq 4$ and $q \in (4, 6)$ if $N = 3$, then for large $\varepsilon > 0$, the problem (P_ε) admits a positive ground state $u_\varepsilon \in H^1(\mathbb{R}^N)$, which is radially symmetric and radially nonincreasing. Moreover, if $N \geq 5$, then for large $\varepsilon > 0$, there exists $\zeta_\varepsilon \in (0, +\infty)$ verifying*

$$\zeta_\varepsilon \sim \varepsilon^{-\frac{2}{(N-2)(q-2)}}$$

such that as $\varepsilon \rightarrow \infty$, the rescaled family of ground states $w_\varepsilon(x) = \zeta_\varepsilon^{\frac{N-2}{2}} u_\varepsilon(\zeta_\varepsilon x)$ converges in $H^1(\mathbb{R}^N)$ to the extremal function V_{ρ_0} with

$$\rho_0 = \left(\frac{A[2N - q(N-2)] \int_{\mathbb{R}^N} |V_1|^q}{2q \int_{\mathbb{R}^N} |V_1|^2} \right)^{\frac{2}{(N-2)(q-2)}}. \quad (2.12)$$

If $N = 4, 3$, then for large $\varepsilon > 0$, there exists $\zeta_\varepsilon \in (0, +\infty)$ verifying

$$\zeta_\varepsilon \sim \begin{cases} (\varepsilon \ln \varepsilon)^{-\frac{1}{q-2}}, & \text{if } N = 4, \\ \varepsilon^{-\frac{1}{q-4}}, & \text{if } N = 3, \end{cases}$$

such that the rescaled family of ground states $\tilde{w}_\varepsilon(x) = \zeta_\varepsilon^{\frac{N-2}{2}} u_\varepsilon(\zeta_\varepsilon x)$ satisfies

$$\|\nabla \tilde{w}_\varepsilon\|_2^2 \sim \|\tilde{w}_\varepsilon\|_q^q \sim \int_{\mathbb{R}^N} (I_\alpha * |\tilde{w}_\varepsilon|^{\frac{N+\alpha}{N-2}}) |\tilde{w}_\varepsilon|^{\frac{N+\alpha}{N-2}} \sim 1, \quad \|\tilde{w}_\varepsilon\|_2^2 \sim \begin{cases} \ln \varepsilon, & \text{if } N = 4, \\ \varepsilon^{\frac{6-q}{2(q-4)}}, & \text{if } N = 3, \end{cases}$$

and as $\varepsilon \rightarrow \infty$, \tilde{w}_ε converges in $D^{1,2}(\mathbb{R}^N)$ and $L^s(\mathbb{R}^N)$ to V_1 for any $s > \frac{N}{N-2}$.

Furthermore, as $\varepsilon \rightarrow \infty$, the least energy m_ε of the ground state satisfies

$$\frac{2+\alpha}{2(N+\alpha)} S_\alpha^{\frac{N+\alpha}{2+\alpha}} - m_\varepsilon \sim \begin{cases} \varepsilon^{-\frac{2N-q(N-2)}{(N-2)(q-2)}}, & \text{if } N \geq 5, \\ (\varepsilon \ln \varepsilon)^{-\frac{4-q}{q-2}}, & \text{if } N = 4, \\ \varepsilon^{-\frac{6-q}{2(q-4)}}, & \text{if } N = 3. \end{cases}$$

Corollary 2.2. Assume (H1) and (H2) hold, $p = \frac{N+\alpha}{N-2}$, $q \in (2, 2^*)$ if $N \geq 4$ and $q \in (4, 6)$ if $N = 3$. Let u_ε be the ground state solution of (P_ε) , then as $\varepsilon \rightarrow \infty$

$$u_\varepsilon(0) \sim \begin{cases} \varepsilon^{\frac{1}{q-2}}, & \text{if } N \geq 5, \\ (\varepsilon \ln \varepsilon)^{\frac{1}{q-2}}, & \text{if } N = 4, \\ \varepsilon^{\frac{1}{2(q-4)}}, & \text{if } N = 3, \end{cases}$$

$$\|u_\varepsilon\|_q^q \sim \begin{cases} \varepsilon^{-\frac{4}{(N-2)(q-2)}}, & \text{if } N \geq 5, \\ (\varepsilon \ln \varepsilon)^{-\frac{4-q}{q-2}}, & \text{if } N = 4, \\ \varepsilon^{-\frac{6-q}{2(q-4)}}, & \text{if } N = 3, \end{cases} \quad \|u_\varepsilon\|_2^2 \sim \begin{cases} \varepsilon^{-\frac{4}{(N-2)(q-2)}}, & \text{if } N \geq 5, \\ \varepsilon^{-\frac{2}{q-2}} (\ln \varepsilon)^{-\frac{4-q}{q-2}}, & \text{if } N = 4, \\ \varepsilon^{-\frac{q-2}{2(q-4)}}, & \text{if } N = 3, \end{cases}$$

$$\|\nabla u_\varepsilon\|_2^2 = S_\alpha^{\frac{N+\alpha}{2+\alpha}} + \begin{cases} O(\varepsilon^{-\frac{2N-q(N-2)}{(N-2)(q-2)}}), & \text{if } N \geq 5, \\ O((\varepsilon \ln \varepsilon)^{-\frac{4-q}{q-2}}), & \text{if } N = 4, \\ O(\varepsilon^{-\frac{6-q}{2(q-4)}}), & \text{if } N = 3, \end{cases}$$

$$\int_{\mathbb{R}^N} (I_\alpha * |u_\varepsilon|^{\frac{N+\alpha}{N-2}}) |u_\varepsilon|^{\frac{N+\alpha}{N-2}} = S_\alpha^{\frac{N+\alpha}{2+\alpha}} + \begin{cases} O(\varepsilon^{-\frac{2N-q(N-2)}{(N-2)(q-2)}}), & \text{if } N \geq 5, \\ O((\varepsilon \ln \varepsilon)^{-\frac{4-q}{q-2}}), & \text{if } N = 4, \\ O(\varepsilon^{-\frac{6-q}{2(q-4)}}), & \text{if } N = 3. \end{cases}$$

Remark 2.2. Corollary 2.2 is a direct consequence of Theorem 2.2. In Theorem 2.2, we need not the assumption that $\alpha > N - 4$ in the case $N \geq 5$, and if $\alpha > N - 4$ holds true, then we can choose $\zeta_\varepsilon = \varepsilon^{-\frac{2}{(N-2)(q-2)}}$.

3. PRELIMINARIES

In this section, we present some preliminary results which are needed in the proof of our main results. Firstly, we consider the following Choquard type equation with combined nonlinearities:

$$-\Delta u + \varepsilon u = (I_\alpha * |u|^p) |u|^p + g(u), \quad \text{in } \mathbb{R}^N, \quad (P_\varepsilon)$$

where $N \geq 3$, $\alpha \in (0, N)$, $p \in [\frac{N+\alpha}{N}, \frac{N+\alpha}{N-2}]$, $\varepsilon > 0$ is a large parameter, and $g(u)$ is a function satisfying the assumptions (H1) and (H2) in the last section.

It has been proved in [16, Theorem 2.1] that any weak solution of (P_ε) in $H^1(\mathbb{R}^N)$ has additional regularity properties, which allows us to establish the Pohožaev identity for all finite energy solutions.

Lemma 3.1. Assume (H1) and (H2) hold. If $u \in H^1(\mathbb{R}^N)$ is a solution of (P_ε) , then $u \in W_{\text{loc}}^{2,r}(\mathbb{R}^N)$ for every $r > 1$. Moreover, u satisfies the Pohožaev identity

$$P_\varepsilon(u) := \frac{N-2}{2} \int_{\mathbb{R}^N} |\nabla u|^2 + \frac{N\varepsilon}{2} \int_{\mathbb{R}^N} |u|^2 - \frac{N+\alpha}{2p} \int_{\mathbb{R}^N} (I_\alpha * |u|^p) |u|^p - N \int_{\mathbb{R}^N} G(u) = 0. \quad (3.1)$$

It is well known that any weak solution of (P_ε) corresponds to a critical point of the action functionals I_ε defined in (1.2), which is well defined and is of C^1 in $H^1(\mathbb{R}^N)$. A nontrivial solution $u_\varepsilon \in H^1(\mathbb{R}^N)$ is called a ground-state if

$$I_\varepsilon(u_\varepsilon) = m_\varepsilon := \inf\{I_\varepsilon(u) : u \in H^1(\mathbb{R}^N) \setminus \{0\} \text{ and } I'_\varepsilon(u) = 0\}. \quad (3.2)$$

In [15, 16] (see also the proof of the main results in [16]), it has been shown that

$$m_\varepsilon = \inf_{u \in \mathcal{M}_\varepsilon} I_\varepsilon(u) = \inf_{u \in \mathcal{P}_\varepsilon} I_\varepsilon(u), \quad (3.3)$$

where \mathcal{M}_ε and \mathcal{P}_ε are the corresponding Nehari and Pohožaev manifolds defined by

$$\mathcal{M}_\varepsilon := \left\{ u \in H^1(\mathbb{R}^N) \setminus \{0\} \mid \int_{\mathbb{R}^N} |\nabla u|^2 + \varepsilon |u|^2 = \int_{\mathbb{R}^N} (I_\alpha * |u|^p) |u|^p + \int_{\mathbb{R}^N} g(u)u \right\}$$

and

$$\mathcal{P}_\varepsilon := \{ u \in H^1(\mathbb{R}^N) \setminus \{0\} \mid P_\varepsilon(u) = 0 \},$$

respectively. Moreover, the following min-max descriptions are valid:

Lemma 3.2. *Let*

$$u_t(x) = \begin{cases} u(\frac{x}{t}), & \text{if } t > 0, \\ 0, & \text{if } t = 0, \end{cases}$$

then

$$m_\varepsilon = \inf_{u \in H^1(\mathbb{R}^N) \setminus \{0\}} \sup_{t \geq 0} I_\varepsilon(tu) = \inf_{u \in H^1(\mathbb{R}^N) \setminus \{0\}} \sup_{t \geq 0} I_\varepsilon(u_t). \quad (3.4)$$

In particular, we have $m_\varepsilon = I_\varepsilon(u_\varepsilon) = \sup_{t > 0} I_\varepsilon(tu_\varepsilon) = \sup_{t > 0} I_\varepsilon((u_\varepsilon)_t)$.

The following well known Hardy-Littlewood-Sobolev inequality can be found in [23].

Lemma 3.3. *Let $p, r > 1$ and $0 < \alpha < N$ with $1/p + (N - \alpha)/N + 1/r = 2$. Let $u \in L^p(\mathbb{R}^N)$ and $v \in L^r(\mathbb{R}^N)$. Then there exists a sharp constant $C(N, \alpha, p)$, independent of u and v , such that*

$$\left| \int_{\mathbb{R}^N} \int_{\mathbb{R}^N} \frac{u(x)v(y)}{|x - y|^{N-\alpha}} \right| \leq C(N, \alpha, p) \|u\|_p \|v\|_r.$$

If $p = r = \frac{2N}{N+\alpha}$, then

$$C(N, \alpha, p) = C_\alpha(N) = \pi^{\frac{N-\alpha}{2}} \frac{\Gamma(\frac{\alpha}{2})}{\Gamma(\frac{N+\alpha}{2})} \left\{ \frac{\Gamma(\frac{N}{2})}{\Gamma(N)} \right\}^{-\frac{\alpha}{N}}.$$

Remark 3.1. By the Hardy-Littlewood-Sobolev inequality, for any $v \in L^s(\mathbb{R}^N)$ with $s \in (1, \frac{N}{\alpha})$, $I_\alpha * v \in L^{\frac{Ns}{N-\alpha s}}(\mathbb{R}^N)$ and

$$\|I_\alpha * v\|_{\frac{Ns}{N-\alpha s}} \leq A_\alpha(N) C(N, \alpha, s) \|v\|_s. \quad (3.5)$$

Lemma 3.4. (P. L. Lions [25]) *Let $r > 0$ and $2 \leq q \leq 2^*$. If (u_n) is bounded in $H^1(\mathbb{R}^N)$ and if*

$$\sup_{y \in \mathbb{R}^N} \int_{B_r(y)} |u_n|^q \rightarrow 0, \text{ as } n \rightarrow \infty,$$

then $u_n \rightarrow 0$ in $L^s(\mathbb{R}^N)$ for $2 < s < 2^$. Moreover, if $q = 2^*$, then $u_n \rightarrow 0$ in $L^{2^*}(\mathbb{R}^N)$.*

Lemma 3.5. (Radial Lemma A.II, H. Berestycki and P. L. Lions [5]) *Let $N \geq 2$, then every radial function $u \in H^1(\mathbb{R}^N)$ is almost everywhere equal to a function $\tilde{u}(x)$, continuous for $x \neq 0$, such that*

$$|\tilde{u}(x)| \leq C_N |x|^{(1-N)/2} \|u\|_{H^1(\mathbb{R}^N)} \quad \text{for } |x| \geq \alpha_N, \quad (3.6)$$

where C_N and α_N depend only on the dimension N .

Lemma 3.6. (Radial Lemma A.III. H. Berestycki and P. L. Lions [5]) *Let $N \geq 3$, then every radial function u in $D^{1,2}(\mathbb{R}^N)$ is almost everywhere equal to a function $\tilde{u}(x)$, continuous for $x \neq 0$, such that*

$$|\tilde{u}(x)| \leq C_N |x|^{(2-N)/2} \|u\|_{D^{1,2}(\mathbb{R}^N)} \quad \text{for } |x| \geq 1, \quad (3.7)$$

where C_N only depends on N .

The following lemma is proved in [27].

Lemma 3.7. *Let $0 < \alpha < N$, $0 \leq f(x) \in L^1(\mathbb{R}^N)$ be a radially symmetric function such that*

$$\lim_{|x| \rightarrow +\infty} f(|x|) |x|^N = 0. \quad (3.8)$$

If $\alpha \leq 1$, we additionally assume that f is monotone non-increasing. Then as $|x| \rightarrow +\infty$, we have

$$\int_{\mathbb{R}^N} \frac{f(y)}{|x-y|^{N-\alpha}} dy = \frac{\|f\|_{L^1}}{|x|^{N-\alpha}} + o\left(\frac{1}{|x|^{N-\alpha}}\right). \quad (3.9)$$

The following Moser iteration lemma is given in [1, Proposition B.1]. See also [21] and [9].

Lemma 3.8. *Assume $N \geq 3$. Let $a(x)$ and $b(x)$ be functions on B_4 , and let $u \in H^1(B_4)$ be a weak solution to*

$$-\Delta u + a(x)u = b(x)u \quad \text{in } B_4. \quad (3.10)$$

Suppose that $a(x)$ and u satisfy that

$$a(x) \geq 0 \quad \text{for a.e. } x \in B_4, \quad (3.11)$$

and

$$\int_{B_4} a(x) |u(x)v(x)| dx < \infty \quad \text{for each } v \in H_0^1(B_4). \quad (3.12)$$

(i) Assume that for any $\varepsilon \in (0, 1)$, there exists $t_\varepsilon > 0$ such that

$$\|\chi_{[|b|>t_\varepsilon]} b\|_{L^{N/2}(B_4)} \leq \varepsilon,$$

where $[|b| > t] := \{x \in B_4 : |b(x)| > t\}$, and $\chi_A(x)$ denotes the characteristic function of $A \subset \mathbb{R}^N$. Then for any $r \in (0, \infty)$, there exists a constant $C(N, r, t_\varepsilon)$ such that

$$\| |u|^{r+1} \|_{H^1(B_1)} \leq C(N, r, t_\varepsilon) \|u\|_{L^{2^*}(B_4)}.$$

(ii) Let $s > N/2$ and assume that $b \in L^s(B_4)$. Then there exists a constant $C(N, s, \|b\|_{L^s(B_4)})$ such that

$$\|u\|_{L^\infty(B_1)} \leq C(N, s, \|b\|_{L^s(B_4)}) \|u\|_{L^{2^*}(B_4)}.$$

Here, the constants $C(N, r, t_\varepsilon)$ and $C(N, s, \|b\|_{L^s(B_4)})$ in (i) and (ii) remain bounded as long as r, t_ε and $\|b\|_{L^s(B_4)}$ are bounded.

4. PROOF OF THEOREM 2.1

In this section, we always assume that $2_\alpha := \frac{N+\alpha}{N}$, $q \in (2, 2 + \frac{4}{N})$ and $\varepsilon > 0$ is a large parameter. The existence and symmetry of ground states in this case follows from the upper bound estimate of the least energy given by Lemma 4.3 below and the argument used in [26]. See also [35, 30] and the references therein.

4.1. **The Case $q \in (2, 2 + \frac{4\alpha}{N(2+\alpha)})$.** In this case, we set

$$w(x) = \varepsilon^{-\frac{2N}{\alpha[4-N(q-2)]}} u(\varepsilon^{-\frac{N(q-2)}{\alpha[4-N(q-2)]}} x). \quad (4.1)$$

Then (P_ε) is reduced to

$$-\varepsilon^{-\sigma} \Delta w + w = (I_\alpha * |w|^{2\alpha}) |w|^{2\alpha-2} w + \varepsilon^{-\sigma-\tau(q-1)} g(\varepsilon^\tau w), \quad (4.2)$$

where

$$\sigma := \frac{4\alpha - N(2+\alpha)(q-2)}{\alpha[4-N(q-2)]} > 0, \quad \tau := \frac{2N}{\alpha[4-N(q-2)]}.$$

The corresponding functional is defined by

$$J_\varepsilon(w) : = \frac{\varepsilon^{-\sigma}}{2} \int_{\mathbb{R}^N} |\nabla w|^2 + \frac{1}{2} \int_{\mathbb{R}^N} |w|^2 - \frac{1}{22\alpha} \int_{\mathbb{R}^N} (I_\alpha * |w|^{2\alpha}) |w|^{2\alpha} - \varepsilon^{-\sigma-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau w).$$

Let $u_\varepsilon \in H^1(\mathbb{R}^N)$ be the ground state for (P_ε) and

$$w_\varepsilon(x) = \varepsilon^{-\frac{2N}{\alpha[4-N(q-2)]}} u_\varepsilon(\varepsilon^{-\frac{N(q-2)}{\alpha[4-N(q-2)]}} x). \quad (4.3)$$

Then w_ε is a ground state of (4.2).

To show that w_ε is bounded in $H^1(\mathbb{R}^N)$, we also make the following

$$w_\varepsilon(x) = \varepsilon^{\frac{N[4\alpha-N(2+\alpha)(q-2)]}{4\alpha[4-N(q-2)]}} v_\varepsilon(\varepsilon^{\frac{4\alpha-N(2+\alpha)(q-2)}{2\alpha[4-N(q-2)]}} x). \quad (4.4)$$

Then $v = v_\varepsilon$ is a ground state of the following equation

$$-\Delta v + v = (I_\alpha * |v|^{2\alpha}) |v|^{2\alpha-2} v + \varepsilon^{-\frac{4-N(q-2)}{4}} \sigma \varepsilon^{-\frac{N(2+\alpha)(q-1)}{4\alpha}} g(\varepsilon^{\frac{N(2+\alpha)}{4\alpha}} v). \quad (4.5)$$

The corresponding functional of (4.5) is defined by

$$\tilde{J}_\varepsilon(v) : = \frac{1}{2} \int_{\mathbb{R}^N} |\nabla v|^2 + |v|^2 - \frac{1}{22\alpha} \int_{\mathbb{R}^N} (I_\alpha * |v|^{2\alpha}) |v|^{2\alpha} - \varepsilon^{-\frac{4-N(q-2)}{4}} \sigma \varepsilon^{-\frac{N(2+\alpha)q}{4\alpha}} \int_{\mathbb{R}^N} G(\varepsilon^{\frac{N(2+\alpha)}{4\alpha}} v).$$

Clearly, the following lemma holds true.

Lemma 4.1. *Let $\varepsilon > 0$, and $w_\varepsilon, v_\varepsilon$ are given in (4.3) and (4.4) respectively. Then*

- (1) $\|w_\varepsilon\|_2^2 = \|v_\varepsilon\|_2^2, \quad \int_{\mathbb{R}^N} (I_\alpha * |w_\varepsilon|^{2\alpha}) |w_\varepsilon|^{2\alpha} = \int_{\mathbb{R}^N} (I_\alpha * |v_\varepsilon|^{2\alpha}) |v_\varepsilon|^{2\alpha},$
- (2) $\varepsilon^{-\sigma} \|\nabla w_\varepsilon\|_2^2 = \|\nabla v_\varepsilon\|_2^2, \quad \varepsilon^{-\frac{N\sigma(q-2)}{4}} \|w_\varepsilon\|_q^q = \|v_\varepsilon\|_q^q,$
- (3) $J_\varepsilon(w_\varepsilon) = \tilde{J}_\varepsilon(v_\varepsilon).$

As $\varepsilon \rightarrow \infty$, it follows from (H2) that the formal limit equation of (4.5) is as follows

$$-\Delta v + v = (I_\alpha * |v|^{2\alpha}) |v|^{2\alpha-2} v. \quad (4.6)$$

Let

$$\tilde{J}_\infty(v) := \frac{1}{2} \int_{\mathbb{R}^N} |\nabla v|^2 + |v|^2 - \frac{1}{22\alpha} \int_{\mathbb{R}^N} (I_\alpha * |v|^{2\alpha}) |v|^{2\alpha},$$

and

$$\tilde{\mathcal{M}}_\infty := \left\{ v \in H^1(\mathbb{R}^N) \setminus \{0\} \mid \int_{\mathbb{R}^N} |\nabla v|^2 + |v|^2 = \int_{\mathbb{R}^N} |v|^{2\alpha} \right\}.$$

Then, from a minimax characterizations for \tilde{J}_∞ similar to (4.9) below, it is easy to see that

$$\tilde{J}_\varepsilon(v_\varepsilon) \leq \inf_{v \in H^1(\mathbb{R}^N) \setminus \{0\}} \sup_{t>0} \tilde{J}_\infty(tv) = \inf_{v \in \tilde{\mathcal{M}}_\infty} \tilde{J}_\infty(v) < \infty.$$

Since v_ε satisfies the Pohožaev identity, we obtain

$$\frac{1}{2^*} \int_{\mathbb{R}^N} |\nabla v_\varepsilon|^2 + \frac{1}{2} \int_{\mathbb{R}^N} |v_\varepsilon|^2 = \frac{1}{2} \int_{\mathbb{R}^N} (I_\alpha * |v_\varepsilon|^{2\alpha}) |v_\varepsilon|^{2\alpha} + \varepsilon^{-\frac{4-N(q-2)}{4}\sigma} \varepsilon^{-\frac{N(2+\alpha)q}{4\alpha}} \int_{\mathbb{R}^N} G(\varepsilon^{\frac{N(2+\alpha)}{4\alpha}} v). \quad (4.7)$$

Therefore, we have

$$\tilde{J}_\varepsilon(v_\varepsilon) = \frac{1}{N} \int_{\mathbb{R}^N} |\nabla v_\varepsilon|^2 + \frac{2\alpha - 1}{22\alpha} \int_{\mathbb{R}^N} (I_\alpha * |v_\varepsilon|^{2\alpha}) |v_\varepsilon|^{2\alpha},$$

and hence $\|\nabla v_\varepsilon\|_2$ and $\int_{\mathbb{R}^N} (I_\alpha * |v_\varepsilon|^{2\alpha}) |v_\varepsilon|^{2\alpha}$ are bounded. Thus, by (4.7), (H2) and the Gagliardo-Nirenberg inequality, we obtain

$$\begin{aligned} \|v_\varepsilon\|_2^2 &\leq \frac{2}{2^*} \|\nabla v_\varepsilon\|_2^2 + \|v_\varepsilon\|_2^2 \\ &= \int_{\mathbb{R}^N} (I_\alpha * |v_\varepsilon|^{2\alpha}) |v_\varepsilon|^{2\alpha} + 2\varepsilon^{-\frac{4-N(q-2)}{4}\sigma} \varepsilon^{-\frac{N(2+\alpha)q}{4\alpha}} \int_{\mathbb{R}^N} G(\varepsilon^{\frac{N(2+\alpha)}{4\alpha}} v) \\ &\leq C + \varepsilon^{-\frac{4-N(q-2)}{4}\sigma} \left(\frac{A}{q} \int_{\mathbb{R}^N} |v_\varepsilon|^q + o_\varepsilon(1) \right) \\ &\leq C + C\varepsilon^{-\frac{4-N(q-2)}{4}\sigma} \|\nabla v_\varepsilon\|_2^{\frac{N(q-2)}{2}} \|v_\varepsilon\|_2^{\frac{2N-q(N-2)}{2}}, \end{aligned}$$

which together with the fact that $\|\nabla v_\varepsilon\|_2$ is bounded and $\frac{2N-q(N-2)}{2} < 2$ yields $\|v_\varepsilon\|_2$ is bounded. Thus, by Lemma 4.1, it follows that w_ε is bounded in $L^2(\mathbb{R}^N)$.

It is also easy to see from Lemma 4.1 that $I_\varepsilon(u) = \varepsilon^{\frac{N+\alpha}{\alpha}} J_\varepsilon(w) = \varepsilon^{\frac{N+\alpha}{\alpha}} \tilde{J}_\varepsilon(v)$ and the energy of the ground states given by

$$m_\varepsilon = \inf_{u \in \mathcal{N}_\varepsilon} I_\varepsilon(u) = \varepsilon^{\frac{N+\alpha}{\alpha}} \inf_{w \in \mathcal{M}_\varepsilon} J_\varepsilon(w) = \varepsilon^{\frac{N+\alpha}{\alpha}} \inf_{v \in \tilde{\mathcal{M}}_\varepsilon} \tilde{J}_\varepsilon(v)$$

are well-defined and positive, where \mathcal{M}_ε and $\tilde{\mathcal{M}}_\varepsilon$ denote the corresponding Nehari manifolds

$$\begin{aligned} \mathcal{M}_\varepsilon &:= \left\{ w \in H^1(\mathbb{R}^N) \setminus \{0\} \mid \begin{aligned} &\int_{\mathbb{R}^N} \varepsilon^{-\sigma} |\nabla w|^2 + |w|^2 = \int_{\mathbb{R}^N} (I_\alpha * |w|^{2\alpha}) |w|^{2\alpha} \\ &+ \varepsilon^{-\sigma-\tau(q-1)} \int_{\mathbb{R}^N} g(\varepsilon^\tau w) w \end{aligned} \right\}, \\ \tilde{\mathcal{M}}_\varepsilon &:= \left\{ v \in H^1(\mathbb{R}^N) \setminus \{0\} \mid \begin{aligned} &\int_{\mathbb{R}^N} |\nabla v|^2 + |v|^2 = \int_{\mathbb{R}^N} (I_\alpha * |v|^{2\alpha}) |v|^{2\alpha} \\ &+ \varepsilon^{-\frac{4-N(q-2)}{4}\sigma} \varepsilon^{-\frac{N(2+\alpha)(q-1)}{4\alpha}} \int_{\mathbb{R}^N} g(\varepsilon^{\frac{N(2+\alpha)}{4\alpha}} v) v \end{aligned} \right\}. \end{aligned}$$

Define the Pohožaev manifold as follows

$$\mathcal{P}_\varepsilon := \{w \in H^1(\mathbb{R}^N) \setminus \{0\} \mid P_\varepsilon(w) = 0\},$$

where

$$\begin{aligned} P_\varepsilon(w) : &= \frac{\varepsilon^{-\sigma}(N-2)}{2} \int_{\mathbb{R}^N} |\nabla w|^2 + \frac{N}{2} \int_{\mathbb{R}^N} |w|^2 \\ &\quad - \frac{N+\alpha}{22\alpha} \int_{\mathbb{R}^N} (I_\alpha * |w|^{2\alpha}) |w|^{2\alpha} - N\varepsilon^{-\sigma-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau w). \end{aligned} \quad (4.8)$$

Then by Lemma 3.1, $w_\varepsilon \in \mathcal{P}_\varepsilon$. Moreover, we have the following minimax characterizations for the least energy level m_ε .

$$\varepsilon^{-\frac{N+\alpha}{\alpha}} m_\varepsilon = \tilde{m}_\varepsilon := \inf_{w \in H^1(\mathbb{R}^N) \setminus \{0\}} \sup_{t \geq 0} J_\varepsilon(tw) = \inf_{w \in H^1(\mathbb{R}^N) \setminus \{0\}} \sup_{t \geq 0} J_\varepsilon(w_t). \quad (4.9)$$

In particular, we have $\varepsilon^{-\frac{N+\alpha}{\alpha}} m_\varepsilon = \tilde{m}_\varepsilon = J_\varepsilon(w_\varepsilon) = \sup_{t>0} J_\lambda(tw_\varepsilon) = \sup_{t>0} J_\varepsilon((w_\varepsilon)_t)$.

Lemma 4.2. *The rescaled family of solutions $\{w_\varepsilon\}$ is bounded in $H^1(\mathbb{R}^N)$, and*

$$\|\nabla w_\varepsilon\|_2^2 = \frac{NA(q-2)}{2q} \|w_\varepsilon\|_q^q + o_\varepsilon(1), \quad (4.10)$$

where and in what follows, $\lim_{\varepsilon \rightarrow \infty} o_\varepsilon(1) = 0$.

Proof. Since $\{w_\varepsilon\}$ is bounded in $L^2(\mathbb{R}^N)$, it suffices to show that $\{w_\varepsilon\}$ is also bounded in $D^{1,2}(\mathbb{R}^N)$. By $w_\varepsilon \in \mathcal{M}_\varepsilon \cap \mathcal{P}_\varepsilon$, we obtain

$$\begin{aligned} \varepsilon^{-\sigma} \int_{\mathbb{R}^N} |\nabla w_\varepsilon|^2 + \int_{\mathbb{R}^N} |w_\varepsilon|^2 &= \int_{\mathbb{R}^N} (I_\alpha * |w_\varepsilon|^{2\alpha}) |w_\varepsilon|^{2\alpha} + \varepsilon^{-\sigma-\tau(q-1)} \int_{\mathbb{R}^N} g(\varepsilon^\tau w_\varepsilon) w_\varepsilon, \\ \frac{\varepsilon^{-\sigma}}{2^*} \int_{\mathbb{R}^N} |\nabla w_\varepsilon|^2 + \frac{1}{2} \int_{\mathbb{R}^N} |w_\varepsilon|^2 &= \frac{N+\alpha}{2Np} \int_{\mathbb{R}^N} (I_\alpha * |w_\varepsilon|^{2\alpha}) |w_\varepsilon|^{2\alpha} + \varepsilon^{-\sigma-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau w_\varepsilon). \end{aligned}$$

Therefore, we have

$$\varepsilon^{-\sigma} \|\nabla w_\varepsilon\|_2^2 = N \varepsilon^{-\sigma-\tau q} \int_{\mathbb{R}^N} \left[\frac{1}{2} g(\varepsilon^\tau w_\varepsilon) \varepsilon^\tau w_\varepsilon - G(\varepsilon^\tau w_\varepsilon) \right]. \quad (4.11)$$

Set $g(s) = As^{q-1} + \tilde{g}(s)$ for $s \geq 0$, then $G(s) = \frac{A}{q}s^q + \tilde{G}(s)$, and

$$\lim_{s \rightarrow \infty} \tilde{g}(s)s^{1-q} = \lim_{s \rightarrow \infty} \tilde{G}(s)s^{-q} = 0.$$

Therefore, for any $\delta > 0$ there exists a constant $C_\delta > 0$ such that

$$\left| \frac{1}{2} \tilde{g}(u)u - \tilde{G}(u) \right| \leq \delta |u|^q + C_\delta |u|^2,$$

and hence

$$\begin{aligned} \varepsilon^{-\tau q} \int_{\mathbb{R}^N} \left| \frac{1}{2} \tilde{g}(\varepsilon^\tau w_\varepsilon) \varepsilon^\tau w_\varepsilon - \tilde{G}(\varepsilon^\tau w_\varepsilon) \right| &\leq \delta \int_{\mathbb{R}^N} |w_\varepsilon|^q + C_\delta \varepsilon^{-\tau(q-2)} \int_{\mathbb{R}^N} |w_\varepsilon|^2 \\ &\leq \delta \|w_\varepsilon\|_q^q + CC_\delta \varepsilon^{-\tau(q-2)}. \end{aligned} \quad (4.12)$$

By the Gagliardo-Nirenberg Inequality, we obtain

$$\|w_\varepsilon\|_q^q \leq C \|\nabla w_\varepsilon\|_2^{\frac{N(q-2)}{2}} \|w_\varepsilon\|_2^{\frac{2N-q(N-2)}{2}} \leq \tilde{C} \|\nabla w_\varepsilon\|_2^{\frac{N(q-2)}{2}}.$$

Therefore, we get

$$\begin{aligned} \|\nabla w_\varepsilon\|_2^2 &= N \varepsilon^{-\tau q} \int_{\mathbb{R}^N} \left[\frac{1}{2} g(\varepsilon^\tau w_\varepsilon) \varepsilon^\tau w_\varepsilon - G(\varepsilon^\tau w_\varepsilon) \right] \\ &\leq \left(\frac{NA(q-2)}{2q} + \delta \right) \int_{\mathbb{R}^N} |w_\varepsilon|^q + C \\ &\leq \tilde{C} \left(\frac{NA(q-2)}{2q} + \delta \right) \|\nabla w_\varepsilon\|_2^{\frac{N(q-2)}{2}} + C, \end{aligned}$$

which together with $2 > \frac{N(q-2)}{2}$ implies that $\{w_\varepsilon\}$ is bounded in $D^{1,2}(\mathbb{R}^N)$.

By virtue of (4.12), for any $\delta > 0$, we have

$$\limsup_{\varepsilon \rightarrow \infty} \varepsilon^{-\tau q} \int_{\mathbb{R}^N} \left| \frac{1}{2} \tilde{g}(\varepsilon^\tau w_\varepsilon) \varepsilon^\tau w_\varepsilon - \tilde{G}(\varepsilon^\tau w_\varepsilon) \right| \leq C\delta,$$

and hence

$$\lim_{\varepsilon \rightarrow \infty} \varepsilon^{-\tau q} \int_{\mathbb{R}^N} \left[\frac{1}{2} \tilde{g}(\varepsilon^\tau w_\varepsilon) \varepsilon^\tau w_\varepsilon - \tilde{G}(\varepsilon^\tau w_\varepsilon) \right] = 0. \quad (4.13)$$

Clearly, (4.11) implies that

$$\|\nabla w_\varepsilon\|_2^2 = \frac{NA(q-2)}{2q} \|w_\varepsilon\|_q^q + N \varepsilon^{-\tau q} \int_{\mathbb{R}^N} \left[\frac{1}{2} \tilde{g}(\varepsilon^\tau w_\varepsilon) \varepsilon^\tau w_\varepsilon - \tilde{G}(\varepsilon^\tau w_\varepsilon) \right],$$

the equality (4.10) then follows from (4.13). The proof is complete. \square

As $\varepsilon \rightarrow \infty$, the limit equation of (4.2) is the Hardy-Littlewood-Sobolev critical equation $w = (I_\alpha * |w|^{2\alpha})|w|^{2\alpha-2}w$, the corresponding functional is defined by

$$J_\infty(w) := \int_{\mathbb{R}^N} |w|^2 - \frac{1}{22\alpha} \int_{\mathbb{R}^N} (I_\alpha * |w|^{2\alpha})|w|^{2\alpha}.$$

We define the Nehari manifolds as follows.

$$\mathcal{M}_\infty = \left\{ w \in H^1(\mathbb{R}^N) \setminus \{0\} \mid \int_{\mathbb{R}^N} |w|^2 = \int_{\mathbb{R}^N} (I_\alpha * |w|^{2\alpha})|w|^{2\alpha} \right\}.$$

Then

$$\tilde{m}_\infty := \inf_{w \in \mathcal{M}_\infty} J_\infty(w)$$

is well-defined and positive. Moreover, as in (4.9), a similar minimax characterizations for the least energy level \tilde{m}_∞ also holds, and J_∞ is attained on \mathcal{M}_∞ and $\tilde{m}_\infty = \frac{\alpha}{2(N+\alpha)} S_1^{\frac{N+\alpha}{\alpha}}$.

For $w \in H^1(\mathbb{R}^N) \setminus \{0\}$, we set

$$\tau_1(w) = \frac{\int_{\mathbb{R}^N} |w|^2}{\int_{\mathbb{R}^N} (I_\alpha * |w|^{2\alpha})|w|^{2\alpha}}. \quad (4.14)$$

Then $(\tau_1(w))^{\frac{N}{2\alpha}} w \in \mathcal{M}_\infty$ for any $w \in H^1(\mathbb{R}^N) \setminus \{0\}$, and $w \in \mathcal{M}_\infty$ if and only if $\tau_1(w) = 1$.

Now, we give the following estimation on $\tau_1(w_\varepsilon)$ and the least energy \tilde{m}_ε :

Lemma 4.3. *Let $N \geq 3$ and $q \in (2, 2 + \frac{4}{N})$. Then $\tau_1(w_\varepsilon) = 1 + O(\varepsilon^{-\sigma})$ and*

$$\frac{\alpha}{2(N+\alpha)} S_1^{\frac{N+\alpha}{\alpha}} - \tilde{m}_\varepsilon = \tilde{m}_\infty - \tilde{m}_\varepsilon \sim \varepsilon^{-\sigma}, \quad (4.15)$$

as $\varepsilon \rightarrow \infty$.

Proof. For any $\delta > 0$ there exists a constant $C_\delta > 0$ such that

$$g(s)s \leq C_\delta |s|^q + \delta |s|^2, \quad \text{for all } s \geq 0.$$

By the Gagliardo-Nirenberg inequality and the fact that $\frac{N(q-2)}{2} < 2$, it follows that

$$\begin{aligned} & \frac{\varepsilon^{-\tau q} \int_{\mathbb{R}^N} g(\varepsilon^\tau w_\varepsilon) \varepsilon^\tau w_\varepsilon - \int_{\mathbb{R}^N} |\nabla w_\varepsilon|^2}{\int_{\mathbb{R}^N} |w_\varepsilon|^2} \\ & \leq \frac{C_\delta \int_{\mathbb{R}^N} |w_\varepsilon|^q - \int_{\mathbb{R}^N} |\nabla w_\varepsilon|^2}{\int_{\mathbb{R}^N} |w_\varepsilon|^2} + \delta \varepsilon^{-\tau(q-2)} \\ & \leq C_\delta \left(\frac{\|\nabla w_\varepsilon\|_2}{\|w_\varepsilon\|_2} \right)^{\frac{N(q-2)}{2}} \|w_\varepsilon\|_2^{q-2} - \left(\frac{\|\nabla w_\varepsilon\|_2}{\|w_\varepsilon\|_2} \right)^2 + \delta \varepsilon^{-\tau(q-2)} \\ & \leq C C_\delta \left(\frac{\|\nabla w_\varepsilon\|_2}{\|w_\varepsilon\|_2} \right)^{\frac{N(q-2)}{2}} - \left(\frac{\|\nabla w_\varepsilon\|_2}{\|w_\varepsilon\|_2} \right)^2 + \delta \varepsilon^{-\tau(q-2)} \\ & \leq C < \infty. \end{aligned}$$

First, since $v_\varepsilon \in \mathcal{M}_\varepsilon$, by (4.14), it follows that

$$\begin{aligned} \tau_1(w_\varepsilon) &= \frac{\int_{\mathbb{R}^N} |w_\varepsilon|^2}{\int_{\mathbb{R}^N} (I_\alpha * |w_\varepsilon|^p) |w_\varepsilon|^p} \\ &= \frac{\int_{\mathbb{R}^N} |w_\varepsilon|^2}{\int_{\mathbb{R}^N} |w_\varepsilon|^{2-\varepsilon^{-\sigma}} \left\{ \varepsilon^{-\tau q} \int_{\mathbb{R}^N} g(\varepsilon^\tau w_\varepsilon) \varepsilon^\tau w_\varepsilon - \int_{\mathbb{R}^N} |\nabla w_\varepsilon|^2 \right\}} \\ &\leq 1 + C \varepsilon^{-\sigma}, \end{aligned}$$

from which we get

$$\int_{\mathbb{R}^N} |w_\varepsilon|^2 = \tau_1(w_\varepsilon) \int_{\mathbb{R}^N} (I_\alpha * |w_\varepsilon|^{2\alpha}) |w_\varepsilon|^{2\alpha} \leq C \tau_1(w_\varepsilon) \left(\int_{\mathbb{R}^N} |w_\varepsilon|^2 \right)^{2\alpha},$$

and hence

$$\int_{\mathbb{R}^N} |w_\varepsilon|^2 \geq (C\tau_1(w_\varepsilon))^{-\frac{N}{\alpha}} \geq C > 0. \quad (4.16)$$

By Lemma 4.1 and the Sobolev inequality, we have

$$\begin{aligned} \tau_1(w_\varepsilon) &= \frac{\int_{\mathbb{R}^N} |w_\varepsilon|^2}{\int_{\mathbb{R}^N} (I_\alpha * |w_\varepsilon|^{2\alpha}) |w_\varepsilon|^{2\alpha}} \\ &= \frac{\int_{\mathbb{R}^N} |w_\varepsilon|^2}{\int_{\mathbb{R}^N} |w_\varepsilon|^{2-\varepsilon-\sigma} \left\{ \varepsilon^{-\tau q} \int_{\mathbb{R}^N} g(\varepsilon^\tau w_\varepsilon) \varepsilon^\tau w_\varepsilon - \int_{\mathbb{R}^N} |\nabla w_\varepsilon|^2 \right\}} \\ &\geq \frac{\int_{\mathbb{R}^N} |w_\varepsilon|^2}{\int_{\mathbb{R}^N} |w_\varepsilon|^{2+\varepsilon-\sigma} \int_{\mathbb{R}^N} |\nabla w_\varepsilon|^2} \\ &\geq 1 - \frac{1}{2} \varepsilon^{-\sigma} \frac{\|\nabla w_\varepsilon\|_2^2}{\|w_\varepsilon\|_2^2} \\ &\geq 1 - C \varepsilon^{-\sigma}, \end{aligned}$$

thus we conclude that $\tau_1(w_\varepsilon) = 1 + O(\varepsilon^{-\sigma})$ as $\varepsilon \rightarrow \infty$.

For $w \in H^1(\mathbb{R}^N)$, let w_t be defined as in Lemma 3.2. Then by Lemma 3.2 and Pohožaev's identity, it is easy to show that $\sup_{t \geq 0} J_\varepsilon((w_\varepsilon)_t) = J_\varepsilon(w_\varepsilon) = \tilde{m}_\varepsilon$. Therefore, we get

$$\begin{aligned} \tilde{m}_\infty &\leq \sup_{t \geq 0} J_\infty((w_\varepsilon)_t) = J_\infty((w_\varepsilon)_t)|_{t=\tau_1((w_\varepsilon))^{1/\alpha}} \\ &\leq \sup_{t \geq 0} J_\varepsilon((w_\varepsilon)_t) + \varepsilon^{-\sigma-\tau q} (\tau_1(w_\varepsilon))^{N/\alpha} \int_{\mathbb{R}^N} G(\varepsilon^\tau w_\varepsilon) \\ &\leq \tilde{m}_\varepsilon + \varepsilon^{-\sigma} (1 + O(\varepsilon^{-\sigma}))^{N/\alpha} \left(\frac{A}{q} \|v_\varepsilon\|_q^q + o_\varepsilon(1) \right) \\ &= \tilde{m}_\varepsilon + O(\varepsilon^{-\sigma}). \end{aligned}$$

For each $\rho > 0$, the family $U_\rho(x) := \rho^{-\frac{N}{2}} U_1(x/\rho)$ are radial ground states of $v = (I_\alpha * |v|^{2\alpha}) v^{2\alpha-1}$, and verify that

$$\|\nabla U_\rho\|_2^2 = \rho^{-2} \|\nabla U_1\|_2^2, \quad \int_{\mathbb{R}^N} |U_\rho|^q = \rho^{N-\frac{N}{2}q} \int_{\mathbb{R}^N} |U_1|^q. \quad (4.17)$$

Let $g_0(\rho) = \frac{A}{q} \int_{\mathbb{R}^N} |U_\rho|^q - \frac{1}{2} \int_{\mathbb{R}^N} |\nabla U_\rho|^2$. Then

$$\begin{aligned} g_0(\rho) &= \frac{A}{q} \rho^{N-\frac{N}{2}q} \int_{\mathbb{R}^N} |U_1|^q - \frac{1}{2} \rho^{-2} \int_{\mathbb{R}^N} |\nabla U_1|^2 \\ &= \rho^{-2} \left\{ \frac{A}{q} \rho^{\frac{4-N(q-2)}{2}} \int_{\mathbb{R}^N} |U_1|^q - \frac{1}{2} \int_{\mathbb{R}^N} |\nabla U_1|^2 \right\}. \end{aligned}$$

Since $q \in (2, 2 + \frac{4}{N})$, we have $\lim_{\rho \rightarrow 0} g_0(\rho) = -\infty$, $\lim_{\rho \rightarrow +\infty} g_0(\rho) = 0$. It is easy to see that there exists $\rho_0 = \rho(q) \in (0, +\infty)$ with

$$\rho_0 = \left(\frac{2q \int_{\mathbb{R}^N} |\nabla U_1|^2}{NA(q-2) \int_{\mathbb{R}^N} |U_1|^q} \right)^{\frac{2}{4-N(q-2)}}$$

such that $g'_0(\rho_0) = 0$ and hence

$$g_0(\rho_0) = \sup_{\rho > 0} g_0(\rho) = \frac{4-N(q-2)}{2N(q-2)} \left(\frac{NA(q-2) \int_{\mathbb{R}^N} |U_1|^q}{2q \int_{\mathbb{R}^N} |\nabla U_1|^2} \right)^{\frac{4}{4-N(q-2)}} \int_{\mathbb{R}^N} |\nabla U_1|^2.$$

Let $U_0 = U_{\rho_0}$, then there exists $t_\varepsilon \in (0, +\infty)$ such that

$$\begin{aligned} \tilde{m}_\varepsilon &\leq \sup_{t \geq 0} J_\varepsilon((U_0)_t) = J_\varepsilon((U_0)_{t_\varepsilon}) \\ &\leq \sup_{t > 0} \left\{ \frac{t^N}{2} \int_{\mathbb{R}^N} |U_0|^2 - \frac{t^{N+\alpha}}{22\alpha} \int_{\mathbb{R}^N} (I_\alpha * |U_0|^{2\alpha}) |U_0|^{2\alpha} \right\} \\ &\quad - \varepsilon^{-\sigma} \left\{ \varepsilon^{-\tau q} t_\varepsilon^N \int_{\mathbb{R}^N} G(\varepsilon^\tau U_0) - \frac{1}{2} t_\varepsilon^{N-2} \int_{\mathbb{R}^N} |\nabla U_0|^2 \right\} \\ &\leq \tilde{m}_\infty - \varepsilon^{-\sigma} \left\{ t_\varepsilon^N \int_{\mathbb{R}^N} \left[\frac{A}{q} |U_0|^q + \varepsilon^{-\tau q} \tilde{G}(\varepsilon^\tau U_0) \right] - \frac{1}{2} t_\varepsilon^{N-2} \int_{\mathbb{R}^N} |\nabla U_0|^2 \right\}. \end{aligned} \quad (4.18)$$

where $t_\varepsilon \in (0, +\infty)$ satisfies

$$\frac{1}{2^*} t_\varepsilon^{N-2} \varepsilon^{-\sigma} \int_{\mathbb{R}^N} |\nabla U_0|^2 + \frac{1}{2} t_\varepsilon^N \int_{\mathbb{R}^N} |U_0|^2 = \frac{1}{2} t_\varepsilon^{N+\alpha} \int_{\mathbb{R}^N} (I_\alpha * |U_0|^{2\alpha}) |U_0|^{2\alpha} + t_\varepsilon^N \varepsilon^{-\sigma-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau U_0).$$

If $t_\varepsilon \geq 1$, then

$$\frac{1}{2^*} \varepsilon^{-\sigma} \int_{\mathbb{R}^N} |\nabla U_0|^2 + \frac{1}{2} \int_{\mathbb{R}^N} |U_0|^2 \geq \frac{1}{2} t_\varepsilon^\alpha \int_{\mathbb{R}^N} (I_\alpha * |U_0|^{2\alpha}) |U_0|^{2\alpha} + \varepsilon^{-\sigma-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau U_0).$$

Hence

$$t_\varepsilon^\alpha \leq \frac{\frac{1}{2^*} \varepsilon^{-\sigma} \int_{\mathbb{R}^N} |\nabla U_0|^2 + \frac{1}{2} \int_{\mathbb{R}^N} |U_0|^2 - \varepsilon^{-\sigma-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau U_0)}{\frac{1}{2} \int_{\mathbb{R}^N} (I_\alpha * |U_0|^{2\alpha}) |U_0|^{2\alpha}},$$

which implies that $\limsup_{\varepsilon \rightarrow \infty} t_\varepsilon^\alpha \leq 1$.

If $t_\varepsilon \leq 1$, then a similar argument yields that

$$t_\varepsilon^\alpha \geq \frac{\frac{1}{2^*} \varepsilon^{-\sigma} \int_{\mathbb{R}^N} |\nabla U_0|^2 + \frac{1}{2} \int_{\mathbb{R}^N} |U_0|^2 - \varepsilon^{-\sigma-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau U_0)}{\frac{1}{2} \int_{\mathbb{R}^N} (I_\alpha * |U_0|^{2\alpha}) |U_0|^{2\alpha}},$$

which implies that $\liminf_{\varepsilon \rightarrow \infty} t_\varepsilon^\alpha \geq 1$.

Therefore, $\lim_{\lambda \rightarrow 0} t_\lambda = 1$ and hence

$$\begin{aligned} \lim_{\varepsilon \rightarrow \infty} \left\{ t_\varepsilon^N \int_{\mathbb{R}^N} \left[\frac{A}{q} |U_0|^q + \varepsilon^{-\tau q} \tilde{G}(\varepsilon^\tau U_0) \right] - \frac{1}{2} t_\varepsilon^{N-2} \int_{\mathbb{R}^N} |\nabla U_0|^2 \right\} \\ = \frac{A}{q} \int_{\mathbb{R}^N} |U_0|^q - \frac{1}{2} \int_{\mathbb{R}^N} |\nabla U_0|^2 = g_0(\rho_0). \end{aligned}$$

Thus, there exists a constant $C > 0$ such that

$$\tilde{m}_\varepsilon \leq \tilde{m}_\infty - C \varepsilon^{-\sigma},$$

for large $\varepsilon > 0$. The proof is complete. \square

Let

$$\mathbb{D}(u) := \int_{\mathbb{R}^N} (I_\alpha * |u|^{\frac{N+\alpha}{N}}) |u|^{\frac{N+\alpha}{N}}.$$

The following result is a special case of the classical Brezis-Lieb lemma [6] for Riesz potentials, for a proof, we refer the reader to [29, Lemma 2.4].

Lemma 4.4. *Let $N \in \mathbb{N}, \alpha \in (0, N)$, and $(w_n)_{n \in \mathbb{N}}$ be a bounded sequence in $L^2(\mathbb{R}^N)$. If $w_n \rightarrow w$ almost everywhere on \mathbb{R}^N as $n \rightarrow \infty$, then*

$$\lim_{n \rightarrow \infty} \mathbb{D}(w_n) - \mathbb{D}(w_n - w_0) = \mathbb{D}(w_0).$$

Lemma 4.5. $\|w_\varepsilon\|_2^2 \sim \mathbb{D}(w_\varepsilon) \sim \|\nabla w_\varepsilon\|_2^2 \sim \|w_\varepsilon\|_q^q \sim 1$ as $\varepsilon \rightarrow \infty$.

Proof. By the definition of $\tau_1(w_\varepsilon)$, Lemma 4.3 and the Hardy-Sobolev-Littlewood inequality, for large $\varepsilon > 0$, we have

$$\|w_\varepsilon\|_2^2 = \tau_1(w_\varepsilon) \int_{\mathbb{R}^N} (I_\alpha * |w_\varepsilon|^{2\alpha}) |w_\varepsilon|^{2\alpha} \leq 2S_1^{-2\alpha} \|w_\varepsilon\|_2^{22\alpha},$$

and thus it follows that

$$\|w_\varepsilon\|_2^2 \geq 2^{-\frac{N}{\alpha}} S_1^{\frac{N+\alpha}{\alpha}},$$

which together with the boundedness of w_ε implies that $\|w_\varepsilon\|_2^2 \sim 1$ as $\varepsilon \rightarrow \infty$.

It follows from Lemma 4.3 that

$$\begin{aligned}\tilde{m}_\infty &\leq J_\infty((\tau_1(w_\varepsilon))^{\frac{N}{2\alpha}} w_\varepsilon) \\ &= \frac{1}{2}(\tau_1(w_\varepsilon))^{\frac{N}{\alpha}} \|w_\varepsilon\|_2^2 - \frac{1}{22\alpha}(\tau_1(w_\varepsilon))^{\frac{N+\alpha}{\alpha}} \int_{\mathbb{R}^N} (I_\alpha * |w_\varepsilon|^{2\alpha}) |w_\varepsilon|^{2\alpha} \\ &\leq (\tau_1(w_\varepsilon))^{\frac{N}{\alpha}} \left[\frac{1}{2} \|w_\varepsilon\|_2^2 - \frac{1}{22\alpha} \int_{\mathbb{R}^N} (I_\alpha * |w_\varepsilon|^{2\alpha}) |w_\varepsilon|^{2\alpha} \right].\end{aligned}\quad (4.19)$$

Since $w_\varepsilon \in \mathcal{N}_\varepsilon$, by Lemma 4.4, we obtain

$$\begin{aligned}\tau_1(w_\varepsilon) &= \frac{\int_{\mathbb{R}^N} |w_\varepsilon|^2}{\int_{\mathbb{R}^N} (I_\alpha * |w_\varepsilon|^{2\alpha}) |w_\varepsilon|^{2\alpha}} \\ &= \frac{\int_{\mathbb{R}^N} |w_\varepsilon|^2}{\int_{\mathbb{R}^N} |w_\varepsilon|^{2+\varepsilon^{-\sigma}} (\int_{\mathbb{R}^N} |\nabla w_\varepsilon|^2 - \varepsilon^{-\tau q} \int_{\mathbb{R}^N} g(\varepsilon^\tau w_\varepsilon) \varepsilon^\tau w_\varepsilon)} \\ &= \frac{\int_{\mathbb{R}^N} |w_\varepsilon|^2}{\int_{\mathbb{R}^N} |w_\varepsilon|^{2-\varepsilon^{-\sigma} \frac{2N-q(N-2)}{2q}} (\int_{\mathbb{R}^N} |w_\varepsilon|^q + o_\varepsilon(1))} \\ &\leq 1 + C_1 \varepsilon^{-\sigma} \|w_\varepsilon\|_q^q.\end{aligned}\quad (4.20)$$

Therefore, by (4.10), (4.19) and (4.20), we obtain

$$\begin{aligned}\tilde{m}_\varepsilon &= \frac{1}{2} \int_{\mathbb{R}^N} \varepsilon^{-\sigma} |\nabla w_\varepsilon|^2 + |w_\varepsilon|_2^2 - \frac{1}{22\alpha} \int_{\mathbb{R}^N} (I_\alpha * |w_\varepsilon|^{2\alpha}) |w_\varepsilon|^{2\alpha} - \varepsilon^{-\sigma-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau w_\varepsilon) \\ &\geq \varepsilon^{-\sigma} \left(\frac{1}{2} \|\nabla w_\varepsilon\|_2^2 - \frac{A}{q} \|w_\varepsilon\|_q^q + o_\varepsilon(1) \right) + \tilde{m}_\infty (\tau_1(w_\varepsilon))^{-\frac{N}{\alpha}} \\ &\geq \varepsilon^{-\sigma} \frac{[N(q-2)-4]A}{4q} (\|w_\varepsilon\|_q^q + o_\varepsilon(1)) + \tilde{m}_\infty - C_1 \varepsilon^{-\sigma} \|w_\varepsilon\|_q^q.\end{aligned}$$

By Lemma 4.3, we have $\tilde{m}_\infty - \tilde{m}_\varepsilon \geq C_2 \varepsilon^{-\sigma}$. Hence, we get

$$\frac{[4 - N(q-2)]A}{4q} \|w_\varepsilon\|_q^q \geq C_2 - C_1 \|w_\varepsilon\|_q^q,$$

which yields

$$\|w_\varepsilon\|_q^q \geq \frac{4qC_2}{[4 - N(q-2)]A + 4qC_1} > 0.$$

Since w_ε is bounded in $H^1(\mathbb{R}^N)$, it follows that $\|w_\varepsilon\|_q^q \sim 1$ as $\varepsilon \rightarrow \infty$.

Since $\|w_\varepsilon\|_2^2 \sim 1$ as $\varepsilon \rightarrow \infty$, the Gagliardo-Nirenberg inequality implies

$$\|w_\varepsilon\|_q^q \leq C \|\nabla w_\varepsilon\|_2^{\frac{N(q-2)}{2}},$$

which together with the boundedness of w_ε in $H^1(\mathbb{R}^N)$ yields $\|\nabla w_\varepsilon\|_2^2 \sim 1$ as $\varepsilon \rightarrow \infty$.

Finally, by the definition of $\tau_1(w_\varepsilon)$, Lemma 4.3 and Lemma 4.4, it follows that

$$\mathbb{D}(w_\varepsilon) = \int_{\mathbb{R}^N} (I_\alpha * |w_\varepsilon|^{\frac{N+\alpha}{N}}) |w_\varepsilon|^{\frac{N+\alpha}{N}} = (\tau_1(w_\varepsilon))^{-1} \|w_\varepsilon\|_2^2 \sim 1,$$

as $\varepsilon \rightarrow \infty$. The proof is complete. \square

Lemma 4.6. *Let $N \geq 3$ and $q \in (2, 2 + \frac{4}{N})$, then for any $\varepsilon_n \rightarrow \infty$, there exists $\rho \in [\rho_0, +\infty)$, such that, up to a subsequence, $w_{\varepsilon_n} \rightarrow U_\rho$ in $L^2(\mathbb{R}^N)$, where*

$$\rho_0 = \rho_0(q) := \left(\frac{2q \int_{\mathbb{R}^N} |\nabla U_1|^2}{NA(q-2) \int_{\mathbb{R}^N} |U_1|^q} \right)^{\frac{2}{4-N(q-2)}}.$$

Moreover, $w_{\varepsilon_n} \rightarrow U_\rho$ in $D^{1,2}(\mathbb{R}^N)$ if and only if $\rho = \rho_0$.

Proof. The proof is similar to that in [27, Lemma 4.8], and for the reader's convenience, we give the detail proof. Note that w_{ε_n} is a positive radially symmetric function, and by Lemma 4.2, $\{w_{\varepsilon_n}\}$ is bounded in $H^1(\mathbb{R}^N)$. Then there exists $w_\infty \in H^1(\mathbb{R}^N)$ such that

$$w_{\varepsilon_n} \rightharpoonup w_\infty \quad \text{weakly in } H^1(\mathbb{R}^N), \quad w_{\varepsilon_n} \rightarrow w_\infty \quad \text{in } L^p(\mathbb{R}^N) \quad \text{for any } p \in (2, 2^*), \quad (4.21)$$

and

$$w_{\varepsilon_n}(x) \rightarrow w_\infty(x) \quad \text{a. e. on } \mathbb{R}^N, \quad w_{\varepsilon_n} \rightarrow w_\infty \quad \text{in } L^2_{loc}(\mathbb{R}^N). \quad (4.22)$$

Observe that

$$J_\infty(w_{\varepsilon_n}) = J_{\varepsilon_n}(w_{\varepsilon_n}) + \varepsilon_n^{-\sigma-\tau q} \int_{\mathbb{R}^N} G(\varepsilon_n^\tau w_{\varepsilon_n}) - \frac{\varepsilon_n^{-\sigma}}{2} \int_{\mathbb{R}^N} |\nabla w_{\varepsilon_n}|^2 = \tilde{m}_{\varepsilon_n} + o_n(1) = \tilde{m}_\infty + o_n(1),$$

and

$$J'_\infty(w_{\varepsilon_n})w = J'_{\varepsilon_n}(w_{\varepsilon_n})w + \varepsilon_n^{-\sigma-\tau(q-1)} \int_{\mathbb{R}^N} g(\varepsilon_n^\tau w_{\varepsilon_n})w - \varepsilon_n^{-\sigma} \int_{\mathbb{R}^N} \nabla w_{\varepsilon_n} \nabla w = o_n(1).$$

Therefore, $\{w_{\varepsilon_n}\}$ is a PS sequence of J_∞ at level $\tilde{m}_\infty = \frac{\alpha}{2(N+\alpha)} S_1^{\frac{N+\alpha}{\alpha}}$.

By Lemma 4.5, we have $w_\infty \neq 0$. By Lemma 4.3 and Lemma 4.4, we have

$$\begin{aligned} o_n(1) &= \tilde{m}_{\varepsilon_n} - \tilde{m}_\infty \\ &\geq \frac{\varepsilon_n^{-\sigma}}{2} \|\nabla w_{\varepsilon_n}\|_2^2 + \frac{1}{2} [\|w_{\varepsilon_n}\|_2^2 - \|w_0\|_2^2] - \frac{N}{2(N+\alpha)} [\mathbb{D}(w_{\varepsilon_n}) - \mathbb{D}(w_\infty)] \\ &\quad - \varepsilon_n^{-\sigma-\tau q} \int_{\mathbb{R}^N} G(\varepsilon_n^\tau w_{\varepsilon_n}) \\ &= \frac{1}{2} \|w_{\varepsilon_n} - w_\infty\|_2^2 - \frac{N}{2(N+\alpha)} \mathbb{D}(w_{\varepsilon_n} - w_\infty) + o_n(1), \\ 0 &= \langle J'_{\varepsilon_n}(w_{\varepsilon_n}) - J'_\infty(w_\infty), w_{\varepsilon_n} - w_\infty \rangle \\ &= \varepsilon_n^{-\sigma} \int_{\mathbb{R}^N} \nabla w_{\varepsilon_n} (\nabla w_{\varepsilon_n} - \nabla w_\infty) + \int_{\mathbb{R}^N} |w_{\varepsilon_n} - w_\infty|^2 \\ &\quad - \int_{\mathbb{R}^N} (I_\alpha * |w_{\varepsilon_n}|^{\frac{N+\alpha}{N}}) w_{\varepsilon_n}^\frac{\alpha}{N} (w_{\varepsilon_n} - w_\infty) + \int_{\mathbb{R}^N} (I_\alpha * |w_\infty|^{\frac{N+\alpha}{N}}) w_\infty^\frac{\alpha}{N} (w_{\varepsilon_n} - w_\infty) \\ &\quad - \varepsilon_n^{-\sigma} \int_{\mathbb{R}^N} |w_{\varepsilon_n}|^{q-2} w_{\varepsilon_n} (w_{\varepsilon_n} - w_\infty) \\ &= \|w_{\varepsilon_n} - w_0\|_2^2 - \mathbb{D}(w_{\varepsilon_n} - w_\infty) + o_n(1). \end{aligned}$$

Hence, it follows that

$$\|w_{\varepsilon_n} - w_\infty\|_2^2 \leq \frac{N}{N+\alpha} \mathbb{D}(w_{\varepsilon_n} - w_\infty) + o_n(1) = \frac{N}{N+\alpha} \|w_{\varepsilon_n} - w_\infty\|_2^2 + o_n(1),$$

and hence

$$\|w_{\varepsilon_n} - w_\infty\|_2 \rightarrow 0, \quad \text{as } \varepsilon_n \rightarrow \infty.$$

By the Hardy-Littlewood-Sobolev inequality and Lemma 4.4, it follows that

$$\lim_{\varepsilon_n \rightarrow \infty} \mathbb{D}(w_{\varepsilon_n}) = \mathbb{D}(w_\infty).$$

Since $\tau_1(w_{\varepsilon_n}) \rightarrow 1$ as $\varepsilon_n \rightarrow \infty$ by Lemma 4.3, it follows that $\tau_1(w_\infty) = 1$ and hence $w_\infty \in \mathcal{M}_\infty$.

On the other hand, by Lemma 4.1 and the boundedness of v_{ε_n} in $H^1(\mathbb{R}^N)$, we have

$$\begin{aligned} \tilde{m}_{\varepsilon_n} &= J_{\varepsilon_n}(w_{\varepsilon_n}) \\ &= \frac{1}{2} \int_{\mathbb{R}^N} \varepsilon_n^{-\sigma} |\nabla w_{\varepsilon_n}|^2 + |w_{\varepsilon_n}|^2 - \frac{1}{2p} \int_{\mathbb{R}^N} (I_\alpha * |w_{\varepsilon_n}|^{\frac{N+\alpha}{N}}) |w_{\varepsilon_n}|^{\frac{N+\alpha}{N}} \\ &\quad - \varepsilon_n^{-\sigma-\tau q} \int_{\mathbb{R}^N} G(\varepsilon_n^\tau w_{\varepsilon_n}) \\ &\geq \frac{1}{2} \int_{\mathbb{R}^N} |w_{\varepsilon_n}|^2 - \frac{1}{2p} \int_{\mathbb{R}^N} (I_\alpha * |w_{\varepsilon_n}|^{\frac{N+\alpha}{N}}) |w_{\varepsilon_n}|^{\frac{N+\alpha}{N}} \\ &\quad - \varepsilon_n^{-\sigma} \{C_\delta \int_{\mathbb{R}^N} |w_{\varepsilon_n}|^q + \delta \varepsilon_n^{-\tau(q-2)} \int_{\mathbb{R}^N} |w_{\varepsilon_n}|^2\}. \end{aligned}$$

Sending $\varepsilon_n \rightarrow \infty$, it then follows from Lemma 4.3 that

$$\tilde{m}_\infty \geq \frac{1}{2} \int_{\mathbb{R}^N} |w_\infty|^2 - \frac{1}{2p} \int_{\mathbb{R}^N} (I_\alpha * |w_\infty|^{\frac{N+\alpha}{N}}) |w_\infty|^{\frac{N+\alpha}{N}} = J_\infty(w_\infty).$$

Therefore, note that $w_\infty \in \mathcal{M}_\infty$, we obtain $J_\infty(w_\infty) = \tilde{m}_\infty$. Thus, $w_\infty = U_\rho$ for some $\rho \in (0, +\infty)$.

Moreover, by (4.10), we obtain

$$\|\nabla w_\infty\|_2^2 \leq \lim_{\varepsilon_n \rightarrow \infty} \|\nabla w_{\varepsilon_n}\|_2^2 = \frac{NA(q-2)}{2q} \int_{\mathbb{R}^N} |w_\infty|^q,$$

from which it follows that

$$\rho \geq \left(\frac{2q \int_{\mathbb{R}^N} |\nabla U_1|^2}{NA(q-2) \int_{\mathbb{R}^N} |U_1|^q} \right)^{\frac{2}{4-N(q-2)}}.$$

If $\rho = \rho_0$, then (4.10) implies that $\lim_{n \rightarrow \infty} \|\nabla w_{\varepsilon_n}\|_2^2 = \|\nabla U_{\rho_0}\|_2^2$, and hence $w_{\varepsilon_n} \rightarrow U_{\rho_0}$ in $D^{1,2}(\mathbb{R}^N)$. \square

Proof of Theorem 2.1 in the case $q \in (2, 2 + \frac{4\alpha}{N(2+\alpha)})$. Let

$$M_\varepsilon = w_\varepsilon(0), \quad z_\varepsilon = M_\varepsilon [U_{\rho_0}(0)]^{-1},$$

where ρ_0 is given in Lemma 4.6. Then

$$z_\varepsilon = \frac{1}{U_{\rho_0}(0)} w_\varepsilon(0) = \frac{1}{U_{\rho_0}(0)} \varepsilon^{-\frac{2N}{\alpha[4-N(q-2)]}} u_\varepsilon(0).$$

Hence

$$\zeta_\varepsilon := z_\varepsilon^{-\frac{2}{N}} \varepsilon^{-\frac{N(q-2)}{\alpha[4-N(q-2)]}} = \left(\frac{U_{\rho_0}(0)}{u_\varepsilon(0)} \right)^{\frac{2}{N}} \varepsilon^{\frac{1}{\alpha}}, \quad z_\varepsilon^{-1} \varepsilon^{-\frac{2N}{\alpha[4-N(q-2)]}} = \varepsilon^{-\frac{N}{2\alpha}} \zeta_\varepsilon^{\frac{N}{2}}.$$

Arguing as in the proof of [27, Theorem 2.1], we can show that

$$\zeta_\varepsilon \sim \varepsilon^{-\frac{N(q-2)}{\alpha[4-N(q-2)]}}$$

and for large $\varepsilon > 0$, the rescaled family of ground states

$$\tilde{w}_\varepsilon(x) = z_\varepsilon^{-1} w_\varepsilon(z_\varepsilon^{-\frac{2}{N}} x) = z_\varepsilon^{-1} \varepsilon^{-\frac{2N}{\alpha[4-N(q-2)]}} u_\varepsilon(z_\varepsilon^{-\frac{2}{N}} \varepsilon^{-\frac{N(q-2)}{\alpha[4-N(q-2)]}} x) = \varepsilon^{-\frac{N}{2\alpha}} \zeta_\varepsilon^{\frac{N}{2}} u_\varepsilon(\zeta_\varepsilon x)$$

satisfies

$$\|\nabla \tilde{w}_\varepsilon\|_2^2 \sim \|\tilde{w}_\varepsilon\|_q^q \sim \int_{\mathbb{R}^N} (I_\alpha * |\tilde{w}_\varepsilon|^{\frac{N+\alpha}{N}}) |\tilde{w}_\varepsilon|^{\frac{N+\alpha}{N}} \sim \|\tilde{w}_\varepsilon\|_2^2 \sim 1,$$

and as $\varepsilon \rightarrow \infty$, \tilde{w}_ε converges in $L^2(\mathbb{R}^N)$ to the extremal function U_{ρ_0} . Then by Lemma 4.6, we also have $\tilde{w}_\varepsilon \rightarrow U_{\rho_0}$ in $D^{1,2}(\mathbb{R}^N)$. Thus we conclude that $\tilde{w}_\varepsilon \rightarrow U_{\rho_0}$ in $H^1(\mathbb{R}^N)$.

Since $w_\varepsilon \in \mathcal{M}_\varepsilon$ and w_ε is bounded in $H^1(\mathbb{R}^N)$, it follows that

$$\begin{aligned} \tilde{m}_\varepsilon &= \frac{2\alpha-1}{22\alpha} \varepsilon^{-\sigma} \int_{\mathbb{R}^N} |\nabla w_\varepsilon|^2 + \frac{2\alpha-1}{22\alpha} \int_{\mathbb{R}^N} |w_\varepsilon|^2 - \varepsilon^{-\sigma-\tau q} \int_{\mathbb{R}^N} [G(\varepsilon^\tau w_\varepsilon) - g(\varepsilon^\tau w_\varepsilon) \varepsilon^\tau w_\varepsilon] \\ &= \frac{\alpha}{2(N+\alpha)} \int_{\mathbb{R}^N} |w_\varepsilon|^2 + O(\varepsilon^{-\sigma}). \end{aligned}$$

Similarly, we also have

$$\tilde{m}_\infty = \frac{\alpha}{2(N+\alpha)} \int_{\mathbb{R}^N} |U_1|^2.$$

Then it follows from Lemma 4.3 that

$$\int_{\mathbb{R}^N} |U_1|^2 - \int_{\mathbb{R}^N} |w_\varepsilon|^2 = \frac{2(N+\alpha)}{\alpha} (\tilde{m}_\infty - \tilde{m}_\varepsilon) + O(\varepsilon^{-\sigma}) = O(\varepsilon^{-\sigma}).$$

Since $\|U_1\|_2^2 = \int_{\mathbb{R}^N} (I_\alpha * |U_1|^p) |U_1|^p = S_1^{\frac{N+\alpha}{\alpha}}$, we conclude that

$$\|\tilde{w}_\varepsilon\|_2^2 = \|w_\varepsilon\|_2^2 = S_1^{\frac{N+\alpha}{\alpha}} + O(\varepsilon^{-\sigma}).$$

Finally, by $w_\varepsilon \in \mathcal{M}_\varepsilon$ and the boundedness of $w_\varepsilon \in H^1(\mathbb{R}^N)$, we also have

$$\int_{\mathbb{R}^N} (I_\alpha * |\tilde{w}_\varepsilon|^{\frac{N+\alpha}{N}}) |\tilde{w}_\varepsilon|^{\frac{N+\alpha}{N}} = \int_{\mathbb{R}^N} (I_\alpha * |w_\varepsilon|^{\frac{N+\alpha}{N}}) |w_\varepsilon|^{\frac{N+\alpha}{N}} = \|w_\varepsilon\|_2^2 + O(\varepsilon^{-\sigma}) = S_1^{\frac{N+\alpha}{\alpha}} + O(\varepsilon^{-\sigma}).$$

The statements on u_ε follow from the corresponding results on v_ε and \tilde{w}_ε , and the proof is complete. \square

4.2. The Case $q \in (2 + \frac{4\alpha}{N(2+\alpha)}, 2 + \frac{4}{N})$. In this case, we set

$$v(x) = \varepsilon^{-\frac{1}{q-2}} u(\varepsilon^{-\frac{1}{2}} x). \quad (4.23)$$

Then (P_ε) is reduced to

$$-\Delta v + v = \varepsilon^{-\frac{N(2+\alpha)(q-2)-4\alpha}{2N(q-2)}} (I_\alpha * |v|^{2\alpha}) |v|^{2\alpha-2} v + \varepsilon^{\frac{1-q}{q-2}} g(\varepsilon^{\frac{1}{q-2}} v). \quad (4.24)$$

Let $u_\varepsilon \in H^1(\mathbb{R}^N)$ be the ground state for (P_ε) and

$$v_\varepsilon(x) = \varepsilon^{-\frac{1}{q-2}} u_\varepsilon(\varepsilon^{-\frac{1}{2}} x).$$

Then v_ε is a ground state of (4.24). Moreover, we have

$$\|u_\varepsilon\|_2^2 = \varepsilon^{\frac{4-N(q-2)}{2(q-2)}} \|v_\varepsilon\|_2^2, \quad \|\nabla u_\varepsilon\|_2^2 = \varepsilon^{\frac{2N-q(N-2)}{2(q-2)}} \|\nabla v_\varepsilon\|_2^2, \quad \|u_\varepsilon\|_q^q = \varepsilon^{\frac{2N-q(N-2)}{2(q-2)}} \|v_\varepsilon\|_q^q. \quad (4.25)$$

The limiting equation of (4.24) reads as follows

$$-\Delta v + v = A |v|^{q-2} v. \quad (4.26)$$

Associated to (4.24) and (4.26), the corresponding functionals are defined by

$$J_\varepsilon(v) := \frac{1}{2} \int_{\mathbb{R}^N} |\nabla v|^2 + |v|^2 - \frac{1}{22\alpha} \varepsilon^{-\frac{N(2+\alpha)(q-2)-4\alpha}{2N(q-2)}} \int_{\mathbb{R}^N} (I_\alpha * |v|^{2\alpha}) |v|^{2\alpha} \\ - \varepsilon^{-\frac{q}{q-2}} \int_{\mathbb{R}^N} G(\varepsilon^{\frac{1}{q-2}} v),$$

and

$$J_\infty(v) := \frac{1}{2} \int_{\mathbb{R}^N} |\nabla v|^2 + |v|^2 - \frac{A}{q} \int_{\mathbb{R}^N} |v|^q,$$

respectively. The associated Nehari manifolds are as follows

$$\mathcal{N}_\varepsilon := \left\{ v \in H^1(\mathbb{R}^N) \mid \begin{aligned} \int_{\mathbb{R}^N} |\nabla v_\varepsilon|^2 + |v_\varepsilon|^2 &= \varepsilon^{-\eta_q} \int_{\mathbb{R}^N} (I_\alpha * |v_\varepsilon|^{2\alpha}) |v_\varepsilon|^{2\alpha} \\ &+ \varepsilon^{\frac{1-q}{q-2}} \int_{\mathbb{R}^N} g(\varepsilon^{\frac{1}{q-2}} v_\varepsilon) v_\varepsilon \end{aligned} \right\},$$

$$\mathcal{N}_\infty := \left\{ v \in H^1(\mathbb{R}^N) \mid \int_{\mathbb{R}^N} |\nabla v|^2 + |v|^2 = A \int_{\mathbb{R}^N} |v|^q \right\},$$

respectively. Denote the least energies by \tilde{m}_ε and \tilde{m}_∞ , then

$$\tilde{m}_\varepsilon = \inf_{v \in \mathcal{N}_\varepsilon} J_\varepsilon(v), \quad \tilde{m}_\infty = \inf_{v \in \mathcal{N}_\infty} J_\infty(v)$$

are well defined and positive. Arguing as in Lemma 4.3, we show that $\tilde{m}_\varepsilon = \tilde{m}_\infty + o_\varepsilon(1)$.

Let $\eta_q := \frac{N(2+\alpha)(q-2)-4\alpha}{2N(q-2)}$, then

$$\tilde{m}_\varepsilon = \frac{1}{2} \int_{\mathbb{R}^N} |\nabla v_\varepsilon|^2 + |v_\varepsilon|^2 - \frac{1}{22_\alpha} \varepsilon^{-\eta_q} \int_{\mathbb{R}^N} (I_\alpha * |v_\varepsilon|^{2\alpha}) |v_\varepsilon|^{2\alpha} - \varepsilon^{-\frac{q}{q-2}} \int_{\mathbb{R}^N} G(\varepsilon^{\frac{1}{q-2}} v_\varepsilon), \quad (4.27)$$

$$\int_{\mathbb{R}^N} |\nabla v_\varepsilon|^2 + |v_\varepsilon|^2 = \varepsilon^{-\eta_q} \int_{\mathbb{R}^N} (I_\alpha * |v_\varepsilon|^{2\alpha}) |v_\varepsilon|^{2\alpha} + \varepsilon^{\frac{1-q}{q-2}} \int_{\mathbb{R}^N} g(\varepsilon^{\frac{1}{q-2}} v_\varepsilon) v_\varepsilon, \quad (4.28)$$

and

$$\frac{1}{2^*} \int_{\mathbb{R}^N} |\nabla v_\varepsilon|^2 + \frac{1}{2} \int_{\mathbb{R}^N} |v_\varepsilon|^2 = \frac{1}{2} \varepsilon^{-\eta_q} \int_{\mathbb{R}^N} (I_\alpha * |v_\varepsilon|^{2\alpha}) |v_\varepsilon|^{2\alpha} + \varepsilon^{-\frac{q}{q-2}} \int_{\mathbb{R}^N} G(\varepsilon^{\frac{1}{q-2}} v_\varepsilon). \quad (4.29)$$

By (4.27) and (4.29), we get

$$\tilde{m}_\infty + o_\varepsilon(1) \geq \tilde{m}_\varepsilon = \frac{1}{N} \int_{\mathbb{R}^N} |\nabla v_\varepsilon|^2 + \frac{2\alpha - 1}{22_\alpha} \varepsilon^{-\eta_q} \int_{\mathbb{R}^N} (I_\alpha * |v_\varepsilon|^{2\alpha}) |v_\varepsilon|^{2\alpha},$$

from which, we conclude that v_ε is bounded in $D^{1,2}(\mathbb{R}^N)$, $\varepsilon^{-\eta_q} \int_{\mathbb{R}^N} (I_\alpha * |v_\varepsilon|^{2\alpha}) |v_\varepsilon|^{2\alpha}$ is bounded. Then by (4.28), it is easy to show that v_ε is also bounded in $L^2(\mathbb{R}^N)$. Therefore, v_ε is bounded in $H^1(\mathbb{R}^N)$, and hence, by the Hardy-Littlewood-Sobolev indetity, $\varepsilon^{-\eta_q} \int_{\mathbb{R}^N} (I_\alpha * |v_\varepsilon|^{2\alpha}) |v_\varepsilon|^{2\alpha} = O(\varepsilon^{-\eta_q})$ and there holds

$$\begin{aligned} \tilde{m}_\varepsilon &= \frac{1}{2} \int_{\mathbb{R}^N} |\nabla v_\varepsilon|^2 + |v_\varepsilon|^2 - \frac{A}{q} \int_{\mathbb{R}^N} |v_\varepsilon|^q + o_\varepsilon(1), \\ \int_{\mathbb{R}^N} |\nabla v_\varepsilon|^2 + |v_\varepsilon|^2 &= A \int_{\mathbb{R}^N} |v_\varepsilon|^q + o_\varepsilon(1), \end{aligned}$$

and

$$\frac{1}{2^*} \int_{\mathbb{R}^N} |\nabla v_\varepsilon|^2 + \frac{1}{2} \int_{\mathbb{R}^N} |v_\varepsilon|^2 = \frac{A}{q} \int_{\mathbb{R}^N} |v_\varepsilon|^q + o_\varepsilon(1).$$

Therefore, we obtain

$$\int_{\mathbb{R}^N} |\nabla v_\varepsilon|^2 = \frac{NA(q-2)}{2q} \int_{\mathbb{R}^N} |v_\varepsilon|^q + o_\varepsilon(1).$$

Put

$$A_\varepsilon := \int_{\mathbb{R}^N} |\nabla v_\varepsilon|^2, \quad B_\varepsilon := \int_{\mathbb{R}^N} |v_\varepsilon|^2, \quad C_\varepsilon := \int_{\mathbb{R}^N} |v_\varepsilon|^q,$$

then

$$\begin{cases} \frac{1}{2} A_\varepsilon + \frac{1}{2} B_\varepsilon - \frac{A}{q} C_\varepsilon &= \tilde{m}_\varepsilon + o_\varepsilon(1), \\ A_\varepsilon + B_\varepsilon - AC_\varepsilon &= o_\varepsilon(1), \\ \frac{1}{2^*} A_\varepsilon + \frac{1}{2} B_\varepsilon - \frac{A}{q} C_\varepsilon &= o_\varepsilon(1). \end{cases}$$

Solving this system, we obtain

$$A_\varepsilon = N\tilde{m}_\varepsilon + o_\varepsilon(1), \quad B_\varepsilon = \frac{2N - q(N-2)}{q-2} \tilde{m}_\varepsilon + o_\varepsilon(1), \quad C_\varepsilon = \frac{2q}{A(q-2)} \tilde{m}_\varepsilon + o_\varepsilon(1).$$

It is easy to show that

$$\begin{aligned} \tilde{m}_\varepsilon &= \tilde{m}_\infty + o_\varepsilon(1) = \frac{q-2}{2q} A^{-\frac{2}{q-2}} S_q^{\frac{q}{q-2}} + o_\varepsilon(1), \\ m_\varepsilon &= \varepsilon^{\frac{2N-q(N-2)}{2(q-2)}} \tilde{m}_\varepsilon = \varepsilon^{\frac{2N-q(N-2)}{2(q-2)}} \left(\frac{q-2}{2q} A^{-\frac{2}{q-2}} S_q^{\frac{q}{q-2}} + o_\varepsilon(1) \right). \end{aligned}$$

Therefore, we conclude that

$$\begin{aligned} A_\varepsilon &= N\tilde{m}_\varepsilon + o_\varepsilon(1) = \frac{N(q-2)}{2q} A^{-\frac{2}{q-2}} S_q^{\frac{q}{q-2}} + o_\varepsilon(1), \\ B_\varepsilon &= \frac{2N-q(N-2)}{q-2} \tilde{m}_\varepsilon + o_\varepsilon(1) = \frac{2N-q(N-2)}{2q} A^{-\frac{2}{q-2}} S_q^{\frac{q}{q-2}} + o_\varepsilon(1), \\ C_\varepsilon &= \frac{2q}{A(q-2)} \tilde{m}_\varepsilon + o_\varepsilon(1) = A^{-\frac{q}{q-2}} S_q^{\frac{q}{q-2}} + o_\varepsilon(1). \end{aligned}$$

Form which, we conclude the proof of Theorem 2.1 in the case $q \in (2 + \frac{4\alpha}{N(2+\alpha)}, 2 + \frac{4}{N})$.

4.3. The Case $q = 2 + \frac{4\alpha}{N(2+\alpha)}$. In this case, we set

$$v(x) = \varepsilon^{-\frac{1}{q-2}} u(\varepsilon^{-\frac{1}{2}} x). \quad (4.30)$$

Then (P_ε) is reduced to

$$-\Delta v + v = (I_\alpha * |v|^{2\alpha}) |v|^{2\alpha-2} v + \varepsilon^{\frac{1-q}{q-2}} g(\varepsilon^{\frac{1}{q-2}} v). \quad (4.31)$$

Let $u_\varepsilon \in H^1(\mathbb{R}^N)$ be the ground state for (P_ε) and

$$v_\varepsilon(x) = \varepsilon^{-\frac{1}{q-2}} u_\varepsilon(\varepsilon^{-\frac{1}{2}} x).$$

Then v_ε is a ground state of (4.26). Moreover, we have

$$\|u_\varepsilon\|_2^2 = \varepsilon^{\frac{4-N(q-2)}{2(q-2)}} \|v_\varepsilon\|_2^2, \quad \|\nabla u_\varepsilon\|_2^2 = \varepsilon^{\frac{2N-q(N-2)}{2(q-2)}} \|\nabla v_\varepsilon\|_2^2, \quad \|u_\varepsilon\|_q^q = \varepsilon^{\frac{2N-q(N-2)}{2(q-2)}} \|v_\varepsilon\|_q^q. \quad (4.32)$$

The limiting equation of (4.31) reads as follows

$$-\Delta v + v = (I_\alpha * |v|^{2\alpha}) |v|^{2\alpha-2} v + A |v|^{q-2} v. \quad (4.33)$$

The rest of proof is similar to the case $q \in (2 + \frac{4\alpha}{N(2+\alpha)}, 2 + \frac{4}{N})$ and is omitted.

5. PROOF OF THEOREM 2.2

In this section, we always assume that $2_\alpha^* := \frac{N+\alpha}{N-2}$ and $q \in (2, 2^*)$ if $N \geq 4$, and $q \in (4, 6)$ if $N = 3$. For large $\varepsilon > 0$, the existence and symmetry of ground states in this case follows from the upper bound estimates of the least energy given by Lemma 5.3 and Lemma 5.4 below and the argument used in [26]. See also [30] and references therein.

5.1. The first scaling. It is easy to see that under the rescaling

$$w(x) = \varepsilon^{-\frac{1}{q-2}} u(\varepsilon^{-\frac{2}{(N-2)(q-2)}} x), \quad (5.1)$$

the equation (P_ε) is reduced to

$$-\Delta w + \varepsilon^{-\sigma} w = (I_\alpha * |w|^{2_\alpha^*}) |w|^{2_\alpha^*-2} w + \varepsilon^{-\sigma-\tau(q-1)} g(\varepsilon^\tau w), \quad (5.2)$$

where $\sigma := \frac{2^*-q}{q-2} = \frac{2N-q(N-2)}{(N-2)(q-2)} > 0$ and $\tau := \frac{1}{q-2}$.

The associated functional is defined by

$$J_\varepsilon(w) := \frac{1}{2} \int_{\mathbb{R}^N} |\nabla w|^2 + \varepsilon^{-\sigma} |w|^2 - \frac{1}{22_\alpha^*} \int_{\mathbb{R}^N} (I_\alpha * |w|^{2_\alpha^*}) |w|^{2_\alpha^*} - \varepsilon^{-\sigma-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau w).$$

Lemma 5.1. *Let $\varepsilon > 0$, $u \in H^1(\mathbb{R}^N)$ and w is the rescaling of u given in (5.1). Then*

$$(1) \quad \|\nabla w\|_2^2 = \|\nabla u\|_2^2, \quad \int_{\mathbb{R}^N} (I_\alpha * |w|^{2_\alpha^*}) |w|^{2_\alpha^*} = \int_{\mathbb{R}^N} (I_\alpha * |u|^{2_\alpha^*}) |u|^{2_\alpha^*},$$

- (2) $\varepsilon^{-\sigma} \|w\|_2^2 = \varepsilon \|u\|_2^2, \quad \varepsilon^{-\sigma} \|w\|_q^q = \|u\|_q^q,$
 (3) $I_\varepsilon(u) = J_\varepsilon(w).$

We define the Nehari manifolds as follows.

$$\mathcal{N}_\varepsilon = \left\{ w \in H^1(\mathbb{R}^N) \setminus \{0\} \mid \begin{array}{l} \int_{\mathbb{R}^N} |\nabla w|^2 + \varepsilon^{-\sigma} \int_{\mathbb{R}^N} |w|^2 \\ = \int_{\mathbb{R}^N} (I_\alpha * |w|^{2_\alpha^*}) |w|^{2_\alpha^*} + \varepsilon^{-\sigma-\tau(q-1)} \int_{\mathbb{R}^N} g(\varepsilon^\tau w) w \end{array} \right\}$$

and

$$\mathcal{N}_\infty = \left\{ w \in H^1(\mathbb{R}^N) \setminus \{0\} \mid \int_{\mathbb{R}^N} |\nabla w|^2 = \int_{\mathbb{R}^N} (I_\alpha * |w|^{2_\alpha^*}) |w|^{2_\alpha^*} \right\}.$$

Then

$$m_\varepsilon := \inf_{w \in \mathcal{N}_\varepsilon} J_\varepsilon(w) \quad \text{and} \quad m_\infty := \inf_{w \in \mathcal{N}_\infty} J_\infty(w)$$

are well-defined and positive. Moreover, J_∞ is attained on \mathcal{N}_∞ .

For $w \in H^1(\mathbb{R}^N) \setminus \{0\}$, we set

$$\tau_2(w) = \frac{\int_{\mathbb{R}^N} |\nabla w|^2}{\int_{\mathbb{R}^N} (I_\alpha * |w|^{2_\alpha^*}) |w|^{2_\alpha^*}}. \quad (5.3)$$

Then $(\tau_2(w))^{\frac{N-2}{2(2+\alpha)}} w \in \mathcal{N}_\infty$ for any $w \in H^1(\mathbb{R}^N) \setminus \{0\}$, and $w \in \mathcal{N}_\infty$ if and only if $\tau_2(w) = 1$.

Define the Pohožaev manifold as follows

$$\mathcal{P}_\varepsilon := \{w \in H^1(\mathbb{R}^N) \setminus \{0\} \mid P_\varepsilon(w) = 0\},$$

where

$$\begin{aligned} P_\varepsilon(w) : &= \frac{N-2}{2} \int_{\mathbb{R}^N} |\nabla w|^2 + \frac{\varepsilon^{-\sigma} N}{2} \int_{\mathbb{R}^N} |w|^2 \\ &\quad - \frac{N+\alpha}{22_\alpha^*} \int_{\mathbb{R}^N} (I_\alpha * |w|^{2_\alpha^*}) |w|^{2_\alpha^*} - N \varepsilon^{-\sigma-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau w). \end{aligned} \quad (5.4)$$

Then by Lemma 3.1, $w_\varepsilon \in \mathcal{P}_\varepsilon$. Moreover, we have a similar minimax characterizations for the least energy level m_ε as in Lemma 3.2. A similar result also holds for m_∞ and J_∞ .

Lemma 5.2. *The rescaled family of solutions $\{w_\varepsilon\}$ is bounded in $H^1(\mathbb{R}^N)$, and*

$$\|w_\varepsilon\|_2^2 = \frac{NA(2^* - q)}{2^* q} \|w_\varepsilon\|_q^q + o_\varepsilon(1), \quad \text{as } \varepsilon \rightarrow \infty. \quad (5.5)$$

Proof. To prove w_ε is bounded in $D^{1,2}(\mathbb{R}^N)$, we let

$$v(x) = \varepsilon^{-\frac{N-2}{4}} u(\varepsilon^{-\frac{1}{2}} x).$$

Then (P_ε) reduces to

$$-\Delta v + v = (I_\alpha * |v|^{2_\alpha^*}) |v|^{2_\alpha^*-2} v + \varepsilon^{-\frac{N+2}{4}} g(\varepsilon^{\frac{N-2}{4}} v),$$

and $v_\varepsilon(x) = \varepsilon^{-\frac{N-2}{4}} u_\varepsilon(\varepsilon^{-\frac{1}{2}} x)$ is a ground state solution, and hence

$$\frac{1}{2^*} \int_{\mathbb{R}^N} |\nabla v_\varepsilon|^2 + \frac{1}{2} \int_{\mathbb{R}^N} |v_\varepsilon|^2 = \frac{N+\alpha}{22_\alpha^* N} \int_{\mathbb{R}^N} (I_\alpha * |v_\varepsilon|^{2_\alpha^*}) |v_\varepsilon|^{2_\alpha^*} + \varepsilon^{-\frac{N}{2}} \int_{\mathbb{R}^N} G(\varepsilon^{\frac{N-2}{4}} v_\varepsilon),$$

and

$$\begin{aligned} m_\infty \geq m_\varepsilon &= \frac{1}{2} \int_{\mathbb{R}^N} |\nabla v_\varepsilon|^2 + |v_\varepsilon|^2 - \frac{1}{22_\alpha^*} \int_{\mathbb{R}^N} (I_\alpha * |v_\varepsilon|^{2_\alpha^*}) |v_\varepsilon|^{2_\alpha^*} - \varepsilon^{-\frac{N}{2}} \int_{\mathbb{R}^N} G(\varepsilon^{\frac{N-2}{4}} v_\varepsilon) \\ &= \frac{1}{N} \int_{\mathbb{R}^N} |\nabla v_\varepsilon|^2 + \frac{\alpha}{22_\alpha^* N} \int_{\mathbb{R}^N} (I_\alpha * |v_\varepsilon|^{2_\alpha^*}) |v_\varepsilon|^{2_\alpha^*}. \end{aligned}$$

Therefore, v_ε is bounded in $D^{1,2}(\mathbb{R}^N)$. Clearly, we have $\|\nabla v_\varepsilon\|_2 = \|\nabla u_\varepsilon\|_2 = \|\nabla w_\varepsilon\|_2$, it follows that $\{w_\varepsilon\}$ is bounded in $D^{1,2}(\mathbb{R}^N)$. Thus, it suffices to show that it is also bounded in $L^2(\mathbb{R}^N)$. By $w_\varepsilon \in \mathcal{N}_\varepsilon \cap \mathcal{P}_\varepsilon$, we obtain

$$\begin{aligned} \int_{\mathbb{R}^N} |\nabla w_\varepsilon|^2 + \varepsilon^{-\sigma} \int_{\mathbb{R}^N} |w_\varepsilon|^2 &= \int_{\mathbb{R}^N} (I_\alpha * |w_\varepsilon|^{2^*_\alpha}) |w_\varepsilon|^{2^*_\alpha} + \varepsilon^{-\sigma-\tau(q-1)} \int_{\mathbb{R}^N} g(\varepsilon^\tau w_\varepsilon) w_\varepsilon, \\ \frac{N-2}{2} \int_{\mathbb{R}^N} |\nabla w_\varepsilon|^2 + \frac{\varepsilon^{-\sigma} N}{2} \int_{\mathbb{R}^N} |w_\varepsilon|^2 &= \frac{N+\alpha}{22^*_\alpha} \int_{\mathbb{R}^N} (I_\alpha * |w_\varepsilon|^{2^*_\alpha}) |w_\varepsilon|^{2^*_\alpha} + N \varepsilon^{-\sigma-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau w_\varepsilon). \end{aligned}$$

Let $g(s) = As^{q-1} + \tilde{g}(s)$ for $s \geq 0$, then $G(s) = \frac{A}{q}s^q + \tilde{G}(s)$, then

$$\frac{1}{N} \int_{\mathbb{R}^N} |w_\varepsilon|^2 = \frac{A(2^* - q)}{2^* q} \int_{\mathbb{R}^N} |w_\varepsilon|^q + \varepsilon^{-\tau q} \int_{\mathbb{R}^N} [\tilde{G}(\varepsilon^\tau w_\varepsilon) - \frac{1}{2^*} \tilde{g}(\varepsilon^\tau w_\varepsilon) \varepsilon^\tau w_\varepsilon]. \quad (5.6)$$

By (H1) and (H2), for any $\delta > 0$ there exists a constant $C_\delta > 0$ such that

$$|\tilde{G}(s) - \frac{1}{2^*} \tilde{g}(s)s| \leq \delta s^q + C_\delta s^2, \quad \text{for all } s \geq 0.$$

Therefore, we get

$$\varepsilon^{-\tau q} \int_{\mathbb{R}^N} |\tilde{G}(\varepsilon^\tau w_\varepsilon) - \frac{1}{2^*} \tilde{g}(\varepsilon^\tau w_\varepsilon) \varepsilon^\tau w_\varepsilon| \leq \delta \int_{\mathbb{R}^N} |w_\varepsilon|^q + C_\delta \varepsilon^{-\tau(q-2)} \int_{\mathbb{R}^N} |w_\varepsilon|^2. \quad (5.7)$$

By (5.5), we have

$$[\frac{1}{N} - C_\delta \varepsilon^{-\tau(q-2)}] \int_{\mathbb{R}^N} |w_\varepsilon|^2 \leq [\frac{A(2^* - q)}{2^* q} + \delta] \int_{\mathbb{R}^N} |w_\varepsilon|^q. \quad (5.8)$$

By the Sobolev embedding theorem and the interpolation inequality, we obtain

$$\begin{aligned} \int_{\mathbb{R}^N} |w_\varepsilon|^q &\leq \left(\int_{\mathbb{R}^N} |w_\varepsilon|^2 \right)^{\frac{2^*-q}{2^*-2}} \left(\int_{\mathbb{R}^N} |w_\varepsilon|^{2^*} \right)^{\frac{q-2}{2^*-2}} \\ &\leq \left(\int_{\mathbb{R}^N} |w_\varepsilon|^2 \right)^{\frac{2^*-q}{2^*-2}} \left(\frac{1}{S} \int_{\mathbb{R}^N} |\nabla w_\varepsilon|^2 \right)^{\frac{2^*(q-2)}{2(2^*-2)}}, \end{aligned}$$

where S is the best Sobolev constant. It then follows from (5.7) and Lemma 5.1 that

$$\frac{1}{2N} \left(\int_{\mathbb{R}^N} |w_\varepsilon|^2 \right)^{\frac{q-2}{2^*-2}} \leq [\frac{A(2^* - q)}{2^* q} + \delta] \left(\frac{1}{S} \int_{\mathbb{R}^N} |\nabla w_\varepsilon|^2 \right)^{\frac{2^*(q-2)}{2(2^*-2)}},$$

which together with the boundedness of w_ε in $D^{1,2}(\mathbb{R}^N)$ implies that w_ε is bounded in $L^2(\mathbb{R}^N)$.

Finally, it follows from (5.7) that

$$\lim_{\varepsilon \rightarrow \infty} \varepsilon^{-\tau q} \int_{\mathbb{R}^N} [\tilde{G}(\varepsilon^\tau w_\varepsilon) - \frac{1}{2^*} \tilde{g}(\varepsilon^\tau w_\varepsilon) \varepsilon^\tau w_\varepsilon] = 0,$$

this together with (5.6) yields (5.5), and completes the proof. \square

Now, we give the following estimation on the least energy:

Lemma 5.3. *If $N \geq 5$ and $q \in (2, 2^*)$, then*

$$m_\infty - m_\varepsilon \sim \varepsilon^{-\sigma} \quad \text{as } \varepsilon \rightarrow \infty.$$

If $N = 4$ and $q \in (2, 4)$, or $N = 3$ and $q \in (4, 6)$, then

$$m_\infty - m_\varepsilon \lesssim \varepsilon^{-\sigma}, \quad \text{as } \varepsilon \rightarrow \infty.$$

Proof. First, we claim that

$$\tau_2(w_\varepsilon) = 1 + O(\varepsilon^{-\sigma}), \quad \text{as } \varepsilon \rightarrow \infty. \quad (5.9)$$

In fact, since $w_\varepsilon \in \mathcal{N}_\varepsilon$, we see that

$$\tau_2(w_\varepsilon) = \frac{\int_{\mathbb{R}^N} |\nabla w_\varepsilon|^2}{\int_{\mathbb{R}^N} (I_\alpha * |w_\varepsilon|^{2^*_\alpha}) |w_\varepsilon|^{2^*_\alpha}} = 1 + \varepsilon^{-\sigma} \frac{\varepsilon^{-\tau(q-1)} \int_{\mathbb{R}^N} g(\varepsilon^\tau w_\varepsilon) w_\varepsilon - \int_{\mathbb{R}^N} |w_\varepsilon|^2}{\int_{\mathbb{R}^N} (I_\alpha * |w_\varepsilon|^{2^*_\alpha}) |w_\varepsilon|^{2^*_\alpha}}.$$

Since

$$\int_{\mathbb{R}^N} |w_\varepsilon|^q \leq \left(\int_{\mathbb{R}^N} |w_\varepsilon|^2 \right)^{\frac{2^*-q}{2^*-2}} \left(\int_{\mathbb{R}^N} |w_\varepsilon|^{2^*} \right)^{\frac{q-2}{2^*-2}},$$

we see that

$$\begin{aligned} \frac{\varepsilon^{-\tau(q-1)} \int_{\mathbb{R}^N} g(\varepsilon^\tau w_\varepsilon) w_\varepsilon - \int_{\mathbb{R}^N} |w_\varepsilon|^2}{\int_{\mathbb{R}^N} |w_\varepsilon|^{2^*}} &\leq \frac{C_\delta \int_{\mathbb{R}^N} |w_\varepsilon|^q - (1 - \delta \varepsilon^{-\tau(q-2)}) \int_{\mathbb{R}^N} |w_\varepsilon|^2}{\int_{\mathbb{R}^N} |w_\varepsilon|^{2^*}} \\ &\leq \zeta_\varepsilon^{\theta_q} (C_\delta - (1 - \delta \varepsilon^{-\tau(q-2)}) \zeta_\varepsilon^{1-\theta_q}). \end{aligned}$$

where

$$\theta_q = \frac{2^* - q}{2^* - 2}, \quad \zeta_\varepsilon = \frac{\int_{\mathbb{R}^N} |w_\varepsilon|^2}{\int_{\mathbb{R}^N} |w_\varepsilon|^{2^*}}.$$

Let $g(x) := x^{\theta_q} (C_\delta - (1 - \delta \varepsilon^{-\tau(q-2)}) x^{1-\theta_q})$. Then

$$\sup_{x \geq 0} g(x) = K_q := C_\delta \theta_q (1 - \theta_q) \left(\frac{C_\delta \theta_q}{1 - \delta \varepsilon^{-\tau(q-2)}} \right)^{\frac{\theta_q}{1-\theta_q}}.$$

Therefore, by the boundedness of w_ε in $D^{1,2}(\mathbb{R}^N)$, we get

$$\begin{aligned} \tau_2(w_\varepsilon) &\leq 1 + \varepsilon^{-\sigma} K_q \frac{\int_{\mathbb{R}^N} |w_\varepsilon|^{2^*}}{\int_{\mathbb{R}^N} (I_\alpha * |w_\varepsilon|^{2^*_\alpha}) |w_\varepsilon|^{2^*_\alpha}} \\ &\leq 1 + \varepsilon^{-\sigma} K_q S^{-\frac{N}{N-2}} \frac{(\int_{\mathbb{R}^N} |\nabla w_\varepsilon|^2)^{\frac{N}{N-2}}}{\int_{\mathbb{R}^N} (I_\alpha * |w_\varepsilon|^{2^*_\alpha}) |w_\varepsilon|^{2^*_\alpha}} \\ &= 1 + \varepsilon^{-\sigma} K_q S^{-\frac{N}{N-2}} \tau_2(w_\varepsilon) (\int_{\mathbb{R}^N} |\nabla w_\varepsilon|^2)^{\frac{2}{N-2}} \\ &\leq 1 + \varepsilon^{-\sigma} C \tau_2(w_\varepsilon), \end{aligned}$$

and hence for large $\varepsilon > 0$, there holds

$$\tau_2(w_\varepsilon) \leq \frac{1}{1 - \varepsilon^{-\sigma} C} = 1 + \varepsilon^{-\sigma} \frac{C}{1 - \varepsilon^{-\sigma} C} \leq 1 + \frac{1}{2} C \varepsilon^{-\sigma}.$$

On the other hand, by (5.3), the Hardy-Littlewood-Sobolev and the Sobolev inequalities, we obtain that

$$\int_{\mathbb{R}^N} |\nabla w_\varepsilon|^2 = \tau_2(w_\varepsilon) \int_{\mathbb{R}^N} (I_\alpha * |w_\varepsilon|^{2^*_\alpha}) |w_\varepsilon|^{2^*_\alpha} \leq C \tau_2(w_\varepsilon) \left(\frac{1}{S} \int_{\mathbb{R}^N} |\nabla w_\varepsilon|^2 \right)^{2^*_\alpha}.$$

Therefore, we get

$$\int_{\mathbb{R}^N} |\nabla w_\varepsilon|^2 \geq \left(\frac{C \tau_2(w_\varepsilon)}{S^{2^*_\alpha}} \right)^{-\frac{1}{2^*_\alpha-1}} \geq C > 0.$$

By (5.3), $w_\varepsilon \in \mathcal{N}_\varepsilon$ and the boundedness of w_ε in $H^1(\mathbb{R}^N)$, we then obtain

$$\begin{aligned}\tau_2(w_\varepsilon) &= \frac{\int_{\mathbb{R}^N} |\nabla w_\varepsilon|^2}{\int_{\mathbb{R}^N} (I_\alpha * |w_\varepsilon|^{2^*}) |w_\varepsilon|^{2^*}} \\ &= \frac{\int_{\mathbb{R}^N} |\nabla w_\varepsilon|^2}{\int_{\mathbb{R}^N} |\nabla w_\varepsilon|^2 + \varepsilon^{-\sigma} \int_{\mathbb{R}^N} |w_\varepsilon|^{2-\varepsilon^{-\sigma-\tau(q-1)}} \int_{\mathbb{R}^N} g(\varepsilon^\tau w_\varepsilon) w_\varepsilon} \\ &\geq \frac{\int_{\mathbb{R}^N} |\nabla w_\varepsilon|^2}{\int_{\mathbb{R}^N} |\nabla w_\varepsilon|^2 + \varepsilon^{-\sigma} \int_{\mathbb{R}^N} |w_\varepsilon|^2} \\ &\geq 1 - \frac{1}{2} \varepsilon^{-\sigma} \frac{\int_{\mathbb{R}^N} |w_\varepsilon|^2}{\int_{\mathbb{R}^N} |\nabla w_\varepsilon|^2} \\ &\geq 1 - C \varepsilon^{-\sigma}.\end{aligned}$$

Thus we get $\tau_2(w_\varepsilon) = 1 + O(\varepsilon^{-\sigma})$ as $\varepsilon \rightarrow \infty$. This proved the claim.

If $N \geq 3$, by Lemma 3.2, (H2) and the boundedness of $\{w_\varepsilon\}$, we find

$$\begin{aligned}m_\infty &\leq \sup_{t \geq 0} J_\varepsilon((w_\varepsilon)_t) + \varepsilon^{-\sigma} t_\varepsilon^N (\varepsilon^{-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau w_\varepsilon) - \frac{1}{2} \int_{\mathbb{R}^N} |w_\varepsilon|^2) \\ &\leq m_\varepsilon + C \varepsilon^{-\sigma},\end{aligned}\tag{5.10}$$

where

$$t_\varepsilon = \left(\frac{\int_{\mathbb{R}^N} |\nabla w_\varepsilon|^2}{\int_{\mathbb{R}^N} (I_\alpha * |w_\varepsilon|^{2^*}) |w_\varepsilon|^{2^*}} \right)^{\frac{1}{2+\alpha}} = (\tau_2(w_\varepsilon))^{\frac{1}{2+\alpha}}.$$

Next, we assume $N \geq 5$. For each $\rho > 0$, the family $V_\rho(x) := \rho^{-\frac{N-2}{2}} V_1(x/\rho)$ are radial ground states of $-\Delta w = (I_\alpha * |w|^{2^*}) w^{2^*-1}$, and verify that

$$\|V_\rho\|_2^2 = \rho^2 \|V_1\|_2^2, \quad \int_{\mathbb{R}^N} |V_\rho|^q = \rho^{N-\frac{N-2}{2}q} \int_{\mathbb{R}^N} |V_1|^q.\tag{5.11}$$

Let $g_0(\rho) = \frac{A}{q} \int_{\mathbb{R}^N} |V_\rho|^q - \frac{1}{2} \int_{\mathbb{R}^N} |V_\rho|^2$. Then there exists $\rho_0 = \rho(q) \in (0, +\infty)$ with

$$\rho_0 = \left(\frac{A[2N - q(N-2)] \int_{\mathbb{R}^N} |V_1|^q}{2q \int_{\mathbb{R}^N} |V_1|^2} \right)^{\frac{2}{(N-2)(q-2)}}$$

such that

$$g_0(\rho_0) = \sup_{\rho > 0} g_0(\rho) = \frac{A(N-2)(q-2)}{4q} \left(\frac{A[2N - q(N-2)] \int_{\mathbb{R}^N} |V_1|^q}{2q \int_{\mathbb{R}^N} |V_1|^2} \right)^{\frac{2N-q(N-2)}{(N-2)(q-2)}} \int_{\mathbb{R}^N} |V_1|^q.$$

Let $V_0 = V_{\rho_0}$, then there exists $t_\varepsilon \in (0, +\infty)$ such that

$$\begin{aligned}m_\varepsilon &\leq \sup_{t \geq 0} J_\varepsilon((V_0)_t) = J_\varepsilon((V_0)_{t_\varepsilon}) \\ &= \frac{t_\varepsilon^{N-2}}{2} \int_{\mathbb{R}^N} |\nabla V_0|^2 - \frac{t_\varepsilon^{N+\alpha}}{22^*} (I_\alpha * |V_0|^{2^*}) |V_0|^{2^*} \\ &\quad - \varepsilon^{-\sigma} t_\varepsilon^N \left\{ \varepsilon^{-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau V_0) - \frac{1}{2} \int_{\mathbb{R}^N} |V_0|^2 \right\} \\ &\leq \sup_{t \geq 0} \left(\frac{t^{N-2}}{2} - \frac{t^{N+\alpha}}{2p} \right) \int_{\mathbb{R}^N} |\nabla V_0|^2 \\ &\quad - \varepsilon^{-\sigma} t_\varepsilon^N \left\{ \varepsilon^{-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau V_0) - \frac{1}{2} \int_{\mathbb{R}^N} |V_0|^2 \right\} \\ &= m_\infty - \varepsilon^{-\sigma} t_\varepsilon^N \left\{ \varepsilon^{-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau V_0) - \frac{1}{2} \int_{\mathbb{R}^N} |V_0|^2 \right\}.\end{aligned}\tag{5.12}$$

Moreover, t_ε satisfies

$$\frac{1}{2^* t^2} \int_{\mathbb{R}^N} |\nabla V_0|^2 = \frac{t^\alpha}{2^*} \int_{\mathbb{R}^N} (I_\alpha * |V_0|^{2^*}) |V_0|^{2^*} + \varepsilon^{-\sigma} \left\{ \varepsilon^{-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau V_0) - \frac{1}{2} \int_{\mathbb{R}^N} |V_0|^2 \right\}.$$

If $t_\varepsilon \geq 1$, then

$$\int_{\mathbb{R}^N} |\nabla V_0|^2 \geq t_\varepsilon^2 \left\{ \int_{\mathbb{R}^N} (I_\alpha * |V_0|^{2^*}) |V_0|^{2^*} + \varepsilon^{-\sigma-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau V_0) - \frac{1}{2} \varepsilon^{-\sigma} \int_{\mathbb{R}^N} |V_0|^2 \right\}.$$

Hence

$$t_\varepsilon^2 \leq \frac{\int_{\mathbb{R}^N} |\nabla V_0|^2}{\int_{\mathbb{R}^N} (I_\alpha * |V_0|^{2_\alpha^*}) |V_0|^{2_\alpha^*} + 2^* \varepsilon^{-\sigma} \{ \varepsilon^{-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau V_0) - \frac{1}{2} \int_{\mathbb{R}^N} |V_0|^2 \}}.$$

If $t_\varepsilon \leq 1$, then

$$\int_{\mathbb{R}^N} |\nabla V_0|^2 \leq t_\varepsilon^2 \left\{ \int_{\mathbb{R}^N} (I_\alpha * |V_0|^{2_\alpha^*}) |V_0|^{2_\alpha^*} + \varepsilon^{-\sigma-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau V_0) - \frac{1}{2} \varepsilon^{-\sigma} \int_{\mathbb{R}^N} |V_0|^2 \right\}.$$

Hence

$$t_\varepsilon^2 \geq \frac{\int_{\mathbb{R}^N} |\nabla V_0|^2}{\int_{\mathbb{R}^N} (I_\alpha * |V_0|^{2_\alpha^*}) |V_0|^{2_\alpha^*} + 2^* \varepsilon^{-\sigma} \{ \varepsilon^{-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau V_0) - \frac{1}{2} \int_{\mathbb{R}^N} |V_0|^2 \}}.$$

Since

$$\int_{\mathbb{R}^N} (I_\alpha * |V_0|^{2_\alpha^*}) |V_0|^{2_\alpha^*} = \int_{\mathbb{R}^N} |\nabla V_0|^2,$$

we conclude that $\lim_{\varepsilon \rightarrow \infty} t_\varepsilon = 1$. By (H2), it follows that

$$\lim_{\varepsilon \rightarrow \infty} \varepsilon^{-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau V_0) - \frac{1}{2} \int_{\mathbb{R}^N} |V_0|^2 = \frac{A}{q} \int_{\mathbb{R}^N} |V_0|^q - \frac{1}{2} \int_{\mathbb{R}^N} |V_0|^2 = g_0(\rho_0),$$

Thus, by (5.12), there exists a constant $C > 0$ such that

$$m_\varepsilon \leq m_\infty - C \varepsilon^{-\sigma},$$

for large $\varepsilon > 0$. The proof is complete. \square

Lemma 5.4. *There exists a constant $\varpi = \varpi(q) > 0$ such that for $\varepsilon > 0$ large,*

$$m_\varepsilon \leq \begin{cases} m_\infty - (\varepsilon \ln \varepsilon)^{-\frac{4-q}{q-2}} \varpi, & \text{if } N = 4, \\ m_\infty - \varepsilon^{-\frac{6-q}{2(q-4)}} \varpi, & \text{if } N = 3 \text{ and } q \in (4, 6). \end{cases}$$

Proof. Let $\rho > 0$ and R be a large parameter, and $\eta_R \in C_0^\infty(\mathbb{R})$ is a cut-off function such that $\eta_R(r) = 1$ for $|r| < R$, $0 < \eta_R(r) < 1$ for $R < |r| < 2R$, $\eta_R(r) = 0$ for $|r| > 2R$ and $|\eta_R'(r)| \leq 2/R$.

For $\ell \gg 1$, a straightforward computation shows that

$$\int_{\mathbb{R}^N} |\nabla(\eta_\ell V_1)|^2 = \begin{cases} \frac{2(N+\alpha)}{2+\alpha} m_\infty + O(\ell^{-2}), & \text{if } N = 4, \\ \frac{2(N+\alpha)}{2+\alpha} m_\infty + O(\ell^{-1}), & \text{if } N = 3. \end{cases}$$

$$\int_{\mathbb{R}^N} (I_\alpha * |\eta_\ell V_1|^{2_\alpha^*}) |\eta_\ell V_1|^{2_\alpha^*} = \frac{2(N+\alpha)}{2+\alpha} m_\infty + O(\ell^{-N}),$$

$$\int_{\mathbb{R}^N} |\eta_\ell V_1|^2 = \begin{cases} \ln \ell (1 + o(1)), & \text{if } N = 4, \\ \ell (1 + o(1)), & \text{if } N = 3. \end{cases}$$

By Lemma 3.2, we find

$$\begin{aligned} m_\varepsilon &\leq \sup_{t \geq 0} J_\varepsilon((\eta_R V_\rho)_t) = J_\varepsilon((\eta_R V_\rho)_{t_\varepsilon}) \\ &\leq \sup_{t \geq 0} \left(\frac{t^{N-2}}{2} \int_{\mathbb{R}^N} |\nabla(\eta_R V_\rho)|^2 - \frac{t^{N+\alpha}}{2p} \int_{\mathbb{R}^N} (I_\alpha * |\eta_R V_\rho|^{2_\alpha^*}) |\eta_R V_\rho|^{2_\alpha^*} \right. \\ &\quad \left. - \varepsilon^{-\sigma} t_\varepsilon^N \left[\varepsilon^{-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau \eta_R V_\rho) - \frac{1}{2} \int_{\mathbb{R}^N} |\eta_R V_\rho|^2 \right] \right) \\ &= (I) - \varepsilon^{-\sigma} (II), \end{aligned} \tag{5.13}$$

where $t_\varepsilon \in (0, +\infty)$ is the unique critical point of the function $g(t)$ defined by

$$g(t) = \frac{t^{N-2}}{2} \int_{\mathbb{R}^N} |\nabla(\eta_R V_\rho)|^2 + \frac{t^N}{2} \varepsilon^{-\sigma} \int_{\mathbb{R}^N} |\eta_R V_\rho|^2 - \frac{t^{N+\alpha}}{22_\alpha^*} \int_{\mathbb{R}^N} (I_\alpha * |\eta_R V_\rho|^{2_\alpha^*}) |\eta_R V_\rho|^{2_\alpha^*} - t^N \varepsilon^{-\sigma-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau \eta_R V_\rho).$$

That is, $t = t_\varepsilon$ solves the equation $\ell_1(t) = \ell_2(t)$, where

$$\ell_1(t) := \frac{1}{2^* t^2} \int_{\mathbb{R}^N} |\nabla(\eta_R V_\rho)|^2$$

and

$$\ell_2(t) := \frac{1}{2^*} t^\alpha \int_{\mathbb{R}^N} (I_\alpha * |\eta_R V_\rho|^{2_\alpha^*}) |\eta_R V_\rho|^{2_\alpha^*} + \varepsilon^{-\sigma-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau \eta_R V_\rho) - \frac{1}{2} \varepsilon^{-\sigma} \int_{\mathbb{R}^N} |\eta_R V_\rho|^2.$$

If $t_\varepsilon \geq 1$, then

$$\begin{aligned} \frac{1}{2^* t_\varepsilon^2} \int_{\mathbb{R}^N} |\nabla(\eta_R V_\rho)|^2 &\geq \frac{1}{2^*} \int_{\mathbb{R}^N} (I_\alpha * |\eta_R V_\rho|^{2_\alpha^*}) |\eta_R V_\rho|^{2_\alpha^*} \\ &\quad + \varepsilon^{-\sigma-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau \eta_R V_\rho) - \frac{1}{2} \varepsilon^{-\sigma} \int_{\mathbb{R}^N} |\eta_R V_\rho|^2, \end{aligned}$$

and hence

$$t_\varepsilon^2 \leq \frac{\int_{\mathbb{R}^N} |\nabla(\eta_R V_\rho)|^2}{\int_{\mathbb{R}^N} (I_\alpha * |\eta_R V_\rho|^{2_\alpha^*}) |\eta_R V_\rho|^{2_\alpha^*} + 2^* \varepsilon^{-\sigma-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau \eta_R V_\rho) - \frac{1}{2} \varepsilon^{-\sigma} \int_{\mathbb{R}^N} |\eta_R V_\rho|^2}. \quad (5.14)$$

If $t_\varepsilon \leq 1$, then

$$\begin{aligned} \frac{1}{2^* t_\varepsilon^2} \int_{\mathbb{R}^N} |\nabla(\eta_R V_\rho)|^2 &\leq \frac{1}{2^*} \int_{\mathbb{R}^N} (I_\alpha * |\eta_R V_\rho|^{2_\alpha^*}) |\eta_R V_\rho|^{2_\alpha^*} \\ &\quad + \varepsilon^{-\sigma-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau \eta_R V_\rho) - \frac{N}{2} \varepsilon^{-\sigma} \int_{\mathbb{R}^N} |\eta_R V_\rho|^2, \end{aligned}$$

and hence

$$t_\varepsilon^2 \geq \frac{\int_{\mathbb{R}^N} |\nabla(\eta_R V_\rho)|^2}{\int_{\mathbb{R}^N} (I_\alpha * |\eta_R V_\rho|^{2_\alpha^*}) |\eta_R V_\rho|^{2_\alpha^*} + 2^* \varepsilon^{-\sigma-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau \eta_R V_\rho) - \frac{1}{2} \varepsilon^{-\sigma} \int_{\mathbb{R}^N} |\eta_R V_\rho|^2}. \quad (5.15)$$

Therefore, set $\ell = R/\rho$, then we obtain

$$\limsup_{\varepsilon \rightarrow \infty} |t_\varepsilon^2 - 1| \leq \left| \frac{\int_{\mathbb{R}^N} |\nabla(\eta_R V_\rho)|^2}{\int_{\mathbb{R}^N} (I_\alpha * |\eta_R V_\rho|^{2_\alpha^*}) |\eta_R V_\rho|^{2_\alpha^*}} - 1 \right| = \left| \frac{\int_{\mathbb{R}^N} |\nabla(\eta_\ell V_1)|^2}{\int_{\mathbb{R}^N} (I_\alpha * |\eta_\ell V_1|^{2_\alpha^*}) |\eta_\ell V_1|^{2_\alpha^*}} - 1 \right| \rightarrow 0,$$

as $\ell \rightarrow \infty$. Therefore, we get

$$\begin{aligned} (I) &= \frac{2+\alpha}{2(N+\alpha)} \frac{(\int_{\mathbb{R}^N} |\nabla(\eta_\ell V_1)|^2)^{\frac{N+\alpha}{2+\alpha}}}{(\int_{\mathbb{R}^N} (I_\alpha * |\eta_\ell V_1|^{2_\alpha^*}) |\eta_\ell V_1|^{2_\alpha^*})^{\frac{N-2}{2+\alpha}}} \\ &= \begin{cases} \frac{2+\alpha}{2(N+\alpha)} \frac{(\frac{2(N+\alpha)}{2+\alpha} m_\infty + O(\ell^{-2}))^{\frac{N+\alpha}{2+\alpha}}}{(\frac{2(N+\alpha)}{2+\alpha} m_\infty + O(\ell^{-4}))^{\frac{N-2}{2+\alpha}}}, & \text{if } N = 4, \\ \frac{2+\alpha}{2(N+\alpha)} \frac{(\frac{2(N+\alpha)}{2+\alpha} m_\infty + O(\ell^{-1}))^{\frac{N+\alpha}{2+\alpha}}}{(\frac{2(N+\alpha)}{2+\alpha} m_\infty + O(\ell^{-3}))^{\frac{N-2}{2+\alpha}}}, & \text{if } N = 3, \end{cases} \\ &= \begin{cases} m_\infty + O(\ell^{-2}), & \text{if } N = 4, \\ m_\infty + O(\ell^{-1}), & \text{if } N = 3. \end{cases} \end{aligned} \quad (5.16)$$

Since

$$\begin{aligned} \phi(\rho) &= \varepsilon^{-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau \eta_R V_\rho) - \frac{1}{2} \int_{\mathbb{R}^N} |\eta_R V_\rho|^2 \\ &= \frac{A}{q} \int_{\mathbb{R}^N} |\eta_R V_\rho|^q - \frac{1}{2} \int_{\mathbb{R}^N} |\eta_R V_\rho|^2 \\ &= \frac{A}{q} \rho^{N-\frac{N-2}{2}q} \int_{\mathbb{R}^N} |\eta_\ell V_1|^q - \frac{1}{2} \rho^2 \int_{\mathbb{R}^N} |\eta_\ell V_1|^2 \end{aligned}$$

take its maximum value $\varphi(\rho_\ell)$ at the unique point

$$\begin{aligned} \rho_\ell : &= \left(\frac{A[2N-q(N-2)] \int_{\mathbb{R}^N} |\eta_\ell V_1|^q}{2q \int_{\mathbb{R}^N} |\eta_\ell V_1|^2} \right)^{\frac{2}{(N-2)(q-2)}} \\ &\sim \begin{cases} (\ln \ell)^{-\frac{2}{(N-2)(q-2)}} & \text{if } N = 4, \\ \ell^{-\frac{2}{(N-2)(q-2)}} & \text{if } N = 3, \end{cases} \end{aligned}$$

we obtain

$$\begin{aligned} \phi(\rho_\ell) &= \sup_{\rho \geq 0} \phi(\rho) \\ &= \frac{4+q(N-2)-2N}{4q} \rho_\ell^{N-\frac{N-2}{2}q} \int_{\mathbb{R}^N} |\eta_\ell V_1|^q \\ &= \frac{4+q(N-2)-2N}{4q} \left(\frac{2N-q(N-2)}{2q} \right)^{\frac{2N-q(N-2)}{(N-2)(q-2)}} \frac{(\int_{\mathbb{R}^N} |\eta_\ell V_1|^q)^{\frac{4}{(N-2)(q-2)}}}{(\int_{\mathbb{R}^N} |\eta_\ell V_1|^2)^{\frac{2N-q(N-2)}{(N-2)(q-2)}}} \\ &\leq \frac{4+q(N-2)-2N}{4q} \left(\frac{2N-q(N-2)}{2q} \right)^{\frac{2N-q(N-2)}{(N-2)(q-2)}} \int_{\mathbb{R}^N} |\eta_\ell V_1|^{2^*} \\ &\rightarrow \frac{4+q(N-2)-2N}{4q} \left(\frac{2N-q(N-2)}{2q} \right)^{\frac{2N-q(N-2)}{(N-2)(q-2)}} \int_{\mathbb{R}^N} |V_1|^{2^*}, \end{aligned}$$

as $\ell \rightarrow +\infty$, where we have used the interpolation inequality

$$\int_{\mathbb{R}^N} |\eta_\ell V_1|^q \leq \left(\int_{\mathbb{R}^N} |\eta_\ell V_1|^2 \right)^{\frac{2^*-q}{2^*-2}} \left(\int_{\mathbb{R}^N} |\eta_\ell V_1|^{2^*} \right)^{\frac{q-2}{2^*-2}}.$$

Since $\int_{\mathbb{R}^N} |\eta_\ell V_1|^q \rightarrow \int_{\mathbb{R}^N} |V_1|^q$, as $\ell \rightarrow +\infty$, it follows that

$$\begin{aligned} \phi(\rho_\ell) &= \frac{(N-2)(q-2)}{4q} A^{\frac{4}{(N-2)(q-2)}} \left(\frac{[2N-q(N-2)] \int_{\mathbb{R}^N} |\eta_\ell V_1|^q}{2q \int_{\mathbb{R}^N} |\eta_\ell V_1|^2} \right)^{\frac{2N-q(N-2)}{(N-2)(q-2)}} \int_{\mathbb{R}^N} |\eta_\ell V_1|^q \\ &= \begin{cases} C(\ln \ell(1+o(1)))^{-\frac{2N-q(N-2)}{(N-2)(q-2)}} & \text{if } N = 4, \\ C(\ell(1+o(1)))^{-\frac{2N-q(N-2)}{(N-2)(q-2)}} & \text{if } N = 3. \end{cases} \end{aligned}$$

Thus, for large $\varepsilon > 0$, we have

$$\begin{aligned} (II) &= \phi(\rho_\ell) + (t_\varepsilon^N - 1)\phi(\rho_\ell) \\ &\sim \begin{cases} (\ln \ell)^{-\frac{2N-q(N-2)}{(N-2)(q-2)}}, & \text{if } N = 4, \\ \ell^{-\frac{2N-q(N-2)}{(N-2)(q-2)}}, & \text{if } N = 3. \end{cases} \end{aligned}$$

It follows that if $N = 4$, then

$$\begin{aligned} m_\varepsilon &\leq (I) - \varepsilon^{-\sigma}(II) \\ &\leq m_\infty + O(\ell^{-2}) - C\varepsilon^{-\sigma}(\ln \ell)^{-\frac{2N-q(N-2)}{(N-2)(q-2)}}. \end{aligned} \tag{5.17}$$

Take $\ell = \varepsilon^M$. Then

$$m_\varepsilon \leq m_\infty + O(\varepsilon^{-2M}) - C\varepsilon^{-\sigma} M^{-\frac{2N-q(N-2)}{(N-2)(q-2)}} (\ln \varepsilon)^{-\frac{2N-q(N-2)}{(N-2)(q-2)}}.$$

If $M > \frac{4-q}{2(q-2)}$, then $2M > \sigma$, and hence

$$m_\varepsilon \leq m_\infty - \varepsilon^{-\sigma} (\ln \varepsilon)^{-\frac{2N-q(N-2)}{(N-2)(q-2)}} \varpi = m_\infty - (\varepsilon \ln \varepsilon)^{-\frac{4-q}{q-2}} \varpi, \tag{5.18}$$

for large $\varepsilon > 0$, where

$$\varpi = \frac{1}{2} C M^{-\frac{2N-q(N-2)}{(N-2)(q-2)}}.$$

If $N = 3$, then

$$\begin{aligned} m_\lambda &\leq (I) - \lambda^\sigma (II) \\ &\leq m_\infty + O(\ell^{-1}) - C\varepsilon^{-\sigma} \ell^{-\frac{2N-q(N-2)}{(N-2)(q-2)}}. \end{aligned} \quad (5.19)$$

Take $\ell = \delta^{-1}\varepsilon^\tau$. Then

$$m_\varepsilon \leq m_\infty + \delta O(\varepsilon^{-\tau}) - C\varepsilon^{-\sigma} \delta^{\frac{2N-q(N-2)}{(N-2)(q-2)}} \varepsilon^{-\tau \frac{2N-q(N-2)}{(N-2)(q-2)}}$$

If $q \in (4, 6)$ and $\tau = \frac{6-q}{2(q-4)}$, then

$$\tau = \frac{2(N-2)}{2+q(N-2)-2N} = \frac{2}{q-4},$$

then

$$m_\varepsilon \leq m_\infty + (\delta O(1) - C\delta^{\frac{2N-q(N-2)}{(N-2)(q-2)}})\varepsilon^{-\frac{6-q}{2(q-4)}}.$$

Since

$$1 > \frac{2N-q(N-2)}{(N-2)(q-2)},$$

it follows that for some small $\delta > 0$, there exists $\varpi > 0$ such that

$$m_\varepsilon \leq m_\infty - \varepsilon^{-\frac{6-q}{2(q-4)}} \varpi.$$

This completes the proof. \square

Combining Lemma 5.3 and Lemma 5.4, we get the following

Lemma 5.5. *Let $\delta_\varepsilon := m_\infty - m_\varepsilon$, then*

$$\varepsilon^{-\sigma} \gtrsim \delta_\varepsilon \gtrsim \begin{cases} \varepsilon^{-\sigma}, & \text{if } N \geq 5, \\ (\varepsilon \ln \varepsilon)^{-\frac{4-q}{q-2}}, & \text{if } N = 4, \\ \varepsilon^{-\frac{6-q}{2(q-4)}}, & \text{if } N = 3 \text{ and } q \in (4, 6). \end{cases}$$

Lemma 5.6. *Assume $N \geq 5$. Then $\|w_\varepsilon\|_q \sim 1$ as $\varepsilon \rightarrow \infty$.*

Proof. By (5.7), we have

$$\begin{aligned} m_\infty &\leq m_\varepsilon + \varepsilon^{-\sigma} (\tau_2(w_\varepsilon))^{\frac{N}{2+\alpha}} \varepsilon^{-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau w_\varepsilon) \\ &\leq m_\varepsilon + \frac{A}{q} \varepsilon^{-\sigma} (\tau_2(w_\varepsilon))^{\frac{N}{2+\alpha}} (\int_{\mathbb{R}^N} |w_\varepsilon|^q + o_\varepsilon(1)). \end{aligned}$$

Therefore, it follows from (5.6) and Corollary 5.5 that

$$\|w_\varepsilon\|_q^q \geq \frac{m_\infty - m_\varepsilon}{(\tau_2(w_\varepsilon))^{\frac{N}{2+\alpha}}} \cdot \frac{q}{A} \varepsilon^\sigma + o_\varepsilon(1) \geq \frac{Cq}{A(\tau_2(w_\varepsilon))^{\frac{N}{2+\alpha}}} \geq C > 0,$$

which together with the boundedness of $\{w_\varepsilon\}$ implies the desired conclusion. \square

Lemma 5.7. *Let $N \geq 5$ and $q \in (2, 2^*)$, then there exists a $\zeta_\varepsilon \in (0, \infty)$ verifying*

$$\zeta_\varepsilon \sim \varepsilon^{-\frac{4}{(N-2)(q-2)}}$$

*such that the rescaled ground states $w_\varepsilon(x) = \zeta_\varepsilon^{\frac{N-2}{2}} u_\varepsilon(\zeta_\varepsilon x)$ converges to V_{ρ_0} in $H^1(\mathbb{R}^N)$ as $\varepsilon \rightarrow \infty$, where V_{ρ_0} is a positive ground state of the equation $-\Delta V = (I_\alpha * |V|^{2_\alpha^*}) V^{2_\alpha^*-1}$ with*

$$\rho_0 = \left(\frac{NA(2^* - q) \int_{\mathbb{R}^N} |V_1|^q}{2^* q \int_{\mathbb{R}^N} |V_1|^2} \right)^{\frac{2}{(N-2)(q-2)}}. \quad (5.20)$$

In the lower dimension cases $N = 4$ and $N = 3$, there exists $\xi_\varepsilon \in (0, +\infty)$ with $\xi_\varepsilon \rightarrow 0$ such that

$$w_\varepsilon - \xi_\varepsilon^{-\frac{N-2}{2}} V_1(\xi_\varepsilon^{-1} \cdot) \rightarrow 0$$

as $\varepsilon \rightarrow \infty$ in $D^{1,2}(\mathbb{R}^N)$ and $L^{2^*}(\mathbb{R}^N)$.

Proof. We adopt a proof of [27, Lemma 5.7]. Note that w_ε is a positive radially symmetric function, and by Lemma 5.2, $\{w_\varepsilon\}$ is bounded in $H^1(\mathbb{R}^N)$. Then there exists $w_\infty \in H^1(\mathbb{R}^N)$ verifying $-\Delta w = (I_\alpha * w^{2_\alpha^*})w^{2_\alpha^*-1}$ such that

$$w_\varepsilon \rightharpoonup w_\infty \quad \text{weakly in } H^1(\mathbb{R}^N), \quad w_\varepsilon \rightarrow w_\infty \quad \text{in } L^p(\mathbb{R}^N) \quad \text{for any } p \in (2, 2^*), \quad (5.21)$$

and

$$w_\varepsilon(x) \rightarrow w_\infty(x) \quad \text{a. e. on } \mathbb{R}^N, \quad w_\varepsilon \rightarrow w_\infty \quad \text{in } L^2_{loc}(\mathbb{R}^N). \quad (5.22)$$

Observe that

$$J_\infty(w_\varepsilon) = J_\varepsilon(w_\varepsilon) + \varepsilon^{-\sigma-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau w_\varepsilon) - \frac{\varepsilon^{-\sigma}}{2} \int_{\mathbb{R}^N} |w_\varepsilon|^2 = m_\varepsilon + o_\varepsilon(1) = m_\infty + o_\varepsilon(1),$$

and

$$J'_\infty(w_\varepsilon)w = J'_\varepsilon(w_\varepsilon)w + \varepsilon^{-\sigma-\tau(q-1)} \int_{\mathbb{R}^N} g(\varepsilon^\tau w_\varepsilon)w - \varepsilon^{-\sigma} \int_{\mathbb{R}^N} w_\varepsilon w = o_\varepsilon(1).$$

Therefore, $\{w_\varepsilon\}$ is a PS sequence of J_ε at level $m_\infty = \frac{2+\alpha}{2(N+\alpha)} S_\alpha^{\frac{N+\alpha}{2+\alpha}}$.

By Lemma 3.4, it is standard to show [34] that there exists $\zeta_\varepsilon^{(j)} \in (0, +\infty)$, $w^{(j)} \in D^{1,2}(\mathbb{R}^N)$ with $j = 1, 2, \dots, k$, k a non-negative integer, such that

$$w_\varepsilon = w_\infty + \sum_{j=1}^k (\zeta_\varepsilon^{(j)})^{-\frac{N-2}{2}} w^{(j)} ((\zeta_\varepsilon^{(j)})^{-1} x) + \tilde{w}_\varepsilon, \quad (5.23)$$

where $\tilde{w}_\varepsilon \rightarrow 0$ in $L^{2^*}(\mathbb{R}^N)$ and $w^{(j)}$ are nontrivial solutions of the limit equation $-\Delta w = (I_\alpha * w^{2_\alpha^*})w^{2_\alpha^*-1}$. Moreover, we have

$$\limsup_{\varepsilon \rightarrow \infty} \|w_\varepsilon\|_{D^1(\mathbb{R}^N)}^2 \geq \|w_\infty\|_{D^1(\mathbb{R}^N)}^2 + \sum_{j=1}^k \|w^{(j)}\|_{D^1(\mathbb{R}^N)}^2 \quad (5.24)$$

and

$$m_\infty = J_\infty(w_\infty) + \sum_{j=1}^k J_0(w^{(j)}). \quad (5.25)$$

Moreover, $J_\infty(w_\infty) \geq 0$ and $J_\infty(w^{(j)}) \geq m_\infty$ for all $j = 1, 2, \dots, k$.

If $N \geq 5$, then by Lemma 5.6, we have $w_\infty \neq 0$ and hence $J_\infty(w_\infty) = m_\infty$ and $k = 0$. Thus $w_\varepsilon \rightarrow w_\infty$ in $L^{2^*}(\mathbb{R}^N)$. Since $J'_\infty(w_\varepsilon) \rightarrow 0$, it follows that $w_\varepsilon \rightarrow w_\infty$ in $D^{1,2}(\mathbb{R}^N)$.

Since $w_\varepsilon(x)$ is radial and radially decreasing, for every $x \in \mathbb{R}^N \setminus \{0\}$, we have

$$w_\varepsilon^2(x) \leq \frac{1}{|B_{|x|}|} \int_{B_{|x|}} |w_\varepsilon|^2 \leq \frac{1}{|x|^N} \int_{\mathbb{R}^N} |w_\varepsilon|^2 \leq \frac{C}{|x|^N},$$

then

$$w_\varepsilon(x) \leq C|x|^{-\frac{N}{2}}, \quad |x| \geq 1. \quad (5.26)$$

If $\alpha > N - 4$, then we have $2_\alpha^* = \frac{N+\alpha}{N-2} > 2$ and hence

$$|w_\varepsilon|^{2_\alpha^*} |x|^N \leq C|x|^{-\frac{N}{2}2_\alpha^*+N} = C|x|^{-\frac{N}{2}(2_\alpha^*-2)} \rightarrow 0, \quad \text{as } |x| \rightarrow +\infty.$$

By virtue of Lemma 3.10, we obtain

$$(I_\alpha * |w_\varepsilon|^{2_\alpha^*})(x) \leq C|x|^{-N+\alpha}, \quad |x| \geq 1.$$

and then

$$(I_\alpha * |w_\varepsilon|^{2_\alpha^*})(x) |w_\varepsilon|^{2_\alpha^*-2}(x) \leq C|x|^{-\frac{N^2-N\alpha+4\alpha}{2(N-2)}}, \quad |x| \geq \tilde{R}. \quad (5.27)$$

As in the proof of [27, Lemma 5.7], for any small $\nu_0 > 0$, by using a comparison argument, we can find $R > 0$ such that for all large $\varepsilon > 0$ there holds

$$w_\varepsilon(x) \lesssim |x|^{-(N-2-\nu_0)}, \quad \text{for } |x| \geq R. \quad (5.28)$$

When $\nu_0 > 0$ is small enough, the domination is in $L^2(\mathbb{R}^N)$ for $N \geq 5$, and this shows, by the dominated convergence theorem, that $w_\varepsilon \rightarrow w_\infty$ in $L^2(\mathbb{R}^N)$. Thus, we conclude that $w_\varepsilon \rightarrow w_0$ in $H^1(\mathbb{R}^N)$. Moreover, by (5.4), we obtain

$$\|w_\infty\|_2^2 = \frac{NA(2^* - q)}{2^*q} \int_{\mathbb{R}^N} |w_\infty|^q,$$

from which it follows that $w_\infty = V_{\rho_0}$ with

$$\rho_0 = \left(\frac{NA(2^* - q) \int_{\mathbb{R}^N} |V_1|^q}{2^*q \int_{\mathbb{R}^N} |V_1|^2} \right)^{\frac{2}{(N-2)(q-2)}}.$$

Therefore, if $\alpha > N - 4$, then the first statement is valid with $\zeta_\varepsilon = \varepsilon^{-\frac{2}{(N-2)(q-2)}}$. If $\alpha \leq N - 4$, then for any $\varepsilon_n \rightarrow \infty$, up to a subsequence, we can assume that $w_{\varepsilon_n} \rightarrow V_\rho$ in $D^{1,2}(\mathbb{R}^N)$ with $\rho \in (0, \rho_0]$. Moreover, $w_{\varepsilon_n} \rightarrow V_\rho$ in $L^2(\mathbb{R}^N)$ if and only if $\rho = \rho_0$. Arguing as in the proof of [27, Theorem 2.1] (see also the proof of Theorem 2.1 in the present paper), we can show that there exists a $\zeta_\varepsilon \sim \varepsilon^{-\frac{2}{(N-2)(q-2)}}$ such that $w_\varepsilon(x) = \zeta_\varepsilon^{\frac{N-2}{2}} u_\varepsilon(\zeta_\varepsilon x)$ converges to V_{ρ_0} in $L^2(\mathbb{R}^N)$, and hence in $H^1(\mathbb{R}^N)$.

If $N = 4$ or 3 . By Fatou's lemma, we have $\|w_\infty\|_2^2 \leq \liminf_{\varepsilon \rightarrow \infty} \|v_\varepsilon\|_2^2 < \infty$, therefore, $w_\infty = 0$ and hence $k = 1$. Thus, we obtain $J_\infty(w^{(1)}) = m_\infty$ and hence $w^{(1)} = V_\rho$ for some $\rho \in (0, +\infty)$. Therefore, we conclude that

$$w_\varepsilon - \xi_\varepsilon^{-\frac{N-2}{2}} V_1(\xi_\varepsilon^{-1} \cdot) \rightarrow 0$$

in $L^{2^*}(\mathbb{R}^N)$ as $\lambda \rightarrow 0$, where $\xi_\varepsilon := \rho \xi_\varepsilon^{(1)} \in (0, +\infty)$ satisfying $\xi_\varepsilon \rightarrow 0$ as $\varepsilon \rightarrow \infty$. Since

$$J'_\infty(w_\varepsilon - \xi_\varepsilon^{-\frac{N-2}{2}} V_1(\xi_\varepsilon^{-1} \cdot)) = J'_\infty(v_\varepsilon) + J'_\infty(V_1) + o_\varepsilon(1) = o_\varepsilon(1)$$

as $\varepsilon \rightarrow \infty$, it follows that $w_\varepsilon - \xi_\varepsilon^{-\frac{N-2}{2}} V_1(\xi_\varepsilon^{-1} \cdot) \rightarrow 0$ in $D^{1,2}(\mathbb{R}^N)$. \square

5.2. The second scaling. In the lower dimension cases $N = 4$ and $N = 3$, we further perform a scaling

$$\tilde{w}(x) = \xi_\varepsilon^{\frac{N-2}{2}} w(\xi_\varepsilon x), \quad (5.29)$$

where $\xi_\varepsilon \in (0, +\infty)$ is given in Lemma 5.7. Then the rescaled equation is as follows

$$-\Delta \tilde{w} + \varepsilon^{-\sigma} \xi_\varepsilon^2 \tilde{w} = (I_\alpha * |\tilde{w}|^{2_\alpha^*}) \tilde{w}^{2_\alpha^*-1} + \varepsilon^{-\sigma} \xi_\varepsilon^{N-\frac{N-2}{2}q} (\varepsilon^{-\tau} \xi_\varepsilon^{\frac{N-2}{2}})^{q-1} g(\varepsilon^\tau \xi_\varepsilon^{-\frac{N-2}{2}} \tilde{w}). \quad (5.30)$$

The corresponding energy functional is given by

$$\begin{aligned} \tilde{J}_\varepsilon(\tilde{w}) : &= \frac{1}{2} \int_{\mathbb{R}^N} |\nabla \tilde{w}|^2 + \varepsilon^{-\sigma} \xi_\varepsilon^2 |\tilde{w}|^2 - \frac{1}{22_\alpha^*} \int_{\mathbb{R}^N} (I_\alpha * |\tilde{w}|^{2_\alpha^*}) |\tilde{w}|^{2_\alpha^*} \\ &\quad - \varepsilon^{-\sigma} \xi_\varepsilon^{N-\frac{N-2}{2}q} (\varepsilon^{-\tau} \xi_\varepsilon^{\frac{N-2}{2}})^q \int_{\mathbb{R}^N} G(\varepsilon^\tau \xi_\varepsilon^{-\frac{N-2}{2}} \tilde{w}). \end{aligned} \quad (5.31)$$

Clearly, we have $\tilde{J}_\varepsilon(\tilde{w}) = J_\varepsilon(w) = I_\varepsilon(u)$.

Furthermore, we have the following lemma.

Lemma 5.8. *Let $u, w, \tilde{w} \in H^1(\mathbb{R}^N)$ satisfy (5.1) and (5.29), then the following statements hold true*

$$(1) \quad \|\nabla \tilde{w}\|_2^2 = \|\nabla w\|_2^2 = \|\nabla u\|_2^2, \quad \int_{\mathbb{R}^N} (I_\alpha * |\tilde{w}|^{2_\alpha^*}) |\tilde{w}|^{2_\alpha^*} = \int_{\mathbb{R}^N} (I_\alpha * |w|^{2_\alpha^*}) |w|^{2_\alpha^*} = \int_{\mathbb{R}^N} (I_\alpha * |u|^{2_\alpha^*}) |u|^{2_\alpha^*},$$

$$(2) \quad \xi_\varepsilon^2 \|\tilde{w}\|_2^2 = \|w\|_2^2 = \varepsilon^{1+\sigma} \|u\|_2^2, \quad \xi_\varepsilon^{N-\frac{N-2}{2}q} \|\tilde{w}\|_q^q = \|w\|_q^q = \varepsilon^\sigma \|u\|_q^q.$$

Set $\tilde{w}_\varepsilon(x) = \xi_\varepsilon^{\frac{N-2}{2}} w_\varepsilon(\xi_\varepsilon x)$, then by Lemma 5.7, we have

$$\|\nabla(\tilde{w}_\varepsilon - V_1)\|_2 \rightarrow 0, \quad \|\tilde{w}_\varepsilon - V_1\|_{2^*} \rightarrow 0, \quad \text{as } \varepsilon \rightarrow \infty. \quad (5.32)$$

Note that the corresponding Nehari and Pohožaev's identities are as follows

$$\begin{aligned} &\int_{\mathbb{R}^N} |\nabla \tilde{w}_\varepsilon|^2 + \varepsilon^{-\sigma} \xi_\varepsilon^2 \int_{\mathbb{R}^N} |\tilde{w}_\varepsilon|^2 \\ &= \int_{\mathbb{R}^N} (I_\alpha * |\tilde{w}_\varepsilon|^{2_\alpha^*}) |\tilde{w}_\varepsilon|^{2_\alpha^*} + \varepsilon^{-\sigma} \xi_\varepsilon^{N-\frac{N-2}{2}q} (\varepsilon^{-\tau} \xi_\varepsilon^{\frac{N-2}{2}})^{q-1} \int_{\mathbb{R}^N} g(\varepsilon^\tau \xi_\varepsilon^{-\frac{N-2}{2}} \tilde{w}_\varepsilon) \tilde{w}_\varepsilon. \end{aligned} \quad (5.33)$$

and

$$\begin{aligned} &\frac{1}{2^*} \int_{\mathbb{R}^N} |\nabla \tilde{w}_\varepsilon|^2 + \frac{1}{2} \varepsilon^{-\sigma} \xi_\varepsilon^2 \int_{\mathbb{R}^N} |\tilde{w}_\varepsilon|^2 \\ &= \frac{1}{2^*} \int_{\mathbb{R}^N} (I_\alpha * |\tilde{w}_\varepsilon|^{2_\alpha^*}) |\tilde{w}_\varepsilon|^{2_\alpha^*} + \varepsilon^{-\sigma} \xi_\varepsilon^{N-\frac{N-2}{2}q} (\varepsilon^{-\tau} \xi_\varepsilon^{\frac{N-2}{2}})^q \int_{\mathbb{R}^N} G(\varepsilon^\tau \xi_\varepsilon^{-\frac{N-2}{2}} \tilde{w}_\varepsilon), \end{aligned} \quad (5.34)$$

it follows that

$$\begin{aligned} \frac{1}{N} \varepsilon^{-\sigma} \xi_\varepsilon^2 \int_{\mathbb{R}^N} |\tilde{w}_\varepsilon|^2 &= \varepsilon^{-\sigma-\tau q} \xi_\varepsilon^N \int_{\mathbb{R}^N} [G(\varepsilon^\tau \xi_\varepsilon^{-\frac{N-2}{2}} \tilde{w}_\varepsilon) - \frac{1}{2^*} g(\varepsilon^\tau \xi_\varepsilon^{-\frac{N-2}{2}} \tilde{w}_\varepsilon) \varepsilon^\tau \xi_\varepsilon^{-\frac{N-2}{2}} \tilde{w}_\varepsilon] \\ &= \varepsilon^{-\sigma} \xi_\varepsilon^{N-\frac{N-2}{2}q} \left[\frac{A(2^*-q)}{2^*q} \int_{\mathbb{R}^N} |\tilde{w}_\varepsilon|^q + o_\varepsilon(1) \right]. \end{aligned}$$

Thus, we obtain

$$\xi_\varepsilon^{\frac{(N-2)(q-2)}{2}} \int_{\mathbb{R}^N} |\tilde{w}_\varepsilon|^2 = \frac{NA(2^*-q)}{2^*q} \int_{\mathbb{R}^N} |\tilde{w}_\varepsilon|^q + o_\varepsilon(1). \quad (5.35)$$

To control the norm $\|\tilde{w}_\varepsilon\|_2$, we note that for any $\varepsilon > 0$, $\tilde{w}_\varepsilon > 0$ satisfies the linear inequality

$$-\Delta \tilde{w}_\varepsilon + \varepsilon^{-\sigma} \xi_\varepsilon^2 \tilde{w}_\varepsilon > 0. \quad (5.36)$$

Lemma 5.9 *There exists a constant $c > 0$ such that*

$$\tilde{w}_\varepsilon(x) \geq c|x|^{-(N-2)} \exp(-\varepsilon^{-\frac{\sigma}{2}} \xi_\varepsilon |x|), \quad |x| \geq 1. \quad (5.37)$$

The proof of the above lemma is similar to that of [28, Lemma 4.8]. As consequences, we have the following lemma.

Lemma 5.10. *If $N = 3$, then $\|\tilde{w}_\varepsilon\|_2^2 \gtrsim \varepsilon^{\frac{\sigma}{2}} \xi_\varepsilon^{-1}$.*

Lemma 5.11. *If $N = 4$, then $\|\tilde{w}_\varepsilon\|_2^2 \gtrsim -\ln(\varepsilon^{-\sigma} \xi_\varepsilon^2)$.*

We remark that \tilde{w}_ε is only defined for $N = 4$ and $N = 3$. But the following discussion also apply to the case $N \geq 5$. To prove our main result, the key point is to show the boundedness of $\|\tilde{w}_\varepsilon\|_q$. We begin with a technique lemma.

Lemma 5.12. *Assume $N \geq 3$, $\alpha > N - 4$ and $2 < q < 2^*$. Then there exist constants $L_0 > 0$ and $C_0 > 0$ such that for any large $\varepsilon > 0$ and $|x| \geq L_0 \varepsilon^{\sigma/2} \xi_\varepsilon^{-1}$,*

$$\tilde{w}_\varepsilon(x) \leq C_0 \varepsilon^{-\sigma(N-2)/4} \xi_\varepsilon^{(N-2)/2} \exp\left(-\frac{1}{2} \varepsilon^{-\sigma/2} \xi_\varepsilon |x|\right).$$

Proof. By (5.26) and (5.27), if $|x| \geq L_0 \varepsilon^{\sigma/2} \xi_\varepsilon^{-1}$ with $L_0 > 0$ being large enough, we have

$$\begin{aligned} (I_\alpha * |\tilde{w}_\varepsilon|^{2^*})(x) |\tilde{w}_\varepsilon|^{2^*-2}(x) &= \xi_\varepsilon^{(N-2)(2^*-1)-\alpha} (I_\alpha * |w_\varepsilon|^{2^*})(\xi_\varepsilon x) |w_\varepsilon|^{2^*-2}(\xi_\varepsilon x) \\ &\leq C \xi_\varepsilon^2 L_0^{-\frac{N^2-N\alpha+4\alpha}{2(N-2)}} \varepsilon^{-\sigma \cdot \frac{N^2-N\alpha+4\alpha}{4(N-2)}} \\ &\leq \frac{1}{4} \varepsilon^{-\sigma} \xi_\varepsilon^2, \end{aligned}$$

here we have used the fact that

$$\frac{N^2 - N\alpha + 4\alpha}{4(N-2)} > 1,$$

which follows from the inequality $N < N + 2 < \frac{4(\alpha+2)}{\alpha-N+4}$, $\forall \alpha \in (N-4, N)$.

By (H1) and (H2), for any $\delta > 0$, there exists $C_\delta > 0$ such that

$$g(s) \leq \delta s + C_\delta s^{q-1}, \quad \text{for } s \geq 0.$$

Therefore, we have

$$(\varepsilon^{-\tau} \xi_\varepsilon^{\frac{N-2}{2}})^{q-1} g(\varepsilon^\tau \xi_\varepsilon^{-\frac{N-2}{2}} \tilde{w}_\varepsilon(x)) \leq \delta (\varepsilon^{-\tau} \xi_\varepsilon^{\frac{N-2}{2}})^{q-2} \tilde{w}_\varepsilon + C_\delta \tilde{w}_\varepsilon^{q-1}.$$

By (5.24) and (5.26), for $|x| \geq L_0 \varepsilon^{\sigma/2} \xi_\varepsilon^{-1}$, we get

$$\begin{aligned} \varepsilon^{-\sigma} \xi_\varepsilon^{N-\frac{N-2}{2}q} (\varepsilon^{-\tau} \xi_\varepsilon^{\frac{N-2}{2}})^{q-1} g(\varepsilon^\tau \xi_\varepsilon^{-\frac{N-2}{2}} \tilde{w}_\varepsilon(x)) &\leq \delta \varepsilon^{-\sigma-\tau(q-2)} \xi_\varepsilon^2 \tilde{w}_\varepsilon + C_\delta \varepsilon^{-\sigma} \xi_\varepsilon^{N-\frac{N-2}{2}q} \tilde{w}_\varepsilon^{q-1} \\ &\leq \frac{1}{4} \varepsilon^{-\sigma} \xi_\varepsilon^2 \tilde{w}_\varepsilon(x). \end{aligned}$$

Therefore, we obtain

$$-\Delta \tilde{w}_\varepsilon(x) + \frac{1}{2} \varepsilon^{-\sigma} \xi_\varepsilon^2 \tilde{w}_\varepsilon(x) \leq 0, \quad \text{for all } |x| \geq L_0 \varepsilon^{\sigma/2} \xi_\varepsilon^{-1}.$$

Arguing as in [1, Lemma 3.2], we prove that

$$\tilde{w}_\varepsilon(x) \leq C L_0^{-\frac{N-2}{2}} e^{L_0/2} \varepsilon^{-\frac{\sigma(N-2)}{4}} \xi_\varepsilon^{\frac{N-2}{2}} e^{-\frac{1}{2} \varepsilon^{-\sigma/2} \xi_\varepsilon |x|},$$

for all $|x| \geq L_0 \varepsilon^{\sigma/2} \xi_\varepsilon^{-1}$. The proof is complete. \square

To obtain the optimal decay estimate of \tilde{w}_ε at infinity, we consider the Kelvin transform of \tilde{w}_ε . For any $w \in H^1(\mathbb{R}^N)$, we denote by $K[w]$ the Kelvin transform of w , that is,

$$K[w](x) := |x|^{-(N-2)} w\left(\frac{x}{|x|^2}\right).$$

It is easy to see that $\|K[\tilde{w}_\lambda]\|_{L^\infty(B_1)} \lesssim 1$ implies that

$$\tilde{w}_\varepsilon(x) \lesssim |x|^{-(N-2)}, \quad |x| \geq 1,$$

uniformly for large $\varepsilon > 0$. Thus, it suffices to show that there exists $\varepsilon_0 > 0$ such that

$$\sup_{\varepsilon \in (\varepsilon_0, \infty)} \|K[\tilde{w}_\varepsilon]\|_{L^\infty(B_1)} < \infty. \quad (5.38)$$

It is easy to verify that $K[\tilde{w}_\varepsilon]$ satisfies

$$\begin{aligned} -\Delta K[\tilde{w}_\varepsilon] + \frac{\varepsilon^{-\sigma} \xi_\varepsilon^2}{|x|^4} K[\tilde{w}_\varepsilon] &= \frac{1}{|x|^4} (I_\alpha * |\tilde{w}_\varepsilon|^{2_\alpha^*}) \left(\frac{x}{|x|^2}\right) \tilde{w}_\varepsilon^{2_\alpha^*-2} \left(\frac{x}{|x|^2}\right) K[\tilde{w}_\varepsilon] \\ &\quad + \frac{1}{|x|^4} \varepsilon^{-\sigma-\tau(q-1)} \xi_\varepsilon^{\frac{N+2}{2}} \frac{g(\varepsilon^\tau \xi_\varepsilon^{-\frac{N-2}{2}} \tilde{w}_\varepsilon(\frac{x}{|x|^2}))}{\tilde{w}_\varepsilon(\frac{x}{|x|^2})} K[\tilde{w}_\varepsilon]. \end{aligned} \quad (5.39)$$

We also see from Lemma 5.12 that if $|x| \leq \varepsilon^{-\sigma/2} \xi_\varepsilon / L_0$, then

$$K[\tilde{w}_\varepsilon](x) \lesssim \frac{1}{|x|^{N-2}} \varepsilon^{\frac{-\sigma(N-2)}{4}} \xi_\varepsilon^{\frac{N-2}{2}} e^{-\frac{1}{2}\varepsilon^{-\sigma/2} \xi_\varepsilon |x|^{-1}}. \quad (5.40)$$

Let $\tilde{a}(x) = \frac{\varepsilon^{-\sigma} \xi_\varepsilon^2}{|x|^4}$ and

$$\tilde{b}(x) = \frac{1}{|x|^4} (I_\alpha * |\tilde{w}_\varepsilon|^{2_\alpha^*}) \left(\frac{x}{|x|^2}\right) \tilde{w}_\varepsilon^{2_\alpha^*-2} \left(\frac{x}{|x|^2}\right) + \frac{1}{|x|^4} \varepsilon^{-\sigma-\tau(q-1)} \xi_\varepsilon^{\frac{N+2}{2}} \frac{g(\varepsilon^\tau \xi_\varepsilon^{-\frac{N-2}{2}} \tilde{w}_\varepsilon(\frac{x}{|x|^2}))}{\tilde{w}_\varepsilon(\frac{x}{|x|^2})}.$$

Then (5.39) reads as

$$-\Delta K[\tilde{w}_\varepsilon] + \tilde{a}(x) K[\tilde{w}_\varepsilon] = \tilde{b}(x) K[\tilde{w}_\varepsilon]. \quad (5.41)$$

Since $g(s) = o(s)$ as $s \rightarrow 0$ and $\lim_{s \rightarrow +\infty} g(s) s^{1-q} = A > 0$, it follows that

$$g(s) \leq C(|s|^{q-1} + |s|), \quad \forall s \geq 0.$$

then we have

$$\begin{aligned} \tilde{b}(x) &\leq \frac{1}{|x|^4} (I_\alpha * |\tilde{w}_\varepsilon|^{2_\alpha^*}) \left(\frac{x}{|x|^2}\right) \tilde{w}_\varepsilon^{2_\alpha^*-2} \left(\frac{x}{|x|^2}\right) \\ &\quad + \frac{1}{|x|^4} \varepsilon^{-\sigma} \xi_\varepsilon^{N-\frac{N-2}{2}q} |\tilde{w}_\varepsilon(\frac{x}{|x|^2})|^{q-2} + \frac{1}{|x|^4} \varepsilon^{-\sigma-\tau(q-2)} \xi_\varepsilon^2 \\ &= \frac{1}{|x|^4} (I_\alpha * |\tilde{w}_\varepsilon|^{2_\alpha^*}) \left(\frac{x}{|x|^2}\right) \tilde{w}_\varepsilon^{2_\alpha^*-2} \left(\frac{x}{|x|^2}\right) \\ &\quad + \frac{1}{|x|^{\gamma q}} \varepsilon^{-\sigma} \xi_\varepsilon^{\gamma q/2} K[\tilde{w}_\varepsilon]^{q-1} + \frac{1}{|x|^4} \varepsilon^{-\sigma-\tau(q-2)} \xi_\varepsilon^2, \end{aligned}$$

here and in what follows, for any $q \in (2, 2^*)$, we set $\gamma_q = 2N - (N-2)q$. Choose $\eta(x) \in [0, 1]$ such that

$$\begin{aligned} \tilde{b}(x) &= \eta(x) \left\{ \frac{1}{|x|^4} (I_\alpha * |\tilde{w}_\varepsilon|^{2_\alpha^*}) \left(\frac{x}{|x|^2}\right) \tilde{w}_\varepsilon^{2_\alpha^*-2} \left(\frac{x}{|x|^2}\right) \right. \\ &\quad \left. + \frac{1}{|x|^{\gamma q}} \varepsilon^{-\sigma} \xi_\varepsilon^{\gamma q/2} K[\tilde{w}_\varepsilon]^{q-1} + \frac{1}{|x|^4} \varepsilon^{-\sigma-\tau(q-2)} \xi_\varepsilon^2 \right\}. \end{aligned} \quad (5.42)$$

Let

$$a(x) = (1 - \varepsilon^{-\tau(q-2)} \eta(x)) \frac{\varepsilon^{-\sigma} \xi_\varepsilon^2}{|x|^4},$$

$$b(x) = \frac{\eta(x)}{|x|^4} (I_\alpha * |\tilde{w}_\varepsilon|^{2_\alpha^*}) \left(\frac{x}{|x|^2}\right) \tilde{w}_\varepsilon^{2_\alpha^*-2} \left(\frac{x}{|x|^2}\right) + \frac{\eta(x)}{|x|^{\gamma q}} \varepsilon^{-\sigma} \xi_\varepsilon^{\gamma q/2} K[\tilde{w}_\varepsilon]^{q-1}.$$

Then (5.41) and (5.42) yields

$$-\Delta K[\tilde{w}_\varepsilon] + a(x) K[\tilde{w}_\varepsilon] = b(x) K[\tilde{w}_\varepsilon]. \quad (5.43)$$

Adopting the Moser iteration, arguing as in [27, Proposition 5.14], we obtain the following

Proposition 5.13. *Assume $N \geq 3$, $\alpha > N - 4$ and $2 < q < 2^*$. Then there exists a constant $C > 0$ such that for large $\varepsilon > 0$, there holds*

$$\tilde{w}_\varepsilon(x) \leq C(1 + |x|)^{-(N-2)}, \quad x \in \mathbb{R}^N.$$

Lemma 5.14. *If $s > \frac{N}{N-2}$, then $\|\tilde{w}_\varepsilon\|_s^s \sim 1$ as $\varepsilon \rightarrow \infty$. Furthermore, $\tilde{w}_\varepsilon \rightarrow V_1$ in $L^s(\mathbb{R}^N)$ as $\varepsilon \rightarrow \infty$.*

Proof. Since $\tilde{w}_\varepsilon \rightarrow V_1$ in $L^{2^*}(\mathbb{R}^N)$, as in [28, Lemma 4.6], using the embeddings $L^{2^*}(B_1) \hookrightarrow L^s(B_1)$ we prove that $\liminf_{\varepsilon \rightarrow \infty} \|\tilde{w}_\varepsilon\|_s^s > 0$.

On the other hand, by virtue of Proposition 5.13, there exists a constant $C > 0$ such that for all large $\varepsilon > 0$,

$$\tilde{w}_\varepsilon(x) \leq \frac{C}{(1 + |x|)^{N-2}}, \quad \forall x \in \mathbb{R}^N,$$

which together with the fact that $s > \frac{N}{N-2}$ implies that \tilde{w}_ε is bounded in $L^s(\mathbb{R}^N)$ uniformly for large $\varepsilon > 0$, and by the dominated convergence theorem $\tilde{w}_\varepsilon \rightarrow V_1$ in $L^s(\mathbb{R}^N)$ as $\varepsilon \rightarrow \infty$. \square

5.3. The Proof of Theorem 2.2. We are in the position to prove Theorem 2.2.

Proof of Theorem 2.2. Firstly, we consider the cases $N = 4$ and $N = 3$. We note that a result similar to Lemma 3.2 holds for \tilde{w}_ε and \tilde{J}_ε . By Lemma 5.8, we also have $\tau_2(\tilde{w}_\varepsilon) = \tau_2(v_\varepsilon)$. Therefore, by (5.35), we get

$$\begin{aligned} m_\infty &\leq \sup_{t \geq 0} \tilde{J}_\varepsilon((\tilde{w}_\varepsilon)_t) \\ &\quad + \varepsilon^{-\sigma} \tau_2(\tilde{w}_\varepsilon)^{\frac{N}{2}} \left\{ \xi_\varepsilon^{N - \frac{N-2}{2}q} (\varepsilon^{-\tau} \xi_\varepsilon^{\frac{N-2}{2}})^q \int_{\mathbb{R}^N} G(\varepsilon^\tau \xi_\varepsilon^{-\frac{N-2}{2}} \tilde{w}_\varepsilon) - \frac{1}{2} \xi_\varepsilon^2 \int_{\mathbb{R}^N} |\tilde{w}_\varepsilon|^2 \right\} \\ &\quad + \varepsilon^{-\sigma} \tau_2(\tilde{w}_\varepsilon)^{\frac{N}{2}} \left\{ \xi_\varepsilon^N \varepsilon^{-\tau q} \int_{\mathbb{R}^N} G(\varepsilon^\tau \xi_\varepsilon^{-\frac{N-2}{2}} \tilde{w}_\varepsilon) - \frac{1}{2} \xi_\varepsilon^2 \int_{\mathbb{R}^N} |\tilde{w}_\varepsilon|^2 \right\} \\ &\leq m_\varepsilon + \varepsilon^{-\sigma} \tau_2(\tilde{w}_\varepsilon)^{\frac{N}{2}} \frac{A}{q} \xi_\varepsilon^{N - \frac{N-2}{2}q} (\int_{\mathbb{R}^N} |\tilde{w}_\varepsilon|^q + o_\varepsilon(1)), \end{aligned} \tag{5.44}$$

which implies that

$$\xi_\varepsilon^{N - \frac{N-2}{2}q} \left(\int_{\mathbb{R}^N} |\tilde{w}_\varepsilon|^q + o_\varepsilon(1) \right) \geq \varepsilon^\sigma \frac{q}{A \tau_2(w_\varepsilon)^{\frac{N}{2}}} \delta_\varepsilon.$$

Hence, by Lemma 5.5, we obtain

$$\xi_\varepsilon^{N - \frac{N-2}{2}q} \left(\int_{\mathbb{R}^N} |\tilde{w}_\varepsilon|^q + o_\varepsilon(1) \right) \gtrsim \varepsilon^\sigma \delta_\varepsilon \gtrsim \begin{cases} (\ln \varepsilon)^{-\frac{4-q}{q-2}}, & \text{if } N = 4, \\ \varepsilon^{-\frac{(6-q)^2}{2(q-2)(q-4)}}, & \text{if } N = 3. \end{cases} \tag{5.45}$$

Therefore, by Lemma 5.14, we have

$$\xi_\varepsilon \gtrsim \begin{cases} (\ln \varepsilon)^{-\frac{1}{q-2}}, & \text{if } N = 4, \\ \varepsilon^{-\frac{6-q}{(q-2)(q-4)}}, & \text{if } N = 3. \end{cases} \tag{5.46}$$

On the other hand, if $N = 3$, then by (5.35) and Lemma 5.10 and Lemma 5.14, we have

$$\varepsilon^{\sigma/2} \xi_\varepsilon^{-1 + \frac{q-2}{2}} \lesssim \xi_\varepsilon^{\frac{q-2}{2}} \|\tilde{w}_\lambda\|_2^2 \lesssim 1.$$

Then, noting that $\sigma = \frac{2^*-q}{q-2} = \frac{6-q}{q-2}$, for $q \in (4, 6)$, we have

$$\xi_\varepsilon^{\frac{q-4}{2}} \lesssim \varepsilon^{-\frac{\sigma}{2}} = \varepsilon^{-\frac{6-q}{2(q-2)}},$$

and hence

$$\xi_\varepsilon \lesssim \varepsilon^{-\frac{6-q}{(q-2)(q-4)}}. \quad (5.47)$$

If $N = 4$, noting that

$$-\ln(\varepsilon^{-\sigma} \xi_\varepsilon^2) = \sigma \ln \varepsilon - 2 \ln \xi_\varepsilon \geq \sigma \ln \varepsilon,$$

by (5.35) and Lemma 5.11 and Lemma 5.14, we have

$$\xi_\varepsilon^{q-2} \ln \varepsilon \lesssim \xi_\varepsilon^{q-2} \|\tilde{w}_\lambda\|_2^2 \lesssim 1.$$

Hence, we obtain

$$\xi_\varepsilon \lesssim (\ln \varepsilon)^{-\frac{1}{q-2}}. \quad (5.48)$$

Thus, it follows from (5.44), (5.47), (5.48) and Lemma 5.14 that

$$\delta_\varepsilon = m_\infty - m_\varepsilon \lesssim \varepsilon^{-\sigma} \xi_\varepsilon^{N-\frac{N-2}{2}q} \lesssim \begin{cases} (\varepsilon \ln \varepsilon)^{-\frac{4-q}{q-2}}, & \text{if } N = 4, \\ \varepsilon^{-\frac{6-q}{2(q-4)}}, & \text{if } N = 3, \end{cases}$$

which together with Lemma 5.5 implies that

$$\delta_\varepsilon \sim \varepsilon^{-\sigma} \xi_\varepsilon^{N-\frac{N-2}{2}q} \sim \begin{cases} (\varepsilon \ln \varepsilon)^{-\frac{4-q}{q-2}}, & \text{if } N = 4, \\ \varepsilon^{-\frac{6-q}{2(q-4)}}, & \text{if } N = 3. \end{cases} \quad (5.49)$$

By (5.31) and (5.35), we get

$$\begin{aligned} m_\varepsilon &= \frac{1}{2} \int_{\mathbb{R}^N} |\nabla \tilde{w}_\varepsilon|^2 - \frac{1}{22_\alpha^*} \int_{\mathbb{R}^N} (I_\alpha * |\tilde{w}_\varepsilon|^{2_\alpha^*}) |\tilde{w}_\varepsilon|^{2_\alpha^*} \\ &\quad - \frac{A(N-2)(q-2)}{4q} \varepsilon^{-\sigma} \xi_\varepsilon^{N-\frac{N-2}{2}q} [\int_{\mathbb{R}^N} |\tilde{w}_\varepsilon|^q + o_\varepsilon(1)]. \end{aligned}$$

By (5.33) and (5.34), we get

$$\int_{\mathbb{R}^N} |\nabla \tilde{w}_\varepsilon|^2 = \int_{\mathbb{R}^N} (I_\alpha * |\tilde{w}_\varepsilon|^{2_\alpha^*}) |\tilde{w}_\varepsilon|^{2_\alpha^*} + \frac{NA(q-2)}{2q} \varepsilon^{-\sigma} \xi_\varepsilon^{N-\frac{N-2}{2}q} [\int_{\mathbb{R}^N} |\tilde{w}_\varepsilon|^q + o_\varepsilon(1)].$$

Therefore, we have

$$m_\varepsilon = \frac{2+\alpha}{2(N+\alpha)} \int_{\mathbb{R}^N} |\nabla \tilde{w}_\varepsilon|^2 - \frac{A\alpha(N-2)(q-2)}{4q(N+\alpha)} \varepsilon^{-\sigma} \xi_\varepsilon^{N-\frac{N-2}{2}q} [\int_{\mathbb{R}^N} |\tilde{w}_\varepsilon|^q + o_\varepsilon(1)].$$

Similarly, we have

$$m_\infty = \frac{2+\alpha}{2(N+\alpha)} \int_{\mathbb{R}^N} |\nabla V_1|^2.$$

Thus, by virtue of (5.49), we obtain

$$\begin{aligned} \|\nabla V_1\|_2^2 - \|\nabla \tilde{w}_\varepsilon\|_2^2 &= \frac{2(N+\alpha)}{2+\alpha} \delta_\varepsilon - \frac{A\alpha(N-2)(q-2)}{2q(2+\alpha)} \varepsilon^{-\sigma} \xi_\varepsilon^{N-\frac{N-2}{2}q} [\int_{\mathbb{R}^N} |\tilde{w}_\varepsilon|^q + o_\varepsilon(1)] \\ &= \begin{cases} O((\varepsilon \ln \varepsilon)^{-\frac{4-q}{q-2}}), & \text{if } N = 4, \\ O(\varepsilon^{-\frac{6-q}{2(q-4)}}), & \text{if } N = 3. \end{cases} \end{aligned} \quad (5.50)$$

By (5.31) and (5.34), we have

$$\begin{aligned} m_\varepsilon &= (\frac{1}{2} - \frac{1}{2^*}) \int_{\mathbb{R}^N} |\nabla \tilde{w}_\varepsilon|^2 + (\frac{1}{2^*} - \frac{1}{22_\alpha^*}) \int_{\mathbb{R}^N} (I_\alpha * |\tilde{w}_\varepsilon|^{2_\alpha^*}) |\tilde{w}_\varepsilon|^{2_\alpha^*} \\ &= \frac{1}{N} \int_{\mathbb{R}^N} |\nabla \tilde{w}_\varepsilon|^2 + \frac{\alpha(N-2)}{2N(N+\alpha)} \int_{\mathbb{R}^N} (I_\alpha * |\tilde{w}_\varepsilon|^{2_\alpha^*}) |\tilde{w}_\varepsilon|^{2_\alpha^*}. \end{aligned}$$

Similarly, we also have

$$m_\infty = \frac{1}{N} \int_{\mathbb{R}^N} |\nabla V_1|^2 + \frac{\alpha(N-2)}{2N(N+\alpha)} \int_{\mathbb{R}^N} (I_\alpha * |V_1|^{2_\alpha^*}) |V_1|^{2_\alpha^*}.$$

Then it follows from (5.49) and (5.50) that

$$\begin{aligned} & \int_{\mathbb{R}^N} (I_\alpha * |V_1|^{2_\alpha^*}) |V_1|^{2_\alpha^*} - \int_{\mathbb{R}^N} (I_\alpha * |\tilde{w}_\varepsilon|^{2_\alpha^*}) |\tilde{w}_\varepsilon|^{2_\alpha^*} \\ &= \frac{2N(N+\alpha)}{\alpha(N-2)} \left[(m_\infty - m_\varepsilon) - \frac{1}{N} \left(\int_{\mathbb{R}^N} |\nabla V_1|^2 - \int_{\mathbb{R}^N} |\nabla \tilde{w}_\varepsilon|^2 \right) \right] \\ &= \frac{2N(N+\alpha)}{\alpha(N-2)} \delta_\varepsilon - \frac{2(N+\alpha)}{\alpha(N-2)} \left(\int_{\mathbb{R}^N} |\nabla V_1|^2 - \int_{\mathbb{R}^N} |\nabla \tilde{w}_\varepsilon|^2 \right) \\ &= \begin{cases} O((\varepsilon \ln \varepsilon)^{-\frac{4-q}{q-2}}), & \text{if } N = 4, \\ O(\varepsilon^{-\frac{6-q}{2(q-4)}}), & \text{if } N = 3. \end{cases} \end{aligned} \quad (5.51)$$

Since $\|\nabla V_1\|_2^2 = \int_{\mathbb{R}^N} (I_\alpha * |V_1|^{2_\alpha^*}) |V_1|^{2_\alpha^*} = S_\alpha^{\frac{N+\alpha}{2+\alpha}}$, it follows from (5.61) and (5.62) that

$$\|\nabla \tilde{w}_\varepsilon\|_2^2 = S_\alpha^{\frac{N+\alpha}{2+\alpha}} + \begin{cases} O((\varepsilon \ln \varepsilon)^{-\frac{4-q}{q-2}}), & \text{if } N = 4, \\ O(\varepsilon^{-\frac{6-q}{2(q-4)}}), & \text{if } N = 3, \end{cases}$$

and

$$\int_{\mathbb{R}^N} (I_\alpha * |\tilde{w}_\varepsilon|^{2_\alpha^*}) |\tilde{w}_\varepsilon|^{2_\alpha^*} = S_\alpha^{\frac{N+\alpha}{2+\alpha}} + \begin{cases} O((\varepsilon \ln \varepsilon)^{-\frac{4-q}{q-2}}), & \text{if } N = 4, \\ O(\varepsilon^{-\frac{6-q}{2(q-4)}}), & \text{if } N = 3. \end{cases}$$

Finally, by (5.35), Lemma 5.10 and Lemma 5.11, we obtain

$$\|\tilde{w}_\varepsilon\|_2^2 \sim \begin{cases} \ln \varepsilon, & \text{if } N = 4, \\ \varepsilon^{\frac{6-q}{2(q-4)}}, & \text{if } N = 3. \end{cases}$$

For $N \geq 5$, the conclusion follows directly from Lemmas 5.5, 5.6 and 5.8. More precisely, Replacing $V_1, \tilde{w}_\varepsilon$ by V_{ρ_0} and v_ε respectively, and setting $\xi_\varepsilon = 1$ in the above arguments, we obtain

$$\|\nabla u_\varepsilon\|_2^2 = \|\nabla v_\varepsilon\|_2^2 = S_\alpha^{\frac{N+\alpha}{2+\alpha}} + O(\varepsilon^{-\frac{2N-q(N-2)}{(N-2)(q-2)}})$$

and

$$\int_{\mathbb{R}^N} (I_\alpha * |u_\varepsilon|^{2_\alpha^*}) |u_\varepsilon|^{2_\alpha^*} = \int_{\mathbb{R}^N} (I_\alpha * |v_\varepsilon|^{2_\alpha^*}) |v_\varepsilon|^{2_\alpha^*} = S_\alpha^{\frac{N+\alpha}{2+\alpha}} + O(\varepsilon^{-\frac{2N-q(N-2)}{(N-2)(q-2)}}).$$

The statements on u_ε follow from the corresponding results on v_ε and \tilde{w}_ε . This completes the proof of Theorem 2.2. \square

6. FINAL REMARKS

Clearly, if $u_\varepsilon \in H^1(\mathbb{R}^N)$ is a ground state of (P_ε) , and for some $a > 0$ there holds

$$\|u_\varepsilon\|_2^2 = a^2, \quad (6.1)$$

then u_ε is a normalized solution of (1.6) with $\lambda = -\varepsilon$. We denote this normalized solution by a pair (u_a, λ_a) with $\lambda_a = -\varepsilon$, or u_a for simplicity.

As direct consequences of Theorem 2.1 and Theorem 2.2, we have the following results.

Proposition 6.1. *Assume that (H1) and (H2) hold. If $p = \frac{N+\alpha}{N}$, $q \in (2, 2 + \frac{4}{N})$, then for any large $a > 0$, the problem (1.6) has at least one positive normalized solution $u_a \in H^1(\mathbb{R}^N)$, which is radially symmetric and radially nonincreasing. Moreover, as $a \rightarrow \infty$, there hold*

$$\|\nabla u_a\|_2^2 \sim a^{\frac{2(2N-q(N-2))}{4-N(q-2)}} \rightarrow +\infty,$$

$$E(u_a) \sim \begin{cases} -a^{\frac{2(N+\alpha)}{N}} \rightarrow -\infty, & \text{if } q \in (2, 2 + \frac{4\alpha}{N(2+\alpha)}), \\ -a^{\frac{2(2N-q(N-2))}{4-N(q-2)}} \rightarrow -\infty, & \text{if } q \in (2 + \frac{4\alpha}{N(2+\alpha)}, 2 + \frac{4}{N}). \end{cases}$$

Proposition 6.2. *Assume that (H1) and (H2) hold. If $p = \frac{N+\alpha}{N-2}$, $q \in (2, 2^*)$ for $N \geq 4$ and $q \in (4, 6)$ for $N = 3$, then for any small $a > 0$, the problem (1.6) has at least one positive normalized solutions $u_a \in H^1(\mathbb{R}^N)$, which are radially symmetric and radially nonincreasing. Moreover, as $a \rightarrow 0$, there hold*

$$\|\nabla u_a\|_2^2 \simeq S_\alpha^{\frac{N+\alpha}{2+\alpha}}, \quad E(u_a) \simeq \frac{2+\alpha}{2(N+\alpha)} S_\alpha^{\frac{N+\alpha}{2+\alpha}}.$$

Acknowledgements. S.M. was supported by National Natural Science Foundation of China (Grant Nos.11571187, 11771182)

REFERENCES

- [1] T. Akahori, S. Ibrahim, N. Ikoma, H. Kikuchi and H. Nawa, *Uniqueness and nondegeneracy of ground states to nonlinear scalar field equations involving the Sobolev critical exponent in their nonlinearities for high frequencies*, Calc. Var. Partial Differential Equations, **58** (2019), Paper No. 120, 32 pp.
- [2] T. Akahori, S. Ibrahim, H. Kikuchi and H. Nawa, *Global dynamics above the ground state energy for the combined power type nonlinear Schrödinger equations with energy critical growth at low frequencies*, Memoirs of the AMS, **272** (2021), 1331.
- [3] T. Akahori, S. Ibrahim, H. Kikuchi, H. Nawa, *Non-existence of ground states and gap of variational values for 3D Sobolev critical nonlinear scalar field equations*, J. Differential Equations, **334** (2022), 25–86.
- [4] C. Alves, M. Souto, M. Montenegro, *Existence of a ground state solution for a nonlinear scalar field equation with critical growth*, Calc. Var. Partial Differ. Equ., **43** (2012), 537–554.
- [5] H. Berestycki and P.-L. Lions, *Nonlinear scalar field equations. I. Existence of a ground state*, Archive for Rational Mechanics and Analysis, **82** (1983), 313–345.
- [6] H. Brezis and E. Lieb, *A relation between pointwise convergence of functions and convergence of functionals*, Proc. Amer. Math. Soc., **88** (1983), 486–490.
- [7] T. Cazenave and P.-L. Lions, *Orbital stability of standing waves for some nonlinear Schrödinger equations*, Comm. Math. Phys., **85** (1982), 549–561.
- [8] M. Coles, S. Gustafson, *Solitary Waves and Dynamics for Subcritical Perturbations of Energy Critical NLS*, arXiv:1707.07219 (to appear in Differ. Integr. Equ.).
- [9] D. Gilbarg and N. S. Trudinger, *Elliptic Partial Differential Equations of Second Order*. Grundlehren der Mathematischen Wissenschaften, Second edn. Springer, Berlin (1983).
- [10] L. Jeanjean, *Existence of solutions with prescribed norm for semilinear elliptic equations*, Nonlinear Anal., **28** (1997), 1633–1659.
- [11] L. Jeanjean and T. Le, *Multiple normalized solutions for a Sobolev critical Schrödinger equation*, Mathematische Annalen, <https://doi.org/10.1007/s00208-021-02228-0>.
- [12] L. Jeanjean, J. Zhang and X. Zhong, *A global branch approach to normalized solutions for the Schrödinger equation*, arXiv:2112.05869v1.
- [13] L. Jeanjean, J. Jendrej, T. Le and N. Visciglia, *Orbital stability of ground states for a Sobolev critical schrödinger equation*, arXiv:2008.12084v1.

- [14] M. Lewin and S. R. Nodari, *The double-power nonlinear Schrödinger equation and its generalizations: uniqueness, non-degeneracy and applications*, arXiv:2006.02809v1.
- [15] Xinfu Li, Shiwang Ma and Guang Zhang, *Existence and qualitative properties of solutions for Choquard equations with a local term*, Nonlinear Analysis: Real World Applications, **45** (2019), 1–25.
- [16] Xinfu Li and Shiwang Ma, *Choquard equations with critical nonlinearities*, Commun. Contemp. Math., **22** (2019), 1950023.
- [17] Xinfu Li, *Existence and symmetry of normalized ground state to Choquard equation with local perturbation*, arXiv:2103.07026v1, 2021.
- [18] Xinfu Li, *Standing waves to upper critical Choquard equation with a local perturbation: multiplicity, qualitative properties and stability*, arXiv:2104.09317v1, 19 Apr 2021.
- [19] Xinfu Li, *Nonexistence, existence and symmetry of normalized ground states to Choquard equations with a local perturbation*, arXiv:2103.07026v2 [math.AP] 7 May 2021.
- [20] Xinfu Li, Jianguang Bao and Wenguang Tang, *Normalized solutions to lower critical Choquard equation with a local perturbation*, arXiv:2207.10377v2 [math.AP] 18 Aug 2022.
- [21] X.Q. Liu, J. Q. Liu and Z.-Q. Wang, *Quasilinear elliptic equations with critical growth via perturbation method*, J. Differential Equations, **254** (2013), 102–124.
- [22] Zeng Liu and V. Moroz, *Limit profiles for singularly perturbed Choquard equations with local repulsion*, arXiv:2107.05065v2.
- [23] E.H. Lieb and M. Loss, *Analysis*, volume 14 of graduate studies in mathematics, American Mathematical Society, Providence, RI, (4) 2001.
- [24] J. Liu, J.-F. Liao, C.-L. Tang, *Ground state solution for a class of Schrödinger equations involving general critical growth term*, Nonlinearity, **30** (2017), 899–911.
- [25] P. H. Lions, *The concentration-compactness principle in the calculus of variations: The locally compact cases, Part I and Part II*, Ann. Inst. H. Poincaré Anal. Non Linéaire, **1** (1984), 223–283.
- [26] S. Ma and V. Moroz, *Asymptotic profiles for a nonlinear Schrödinger equation with critical combined powers nonlinearity*, Mathematische Zeitschrift, (2023) 304:13.
- [27] S. Ma and V. Moroz, *Asymptotic profiles for Choquard equations with combined attractive nonlinearities*, arXiv:2302.13727v1 [math.AP] 27 Feb 2023.
- [28] V. Moroz and C. B. Muratov, *Asymptotic properties of ground states of scalar field equations with a vanishing parameter*, J. Eur. Math. Soc., **16** (2014), 1081–1109.
- [29] V. Moroz and J. Van Schaftingen, *Groundstates of nonlinear Choquard equations: Existence, qualitative properties and decay asymptotics*, J. Functional Analysis, **265** (2013), 153–184.
- [30] V. Moroz and J. Van Schaftingen, *A guide to the Choquard equation*, J. Fixed Point Theory Appl. **19** (2017) 773–813.
- [31] N. Soave, *Normalized ground states for the NLS equation with combined nonlinearities: The Sobolev critical case*, J. Functional Analysis, **279** (2020), 108610.
- [32] N. Soave, *Normalized ground state for the NLS equations with combined nonlinearities*, J. Differential Equations, **269** (2020), 6941–6987.
- [33] T. Tao, M. Visan and X. Zhang, *The nonlinear Schrödinger equation with combined power-type nonlinearities*, Commun. Partial Differ. Equ., **32** (2007), 1281–1343.
- [34] K. Tintarev and K.-H. Fieseler, *Concentration Compactness: functional-Analytic Grounds and Applications*, Imperial College Press, 57 Shelton Street, London, 2007.
- [35] J. Van Schaftingen and Jiankang Xia, *Groundstates for a local nonlinear perturbation of the Choquard equations with lower critical exponent*, J. Math. Anal. Appl., **464** (2018), 1184–1202.
- [36] J. Wei and Y. Wu, *Normalized solutions for Schrödinger equations with critical Sobolev exponent and mixed nonlinearities*, J. Functional Analysis, **283** (2022) 109574.
- [37] J. Wei and Y. Wu, *On some nonlinear Schrödinger equations in \mathbb{R}^N* , Proc. Royal Soc. Edinburgh, page 1 of 26, DOI:10.1017/prm.2022.56
- [38] Shuai Yao, Juntao Sun and Tsung-fang Wu, *Normalized solutions for the Schrödinger equation with combined Hartree type and power nonlinearities*, arXiv:2102.10268v1, 2021.
- [39] Shuai Yao, Haibo Chen, V. D. Rădulescu and Juntao Sun, *Normalized solutions for lower critical Choquard equations with critical Sobolev perturbation*, SIAM J. Math. Anal., **54** (2022), 3696–3723.
- [40] J. Zhang, W. Zou, *A Berestycki-Lions theorem revisited*, Commun. Contemp. Math., **14** (2012), 1250033.