

EFFECTIVE VOLUME GROWTH OF THREE-MANIFOLDS WITH POSITIVE SCALAR CURVATURE

YIPENG WANG

ABSTRACT. In this note, we prove an effective linear volume growth for complete three-manifolds with non-negative Ricci curvature and uniformly positive scalar curvature. This recovers the results obtained by Munteanu-Wang [7]. Our method builds upon recent work by Chodosh-Li-Stryker [4], which utilizes the technique of μ -bubbles and the almost-splitting theorem by Cheeger-Colding.

1. INTRODUCTION

The Bishop-Gromov volume comparison theorem asserts that a complete Riemannian manifold M^n with non-negative Ricci curvature exhibits at most Euclidean volume growth: specifically, there exists a universal constant $C(n)$ such that $\text{vol}(B_r(p)) \leq Cr^n$ for any point $p \in M$ and all $r > 0$. A well known conjecture of Gromov [5] proposes that if M^n additionally possesses uniformly positive scalar curvature $R \geq 1$, then there should be a universal constant $C(n)$ such that $\text{vol}(B_r(p)) \leq Cr^{n-2}$.

In this note, we explore Gromov's conjecture within the three-dimensional setting to establish an effective linear volume growth result. It is important to mention that, according to Yau's linear volume growth theorem for manifolds with non-negative Ricci curvature [9], Gromov's conjecture would provide a precise characterization of volume growth for three-manifolds with $\text{Ric}_g \geq 0$ and $R \geq 1$.

Theorem 1.1. *There exists a universal constant C such that the following statement is true. Let (M, g) be a complete Riemannian 3-manifold with $\text{Ric}_g \geq 0$. For all $p \in M$ and $r > 0$, if $R_g \geq 1$ in $B_r(p)$, then*

$$\text{vol}(B_r(p)) \leq Cr$$

This problem has recently been examined by Chodosh, Li and Stryker [4] in the case where M is non-compact. Utilizing the technique of μ -bubbles and Cheeger-Colding theory, they have demonstrated that

$$\text{vol}(B_r(p)) \leq C(p, M, g)r$$

for all $p \in M$ and $r > 0$, where the constant may depend on the manifold. Theorem 1.1 can be considered as an effective version of the main results considered in [4].

An alternative method has been developed by Munteanu-Wang [7], which relies on analysis of level sets of harmonic functions. We refer some other related works for this problem in higher dimensions [8],[10].

Acknowledgment. The author wishes to express sincere gratitude to his advisor, Simon Brendle, for inspiring discussions and continuing support. Additionally, the author is thankful to Chao Li for explaining the ideas presented in [4].

2. MAIN INGREDIENTS OF THE PROOF

Now let us describe the main techniques employed in the proof of Theorem 1.1. The method we use is similar to that in [4]. The Ricci curvature condition enables the application of the Cheeger-Colding almost splitting theorem [1], which ensures that geodesic balls up to a certain scale, centered at the midpoint of a long geodesic, is Gromov-Hausdorff close to a ball with same radius in $N \times \mathbb{R}$, where N as a length space can be constructed as the level set of certain harmonic function that is closed to the distance function. Additionally, the scalar curvature condition allows us to construct a specific surface, Σ (called the μ -bubble) around N , so that each connected component of Σ maintains a uniform diameter bound. We should note that Σ generally possesses numerous connected components, but a more careful analysis of the Cheeger-Colding estimates ensures that N is closed to one specific component of Σ , therefore having a uniform diameter bound.

By considering the universal cover of M , we could assume that M is simply connected. The first key ingredient in our analysis is the geometric estimates of μ -bubbles, which we outline in the following Lemma. The concept of μ -bubble was initially introduced by Gromov, and we refer to [6] for a general introduction to this technique.

Throughout this note, we will write

$$N_R(\Gamma) := \{x : d(x, \Gamma) < R\}$$

to denote the tubular neighborhood of a given closed subset Γ and

$$\beta_{a,b} := \beta([a,b]) \subset M$$

for any unit speed minimizing geodesic β and $0 \leq a < b < \infty$.

Lemma 2.1 (Chodosh-Li [2],[4], Chodosh-Li-Stryker [3]). *There exists constants L and c such that the following is true:*

Let (X^3, g) be a 3-manifold with boundary. If there exists some $p \in X$ such that $d_X(p, \partial X) > L$ and $R_g \geq 1$ in $N_L(\partial X)$, then there exists an open subsets $\Omega \subset N_{\frac{L}{2}}(\partial X) \cap X$ and a smooth surface Σ with $\partial\Omega = \Sigma \sqcup \partial X$ and each connected component of Σ has diameter bounded by c .

Throughout the remainder of this note, we will let L and c to be the universal constants from Lemma 2.1, and without loss of generality, we may assume that $L > 4c$.

Fix a point $p \in M$ and a large constant ℓ , we assume that $R_g \geq 1$ within $B_\ell(p)$. Let $\gamma : [0, \ell] \rightarrow M$ to be a unit speed minimizing geodesic with $\gamma(0) = p$. For each $k \geq 1$ where $(k+1)L < \ell$, we apply Lemma 2.1 to $M \setminus \overline{B_{kL}(p)}$. This yields a smooth surface $\tilde{\Sigma}_k$ that is homologous to $\partial B_{kL}(p)$, with the following properties:

- $\tilde{\Sigma}_k \subset N_{\frac{L}{2}}(\partial B_{kL}(p)) \cap \left(M \setminus \overline{B_{kL}(p)}\right)$.
- The diameter of each connected component of $\tilde{\Sigma}_k$ is bounded by c .

Note that for all positive integers k with $(k+1)L < \ell$, we must have $\gamma \cap \tilde{\Sigma}_k \neq \emptyset$. We set $\Sigma_k \subset \tilde{\Sigma}_k$ as a connected component that intersects with γ . Finally, we let $t_k \in [0, \ell]$ such that $p_k := \gamma(t_k) \in \Sigma_k$.

Lemma 2.2. *We have $d(p_k, p_{k+1}) = |t_k - t_{k+1}| < 2L$.*

Proof. Given that $\Sigma_k \subset N_{\frac{L}{2}}(\partial B_{kL}(p)) \cap \left(M \setminus \overline{B_{kL}(p)}\right)$, it follows that

$$kL \leq t_k < (k+1)L.$$

Therefore $|t_k - t_{k+1}| < 2L$. \square

Lemma 2.3. *For any $0 < s_0 < t_k < s_1 < \ell$, if γ' is any continuous path joining $\gamma(s_0)$ and $\gamma(s_1)$, then $\gamma' \cap \Sigma_k \neq \emptyset$.*

Proof. Suppose not, then γ_{s_0, s_1} and γ' form a loop that has a non-trivial intersection number with Σ_k , contradicting the fact that M is simply connected. \square

2.1. Results from Cheeger-Colding Theory. The second key component of our proof is the almost-splitting theorem, which we outline as follows. Let $S = 5c + 2L$. We let $r_0 \gg S$ as a large constant to be determined later. We denote $\Psi = \Psi(R) : \mathbb{R}^+ \rightarrow \mathbb{R}$ as a continuous function that may vary from line by line, with the property that $\lim_{R \rightarrow \infty} \Psi(R) = 0$.

Let $R \geq r_0$ and $k \in \mathbb{N}^+$ with $R < t_k < \ell - R$. We set up some notations: First we define $B_k := B_S(p_k)$ as the region to apply the almost splitting theorem. Let b to be the Buseman function associate with the geodesic γ

$$b(y) := d(y, \gamma(t_k + R)) - R$$

We then define h as the harmonic replacement of b in a larger ball:

$$\begin{cases} \Delta h = 0, & \text{in } B_{16S}(p_k) \\ h = b, & \text{on } \partial B_{16S}(p_k) \end{cases}$$

By perturbing an arbitrarily small amount along γ , we assume $h(p_k)$ is a regular value of h . We define $\Gamma_k := h^{-1}\{h(p_k)\}$ to be the level set of h at $h(p_k)$. For any $x \in B_k$, we denote x' to be a point in Γ_k that minimizes the distance to x among all points in Γ_k .

Lemma 2.4 (Cheeger-Colding [1]). *Suppose $x, y, z \in B_{2S}(p_k)$, with $h(x) = h(z)$, and z minimizes the distance from y over the level set $h^{-1}\{h(z)\}$, then*

$$\begin{aligned} (1) \quad & |h(x) - b(x)| \leq \Psi \\ (2) \quad & |d(y, z) - |h(y) - h(z)|| \leq \Psi \\ (3) \quad & |d(x, y)^2 - d(x, z)^2 - d(y, z)^2| \leq \Psi \end{aligned}$$

Corollary 2.5. *For all t with $x = \gamma(t) \in B_k$, we have*

$$d(x, x') \leq |t - t_k| + \Psi, \quad d(p_k, x') \leq \Psi$$

Proof. It is clear that for all $x \in B_k$ we have $x' \in B_{2S}(p_k)$. It then follows from (1) that $|h| \leq \Psi$ on Γ_k . With (2) together we obtain

$$\begin{aligned} |d(x, x') - |t - t_k|| &= |d(x, x') - |b(x)|| \\ &\leq |d(x, x') - |h(x)|| + \Psi \\ &\leq |d(x, x') - |h(x) - h(x')|| + \Psi \\ &\leq \Psi \end{aligned}$$

This establishes the first inequality; the second inequality then follows from (3). \square

Lemma 2.6. *There exists some $r_0 = r_0(c, L, S)$ such that for all $R \geq r_0$ the following statement is true. For all k with $R < t_k < \ell - R$, we have*

$$\text{diam}(\Gamma_k \cap B_k) \leq 3c$$

Proof. Suppose, instead, that $\text{diam}(\Gamma_k \cap B_k) > 3c$. Then, there exists some $y \in \Gamma_k \cap B_k$ such that $d(y, B_c(p_k)) > 2c$. In particular, since $\text{diam}(\Sigma_k) \leq c$, we obtain that $d(y, \Sigma_k) > 2c$.

Now we take σ_{\pm} to be the unit speed minimizing geodesic joining $\gamma(t_k \pm \frac{S}{2})$ to y . By Lemma 2.3, we must have at least one of σ_{\pm} intersect with Σ_k . For simplicity, we denote σ as the minimizing geodesic such that

$$\sigma(s_0) = \gamma(\bar{t}) = q, \quad \sigma(s_1) \in \Sigma_k, \quad \sigma(s_2) = y$$

where $s_0 < s_1 < s_2$ and $|\bar{t} - t_k| = \frac{S}{2}$. By Corollary 2.5, we know that

$$d(q, q') \leq \frac{S}{2} + \Psi, \quad d(q', p_k) \leq \Psi$$

Next, we apply (3) to obtain

$$\begin{aligned} (s_2 - s_0)^2 &\leq d(q, q')^2 + d(q', y)^2 + \Psi \\ &\leq \left(\frac{S}{2} + \Psi\right)^2 + \left(d(p_k, y) + d(q', p_k)\right)^2 + \Psi \\ &\leq \left(\frac{S}{2}\right)^2 + d(p_k, y)^2 + \Psi \end{aligned}$$

Hence, by the definition of Ψ , there exists some $r_0 = r_0(c, L, S)$ such that for all $R \geq r_0$, we have

$$(4) \quad (s_2 - s_0)^2 \leq d(p_k, y)^2 + \left(\frac{S}{2}\right)^2 + 2c^2$$

On the other hand, since $d(\sigma(s_1), p_k) \leq c$, we apply the triangle inequality.

$$\begin{aligned} (s_2 - s_0)^2 &= [d(q, \sigma(s_1)) + d(\sigma(s_1), y)]^2 \\ &\geq [d(q, p_k) + d(y, p_k) - 2c]^2 \\ &= d(q, p_k)^2 + d(y, p_k)^2 + 4c^2 \\ &\quad + d(q, p_k)(d(y, p_k) - 2c) + d(y, p_k)(d(q, p_k) - 2c) \end{aligned}$$

Given that $d(y, p_k) \geq 2c$ and $d(q, p_k) = \frac{S}{2} \geq 2c$, this implies

$$\begin{aligned} (s_2 - s_0)^2 &\geq d(q, p_k)^2 + d(y, p_k)^2 + 4c^2 \\ &= \left(\frac{S}{2}\right)^2 + d(y, p_k)^2 + 4c^2 \end{aligned}$$

which contradicts (4). \square

Lemma 2.7. *There exists some $r_0 = r_0(c, L, S)$ such that for all $R \geq r_0$ the following statement is true. For all k with $R < t_k < \ell - R$, we have $B_k \subset N_{4c}(\gamma_{0, \ell})$.*

Proof. Suppose that $d(x, p_k) < S$, and denote $\Gamma_x = h^{-1}\{h(x)\}$ as the level set of h at $h(x)$.

Using the estimates in Lemma 2.4, we obtain:

$$\begin{aligned} &|d(x, p_k)^2 - d(x', p_k)^2 - (h(x) - h(x'))^2| \\ &\leq |d(x, x')^2 - (h(x) - h(x'))^2| + \Psi \\ &= |d(x, x') - |h(x) - h(x')|| \cdot (d(x, x') + |h(x) - h(x')|) + \Psi \\ &\leq \Psi(2d(x, x') + \Psi) + \Psi \\ &\leq \Psi \end{aligned}$$

Let $\hat{p}_k \in \Gamma_x$ that minimizes the distance from p_k among all the points in Γ_x . Then the same argument shows that

$$|d(x, p_k)^2 - d(x, \hat{p}_k)^2 - (h(p_k) - h(\hat{p}_k))^2| \leq \Psi$$

Given that $h(x) = h(\hat{p}_k)$ and $h(p_k) = h(x')$, we deduce:

$$(5) \quad |d(x, \hat{p}_k)^2 - d(x', p_k)^2| \leq \Psi$$

We know that $|h(x)| \leq S + \Psi$ and $[-2S + \Psi, 2S - \Psi] \subset h(\gamma_{t_k - 2S, t_k + 2S})$. Therefore by choosing r_0 large enough, one can make sure that for $R \geq r_0$ we would have

$$|h(x)| < \frac{3}{2}S, \quad \left[\frac{3}{2}S, \frac{3}{2}S\right] \subset h(\gamma_{t_k - 2S, t_k + 2S})$$

Thus we could consider some $\gamma(s) \in \gamma_{t_k-2S, t_k+2S} \cap \Gamma_x$. We find:

$$\begin{aligned} |d(p_k, \hat{p}_k) - d(p_k, \gamma(s))| &= |d(p_k, \hat{p}_k) - |s - t_k|| \\ &\leq |h(\hat{p}_k) - |s - t_k|| + \Psi \\ &= |h(\gamma(s)) - |s - t_k|| + \Psi \\ &\leq \Psi \end{aligned}$$

Using (3), we obtain:

$$d(\hat{p}_k, \gamma(s))^2 \leq d(p_k, \gamma(s))^2 - d(p_k, \hat{p}_k)^2 + \Psi \leq \Psi$$

Applying the triangle inequality and combining with (5), we find:

$$\begin{aligned} d(x, \gamma(s))^2 &\leq (d(x, \hat{p}_k) + d(\hat{p}_k, \gamma(s)))^2 \\ &\leq d(x, \hat{p}_k)^2 + \Psi \\ &\leq d(x', p_k)^2 + \Psi \end{aligned}$$

Finally, from Lemma 2.6, we know $d(x', p_k) \leq 3c$ for sufficiently large r_0 . Replacing r_0 with a larger value if necessary, we conclude that if $R \geq r_0$, then $d(x, \gamma(s))^2 \leq 16c^2$. This guarantees that $x \in N_{4c}(\gamma_{0,\ell})$. \square

Now we fix r_0 to be the constant from Lemma 2.7 and we assume $\ell > 2r_0$.

Proposition 2.8. *We have $N_{5c}(\gamma_{r_0, \ell-r_0}) \subset N_{4c}(\gamma_{0,\ell})$.*

Proof. Consider a point $x \in N_{5c}(\gamma_{r_0, \ell-r_0})$ and suppose

$$d(x, \gamma(\hat{t})) = \inf_{t \in [r_0, \ell-r_0]} \gamma(t)$$

for some $\hat{t} \in [r_0, \ell - r_0]$. Then by assumption, we have $d(x, \gamma(\hat{t})) < 5c$. Furthermore, since $|t_{k+1} - t_k| \leq 2L$ by Lemma 2.2, there exists some $t_k \in [r_0, \ell - r_0]$ such that $|\hat{t} - t_k| < 2L$. Applying the triangle inequality gives

$$d(x, p_k) < d(x, \gamma(\hat{t})) + 2L \leq S$$

Therefore, $x \in B_k$. By Lemma 2.7, for $t_k \in [r_0, \ell - r_0]$, we have $B_k \subset N_{4c}(\gamma_{0,\ell})$. \square

Proposition 2.9. *Given $x \in M$, let $\hat{t} \in [0, \ell]$ such that $d(x, \gamma(\hat{t})) = d(x, \gamma_{0,\ell})$. If $\hat{t} \in [r_0, \ell - r_0]$, then $d(x, \gamma(\hat{t})) \leq 4c$.*

Proof. Suppose the assertion is false and $d(x, \gamma(\hat{t})) > 4c$ with $\hat{t} \in [r_0, \ell - r_0]$. According to Proposition 2.8, we would then have $d(x, \gamma(\hat{t})) > 5c$. By continuity, there exists a point \hat{x} on the geodesic segment joining x to $\gamma(\hat{t})$ where $d(\hat{x}, \gamma(\hat{t})) = 5c$. Applying Proposition 2.8 again, we find $d(\hat{x}, \gamma_{0,\ell}) \leq 4c$. However, since \hat{x} is on the minimizing geodesic between x and $\gamma(\hat{t})$, we must have

$$4c \geq d(\hat{x}, \gamma_{0,\ell}) = d(\hat{x}, \gamma(\hat{t})) = 5c$$

This is a contradiction. \square

Corollary 2.10. *For any s with $\ell > 4s + 2r_0$, one has*

$$N_s(\gamma_{2s+r_0, \ell-2s-r_0}) \subset N_{4c}(\gamma_{0, \ell})$$

Proof. For $x \in N_s(\gamma_{2s+r_0, \ell-2s-r_0})$, let $\hat{t} \in [0, \ell]$ such that $d(x, \gamma(\hat{t})) = d(x, \gamma_{0, \ell})$. We must then have $\hat{t} \in [r_0, \ell - r_0]$. The claim now follows directly from Proposition 2.9. \square

Corollary 2.11. *For $x \notin N_{4c}(\gamma_{0, \ell})$, let $\hat{t} \in [0, \ell]$ such that $d(x, \gamma(\hat{t})) = d(x, \gamma)$. If $d(x, \gamma(t)) > d(x, \gamma)$ for all $t > \ell - r_0$, then $\hat{t} \leq r_0$. Therefore, $\hat{t} \leq r_0$ if either of the following conditions holds:*

- $d(x, \gamma(0)) < \frac{\ell - r_0}{2}$.
- $d(x, \gamma(0)) < d(q, \gamma(\ell)) - r_0$.

Proof. Since $x \notin N_{4c}(\gamma_{0, \ell})$, Proposition 2.9 implies that if $\hat{t} \notin [\ell - r_0, \ell]$, then \hat{t} must be within $[0, r_0]$. Given $t > \ell - r_0$, we consider the following scenarios:

Under the first assumption:

$$d(x, \gamma(t)) \geq \ell - r_0 - d(x, \gamma(0)) > \frac{\ell - r_0}{2} > d(x, \gamma(0))$$

Under the second assumption:

$$d(x, \gamma(t)) \geq d(x, \gamma(\ell)) - r_0 > d(x, \gamma(0))$$

In both cases, this ensures that $\hat{t} \in [0, r_0]$. \square

3. PROOF OF THEOREM 1.1

Now, let us fix $p \in M$ and $r > 0$. We assume $R_g \geq 1$ within $B_r(p)$. We consider a minimizing geodesic $\gamma : [0, r] \rightarrow M$ with $\gamma(0) = p$. Let r_0 be the universal constant from Lemma 2.7, and without loss of generality, we assume that $r_0 > 10S$ and $r > 32r_0$.

We define the region

$$U = N_{4c}(\gamma_{0, r}) \cup B_{6r_0}(p), \quad V = B_{\frac{r}{16}}(p) \setminus U$$

Lemma 3.1. *We have $\text{vol}(U) \leq C(r_0, c)r$.*

Proof. The tubular neighborhood $N_{4c}(\gamma_{0, r})$ can be covered by r/c geodesic balls of radius $4c$. Bishop-Gromov then implies that

$$\text{vol}(U) \leq \text{vol}(N_{4c}(\gamma_{0, r})) + \text{vol}(B_{6r_0}(p)) \leq C\left(\frac{r}{c}\right)c^3 + Cr_0^3 \leq Cr$$

where C depends only on c and r_0 . \square

Let us from now assume that $V \neq \emptyset$. For any $q \in V$, we let $\gamma^q : [0, \ell^q]$ to be the unit speed minimizing geodesic joining q and $\gamma(\frac{r}{4})$. Note that it follows from the triangle inequality that $\frac{3}{16}r \leq \ell^q \leq \frac{5}{16}r$.

Proposition 3.2. *For any $q \in V$, there exists some $t^q \leq \frac{r}{8}$ such that*

$$d(p, \gamma^q) = d(p, \gamma^q(t^q)) \leq 4c.$$

Proof. First notice that $d(p, q) < \frac{r-r_0}{2}$ and $q \notin V$, hence Corollary 2.11 implies that

$$d(\gamma, \gamma^q(0)) \geq d(p, q) - r_0 > 5c$$

Since $d(\gamma, \gamma^q(\ell^q)) = 0$, by continuity there exists some $\tilde{t} \in [0, \ell^q]$ such that $d(\gamma, \gamma^q(\tilde{t})) = 5c$. It is clear that

$$\min\{d(\gamma^q(\tilde{t}), \gamma^q(0)), d(\gamma^q(\tilde{t}), \gamma^q(\ell^q))\} \leq \frac{\ell^q}{2}$$

thus,

$$d(p, \gamma^q(\tilde{t})) \leq \frac{\ell^q}{2} + \max\{d(p, \gamma^q(0)), d(p, \gamma^q(\ell^q))\} \leq \frac{13}{32}r < \frac{r-r_0}{2}$$

Applying Corollary 2.11 again, we find that $\gamma^q(\tilde{t}) \in N_{5c}(\gamma_0, r_0)$. We therefore conclude that $d(p, \gamma^q(\tilde{t})) < \frac{3}{2}r_0$. This implies for all $t \in [0, \ell^q]$, we have

$$d(p, \gamma^q(t)) \leq d(p, \gamma^q(\tilde{t})) + |t - \tilde{t}| < \frac{3}{2}r_0 + |t - \tilde{t}|$$

But by assumption $d(p, \gamma^q(0)), d(p, \gamma^q(\ell^q)) \geq 6r_0$, and hence $\tilde{t} \in [\frac{9}{2}r_0, \ell^q - \frac{9}{2}r_0]$ and $p \in N_{\frac{3}{2}r_0}(\gamma_{4r_0, \ell^q-4r_0}^q)$. Corollary 2.10 then implies that

$$d(p, \gamma^q(t^q)) := \min_{t \in [0, \ell^q]} d(p, \gamma^q(t)) \leq 4c.$$

To control t^q , we observe that

$$\ell^q - t^q \geq d(p, \gamma^q(\ell^q)) - d(p, \gamma^q(t^q)) = \frac{r}{4} - d(p, \gamma^q(t^q))$$

But given $\ell^q \leq \frac{5}{16}r$ and $d(p, \gamma^q(t^q)) \leq 4c$, it follows that $t^q \leq \frac{r}{8}$. \square

Let us now pick some $q_0 \in V$ such that $d(q, \gamma(\frac{r}{4})) \leq d(q_0, \gamma(\frac{r}{4})) + r_0$ for all $q \in V$.

Proposition 3.3. *We have $V \subset N_{4c}(\gamma_{0, \ell^{q_0}}^{q_0}) \cup B_{6r_0}(q_0)$.*

Proof. Suppose the statement is false and there exists some $q \in V$ such that $q \notin N_{4c}(\gamma_{0, \ell^{q_0}}^{q_0}) \cup B_{6r_0}(q_0)$. Let $\hat{t} \in [0, \ell^{q_0}]$ so that $d(q, \gamma^{q_0}(\hat{t})) = d(q, \gamma^{q_0})$. We first observe that

$$d(q_0, q) < 2 \cdot \frac{r}{16} < \ell^q - r_0 = d(q, \gamma^{q_0}(\ell^{q_0})) - r_0$$

Then, with $q \notin N_{4c}(\gamma_{0, \ell^{q_0}}^{q_0})$, Corollary 2.11 would imply that $\hat{t} \in [0, r_0]$. This implies:

$$d(q, \gamma^{q_0}) \geq d(q, q_0) - d(q_0, \gamma^{q_0}(\hat{t})) \geq d(q_0, q) - r_0$$

As we are assuming $q \notin B_{6r_0}(q_0)$, it follows that $d(q, \gamma^{q_0}) > 5r_0$. By continuity, along γ^q there exists some $t^* \in [0, \ell^q]$ such that

$$t^* := \inf\{t \in [0, \ell^q] : d(\gamma^q(t), \gamma^{q_0}) = r_0\}$$

By Proposition 3.2, there exists some $t^{q_0}, t^q \in [0, \frac{r}{8}]$ satisfying

$$d(p, \gamma^{q_0}(t^{q_0})) \leq 4c, \quad d(p, \gamma^q(t^q)) \leq 4c$$

This implies $d(\gamma^{q_0}(t^{q_0}), \gamma^q(t^q)) \leq 8c < r_0$, thus the minimality assumption of t^* gives $t^* \leq \frac{r}{8}$. However, this shows that $\ell^q - t^* > \frac{r}{16} > 2r_0$. Hence for all $t > \ell^{q_0} - r_0$

$$d(\gamma^q(t^*), \gamma^{q_0}(t)) \geq d(\gamma^q(t^*), \gamma^q(\ell^q)) - r_0 > r_0$$

and we must have $\gamma^q(t^*) \in N_{r_0}(\gamma_{0,r_0}^{q_0})$ by Corollary 2.11. In particular, this implies $d(q_0, \gamma^q(t^*)) \leq 2r_0$. Using the triangle inequality

$$\begin{aligned} d\left(q, \gamma\left(\frac{r}{4}\right)\right) &\geq d(q, q_0) + d\left(q_0, \gamma\left(\frac{r}{4}\right)\right) - 2d(q_0, \gamma^q(t^*)) \\ &\geq d(q, q_0) + d\left(q_0, \gamma\left(\frac{r}{4}\right)\right) - 4r_0 \end{aligned}$$

But the assumption of q_0 implies $d(q_0, \gamma(\frac{r}{4})) \geq d(q, \gamma(\frac{r}{4})) - r_0$ and we obtain that $d(q, q_0) \leq 5r_0$. This contradicts with the assumption that $q \notin B_{6r_0}(q_0)$. \square

Corollary 3.4. *We have $\text{vol}(V) \leq Cr$, and hence $\text{vol}(B_{\frac{r}{16}}(p)) < Cr$.*

Proof. This follows from the same argument as the proof of Lemma 3.1. \square

Now, we will prove the main result of this note.

Proof of Theorem 1.1. Given $p \in M$ and $r > 0$, we suppose that $r > 64r_0$. Otherwise, we have $\text{vol}(B_r(p)) \leq Cr_0^3 \leq Cr_0$. We can also assume that there exists a unit speed minimizing geodesic $\gamma : [0, r] \rightarrow M$ with $\gamma(0) = p$. It then follows from Corollary 3.4 and the Bishop-Gromov volume comparison theorem again

$$\text{vol}(B_r(p)) \leq C \text{vol}(B_{\frac{r}{16}}(p)) \leq Cr$$

This proves Theorem 1.1. \square

REFERENCES

- [1] Jeff Cheeger and Tobias H. Colding, *Lower bounds on Ricci curvature and the almost rigidity of warped products*, Ann. of Math. (2) **144** (1996), no. 1, 189–237. MR 1405949
- [2] Otis Chodosh and Chao Li, *Generalized soap bubbles and the topology of manifolds with positive scalar curvature*, Ann. of Math. (2) **199** (2024), no. 2, 707–740. MR 4713021
- [3] Otis Chodosh, Chao Li, and Douglas Stryker, *Complete stable minimal hypersurfaces in positively curved 4-manifolds*, to appear in J. Eur. Math. Soc. (2022).
- [4] Otis Chodosh, Chao Li, and Douglas Stryker, *Volume growth of 3-manifolds with scalar curvature lower bounds*, Proc. Amer. Math. Soc. **151** (2023), no. 10, 4501–4511. MR 4643334
- [5] M. Gromov, *Large Riemannian manifolds*, Curvature and topology of Riemannian manifolds (Katata, 1985), Lecture Notes in Math., vol. 1201, Springer, Berlin, 1986, pp. 108–121. MR 859578
- [6] Misha Gromov, *Four lectures on scalar curvature*, Perspectives in scalar curvature. Vol. 1, World Sci. Publ., Hackensack, NJ, [2023] ©2023, pp. 1–514. MR 4577903
- [7] Ovidiu Munteanu and Jiaping Wang, *Geometry of three-dimensional manifolds with scalar curvature lower bound*, 2022. arXiv:2201.05595

- [8] Jinmin Wang, Zhizhang Xie, Bo Zhu, and Xingyu Zhu, *Positive scalar curvature meets ricci limit spaces*, 2023. arXiv:2212.10416
- [9] Shing Tung Yau, *Some function-theoretic properties of complete Riemannian manifold and their applications to geometry*, Indiana Univ. Math. J. **25** (1976), no. 7, 659–670. MR 417452
- [10] Bo Zhu, *Geometry of positive scalar curvature on complete manifold*, Journal für die reine und angewandte Mathematik (Crelles Journal) **2022** (2022), no. 791, 225–246.

COLUMBIA UNIVERSITY, 2990 BROADWAY, NEW YORK NY 10027, USA