

On the enumeration of permutations avoiding chains of patterns

Robin D.P. Zhou*, Yongchun Zang

College of Mathematics Physics and Information
Shaoxing University
Shaoxing 312000, P.R. China

Abstract. In 2019, Bóna and Smith introduced the notion of strong pattern avoidance, saying that a permutation π strongly avoids a pattern σ if π and π^2 both avoid σ . Recently, Archer and Geary generalized the idea of strong pattern avoidance to chain avoidance, in which a permutation π avoids a chain of patterns $(\tau^{(1)} : \tau^{(2)} : \dots : \tau^{(k)})$ if the i -th power of the permutation avoids the pattern $\tau^{(i)}$ for $1 \leq i \leq k$. In this paper, we give explicit formulae for the number of sets of permutations avoiding certain chains of patterns. Our results give affirmative answers to two conjectures proposed by Archer and Geary.

Keywords: pattern avoidance, strong pattern avoidance, chain avoidance.

AMS Subject Classifications: 05A05, 05C30

1 Introduction

Let $\mathcal{S}_n$ denote the symmetry group consisting of all permutations of $[n] = \{1, 2, \dots, n\}$, which we can view as bijections from $[n]$ to $[n]$. A permutation π can also be viewed as a word by associating π with the word $\pi(1)\pi(2)\cdots\pi(n)$. Throughout this paper, we always write a permutation $\pi \in \mathcal{S}_n$ as a word $\pi = \pi_1\pi_2\cdots\pi_n$, where $\pi_i = \pi(i)$ for $1 \leq i \leq n$.

Given a permutation $\pi \in \mathcal{S}_n$ and a permutation $\sigma \in \mathcal{S}_k$, an *occurrence* of σ in π is a subsequence $\pi_{i_1}\pi_{i_2}\cdots\pi_{i_k}$ of π that is order isomorphic to σ . We say π *contains* the pattern σ if π contains an occurrence of σ . Otherwise, we say π *avoids* the pattern σ or π is σ -*avoiding*. For example, the permutation 1534627 avoids the pattern 3142 while it contains the pattern 12345 corresponding to the subsequence 13467.

The study of pattern avoidance can be traced back to the work of MacMahon [8] and has become a research focus in enumerative combinatorics over the past half century. A recent survey on permutation patterns can be found in the books [2] and

*Corresponding author.

E-mail address: dapao2012@163.com.

[6]. The notion of pattern avoidance has arisen many further variations. Bóna and Smith [3] initiated the study of *strong pattern avoidance*, in which a permutation π strongly *avoids* a pattern σ if π and π^2 both avoid σ . The study of strong pattern avoidance has attracted the interest of combinatorial scholars such as [4, 9].

Recently, Archer and Geary [1] generalized the idea of strong pattern avoidance to *chain avoidance*, in which a permutation π *avoids* a chain (of patterns) $(\tau^{(1)} : \tau^{(2)} : \dots : \tau^{(k)})$ if the i -th power of the permutation (i.e., π^i) avoids the pattern $\tau^{(i)}$ for $1 \leq i \leq k$. Let $\mathcal{S}_n(\tau_1 : \tau_2 : \dots : \tau_k)$ denote the set of $(\tau_1 : \tau_2 : \dots : \tau_k)$ -avoiding permutations in $\mathcal{S}_n$. This definition can be extended to include sets of patterns, which we will separate the patterns in a set with commas. For example, if π avoids the chain $(\sigma, \rho : \tau)$, then π avoids both σ and ρ and π^2 avoids τ . For example, let $\pi = 1325467$. Then we have $\pi \in \mathcal{S}_7(231, 1432 : 231)$ because π avoids both 231 and 1432, $\pi^2 = 1234567$ avoids 231.

It would also be natural to consider *consecutive patterns* as part of the chain. We say π *contains* a consecutive pattern $\bar{\sigma} = \overline{\sigma_1 \sigma_2 \dots \sigma_k}$ if there is an occurrence of σ corresponding to a consecutive subsequence $\pi_i \pi_{i+1} \dots \pi_{i+k-1}$ ($1 \leq i \leq n - k + 1$) of π . Otherwise, we say π *avoids* the (consecutive) pattern $\bar{\sigma}$ or π is $\bar{\sigma}$ -*avoiding*. For example, the permutation $\pi = 1534627$ avoids the pattern $\overline{321}$ while π contains the pattern $\overline{213}$ corresponding to the consecutive subsequence 627.

Archer and Geary [1] obtained the explicit formulae for the number of sets $\mathcal{S}_n(213, 312 : \tau)$ for $\tau \in \mathcal{S}_3$ and further posed the following conjectures.

Conjecture 1.1 *For $n \geq 1$, we have $|\mathcal{S}_n(231, 1432 : 231)| = L_{n+1} - \lfloor \frac{n}{2} \rfloor - 1$, where L_{n+1} is the $(n+1)$ -th Lucas number (A000032, [7]).*

Conjecture 1.2 *For $n \geq 2$, we have $|\mathcal{S}_n(213, 312 : \overline{213})| = 2^{n-2} + n - 1$.*

The objective of this paper is to prove Conjecture 1.1 and Conjecture 1.2. Our proofs are based on the recursive decompositions of the permutations in $\mathcal{S}_n(231, 1432 : 231)$ and the permutations in $\mathcal{S}_n(213, 312 : \overline{213})$, respectively.

2 Avoiding the chain $(231, 1432 : 231)$

This section is devoted to the proof of Conjecture 1.1. This will be accomplished by showing that the both sides of the equation in Conjecture 1.1 satisfy the same recurrence and initial conditions. The following lemma is crucial for our proof.

Lemma 2.1 *Let $n \geq 3$ and $\pi = \pi_1\pi_2\cdots\pi_n \in \mathcal{S}_n(231, 1432 : 231)$. Then we have either $\pi_1 = n$, $\pi_n = n$ or $\pi_{n-1}\pi_n = n(n-1)$.*

Proof. The result is trivial for $n = 3$. Now we let $n \geq 4$. Assume that $\pi_1 \neq n$, $\pi_n \neq n$ and $\pi_{n-1}\pi_n \neq n(n-1)$. Let $\pi_k = n$. Then $2 \leq k \leq n-1$. Combining the fact π is 231-avoiding, we have that π can be written in the form $\pi = \sigma n \tau$, where σ and τ are not empty and all the elements in σ are less than the elements in τ . Notice that $\pi_{n-1}\pi_n \neq n(n-1)$. It is easily seen that τ contains at least two elements, namely, $k \leq n-2$. We claim that the elements in τ are increasing. If not, there exists some $i > k$ such that $\pi_i > \pi_{i+1}$. It follows that the subsequence $\pi_1 n \pi_i \pi_{i+1}$ forms an occurrence of the pattern 1432 in π , contradicting to the fact that π is 1432-avoiding. This proves the claim. Hence π is of the form $\pi = \sigma n k(k+1)\cdots(n-1)$. Now consider π^2 . It is easily checked that $\pi^2(k) = \pi(\pi(k)) = \pi(n) = n-1$, $\pi^2(k+1) = \pi(\pi(k+1)) = \pi(k) = n$ and $\pi^2(n) = \pi(\pi(n)) = \pi(n-1) = n-2$. Then $\pi^2(k)\pi^2(k+1)\pi^2(n) = (n-1)n(n-2)$ forms an occurrence of the pattern 231 in π^2 , contradicting to the fact that π^2 is 231-avoiding. This completes the proof. $\blacksquare$

Given $n \geq 3$, Lemma 2.1 enables us to divide the set $\mathcal{S}_n(231, 1432 : 231)$ into the following three disjoint subsets:

$$\begin{aligned}\mathcal{P}_n^1 &:= \{\pi \mid \pi = \pi_1\pi_2\cdots\pi_n \in \mathcal{S}_n(231, 1432 : 231) \text{ and } \pi_1 = n\}, \\ \mathcal{P}_n^2 &:= \{\pi \mid \pi = \pi_1\pi_2\cdots\pi_n \in \mathcal{S}_n(231, 1432 : 231) \text{ and } \pi_n = n\}, \\ \mathcal{P}_n^3 &:= \{\pi \mid \pi = \pi_1\pi_2\cdots\pi_n \in \mathcal{S}_n(231, 1432 : 231) \text{ and } \pi_{n-1}\pi_n = n(n-1)\}.\end{aligned}$$

We shall characterize the permutations in $\mathcal{P}_n^1$, $\mathcal{P}_n^2$ and $\mathcal{P}_n^3$, respectively. The following theorem is needed for the characterization of $\mathcal{P}_n^1$.

Theorem 2.2 ([3], Theorem 3.1) *For any permutation π ending in 1, the following two statements are equivalent.*

- (A) *The permutation π is strongly 312-avoiding.*
- (B) *The permutation π has form $\pi = (k+1)(k+2)\cdots nk(k-1)(k-2)\cdots 1$ where $\lceil \frac{n}{2} \rceil \leq k \leq n-1$.*

Lemma 2.3 *Let $n \geq 3$ and $\pi = \pi_1\pi_2\cdots\pi_n$. Then we have*

$$\mathcal{P}_n^1 = \{\pi \mid \pi = n(n-1)\cdots(n-k+1)12\cdots(n-k), \lceil \frac{n}{2} \rceil \leq k \leq n-1\}.$$

Proof. Let $\mathcal{A}_n := \{\pi \mid \pi = \pi_1\pi_2\cdots\pi_n \in \mathcal{S}_n(312 : 312), \pi_n = 1\}$ and $\mathcal{B}_n := \{\pi \mid \pi = \pi_1\pi_2\cdots\pi_n \in \mathcal{S}_n(231 : 231), \pi_1 = n\}$. By the definition of $\mathcal{P}_n^1$, we have $\mathcal{P}_n^1 \subseteq \mathcal{B}_n$. As

the pattern 231 is the inverse of 312, it is straightforward to see that π is strongly 231-avoiding if and only if π^{-1} is strongly 312-avoiding. Note that a permutation π begins with n if and only if π^{-1} ends in 1. Therefore, the inverse map is a bijection between $\mathcal{A}_n$ and $\mathcal{B}_n$. From Theorem 2.2, we obtain that

$$\mathcal{A}_n = \{\pi \mid \pi = (k+1)(k+2)\cdots nk(k-1)(k-2)\cdots 1, \lceil \frac{n}{2} \rceil \leq k \leq n-1\}.$$

It can be checked that the permutation $(k+1)(k+2)\cdots nk(k-1)(k-2)\cdots 1$ is the inverse of $n(n-1)\cdots(n-k+1)12\cdots(n-k)$. It yields that

$$\mathcal{B}_n = \{\pi \mid \pi = n(n-1)\cdots(n-k+1)12\cdots(n-k), \lceil \frac{n}{2} \rceil \leq k \leq n-1\}.$$

Observe that all the permutations of the form $n(n-1)\cdots(n-k+1)12\cdots(n-k)$ are 1432-avoiding. We derive that $\mathcal{B}_n \subseteq \mathcal{P}_n^1$. Hence we have $\mathcal{P}_n^1 = \mathcal{B}_n$. This completes the proof. $\blacksquare$

Given $\sigma \in \mathcal{S}_k$ and $\tau \in \mathcal{S}_m$, let $\sigma \oplus \tau$ denote the *direct sum* of σ and τ defined by

$$(\sigma \oplus \tau)(i) = \begin{cases} \sigma(i), & \text{if } 1 \leq i \leq k, \\ \tau(i-k) + k, & \text{if } k+1 \leq i \leq k+m. \end{cases}$$

For example, $231 \oplus 21 = 23154$. The following fact is straightforward from the definition of direct sum.

$$(\sigma \oplus \tau)^2 = \sigma^2 \oplus \tau^2. \quad (2.1)$$

Lemma 2.4 *Let $n \geq 3$ and $\pi = \pi_1\pi_2\cdots\pi_n \in \mathcal{S}_n$. Then $\pi \in \mathcal{P}_n^2$ if and only if $\pi = \sigma \oplus 1$ for some $\sigma \in \mathcal{S}_{n-1}(231, 1432 : 231)$.*

Proof. If $\pi \in \mathcal{P}_n^2$, then we have $\pi_n = n$. By the definition of the direct sum, we have that $\pi = \sigma \oplus 1$ for some $\sigma \in \mathcal{S}_{n-1}$. From (2.1) we have $\pi^2 = \sigma^2 \oplus 1$. This means that σ (resp. σ^2) is a subsequence of π (resp. π^2). Since π avoids the chain $(231, 1432 : 231)$, we deduce that σ also avoids the chain $(231, 1432 : 231)$, that is, $\sigma \in \mathcal{S}_{n-1}(231, 1432 : 231)$.

Conversely, if $\pi = \sigma \oplus 1$ for some $\sigma \in \mathcal{S}_{n-1}(231, 1432 : 231)$, we have $\pi_n = n$ and $\pi^2 = \sigma^2 \oplus 1$. To prove $\pi \in \mathcal{P}_n^2$, it suffices to show that π avoids the chain $(231, 1432 : 231)$. Observe that the element $\pi_n = n$ can not appear in any occurrences of the pattern 231 (resp. 1432) in π . Combining the fact that σ avoids the chain $(231, 1432 : 231)$, we deduce that π avoids both 231 and 1432. By a similar argument as above, we have π^2 is 231-avoiding. In conclusion, we have π avoids the chain $(231, 1432 : 231)$, as desired. $\blacksquare$

Lemma 2.5 *Let $n \geq 3$ and $\pi = \pi_1\pi_2\cdots\pi_n \in \mathcal{S}_n$. Then $\pi \in \mathcal{P}_n^3$ if and only if $\pi = \sigma \oplus 21$ for some $\sigma \in \mathcal{S}_{n-2}(231, 1432 : 231)$.*

Proof. If $\pi \in \mathcal{P}_n^3$, then we have $\pi_{n-1}\pi_n = n(n-1)$. By the definition of the direct sum, we have that $\pi = \sigma \oplus 21$ for some $\sigma \in \mathcal{S}_{n-2}$. Then we obtain $\pi^2 = \sigma^2 \oplus 12$ from (2.1). This means that σ (resp. σ^2) is a subsequence of π (resp. π^2). Since π avoids the chain $(231, 1432 : 231)$, we deduce that σ also avoids the chain $(231, 1432 : 231)$, that is, $\sigma \in \mathcal{S}_{n-2}(231, 1432 : 231)$.

Conversely, assume that $\pi = \sigma \oplus 21$ for some $\sigma \in \mathcal{S}_{n-2}(231, 1432 : 231)$. By the definition of direct sum and (2.1), we have $\pi_{n-1}\pi_n = n(n-1)$ and $\pi^2 = \sigma^2 \oplus 12$. To prove $\pi \in \mathcal{P}_n^3$, it suffices to show that π avoids the chain $(231, 1432 : 231)$. As $\pi_{n-1}\pi_n = n(n-1)$, we deduce that neither n nor $n-1$ can appear in any occurrences of the pattern 231 (resp. 1432) in π . Combining the fact that σ avoids the chain $(231, 1432 : 231)$, we deduce that π avoids both 231 and 1432. By a similar argument as above, we have π^2 is 231-avoiding. Therefore π avoids the chain $(231, 1432 : 231)$, as desired. ■

Proof of Conjecture 1.1. Let $n \geq 1$ and $f(n) = |\mathcal{S}_n(231, 1432 : 231)|$. Then we have

$$f(n) = |\mathcal{P}_n^1| + |\mathcal{P}_n^2| + |\mathcal{P}_n^3|$$

for $n \geq 3$. Lemmas 2.3, 2.4 and 2.5 tell us that $|\mathcal{P}_n^1| = n - \lceil \frac{n}{2} \rceil = \lceil \frac{n-1}{2} \rceil$, $|\mathcal{P}_n^2| = f(n-1)$ and $|\mathcal{P}_n^3| = f(n-2)$, respectively. Therefore we have

$$f(n) = \lceil \frac{n-1}{2} \rceil + f(n-1) + f(n-2)$$

for $n \geq 3$ with the initial conditions $f(1) = 1$ and $f(2) = 2$. Notice that the Lucas numbers L_n satisfy that $L_n = L_{n-1} + L_{n-2}$ for $n \geq 3$ with the initial conditions $L_1 = 1$ and $L_2 = 3$. It is routine to check that $f(n)$ and $L_{n+1} - \lceil \frac{n}{2} \rceil - 1$ satisfy the same recurrence and initial conditions. Thus $f(n) = L_{n+1} - \lceil \frac{n}{2} \rceil - 1$ for $n \geq 1$. This completes the proof. ■

3 Avoiding the chain $(213, 312 : \overline{213})$

This section is devoted to the proof of Conjecture 1.2. To this end, we need to give a characterization of the permutations in $\mathcal{S}_n(213, 312 : \overline{213})$. Note that the permutations in $\mathcal{S}_n(213, 312)$ are exactly the set of unimodal permutations, i.e., those permutations $\pi = \pi_1\pi_2\cdots\pi_n$ with $\pi_1 < \pi_2 < \cdots < \pi_k > \pi_{k+1} > \cdots > \pi_n$ for some $1 \leq k \leq n$ [5]. For a unimodal permutation $\pi = \pi_1\pi_2\cdots\pi_n$, we have either $\pi_1 = 1$ or

$\pi_n = 1$. Hence, for $n \geq 2$, we can divide the set $\mathcal{S}_n(213, 312 : \overline{213})$ into the following two disjoint subsets:

$$\begin{aligned}\mathcal{Q}_n^1 &:= \{\pi \mid \pi = \pi_1\pi_2\cdots\pi_n \in \mathcal{S}_n(213, 312 : \overline{213}) \text{ and } \pi_1 = 1\}, \\ \mathcal{Q}_n^2 &:= \{\pi \mid \pi = \pi_1\pi_2\cdots\pi_n \in \mathcal{S}_n(213, 312 : \overline{213}) \text{ and } \pi_n = 1\}.\end{aligned}$$

We will give characterizations of the unimodal permutations in $\mathcal{Q}_n^1$ and $\mathcal{Q}_n^2$, respectively.

Lemma 3.1 *Let $n \geq 3$ and $\pi = \pi_1\pi_2\cdots\pi_n \in \mathcal{S}_n$. Then we have $\pi \in \mathcal{Q}_n^1$ if and only if $\pi = 1 \oplus \sigma$ for some $\sigma \in \mathcal{S}_{n-1}(213, 312 : \overline{213})$.*

Proof. If $\pi \in \mathcal{Q}_n^1$, then we have $\pi_1 = 1$. By the definition of direct sum, the permutation π can be written as $\pi = 1 \oplus \sigma$ for some $\sigma \in \mathcal{S}_{n-1}$. From (2.1), we have $\pi^2 = 1 \oplus \sigma^2$. Hence σ (resp. σ^2) is order isomorphic to $\pi_2\pi_3\cdots\pi_n$ (resp. $\pi^2(2)\pi^2(3)\cdots\pi^2(n)$). Then $\sigma \in \mathcal{S}_{n-1}(213, 312 : \overline{213})$ follows directly from the fact that π avoids the chain $(213, 312 : \overline{213})$.

Conversely, if $\pi = 1 \oplus \sigma$ for some $\sigma \in \mathcal{S}_{n-1}(213, 312 : \overline{213})$. Then we have $\pi_1 = 1$ and $\pi^2 = 1 \oplus \sigma^2$. To prove $\pi \in \mathcal{Q}_n^1$, it suffices to show that π avoids the chain $(213, 312 : \overline{213})$. Observe that the element $\pi_1 = 1$ can not appear in any occurrences of the pattern 213 (resp. 312) in π . Combining the fact that σ avoids the chain $(213, 312 : \overline{213})$, we deduce that π avoids both 213 and 312. By a similar argument as above, we have π^2 is $\overline{213}$ -avoiding. Therefore the permutation π avoids the chain $(213, 312 : \overline{213})$, as desired. ■

The following lemma is needed for the characterization of permutations in $\mathcal{Q}_n^2$.

Lemma 3.2 *Let $n \geq 3$ and $\pi = \pi_1\pi_2\cdots\pi_n \in \mathcal{S}_n(213, 312)$. If $\pi^2(i-1)\pi^2(i)\pi^2(i+1)$ is an occurrence of the consecutive pattern $\overline{213}$ in π^2 , then we have $\pi_i = n$.*

Proof. We have known that the permutations in $\mathcal{S}_n(213, 312)$ are unimodal permutations. If $\pi_i \neq n$, then we have either $\pi_{i-1} < \pi_i < \pi_{i+1}$ or $\pi_{i-1} > \pi_i > \pi_{i+1}$. In the former case, we have $\pi(\pi_{i-1})\pi(\pi_i)\pi(\pi_{i+1}) = \pi^2(i-1)\pi^2(i)\pi^2(i+1)$ is a subsequence of π . Since $\pi^2(i-1)\pi^2(i)\pi^2(i+1)$ is an occurrence of the consecutive pattern $\overline{213}$ in π^2 , we obtain that π contains the pattern 213, contradicting to the fact that π is 213-avoiding. In the later case, we have $\pi(\pi_{i+1})\pi(\pi_i)\pi(\pi_{i-1}) = \pi^2(i+1)\pi^2(i)\pi^2(i-1)$ is a subsequence of π . Since $\pi^2(i-1)\pi^2(i)\pi^2(i+1)$ is an occurrence of the consecutive pattern $\overline{213}$ in π^2 , we have $\pi^2(i+1)\pi^2(i)\pi^2(i-1)$ is an occurrence of the pattern 312 in π , contradicting to the fact that π is 312-avoiding. This completes the proof. ■

Lemma 3.3 For $n \geq 3$, we have

$$\mathcal{Q}_n^2 = \{\pi = \pi_1\pi_2\cdots\pi_n \mid \pi = 23\cdots n1 \text{ or } \pi = \sigma n(n-1)\tau 1\},$$

where σ (possibly empty) is increasing and τ (possibly empty) is decreasing.

Proof. Let $\pi = \pi_1\pi_2\cdots\pi_n$ be a unimodal permutation with $\pi_k = n$ and $\pi_n = 1$. Then we have either $\pi_{k-1} = n-1$ or $\pi_{k+1} = n-1$.

Case (i). $\pi_{k-1} = n-1$.

First we consider the permutation $\pi = 23\cdots n1$. We have $\pi^2 = 34\cdots n12$. By the definition of $\mathcal{Q}_n^2$, one can easily check that $\pi \in \mathcal{Q}_n^2$. Now assume that $\pi \neq 23\cdots n1$. It yields that there exists at least one element between n and 1 in π . We claim that $\pi \notin \mathcal{Q}_n^2$. We will prove the claim by showing that $\pi^2(k-1)\pi^2(k)\pi^2(k+1)$ is an occurrence of the consecutive pattern $\overline{213}$ in π^2 . Since π is unimodal with $\pi_k = n$ and $\pi_n = 1$, we have $\pi_i > i$ for $i \in [1, k]$ and $\pi_{k+1} \geq \pi_j \geq \pi_{n-1}$ for $j \in [k+1, n-1]$. It is routine to check that $\pi^2(k) = \pi(\pi_k) = \pi_n = 1$ and $\pi^2(k-1) = \pi(\pi_{k-1}) = \pi_{n-1}$. If $\pi_{k+1} \leq k$, we have $\pi^2(k+1) = \pi(\pi_{k+1}) > \pi_{k+1}$. As $\pi_{k+1} \geq \pi_{n-1} = \pi^2(k-1)$, we have $\pi^2(k+1) > \pi^2(k-1)$. It follows that $\pi^2(k-1)\pi^2(k)\pi^2(k+1)$ is an occurrence of the pattern $\overline{213}$ in π^2 , as desired. Now suppose that $k < \pi_{k+1} < n-1$. We have $\pi^2(k-1) = \pi(n-1) < \pi(\pi_{k+1}) = \pi^2(k+1)$. Again $\pi_{k-1}^2\pi_k^2\pi_{k+1}^2$ is an occurrence of the pattern $\overline{213}$ in π^2 , as desired. Hence, the claim is verified. In conclusion, there exists only one permutation $\pi = 23\cdots n1 \in \mathcal{Q}_n^2$ if $\pi_{k-1} = n-1$.

Case (ii). $\pi_{k+1} = n-1$.

Recall that π is unimodal. In this case, the permutation π can be written in the form $\pi = \sigma n(n-1)\tau 1$ where σ (possibly empty) is increasing and τ (possibly empty) is decreasing. We need to show that all such permutations are contained in the set $\mathcal{Q}_n^2$. If σ is empty, it is straightforward to check that $\pi = n(n-1)\cdots 1 \in \mathcal{Q}_n^2$, as desired. Now assume that σ is nonempty. To prove $\pi \in \mathcal{Q}_n^2$, we need to show that π^2 avoids the consecutive pattern $\overline{213}$. From Lemma 3.2, it is sufficient to show that $\pi^2(k-1)\pi^2(k)\pi^2(k+1)$ is not an occurrence of $\overline{213}$ in π^2 . Observe that $k \leq \pi_{k-1} < n-1$. Since π is unimodal, we have $\pi^2(k-1) = \pi(\pi_{k-1}) > \pi(n-1) = \pi^2(k+1)$. Thus, $\pi^2(k-1)\pi^2(k)\pi^2(k+1)$ is not an occurrence of $\overline{213}$ in π^2 , as desired. ■

Proof of Conjecture 1.2. Let $n \geq 2$ and $g(n) = |\mathcal{S}_n(213, 312 : \overline{213})|$. Then we have

$$g(n) = |\mathcal{Q}_n^1| + |\mathcal{Q}_n^2|.$$

By Lemma 3.1, we have $|\mathcal{Q}_n^1| = g(n-1)$ for $n \geq 3$. Lemma 3.3 tells us that

$$\mathcal{Q}_n^2 = \{\pi = \pi_1\pi_2\cdots\pi_n \mid \pi = 23\cdots n1 \text{ or } \pi = \sigma n(n-1)\tau 1\},$$

where σ (possibly empty) is increasing and τ (possibly empty) is decreasing. A permutation $\pi \in \mathcal{Q}_n^2$ with the form $\pi = \sigma n(n-1)\tau 1$ is uniquely determined by the

set consisting of the elements in σ . We can choose the elements of σ in $\binom{n-3}{r}$ way if σ contains r elements. Then we deduce that

$$|\mathcal{Q}_n^2| = 1 + \sum_{r=0}^{n-3} \binom{n-3}{r} = 2^{n-3} + 1$$

for $n \geq 3$. Therefore we have

$$g(n) = g(n-1) + 2^{n-3} + 1$$

for $n \geq 3$. Taking into account the initial conditions $g(2) = 2$, the unique solution is $g(n) = 2^{n-2} + n - 1$. This completes the proof. ■

Acknowledgments

The work was supported by the National Natural Science Foundation of China (11801378).

References

- [1] K. Archer, A. Geary, Powers of permutations that avoid chains of patterns, arXiv:2312.14351v1 [math.CO].
- [2] M. Bóna, Combinatorics of Permutations, second ed., CRC Press, 2012.
- [3] M. Bóna, R. Smith, Pattern avoidance in permutations and their squares, *Discrete Math.*, **342(11)** (2019), 3194–3200.
- [4] A. Burcroff and C. Defant, Pattern-avoiding permutation powers, *Discrete Math.*, **343(11)** (2020), 112017.
- [5] T. Gannon, The cyclic structure of unimodal permutations, *Discrete Math.*, **237** (2001), 149–161.
- [6] S. Kitaev, Patterns in permutations and words. Springer Verlag (EATCS monographs in Theoretical Computer Science book series), 2011.
- [7] N.J.A. Sloane, The on-line encyclopedia of integer sequences, <http://oeis.org>.
- [8] P.A. MacMahon, Combinatory Analysis, Cambridge University Press, London, 1915/16. 1.
- [9] J. Pan, On a conjecture about strong pattern avoidance, *Graphs and Combin.*, **39(2)** (2023), <https://doi.org/10.1007/s00373-022-02602-y>.