Probing the spectrum of the magnetar 4U 0142+61 with JWST

JEREMY HARE,^{1,2,3} GEORGE G. PAVLOV,⁴ BETTINA POSSELT,^{5,4} OLEG KARGALTSEV,⁶ TEA TEMIM,⁷ AND STEVEN CHEN⁶

¹NASA Goddard Space Flight Center, Greenbelt, Maryland, 20771, USA

²Center for Research and Exploration in Space Science and Technology, NASA/GSFC, Greenbelt, Maryland 20771, USA

³ The Catholic University of America, 620 Michigan Ave., N.E. Washington, DC 20064, USA

⁴Department of Astronomy & Astrophysics, Pennsylvania State University, 525 Davey Lab, University Park, PA 16802, USA

⁵Department of Astrophysics, University of Oxford, Denys Wilkinson Building, Keble Road, Oxford OX1 3RH, UK

⁶Department of Physics, The George Washington University, 725 21st St. NW, Washington, DC 20052

⁷Princeton University, 4 Ivy Lane, Princeton, NJ 08544, USA

ABSTRACT

JWST observed the magnetar 4U 0142+61 with the MIRI and NIRCam instruments within a 77 min time interval on 2022 September 20–21. The low-resolution MIRI spectrum and NIRCam photometry show that the spectrum in the wavelength range $1.4-11 \ \mu$ m range can be satisfactorily described by an absorbed power-law model, $f_{\nu} \propto \nu^{-\alpha}$, with a spectral slope $\alpha = 0.96 \pm 0.02$, interstellar extinction $A_V = 3.9 \pm 0.2$, and normalization $f_0 = 59.4 \pm 0.5 \ \mu$ Jy at $\lambda = 8 \ \mu$ m. These observations do not support the passive disk model proposed by Wang et al. (2006), based on the Spitzer photometry, which was interpreted as evidence for a fallback disk from debris formed during the supernova explosion. We suggest a nonthermal origin for this emission and source variability as the most likely cause of discrepancies between the JWST data and other IR-optical observing campaigns. However, we cannot firmly exclude the presence of a large disk with a different dependence of the effective disk temperature on distance from the magnetar. Comparison with the power-law fit to the hard X-ray spectrum above 10 keV, measured by NuSTAR contemporaneously with JWST, shows that the X-ray spectrum is significantly harder. This may imply that the X-ray and IR nonthermal emission come from different sites in the magnetosphere of the magnetar.

1. INTRODUCTION

Magnetars are neutron stars with extremely large inferred magnetic fields $(B \sim 10^{14} - 10^{15} \text{ G})$ and long spin periods $(P = 1 - 12 \text{ s}; \text{Olausen } \& \text{ Kaspi } 2014^1)$. Their persistent X-ray luminosities, substantially exceed their spin-down energy loss rates, suggesting that magnetar emission is powered by their strong magnetic fields. Soft X-ray (0.5–10 keV) spectra of magnetars are often described by a two-component, blackbody (BB) + powerlaw (PL), model with $kT \sim 0.4$ –0.6 keV, and spectral slope $\alpha \sim 1-3$ (energy flux density $f_{\nu} \propto \nu^{-\alpha}$). Magnetars have been also detected in hard X-rays (10-100 keV), where their spectra show an upturn (νf_{ν} increases with increasing photon energy). For several magnetars observed multiple times in the NIR, variability has been reported (e.g., Durant & van Kerkwijk 2005; Hulleman et al. 2004).

Although the isolated and non-accreting magnetar interpretation is widely accepted, other hypotheses have also been discussed. One alternative explanation attributes the quiescent emission of magnetars to low-rate accretion, possibly from a "fallback" disk (e.g., Chatterjee & Hernquist 2000; Perna et al. 2000; Ekşı & Alpar 2003). Fallback in supernovae occurs when supernova ejecta is accreted by the newly formed compact object (Colgate 1971, 1988; Chevalier 1989). This would naturally explain the long magnetar spin periods but can hardly explain the powerful outbursts and high-energy tails (although, see e.g., Kylafis et al. 2014, Zezas et al. 2015). However, even if the fallback disks are not responsible for all the X-ray behavior of magnetars, they could still exist and manifest themselves in different ways and/or at other wavelengths.

Being one of the brightest magnetars, the "anomalous X-ray pulsar" 4U 0142+61 was detected in X-rays with Uhuru (Forman et al. 1978), and its X-ray period, $P \simeq 8.7$ s, was discovered with EXOSAT (Israel et al. 1994). The energy loss rate (or spin-down power) of the magnetar, $\dot{E} = 1.2 \times 10^{32}$ erg s⁻¹, is about 3 or-

jeremy.hare@nasa.gov

¹ See the McGill magnetar catalog at http://www.physics.mcgill. ca/~pulsar/magnetar/main.html

ders of magnitude lower than its quiescent X-ray luminosity, $L_{2-10 \text{ keV}} \approx 1 \times 10^{35}$ erg s⁻¹ at the the most likely distance d = 3.6 kpc. 4U 0142+61 exhibits a typical magnetar-like X-ray spectrum and energy-dependent pulse profiles with a pulsed fraction of $\approx 4\%$ at energies of a few keV (Woods & Thompson 2006). On 2007 February 8 4U 0142+61 produced a strong outburst detected in RXTE monitoring data (Gavriil et al. 2008). During the outburst, the peak flux in the 2–60 keV band exceeded the quiescent level by a factor of 500–1000. Recent X-ray polarization measurements (Taverna et al. 2022) are consistent with a model in which thermal radiation from the magnetar surface is reprocessed by scattering off charged particles in the magnetosphere.

At longer wavelengths, an optical source having peculiar colors was detected in Keck observations by Hulleman et al. (2000) at the X-ray position of 4U0142+61, and it was the first identified optical counterpart of a magnetar. Its faintness ($R \approx 25$ in 1999 September) argued against an active accretion disk as the source of the observed optical emission. In addition, strong optical pulsations, with a pulsed fraction of $\approx 28\%$ (Kern & Martin 2002; Dhillon et al. 2005), were detected at the X-ray period, suggesting that most of the optical emission originates in the magnetar's magnetosphere (but see Ertan & Cheng 2004; Ertan et al. 2007). The optical and X-ray pulse profiles show similar morphologies and are approximately in phase with each other. The NIR emission of 4U 0142+61 was first investigated by Hulleman et al. (2004) who found the $K(K_s)$ magnitudes to be in 19.6–20 mag range from 3 observations with NIRC on Keck I. Hulleman et al. (2004) inferred significant variability in NIR, later confirmed by Durant & van Kerkwijk (2006) who re-analyzed all the NIR-optical data. However, the optical-NIR variability was later disputed by Muñoz-Darias et al. (2016) who re-analyzed some of the archival data and claimed that the source was not variable in optical data over a 12 year timespan. 4U 0142+61 was also observed with HST WFC3 in NIR in 2018 and detected at 22.64 ± 0.01 and 22.07 ± 0.01 mag (in the AB magnitude system) in F125W and F160W filters, respectively (Chrimes et al. 2022), which is slightly brighter than inferred from previous ground-based observations in J and H filters.

As part of a systematic search for supernova debris disks around young NSs, predicted by current supernova models (Heger et al. 2003), 4U 0142+61's mid-IR counterpart was detected with Spitzer IRAC in the $4.5 \,\mu\text{m}$ and $8.0 \,\mu\text{m}$ bands by Wang et al. (2006). From these measurements and earlier optical-NIR observations, which were not contemporaneous with the Spitzer observations, Wang et al. (2006) inferred an IR excess and concluded that this emission emerges from a passive, non-accreting disk, illuminated by the magnetar's X-rays. After correcting for the interstellar reddening (assuming $A_V = 3.5$), Wang et al. (2006) found that the intrinsic IR spectrum was consistent with a 920 K blackbody or, even better, with a multi-temperature (700-1200 K) thermal disk model. They proposed that the disk is a dusty remnant of fallback material from the supernova that created the magnetar, which would make it the first supernova debris disk directly detected around a young NS. Four additional IRAC observations, taken 2–3 weeks after 4U 0142+61 showed an X-ray burst on 2007 February 7, showed virtually the same fluxes in the 4.5 um and 8.0 um bands (Wang & Kaspi 2008).

In a subsequent paper, Wang et al. (2008) analyzed Spitzer IRS data and argued that the spectrum may contain a 9.7 μ m emission feature, which could be a signature of dust emission from silicate grains. From a Spitzer MIPS observation, Wang et al. (2008) also reported an upper limit at 24 μ m, showing a turndown of the magnetar's spectrum at longer wavelengths.

These findings motivated us to propose contemporaneous low-resolution spectroscopy in IR and photometry in NIR with JWST, supplemented by Swift-XRT and NuSTAR observations in X-rays (Pavlov et al. 2021). Section 2 provides a description of the MIRI, NIRCam, Swift-XRT, and NuSTAR observations and their analyses. We discuss our findings in Section 3 and conclude with a summary in Section 4.

2. OBSERVATIONS AND DATA ANALYSIS

2.1. JWST NIRCam imaging and photometry

The Near Infrared Camera (NIRCam; Rieke et al. 2023) was used to image the magnetar and measure its flux in two spectral bands. The NIRCam observations were carried out on 2022 September 21 from 00:34:19 to 00:43:48 UTC. The F250M and F140M filters were employed in the long and short-wavelength NIRCam channels, using the NRCBLONG and NRCB1 detectors, respectively, in the SUB400P subarray configuration. The pivot wavelengths and bandwidths for these filters are listed in Appendix A and Figure 1 shows the wavelength dependencies of the filter throughputs.

We used the RAPID readout pattern, having seven total dithered integrations and 10 groups per integration. The effective scientific exposure time in each of the F250M and F140M filters is 115.9 s. We use the pipelineprocessed data (calibration software version 1.11.4). In the drizzle process, a pixel shrinking of 1.0, pixel scale ratio of 1.0, and inverse variance map (IVM) weighting scheme were used for the final resampling of the data. The nominal pixel sizes are 62.7 mas and 30.7 mas for the F250M and F140M images, respectively.

Both the 250M and F140M images show a strongly nonuniform background, dominated by the so-called 1/f noise patterns (Schlawin et al. 2020). The details of the source photometry for these noisy images are described in Appendix A. The net source flux densities in the F250M and F140M filters are $f_{\nu} = 14.5 \pm 0.4 \,\mu$ Jy and $5.1 \pm 0.3 \,\mu$ Jy, respectively. The statistical uncertainties here and throughout the paper are reported at the 1σ level unless noted otherwise.

We compared the radial profile of 4U 0142+61 with those of other (likely stellar) sources in the field and found them to be consistent with each other. Hence, we conclude that there is no indication of any extended NIR emission around the magnetar.

2.2. JWST MIRI spectroscopy with Low Resolution Spectrometer

$2.2.1. \quad Observation \ and \ data \ reduction$

The Mid-Infrared Instrument (MIRI; Rieke et al. 2015) on JWST observed 4U 0142+61 on 2022 September 20 starting at 23:26:56 UTC. We employed the MIRI Low Resolution Spectrometer (LRS; Kendrew et al. 2015), which has a resolving power $\mathcal{R} \equiv \lambda/\Delta\lambda \approx 24(\lambda/1\,\mu\text{m}) - 80$ in the 5–12 μm range. A two-point dither ALONG SLIT NOD, which places the source at two different positions in the slit so that the sky background could be subtracted, was used. The MIRI LRS observed 4U 0142+61 for 1975.8 s, and the data were readout using the FASTR1 mode. An acquisition image was obtained in the MIRI F560W filter (using the FAST-GRPAVG readout) with an exposure time of 44.4 s.

We performed several different extractions of the 1D spectrum using the JWST pipeline to assess the impact of using non-default extraction parameters (e.g., extraction center and width, linear versus polynomial trace, different background regions). For instance, some MIRI LRS spectra have been found to be offset from the aperture position used by the standard pipeline, which can lead to poor extractions of the 1D spectrum (see, e.g., DerKacy et al. 2023). These additional extractions did not exhibit any significant shift in the normalization or shape of the spectrum compared to the spectrum extracted by the standard pipeline. Therefore, we simply used the JWST pipeline extraction of the 1D spectrum for analysis, which was extracted using version 1.13.3 of the calibration pipeline (Bushouse et al. 2023). We analvzed the spectrum in the 5–11 μ m wavelength range. since that is where the source dominates over the background.

Several issues remained with the spectrum after reducing it with the standard JWST pipeline. The first is that there was a hot pixel in one of the MIRI LRS observation nods that was not removed by the pipeline. The hot pixel appears as a very bright pixel in the 2D spectrum and as a single, bright but narrow line-like feature in the 1D spectrum at 9.9 μ m. To remove this feature, we took the mean of the nearest 4 points (spanning ~ 0.07 μ m), using the values of the two next longer and shorter wavelengths, and replaced the hot pixel value with this mean value. The second issue is that two points near 6.5 μ m did not have any associated uncertainties, so we adopted the error bars from neighboring points for these two points.

Another issue is that the uncertainties of the spectral data points appear to be underestimated (e.g., neighboring values often fluctuate by a much larger amount than the pipeline-calculated uncertainties). Because of this error underestimation, one has to add hypothetical systematic errors to get acceptable spectral fits with reasonable models (see Section 2.2.2).

2.2.2. Modeling the JWST spectrum of 4U0142+61

We start from fitting the MIRI LRS spectrum with two simple models. Firstly, following Wang et al. (2006), we use an absorbed blackbody (BB) model to look for a thermal IR source, such as a small disk with a nearly constant temperature. To account for absorption, here and in other fits and estimates, we use the dust_extinction package (Gordon et al. 2023), with the V band extinction A_V as a fitting parameter. We found that the best-fit BB model has a temperature $T = 562.2 \pm 0.8$ K and an equivalent sphere radius $R = (11.40 \pm 0.04) R_{\odot}$ (at d = 3.6 kpc), but the fit is clearly unacceptable (see Figure 1), with a large reduced chi-squared, $\chi^2_{\nu} = 14$ for $\nu = 184$ degrees of freedom (dof). Additionally, the best-fit absorption in this model is $A_V = 0$, which is inconsistent with the previously reported value of $A_V = 3.5 \pm 0.4$ (Durant & van Kerkwijk 2006). The best-fit BB model and residuals to the LRS spectrum are shown in the top panel of Figure 1. Note that the NIRCam points are not used in the fit (or χ^2 calculation), but the extrapolation to the NIR-Cam wavelengths also show this model is incompatible with the data. Thus, we conclude that the 5–11 μm emission cannot be explained by a single-temperature thermal model.

The second model used is an absorbed powerlaw (PL) model, i.e., $f_{\nu} = f_0(\lambda/\lambda_0)^{\alpha}10^{-0.4A_{\lambda}} = f_0(\nu/\nu_0)^{-\alpha}10^{-0.4A_{\nu}}$. Fitting with this model yields an extinction parameter $A_V = 3.94 \pm 0.07$, spectral index $\alpha = 0.963 \pm 0.005$, and normalization $f_0 = 59.4 \pm 0.2 \,\mu$ Jy,





A_V<0.002

T=562.2±0.8 K

Figure 1. Spectral fits and fit residuals of the observed MIRI LRS spectrum (blue points) with absorbed blackbody (top) and power-law (bottom) models. The NIRCam F140M and F250M flux densities (black) are shown but not included in the fits. The semi-transparent red areas show wavelength dependencies of the throughputs of these filters. The residuals are defined as $\Delta \chi = (\text{data} - \text{model})/\text{error}$, where the values of 'data' and 'error' are supplied by the JWST calibration pipeline.



Figure 2. 1σ and 3σ confidence contours in the extinction - spectral slope plane for the absorbed PL fits to the MIRI LRS only and MIRI LRS+NIRCam data. Two sets of contours are shown, one set shows the contours using the pipeline-produced MIRI LRS uncertainties (err1), while the other shows the contours when inflating the MIRI LRS uncertainties by a factor of 3 (err3; see Section 2.2.2). Note that the NIRCam uncertainties are not inflated in the latter case.

at $\lambda_0 = 8 \ \mu\text{m}$. The fit is significantly better than the absorbed BB model fit (see Figure 1), but $\chi^2_{\nu} = 7.1$ for $\nu = 184$ is still too large to make it formally acceptable. This is likely due to the uncertainties being underestimated, particularly at shorter wavelengths. To make the fit formally acceptable, one has to significantly increase the pipeline-produced uncertainties. For instance, we find that multiplying each of the pipeline-produced uncertainties by a factor of 3, gives a $\chi^2_{\nu} = 0.79$. This does not change the best-fit values but increases the uncertainties of the fitting parameters: $A_V = 3.9 \pm 0.2$, $\alpha = 0.963 \pm 0.015$, and $f_0 = 59.4 \pm 0.5 \ \mu \text{Jy}^2$.

The best-fit PL model and residuals to the LRS spectrum (using the pipeline-produced uncertainties) are shown in the bottom panel of Figure 1. Again we note that the NIRCam points are not used in the fit (or χ^2 calculation), but the extrapolation to the NIRCam wavelengths also shows good agreement with the data. The residuals of the PL fit show a hint of a spectral feature around 6 μ m. We, however, do not have a plausible interpretation of this feature and suspect it is due to calibration errors.

In Figure 2 we show the 1σ and 3σ confidence contours in the A_V - α plane, using both the pipeline-produced uncertainties (err1 in Figure 2) and the uncertainties multiplied by a factor of 3 (err3 in Figure 2). We also plot the confidence contours both including and excluding the NIRCam data points. The best-fit parameters do not change much with the inclusion of the NIRCam points.

We note that the pipeline-produced LRS spectrum, used in our fits, is oversampled – the number of wavelength bins (187 in the 5–11 μ m band) is larger than the number of resolution elements (~ 80 in the same band), which means that the original bins can be combined into broader bins without a significant loss of resolution. We have checked, however, that a moderate binning (e.g., by a factor of 3) does not change the best-fit parameter values and only very slightly increases their uncertainties.

We also note that throughout the analysis of the MIRI LRS spectrum several calibration updates have been released and have occasionally caused shifts larger than the formal uncertainties in the best-fit parameters. For instance, using an earlier calibration version, we found best-fit parameters of $A_V = 3.3 \pm 0.2$, spectral index $\alpha = 1.034 \pm 0.013$, and normalization $f_0 = 55.6 \pm 0.4 \,\mu\text{J}$ with a similar $\chi^2_{\nu} = 0.76$ (multiplying the data uncertainties by a factor of 3). This shift due to calibration is much larger than the uncertainties obtained for a given calibration version. However, the calibration of the LRS should be continually improving throughout the mission, so we use the best-fit values from the latest calibration for the remainder of this paper, but caution that a later calibration update may cause further changes in the best-fit parameters.

2.3. X-ray observations

To examine a contemporaneous spectral energy distribution (SED) from IR to hard X-ray, we carried out nearly simultaneous observations of the magnetar with the NuSTAR and Swift X-ray observatories.

2.3.1. Neil Gehrels Swift X-ray Telescope (XRT)

Swift-XRT observed 4U 0142+61 starting at 01:11:36 UTC on 2022-09-21. This observation overlapped with the NuSTAR observation but occurred about 25 minutes after the end of the JWST observation. The source was observed using windowed timing (WT) mode and having an exposure time of 2.2 ks. The source spectrum was extracted using the online Build XRT products pipeline³ (Evans et al. 2009). We binned the spectrum to have a

² We note that increasing the uncertainties for the BB model gives $\chi^2_{\nu} = 1.55$, but there are still strong systematic residuals, particularly at short wavelengths, while the NIRCam points remain discrepant with this model.

³ https://www.swift.ac.uk/user_objects/

minimum signal-to-noise ratio of five and fit it in the 0.5–10 keV energy range.

2.3.2. NuSTAR

The Nuclear Spectroscopic Array (NuSTAR) observed 4U 0142+61 simultaneously with JWST for 40.8 ks on 2022-09-20 (ObsID 30801028002) with the observation starting at 23:01:09 UTC (i.e., about 30 minutes before the start of the JWST observation). The data were processed and reduced using HEASOFT version 6.31.1, with version 2.1.2 of the NuSTAR data analysis pipeline using calibration version 20221115. The source spectrum was extracted from a 70'' radius circle centered on the source. The background spectrum was extracted from a 70'' circle placed in a source-free region on the same detector as the source. The source count rate was about 0.6 cts s⁻¹ for FPMA/B in the 3-78 keV energy range. The spectra were binned to have a minimum signal-to-noise ratio of five and fit in the 3-78 keV energy range. The X-ray spectra were fit with XSPEC version 12.13.0c (Arnaud 1996). We used the tbabs absorption model with wilms abundances (Wilms et al. 2000).

2.4. X-ray Spectral Modelling

We jointly fit the Swift and NuSTAR spectrum of 4U0142+61 to determine whether or not the hard PL tail connects smoothly to the IR spectrum, thus may be produced by the same particle population. To fit these spectra, we use an empirical absorbed BB+PL+PL model (model 1 hereafter), multiplied by a constant (frozen to 1 for Swift-XRT), to account for any cross-calibration differences between Swift and NuSTAR FPMA/B (i.e., const×tbabs×(bbodyrad+pow+pow) in XSPEC). We also fit an empirical BB+BB+PL (model 2 hereafter) also multiplied by a constant (i.e., const×tbabs×(bbodyrad+bbodyrad+pow)), as the previous work by Tendulkar et al. (2015) showed that it can fit the spectra equally well, but provides a significantly different photon index for the hard powerlaw tail.

We report the best-fit values for both models in Table 1. Statistically acceptable fits were achieved for both models, although the BB+PL+PL model fits the data better than the BB+BB+PL model. Overall, the bestfit parameters for model 1 from our dataset agree with the values found by Tendulkar et al. (2015). However, we find higher temperatures for both thermal components in model 2 and a harder photon index (i.e., 0.83 for our fits versus 1.03 from Tendulkar et al. 2015). These differences may be due to changes in the NuSTAR calibration between the NuSTAR observations and analyses. In any case, we are primarily focused on the hard PL component of the spectrum, as it is the most likely to connect to the IR. We find that the photon index of the hard power-law tail is $\Gamma_h = 0.35 \pm 0.07$ ($\alpha_h = -0.65 \pm 0.07$) or $\Gamma_h = 0.83 \pm 0.06$ ($\alpha_h = -0.17 \pm 0.06$) for model 1 and model 2, respectively. Figure 3 shows the best-fit for model 1, the fit residuals, and the unfolded spectrum and individual model components.

3. DISCUSSION

The JWST observations have shown that the spectrum of 4U 0142 in the 1.4–11 μ m wavelength range can be described by an absorbed PL model, $f_{\nu} \propto \nu^{-\alpha}$, with a slope $\alpha \approx 1$ and extinction $A_V \approx 3.9$. This spectrum corresponds the observed and dereddened fluxes $F_{1.4-11\,\mu\text{m}} \simeq 4.2 \times 10^{-14}$ and $F_{1.4-11\,\mu\text{m}}^{\text{dered}} \simeq 4.7 \times 10^{-14}$ erg cm⁻² s⁻¹, and "isotropic luminosity" $L_{1.4-11\,\mu\text{m}} \equiv$ $4\pi d^2 F_{1.4-11\,\mu\text{m}}^{\text{dered}} \simeq 7.3 \times 10^{31} d_{3.6}^2$ erg s⁻¹, where $d_{3.6} =$ d/3.6 kpc. This IR luminosity is close to the magnetar's spin-down power, $L_{1.4-11\,\mu\text{m}} = 0.61 \dot{E} d_{3.6}^2$, but it is much lower than the quiescent X-ray luminosity, $L_{1.4-11\,\mu\text{m}}/L_{0.5-80\,\text{keV}} \sim 6 \times 10^{-5}$ or 2.8×10^{-4} for the BB+PL+PL and BB+BB+PL X-ray spectral models, respectively.

In this section we compare the JWST results with the results of previous IR-optical observations and discuss the multi-wavelength spectrum of the magnetar and the nature of its IR-optical emission.

3.1. Comparison with the Spitzer results

Wang et al. (2006) reported on a Spitzer Infrared Array Camera (IRAC) observation of 4U0142+61, taken on 2005 January 17, however, Wang & Kaspi (2008) reanalyzed these data and reported more accurate spectral fluxes (based on PSF fitting photometry instead of aperture photometry) $f_{\nu} = 32.1 \pm 1.2$ and 48.8 ± 7.6 μ Jy at 4.5 and 8 μ m, respectively. Spitzer also observed the source four times after it underwent an Xray outburst on 2007 February 7. The IRAC observations spanned 2007 February 14-21 and measured average spectral fluxes of $f_{\nu} = 32.1 \pm 2.0$ and $59.8 \pm 8.5 \ \mu$ Jy at 4.5 and 8 μ m, respectively (Wang & Kaspi 2008), which are consistent with the observation from 2005. These points agree well with the JWST NIRCam photometry and MIRI spectrum, obtained almost 18 years later, with the largest offset between all of the Spitzer IRAC photometry and our PL fit being only about 1.5σ for the 4.5μ m flux measured from the 2005 observation (see Figures 4, 5, and 6).

Spitzer also observed the source with the Infrared Spectrograph (IRS) on 2006 January 22. The analysis of these data were reported by Wang et al. (2008), who discuss the possible detection of a silicate emission feature at 9.7 μ m. Unfortunately, the Spitzer IRS spectrum is too noisy for a meaningful comparison with the



Figure 3. Left: The observed Swift-XRT and NuSTAR X-ray spectra and best-fit BB+PL+PL model (model 1). The best-fit parameters are shown in Table 1. The residuals are shown in the bottom panel. Right: The same as left panel but for the unfolded νf_{ν} spectra. The model components are shown as dashed lines.

Table 1. Best-fit parameters for BB+PL+PL (model 1) and BB+BB+PL (model 2; see Section 2.4) fit to the near simultaneous Swift+NuSTAR X-ray spectrum of 4U 0142+61.

Model	Const.	$N_{\rm H}$	kT_1	$Norm_1^{b}$	Γ_s/kT_2	$Norm_2^{b,c}$	Γ_h	$Norm_3$ ^C	$F_X d$	χ^2/dof
#	$FPMA/FPMB^a$	$10^{22}~{\rm cm}^{-2}$	keV		$\dots/{\rm keV}$			10^{-5}	10^{-10} c.g.s.	
1	$1.00^{+0.04}_{-0.03}/0.99^{+0.04}_{-0.03}$	1.58(8)	0.47(1)	138^{+20}_{-18}	3.90(8)	0.22(3)	0.35(7)	$3.5^{+0.8}_{-0.7}$	7.7(5)	596.1/610
2	0.96(3)/0.95(3)	0.54(4)	0.464(8)	302^{+30}_{-26}	1.09(3)	1.2(2)	0.83(6)	20(3)	1.68(8)	634.0/610

^aNormalization constant to account for differences in normalization due to calibration uncertainties between Swift-XRT and NuSTAR. The constant value is frozen to 1 for the Swift-XRT spectrum and fit for the NuSTAR spectra.

^bbbodyrad normalization, $R_{\rm km}^2/d_{10}^2$, where $R_{\rm km}$ is the source radius in km and d_{10} is the source distance in units of 10 kpc.

^c Power-law normalization in photons keV⁻¹ cm⁻² s⁻¹ at 1 keV.

 d Unabsorbed X-ray flux in the 0.5-80 keV energy band.

MIRI LRS spectrum. However, we found no evidence of the 9.7 μ m emission feature in the LRS spectrum.

From the Spitzer observation with the Multiband Imaging Photometer (MIPS) on 2006 February 19, about a year after the Spitzer IRAC observations, Wang et al. (2008) reported a 3σ upper-limit of 38 μ Jy at 24 μ m. Posselt et al. (2011) re-analyzed this MIPS observation and found a possible source at the position of 4U 0142+61 in two independent, re-processed AORs (Astronomical Observation Requests), with a flux of 41 μ Jy, but it was only a 2σ significance detection. We use the latter measurement for this paper, since the analysis was performed using an updated Spitzer data reduction pipeline. This upper limit (or possible detection) from the MIPS observation strongly disagrees with the PL fit from the JWST data (see Figure 4).

3.2. Previous optical-NIR observations

There have been numerous optical and NIR observations of 4U 0142+61 (e.g., Hulleman et al. 2000, 2004; Dhillon et al. 2005; Morii et al. 2005; Durant & van Kerkwijk 2006; Muñoz-Darias et al. 2016; Chrimes et al. 2022). Wang et al. (2006) used published results from Hulleman et al. (2004) and Israel et al. (2004), which included four optical (B, V, R, I) and three NIR (J, H, K_s) points (see Figure 4). We also have gathered the results using different observations from those used in Wang et al. (2006) and plotted them in Figure 5 (to avoid overcrowding the plots due to many data points measured in similar bands), together with the best PL fit of the JWST data. In Figure 5, the 2007 Spitzer IRAC fluxes from Wang & Kaspi (2008) are plotted. The J, H, and K_s points, shown by filled triangles, were taken from the Gemini observations of 2004 November 2, reported by Durant & van Kerkwijk (2006). We added two square points that span the minimum and maximum K-band values reported by Durant & van Kerk-



Figure 4. Comparison of the dereddened (with $A_V = 3.9$) JWST measurements and model spectra with the results from Wang et al. (2006). The JWST MIRI LRS spectrum is plotted in blue, and the NIRCam points are plotted in red. The 2005 Spitzer IRAC fluxes (Wang & Kaspi 2008; see Section 3.1) and MIPS upper limit (Wang et al. 2008) are shown in yellow and green, respectively. The brown points show NIR (J, H, K_s) and optical (I, R, V, B) photometry collected by Wang et al. (2006) from previous publications. Note that the points from Wang et al. (2006) and Spitzer were dereddened with $A_V = 3.5$, but this difference causes only a small shift downward in the JWST points, and the large gap between the F140M and NIR points still remains. The black dashed line is the best PL fit ($\alpha = 0.963$, $A_V = 3.9$) to the MIRI LRS spectrum. The solid black line shows the model adopted by Wang et al. (2006), which is the sum of the spectrum of a disk (cyan) with $T \propto r^{-3/7}$, $R_{\rm in} = 2.9R_{\odot}$, $R_{\rm out} = 9.7R_{\odot}$, and a PL with $\alpha = -0.3$ (violet). The MIRI LRS and NIRCam points strongly disagree with this model.

wijk (2006). The F125W and F160W fluxes, in bands centered at 1.25 and 1.6 μ m, were obtained by Chrimes et al. (2022) from HST WFC3/IR observations of 2018 January 1, and the optical GTC/OSIRIS fluxes in the SDSS z, r, i, g bands (which were all observed on the same night) were adopted from the 2013 observations reported by Muñoz-Darias et al. (2016). We chose these datasets as they consist of observations across several filters which were obtained in the same observing run (i.e., nearly simultaneous for each individual dataset).

As shown in Figures 4 and 5, the results of these different observing campaigns exhibit offsets of different sizes from the PL fit to the MIRI LRS data. Notably, the points in Figure 4 show a strong discrepancy with the high-frequency extension of the LRS PL fit, as well as with the results of more recent NIR-optical observations. However, the strong discrepancies between the NIR data and LRS PL model still remain even in the more recent observations. The discrepancies, defined as $\Delta \chi = (\text{data-model})/\sigma$ between the broadband data points from Figure 5 and the PL fit to the LRS spectrum, are shown in Figure 6.

The largest discrepancy between the model and photometry ($\approx 15\sigma - 25\sigma$) occurs in the HST bands, primarily due to the small uncertainties on the HST points. The second largest discrepancy occurs in the K_s band, where the model overpredicts the K_s flux by a factor of about 1.7 (or $\sim 11\sigma$). Additionally, the H and J band points lie below the model but are less discrepant than the K_s band point (at about 10σ and 5σ , respectively). The two HST points, which lie above the nearby J and H band points, are still about a factor of 1.2 smaller than flux predicted by the best-fit model to the LRS data. Surprisingly, the NIRCam points agree remarkably well with the best-fit PL model, even though these points were not included when fitting the model. One



Figure 5. Plot of the dereddened (top) and observed (bottom) broadband spectral energy distribution of $4U\,0142+61$. The points in the top panel were dereddened with the best-fit $A_V = 3.9$ from the absorbed power-law model. The extrapolated best-fit absorbed PL model from the fit to the MIRI LRS spectrum is shown as a black line. Note that the Spitzer IRAC fluxes shown here are the averages from the 2007 observations reported by Wang & Kaspi (2008), whereas those shown in Figure 4 were taken from the 2005 observation reported in Wang & Kaspi (2008), thus are slightly different. The IR to optical photometry is taken from multiple publications (see Section 3.2). The black squares in the K_s band show the minimum and maximum K_s band fluxes reported by Durant & van Kerkwijk (2006).



Figure 6. Discrepancies between the Spitzer IRAC, Gemini NIRI, HST WFC3, and GTC OSIRIS observations of 4U 0142+61 and the best-fit absorbed PL fit to the JWST MIRI LRS spectrum, in units of σ . All points are shown as black triangles, with the exception of the HST points, which are shown as squares. The error bars shown correspond to $\pm 1\sigma$. Note that the Spitzer IRAC fluxes used here are the averages from the 2007 observations reported by Wang & Kaspi (2008).

of these points lies a bit redward of the K_s band point, while the other is between the H and J bands. JWST is supposed to have an absolute flux calibration that is much better than 10% (typically better than 2%), making calibration an unlikely cause for the discrepancies (Gordon et al. 2022). The optical points from Muñoz-Darias et al. (2016) also agree well with the extrapolated PL spectrum, but these points are the most reliant on the best-fit A_V , which has changed throughout the various calibration updates (see Section 2.2.2). We discuss potential causes for these discrepancies, such as variability and/or different emission sites for the IR and optical emission, in the following subsections.

3.3. Variability

Since the IR-optical observations of the magnetar were taken at different epochs, an obvious explanation for the discrepancies between the MIRI LRS spectrum (and NIRCam photometry) and previous broadband photometry is source variability. Originally, Hulleman et al. (2004) reported variability at NIR wavelengths, particularly in the K band, but noted that the source did not appear to be variable at optical wavelengths. Kband variability of greater than one magnitude, over a timescale of a few days, was reported by Durant & van Kerkwijk (2006). These authors also found that the source was possibly variable at optical wavelengths, by about half a magnitude in the I band. On the other hand, Muñoz-Darias et al. (2016) reported a lack of variability in the optical z, i, r and q bands, but they did not comment on the possibility of NIR variability. Chrimes et al. (2022) found that the NIR HST fluxes were consistent with the source being variable when compared to the NIR results of Hulleman et al. (2004) and Durant & van Kerkwijk (2006). Additionally, using AKARI observations, Kohmura et al. (2013) found a 64% lower flux at 4 μ m compared to the Spitzer 4.5 μ m flux and suggested that this discrepancy was due to variability.

There is also the MIPS 24 μ m upper limit, which is about a factor of three lower than the expected flux extrapolated from the best PL fit to the MIRI data. The factor of three discrepancy in flux is consistent with the extent of the variability reported by Durant & van Kerkwijk (2006) in the NIR (demonstrated in Figure 5 for the K_s band). Therefore, we cannot exclude the possibility that the mid-IR and NIR fluxes vary by about a factor of three, giving rise to the observed discrepancies. However, the good agreement of the Spitzer IRAC fluxes, from both Wang et al. 2006 and Wang & Kaspi 2008, with the JWST LRS + NIRCam spectrum, observed 17+ years later, as well as the lack of (or small scale) variability observed at optical wavelengths challenge this scenario. Additionally, Table 1 in Wang & Kaspi (2008) shows that the IRAC flux of 4U 0142+61 did not vary over the week long observing campaign.

It could also be the case that the IR and optical emission are produced by different mechanisms and/or at different sites near the magnetar, leading to variability in the IR that is not observed in the optical. However, there is a fair agreement between the optical observations and extrapolated best-fit PL model, suggesting that both emission components are produced by the same mechanism. Thus, the question of the origin and true extent of the possibly wavelength-dependent variability remains open.

3.4. Possible contribution from a fallback disk

Another explanation of the possibly different IR and optical spectra of $4U\,0142+61$ is the presence of a fall-back disk whose contribution to the observed emission

may be different in different wavelength ranges. The discovery of optical pulsations (Kern & Martin 2002; Dhillon et al. 2005), with a pulsed fraction of about 25%-30%, significantly higher than ~ 4% in X-rays, has led to perception that the optical emission comes from the magnetar's magnetosphere. If true, it means that the hypothetical fallback disk would be more easily detected in the IR. Wang et al. (2006) noticed that the previously reported V, R, I fluxes are consistent with a PL model with $\alpha = -0.3$, at an assumed $A_V = 3.5$. They used this PL in combination with a disk model to fit the IR + optical data (see Figure 4). That model assumes that the optically thick disk is heated by the X-ray emission from the magnetar, the disk's local effective temperature decreases with radial distance from the magnetar as $T \propto r^{-3/7}$, and each point on the disk surface emits BB radiation with this temperature. Wang et al. (2006) found that the best-fit disk model had inner and outer disk temperatures $T_{\rm in} = 1200$ K and $T_{\rm out} = 715$ K, and inner and outer disk radii $r_{\rm in} = 2.9 R_{\odot}$ and $r_{\rm out} = 9.7 R_{\odot}$, respectively, at d = 3.6 kpc and $\cos i = 0.5$, where i is the disk inclination. Because of the small difference between the inner and outer temperatures, the disk spectrum resembles a BB spectrum with a temperature of 920 K.

We reproduced the spectrum from Wang et al. (2006) using a simple disk spectrum model given by Equation (1). Figure 4 shows the model disk+PL spectrum together with the results of the optical/IR photometry (adopted from Wang et al. 2006 with updated IRAC fluxes from Wang et al. 2008) and the JWST data. It is immediately apparent that the JWST data are strongly inconsistent with this model. The MIRI spectrum does not have the anticipated curvature, and it overshoots the model at the long wavelength end. Additionally, the NIRCam F140M point greatly exceeds the anticipated model flux. Moreover, the most recent optical points from Muñoz-Darias et al. (2016) do not follow a PL with $\alpha = -0.3$, at least for the best-fit $A_V = 3.9$ found from the PL fit of the MIRI LRS data, as we see from Figure 5.

Although the PL+disk model used by Wang et al. (2006) is inconsistent with the JWST data, it does not mean that another disk model, perhaps in combination with a PL component, cannot describe the IR or IR+optical data. For instance, in the approximation of a multi-temperature BB flat disk, the flux density spectrum is

$$f_{\nu} = \frac{2\pi\cos i}{d^2} \frac{2h\nu^3}{c^2} \int_{r_{\rm in}}^{r_{\rm out}} \frac{r\,dr}{\exp[h\nu/kT(r)] - 1} \,. \tag{1}$$

If the radial dependence of the local effective temperature of a disk obeys a PL, $T(r) = T_{\rm in}(r/r_{\rm in})^{-\beta} =$ $T_{\rm out}(r/r_{\rm out})^{-\beta}$, then the disk spectrum is close to a PL with the slope $\alpha = 2/\beta - 3$, i.e.,

$$f_{\nu} \propto \nu^{3-2/\beta}$$
 at $(2/\beta - 1)kT_{\text{out}} \ll h\nu \ll kT_{\text{in}}$. (2)

Thus, the spectrum of a disk with a sufficiently large ratio of the inner and outer temperatures, $T_{\rm in}/T_{\rm out} \gg 2/\beta - 1$ (i.e., the large ratio of the outer and inner radii, $r_{\rm out}/r_{\rm in} \gg (2/\beta - 1)^{1/\beta}$) has a broad PL part. If we assume that the JWST PL spectrum is due to such a disk, then $\beta = 2/(\alpha + 3) = 0.505$ for $\alpha = 0.96$ (or $\beta = 0.5$ for $\alpha = 1$).

In a self-consistent disk model, the radial dependence of temperature should be derived from a balance between heating and emitted energies. In original models for a passive protostellar flat disk, illuminated by a central star, the dependence $T \propto r^{-3/4}$ was derived, which corresponds to $\alpha = -1/3$ (see Adams et al. 1987, and references therein). However, most protostellar disks exhibit flattish SEDs in the IR range, with $\alpha \approx 0.25$ –1 (e.g., Beckwith et al. 1990), which corresponds to $\beta \approx 0.5$ –0.6, similar to the slope we found for 4U0142+61. The difference between the observed and model temperature dependencies is likely due to some unrealistic assumptions in the original models. In particular, realistic disks should be flared rather than flat, and the local emission spectra are not BBs because the temperature of an irradiated passive disk should decrease inward (toward the disk mid-plane), and the disks become optically thin at large wavelengths (Chiang & Goldreich 1997). Therefore, the spectrum given by Equation (1)may be a crude (albeit useful) approximation.

The entire spectrum of an optically thick, multitemperature disk has three distinct regions. For the flat disk spectrum given by Equation (1), these are a Rayleigh-Jeans tail at $h\nu \ll kT_{\rm out}$, where $f_{\nu} \propto \nu^2$, a PL part given by Equation (2), and a high-frequency tail resembling a Wien spectrum, $f_{\nu} \propto \nu^2 \exp(-h\nu/kT_{\rm in})$ at $h\nu \gg (2/\beta - 1)kT_{\rm in}$. One could assume that the highfrequency part might correspond to the cutoff of the optical spectrum between the V and B bands (Hulleman et al. 2004), seen in Figure 4. However, such an interpretation requires a rather high temperature $T_{\rm in} \sim 7000$ K, which is well above the dust sublimation temperature for grains of any chemical composition (see e.g., Koch-Miramond et al. 2002; Temple et al. 2021). This means that it would be a gaseous disk. Ertan et al. (2007) argue that the Spitzer NIR data, together with the data from preceding NIR-optical observations, can be interpreted as emission of a gaseous fallback disk with temperatures $T_{\rm in} \approx 6500$ K, $T_{\rm out} \approx 360$ K, and radii $R_{\rm in} \approx 7 \times 10^9$ cm, $R_{\rm out} \approx 1.9 \times 10^{12}$ cm. The T(r) dependence provided in their Table 1 can be approximated

by a PL, $T(r) \propto r^{-0.51}$, which leads to the spectral slope $\alpha = 0.92$, virtually coinciding with our measurement. Ertan & Cheng (2004) and Ertan et al. (2007) discuss how such a disk could be formed and how its emission could be strongly pulsed.

Looking at Figure 5, one might assume that the highfrequency cutoff of the disk spectrum corresponds to the K band, which would not require such high values of $T_{\rm in}$. However, such an assumption does not look plausible because the K band is in between the simultaneously observed NIRCam 2.5 μ m and 1.4 μ m points that lie on the extrapolation of the LRS spectrum towards NIR. Although it is somewhat suspicious that all the K-band measurements are below the LRS spectrum, it is likely due to a chance coincidence.

One of appealing features of the disk interpretation of the (N)IR spectrum is that it could successfully capture the MIPS upper limit. Therefore, we also attempted to fit the disk model given by Equation (1) to the MIRI LRS spectrum, including the 2σ MIPS upper limit. We find, however, that this disk model is not able to account for the MIPS upper limit while also matching the slope of the LRS spectrum, particularly the points at wavelengths longer than 8μ m. Specifically, the rollover from the Rayleigh-Jeans tail to the flat part of the disk spectrum is not sharp enough to capture the longer wavelength portion of the LRS spectrum, so it undershoots the LRS data. Additionally, the slope of the flat part of this disk model is too soft and under-predicts the NIR-Cam points. If the MIPS point is ignored in the fit, then the best-fit disk model can successfully capture the MIRI LRS and NIRCam data, but it overpredicts the MIPS upper-limit by a factor of $\sim 2-3$.

It is important to note, however, that at long wavelengths the disk may become optically thin, so its flux at a given λ becomes smaller by a factor of $1 - \exp(-\tau_{\lambda})$, where the disk's optical thickness τ_{λ} decreases with increasing wavelength (e.g., Draine & Lee 1984). Therefore, we cannot exclude the possibility that a fallback disk with $T_{\rm in}/T_{\rm out} \gtrsim 15$ (hence, $r_{\rm out}/r_{\rm in} \gtrsim 200$) significantly contributes to at least the IR spectrum of $4U\,0142+61$.

Overall, variability of nonthermal (magnetospheric) emission seems to be a more natural explanation of the MIPS upper limit discrepancy with the MIRI + NIR-Cam PL spectrum. However, the true cause of this discrepancy and the origin of the 4U 0142+61's IR-optical emission can hardly be firmly established without nearly simultaneous observation with JWST and HST.

3.5. Comparison between IR and X-ray spectra

As we show in Section 2.2.2, the NIRCam and MIRI LRS data in the $\lambda = 1.4$ –11 µm band are well fit by an absorbed PL model with $\alpha \simeq 0.96$. Since PL spectra are typical for emission of relativistic particles, this might suggest that the observed IR spectrum comes from the magnetosphere of 4U0142+61 and can be connected with a nonthermal component of the X-ray spectrum, particularly, with the hard X-ray component, as suggested by Muñoz-Darias et al. (2016). However, the slopes of the IR and hard X-ray spectra are quite different, i.e., they cannot be connected without a large spectral break, $\Delta \alpha \simeq 1.6$ or $\Delta \alpha \simeq 1.1$ for models 1 and 2, respectively (see Figure 7). This suggests that even if the IR originates from the magnetosphere, the IR and hard X-ray emissions either come from two separate particle populations or are produced by different emission mechanisms. Future NIRCam timing observations can help to solidify the location of these IR/NIR emitting particles, the emission mechanism, and, depending on the IR spectrum at longer wavelengths, the strength of IR/NIR pulsations. Additionally, measuring the phaseshifts between the (N)IR and X-ray emission could help to determine the relative locations of the emission sites.

4. SUMMARY AND CONCLUSIONS

The JWST MIRI and NIRCam observations of the brightest magnetar 4U 0142+61, supplemented by Swift-XRT and NuSTAR observations in X-rays, have shown the following.

- The spectrum extracted from the MIRI LRS data in the 5–11 μ m wavelength range can be described with an absorbed PL model with the spectral slope $\alpha = 0.963 \pm 0.015$, extinction $A_V = 3.9 \pm 0.2$, and normalization $f_0 = 59.4 \pm 0.5 \ \mu$ Jy at $\lambda = 8 \ \mu$ m.
- The NIRCam F140M and F250M photometric data points, with flux densities $f_{\nu} = 5.1 \pm 0.3 \mu$ Jy and $14.5 \pm 0.4 \mu$ Jy at 1.4μ m and 2.5μ m, respectively, agree very well (within 1σ) with the extrapolation of this absorbed PL model.
- Although the results of Spitzer IRAC photometry at 4.5 μm and 8 μm, presented by Wang et al. (2006) (and re-analyzed by Wang & Kaspi 2008), agree with the JWST spectrum, the PL+disk model, suggested by Wang et al. (2006), strongly disagrees with the JWST results.
- The extrapolated absorbed PL model also agrees reasonably well with the optical data from Muñoz-Darias et al. (2016). However, there is still strong disagreement with older NIR-optical observations, particularly in the NIR bands, and with the Spitzer MIPS 24 μ m upper limit.



Figure 7. Broadband spectral energy distribution extending from IR to hard X-rays plotted with X-ray spectral model 1 (M1; left) and model 2 (M2; right, see Table 1). The broadband photometry is the same as plotted in Figure 5. The black lines show the best-fit absorbed PL model (with uncertainties on the slope) to the MIRI LRS spectrum. The blue stripe shows the X-ray energy covered by Swift-XRT and NuSTAR. The model components on the left (right) are from the best-fit unabsorbed BB+PL1+PL2 (BB1+BB2+PL) model (i.e., model 1 and 2, respectively) to the X-ray data and are shown in red, orange, and green respectively. The black dashed lines in both panels show the sum of the unabsorbed model components. This plot clearly shows that the hard X-ray power-law component of the spectrum cannot continuously extend into the IR without a large spectral break for either model 1 or 2.

- We favor variability as the explanation for these discrepancies, but we cannot exclude the possibility that a contribution from a fallback disk plays a role, at least in the IR part of the spectrum. The origin of the IR-optical emission of 4U 0142+61 can be tested with nearly simultaneous deep JWST and HST observations across a number of bands in the broad IR-optical range.
- We find that the MIR-NIR spectrum requires a large spectral break ($\Delta \alpha > 1$) to be consistent with the hard power law observed in X-rays, suggesting that the IR and hard X-ray emission are likely being emitted from two different particle populations.

Future approved JWST timing observations of 4U 0142+61 will help to further elucidate the nature of the IR emission.

Acknowledgments: We thank Sarah Kendrew, Brian Brooks, Bryan Hilbert for their useful advice on JWST data reduction. JH thanks George Younes and Zorawar Wadiasingh for useful discussions related to this project. This work is based on observations made with the NASA/ESA/CSA James Webb Space Telescope. The data were obtained from the Mikulski Archive for Space Telescopes at the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-03127 for JWST. These observations are associated with program #2635. Support for program #2635 was provided by NASA through a grant from the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-03127. JH acknowledges support from NASA under award number 80GSFC21M0002 and partial support through NuSTAR award NNH21ZDA001N-NUSTAR. This work made use of data supplied by the UK Swift Science Data Centre at the University of Leicester.

APPENDIX

A. MEASURING THE NIRCAM FLUX DENSITIES

The F250M and F140M images, shown in Figure 8, exhibit strong 1/f noise patterns (see Schlawin et al. 2020 for additional details). We use the approach of many "empty background apertures" (e.g., Skelton et al. 2014; Abramkin et al. 2022). By placing our 55 (30) overlapping background apertures in the same 1/f noise bands as the source aperture for F250M (F140M), we obtained error estimates of the net source counts, N_s . The uncertainty of these net counts,





Figure 8. The NIRCam F250M (left) and F140M (right) images of the magnetar. North is up, East to the left. The 1/f noise pattern is clearly visible in both images. For F250M (F140M), we use 55 (30) white background apertures with the same size as the source aperture in the "empty aperture" approach, see text.



Figure 9. Aperture measurements in the NIRCam F250M and F140M filters with total, net, and background counts (N_t, N_s, N_b) as indicated. The background apertures always have the same size as the source apertures. Considering the 1/f noise seen in Figure 8, and (corrected) flux uncertainties, 0.31" and 0.2" are chosen as the optimal apertures for F250M and F140M, respectively.

 σ_{N_s} , consists of contributions from the net counts themselves (estimated as a Poisson error) and the background noise contribution (the variance $\sigma_{N_b}^2$):

$$\sigma_{N_s} = \left(\sigma_{N_b}^2 + N_s/g\right)^{1/2},\tag{A1}$$

where g is the gain⁴, which converts the image counts (ADU) into detected electrons (see Table 2).

We choose the same size of source and background apertures. The source aperture is centered on the position obtained from a 2D Gaussian fit in the image. Figure 9 shows the measured total counts, median background counts, and net counts as functions of aperture radius. Aiming to minimize the noise to count ratio and considering the size of the 1/f noise band, we choose 0''.31 (or 5 pixels) and 0''.20 (or 6.5 pixels) as optimal aperture radii for the F250M and F140M images, respectively. Applying the respective (linearly interpolated) encircled energy fractions (versions ETCv2) ⁵ and the image unit conversion, we calculated the aperture-corrected net source counts and flux densities f_{ν} (see Table 2). Note that for both filters the uncertainties are mostly due to the nonuniform 1/f background, i.e., $\sigma_{N_s} \approx \sigma_{N_b}$.

⁴ See https://jwst-docs.stsci.edu/jwst-near-infrared-camera/ nircam-instrumentation/nircam-detector-overview/ nircam-detector-performance

Table 2. Pulsar NIRCam photometry using many background apertures

Band	$\lambda_{ m piv}$	$\delta\lambda$	$\mathcal{P}_{ u}$	g	$t_{\rm exp}$	$r_{\rm ap}$	ϕ N_t \mathcal{N}_b		$\overline{N}_b \pm \sigma_{N_b}$	N_s	$f_{ u}$	
	$\mu { m m}$	$\mu { m m}$	$\mu \rm{Jy}~s/cnt$	$\mathrm{e}^{-}/\mathrm{cnt}$	sec	"		cnt		cnt	cnt	μJy
F140M	1.404	0.142	0.121	2.05	115.9	0.2	0.847	4744	30	595 ± 231	4150 ± 236	5.1 ± 0.3
F250M	2.503	0.181	0.152	1.82	115.9	0.31	0.857	7898	55	-1602 ± 241	9501 ± 251	14.5 ± 0.4

NOTE— λ_{piv} and $\delta\lambda$ are the pivot wavelength and bandwidth of the filter, \mathcal{P}_{ν} is the filter's inverse sensitivity, g is the gain, t_{exp} is the exposure time, ϕ is the encircled count fraction in the source aperture with radius r_{ap} , N_t indicates the total (source plus background) counts in the source aperture, \mathcal{N}_b is the number of background apertures considered, \overline{N}_b and σ_{N_b} are the median and variance of these \mathcal{N}_b background measurements, $N_s = (N_t - \overline{N}_b) \pm \sigma_{N_s}$ lists the net (source) number of counts, $f_{\nu} = \mathcal{P}_{\nu} N_s (\phi t_{\text{exp}})^{-1}$ is the spectral flux density at the pivot wavelength.

REFERENCES

- Abramkin, V., Pavlov, G. G., Shibanov, Y., et al. 2022, ApJ, 924, 128. doi:10.3847/1538-4357/ac3a6f
- Adams, F. C., Lada, C. J., & Shu, F. H. 1987, ApJ, 312, 788. doi:10.1086/164924
- Arnaud, K. A. 1996, Astronomical Data Analysis Software and Systems V, 101, 17
- Beckwith, S. V. W., Sargent, A. I, Chini, R. S., & Guesten, R. 1990, AJ, 99, 924. doi:10.1086/115385
- Bushouse, H., Eisenhamer, J., Dencheva, N., et al. 2023, Zenodo
- Chatterjee, P. & Hernquist, L. 2000, ApJ, 543, 368. doi:10.1086/317097
- Chevalier, R. A. 1989, ApJ, 346, 847. doi:10.1086/168066
- Chiang, E. I. & Goldreich, P. 1997, ApJ, 490, 368. doi:10.1086/304869
- Chrimes, A. A., Levan, A. J., Fruchter, A. S., et al. 2022, MNRAS, 512, 6093. doi:10.1093/mnras/stac870
- Colgate, S. A. 1971, ApJ, 163, 221. doi:10.1086/150760
- Colgate, S. A. 1988, Supernova 1987A in the Large Magellanic Cloud, 341
- DerKacy, J. M., Ashall, C., Hoeflich, P., et al. 2023, ApJL, 945, L2. doi:10.3847/2041-8213/acb8a8
- Dhillon, V. S., Marsh, T. R., Hulleman, F., et al. 2005, MNRAS, 363, 609. doi:10.1111/j.1365-2966.2005.09465.x
- Draine, B. T. & Lee, H. M. 1984, ApJ, 285, 89
- Durant, M. & van Kerkwijk, M. H. 2005, ApJ, 627, 376. doi:10.1086/429714
- Durant, M. & van Kerkwijk, M. H. 2006, ApJ, 652, 576. doi:10.1086/507605
- Ekşı, K. Y. & Alpar, M. A. 2003, ApJ, 599, 450. doi:10.1086/379193

- Ertan, Ü., & Cheng, K. S. 2004, ApJ, 605, 840. doi:10.1086/382502
- Ertan, Ü., Erkut, M. H., Ekşı, K. Y., & Alpar, M. A. 2007. doi:10.1086/510303
- Evans, P. A., Beardmore, A. P., Page, K. L., et al. 2009, MNRAS, 397, 1177. doi:10.1111/j.1365-2966.2009.14913.x
- Forman, W., Jones, C., Cominsky, L., et al. 1978, ApJS, 38, 357. doi:10.1086/190561
- Gavriil, F. P., Dib, R., & Kaspi, V. M. 2008, 40 Years of Pulsars: Millisecond Pulsars, Magnetars and More, 983, 234. doi:10.1063/1.2900150
- Green, G. M., Schlafly, E., Zucker, C., et al. 2019, ApJ, 887, 93. doi:10.3847/1538-4357/ab5362
- Gordon, K. D., Bohlin, R., Sloan, G. C., et al. 2022, AJ, 163, 267. doi:10.3847/1538-3881/ac66dc
- Gordon, K. D., Clayton, G. C., Decleir, M., et al. 2023, ApJ, 950, 86. doi:10.3847/1538-4357/accb59
- Gordon, K. 2023, Zenodo
- Harrison, F. A., Craig, W. W., Christensen, F. E., et al. 2013, ApJ, 770, 103. doi:10.1088/0004-637X/770/2/103
- Heger, A., Fryer, C. L., Woosley, S. E., et al. 2003, ApJ, 591, 288. doi:10.1086/375341
- Hulleman, F., van Kerkwijk, M. H., & Kulkarni, S. R. 2000, Nature, 408, 689. doi:10.1038/35047024
- Hulleman, F., van Kerkwijk, M. H., & Kulkarni, S. R. 2004, A&A, 416, 1037. doi:10.1051/0004-6361:20031756
- Israel, G. L., Mereghetti, S., & Stella, L. 1994, ApJL, 433, L25. doi:10.1086/187539
- Israel, G., Stella, L., Covino, S., et al. 2004, Young Neutron Stars and Their Environments, 218, 247. doi:10.48550/arXiv.astro-ph/0310482

- Kaastra, J. S. & Bleeker, J. A. M. 2016, A&A, 587, A151. doi:10.1051/0004-6361/201527395
- Kendrew, S., Scheithauer, S., Bouchet, P., et al. 2015, PASP, 127, 623. doi:10.1086/682255
- Kern, B. & Martin, C. 2002, Nature, 417, 527. doi:10.1038/417527a
- Koch-Miramond, L., Haas, M., Pantin, E., et al. 2002, A&A, 387, 233. doi:10.1051/0004-6361:20020386
- Kohmura, T., Kaneko, K., Ikeda, S., et al. 2013, Neutron Stars and Pulsars: Challenges and Opportunities after 80 years, 291, 422. doi:10.1017/S1743921312024362
- Kylafis, N. D., Trümper, J. E., & Ertan, Ü. 2014, å, 562, 61. doi:10.1051/0004-6361/201322303
- Levan, A., Kouveliotou, C., & Fruchter, A. 2018, ApJ, 854, 161. doi:10.3847/1538-4357/aaa88d
- Morii, M., Kawai, N., Kataoka, J., et al. 2005, Advances in Space Research, 35, 1177. doi:10.1016/j.asr.2005.04.051
- Muñoz-Darias, T., de Ugarte Postigo, A., & Casares, J. 2016, MNRAS, 458, L114. doi:10.1093/mnrasl/slw024
- Newville, M., Stensitzki, T., Allen, D. B., et al. 2014, Zenodo
- Olausen, S. A. & Kaspi, V. M. 2014, ApJS, 212, 6. doi:10.1088/0067-0049/212/1/6
- Pavlov, G. G., Kargaltsev, O., Hare, J., & Posselt, B. 2021, JWST Proposal. Cycle 1, ID. #2635
- Perna, R., Hernquist, L., & Narayan, R. 2000, ApJ, 541, 344. doi:10.1086/309404
- Posselt, B., Schreyer, K., Ertan, Ü., et al. 2011, AstroPhysics of Neutron Stars 2010: A Conference in Honor of M. Ali Alpar, 1379, 152. doi:10.1063/1.3629504

- Rieke, G. H., Wright, G. S., Böker, T., et al. 2015, PASP, 127, 584. doi:10.1086/682252
- Rieke, M. J., Kelly, D. M., Misselt, K., et al. 2023, PASP, 135, 028001. doi:10.1088/1538-3873/acac53
- Schlawin, E., Leisenring, J., Misselt, K., et al. 2020, AJ, 160, 231. doi:10.3847/1538-3881/abb811
- Skelton, R. E., Whitaker, K. E., Momcheva, I. G., et al. 2014, ApJS, 214, 24. doi:10.1088/0067-0049/214/2/24
- Taverna, R., Turolla, R., Muleri, F., et al. 2022, Science, 378, 646. doi:10.1126/science.add0080
- Temple, M. J., Banerji, M., Hewett, P. C., et al. 2021, MNRAS, 501, 3061. doi:10.1093/mnras/staa3842
- Tendulkar, S. P., Hascöet, R., Yang, C., et al. 2015, ApJ, 808, 32. doi:10.1088/0004-637X/808/1/32
- Vrtilek, S. D., Raymond, J. C., Garcia, M. R., et al. 1990, A&A, 235, 162
- Wang, Z., Chakrabarty, D., & Kaplan, D. L. 2006, Nature, 440, 772. doi:10.1038/nature04669
- Wang, Z., Chakrabarty, D., & Kaplan, D. L. 2008, 40 Years of Pulsars: Millisecond Pulsars, Magnetars and More, 983, 274. doi:10.1063/1.2900161
- Wang, Z., Bassa, C., Kaspi, V. M., et al. 2008, ApJ, 679, 1443. doi:10.1086/587505
- Wang, Z. & Kaspi, V. M. 2008, ApJ, 675, 695. doi:10.1086/526449
- Wilms, J., Allen, A., & McCray, R. 2000, ApJ, 542, 914. doi:10.1086/317016
- Woods, P. M. & Thompson, C. 2006, Compact stellar X-ray sources, 547. doi:10.48550/arXiv.astro-ph/0406133
- Zezas, A., Trümper, J. E., & Kylafis, N. D. 2015, MNRAS, 454, 3366. doi:10.1093/mnras/stv1918