On small densities defined without pseudorandomness

Thomas Karam^{*}

May 8, 2024

Abstract

We identify an assumption on linear forms $\phi_1, \ldots, \phi_k : \mathbb{F}_p^n \to \mathbb{F}_p$ that is much weaker than approximate joint equidistribution on the Boolean cube $\{0, 1\}^n$ and is in a sense almost as weak as linear independence, but which guarantees that every subset of $\{0, 1\}^n$ on which none of ϕ_1, \ldots, ϕ_k has full image has a density which tends to 0 with k. This density is at most quasipolynomially small in k, a bound that is necessarily close to sharp.

Contents

1	Introduction		2
	1.1	Background on mod- p linear forms and basic difficulties	2
	1.2	Main results	4
2	Constant separation is much weaker than approximate independence		7
	2.1	Adding a new condition with small or absent effects	8
	2.2	A density with a power lower bound	9
3	Proof of the main result		11
	3.1	Exponentially small densities from the sunflower structure	12
	3.2	From sunflowers to small balls	13
	3.3	Finishing the proof	13
4	Open questions		14
	4.1	Quantitative bounds	15
	4.2	Remaining qualitative questions	15

*Mathematical Institute, University of Oxford. Email: thomas.karam@maths.ox.ac.uk.

1 Introduction

1.1 Background on mod-*p* linear forms and basic difficulties

All our statements in this paper will be uniform in the integer n, which will not play any significant role.

The present paper will involve restrictions of mod-p forms (that is, of linear forms $\mathbb{F}_p^n \to \mathbb{F}_p$) to subsets of \mathbb{F}_p^n such as $\{0,1\}^n$ and more generally S^n for some non-empty subset S of \mathbb{F}_p . These restrictions have been studied in several recent works, such as [5], [6], [11] (by Gowers and the author, for the first two) in the more general settings of polynomials and of abelian group homomorphisms, as well as in [10], which focused on the question of how many dense subsets of the cube $\{0,1\}^n$ could be built so that any two of them can be strongly distinguished by some mod-p form, in various senses.

Another context in which these restrictions of mod-p forms appear is that of obstructions to combinatorial statements of Hales-Jewett type, such as those discussed in [4]. We refer the reader to the introductions of [5], [6], [10] for a discussion of that connection.

Restrictions of \mathbb{F}_p^n to $\{0, 1, 2\}^n$ also arise in the recent work of Bhangale, Khot and Minzer [2] where they considerably improve the best known bounds on the size of subsets of \mathbb{F}_p^n which do not contain any length three arithmetic progression $\{x, x+a, x+2a\}$ where the common difference *a* is furthermore required to be an element of $\{0, 1, 2\}^n \setminus \{0\}$. As mentioned in their paper, the slightly stronger variant where $\{0, 1, 2\}$ is replaced by $\{0, 1\}$ had been asked by Hazla, Holenstein and Mossel [9] and emphasized by Green [7].

In this paper we continue the development of the basic theory of sets defined by restricted mod-p forms, in a new direction that focuses on a perhaps surprisingly weak condition - in particular, much weaker than the condition used in previous works [6], [10] - which nonetheless suffices to guarantee that a subset of $\{0, 1\}^n$ defined by a large conjunction of non-trivial conditions involving mod-p forms is not dense.

The distributions of mod-p forms and k-tuples of mod-p forms on $\{0,1\}^n$ for some integer k present additional difficulties compared to their counterparts on the whole of \mathbb{F}_p^n . On the latter, we have an equivalence between linear independence and joint equidistribution: a k-tuple of mod-p forms is jointly equidistributed (that is, it takes each value of \mathbb{F}_p^k with probability p^{-k}) if and only if it is a linearly independent family. On $\{0,1\}^n$, substantially more complicated behaviour arises (provided that $p \geq 3$): some forms such as x_1 do not even have full range, so even for k = 1 there is something to be said: if the mod-p form depends on a large number r of coordinates, then as we will recall in Proposition 1.2 and its proof a Fourier-analytic calculation shows that the form is approximately equidistributed, with an error term decreasing exponentially with r. If on the other hand the mod-p form only depends on some r coordinates, then divisibility arguments provide a lower bound on the distance to equidistribution (using that $p \geq 3$): the probability that the form takes any given value is an integer multiple of 2^{-r} , so its difference with 1/p is at least $1/2^r p$ in absolute value.

Moving from one to two mod-p forms ϕ_1 , ϕ_2 , we again do not have approximate joint

equidistribution nor even approximate independence of the events

$$\phi_1(x) = y, \phi_2(x) = z$$

for general linearly independent ϕ_1 , ϕ_2 , and general y, z when x is chosen at random in $\{0,1\}^n$ even if ϕ_1, ϕ_2 both depend on many coordinates: this can be seen by taking $\phi_2 = \phi_1 + x_1$ for an arbitrary choice of ϕ_1 . Rather, Fourier analysis there again shows (by a short calculation, of the kind that we shall do in a moment when proving Proposition 1.2) that a sufficient condition for $\phi_2(x)$ to be approximately equidistributed conditionally on any value of $\phi_1(x)$ is that $\phi_2 - b\phi_1$ has large support for any $b \in \mathbb{F}_p$, and that a sufficient condition for the pair $(\phi_1(x), \phi_2(x))$ to be approximately equidistributed is that $a\phi_1 + b\phi_2$ has large support for any $(a, b) \in \mathbb{F}_p^2 \setminus \{0\}$. That condition, in turn, extends to k-tuples of mod-p forms for any $k \geq 1$, and was already studied in [6]. We recall the relevant definition, statement and proof as starting points.

Definition 1.1. Let p be a prime. If $\phi : \mathbb{F}_p^n \to \mathbb{F}_p$ is a mod-p form defined by

$$\phi(x) = a_1 x_1 + \dots + a_n x_n,$$

then we say that the support $Z(\phi)$ is the set

$$\{i \in [n] : a_i \neq 0\},\$$

and that the support size of ϕ is the size of $Z(\phi)$.

If $k \ge 1$, $r \ge 0$ are integers, then we say that mod-p forms ϕ_1, \ldots, ϕ_k are r-separated if the support size of the linear combination

$$a_1\phi_1 + \cdots + a_k\phi_k$$

is at least r for every $(a_1, \ldots, a_k) \in \mathbb{F}_n^k \setminus \{0\}$.

Proposition 1.2 ([6], Proposition 2.4). Let p be a prime, let S be a subset of \mathbb{F}_p with size at least 2, and let $k \ge 1$, $r \ge 0$ be integers. If $\phi_1, \ldots, \phi_k : \mathbb{F}_p^n \to \mathbb{F}_p$ are r-separated mod-p forms, then for every $(y_1, \ldots, y_k) \in \mathbb{F}_p^k$ we have

$$|\mathbb{P}_{x \in \{0,1\}^n}(\phi_1(x) = y_1, \dots, \phi_k(x) = y_k) - p^{-k}| \le (1 - p^{-2})^r.$$

Proof. The probability on the left-hand side can be expressed as

$$\mathbb{E}_{x\in S^n} 1_{\phi_1(x)=y_1} \dots 1_{\phi_k(x)=y_k} = \mathbb{E}_{x\in S^n} \left((\mathbb{E}_{a_1\in\mathbb{F}_p} \omega_p^{a_1(\phi_1(x)-y_1)}) \dots (\mathbb{E}_{a_k\in\mathbb{F}_p} \omega_p^{a_k(\phi_k(x)-y_k)}) \right)$$
$$= \mathbb{E}_{a\in\mathbb{F}_n^k} F(a)$$

where F(a) is the Fourier coefficient

$$\mathbb{E}_{x\in S^n}\omega_p^{a_1(\phi_1(x)-y_1)+\dots+a_k(\phi_k(x)-y_k)}.$$

The contributions of each coordinate of x are independent, so for a fixed $a \in \mathbb{F}_p^k$ we may decompose

$$F(a) = \omega_p^{-(a_1y_1 + \dots + a_ky_k)} \prod_{z=1}^n \mathbb{E}_{x_z \in S} \omega_p^{(a_1\phi_{1,z} + \dots + a_k\phi_{k,z})x_z}$$

where $\phi_{i,z} \in \mathbb{F}_p$ is the coefficient of the form ϕ_i at the coordinate z for every $i \in [k]$ and every $z \in [n]$. It is not difficult to show (see for instance, [6, Lemma 2.1] and its proof) that whenever $t \in \mathbb{F}_p^*$ we have $|\mathbb{E}_{u \in S} \omega_p^{tu}| \leq 1 - p^{-2}$. Therefore

$$|F(a)| \le (1 - p^{-2})^{|Z(a_1\phi_1 + \dots + a_k\phi_k)|}$$

for every $a \in \mathbb{F}_p^k$. Using $F(0) = p^{-k}$ and the assumption, the desired inequality follows. \Box

The reader not familiar with mod-p linear forms restricted to $\{0,1\}^n$ may also want to view the previously described behaviours as analogous to those that arise when considering quadratic forms on the whole of \mathbb{F}_p^n . Indeed, for $p \geq 3$ a non-zero quadratic form $\mathbb{F}_p^n \to \mathbb{F}_p$ is not necessarily approximately equidistributed (or even surjective), but if it has sufficiently high rank then it is. Likewise, linearly independent k-tuples of quadratic forms (q_1, \ldots, q_k) are not approximately equidistributed in general, but if every linear combination

$$a_1q_1 + \cdots + a_kq_k$$

of them has large enough rank then they are. The first of these facts follows from diagonalising the quadratic form, and a generalisation to polynomials is discussed in a landmark paper [8] of Green and Tao. The second fact reduces to the first by a standard Fourieranalytic calculation.

1.2 Main results

Previous works [6], [10] relied primarily on Proposition 1.2 to show that a set defined by linear mod-p conditions has small density. One quantitative way in which Proposition 1.2 may be viewed as not so strong is that in order to immediately deduce from Proposition 1.2 an upper bound of the type

$$|\mathbb{P}_{x \in \{0,1\}^n}(\phi_1(x) = y_1, \dots, \phi_k(x) = y_k) - p^{-k}| \le cp^{-k}$$

for some fixed c > 0, the lower bound r on the supports of the linear combinations

$$a_1\phi_1 + \cdots + a_k\phi_k$$

with $a \in \mathbb{F}_p^k \setminus \{0\}$ that we must take grows linearly in k.

In particular, if we consider strict subsets E_1, \ldots, E_k of \mathbb{F}_p and the set Λ of $x \in \{0, 1\}^n$ satisfying all conditions

$$\phi_1(x) \in E_1, \dots, \phi_k(x) \in E_k,$$

as has been done in [6], then it suffices for E_1, \ldots, E_k to have size 2 or more for the upper bound on the density of Λ that comes from applying the bounds from Proposition 1.2 to be

$$2^k(1-p^{-2})^r$$

and for that upper bound to be meaningful for large k we must then require r to be at least linear in k.

Our main contribution in the present paper will be to show that there exists some lower bound on r that is *uniform* in k, and which suffices to guarantee that the density of Λ tends to 0 as k tends to infinity. We will state and prove our results in the general case where the alphabet of the variables is an arbitrary subset S of \mathbb{F}_p containing at least two elements, as the proofs in this setting are the same as those of the special case $S = \{0, 1\}$.

Definition 1.3. Let p be a prime, let S be a subset of \mathbb{F}_p with size at least 2, and let E be a strict subset of \mathbb{F}_p . Then there exists a smallest nonnegative integer

$$L = L(S, E) \le \lfloor \frac{|E| - 1}{|S| - 1} \rfloor + 1 \le |E|$$
(1)

such that for any $a_1, \ldots, a_L \in \mathbb{F}_p^*$ the set

$$a_1 S + \dots + a_L S \tag{2}$$

is not contained in E. We will write L(S) for $L(S, \mathbb{F}_p)$.

The existence of L follows from the k = 1 case of Proposition 1.2, but it also follows immediately from the Cauchy-Davenport inequality, which furthermore provides the inequality (1). We note that if $S = \{0, 1, ..., |S| - 1\}$ and $E = \{0, 1, ..., |E| - 1\}$, then (1) becomes an equality. We are now ready to state our main result in this paper.

Theorem 1.4. Let p be a prime, and let S be a subset of \mathbb{F}_p with size at least 2. Then there exists some a = a(p, S) > 0 such that the following holds. If $k \ge 1$ is an integer, E_1, \ldots, E_k are strict subsets of \mathbb{F}_p , and ϕ_1, \ldots, ϕ_k are mod-p forms satisfying

$$|Z(\phi_j - \phi_i)| \ge 2 \max_{1 \le t \le k} L(S, E_t) - 1$$
(3)

for any $1 \leq i < j \leq k$, then the density of points $x \in S^n$ satisfying

$$\phi_1(x) \in E_1, \dots, \phi_k(x) \in E_k$$

is at most $O(k^{-a/\log \log k})$ as k tends to infinity.

In the case where the sets E_1, \ldots, E_k all have size bounded above by some common value strictly less than p, Theorem 1.4 specialises as follows thanks to the inequality (1).

Corollary 1.5. Let p be a prime, let S be a subset of \mathbb{F}_p with size at least 2, and let $1 \leq r \leq p-1$ be an integer. Then there exists some a = a(p, S, r) > 0 such that the following holds. If $k \geq 1$ is an integer, E_1, \ldots, E_k are strict subsets of \mathbb{F}_p each with size at most r, and ϕ_1, \ldots, ϕ_k are mod-p forms satisfying

$$|Z(\phi_j - \phi_i)| \ge 2\lfloor \frac{r-1}{|S|-1} \rfloor + 1$$

for any $1 \leq i < j \leq k$, then the density of points $x \in S^n$ satisfying

 $\phi_1(x) \in E_1, \ldots, \phi_k(x) \in E_k$

is at most $O(k^{-a/\log \log k})$ as k tends to infinity.

The bound in the assumption of Theorem 1.4 is optimal, at least in the case where the sets E_1, \ldots, E_k are the same set E: indeed, by definition of L = L(S, E) we can find a_1, \ldots, a_{L-1} such that

$$a_1S + \dots + a_{L-1}S \subset E.$$

Taking ϕ_1, \ldots, ϕ_k to be mod-*p* forms with pairwise disjoint supports, and with non-zero coefficients equal to a_1, \ldots, a_{L-1} (respecting multiplicities), we have that $\phi_i(S^n) \subset E$ for every $i \in [k]$, and that

$$|Z(\phi_j - \phi_i)| = |Z(\phi_j)| + |Z(\phi_i)| = 2L - 2$$

for any $1 \le i < j \le k$.

We are about to use the following proposition, proved in [6], which informally states that the probability that a given mod-p form takes any prescribed value is either 0 or bounded below by some positive quantity. Proposition 1.6 will again be used in Section 3.

Proposition 1.6 ([6], Proposition 2.5). Let p be a prime, let S be a non-empty subset of \mathbb{F}_p , and let $\phi : \mathbb{F}_p^n \to \mathbb{F}_p$ be a linear form. Then for every $y \in \mathbb{F}_p$ we have

$$\mathbb{P}_{x\in S^n}(\phi(x)=y)=0 \text{ or } \mathbb{P}_{x\in S^n}(\phi(x)=y) \ge \beta(p,S)$$

where $\beta(p, S) = |S|^{-\lceil (p-1)/(|S|-1) \rceil}$.

We note that the bounds in Theorem 1.4 and in Corollary 1.5 cannot be too far from tight, because there are simple examples satisfying power lower bounds in the other direction (and furthermore, uniformly with respect to $r \ge 1$).

Example 1.7. Let $r, t \ge 1$ be integers and let $k = p^t$. Let $\psi_1, \ldots, \psi_t : \mathbb{F}_p^n \to \mathbb{F}_p$ be linear forms with pairwise disjoint supports each with size at least r. The k linear combinations ϕ_1, \ldots, ϕ_k of ψ_1, \ldots, ψ_t then satisfy $|Z(\phi_j - \phi_i)| \ge r$ for any $1 \le i < j \le k$, but the set

$${x \in {0,1}^n : \phi_1(x) = 0, \dots, \phi_k(x) = 0}$$

is the same as the set

$${x \in {0,1}^n : \psi_1(x) = 0, \dots, \psi_t(x) = 0}$$

so by Proposition 1.6 has density at least $2^{-(p-1)t} = k^{-a}$ where $a = \log(2^{p-1})/\log p$.

As Example 1.7 illustrates, the assumption that the difference of any two distinct modp forms of some family has support size bounded below by some integer is in particular insufficient on its own to guarantee that the joint distribution of the family of forms is anywhere near uniform. Towards the end of Section 2, we will discuss an example (Example 2.4) where a power lower bound on the density can be obtained even for a set defined in terms of conditions where the linear forms satisfy the much stronger condition of being r-separated. (However, the bound will not be uniform with respect to large r this time.) This will in turn illustrate that this stronger assumption does not remotely come close to ensuring approximate equidistribution either.

More generally, Section 2 will as a whole focus on r-separation for a fixed r and on illustrating the various ways that it comes short of guaranteeing several properties that are guaranteed by the assumption of Proposition 1.2. By contrast, in Section 3 we shall prove Theorem 1.4, the assumption of which is qualitatively weaker than separation. Finally, Section 4 will be devoted to some open questions.

Throughout, we will often write (ϕ, E) for a condition $\phi(x) \in E$ where x is an element of S^n , as was done in [6]. If ϕ, ψ are two mod-p forms, then we will say that the *support distance* (or more simply *distance*) between ϕ and ψ is the support size of $\phi - \psi$. If ϕ_0 is a mod-p form and $r \ge 0$ is an integer, then we will say that the *ball* with *radius* r centered at ϕ_0 is the set of mod-p forms ϕ such that the support size of $\phi - \phi_0$ is at most r.

Acknowledgements

The author is grateful to Timothy Gowers for helpful discussions in Spring 2019, when the content of the present paper arose. The author also thanks Noga Alon and Swastik Kopparty for encouraging comments on related projects involving restrictions to $\{0,1\}^n$ of objects defined on \mathbb{F}_p^n and for their optimism about this area. This research received funding from an EPSRC International Doctoral Scholarship (project number 2114524).

2 Constant separation is much weaker than approximate independence

Throughout this section we fix a prime $p \geq 3$. We will construct examples of situations where linear forms that are *r*-separated for some fixed integer *r* (or, in Subsection 2.1, rather satisfy a variant of this assumption that is focused on the last mod-*p* form) nonetheless exhibit behaviour that is far from joint independence in the probabilistic sense, in various ways. We will take $S = \{0, 1\}$ for simplicity, but the examples can be adapted rather straightforwardly to any non-empty $S \neq \mathbb{F}_p$.

Example 2.1. The most basic example is the case r = 1, that is, the assumption that the linear forms are merely linearly independent. The linear forms $x_1 + x_i$ with $i \neq 1$ are not approximately independent in the probabilistic sense, and that is even more striking if

we furthermore add the linear form x_1 to the family, since the value of x_1 then forces the values of all $x_1 + x_i$ to belong to $\{x_1\} + \{0, 1\}$. This example, and examples of sunflower type more generally, were the starting point of the paper [6] and have been foundational for its proofs.

2.1 Adding a new condition with small or absent effects

Let $r \geq 1$ be any fixed integer. In this subsection we focus on the effect of an additional condition, and correspondingly we will not work under the full assumption that the mod-p forms that we consider are r-separated, but that, informally speaking, the last condition is r-separated from all previous ones. More precisely our setting will be to consider "pre-existing" mod-p forms ϕ_1, \ldots, ϕ_k , and a "new" mod-p form ϕ satisfying

$$|Z(\phi - \sum_{i=1}^{k} a_i \phi_i)| \ge r \tag{4}$$

for every $a \in \mathbb{F}_p^k$. We begin by an example showing that it is possible to find an integer $k \geq 1$, mod-*p* forms ϕ_1, \ldots, ϕ_k , and a mod-*p* form ϕ satisfying (4) for every $a \in \mathbb{F}_p^k$ and yet still have that for some subsets E_1, \ldots, E_k, E with $E \neq \mathbb{F}_p$ the condition $\phi(x) \in E$ is implied by the conditions $\phi_i(x) \in E_i$ in the sense that every $x \in \{0, 1\}^n$ satisfying the latter also satisfies the former.

Example 2.2. Let q = (p-1)/2. We take $k = \lceil r/q \rceil$. For each $i \in [k]$ we define

$$\psi_{i} = x_{(2i-2)q+1} + \dots + x_{2(i-1)q}$$

$$\rho_{i} = x_{(2i-1)q+1} + \dots + x_{2iq}$$

$$\phi_{i} = \psi_{i} + \rho_{i}$$

and then further define

 $\phi = \psi_1 + \dots + \psi_k.$

For every $a \in \mathbb{F}_p^k$ the linear combination

$$\phi - \sum_{i=1}^{k} a_i \phi_i \tag{5}$$

has support size at least $kq \ge r$. Indeed, letting $J(a) \subset [k]$ be the set

$$\{i \in [k] : a_i \neq 0\},\$$

the support of the linear combination (5) contains the disjoint union

$$\bigcup_{i \in J(a)} Z(\rho_i) \bigcup \bigcup_{i \in [k] \setminus J(a)} Z(\psi_i),$$

which has size kq. But if $x \in \{0, 1\}^n$ satisfies $\phi_i(x) = 0$ for each $i \in [k]$, then $\psi_i(x) = 0$ for each $i \in [k]$ and hence $\phi(x) = 0$.

We now discuss another example, showing that even if some conditions (ϕ_i, E_i) do not imply some condition (ϕ, E) in the sense from the previous example, the ratio of the density of the subset

$$\{x \in \{0,1\}^n : \phi_1(x) \in E_1, \dots, \phi_k(x) \in E_k, \phi(x) \in E\}$$

over the density of the subset

$${x \in {0,1}^n : \phi_1(x) \in E_1, \dots, \phi_k(x) \in E_k}$$

can be arbitrarily close to 1.

Example 2.3. Let k be a fixed integer which we will later take to be large enough. We let $\psi_0, \psi_1, \ldots, \psi_k$ be the mod-p forms defined by

$$\psi_i = x_{ir+1} + \dots + x_{(i+1)r}$$

for every $i \in \{0\} \cup [k]$. We then take $\phi_i = \psi_0 + \psi_i$ for each $i \in [k]$ and $\phi = \psi_0$. By definition the linear forms ψ_1, \ldots, ψ_k all have the same distribution D on $\{0, 1\}^n$. The probability mass that this distribution puts on any element of \mathbb{F}_p is an integer multiple of 2^{-r} , and that cannot be 1/p. This shows in particular that D is not perfectly uniform.

Our construction is designed to amplify the effect of the lack of uniformity of D on the distribution of ϕ given suitable values of ϕ_1, \ldots, ϕ_k . We assume for simplicity that $r \ge p - 1$, so that

$$\mathbb{P}(\phi = y) > 0, D(v) > 0$$

for all $y, v \in \mathbb{F}_p$. We have

$$\mathbb{P}(\phi = y, \phi_1 = u, \dots, \phi_k = u) = \mathbb{P}(\phi = y)\mathbb{P}(\psi_1 = u - y)\dots\mathbb{P}(\psi_k = u - y)$$
$$= \mathbb{P}(\phi = y)D(u - y)^k.$$

for all $y, u \in \mathbb{F}_p$. Let E be the set of $y \in \mathbb{F}_p$ which minimise D(u-y). The ratio of the density of

$$\{x \in \{0,1\}^n : \phi_1(x) = u, \dots, \phi_k(x) = u, \phi(x) \in E\}$$

over the density of

$$\{x \in \{0,1\}^n : \phi_1(x) = u, \dots, \phi_k(x) = u\}$$

then becomes exponentially close to 1 as k tends to infinity.

2.2 A density with a power lower bound

We now return from the effect of one additional condition to what occurs more globally for a system of conditions. In the next example we will show that no integer $r \ge 1$ guarantees that the set of points of $\{0, 1\}^n$ satisfying a family of r-separated conditions has a density that (uniformly in n) is at most exponential in the number of conditions, and that a power lower bound on the density is even possible. **Example 2.4.** Let k be a fixed integer, let T = T(r, k) be an integer that we will choose later, let ψ_1, \ldots, ψ_k be mod-p forms supported inside

$$\mathcal{T} = \{k+1, k+2, \dots, k+T(r, k)\}$$

which we will also construct later, and let $\phi_i = \psi_i + x_i$ for each $i \in [k]$.

For the forms ϕ_1, \ldots, ϕ_k to be *r*-separated, it suffices that every linear combination

$$a_1\phi_1 + \cdots + a_k\phi_k$$

where at most r-1 values a_1, \ldots, a_k are non-zero has support size at least r, as otherwise that follows immediately from considering the contributions of x_1, \ldots, x_k to the support. In turn, for that it suffices that for these $a \in \mathbb{F}_p^k$ the linear combinations

$$a_1\psi_1 + \cdots + a_k\psi_k$$

have support size at least r.

To satisfy that we now choose the forms ψ_1, \ldots, ψ_k by induction: we begin by choosing some ψ_1 with $|Z(\psi_1)| \ge r$, and for any u < k, assuming that ψ_1, \ldots, ψ_u have been fixed we choose ψ_{u+1} to be some form supported inside \mathcal{T} and outside the union U_u of all balls with radius r-1 centered at any linear combination

$$a_1\psi_1 + \dots + a_u\psi_u$$

where at most r-1 values a_1, \ldots, a_u are non-zero. The number of such $a \in \mathbb{F}_p^u$ is at most

$$\sum_{i=0}^{r-1} p^i \binom{u}{i} \le p^r \binom{k}{r} \tag{6}$$

for k large enough (depending on r) and the number of mod-p forms supported inside \mathcal{T} and with support size at most r-1 is at most

$$\sum_{j=0}^{r-1} p^j \binom{T}{j} \le p^r \binom{T}{r}$$

for T large enough (depending on r). The total number of mod-p forms in U_u is hence at most

$$p^{2r}\binom{k}{r}\binom{T}{r} \le p^{2r}(kT)^r.$$

Meanwhile the number of possible mod-p forms supported inside \mathcal{T} is p^T , so provided that

$$p^T > p^{2r} (kT)^r \tag{7}$$

we may inductively choose ψ_1, \ldots, ψ_k as desired. As for (7), we may (for fixed p and r) choose T which satisfies (7) and which has a growth rate $T = O(\log k)$ in the value of k.

If $x \in \{0,1\}^n$ is such that $x_z = 0$ for every $z \in \mathcal{T}$, then for every $i \in [k]$ we have $\psi_i(x) = 0$ and hence $\phi_i(x) \in \{0,1\}$. The set of points $x \in \{0,1\}^n$ satisfying $\phi_i(x) \in \{0,1\}$ for every $i \in [k]$ therefore has density at least 2^{-T} , which is at least some negative power of k (that depends on p and r). A back-of-envelope calculation shows that for any $\epsilon > 0$ this density is at least

$$k^{-((1+\epsilon)\log 2/\log p)r}$$

once k is large enough depending on ϵ .

As Example 2.4 shows, r-separation for a fixed r is in a way not much stronger than linear independence, since linear independence of a family of $k \mod p$ forms implies that the union of the supports of the k forms has size at least k, whereas in this example it has size $k + O(\log k)$. It is hence all the more remarkable that Theorem 1.4 holds with its current assumptions.

Remark 2.5. The mod-p forms used in Example 2.4 satisfy an assumption that is much stronger than that of Theorem 1.4: not only does the support of the difference of any two of these forms have size at least r, but the support of any linear combination of them has size at least r. It is hence natural to wonder for a moment whether we could not improve the power lower bound still much further and contradict Theorem 1.4, by considering the linear combinations of all forms involved in Example 2.4 (as they still satisfy the assumption of Theorem 1.4). Where this line of thought hits a wall is that if

$$\phi = a_1\phi_1 + \dots + a_k\phi_k$$

is a linear combination with at least p-1 non-zero coefficients a_1, \ldots, a_k then the image of the set

$$\{x \in \{0,1\}^n : \forall z \in \mathcal{T}, x_z = 0\}$$

by ϕ is the full of \mathbb{F}_p , as can be seen by considering the contribution of the coordinates inside [k], which is independent from that of the other coordinates. Therefore, we cannot use that linear form to build a condition (ϕ, E) with $E \neq \mathbb{F}_p$ that is satisfied by this set. Since the number of other possibilities for (a_1, \ldots, a_k) is at most $p^p \binom{k}{p-1}$ (by an estimate similar to (6)), so at most a fixed power of k, the best that can be achieved by taking this enlarged set of forms is still a power lower bound in k (with a power divided by p-1compared to that of Example 2.4).

3 Proof of the main result

In this section we prove Theorem 1.4. Although there are several ways of writing this proof, the choice that we ultimately made was to do so in a way that mirrors the high-level structure of that of [6, Theorem 1.2] (performed in Section 3 of that paper). In both situations the proof proceeds in three successive stages establishing the desired result, in the cases where the linear forms that we start with (i) constitute a sunflower, (ii) are contained in a ball with bounded radius, and finally (iii) are completely arbitrary. Let us begin by recalling the relevant formal definition of a sunflower.

Definition 3.1. Let p be a prime, let I be a finite set, let $\phi_i : \mathbb{F}_p^n \to \mathbb{F}_p$ be a linear form for every $i \in I$, and let ϕ_0 be an additional such form. We say that the forms $\phi_0 + \phi_i$ with $i \in I$ constitute a sunflower with center ϕ_0 if the supports of the forms ϕ_i with $i \in I$ are pairwise disjoint.

The assumption of Theorem 1.4 is that any two distinct forms differ by at least some number M of coefficients. Our intermediate statements in stages (i) and (ii) will be more convenient to use and to prove with a modified assumption: that after subtracting the centre of the sunflower or of the ball to all forms, the resulting forms all have support size bounded below by some integer r. Both assumptions are of course closely related: the triangle inequality on the support distance immediately shows that for $r = \lceil M/2 \rceil$ the first implies the second except for possibly one of the forms.

3.1 Exponentially small densities from the sunflower structure

The most basic structure on mod-p forms that we will use to ensure that a set of points of S^n satisfying many conditions involving these forms has small density - exponentially small in the number of forms for now, although not for long - is a sunflower.

Proposition 3.2. Let p be a prime and let $k \ge 1$ be an integer. Suppose that E_1, \ldots, E_k are strict subsets of \mathbb{F}_p and that $\phi_0, \phi_1, \ldots, \phi_k : \mathbb{F}_p^n \to \mathbb{F}_p$ are linear forms such that $\phi_0 + \phi_1, \ldots, \phi_0 + \phi_k$ constitute a sunflower with center ϕ_0 and ϕ_1, \ldots, ϕ_k each have support size at least $\max_{1\le i\le k} L(S, E_i)$. Then the set

$$\{x \in S^n : (\phi_0 + \phi_1)(x) \in E_1, \dots, (\phi_0 + \phi_k)(x) \in E_k\}$$

has density at most $p(1 - \beta(p, S))^k$ inside S^n .

Proof. For every $y \in \mathbb{F}_p$ the probability

 $\mathbb{P}_{x \in S^n}((\phi_0 + \phi_1)(x) \in E_1, \dots, (\phi_0 + \phi_k)(x) \in E_k, \phi_0(x) = y)$

can be rewritten as

$$\mathbb{P}_{x \in S^n}(\phi_1(x) \in E_1 - \{y\}, \dots, \phi_k(x) \in E_k - \{y\}, \phi_0(x) = y),$$

which in turn is at most

$$\mathbb{P}_{x \in S^n}(\phi_1(x) \in E_1 - \{y\}, \dots, \phi_k(x) \in E_k - \{y\}).$$

Because the supports of ϕ_1, \ldots, ϕ_k are pairwise disjoint, the last expression is equal to the product

$$\mathbb{P}_{x \in S^n}(\phi_1(x) \in E_1 - \{y\}) \dots \mathbb{P}_{x \in S^n}(\phi_k(x) \in E_k - \{y\}).$$

The sets $E_1 - \{y\}, \ldots, E_k - \{y\}$ have the same sizes as E_1, \ldots, E_k respectively, and every term of the product is strictly less than 1 by definition of $L(S, E_1), \ldots, L(S, E_k)$, so is at most $1 - \beta(p, S)$ by Proposition 1.6. We obtain

$$\mathbb{P}_{x \in S^n}((\phi_0 + \phi_1)(x) \in E_1, \dots, (\phi_0 + \phi_k)(x) \in E_k, \phi_0(x) = y) \le (1 - \beta(p, S))^k$$

for every $y \in \mathbb{F}_p$ and conclude by the law of total probability.

3.2 From sunflowers to small balls

We next tackle the case where the assumption that the mod-p forms constitute a sunflower is relaxed to the assumption that they are contained in a ball of bounded radius. This assumption together with the other assumption of Proposition 3.2 say that the supports of ϕ_1, \ldots, ϕ_k are bounded both below and above. The main proof step is that such a family that has many forms necessarily contains a sunflower with many forms. The argument is quite standard: it is a variant of that establishing the Erdős-Rado sunflower theorem (proved in [3]).

Proposition 3.3. Let p be a prime, let $t \ge 1, k \ge 1, r \ge 0$ be integers. Assume that $\phi_0 + \phi_1, \ldots, \phi_0 + \phi_k : \mathbb{F}_p^n \to \mathbb{F}_p$ are linear forms and that ϕ_1, \ldots, ϕ_k have supports at most r. If $k \ge p^r r! t^r$ then we can find $I \subset [k]$ with $|I| \ge t$ such that the linear forms $\phi_0 + \phi_i$ with $i \in I$ constitute a sunflower with center ϕ_0 .

Proof. We proceed by induction on r. If r = 0 then the result is immediate. Suppose that for some $r \ge 1$ the result holds for r - 1. Let $M \subset [k]$ be a maximal set such that the linear forms ϕ_i with $i \in M$ have pairwise disjoint supports. If $|M| \ge t$ then we take I = Mand we are done. Otherwise, for every $i \in [k]$ the support of ϕ_i intersects the union

 $\cup_{j\in M} Z(\phi_j)$

of supports, which has size at most rt, in at least one element. By the pigeonhole principle we can find $J \subset [k]$ with size at least k/(p-1)rt such that all forms ϕ_i with $i \in J$ have a common element z in their support, and furthermore have the same coefficient $u \in \mathbb{F}_p$ at that element of the support. The forms $\phi_i - ux_z$ with $i \in J$ then all have supports with size at most r-1. We then conclude by the inductive hypothesis.

Our result in the case of forms contained in balls of bounded radius then follows from applying Proposition 3.3 and Proposition 3.2 successively.

Corollary 3.4. Let p be a prime, let $k \ge 1$ and $r \ge 0$ be integers. If $\phi_0, \phi_1, \ldots, \phi_k : \mathbb{F}_p^n \to \mathbb{F}_p$ are linear forms, and E_1, \ldots, E_k are strict subsets of \mathbb{F}_p satisfying

$$\max_{1 \le i \le k} L(S, E_i) \le |Z(\phi_i)| \le r$$

for every $i \in [k]$ then

$$\mathbb{P}_{x \in S^n}((\phi_0 + \phi_1)(x) \in E_1, \dots, (\phi_0 + \phi_k)(x) \in E_k) \le p(1 - \beta(p, S))^{(k/p^r r!)^{1/r}}$$

3.3 Finishing the proof

We are now ready to finish the proof of Theorem 1.4. Given an arbitrary family of mod-p forms, either this family contains a large family of forms that jointly is approximately equidistributed, in which case we are done, or it does not, and we then reduce to the case of forms contained in a ball with bounded radius.

Proof of Theorem 1.4. Let $\epsilon > 0$. We distinguish two cases. If we can find a set $I \subset [k]$ with size $u = \lceil p \log(2\epsilon^{-1}) \rceil$ satisfying

$$|Z(\sum_{i\in I} a_i\phi_i)| \ge 5p^4\log(2\epsilon^{-1})$$

for every $a \in \mathbb{F}_p^I \setminus \{0\}$ then by Proposition 1.2 we have

$$\mathbb{P}_{x \in S^n} (\forall i \in I, \phi_i(x) = y_i) \le 2p^{-u}$$

for every $y \in \mathbb{F}_p^I$, and hence

$$\mathbb{P}_{x \in S^n} (\forall i \in I, \phi_i(x) \in E_i) \le 2(1 - p^{-1})^u \le \epsilon.$$

If on the other hand we cannot find a set I as above, then we can partition [k] into at most p^u sets such that all forms $\phi_0 + \phi_i$ with index i in a given set are contained in some ball of radius at most $r = 5p^4 \log(2\epsilon^{-1})$. By the pigeonhole principle we can find such a set containing at least k/p^u forms. The assumption (3) together with the triangle inequality show that all but at most one of these forms is at a distance at least $\max_{1 \le i \le k} L(S, E_i)$ from the center of the ball. Using that

$$1 - \beta(p, S) \le \exp(-\beta(p, S)),$$

Corollary 3.4 then provides the upper bound

$$\mathbb{P}_{x\in S^n}(\forall i\in I, \phi_i(x)\in E_i) \le p\exp\left(-\beta(p,S)((k-p^u)/p^up^rr!)^{1/r}\right).$$

For the previous right-hand side to be at most ϵ , it suffices that

$$k \le 2p^u p^r r! (\beta(p, S)^{-1} \log(p\epsilon^{-1}))^r.$$

Plugging in the values of u and r in terms of ϵ , that becomes

$$k \le 2p^{p\log(2\epsilon^{-1})+1}p^{5p^4\log(2\epsilon^{-1})}(5p^4\log(2\epsilon^{-1}))!(\beta(p,S)^{-1}\log(p\epsilon^{-1}))^{5p^4\log(2\epsilon^{-1})}.$$
(8)

As ϵ tends to 0, the right-hand side (the dominant terms of which are the last two) grows as $\Omega(\epsilon^{-c(p)\log\log \epsilon^{-1}})$ for some c(p) > 0.

4 Open questions

Let us finish by discussing a few remaining questions left open by the present results and proofs.

4.1 Quantitative bounds

Example 2.4 shows in particular that the bounds obtained in the conclusion of Theorem 1.4 (and Corollary 1.5) cannot be improved to anything better than inverse power bounds in k. The bounds from Theorem 1.4 and Corollary 1.5 are therefore not too far from the optimal bounds, which leads immediately to our first question.

Question 4.1. What are the optimal bounds in Theorem 1.4 and Corollary 1.5 ? In particular, can they be taken to be a negative power of k ?

To obtain such bounds, it would suffice to make the last two terms of (8) each have a power dependence in ϵ^{-1} , as the first two already do. The third term would be replaced by a power of ϵ^{-1} assuming the following slight variant of the notorious sunflower conjecture, which would then be used instead of Proposition 3.3 and which could be of independent interest. (The sunflower conjecture itself was first formulated by Erdős and Rado in [3], with recent celebrated progress by Alweiss, Lovett, Wu and Zhang in [1].)

Conjecture 4.2. Let p be a (not necessarily prime) integer, and let $r \ge 1$. Then there exist $c_1(p), c_2(p) > 0$ such that if $k, t \ge 1$ are integers satisfying $k \ge c_1(p)t^{c_2(p)}$ and v_1, \ldots, v_k are elements of \mathbb{Z}_p^n each with at most r non-zero coordinates, then we can find a subset $I \subset [k]$ with size t and pairwise disjoint subsets $X_i \subset [n]$ with $i \in \{0\} \cup I$ such that for every $i \in I$ the set of non-zero coordinates of v_i is $X_0 \cup X_i$, and the restrictions of v_1, \ldots, v_k to their coordinates in X_0 coincide.

The last term of (8) presents, however, another difficulty on its own: fundamentally the extra $\log \log \epsilon^{-1}$ term in the exponent is due to the fact that the parameter u in that proof grows with ϵ^{-1} , and this dependence in turn appears to be quite basic and difficult to get around. This suggests that a proof which manages to obtain a power bound in Theorem 1.4 would involve substantial extra ideas compared to the current proof.

4.2 Remaining qualitative questions

In another direction we can try to understand better which sets of conditions

$$\{(\phi_i, E_i) : i \in I\}$$

lead to sets

$$\{x \in S^n : \forall i \in I, \phi_i(x) \in E_i\}$$

that are dense inside S^n and which do not. This question is perhaps best formulated in an infinitary manner. We consider a family of linear forms from $\mathbb{F}_p^{\mathbb{N}}$ to \mathbb{F}_p with finite support, that is, a family of linear forms of the type

$$x \mapsto a_1 x_1 + \dots + a_n x_n$$

for some finite n (that depends on the form) and some coefficients $a_1, \ldots, a_n \in \mathbb{F}_p$. If $k \ge 1$ is an integer, ϕ_1, \ldots, ϕ_k are such linear forms, and E_1, \ldots, E_k are subsets of \mathbb{F}_p , then we

say that the density of the satisfying set of the conditions $\phi_1(x) \in E_1, \ldots, \phi_k(x) \in E_k$ is the density of the set

$$\{x \in S^N : \phi_1^N(x) \in E_1, \dots, \phi_k^N(x) \in E_k\}$$

inside S^N , where N is a value of n that works for all linear forms ϕ_1, \ldots, ϕ_k , and the forms $\phi_1^N, \ldots, \phi_k^N : \mathbb{F}_p^N \to \mathbb{F}_p$ are defined by

$$\phi_i^N(x_1,\ldots,x_N) = \phi_i(x)$$

for every $i \in [k]$ and every $x \in \mathbb{F}_p^{\mathbb{N}}$. We say that a family of conditions $\{(\phi_i, E_i) : i \in I\}$ with $\phi_i : \mathbb{F}_p^{\mathbb{N}} \to \mathbb{F}_p$ and $E_i \subset \mathbb{F}_p$ for every $i \in I$ is *density-reducing* if there exists a function $F : \mathbb{N} \to \mathbb{N}$ tending to 0 at infinity such that whenever $I' \subset I$ is a finite set, then the density of the satisfying set of the conditions $\{(\phi_i, E_i) : i \in I'\}$ is at most F(|I'|).

For instance, the family of conditions

$$\{x_i = 0 : i \ge 1\}$$

is density-reducing for $S = \{0, 1\}$, since we may take $F(k) = 2^{-k}$, whereas the family of conditions

$$\{x_1 + x_i \in \{0, 1\} : i \ge 2\}$$

is not density-reducing for $S = \{0, 1\}$, as discussed in Example 2.1.

With this definition, the qualitative side of the present paper's contribution can be reformulated in terms of properties which ensure that a family of conditions $\{(\phi_i, E_i) : i \in I\}$ with $E_i \neq \mathbb{F}_p$ is density-reducing. While Proposition 1.2 showed that it suffices that for every integer r the set of forms $\{\phi_i : i \in I\}$ cannot be contained in finitely many balls with radius r, we now know by our Theorem 1.4 that it suffices to guarantee that for the single value r = 2p - 1. A definitive characterisation remains open.

Question 4.3. Let p be a prime, and let S be a non-empty subset of \mathbb{F}_p . Can we characterise families of conditions $\{(\phi_i, E_i) : i \in I\}$ that are density-reducing in S^n , where $\phi_i : \mathbb{F}_p^{\mathbb{N}} \to \mathbb{F}_p$ are linear forms with finite support and E_i are strict subsets of \mathbb{F}_p ?

In the special case $S = \mathbb{F}_p$, this question has a simple answer: the family is densityreducing if any only if we are in one of the following two cases: (i) the linear subspace $\langle \phi_i : i \in I \rangle$ has infinite dimension, or (ii) (the family involves finitely many pairwise distinct conditions and) and no $x \in S^n$ satisfies them, that is, informally speaking, the conditions are incompatible.

On the broader topic of understanding the behaviour of mod-*p* forms on $\{0, 1\}^n$ and on S^n we may want to better understand how conditions may be deduced formally from one another. Given a non-empty subset S of \mathbb{F}_p , if $\phi_1, \phi_2 : \mathbb{F}_p^n \to \mathbb{F}_p$ are two linear forms, and E_1, E_2 are two subsets of \mathbb{F}_p , then $\{(\phi_1 + \phi_2, E_1 + E_2)\}$ is implied by $\{(\phi_1, E_1), (\phi_2, E_2)\}$ in the sense of the inclusion

$$\{x \in S^n : \phi_1(x) \in E_1, \phi_2(x) \in E_2\} \subset \{x \in S^n : (\phi_1 + \phi_2)(x) \in E_1 + E_2\}$$

but it is also the case that we can deduce $(\phi_1 + \phi_2)(x) \in E_1 + E_2$ formally from $\phi_1(x) \in E_1$ and $\phi_2(x) \in E_2$. **Question 4.4.** Given a non-empty subset S of \mathbb{F}_p and a family $\{(\phi_i, E_i) : i \in I\}$ of mod -p conditions, is there a class of formal operations which is simple to describe and such that whenever (ϕ, E) is a mod-p condition satisfying

 $\{x \in S^n : \forall i \in I, \phi_i(x) \in E_i\} \subset \{x \in S^n : \phi(x) \in E\},\$

then (ϕ, E) can be formally deduced from the conditions (ϕ_i, E_i) with $i \in I$ using these operations?

References

- R. Alweiss, S. Lovett, K. Wu, J. Zhang, Improved bounds for the sunflower lemma, Ann. of Math. 194 (2021), 795-815.
- [2] A. Bhangale, S. Khot, D. Minzer, *Effective bounds for restricted 3-arithmetic progressions in* \mathbb{F}_p^n , arXiv:2308.06600 (2023).
- [3] P. Erdős and R. Rado, Intersection theorems for systems of sets, Journal of the London Mathematical Society, 35 (1960), 85–90.
- W. T. Gowers, The first unknown case of polynomial DHJ, https://gowers.wordpress.com/2009/11/14/the-first-unknown-case-of-polynomialdhj. (Blog post.)
- [5] W. T. Gowers and T. Karam, Equidistribution of high-rank polynomials with variables restricted to subsets of \mathbb{F}_p , arXiv:2209.04932 (2022).
- [6] W. T. Gowers and T. Karam, Low-complexity approximations for sets defined by generalizations of affine conditions, arXiv:2306.00747 (2023).
- [7] B. Green, 100 open problems, manuscript.
- [8] B. Green and T. Tao, The distribution of polynomials over finite fields, with applications to the Gowers norms. Contributions to Discrete Mathematics, 4 (2009), no. 2, 1-36.
- [9] Jan Hazla, Thomas Holenstein, Elchanan Mossel, Product space models of correlation: Between noise stability and additive combinatorics, Discrete Anal. **19** (2018).
- [10] T. Karam, On the expressive power of mod-p linear forms on the Boolean cube, arXiv:2309.14229 (2023).
- [11] T. Karam, Ranges of polynomials control degree ranks of Green and Tao over finite prime fields, arXiv:2305.11088 (2023).