

THE WEDDERBURN-ARTIN THEOREM

MATEJ BREŠAR

ABSTRACT. The celebrated Wedderburn-Artin theorem states that a simple left artinian ring is isomorphic to the ring of matrices over a division ring. We give a short and self-contained proof which avoids the use of modules.

1. INTRODUCTION

The following theorem can be considered the fundamental theorem of noncommutative algebra.

Theorem 1.1. (Wedderburn-Artin Theorem) *If R is a simple left artinian ring, then there exist a division ring D and a positive integer n such that $R \cong M_n(D)$.*

Joseph Wedderburn proved Theorem 1.1 for finite-dimensional algebras in 1908 [4]. Some twenty years later, Emil Artin generalized Wedderburn's seminal result to rings satisfying chain conditions [1].

We will present a proof that requires only basic knowledge of undergraduate algebra and does not involve the concept of a module. The standard module-theoretic approach is certainly efficient and yields further insight into the structure theory of rings, but requires more prerequisites and may be a bit difficult to understand for a beginner.

This paper is an adaptation and revision of the author's earlier paper [2] in which a proof of Wedderburn's classical version of Theorem 1.1 is given. This proof is also presented in the book [3]. The proof in the present paper is based on the same idea, but is more transparent and covers the general case.

In Section 2, we give all necessary definitions, provide basic examples, and prove several lemmas. The proof of Theorem 1.1, in fact of a slightly more general version of this theorem, is given in Section 3.

2. PRELIMINARIES

We assume the reader is familiar with the concept of a (not necessarily commutative) ring. Our rings will be assumed to be unital, i.e., they contain an element 1, called a unity, that satisfies $x = 1x = x1$ for every ring element x . We remark that, like in [2], the existence of 1 could be avoided, but at the cost of simplicity of proofs.

To refresh the reader's memory, we start by recalling some basic definitions and facts.

An element y is a left inverse of the element x if $yx = 1$. A right inverse z is defined analogously. If x has both a left inverse y and a right inverse z , then $y = y(xz) = (yx)z =$

2020 *Mathematics Subject Classification.* 16K40, 16N60.

Key words and phrases. Wedderburn-Artin theorem, division ring, simple ring, prime ring, left artinian ring, minimal left ideal, idempotent, matrix unit.

Partially supported by ARIS Grant P1-0288.

z . A left inverse of x thus coincides with a right inverse of x . We call it the inverse of x and denote it by x^{-1} . An element having an inverse is said to be invertible.

A ring is nonzero if 0 is not its only element (equivalently, $0 \neq 1$). A nonzero ring in which every nonzero element is invertible is called a division ring. Such a ring may not be commutative. For example, the ring of quaternions \mathbb{H} , with which the reader is presumably familiar, is an example of a noncommutative division ring. A commutative division ring is called a field. Standard examples are the fields of rational numbers \mathbb{Q} , real numbers \mathbb{R} , and complex numbers \mathbb{C} .

A (ring) isomorphism is bijective map φ from a ring R to a ring R' that satisfies $\varphi(a+b) = \varphi(a) + \varphi(b)$ and $\varphi(ab) = \varphi(a)\varphi(b)$ for all $a, b \in R$. We say that R and R' are isomorphic if there exists an isomorphism from R to R' . In this case, we write $R \cong R'$.

An additive subgroup I of a ring R is called a left ideal (resp. right ideal) if $ux \in I$ (resp. $xu \in I$) for all $u \in I$ and $x \in R$. If I is both a left and a right ideal, then it is called an ideal. In commutative rings, left and right ideals are automatically ideals.

We introduce some notation. For a subset S of a ring R and elements $a, b \in R$, write $Sa = \{sa \mid s \in S\}$, $aS = \{as \mid s \in S\}$, and $aSb = \{asb \mid s \in S\}$. By RS we denote the set of all sums of elements of the form xs with $x \in R$ and $s \in S$. Note that RS is the left ideal of R generated by S (i.e., the smallest left ideal containing S). Taking $S = aR$, we obtain RaR which is the ideal of R generated by a . If I and J are left ideals of R , then IJ denotes the set of all sums of elements uv with $u \in I$ and $v \in J$. Observe that IJ is again a left ideal (and is an ideal if I and J are ideals). If $I = J$, we write I^2 for IJ .

In the next subsections, we introduce several notions that are not always treated in basic algebra courses.

2.1. Simple Rings. A nonzero ring R whose only ideals are $\{0\}$ and R is called a **simple ring**.

Examples 2.1. (a) Obvious examples of simple rings are division rings, which do not even contain left (or right) ideals different from $\{0\}$ and R . This is because if a left ideal I of any ring R contains an invertible element a , then $I = R$. Indeed, $x = (xa^{-1})a \in I$ for every $x \in R$.

(b) Let D be a ring and $n \in \mathbb{N}$. The set of all $n \times n$ matrices with entries in D is a ring under the usual matrix addition and multiplication. We denote it by $M_n(D)$ (for $n = 1$, this is just D). Let E_{ij} denote the matrix whose (i, j) entry is 1 and all other entries are 0. We call E_{ij} , $1 \leq i, j \leq n$, the **standard matrix units**. By aE_{ij} we denote the matrix whose (i, j) entry is $a \in D$ and other entries are 0. Observe that for every $A = (a_{ij}) \in M_n(D)$,

$$E_{ij}AE_{kl} = a_{jk}E_{il} \quad \text{for all } 1 \leq i, j, k, l \leq n.$$

Now assume that D is a division ring. Let I be a nonzero ideal of $M_n(D)$. Take a nonzero $(a_{ij}) \in I$ and choose j and k such that $a_{jk} \neq 0$. From the displayed formula we see that $a_{jk}E_{il} \in I$ for all i and l , and hence also $(da_{jk}^{-1})E_{ii} \cdot a_{jk}E_{il} = dE_{il} \in I$ for every $d \in D$. Since every matrix in $M_n(D)$ is a sum of matrices of the form dE_{il} , $I = M_n(D)$. We have thus proved that $M_n(D)$ is a simple ring for every $n \in \mathbb{N}$ and every division ring D .

Remark 2.2. If $a \neq 0$ is an element of a ring R that does not have a left inverse, then the left ideal Ra does not contain 1 and is therefore different from $\{0\}$ and R . This

readily implies that division rings are actually the only rings without proper nonzero left ideals. In particular, the ring $M_n(D)$ has proper nonzero left ideals for every $n \geq 2$.

2.2. Prime Rings. A ring R is called a **prime ring** if for all ideals I and J of R , $IJ = \{0\}$ implies $I = \{0\}$ or $J = \{0\}$. Obviously, every simple ring is prime, and so is every ring without zero-divisors (i.e., a ring in which $ab = 0$ implies $a = 0$ or $b = 0$). The ring of integers \mathbb{Z} is thus prime, but not simple since $n\mathbb{Z}$ is an ideal of \mathbb{Z} for every $n \in \mathbb{N}$.

Lemma 2.3. *Let R be a ring. The following conditions are equivalent:*

- (i) R is prime.
- (ii) For all $a, b \in R$, $aRb = \{0\}$ implies $a = 0$ or $b = 0$.
- (iii) For all left ideals I and J of R , $IJ = \{0\}$ implies $I = \{0\}$ or $J = \{0\}$.

Proof. (i) \implies (ii). Observe that $aRb = \{0\}$ yields $(RaR)(RbR) = \{0\}$, which by assumption implies $RaR = \{0\}$ and hence $a = 0$, or $RbR = \{0\}$ and hence $b = 0$.

(ii) \implies (iii). If left ideals I and J satisfy $IJ = \{0\}$, then $aRb = \{0\}$ for every $a \in I$ and every $b \in J$. Therefore, by assumption, $a = 0$ or $b = 0$.

(iii) \implies (i). Trivial. □

2.3. Left Artinian Rings. A ring R is said to be an **algebra** over a field F if R is also a vector space over F and $\lambda(xy) = (\lambda x)y = x(\lambda y)$ holds for all $\lambda \in F$ and $x, y \in R$. A finite-dimensional algebra is an algebra which is finite-dimensional as a vector space.

Example 2.4. The ring $M_n(F)$ becomes a finite-dimensional algebra if we endow it with the usual scalar multiplication $\lambda(a_{ij}) = (\lambda a_{ij})$. The standard matrix units E_{ij} form its basis, so its dimension is n^2 .

The notion of a left artinian ring is a generalization of the notion of a finite-dimensional algebra. It is defined as follows: R is a **left artinian ring** (or is said to satisfy the **descending chain condition** on left ideals) if for every descending chain $L_1 \supseteq L_2 \supseteq L_3 \supseteq \dots$ of left ideals of R there exists an $n_0 \in \mathbb{N}$ such that $L_n = L_{n_0}$ for every $n \geq n_0$.

Examples 2.5. (a) Since a left ideal of an algebra is a vector subspace (because if $\lambda \in F$ and $u \in L$ then $\lambda u = (\lambda 1)u \in L$), a finite-dimensional algebra is indeed left-artinian.

(b) Another obvious example of a left artinian ring is every division ring. Indeed, we saw above that such a ring does not even have proper nonzero left ideals. We claim that, more generally, the matrix ring $M_n(D)$ is left artinian for every $n \in \mathbb{N}$ and every division ring D . If D is commutative, this follows from (a) and Example 2.4. It turns out that essentially the same proof works if D is not commutative. We omit details, but only indicate why this is true. One defines a vector space over a division ring in exactly the same way as a vector space over a field (but we call it more precisely a left vector space), and shows that every such vector space has a basis and that all bases have the same cardinality. This enables one to define the concept of dimension in the usual way. Now, the ring $M_n(D)$ is a vector space of dimension n^2 over D , and its left ideals are automatically vector subspaces. This implies that $M_n(D)$ is left artinian. (Incidentally, we do not define algebras over noncommutative division rings, so $M_n(D)$ is a ring and a vector space over D , but not an algebra over D .)

(c) Being left artinian is a rather special property of a ring. For example, the ring \mathbb{Z} has an infinite strictly descending chain of ideals $\mathbb{Z} \supsetneq 2\mathbb{Z} \supsetneq 4\mathbb{Z} \supsetneq 8\mathbb{Z} \supsetneq \dots$

Remark 2.6. The ring $M_n(D)$ is thus simple and left artinian. The Wedderburn-Artin Theorem states that the converse of this observation is true.

Remark 2.7. A right artinian ring is defined analogously via right ideals. However, we will consider only left artinian rings. This is just a matter of choice, the treatment of right artinian rings is essentially no different. In particular, we can substitute right for left in Theorem 1.1. (A non-simple left artinian ring, however, is not always right artinian.)

2.4. Idempotents. An element e of a ring R is called an **idempotent** if $e^2 = e$. Idempotents e and f are said to be **orthogonal** if $ef = fe = 0$. In this case, $e + f$ is also an idempotent.

Examples 2.8. (a) The simplest pair of orthogonal idempotents is $0, 1$. An idempotent different from 0 and 1 is called a **nontrivial idempotent**.

(b) If e is an idempotent, then so is $1 - e$, and e and $1 - e$ are orthogonal.

(c) The standard matrix units $E_{11}, E_{22}, \dots, E_{nn}$ are pairwise orthogonal idempotents.

Lemma 2.9. *If e and $f \neq 0$ are orthogonal idempotents in a ring R , then $R(1 - e) \supsetneq R(1 - e - f)$.*

Proof. Note that $(1 - e - f)(1 - e) = 1 - e - f$, and hence

$$x(1 - e - f) = x(1 - e - f)(1 - e) \in R(1 - e)$$

for every $x \in R$. This proves that $R(1 - e) \supseteq R(1 - e - f)$.

We have $f = f(1 - e) \in R(1 - e)$, while $f = x(1 - e - f)$ with $x \in R$ implies

$$0 = f(1 - f) = x(1 - e - f)(1 - f) = x(1 - e - f) = f,$$

a contradiction. Therefore, $R(1 - e) \neq R(1 - e - f)$. \square

If e is an idempotent in a ring R , then eRe is clearly a ring with unity e .

Lemma 2.10. *Let e be an idempotent in a ring R .*

(a) *If R is left artinian, then so is eRe .*

(b) *If R is prime, then so is eRe .*

Proof. (a) Let $L_1 \supseteq L_2 \supseteq L_3 \supseteq \dots$ be a chain of left ideals of eRe . Then $RL_1 \supseteq RL_2 \supseteq RL_3 \supseteq \dots$ is a chain of left ideals of R . Since R is left artinian, there exists an $n_0 \in \mathbb{N}$ such that $RL_n = RL_{n_0}$, and hence $eRL_n = eRL_{n_0}$, for every $n \geq n_0$. However, since L_n is a left ideal of eRe , $eRL_n = L_n$ for each n . Therefore, $L_n = L_{n_0}$ for $n \geq n_0$.

(b) We verify that eRe satisfies condition (ii) of Lemma 2.3. Let $eae, ebe \in eRe$ be such that $(eae)eRe(ebe) = \{0\}$. Since R is prime, it follows that $eae = (eae)e = 0$ or $ebe = e(ebe) = 0$, as desired. \square

2.5. Minimal Left Ideals. A nonzero left ideal L of a ring R is called a **minimal left ideal** if L does not properly contain a nonzero left ideal of R . That is to say, if I is a left ideal such that $I \subseteq L$, then $I = \{0\}$ or $I = L$.

Example 2.11. Let $R = M_n(D)$ with D a division ring, and let L be the set of matrices in R that have arbitrary entries in the i th column and zeros in all other columns. It is an easy exercise to show that L is a minimal left ideal of R . Observe that $L = RE_{ii}$ where E_{ii} is the standard matrix unit, and $E_{ii}RE_{ii} = \{dE_{ii} \mid d \in D\}$, so $E_{ii}RE_{ii} \cong D$.

Lemma 2.12. *Let L be a minimal left ideal of a ring R . If $L^2 \neq \{0\}$, then there exists an idempotent $e \in L$ such that $L = Re$ and eRe is a division ring.*

Proof. By assumption, there exists a $y \in L$ such that $Ly \neq \{0\}$. As Ly is a left ideal of R contained in L , $Ly = L$ since L is minimal. Therefore, there exists an $e \in L$ such that $ey = y$. Hence, $e^2y = ey$, meaning that $e^2 - e$ lies in the set $J = \{z \in L \mid zy = 0\}$. Observe that J is a left ideal of R contained in L . Since $Ly \neq \{0\}$ and so $J \neq L$, it follows that $J = \{0\}$. In particular, $e^2 = e$. We have $Re \subseteq L$ since $e \in L$, and since $0 \neq e \in Re$ it follows from the minimality assumption that $L = Re$.

Now consider the ring eRe . Let $a \in R$ be such that $ae \neq 0$. We must prove that ae is invertible in eRe . We have $\{0\} \neq Reae \subseteq Re = L$, and so $Reae = L$. Hence, $beae = e$ for some $b \in R$, which gives $(ebe)(eae) = e$. Since ebe is a nonzero element in eRe , by the same argument there exists a $c \in R$ such that $(ece)(ebe) = e$. But a left inverse coincides with a right inverse, so $ae = ece$ is invertible in eRe , with inverse ebe . \square

Example 2.13. Not every ring has a minimal left ideal. For example, the nonzero ideals of the ring \mathbb{Z} are $n\mathbb{Z}$, $n \in \mathbb{N}$, and none of them is minimal since, say, $2n\mathbb{Z} \subsetneq n\mathbb{Z}$.

Lemma 2.14. *A nonzero left artinian ring R has a minimal left ideal.*

Proof. Choose a nonzero left ideal L_1 of R (e.g., $L_1 = R$). If L_1 is not minimal, there exists a nonzero left ideal L_2 such that $L_1 \supsetneq L_2$. If L_2 is not minimal, then $L_2 \supsetneq L_3$ for some nonzero left ideal L_3 . Continuing this process we arrive in a finite number of steps at a minimal left ideal; for if not, there would exist an infinite chain $L_1 \supsetneq L_2 \supsetneq L_3 \supsetneq \dots$ \square

2.6. Matrix Units. Above, we defined the standard matrix units E_{ij} . Now we will consider their abstract generalization.

Let R be a ring and let $n \in \mathbb{N}$. We call $\{e_{ij} \in R \mid 1 \leq i, j \leq n\}$ a **set of $n \times n$ matrix units** if

$$e_{11} + e_{22} + \dots + e_{nn} = 1 \quad \text{and} \quad e_{ij}e_{kl} = \delta_{jk}e_{il}$$

for all $1 \leq i, j, k, l \leq n$. Here, δ_{jk} stands for the Kronecker delta.

Example 2.15. By taking the identity matrix for A in the displayed formula in Example 2.1 (b) we see that the standard matrix units E_{ij} are indeed matrix units of $M_n(D)$. There are others. For example, if $S \in M_n(D)$ is invertible, then $SE_{ij}S^{-1}$ are also matrix units.

Remark 2.16. If e_{ij} are matrix units, then each ring $e_{ii}Re_{ii}$ is isomorphic to $e_{11}Re_{11}$. Indeed, it is an easy exercise to verify that

$$e_{ii}ae_{ii} \mapsto e_{1i}(e_{ii}ae_{ii})e_{i1} = e_{11}(e_{1i}ae_{i1})e_{11}$$

is an isomorphism. Although not directly used in our proof of the Wedderburn-Artin Theorem, this helps in understanding it better. In particular, it explains why the next lemma involves $e_{11}Re_{11}$.

Lemma 2.17. *If a ring R contains a set of $n \times n$ matrix units e_{ij} , then $R \cong M_n(e_{11}Re_{11})$.*

Proof. For every $a \in R$, write $a_{ij} = e_{1i}ae_{j1}$. Observe that $a_{ij} = e_{11}a_{ij}e_{11} \in e_{11}Re_{11}$. Define $\varphi : R \rightarrow M_n(e_{11}Re_{11})$ by

$$\varphi(a) = (a_{ij}).$$

Our goal is to show that φ is an isomorphism.

It is clear that $\varphi(a + b) = \varphi(a) + \varphi(b)$ for all $a, b \in R$. The (i, j) entry of $\varphi(a)\varphi(b)$ is equal to

$$\sum_{k=1}^n e_{1i} a e_{k1} e_{1k} b e_{j1} = e_{1i} a \left(\sum_{k=1}^n e_{kk} \right) b e_{j1} = e_{1i} a b e_{j1},$$

which is the (i, j) entry of $\varphi(ab)$. Therefore, $\varphi(ab) = \varphi(a)\varphi(b)$. If $a_{ij} = 0$ for all i, j , then $e_{ii} a e_{jj} = e_{i1} a_{ij} e_{1j} = 0$, and so $a = 0$ since the sum of all e_{ii} is 1. Thus, φ is injective. Finally, observe that $\varphi(e_{k1} a e_{1l})$ is the matrix whose (k, l) entry is $e_{1l} a e_{11}$ and all other entries are 0. Since every matrix in $M_n(e_{11} R e_{11})$ is a sum of such matrices, φ is surjective. \square

Lemma 2.17 tells us that in order to show that a ring R is a matrix ring, it is enough to find matrix units in R . The next lemma simplifies this task.

Lemma 2.18. *If a ring R contains pairwise orthogonal idempotents e_{ii} , $i = 1, \dots, n$, whose sum is 1, and elements $e_{1i} \in e_{11} R e_{ii}$ and $e_{i1} \in e_{ii} R e_{11}$, $i = 2, \dots, n$, satisfying $e_{1i} e_{i1} = e_{11}$ and $e_{i1} e_{1i} = e_{ii}$, then the set $\{e_{ii}, e_{1i}, e_{i1} \mid i = 1, \dots, n\}$ can be extended to a set of $n \times n$ matrix units. Accordingly, $R \cong M_n(e_{11} R e_{11})$.*

Proof. For $i \neq j$ and $i, j \neq 1$, define $e_{ij} = e_{i1} e_{1j}$. Observe that our assumptions imply that $e_{ij} = e_{i1} e_{1j}$ holds in any case, that is, also if $i = j$ or one of i, j is 1. Note also that $e_{1j} e_{k1} = \delta_{jk} e_{11}$ holds for all j and k . Consequently, for all i, j, k, l we have

$$e_{ij} e_{kl} = e_{i1} e_{1j} e_{k1} e_{l1} = \delta_{jk} e_{i1} e_{11} e_{l1} = \delta_{jk} e_{i1} e_{l1} = \delta_{jk} e_{il},$$

proving that e_{ij} are matrix units. Therefore, $R \cong M_n(e_{11} R e_{11})$ by Lemma 2.17. \square

We need two more lemmas.

Lemma 2.19. *Let e, f be a pair of orthogonal idempotents in a prime ring R such that eRe and fRf are division rings. Then:*

- (a) *There exist $u \in eRf$ and $v \in fRe$ such that $uv = e$ and $vu = f$.*
- (b) *$(e + f)R(e + f) \cong M_2(eRe)$.*
- (c) *$eRe \cong fRf$.*

Proof. (a) Using condition (ii) of Lemma 2.3 twice we infer that $ea f b e \neq 0$ for some $a, b \in R$. As eRe is a division ring with unity e , there is a $c \in R$ such that $(ea f b e)(ece) = e$. Thus, $u = eaf \in eRf$ and $v = fbece \in fRe$ satisfy $uv = e$. Suppose $vu \neq f$. Then $vu - f$ is a nonzero, and hence invertible element of the division ring fRf . Observe that $vuv = ve = v$ and therefore $(vu - f)v = 0$. Multiplying this equation from the left by the inverse of $vu - f$, we arrive at a contradiction $v = fv = 0$. Thus, $vu = f$.

(b) Observe that $e_{11} = e$, $e_{12} = u$, $e_{21} = v$, $e_{22} = f$ form a set of 2×2 matrix units of $(e + f)R(e + f)$, so we can apply Lemma 2.17.

(c) Use Remark 2.16. \square

We actually only need the assertion (a). However, (b) and (c) give a more clear picture. We also remark that we can replace the assumption that fRf is a division ring by a milder assumption that $f \neq 0$ and fRf has no nontrivial idempotents. This is because $uv = e$ with $u \in eRf$ and $v \in fRe$ implies $(vu)^2 = v(uv)u = veu = vu$.

Lemma 2.20. *If a prime ring R contains pairwise orthogonal idempotents e_1, \dots, e_n such that their sum is 1 and $e_i Re_i$ is a division ring for each i , then $R \cong M_n(e_1 Re_1)$.*

Proof. Write e_{ii} for e_i . Applying Lemma 2.19 (a) to the idempotents e_{11} and e_{ii} , $i \geq 2$, we see that there exist elements $e_{1i} \in e_{11} Re_{ii}$ and $e_{i1} \in e_{ii} Re_{11}$ satisfying $e_{1i} e_{i1} = e_{11}$ and $e_{i1} e_{1i} = e_{ii}$. Therefore, $R \cong M_n(e_{11} Re_{11})$ by Lemma 2.18. \square

3. PROOF OF THE WEDDERBURN-ARTIN THEOREM

The standard, classical version of the Wedderburn-Artin Theorem was stated in Introduction. We will prove a more general version in which the assumption that R is simple is replaced by the assumption that R is prime. The main reason for this is that prime rings are more suitable for our proof than simple rings. On the other hand, it is interesting in its own right that nonzero prime left artinian rings are automatically simple.

Theorem 3.1. *If R is a nonzero prime left artinian ring, then there exist a division ring D and a positive integer n such that $R \cong M_n(D)$.*

Proof. Since R is left artinian, R contains a minimal left ideal L (by Lemma 2.14), and since R is prime, $L^2 \neq \{0\}$ (by Lemma 2.3). Lemma 2.12 therefore implies that R contains an idempotent e_1 such that $e_1 Re_1$ is a division ring. By Lemma 2.9 (applied to $e = 0$), $R \supsetneq R(1 - e_1)$. If $e_1 \neq 1$, then $(1 - e_1)R(1 - e_1)$ is a nonzero ring, which is also prime and left artinian by Lemma 2.10. We can thus repeat the argument from the beginning of the proof to conclude that $(1 - e_1)R(1 - e_1)$ contains an idempotent e_2 such that $e_2(1 - e_1)R(1 - e_1)e_2$ is a division ring. Observe that $e_2 \in (1 - e_1)R(1 - e_1)$ implies that e_1 and e_2 are orthogonal. Therefore, $e_2(1 - e_1)R(1 - e_1)e_2 = e_2 Re_2$ and $R(1 - e_1) \supsetneq R(1 - e_1 - e_2)$ by Lemma 2.9. If $e_1 + e_2 \neq 1$, the same argument shows that there exists an idempotent e_3 belonging to $(1 - e_1 - e_2)R(1 - e_1 - e_2)$, and hence orthogonal to e_1 and e_2 , such that $e_3(1 - e_1 - e_2)R(1 - e_1 - e_2)e_3 = e_3 Re_3$ is a division ring and $R(1 - e_1 - e_2) \supsetneq R(1 - e_1 - e_2 - e_3)$. We continue this process, which must stop after finitely many steps since R is left artinian. Therefore, there exist pairwise orthogonal idempotents $e_1, \dots, e_n \in R$ such that $e_i Re_i$ is a division ring and $e_1 + \dots + e_n = 1$. Lemma 2.20 now gives the desired conclusion that $R \cong M_n(e_1 Re_1)$. \square

REFERENCES

- [1] E. Artin, Zur Theorie der hyperkomplexen Zahlen, *Abh. Math. Sem. Univ. Hamburg* **5** (1927), 251–260.
- [2] M. Brešar, An elementary approach to Wedderburn’s structure theory, *Expo. Math.* **28** (2010), 79–83.
- [3] M. Brešar, *Introduction to Noncommutative Algebra*, Universitext, Springer, 2014.
- [4] J. H. M. Wedderburn, On hypercomplex numbers, *Proc. London Math. Soc.* **2-6** (1908), 77–118.

FACULTY OF MATHEMATICS AND PHYSICS, UNIVERSITY OF LJUBLJANA & FACULTY OF NATURAL SCIENCES AND MATHEMATICS, UNIVERSITY OF MARIBOR & IMFM, LJUBLJANA, SLOVENIA

Email address: `matej.bresar@fmf.uni-lj.si`