

UNIFORM HANSON-WRIGHT TYPE DEVIATION INEQUALITIES FOR α -SUBEXPONENTIAL RANDOM VECTORS

GUOZHENG DAI AND ZHONGGEN SU

ABSTRACT. This paper is devoted to uniform versions of the Hanson-Wright inequality for a random vector with independent centered α -subexponential entries, $0 < \alpha \leq 1$. Our method relies upon a novel decoupling inequality and a comparison of weak and strong moments. As an application, we use the derived inequality to prove the restricted isometry property of partial random circulant matrices generated by standard α -subexponential random vectors, $0 < \alpha \leq 1$.

1. INTRODUCTION AND MAIN RESULT

Let $A = (a_{ij})_{n \times n}$ be an $n \times n$ fixed symmetric matrix and $S_A(\xi) := \sum_{i,j} a_{ij} \xi_i \xi_j$, where $\xi = (\xi_1, \dots, \xi_n)$ is a random vector with independent centered entries. A well-known concentration property of $S_A(\xi)$ is due to Hanson and Wright, claiming that if ξ_i are independent centered subgaussian variables with $\|\xi_i\|_{\Psi_2} \leq L$, then for all $t \geq 0$ (a modern version in [12])

$$\mathbb{P}\{|S_A(\xi) - \mathbb{E}S_A(\xi)| \geq t\} \leq 2 \exp\left(-c \min\left\{\frac{t^2}{L^4 \|A\|_F^2}, \frac{t}{L^2 \|A\|_{l_2 \rightarrow l_2}}\right\}\right).$$

Here, $\|\cdot\|_F$ and $\|\cdot\|_{l_2 \rightarrow l_2}$ are the Frobenius norm and the spectral norm of a matrix (see (2.1) below), respectively. The Hanson-Wright type inequalities have found numerous applications in non-asymptotic theory of random matrices, statistics, and so on [1, 2, 6, 8, 15].

An important extension of such results is to consider the behaviour of random quadratic forms simultaneously for a family of $n \times n$ fixed symmetric matrices \mathcal{A} , that is the concentration property of $Z_{\mathcal{A}}(\xi) := \sup_{A \in \mathcal{A}} \sum_{i,j} a_{ij} \xi_i \xi_j$. When ξ_i are independent Rademacher variables (take values ± 1 with equal probabilities) and the diagonal entries of A are 0, Talagrand's celebrated work [13] yields for $t \geq 0$

$$(1.1) \quad \mathbb{P}\{|Z_{\mathcal{A}}(\xi) - \mathbb{E}Z_{\mathcal{A}}(\xi)| \geq t\} \leq 2 \exp\left(-c \min\left\{\frac{t^2}{(\mathbb{E} \sup_{A \in \mathcal{A}} \|A\xi\|_2)^2}, \frac{t}{\sup_{A \in \mathcal{A}} \|A\|_{l_2 \rightarrow l_2}}\right\}\right).$$

Recently, Klochov and Zhivotovskiy [5] obtained a uniform Hanson-Wright type inequality for more general random variables via the entropy method. In particular, let ξ_i be independent centered subgaussian variables and $L' = \|\max_i |\xi_i|\|_{\Psi_2}$, then

Key words and phrases. α -subexponential random variable, chaining argument, uniform Hanson-Wright inequality, restricted isometry property.

(without the condition $a_{ii} = 0$)

(1.2)

$$\mathbb{P}\{Z_{\mathcal{A}}(\xi) - \mathbb{E}Z_{\mathcal{A}}(\xi) \geq t\} \leq \left(-c \min \left\{ \frac{t^2}{L'^2(\mathbb{E} \sup_{A \in \mathcal{A}} \|A\xi\|_2)^2}, \frac{t}{L'^2 \sup_{A \in \mathcal{A}} \|A\|_{l_2 \rightarrow l_2}} \right\} \right),$$

where $t \geq \max\{L'\mathbb{E} \sup_{A \in \mathcal{A}} \|A\xi\|, L'^2 \sup_{A \in \mathcal{A}} \|A\|_{l_2 \rightarrow l_2}\}$.

We usually call the results of the form (1.1) and (1.2) (one-sided) concentration inequalities. Another similar bounds are one-sided and have a multiplicative constant before $\mathbb{E}Z_{\mathcal{A}}(\xi)$ (or replace $\mathbb{E}Z_{\mathcal{A}}(\xi)$ with a suitable upper bound). This type of bound is called a deviation inequality. We next introduce some known uniform Hanson-Wright type deviation inequalities.

We first introduce some notations. Denote $(\mathbb{E}|\xi_1|^p)^{1/p}$ by $\|\xi_1\|_{L_p}$ for a random variable ξ_1 . Set $\alpha^* = \alpha/(\alpha - 1)$ as the conjugate exponent of α . Define $M_F(\mathcal{A}) := \sup_{A \in \mathcal{A}} \|A\|_F$ and $M_{l_{p_1} \rightarrow l_{p_2}}(\mathcal{A}) := \sup_{A \in \mathcal{A}} \|A\|_{l_{p_1} \rightarrow l_{p_2}}$ (see (2.1) below for the definition of the norm). We also denote Talagrand's γ_α -functional with respect to $(\mathcal{A}, \|\cdot\|_{l_2 \rightarrow l_{\alpha^*}})$ by $\gamma_\alpha(\mathcal{A}, \|\cdot\|_{l_2 \rightarrow l_{\alpha^*}})$ (see (2.2) below). Let

$$\Gamma(\alpha, \beta, \mathcal{A}) = \gamma_2(\mathcal{A}, \|\cdot\|_{l_2 \rightarrow l_2}) + \gamma_\alpha(\mathcal{A}, \|\cdot\|_{l_2 \rightarrow l_\beta})$$

and

$$\begin{aligned} U_1(\alpha, \beta) &= \Gamma(\alpha, \beta, \mathcal{A})(\Gamma(\alpha, \beta, \mathcal{A}) + M_F(\mathcal{A})), \\ U_2(\alpha, \beta) &= M_{l_2 \rightarrow l_2}(\mathcal{A})\Gamma(\alpha, \beta, \mathcal{A}) + \sup_{A \in \mathcal{A}} \|A^\top A\|_F, \\ U_3(\alpha, \beta) &= M_{l_2 \rightarrow l_\beta}(\mathcal{A})\Gamma(\alpha, \beta, \mathcal{A}). \end{aligned}$$

A widely used deviation inequality in the field of compressive sensing [1, 6, 8] was obtained by Krahmer et al [8]. Let ξ_i be independent centered subgaussian variables and $L = \max_i \|\xi_i\|_{\Psi_2}$. They proved for $t \geq 0$

(1.3)

$$\mathbb{P}\left\{ \sup_{A \in \mathcal{A}} \left| \|A\xi\|_2^2 - \mathbb{E}\|A\xi\|_2^2 \right| \geq CL^2(U_1(2, 2) + t) \right\} \leq 2 \exp\left(- \min\left\{ \frac{t^2}{U_2^2(2, 2)}, \frac{t}{M_{l_2 \rightarrow l_2}^2(\mathcal{A})} \right\} \right),$$

where \mathcal{A} is a family of $m \times n$ fixed matrices.

Dai et al [3] extended (1.3) to the case where ξ_i are independent centered α -subexponential variables, $1 \leq \alpha \leq 2$. Recall that a random variable ξ_1 is α -subexponential (or α -sub-Weibull) if satisfying

$$\mathbb{P}\{|\xi_1| \geq Kt\} \leq ce^{-ct^\alpha}, \quad t \geq 0,$$

where K is a parameter and c is a universal constant. The α -subexponential (quasi-)norm of ξ_1 is defined as follows:

$$\|\xi_1\|_{\Psi_\alpha} := \inf\{t > 0 : \mathbb{E} \exp\left(\frac{|\xi_1|^\alpha}{t^\alpha}\right) \leq 2\}.$$

There are many interesting choices of random variables of this type, such as subexponential variables ($\alpha = 1$), subgaussian variables ($\alpha = 2$), or bounded variables (for any $\alpha > 0$).

Let ξ_i be independent centered α -subexponential variables and $L = L(\alpha) = \max_i \|\xi_i\|_{\Psi_\alpha}$. Dai et al (see Corollary 1.2* in [3]) proved for $t \geq 0$

$$\mathbb{P}\left\{ \sup_{A \in \mathcal{A}} \left| \|A\xi\|_2^2 - \mathbb{E}\|A\xi\|_2^2 \right| > C(\alpha)L^2(U_1(\alpha, \alpha^*) + t) \right\}$$

$$(1.4) \quad \leq C_1(\alpha) \exp \left(- \min \left\{ \left(\frac{t}{U_2(\alpha, \alpha^*)} \right)^2, \left(\frac{t}{U_3(\alpha, \alpha^*)} \right)^\alpha, \left(\frac{t}{M_{l_2 \rightarrow l_2}^2(\mathcal{A})} \right)^{\alpha/2} \right\} \right),$$

where \mathcal{A} is a family of $m \times n$ fixed matrices.

Our main result shows a uniform Hanson-Wright type deviation inequality for a random vector with independent centered α -subexponential variables ($0 < \alpha \leq 1$), reading as follows:

Theorem 1.1. *Let $\xi = (\xi_1, \dots, \xi_n)$ be a random vector with independent centered α -subexponential entries ($0 < \alpha \leq 1$) and \mathcal{A} be a family of fixed $m \times n$ matrices. Then, we have for $t \geq 0$*

$$\begin{aligned} & P \left\{ \sup_{A \in \mathcal{A}} \left| \|A\xi\|_2^2 - E\|A\xi\|_2^2 \right| > C(\alpha)L^2(U_1(\alpha, \infty) + t) \right\} \\ & \leq C_1(\alpha) \exp \left(- \min \left\{ \left(\frac{t}{U_2(\alpha, \infty)} \right)^2, \left(\frac{t}{U_3(\alpha, \infty)} \right)^\alpha, \left(\frac{t}{M_{l_2 \rightarrow l_2}^2(\mathcal{A})} \right)^{\alpha/2} \right\} \right), \end{aligned}$$

where $L = L(\alpha) = \max_i \|\xi_i\|_{\Psi_\alpha}$.

Remark 1.1. (i) In the setting of Theorem 1.1, Götze et al showed for $t \geq 0$ (see Proposition 1.1 in [4])

$$P \left\{ \left| \|A\xi\|_2^2 - E\|A\xi\|_2^2 \right| > C(\alpha)L^2t \right\} \leq C_1(\alpha) \exp \left(- \min \left\{ \left(\frac{t}{\|A^\top A\|_F} \right)^2, \left(\frac{t}{\|A^\top A\|_{l_2 \rightarrow l_2}} \right)^{\alpha/2} \right\} \right).$$

Theorem 1.1 recovers this result when \mathcal{A} contains only one matrix.

(ii) When $\alpha = 1$, Theorem 1.1 yields the same result as (1.4).

In the case $\alpha = 2$, the uniform Hanson-Wright deviation inequality (see (1.3)) is derived through the application of the classic decoupling inequality (see Theorem 2.5 in [8]) and the majorizing measure theorem (an optimal bound for the suprema of the stochastic process, e.g., (2.2) in [8]). To extend the deviation inequality to the case $1 \leq \alpha \leq 2$, Dai et al [3] establish a novel decoupling inequality (see Lemma 2.6 below) and then obtain the desired result via the corresponding majorizing measure theorem (see Lemma 2.7 in [3]). For the case $0 < \alpha \leq 1$, although the novel decoupling inequality is valid (see Remark 2.3 below), the majorizing measure theorem is absent, presenting a significant challenge to obtaining the deviation inequalities.

The main contribution of this paper is to show a uniform Hanson-Wright type deviation inequality for α -subexponential ($0 < \alpha \leq 1$) entries. Our method of proof is based on a combination of the novel decoupling inequality with a comparison of weak and strong moments (see Lemma 2.5), the latter of which plays the same role as the majorizing measure theorem.

The remainder of this paper is structured as follows: Section 2 will present some notations and auxiliary lemmas. In Section 3, we shall prove our main result, Theorem 1.1. Finally, we will give an application of our main result in Section 4. In particular, we show the R.I.P. for partial random circulant matrices generated by standard α -subexponential ($0 < \alpha \leq 1$) random vectors.

2. PRELIMINARIES

2.1. Notations. For a fixed vector $x = (x_1, \dots, x_n)^\top \in \mathbb{R}^n$, we denote by $\|x\|_p = (\sum |x_i|^p)^{1/p}$ the l_p norm. Set $S_p^{n-1} = \{x \in \mathbb{R}^n : \|x\|_p = 1\}$ and $B_p^n = \{x \in \mathbb{R}^n :$

$\|x\|_p \leq 1\}$. We use $\|\xi\|_{L_p} = (\mathbb{E}|\xi|^p)^{1/p}$ for the L_p norm of a random variable ξ . As for an $m \times n$ matrix $A = (a_{ij})$, recall the following notations of norms

$$\begin{aligned} \|A\|_F &= \sqrt{\sum_{i,j} |a_{ij}|^2}, \quad \|A\|_\infty = \max_{i,j} |a_{ij}|, \quad \|A\|_{l_p(l_2)} = (\sum_{i \leq m} (\sum_{j \leq n} |a_{ij}|^2)^{p/2})^{1/p} \\ (2.1) \quad \|A\|_{l_{p_1} \rightarrow l_{p_2}} &= \sup\{|\sum a_{ij} x_j y_i| : \|x\|_{p_1} \leq 1, \|y\|_{p_2^*} \leq 1\}, \end{aligned}$$

where $p_2^* = p_2/(p_2 - 1)$. We remark that $\|A\|_{l_2 \rightarrow l_2}$ is the important spectral norm of A and $\|A\|_{l_1 \rightarrow l_\infty} = \|A\|_\infty$.

Unless otherwise stated, we denote by C, C_1, c, c_1, \dots universal constants, and by $C(\delta), c(\delta), \dots$ constants that depend only on the parameter δ . For convenience, we write $f \lesssim g$ if $f \leq Cg$ for some universal constant C and write $f \lesssim_\delta g$ if $f \leq C(\delta)g$ for some constant $C(\delta)$. We also say $f \asymp g$ if $f \lesssim g$ and $g \lesssim f$, so does $f \asymp_\delta g$.

We say ξ is a symmetric Weibull variable with the scale parameter 1 and the shape parameter α if $-\log \mathbb{P}\{|\xi| > x\} = x^\alpha, x \geq 0$. For convenience, we shall write $\xi \sim \mathcal{W}_s(\alpha)$. Given a nonempty set $T \subset l_2 := \{t : \sum_i t_i^2 < \infty\}$, we call $\{S_t : t \in T\}$ a canonical process, where $S_t = \sum_{i=1}^\infty t_i \xi_i$ and $\{\xi_i\}$ are independent random variables.

For a metric space (T, d) , a sequence of subsets $\{T_r : r \geq 0\}$ of T is admissible if for every $r \geq 1, |T_r| \leq 2^{2^r}$ and $|T_0| = 1$. For any $0 < \alpha < \infty$, the definition of Talagrand's γ_α -functional of (T, d) is as follows:

$$(2.2) \quad \gamma_\alpha(T, d) = \inf \sup_{t \in T} \sum_{r \geq 0} 2^{r/\alpha} d(t, T_r),$$

where the infimum is taken concerning all admissible sequences of T .

Given a set T , a sequence of partitions $(\mathcal{T}_n)_{n \geq 0}$ of T is admissible if satisfies $|\mathcal{T}_0| = 1, |\mathcal{T}_n| \leq 2^{2^n}$ for $n \geq 1$ and every set of \mathcal{T}_{n+1} is contained in a set of \mathcal{T}_n (increasing partitions). Denote the unique element of \mathcal{T}_n which contains t by $T_n(t)$ and the diameter of the set T by $\Delta_d(T)$. Let

$$\gamma'_\alpha(T, d) = \inf_{\mathcal{T}} \sup_{t \in T} \sum_{n \geq 0} 2^{n/\alpha} \Delta_d(T_n(t)),$$

where the infimum is taken over all admissible partitions \mathcal{T} of T . Talagrand [14] showed $\gamma_\alpha(T, d) \leq \gamma'_\alpha(T, d) \leq C(\alpha) \gamma_\alpha(T, d)$.

2.2. Tails and Moments. In this subsection, we shall present some properties about the tails and the moments of random variables.

Lemma 2.1 (Lemma 4.6 in [10]). *Let ξ be a random variable such that*

$$\theta_1 p^{1/\alpha} \leq \|\xi\|_{L_p} \leq \theta_2 p^{1/\alpha} \quad \text{for all } p \geq 2.$$

Then there exist constants c_1, c_2 depending only on θ_1, θ_2 and α such that

$$c_1 e^{-t^\alpha/c_1} \leq \mathbb{P}\{|\xi| \geq t\} \leq c_2 e^{-t^\alpha/c_2} \quad \text{for all } t \geq 0.$$

Lemma 2.2 (Lemma 2.1 in [3]). *Assume that a random variable ξ satisfies for $p \geq p_0$*

$$\|\xi\|_{L_p} \leq \sum_{k=1}^m C_k p^{\beta_k} + C_{m+1},$$

where $C_1, \dots, C_{m+1} > 0$ and $\beta_1, \dots, \beta_m > 0$. Then we have for any $t > 0$,

$$P\{|\xi| > e(mt + C_{m+1})\} \leq e^{p_0} \exp\left(-\min\left\{\left(\frac{t}{C_1}\right)^{1/\beta_1}, \dots, \left(\frac{t}{C_m}\right)^{1/\beta_m}\right\}\right)$$

and

$$P\left\{|\xi| > e\left(\sum_{k=1}^m C_k t^{\beta_k} + C_{m+1}\right)\right\} \leq e^{p_0} e^{-t}.$$

Lemma 2.3 (Example 3 in [7]). Let $\xi_1, \dots, \xi_n \stackrel{i.i.d.}{\sim} \mathcal{W}_s(\alpha)$. Assume $0 < \alpha \leq 1$, then we have for $p \geq 2$

(i)

$$\left\| \sum_{i \leq n} a_i \xi_i \right\|_{L_p} \prec_{\alpha} p^{1/2} \|a\|_2 + p^{1/\alpha} \|a\|_{\infty},$$

where $a = (a_1, \dots, a_n)^{\top}$ is a fixed vector.

(ii)

(2.3)

$$\left\| \sum_{i,j} a_{ij} \xi_i \tilde{\xi}_j \right\|_{L_p} \prec_{\alpha} p^{1/2} \|A\|_F + p \|A\|_{l_2 \rightarrow l_2} + p^{(\alpha+2)/2\alpha} \|A\|_{l_2 \rightarrow l_{\infty}} + p^{2/\alpha} \|A\|_{\infty},$$

where $A = (a_{ij})$ is a fixed symmetric matrix and $\{\tilde{\xi}_i, i \leq n\}$ are independent copies of $\{\xi_i, i \leq n\}$.

Remark 2.1. Observe that

$$\|A\|_{\infty} \leq \|A\|_{l_2 \rightarrow l_{\infty}} \leq \|A\|_{l_2 \rightarrow l_2}.$$

Hence, (2.3) yields the following simplified bound

$$\left\| \sum_{i,j} a_{ij} \xi_i \tilde{\xi}_j \right\|_{L_p} \lesssim_{\alpha} p^{1/2} \|A\|_F + p^{2/\alpha} \|A\|_{l_2 \rightarrow l_2}.$$

2.3. Contraction Principle. In this subsection, we shall present a well-known contraction principle.

Lemma 2.4 (Lemma 4.6 in [11]). Let $F : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ be a convex function. Let further $\{\eta_i, i \leq n\}$ and $\{\xi_i, i \leq n\}$ be two symmetric sequences of independent random variables such that for some constant $K \geq 1$ and all $i \leq n$ and $t > 0$

$$P\{|\eta_i| > t\} \leq K P\{|\xi_i| > t\}.$$

Then, for any finite sequence $\{a_i, i \leq n\}$ in a Banach space,

$$EF\left(\left\| \sum_{i=1}^n \eta_i a_i \right\|\right) \leq EF\left(K \left\| \sum_{i=1}^n \xi_i a_i \right\|\right).$$

Corollary 2.1. Let T be a nonempty subset of \mathbb{R}^n . $\{\eta_i\}$ and $\{\xi_i\}$ are independent random variables as in Lemma 2.4. Then, we have for $p \geq 1$

$$E \sup_{t \in T} \left| \sum_{i=1}^n \eta_i t_i \right|^p \leq K^p E \sup_{t \in T} \left| \sum_{i=1}^n \xi_i t_i \right|^p.$$

Proof. We first assume that $\text{span}(T) = \mathbb{R}^n$. Then, define the following norm on \mathbb{R}^n :

$$\|x\| = \sup_{t \in T} \left| \sum_{i=1}^n x_i t_i \right|, \quad x \in \mathbb{R}^n.$$

Let $e_i, i \geq 1$ be vectors in \mathbb{R}^n such that the i -th coordinate is 1 and the other coordinates are 0. Then Lemma 2.4 yields that

$$\mathbb{E}F\left(\left\|\sum_{i=1}^n \eta_i e_i\right\|\right) \leq \mathbb{E}F\left(K\left\|\sum_{i=1}^n \xi_i e_i\right\|\right).$$

We obtain the desired result by taking $F(\cdot) = |\cdot|^p$.

As for a general T , we let $T_\delta = T \cup \delta B_2^n$. Note that, $\text{span}(T_\delta) = \mathbb{R}^n$. Hence, Corollary 2.1 is valid for T_δ . Then, go with $\delta \rightarrow 0$ and obtain the desired result. \square

2.4. Comparison of Weak and Strong Moments. Let $\{S_t = \sum t_i \xi : t \in T\}$ be a canonical process. There is a trivial lower estimate:

$$\left\| \sup_{t \in T} |S_t| \right\|_{L_p} \geq \max \left\{ \mathbb{E} \sup_{t \in T} |S_t|, \sup_{t \in T} \|S_t\|_{L_p} \right\}.$$

In some situations, the lower bound may be reversed:

$$(2.4) \quad \left\| \sup_{t \in T} |S_t| \right\|_{L_p} \lesssim \mathbb{E} \sup_{t \in T} |S_t| + \sup_{t \in T} \|S_t\|_{L_p}.$$

We call a result of the form (2.4) a comparison of weak and strong moments. In this subsection, we shall introduce a comparison of weak and strong moments (maybe the most general version currently), which was proved by Latała and Strzelecka [9].

Lemma 2.5 (Theorem 1.1 in [9]). *Let ξ_1, \dots, ξ_n be independent centered random variables with finite moments satisfying*

$$\|\xi_i\|_{L_{2p}} \leq \alpha \|\xi_i\|_{L_p}, \quad \forall p \geq 2, \quad i = 1, \dots, n,$$

where α is a finite positive constant. Then we have for any $p \geq 1$ and any non-empty set $T \subset \mathbb{R}^n$

$$\left\| \sup_{t \in T} \left| \sum_{i=1}^n t_i \xi_i \right| \right\|_{L_p} \lesssim_\alpha \mathbb{E} \sup_{t \in T} \left| \sum_{i=1}^n t_i \xi_i \right| + \sup_{t \in T} \left\| \sum_{i=1}^n t_i \xi_i \right\|_{L_p}.$$

Remark 2.2. Let $\xi_1 \sim \mathcal{W}_s(\alpha), 0 < \alpha \leq 1$. A direct integration yields for $p \geq 1$

$$\mathbb{E}|\xi_1|^p = \frac{p}{\alpha} \Gamma\left(\frac{p}{\alpha}\right),$$

where $\Gamma(\cdot)$ is the Gamma function. Hence, ξ_1 satisfies the condition in Lemma 2.5.

2.5. Decoupling Inequality. Let $S_A = \sum_{i,j \leq n} a_{ij} \xi_i \xi_j$, where ξ_i are independent centered random variables and $A = (a_{ij})$ is a fixed matrix. The decoupling technique is applied to control S_A by its decoupled version \tilde{S}_A of the form: $\tilde{S}_A = \sum_{i,j \leq n} a_{ij} \xi_i \tilde{\xi}_j$. We next introduce a decoupling inequality.

Lemma 2.6 (Proposition 3.1 in [3]). *Let F be a convex function satisfying $F(x) = F(-x), \forall x \in \mathbb{R}$. Assume $\xi = (\xi_1, \dots, \xi_n)^\top$ and $\eta = (\eta_1, \dots, \eta_n)^\top$ are random vectors with independent centered entries. Let for any $t > 0$ and $i \geq 1$, the independent random variables ξ_i, η_i satisfy for some $c \geq 1$*

$$(2.5) \quad P\{\xi_i^2 \geq t\} \leq cP\{c|\eta_i \tilde{\eta}_i| \geq t\},$$

where $\tilde{\eta}_i$ is an independent copy of η_i . Then, there exists a constant C depending only on c such that

$$E \sup_{A \in \mathcal{A}} F(\xi^\top A \xi - E \xi^\top A \xi) \leq E \sup_{A \in \mathcal{A}} F(C \eta^\top A \tilde{\eta})$$

where \mathcal{A} is a set of $n \times n$ fixed matrices.

Remark 2.3. Let ξ_i be independent centered α -subexponential variables and $\eta_1, \dots, \eta_n \stackrel{i.i.d.}{\sim} \mathcal{W}_s(\alpha)$. Note that for $i \leq n$ and $t \geq 0$

$$P\{\xi_i^2 \geq t\} \leq ce^{-ct^{\alpha/2}} \leq cP\{|\eta_i \tilde{\eta}_i| \geq (\frac{c}{2})^{2/\alpha} t\} \leq c_1 P\{c_1 |\eta_i \tilde{\eta}_i| \geq t\},$$

where $c_1 = \max\{c, (2/c)^{2/\alpha}\}$ and $\tilde{\eta}_i$ is an independent copy of η_i . Hence, ξ_i and η_i mentioned above satisfy the condition in Lemma 2.6.

3. PROOF FOR THEOREM 1.1

In this section, we shall prove our main result, a uniform Hanson-Wright type deviation inequality. For convenience of reading, we will divide the proof of the theorem into two parts. We first prove the following proposition, a p -th moment bound for the suprema of decoupled bilinear random forms.

Proof. Let $\zeta_1, \dots, \zeta_n \stackrel{i.i.d.}{\sim} \mathcal{W}_s(\alpha)$, $0 < \alpha \leq 1$. Then, we have for $p \geq 1$

$$(3.1) \quad \left\| \sup_{A \in \mathcal{A}} |\zeta^\top A^\top A \tilde{\zeta}| \right\|_{L_p} \lesssim_\alpha \sup_{A \in \mathcal{A}} \|\zeta^\top A^\top A \tilde{\zeta}\|_{L_p} + \left\| \sup_{A \in \mathcal{A}} \|A \tilde{\zeta}\|_2 \right\|_{L_p} (\gamma_2(\mathcal{A}, \|\cdot\|_{l_2 \rightarrow l_2}) + \gamma_\alpha(\mathcal{A}, \|\cdot\|_{l_2 \rightarrow l_\infty})),$$

where $\zeta = (\zeta_1, \dots, \zeta_n)^\top$ and $\tilde{\zeta}$ is an independent copy of ζ .

Proof. For convenience, denote by d_2 and d_∞ the distance on \mathcal{A} induced by $\|\cdot\|_{l_2 \rightarrow l_2}$ and $\|\cdot\|_{l_2 \rightarrow l_\infty}$. By the definition of γ'_α functionals, we can select two admissible sequences of partitions $\mathcal{T}^{(1)} = (\mathcal{T}_n^{(1)})_{n \geq 0}$ and $\mathcal{T}^{(2)} = (\mathcal{T}_n^{(2)})_{n \geq 0}$ of \mathcal{A} such that

$$\sup_{A \in \mathcal{A}} \sum_{n \geq 0} 2^{n/2} \Delta_{d_2}(\mathcal{T}_n^{(1)}(A)) \leq 2\gamma'_2(\mathcal{A}, d_2), \quad \sup_{A \in \mathcal{A}} \sum_{n \geq 0} 2^{n/\alpha} \Delta_{d_\infty}(\mathcal{T}_n^{(2)}(A)) \leq 2\gamma'_\alpha(\mathcal{A}, d_\infty).$$

Let $\mathcal{T}_0 = \{\mathcal{A}\}$ and

$$\mathcal{T}_n = \{T^{(1)} \cap T^{(2)} : T^{(1)} \in \mathcal{T}_{n-1}^{(1)}, T^{(2)} \in \mathcal{T}_{n-1}^{(2)}\}, \quad n \geq 1.$$

Then, $\mathcal{T} = (\mathcal{T}_n)_{n \geq 0}$ is a sequence of increasing partitions and

$$|\mathcal{T}_n| \leq |\mathcal{T}_{n-1}^{(1)}| |\mathcal{T}_{n-1}^{(2)}| \leq 2^{2^{n-1}} 2^{2^{n-1}} = 2^{2^n}.$$

We next define a subset \mathcal{A}_n of \mathcal{A} by selecting exactly one point from each $T \in \mathcal{T}_n$. By this means, we build an admissible sequence $(\mathcal{A}_n)_{n \geq 0}$ of subsets of \mathcal{A} . Let $\pi = \{\pi_r, r \geq 0\}$ be a sequence of functions $\pi_n : \mathcal{A} \rightarrow \mathcal{A}_n$ such that $\pi_n(A) = \mathcal{A}_n \cap T_n(A)$, where $T_n(A)$ is the element of \mathcal{T}_n containing A . Let l be the largest integer satisfying $2^l \leq p$.

We first make the following decomposition:

$$(3.2) \quad \left\| \sup_{A \in \mathcal{A}} |\zeta^\top A^\top A \tilde{\zeta}| \right\|_{L_p} \leq \left\| \sup_{A \in \mathcal{A}} |\zeta^\top A^\top A \tilde{\zeta} - \zeta^\top \pi_l(A)^\top \pi_l(A) \tilde{\zeta}| \right\|_{L_p} + \left\| \sup_{A \in \mathcal{A}} |\zeta^\top \pi_l(A)^\top \pi_l(A) \tilde{\zeta}| \right\|_{L_p}.$$

We have for any fixed $A \in \mathcal{A}$,

$$(3.3) \quad |\zeta^\top A^\top A \tilde{\zeta} - \zeta^\top \pi_l(A)^\top \pi_l(A) \tilde{\zeta}| \leq \sum_{r \geq l} |\zeta^\top \Lambda_{r+1}(A)^\top \pi_{r+1}(A) \tilde{\zeta}| + \sum_{r \geq l} |\zeta^\top \pi_r(A)^\top \Lambda_{r+1}(A) \tilde{\zeta}|,$$

where $\Lambda_{r+1}A = \pi_{r+1}(A) - \pi_r(A)$.

Conditionally on $\tilde{\zeta}$, we have by Lemma 2.3

$$\|\zeta^\top \Lambda_{r+1}(A)^\top \pi_{r+1}(A) \tilde{\zeta}\|_{L_p} \lesssim_\alpha p^{1/2} \|\Lambda_{r+1}(A)^\top \pi_{r+1}(A) \tilde{\zeta}\|_2 + p^{1/\alpha} \|\Lambda_{r+1}(A)^\top \pi_{r+1}(A) \tilde{\zeta}\|_\infty.$$

Then, we have by Lemma 2.2

$$\mathbb{P}_\zeta \left\{ |\zeta^\top S_{r+1}(A, \tilde{\zeta})| \geq C(\alpha) (\sqrt{t} \|S_{r+1}(A, \tilde{\zeta})\|_2 + t^{1/\alpha} \|S_{r+1}(A, \tilde{\zeta})\|_\infty) \right\} \leq e^2 e^{-t}.$$

Here, $S_{r+1}(A, \tilde{\zeta}) = \Lambda_{r+1}(A)^\top \pi_{r+1}(A) \tilde{\zeta}$. By the definition of the operator norms, we have

$$\|S_{r+1}(A, \tilde{\zeta})\|_2 \leq \|\Lambda_{r+1}(A)\|_{l_2 \rightarrow l_2} \sup_{A \in \mathcal{A}} \|A \tilde{\zeta}\|_2$$

and

$$\|S_{r+1}(A, \tilde{\zeta})\|_\infty \leq \|\Lambda_{r+1}(A)\|_{l_2 \rightarrow l_\infty} \sup_{A \in \mathcal{A}} \|A \tilde{\zeta}\|_2.$$

Note that $|\{\pi_r(A) : A \in \mathcal{A}\}| \leq |\mathcal{A}_r| \leq 2^{2^r}$. Then,

$$|\{\Lambda_{r+1}(A)^\top \pi_{r+1}(A) : A \in \mathcal{A}\}| \leq 2^{2^{r+3}}.$$

Denote $\Omega_{t,p}$ by the event

$$\bigcap_{r \geq l} \bigcap_{A \in \mathcal{A}} \left\{ |\zeta^\top S_{r+1}(A, \tilde{\zeta})| \leq C(\alpha) \sup_{A \in \mathcal{A}} \|A \tilde{\zeta}\|_2 (\sqrt{t} 2^{\frac{r}{2}} \|\Lambda_{r+1}(A)\|_{l_2 \rightarrow l_2} + t^{\frac{1}{\alpha}} 2^{\frac{r}{\alpha}} \|\Lambda_{r+1}(A)\|_{l_2 \rightarrow l_\infty}) \right\}.$$

For $t > 16$, we have by a union bound

$$\mathbb{P}_\zeta \{\Omega_{t,p}^c\} \leq \sum_{r \geq l} 2^{2^{r+3}} e^2 e^{-2^r t} \leq C_1(\alpha) \exp(-c(\alpha) p t).$$

Assuming the event $\Omega_{t,p}$ occurs, we have

$$\sum_{r \geq l} |\zeta^\top S_{r+1}(A, \tilde{\zeta})| \leq C(\alpha) \sup_{A \in \mathcal{A}} \|A \tilde{\zeta}\|_2 \left(\sum_{r \geq l} \sqrt{t} 2^{\frac{r}{2}} \|\Lambda_{r+1}A\|_{l_2 \rightarrow l_2} + \sum_{r \geq l} t^{\frac{1}{\alpha}} 2^{\frac{r}{\alpha}} \|\Lambda_{r+1}A\|_{l_2 \rightarrow l_\infty} \right).$$

Note that we have $\pi_{r+1}(A), \pi_r(A) \in T_r(A) \subset T_{r-1}^{(1)}(A)$ and so

$$(3.4) \quad \|\Lambda_{r+1}A\|_{l_2 \rightarrow l_2} \leq \Delta_{d_2}(T_{r-1}^{(1)}(A)).$$

Hence, we have by the definition of $\mathcal{T}^{(1)}$

$$\sum_{r \geq l} 2^{\frac{r}{2}} \|\Lambda_{r+1}A\|_{l_2 \rightarrow l_2} \leq \sum_{r \geq l} 2^{\frac{r}{2}} \Delta_{d_2}(T_{r-1}^{(1)}(A)) \leq 2\sqrt{2} \gamma'_2(\mathcal{A}, d_2).$$

Analogously, we have

$$\sum_{r \geq l} 2^{\frac{r}{\alpha}} \|\Lambda_{r+1}A\|_{l_2 \rightarrow l_{\alpha^*}} \leq 2 \cdot 2^{\frac{1}{\alpha}} \gamma'_\alpha(\mathcal{A}, d_\infty).$$

Thus,

$$\sup_{A \in \mathcal{A}} \sum_{r \geq l} |\zeta^\top S_{r+1}(A, \tilde{\zeta})| \leq 4C(\alpha) \sup_{A \in \mathcal{A}} \|A \tilde{\zeta}\|_2 (\sqrt{t} \gamma'_2(\mathcal{A}, d_2) + t^{\frac{1}{\alpha}} \gamma'_\alpha(\mathcal{A}, d_\infty)).$$

As a consequence, we have for $t > 16$

$$\mathbb{P}\left\{\sup_{A \in \mathcal{A}} \sum_{r \geq l} |\zeta^\top S_{r+1}(A, \tilde{\zeta})| > C_2(\alpha) \sup_{A \in \mathcal{A}} \|A\tilde{\zeta}\|_2 t (\gamma'_2(\mathcal{A}, d_2) + \gamma'_\alpha(\mathcal{A}, d_\infty))\right\} \leq C_1(\alpha) \exp(-c(\alpha)pt).$$

A direct integration and taking expectation with respect to variables $\{\tilde{\eta}_i, i \leq n\}$ lead to

$$\left\| \sup_{A \in \mathcal{A}} \sum_{r \geq l} |\zeta^\top S_{r+1}(A, \tilde{\zeta})| \right\|_{L_p} \lesssim_\alpha \left\| \sup_{A \in \mathcal{A}} \|A\tilde{\zeta}\|_2 \right\|_{L_p} (\gamma_2(\mathcal{A}, d_2) + \gamma_\alpha(\mathcal{A}, d_\infty)).$$

Following the same line, one can give a similar bound for $\sup_{A \in \mathcal{A}} \sum_{r \geq l} |\zeta^\top \pi_r(A)^\top \Lambda_{r+1}(A) \tilde{\zeta}|$. By virtue of (3.3), we have for $p \geq 1$

$$\left\| \sup_{A \in \mathcal{A}} |\zeta^\top A^\top A \tilde{\zeta} - \eta^\top \pi_l(A)^\top \pi_l(A) \tilde{\zeta}| \right\|_{L_p} \lesssim_\alpha \left\| \sup_{A \in \mathcal{A}} \|A\tilde{\zeta}\|_2 \right\|_{L_p} (\gamma_2(\mathcal{A}, d_2) + \gamma_\alpha(\mathcal{A}, d_\infty)).$$

As for the other part of (3.2), we have for $p \geq 1$

$$\mathbb{E} \sup_{A \in \mathcal{A}} |\zeta^\top \pi_l(A)^\top \pi_l(A) \tilde{\eta}|^p \leq \sum_{B \in T_l} \mathbb{E} |\zeta^\top B^\top B \tilde{\zeta}|^p \leq e^p \sup_{A \in \mathcal{A}} \mathbb{E} |\zeta^\top A^\top A \tilde{\zeta}|^p.$$

Thus

$$\left\| \sup_{A \in \mathcal{A}} |\zeta^\top \pi_l(A)^\top \pi_l(A) \tilde{\zeta}| \right\|_{L_p} \leq e \sup_{A \in \mathcal{A}} \|\zeta^\top A^\top A \tilde{\zeta}\|_{L_p}.$$

Then, we conclude the proof by virtue of (3.2). \square

Proof of Theorem 1.1. Let $\zeta = (\zeta_1, \dots, \zeta_n)$ and $\zeta_1, \dots, \zeta_n \stackrel{i.i.d.}{\sim} \mathcal{W}_s(\alpha)$, $0 < \alpha \leq 1$. We have by Remark 2.3 and Proposition 3

$$\left\| \sup_{A \in \mathcal{A}} \left| \|A\tilde{\zeta}\|_2^2 - \mathbb{E} \|A\tilde{\zeta}\|_2^2 \right| \right\|_{L_p} \lesssim_{L, \alpha} \sup_{A \in \mathcal{A}} \|\zeta^\top A^\top A \tilde{\zeta}\|_{L_p} + \left\| \sup_{A \in \mathcal{A}} \|A\tilde{\zeta}\|_2 \right\|_{L_p} (\gamma_2(\mathcal{A}, \|\cdot\|_{l_2 \rightarrow l_2}) + \gamma_\alpha(\mathcal{A}, \|\cdot\|_{l_2 \rightarrow l_\infty}))$$

Set $S = \{A^\top x : x \in B_2^n, A \in \mathcal{A}\}$. Lemmas 2.3 and 2.5 yield that

$$\begin{aligned} \left\| \sup_{A \in \mathcal{A}} \|A\tilde{\zeta}\|_2 \right\|_{L_p} &= \left(\mathbb{E} \sup_{A \in \mathcal{A}, x \in B_2^n} |x^\top A \tilde{\zeta}|^p \right)^{1/p} = \left(\mathbb{E} \sup_{u \in S} |u^\top \tilde{\zeta}|^p \right)^{1/p} \lesssim_\alpha \mathbb{E} \sup_{u \in S} |u^\top \tilde{\zeta}| + \sup_{u \in S} \|u^\top \tilde{\zeta}\|_{L_p} \\ &\lesssim_\alpha \mathbb{E} \sup_{u \in S} |u^\top \tilde{\zeta}| + \sqrt{p} \sup_{u \in S} \|u\|_2 + p^{1/\alpha} \sup_{u \in S} \|u\|_\infty \\ (3.5) \quad &= \mathbb{E} \sup_{A \in \mathcal{A}} \|A\tilde{\zeta}\|_2 + \sqrt{p} M_{l_2 \rightarrow l_2}(\mathcal{A}) + p^{1/\alpha} M_{l_2 \rightarrow l_\infty}(\mathcal{A}). \end{aligned}$$

We have by Lemma 2.6 and Proposition 3

$$\begin{aligned} \mathbb{E} \sup_{A \in \mathcal{A}} \|A\tilde{\zeta}\|_2^2 &\leq \mathbb{E} \sup_{A \in \mathcal{A}} \left| \|A\tilde{\zeta}\|_2^2 - \mathbb{E} \|A\tilde{\zeta}\|_2^2 \right| + \sup_{A \in \mathcal{A}} \mathbb{E} \|A\tilde{\zeta}\|_2^2 \lesssim_\alpha \mathbb{E} \sup_{A \in \mathcal{A}} |\zeta^\top A^\top A \tilde{\zeta}| + M_F^2(\mathcal{A}) \\ &\lesssim_\alpha \mathbb{E} \sup_{A \in \mathcal{A}} \|A\tilde{\zeta}\|_2 (\gamma_2(\mathcal{A}, \|\cdot\|_{l_2 \rightarrow l_2}) + \gamma_\alpha(\mathcal{A}, \|\cdot\|_{l_2 \rightarrow l_\infty})) + M_F^2(\mathcal{A}). \end{aligned}$$

Therefore,

$$\mathbb{E} \sup_{A \in \mathcal{A}} \|A\tilde{\zeta}\|_2 \leq \left\| \sup_{A \in \mathcal{A}} \|A\tilde{\zeta}\|_2 \right\|_{L_2} \lesssim_\alpha \gamma_2(\mathcal{A}, \|\cdot\|_{l_2 \rightarrow l_2}) + \gamma_\alpha(\mathcal{A}, \|\cdot\|_{l_2 \rightarrow l_\infty}) + M_F(\mathcal{A}).$$

Then, we conclude the proof by Remark 2.1 and Lemma 2.2. \square

4. APPLICATIONS

An $m \times n$ matrix A has the restricted isometry property (R.I.P.) with parameters δ and s if satisfying for all s -sparse x (i.e., $|\{l : x_l \neq 0\}| \leq s$)

$$(4.1) \quad (1 - \delta)\|x\|_2^2 \leq \|Ax\|_2^2 \leq (1 + \delta)\|x\|_2^2.$$

The restricted isometry constant δ_s is defined as the smallest number satisfying (4.1). In this section, we shall prove the R.I.P. of partial random circulant matrices (see the definition below).

Recall the cyclic subtraction: $j \ominus k = (j - k) \bmod n$. Denote by $z * x = ((z * x)_1, \dots, (z * x)_n)^\top$ the circular convolution of two vectors $x, y \in \mathbb{R}^n$, where

$$(z * x)_j := \sum_{k=1}^n z_{j \ominus k} x_k.$$

The circulant matrix $H = H_z$ generated by z is a $n \times n$ matrix with entries $H_{jk} = z_{j \ominus k}$. Equivalently, $Hx = z * x$ for every $x \in \mathbb{R}^n$.

Let $\Omega \subset \{1, \dots, n\}$ be a fixed set with $|\Omega| = m$. Denote by $R_\Omega : \mathbb{R}^n \rightarrow \mathbb{R}^m$ the operator restricting a vector $x \in \mathbb{R}^n$ to its entries in Ω . The partial circulant matrix generated by z is defined as follows

$$\Phi = \frac{1}{\sqrt{m}} R_\Omega H_z.$$

We say Φ a partial random circulant matrix when z is a random vector.

Krahmer et al [8] showed the R.I.P. of partial random circulant matrices generated by standard subgaussian random vectors (i.e., random vectors with independent standard subgaussian entries). Dai et al [3] extended this result to the case where the generating random vectors are standard α -subexponential vectors, $1 \leq \alpha \leq 2$. The following result extends this result to the case $0 < \alpha \leq 1$.

Theorem 4.1. *Let Φ be an $m \times n$ partial random circulant matrix generated by a standard α -subexponential random vector η , $0 < \alpha \leq 1$. Set $L = \max_{i \leq n} \|\eta_i\|_{\Psi_\alpha}$. Then, under the condition $m \geq c_1(\alpha, L) \delta^{-2} s^{2/\alpha} \log^{4/\alpha} n$, the restricted isometry constant δ_s satisfies*

$$P\{\delta_s \leq \delta\} \geq 1 - \exp(-c_0(\alpha, L) s^{2/\alpha} \log^{4/\alpha} n).$$

Proof of Theorem 4.1. Let Ω be a subset of $\{1, \dots, n\}$ with $|\Omega| = m$ and $P_\Omega = R_\Omega^\top R_\Omega$, where $R_\Omega : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is the operator restricting a vector $x \in \mathbb{R}^n$ to its entries in Ω . In particular, $P_\Omega : \mathbb{R}^n \rightarrow \mathbb{R}^n$, is a projection operator such that

$$(P_\Omega x)_l = x_l, \quad l \in \Omega; \quad (P_\Omega x)_l = 0, \quad l \notin \Omega.$$

Let $D_{s,n} = \{x \in \mathbb{R}^n : \|x\|_2 \leq 1, \|x\|_0 \leq s\}$. We have by (4.1),

$$\begin{aligned} \delta_s &= \sup_{x \in D_{s,n}} \left| \frac{1}{m} \|\Phi x\|_2^2 - \|x\|_2^2 \right| = \sup_{x \in D_{s,n}} \left| \frac{1}{m} \|P_\Omega(\eta * x)\|_2^2 - \|x\|_2^2 \right| \\ &= \sup_{x \in D_{s,n}} \left| \frac{1}{m} \|P_\Omega(x * \eta)\|_2^2 - \|x\|_2^2 \right| = \sup_{x \in D_{s,n}} \left| \|V_x \eta\|_2^2 - \|x\|_2^2 \right| \\ (4.2) \quad &= \sup_{x \in D_{s,n}} \left| \|V_x \eta\|_2^2 - \mathbb{E} \|V_x \eta\|_2^2 \right|, \end{aligned}$$

where the last equality is due to $\mathbb{E} \|V_x \eta\|_2^2 = \|x\|_2^2$.

Let $\mathcal{A} = \{V_x : x \in D_{s,n}\}$. To bound δ_s , we need to control the quantities $M_F(\mathcal{A})$, $M_{l_2 \rightarrow l_\beta}(\mathcal{A})$ and $\gamma_\alpha(\mathcal{A}, \|\cdot\|_{l_2 \rightarrow l_\beta})$ ($\beta = 2, \infty$) by virtue of Theorem 1.1.

Observe that the matrices V_x have shifted copies of x in all their m nonzero rows. Thus, the l_2 -norm of each nonzero row is $m^{-1/2}\|x\|_2$. Then, for $x \in D_{s,n}$, we have $\|V_x\|_F = \|x\|_2 \leq 1$, and thus $M_F(\mathcal{A}) = 1$.

Note that

$$\|V_x\|_{l_2 \rightarrow l_\beta} \leq \|V_x\|_{l_2 \rightarrow l_2} \leq \sup_{\|y\|_2, \|z\|_2 \leq 1} \frac{1}{\sqrt{m}} |y^\top V_x z| \leq \frac{1}{\sqrt{m}} \|x\|_1 \leq \sqrt{\frac{s}{m}} \|x\|_2 \leq \sqrt{\frac{s}{m}},$$

It follows immediately from that $M_{l_2 \rightarrow l_\beta}(\mathcal{A}) \leq \sqrt{s/m}$ ($\beta = 2, \infty$).

Lastly, we use the covering number to bound $\gamma_\alpha(\mathcal{A}, \|\cdot\|_{l_2 \rightarrow l_\beta})$. In particular, we have

$$\gamma_\alpha(\mathcal{A}, \|\cdot\|_{l_2 \rightarrow l_\beta}) \lesssim_\alpha \int_0^\infty (\log N(T, \|\cdot\|_{l_2 \rightarrow l_\beta}, u))^{1/\alpha} du,$$

where $N(T, d, u)$ is the covering number of T with the distance d and the radius u . Note that, Krahmer et al [8] showed the following bounds for $u \geq 1/\sqrt{m}$

$$\log N(\mathcal{A}, \|\cdot\|_{l_2 \rightarrow l_2}, u) \lesssim \frac{s}{m} \left(\frac{\log n}{u}\right)^2$$

and for $u \leq 1/\sqrt{m}$

$$\log N(\mathcal{A}, \|\cdot\|_{l_2 \rightarrow l_2}, u) \lesssim s \log\left(\frac{en}{su}\right).$$

A direct integration (Section 5.2 in [3] contains a detailed calculation) yields that for $0 < \alpha \leq 1$

$$\gamma_\alpha(\mathcal{A}, \|\cdot\|_{l_2 \rightarrow l_\infty}) \leq \gamma_\alpha(\mathcal{A}, \|\cdot\|_{l_2 \rightarrow l_2}) \lesssim_\alpha \frac{s^{1/\alpha}}{\sqrt{m}} \log^{2/\alpha} n$$

and for $\alpha = 2$

$$\gamma_2(\mathcal{A}, \|\cdot\|_{l_2 \rightarrow l_2}) \lesssim \sqrt{\frac{s}{m}} \log s \log n.$$

Assume for any $0 < \delta, \alpha \leq 1$

$$m \geq c_1(\alpha, L) \delta^{-2} \max \left\{ (s^{2/\alpha} \log^{4/\alpha} n), (s \log^2 s \log^2 n) \right\} = c_1(\alpha, L) \delta^{-2} s^{2/\alpha} \log^{4/\alpha} n.$$

Then, we have

$$\gamma_2(\mathcal{A}, \|\cdot\|_{l_2 \rightarrow l_2}), \gamma_\alpha(\mathcal{A}, \|\cdot\|_{l_2 \rightarrow l_\infty}) \lesssim_{\alpha, L} \delta.$$

Let $c_1(\alpha, L)$ be an appropriately constant such that $C(\alpha) L^2 U_1(\alpha, \infty) \leq \delta/2$, where $C(\alpha)$ is the constant in Theorem 1.1. Then, we have

$$\mathbb{P}\{\delta_s \geq \delta\} \leq \mathbb{P}\{\delta_s \geq C(\alpha) L^2 U_1(\alpha, \infty) + \delta/2\} \leq C_1(\alpha) \exp(-c_2(\alpha, L)(m/s)^{\alpha/2} \delta^\alpha),$$

which concludes the proof. \square

Acknowledgment The work was partly supported by the National Natural Science Foundation of China (12271475, U23A2064).

REFERENCES

- [1] Bourgain, J., Dirksen, S., Nelson, J. (2015). Toward a unified theory of sparse dimensionality reduction in Euclidean space. *Geom. Funct. Anal.* **25**(4) 1009–1088.
- [2] Chen, X. H., Yang, Y. (2021). Hanson–Wright inequality in Hilbert spaces with application to K-means clustering for non-Euclidean data. *Bernoulli* **27**(1) 586–614.
- [3] Dai, G.Z., Su, Z.G., Ulyanov, V., Wang, H.C. (2024). On log-concave-tailed chaos and the restricted isometry property. arXiv: 2401.14860.
- [4] Götze, F., Sambale, H., Sinulis, A. (2021) Concentration inequalities for polynomials in α -sub-exponential random variables. *Electron. J. Probab.* **48**(26) 1–22.
- [5] Klochov, Y., Zhivotovskiy, N. (2020) Uniform Hanson-Wright type concentration inequalities for unbounded entries via the entropy method. *Electron. J. Probab.* **25**(22) 1–30.
- [6] Koep, N., Behboodi, A., Mathar, R. (2022). The restricted isometry property of block diagonal matrices for group-sparse signal recovery. *Appl. Comput. Harmon. Anal.* **60** 333–367.
- [7] Kolesko, K., Latała, R. (2015). Moment estimates for chaos generated by symmetric random variables with logarithmically convex tails. *Statist. Probab. Lett.* **107** 210–214.
- [8] Krahmer, F., Mendelson, S., Rauhut, H. (2014). Suprema of chaos processes and the restricted isometry property. *Comm. Pure Appl. Math.* **67** 1877–1904.
- [9] Latała, R., Strzelecka, M. (2018). Comparison of weak and strong moments for vectors with independent coordinates. *Mathematika* **64**(1) 211–229.
- [10] Latała, R., van Handel, R. and Youssef, P. (2018). The dimension-free structure of nonhomogeneous random matrices. *Invent. math.* **214** 1031–1080.
- [11] Ledoux, M., Talagrand, M. (1991). Probability in Banach spaces. Isoperimetry and Processes. *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)*, 23. Springer, Berlin.
- [12] Rudelson, M., Vershynin, R. (2013). Hanson-Wright inequality and sub-gaussian concentration. *Electron. Commun. Probab.* **18** 82–91.
- [13] Talagrand, M. (1996). New concentration inequalities in product spaces. *Invent. math.* **126** 505–563.
- [14] Talagrand, M. (2001). Majorizing measures without measures *Ann. Prob.* **29**(1) 411–417.
- [15] Zhou, S.H. (2019). Sparse Hanson–Wright inequalities for subgaussian quadratic forms. *Bernoulli* **25**(3) 1603–1639.

ZHEJIANG UNIVERSITY, HANGZHOU, 310027, CHINA.
 Email address: 11935022@zju.edu.cn

ZHEJIANG UNIVERSITY, HANGZHOU, 310027, CHINA.
 Email address: suzhonggen@zju.edu.cn