HOCHSTER'S TYPE FORMULAE

FERNANDO SANCHO DE SALAS AND ALEJANDRO TORRES SANCHO

ABSTRACT. We give an elementary proof and generalize some Hochsters's type formulae on local cohomology and Ext's of squarefree modules.

INTRODUCTION

Let k be a commutative ring and $R = k[x_1, \ldots, x_n]$ the polynomial ring over k. Let $\mathbb{A}_{F_1}^n$ be the finite poset of subsets of the finite set $\Delta_n = \{1, \ldots, n\}$ and let us denote by $\mathrm{Shv}_k(\mathbb{A}_{F_1}^n)$ the category of sheaves of k-modules on $\mathbb{A}_{F_1}^n$. In [8], Yanagawa shows that a sheaf of k-modules on $\mathbb{A}_{F_1}^n$ has an associated squarefree R-module; this defines an exact functor:

$$\pi^{\star} \colon \operatorname{Shv}_k(\mathbb{A}^n_{\mathbf{F}_1}) \longrightarrow R - \operatorname{mod}$$

(where R - mod. denotes the category of all R-modules) which maps each sheaf F on $\mathbb{A}_{F_1}^n$ to its associated squarefree module. This functor has an exact right adjoint

$$\pi_{\bigstar} \colon R - \text{mod.} \longrightarrow \text{Shv}_k(\mathbb{A}^n_{\mathbf{F}_1}).$$

Let D(R) be the derived category of complexes of R-modules and $D(\mathbb{A}_{F_1}^n)$ the derived category of complexes of sheaves of k-modules on $\mathbb{A}_{F_1}^n$. A complex $F \in D(\mathbb{A}_{F_1}^n)$ is called *pointwise perfect* if its stalk F_p at any point $p \in \mathbb{A}_{F_1}^n$ is a perfect complex of k-modules, i.e., F_p is quasi-isomorphic to a bounded complex of finitely generated projective k-modules.

Let $k_{\{1\}}$ be the constant sheaf k on $\mathbb{A}_{F_1}^n$ supported at the point $\mathbf{1} \in \mathbb{A}_{F_1}^n$ that represents the total subset of Δ_n . For any $F \in D(\mathbb{A}_{F_1}^n)$, let us denote

$$D(F) = \mathbb{R}\mathcal{H}om^{\bullet}_{\mathbb{A}^n_{\mathbf{F}_{\bullet}}}(F, k_{\{\mathbf{1}\}})$$

and for any $M \in D(R)$ let us denote

$$D(M) = \mathbb{R} \operatorname{Hom}^{\bullet}_{R}(M, \Omega^{n}_{R/k})$$

where $\Omega_{R/k}^n$ is the module of highest Kähler differentials of R over k. The main theorem of this paper is the following:

Theorem. [Thm. 2.1] For any $M \in D(R)$, and any pointwise perfect complex $F \in D(\mathbb{A}^n_{\mathbb{F}_1})$, one has

$$\mathbb{R}\operatorname{Hom}_{R}^{\bullet}(M, \pi^{\star}F) = \mathbb{R}\operatorname{Hom}_{\mathbb{A}_{F_{\star}}^{n}}^{\bullet}(D(F), \pi_{\star}D(M)).$$

We shall see that this theorem yields Hochster's type formulae whenever one is able to decompose $\pi_{\star} D(M)$ as a direct sum, and this of course highly depends on the choice of M. The

²⁰²⁰ Mathematics Subject Classification. 13F55, 13D45.

Key words and phrases. Hochster formula, Stanley-Reisner theory.

Work supported by Grant PID2021-128665NB-I00 funded by MCIN/AEI/ 10.13039/501100011033 and, as appropriate, by "ERDF A way of making Europe".

first example of this is given in Theorem 2.4, obtaining a Hochster formula when k is a field and D(M) = I[r] for some injective R-module I and some integer r.

Let us denote $\mathfrak{m} = (x_1, \ldots, x_n)$ and $\mathfrak{m}_l = (x_1^l, \ldots, x_n^l)$. In Theorem 2.6 we give a Hochster formula for the local cohomology modules $H^i_{\mathfrak{m}}(\pi^{\star}F)$, which generalizes Yanagawa's results, stated for a sheaf F of finite dimensional vector k-spaces, where k is a field ([7, Thm. 3.3, Thm. 3.5], see also [2, Thm. 1.1]) for any ring k and any pointwise perfect complex F of k-modules. In this case, the formula is a consequence of Theorem 2.1 and of the decomposition of $\pi_{\star}H^n_{\mathfrak{m}}(\Omega^n_{R/k})$ given in Proposition 2.5, (a). We also give the explicit R-module structure, in terms of residue maps, of the right hand term $\bigoplus_{p \in \text{supp}(F)} H^{i-c_p}_p(U_p, F) \otimes_k H^{c_p}_{\mathfrak{m}}(R_{C_p})$, in Remarks 2.7, (2).

In Theorem 2.8 we give a Hochster formula for $\operatorname{Ext}_{R}^{i}(R/\mathfrak{m}_{l}, \pi^{\star}F)$ for any pointwise perfect complex F. As a particular case, we obtain Miyazaki's result ([4, Thm. 1]). In this case, the formula is a consequence of Theorem 2.1 and of the decomposition of $\pi_{\star} \operatorname{Hom}_{R}(R/\mathfrak{m}_{l}, H^{n}_{\mathfrak{m}}(\Omega^{n}_{R/k}))$ given in Proposition 2.5, (b).

Finally, let us mention that the decompositions of Proposition 2.5 are easily obtained by reduction to the case n = 1, thanks to the additivity of π_{\star} (Proposition 1.5).

In the last thirty years, many Hochster's type formula have appeared in the literature (see specially [1], where many Hochster's type formulae are obtained and compared with previous work on the subject), essentially providing Hochster's type formulae for arbitrary posets. We shall see at the end of the paper how these results essentially differ from ours.

1. Preliminaries and notations

1.1 (The finite poset $\mathbb{A}_{F_1}^n$). Let $\Delta_n = \{1, \ldots, n\}$ be the finite set with n elements.

Definition 1.2. We shall denote

$$\mathbb{A}^n_{\mathbf{F}_1} := \mathcal{P}(\Delta_n)$$

the set of subsets of Δ_n , which is a finite topological space (a finite poset) with the topology given by the partial order defined by inclusion: $p \leq q \Leftrightarrow p \subseteq q$. Any finite poset (X, \leq) will be considered as a topological space with its Alexandroff topology: a subset U of X is open if $p \in U$ and $q \geq p$ implies $q \in U$.

We shall denote:

- $\mathbf{0} \in \mathbb{A}_{F_1}^n$ the element representing the empty subset of Δ_n . It is the unique closed point of $\mathbb{A}_{F_1}^n$.
- $\mathbf{1} \in \mathbb{A}_{F_1}^n$ the generic point of $\mathbb{A}_{F_1}^n$, i.e., the element representing the total subset of Δ_n . It is the unique open point of $\mathbb{A}_{F_1}^n$.
- For each $p \in \mathbb{A}_{F_1}^n$, we denote

 $U_p = \text{ smallest open subset containing } p = \{q \in \mathbb{A}^n_{\mathbb{F}_1} : q \ge p\}$

 $C_p = \text{closure of } p = \{q \in \mathbb{A}^n_{\mathbb{F}_1} : q \le p\}$

- For each $i \in \Delta_n$, we shall denote by U_i the open subset of $\mathbb{A}^n_{\mathbb{F}_1}$ whose elements are the subsets of Δ_n containing i; in other words $U_i = U_{\{i\}}$.
- We shall denote $H_i := \mathbb{A}_{F_1}^n U_i$ the hyperplane of $\mathbb{A}_{F_1}^n$ whose elements are the subsets of Δ_n contained in $\Delta_n \{i\}$.
- For any $p \in \mathbb{A}_{F_1}^n$, \hat{p} denotes the complement subset: $\hat{p} = \Delta_n p$.

A finite simplicial complex K (with at most n vertices) is just a closed subset of $\mathbb{A}_{\mathbf{F}_1}^n$. Recall that for any simplical complex $K \subseteq \mathbb{A}^n_{\mathbb{F}_1}$ and any $p \in K$, the link of p in K is:

$$\operatorname{Link}_{K}(p) = \{ q \in K : q \neq \mathbf{0}, p \cap q = \mathbf{0} \text{ and } p \cup q \in K \}$$

which is a closed subset of $K^* = K - \{0\}$. If we denote $U_p^* = U_p - \{p\}$, one has an homeomorphism (equivalently, an order preserving bijection)

$$\operatorname{Link}_{K}(p) \xrightarrow{\sim} U_{p}^{*} \cap K$$
$$q \longmapsto p \cup q \quad .$$
$$x - p \longleftrightarrow x$$

We shall denote by $\operatorname{Shv}_k(\mathbb{A}^n_{\mathbb{F}_1})$ the category of sheaves of k-modules on $\mathbb{A}^n_{\mathbb{F}_1}$ and by $D(\mathbb{A}^n_{\mathbb{F}_1})$ the derived category of complexes of sheaves of k-modules on $\mathbb{A}^n_{\mathbb{F}_1}$. We shall use the standard notations of the derived category of sheaves on a topological space. As it is well known, a sheaf F of k-modules on $\mathbb{A}_{F_1}^n$ is equivalent to give, for each $p \in \mathbb{A}_{F_1}^n$, a k-module F_p , and for each $p \leq q$ a morphism of k-modules $r_{pq}: F_p \to F_q$, such that $r_{pp} = \operatorname{id}_{F_p}$ and $r_{ql} \circ r_{pq} = r_{pl}$ for any $p \le q \le l$. The constant sheaf k on $\mathbb{A}_{F_1}^n$ is still denoted by k. For any $p \in \mathbb{A}_{F_1}^n$ we denote

$$k_{U_p}$$
 , k_{C_p} , $k_{\{p\}}$

the constant sheaf k supported, respectively, on U_p , C_p and $\{p\}$. We shall use the basic formulae

$$\mathbb{R} \operatorname{Hom}_{\mathbb{A}_{F_{1}}^{n}}^{\mathbb{A}}(k_{U_{p}}, F) = F_{p}$$
$$\mathbb{R} \operatorname{Hom}_{\mathbb{A}_{F_{1}}^{n}}^{\mathbb{A}}(F, k_{C_{p}}) = \mathbb{R} \operatorname{Hom}_{k}^{\mathbb{A}}(F_{p}, k)$$
$$\mathbb{R} \operatorname{Hom}_{\mathbb{A}_{F_{1}}^{n}}^{\mathbb{A}}(k_{\{p\}}, F) = \mathbb{R}\Gamma_{p}(U_{p}, F),$$

where \mathbb{R} Hom[•] is the right derived functor of Hom[•] (the complex of homomorphisms) and $\Gamma_p(U_p, \dots)$ denotes the functor of sections with support in p.

One has a natural isomorphism

$$\mathbb{A}^n_{\mathbb{F}_1} = \mathbb{A}^1_{\mathbb{F}_1} imes \stackrel{n}{\cdots} imes \mathbb{A}^1_{\mathbb{F}_1}$$

so an element $p \in \mathbb{A}_{F_1}^n$ is identified with a sequence $p = (p_1, \ldots, p_n)$ with $p_i \in \mathbb{A}_{F_1}^1 = \{0, 1\}$. Explicitly, $p_i = \mathbf{0}$ if $i \notin p$ and $p_i = \mathbf{1}$ if $i \in p$.

For each i = 1, ..., n, we have the *i*-th projection

$$\phi_i \colon \mathbb{A}^n_{\mathbf{F}_1} \longrightarrow \mathbb{A}^1_{\mathbf{F}_1}$$

defined by $\phi_i^{-1}(\mathbf{0}) = H_i$ (and then $\phi_i^{-1}(\mathbf{1}) = U_i$).

If F_1, \ldots, F_n are sheaves on $\mathbb{A}^1_{\mathbb{F}_1}$ we have a sheaf on $\mathbb{A}^n_{\mathbb{F}_1}$:

$$F_1 \boxtimes \cdots \boxtimes F_n := \phi_1^{-1} F_1 \otimes_k \cdots \otimes_k \phi_n^{-1} F_n$$

whose stalkwise description is

 $(F_1 \boxtimes \cdots \boxtimes F_n)_p = (F_1)_{p_1} \otimes_k \cdots \otimes_k (F_n)_{p_n}$

with $p = (p_1, \ldots, p_n)$, via the equality $\mathbb{A}_{\mathbb{F}_1}^n = \mathbb{A}_{\mathbb{F}_1}^1 \times \cdots \times \mathbb{A}_{\mathbb{F}_1}^1$.

1.3 (Pointwise perfect complexes and duality). A complex $F \in D(\mathbb{A}_{F_1}^n)$ is called *pointwise* perfect if F_p is a perfect complex of k-modules for any $p \in \mathbb{A}_{F_1}^n$. For example, if k is a field, then F is pointwise perfect if and only if F_p is a complex of k-vector spaces with bounded and finite dimensional cohomology. We shall denote by $D_{perf}(\mathbb{A}_{F_1}^n)$ the subcategory of pointwise perfect complexes.

For any ring k, D(k) denotes the derived category of complexes of k-modules. For any $E \in D(k)$, we denote $E^{\vee} = \mathbb{R} \operatorname{Hom}_{k}^{\bullet}(E,k)$. One has a natural morphism $E \to E^{\vee\vee}$ which is an isomorphism if E is perfect. For any $E \in D(k)$, we still denote by E the constant complex in $D(\mathbb{A}_{F_1}^n)$ defined by $E_p = E$ for any $p \in \mathbb{A}_{F_1}^n$ and the identity morphism $E_p \to E_q$ for any $p \leq q$. For each $F \in D(\mathbb{A}_{F_1}^n)$ we shall denote

 $E \in D(\mathbb{A}_{F_1})$ we shall denote

$$D(F) := \mathbb{R}\mathcal{H}om^{\bullet}_{\mathbb{A}^n_{\mathsf{F}_1}}(F, k_{\{\mathbf{1}\}}),$$

where $\mathcal{H}om^{\bullet}_{\mathbb{A}^n_{\mathbb{F}_1}}$ denotes the complex of sheaves of homomorphisms and $\mathbb{R}\mathcal{H}om^{\bullet}_{\mathbb{A}^n_{\mathbb{F}_1}}$ its right derived functor.

We shall use the following property (see [5]):

If F is pointwise perfect, so is D(F), and for any $p \in \mathbb{A}_{F_1}^n$ and any $E \in D(k)$, one has:

$$\mathbb{R}\operatorname{Hom}_{\mathbb{A}_{\mathbf{F}_{1}}^{n}}^{\bullet}(D(F), E \otimes_{k} k_{C_{p}}) = \mathbb{R}\Gamma_{p}(U_{p}, F) \overset{\mathbb{L}}{\otimes}_{k} E[n - c_{p}].$$
(1.3.1)

The support of a complex $F \in D(\mathbb{A}^n_{F_1})$ is the set

$$\operatorname{supp}(F) = \{ p \in \mathbb{A}_{\mathbb{F}_1}^n : F_p \neq 0 \text{ (in } D(k)) \}.$$

We shall denote by $\overline{\operatorname{supp}(F)}$ the closure of $\operatorname{supp}(F)$. Notice that $F_{|U_p|} = 0$ if $p \notin \overline{\operatorname{supp}(F)}$.

1.4 (Yanagawa's functor and its right adjoint). Let $R = k[x_1, \ldots, x_n]$. If M is a \mathbb{Z}^n -graded R-module (or k-module or a sheaf of k-modules) and $\alpha = (\alpha_1, \ldots, \alpha_n) \in \mathbb{Z}^n$, then $M(\alpha)$ denotes the module M shifted by α , that is, it is the same module M but with the gradation

$$[M(\alpha)]_{(m_1,...,m_n)} = M_{(m_1+\alpha_1,...,m_n+\alpha_n)}.$$

One has that $M(\alpha) = M \otimes_R R(\alpha)$. Hence, for any (ungraded) *R*-module *M*, we shall denote $M(\alpha) := M \otimes_R R(\alpha)$. For each $p \in \mathbb{A}^n_{\mathbf{F}_1}$, $l \in \mathbb{Z}$, we shall denote

$$l_p = (\alpha_1, \dots, \alpha_n), \text{ with } \alpha_i = \begin{cases} l, \text{ if } i \in p \\ 0, \text{ if } i \notin p \end{cases}$$

For any $p \in \mathbb{A}_{F_1}^n$, \mathbf{x}^p denotes the squarefree monomial $\mathbf{x}^p = \prod_{i \in p} x_i$. One has a natural isomorphism of \mathbb{Z}^n -graded *R*-modules $(\mathbf{x}^p) = R(-1_p)$. For any closed subset $K \subseteq \mathbb{A}_{F_1}^n$, I_K denotes the Stanley–Reisner ideal associated to K, i.e., the ideal generated by the monomials \mathbf{x}^p with $p \notin K$. We shall denote $R_K = R/I_K$, the Stanley–Reisner ring associated to K. In particular, for each $p \in \mathbb{A}_{F_1}^n$, R_{C_p} is the polynomial ring in the variables x_i , $i \in p$.

Let us consider Yanagawa's functor ([8])

$$\pi^{\star} \colon \operatorname{Shv}_k(\mathbb{A}^n_{\mathbf{F}_1}) \longrightarrow R - \operatorname{mod}$$

which associates, to each sheaf F, a squarefree module $\pi^{\star}F$. If $F \to F'$ is a morphism of sheaves, then $\pi^{\star}F \to \pi^{\star}F'$ is a homogeneous (i.e. \mathbb{Z}^n -degree preserving) morphism of \mathbb{Z}^n -graded *R*-modules. We shall not use Yanagawa's original definition. Instead, we shall use an alternative construction given in [5]: the functor π^{\star} is defined as the unique left exact functor

 π^{\star} : Shv_k($\mathbb{A}^n_{\mathbb{F}_1}$) $\longrightarrow R$ – mod that maps k_{U_p} to (\mathbf{x}^p) for any $p \in \mathbb{A}^n_{\mathbb{F}_1}$ and the natural inclusion

 $k_{U_q} \hookrightarrow k_{U_p}$ to the natural inclusion $(\mathbf{x}^q) \hookrightarrow (\mathbf{x}^p)$ for any $p \leq q$. Further properties of π^* are: It is an exact functor and it satisfies:

$$\pi^{\bigstar}(k_K) = R_K$$

for any closed subset $K \subseteq \mathbb{A}_{F_1}^n$. Also notice that,

$$\pi^{\star}(k_{\{1\}}) = (\mathbf{x}^{1}) = \Omega_{R/k}^{n}$$
$$\mathbf{x}^{1} \mapsto \mathrm{d}\, x_{1} \wedge \dots \wedge \mathrm{d}\, x_{n}$$

where $\Omega^n_{\mathbb{A}^n/k}$ is the module of highest Kähler differentials.

The functor π^{\star} has a right adjoint (because it is exact and commutes with direct limits)

 $\pi_{\star} \colon R - \mathrm{mod} \longrightarrow \mathrm{Shv}_k(\mathbb{A}^n_{\mathsf{F}_1})$

whose stalkwise description is the following: for any *R*-module *M* and any $p \in \mathbb{A}_{F_1}^n$,

$$(\pi_{\star}M)_p = \operatorname{Hom}_R((\mathbf{x}^p), M) = M \otimes_R R(1_p)$$

and for any $p \leq q$, the morphism $(\pi_{\star}M)_p \to (\pi_{\star}M)_q$ is obtained by applying $\operatorname{Hom}_R(-,M)$ to the inclusion $(\mathbf{x}^q) \hookrightarrow (\mathbf{x}^p)$. Then, π_{\star} is an exact functor. Moreover, if M is a \mathbb{Z}^n -graded R-module, then $\pi_{\star}M$ is a sheaf of \mathbb{Z}^n -graded k-modules (in fact, R-modules) on $\mathbb{A}^n_{\mathbb{F}_1}$.

Proposition 1.5 (Additivity of π_{\star}). Let $R = k[x_1, \ldots, x_n]$. For each $i = 1, \ldots, n$, let M_i be a $k[x_i]$ -module. Then $M_1 \otimes_k \cdots \otimes_k M_n$ is an R-module and:

$$\pi_{\bigstar}(M_1 \otimes_k \cdots \otimes_k M_n) = (\pi_{\bigstar} M_1) \boxtimes \cdots \boxtimes (\pi_{\bigstar} M_n),$$

If each M_i is a \mathbb{Z} -graded $k[x_i]$ -module, then this equality is an isomorphism of \mathbb{Z}^n -graded sheaves of k-modules.

Proof. Let $p = (p_1, \ldots, p_n) \in \mathbb{A}_{\mathbb{F}_1}^n$. Then $(\mathbf{x}^p) = (x_1^{p_1}) \otimes_k \cdots \otimes_k (x^{p_n})$, where $x_i^{p_i} = 1$ if $p_i = \mathbf{0}$ and $x_i^{p_i} = x_i$ if $p_i = \mathbf{1}$. Then

$$[\pi_{\bigstar}(M_1 \otimes_k \cdots \otimes_k M_n)]_p = (M_1 \otimes_k \cdots \otimes_k M_n) \otimes_R R(1_p)$$

= $M_1(1_{p_1}) \otimes_k \cdots \otimes_k M_n(1_{p_n})$, with $1_{p_i} = \begin{cases} 0, \text{ if } p_i = \mathbf{0} \\ 1, \text{ if } p_i = \mathbf{1} \end{cases}$
= $(\pi_{\bigstar}M_1)_{p_1} \otimes_k \cdots \otimes_k (\pi_{\bigstar}M_n)_{p_n} = [(\pi_{\bigstar}M_1) \boxtimes \cdots \boxtimes (\pi_{\bigstar}M_n)]_p.$

Since π^{\star} and π_{\star} are exact functors, they induce functors in the derived categories

$$\pi^{\bigstar} : D(\mathbb{A}^n_{\mathbb{F}_1}) \to D(R) \quad , \quad \pi_{\bigstar} : D(R) \to D(\mathbb{A}^n_{\mathbb{F}_1})$$

which are still adjoint.

2. Main theorem and applications

For each $M \in D(R)$, let us denote

$$D(M) = \mathbb{R} \operatorname{Hom}_{R}^{\bullet}(M, \Omega_{R/k}^{n}).$$

Recall that for any $F \in D(\mathbb{A}^n_{\mathbb{F}_1})$, D(F) denotes the complex $\mathbb{RHom}^{\bullet}_{\mathbb{A}^n_{\mathbb{F}_1}}(F, k_{\{1\}})$. Then:

Theorem 2.1. For any $M \in D(R)$, and any pointwise perfect $F \in D_{perf}(\mathbb{A}^n_{F_1})$, one has

$$\mathbb{R}\operatorname{Hom}_{R}^{\bullet}(M, \pi^{\star}F) = \mathbb{R}\operatorname{Hom}_{\mathbb{A}_{\mathsf{F}_{*}}^{n}}^{\bullet}(D(F), \pi_{\star}D(M)).$$

Proof. Let us first see that $\pi^* F$ is a perfect complex of *R*-modules. Let us consider the standard resolution (see [3])

$$0 \to C_n F \to \dots \to C_1 F \to C_0 F \to F \to 0$$

with $C_i F = \bigoplus_{p_0 < \cdots < p_i} (F_{p_0}) \otimes_k k_{U_{p_i}}$. If suffices to see that $\pi^{\star} C_i F$ is perfect. Hence, we are reduced to prove that $\pi^{\star} (E \otimes_k k_{U_p})$ is perfect, for any $E \in D(k)$ perfect and $p \in \mathbb{A}^n_{F_1}$. In this case, $\pi^{\star} (E \otimes_k k_{U_p}) = E \otimes_k \pi^{\star} (k_{U_p}) = E \otimes_k (\mathbf{x}^p)$ is perfect.

Now, since $\pi^{\star}F$ is perfect and $\Omega_{R/k}^{n}$ is an invertible *R*-module, one has

$$\mathbb{R}\operatorname{Hom}_{B}^{\bullet}(M, \pi^{\star}F) = \mathbb{R}\operatorname{Hom}_{B}^{\bullet}(D(\pi^{\star}F), D(M)).$$

Moreover, $D(\pi^{\star}F) = \pi^{\star}D(F)$ (see [5, Thm 5.1]). Then

$$\mathbb{R}\operatorname{Hom}^{\bullet}_{R}(M, \pi^{\star}F) = \mathbb{R}\operatorname{Hom}^{\bullet}_{R}(\pi^{\star}D(F), D(M)) = \mathbb{R}\operatorname{Hom}^{\bullet}_{\mathbb{A}^{n}_{F_{1}}}(D(F), \pi_{\star}D(M)).$$

Remark 2.2. As we shall see, Hochster's formulae appear when one is able to decompose $\pi_{\star}D(M)$, and this highly depends on of the choice of M. A first example will be Theorem 2.4.

Lemma 2.3. Let J be a sheaf of k-vector spaces on a finite poset X. If J is injective, then it decomposes as

$$J \simeq \bigoplus_{x \in X} J(x) \otimes_k k_{C_x},$$

with $J(x) = \operatorname{Hom}_X(k_{\{x\}}, J)$.

Proof. We proceed by induction on r = number of points of X. If r = 1, it is immediate. Assume r > 1, let $x_0 \in X$ be a closed point and $U := X - \{x_0\} \stackrel{j}{\hookrightarrow} X$. Taking $\mathcal{H}om_X(\ , J)$ in the exact sequence $0 \to k_U \to k \to k_{\{x_0\}} \to 0$ (where $\mathcal{H}om_X$ is the sheaf of homomorphisms), we obtain

$$0 \to J(x_0) \otimes_k k_{\{x_0\}} \to J \to j_* J_{|U} \to 0.$$

Since $J(x_0) \otimes_k k_{\{x_0\}}$ is injective, this sequence splits, so $J \simeq J(x_0) \otimes_k k_{\{x_0\}} \oplus j_* J_{|U}$. One concludes by induction because $J_{|U}$ is an injective sheaf on U.

Theorem 2.4. Assume that k is a field. Let $M \in D(R)$ such that D(M) = I[r] for some injective R-module I and some integer r. Then, for any pointwise perfect $F \in D_{perf}(\mathbb{A}^n_{F_1})$ one has a (non-canonical) decomposition

$$\operatorname{Ext}_{R}^{i}(M, \pi^{\star}F) \simeq \bigoplus_{p \in \overline{\operatorname{supp}(F)}} H_{p}^{i+n+r-c_{p}}(U_{p}, F) \otimes_{k} \operatorname{Hom}_{R}(R_{C_{p}}, I)(1_{p}), \qquad c_{p} = \dim C_{p}.$$

If M is \mathbb{Z}^n -graded, this decomposition is an isomorphism of \mathbb{Z}^n -graded k-vector spaces. Proof. Since π^* is exact, $J := \pi_* I$ is an injective sheaf on $\mathbb{A}^n_{\mathsf{F}_1}$, so it decomposes (by Lemma 2.3)

$$J \simeq \bigoplus_{p \in \mathbb{A}_{\mathbb{F}_1}^n} J(p) \otimes_k k_{C_p}$$

with $J(p) = H_p^0(U_p, J) = \operatorname{Hom}_{\mathbb{A}_{F_1}^n}(k_{\{p\}}, J)$. Then

$$\mathbb{R}\operatorname{Hom}_{R}^{\bullet}(\mathcal{M}, \pi^{\star}F) \stackrel{2.1}{=} \mathbb{R}\operatorname{Hom}_{\mathbb{A}_{F_{1}}^{n}}(D(F), \pi_{\star}D(M)) = \mathbb{R}\operatorname{Hom}_{\mathbb{A}_{F_{1}}^{n}}(D(F), \pi_{\star}I)[r]$$
$$\simeq \bigoplus_{p \in \mathbb{A}_{F_{1}}^{n}} \mathbb{R}\operatorname{Hom}_{\mathbb{A}_{F_{1}}^{n}}^{\bullet}(D(F), J(p) \otimes_{k} k_{C_{p}})[r].$$

Now, since k is a field, $\overset{\mathbb{L}}{\otimes}_{k} = \bigotimes_{k}$, and then $\mathbb{R} \operatorname{Hom}_{\mathbb{A}_{F_{1}}^{n}}^{\bullet}(D(F), J(p) \otimes_{k} k_{C_{p}}) \overset{1.3.1}{=} \mathbb{R}\Gamma_{p}(U_{p}, F)[n - c_{p}] \otimes_{k} J(p)$, which is zero if $p \notin \overline{\operatorname{supp}(F)}$. To conclude, it suffices to see that $J(p) = \operatorname{Hom}_{R}(R_{C_{p}}, I)(1_{p})$. By definition

$$J(p) = \operatorname{Hom}_{\mathbb{A}_{F_1}^n}(k_{\{p\}}, J) = \operatorname{Hom}_R(\pi^{\star}k_{\{p\}}, I).$$

Applying π^{\star} to the exact sequence

$$\bigoplus_{i \notin p} k_{U_{p \cup \{i\}}} \to k_{U_p} \to k_{\{p\}} \to 0$$

one obtains that $\pi^{\star}k_{\{p\}} = R_{C_p}(-1_p)$, and then $J(p) = \operatorname{Hom}_R(R_{C_p}, I)(1_p)$.

Proposition 2.5. Let
$$R = k[x_1, \ldots, x_n]$$
, $\mathfrak{m} = (x_1, \ldots, x_n)$ and $\mathfrak{m}_l = (x_1^l, \ldots, x_n^l)$. One has
(a) $\pi_{\star} H^n_{\mathfrak{m}}(\Omega^n_{R/k}) = \bigoplus_{p \in \mathbb{A}^n_{\mathbb{F}_1}} H^{c_p}_{\mathfrak{m}}(R_{C_p}) \otimes_k k_{C_p}$, with $c_p = \dim C_p$.
(b) $\pi_{\star} \operatorname{Hom}_R(R/\mathfrak{m}_l, H^n_{\mathfrak{m}}(\Omega^n_{R/k})) = \bigoplus_{q \ge p \in \mathbb{A}^n_{\mathbb{F}_1}} (R_{C_p}/\mathfrak{m}_{l-1}) \otimes_k k_{C_q} \otimes_k k_{U_{q-p}}(l_q)$.

Proof. For n = 1 it is a direct computation. The general case follows from the case n = 1 and the additivity of π_{\star} (Proposition 1.5). Let us give the details.

(a). Let us first consider the case n = 1. In this case,

$$R = k[x], \quad \mathfrak{m} = (x), \quad I := H^1_{\mathfrak{m}}(\Omega_{k[x]/k}) = k[x^{-1}] \frac{\mathrm{d}x}{x}.$$

One has a canonical isomorphism

$$k = \operatorname{Hom}_R(R/\mathfrak{m}, I), \quad 1 \mapsto \frac{\mathrm{d}\,x}{x}$$

and the natural inclusion $k \hookrightarrow I$ has a retract, the residue map:

$$\operatorname{Res}_0 \colon I \longrightarrow k$$
$$\omega \mapsto \operatorname{Res}_0(\omega).$$

Now, $\pi_{\star}I$ is the sheaf on $\mathbb{A}^1_{\mathbf{F}_1}$ given by

$$(\pi_{\bigstar}I)_{\mathbf{0}} = I, \quad (\pi_{\bigstar}I)_{\mathbf{1}} = \operatorname{Hom}_{R}((x), I) = I(1)$$

and the morphism $I = (\pi_{\star}I)_{\mathbf{0}} \to (\pi_{\star}I)_{\mathbf{1}} = I(1)$ is the multiplication by x. Taking $\mathcal{H}om_{\mathbb{A}^n_{\mathbb{F}_1}}(\ , \pi_{\star}I)$ in the exact sequence of sheaves on $\mathbb{A}^1_{\mathbb{F}_1}$

$$0 \to k_{\{\mathbf{1}\}} \to k \to k_{\{\mathbf{0}\}} \to 0$$

we obtain the exact sequence

$$0 \to k_{\{\mathbf{0}\}} \to \pi_{\bigstar} I \to I(1) \to 0 \tag{2.5.1}$$

where I(1) is the constant sheaf I(1) on $\mathbb{A}^1_{\mathbb{F}_1}$; indeed, let us first see that $\mathcal{H}om_{\mathbb{A}^n_{\mathbb{F}_1}}(k_{\{0\}}, \pi_{\star}I) = k_{\{0\}}$. Since $\mathcal{H}om_{\mathbb{A}^n_{\mathbb{F}_1}}(k_{\{0\}}, \pi_{\star}I)$ is supported at the closed point **0**, one has $\mathcal{H}om_{\mathbb{A}^n_{\mathbb{F}_1}}(k_{\{0\}}, \pi_{\star}I) = M \otimes_k k_{\{0\}}$ for some k-module M. Taking the stalk at **0** (i.e., global sections), one obtains $M = \operatorname{Hom}_{\mathbb{A}^n_{\mathbb{F}_1}}(k_{\{0\}}, \pi_{\star}I) = \operatorname{Hom}_R(\pi^{\star}k_{\{0\}}, I) = \operatorname{Hom}_R(R/\mathfrak{m}, I) = k$. Now, let us see that $\mathcal{H}om_{\mathbb{A}^n_{\mathbb{F}_1}}(k_{\{1\}}, \pi_{\star}I) = I(1)$. Let us denote $j: \{1\} \hookrightarrow \mathbb{A}^1_{\mathbb{F}_1}$. Then

$$\mathcal{H}om_{\mathbb{A}_{\mathbb{F}_{1}}^{n}}(k_{\{\mathbf{1}\}}, \pi_{\star}I) = j_{*}j^{-1}\pi_{\star}I = j_{*}(\pi_{\star}I)_{\mathbf{1}} = j_{*}I(1) = I(1).$$

The sequence (2.5.1) splits canonically, since a retract of $k_{\{0\}} \to \pi_{\star} I$ is given by the morphism $\pi_{\star} I \to k_{\{0\}}$ corresponding to the residue map Res₀: $I \to k$ under the equality

$$\begin{split} \operatorname{Hom}_{\mathbb{A}_{\mathbb{F}_{1}}^{1}}(\pi_{\bigstar}I,k_{\{\mathbf{0}\}}) &= \operatorname{Hom}_{k}(I,k) \\ f \mapsto f_{\mathbf{0}} &= \Gamma(\mathbb{A}_{\mathbb{F}_{1}}^{1},f) \end{split}$$

Also, the section of $\pi_{\star}I \to I(1)$ is the morphism $I(1) \to \pi_{\star}I$ corresponding to the morphism $I(1) \xrightarrow{\cdot 1/x} I$ under the equality

$$\operatorname{Hom}_{\mathbb{A}_{\mathbb{F}_{1}}^{1}}(I(1), \pi_{\star}I) = \operatorname{Hom}_{k}(I(1), I)$$
$$f \mapsto f_{\mathbf{0}} = \Gamma(\mathbb{A}_{\mathbb{F}_{1}}^{1}, f)$$

Since the sequence splits, one has: $\pi_{\star}I = k_{\{0\}} \oplus I(1)$ and (a) is proved for n = 1 (notice that $I(1) = H^1_{\mathfrak{m}}(R)$, since $\Omega_{R/k}(1) = R$).

The general case of (a) follows from the case n = 1 by additivity, as follows: now, $R = k[x_1, \ldots, x_n]$ and one has an isomorphism

$$H^n_{\mathfrak{m}}(\Omega^n_{R/k}) = H^1_{(x_1)}(\Omega_{k[x_1]/k}) \otimes_k \cdots \otimes_k H^1_{(x_n)}(\Omega_{k[x_n]/k}))$$

and then (Proposition 1.5)

$$\pi_{\bigstar} H^n_{\mathfrak{m}}(\Omega^n_{R/k}) = (k_{\{\mathbf{0}\}} \oplus H^1_{(x_1)}(k[x_1])) \boxtimes \cdots \boxtimes (k_{\{\mathbf{0}\}} \oplus H^1_{(x_n)}(k[x_n])).$$

This yields the result, taking into account that

$$k_{C_p} = \bigotimes_{i \notin p} k_{H_i} = \bigotimes_{i \notin p} \phi_i^{-1} k_{\{\mathbf{0}\}}, \qquad \phi_i \colon \mathbb{A}^n_{\mathbf{F}_1} \to \mathbb{A}^1_{\mathbf{F}_1} \text{ the } i\text{-th projection}$$

and

$$H^{c_p}_{\mathfrak{m}}(R_{C_p}) = \bigotimes_{i \in p} H^1_{(x_i)}(k[x_i]).$$

(b). Again, let us first consider the case n = 1. In this case, R = k[x], $\mathfrak{m}_l = (x^l)$ and we denote

$$I_l := \operatorname{Hom}_R(R/\mathfrak{m}_l, H^1_\mathfrak{m}(\Omega_{R/k})) = (R/\mathfrak{m}_l)^* = k \frac{\mathrm{d}\,x}{x} \oplus \cdots \oplus k \frac{\mathrm{d}\,x}{x^l}$$

The sheaf $\pi_{\star}I_l$ on $\mathbb{A}^1_{\mathbb{F}_1}$ is described by

$$(\pi_{\star}I_l)_{\mathbf{0}} = I_l, \quad (\pi_{\star}I_l)_{\mathbf{1}} = I_l(1), \quad I_l = (\pi_{\star}I)_{\mathbf{0}} \stackrel{\cdot x}{\to} (\pi_{\star}I)_{\mathbf{1}} = I_l(1).$$

Let us see that

$$\pi_{\star}I_l = k_{\{\mathbf{0}\}} \oplus k_{\{\mathbf{1}\}}(l) \oplus (R/\mathfrak{m}_{l-1})(l).$$

Taking $\mathcal{H}om_{\mathbb{A}_{\mathbf{F}_{1}}^{n}}(\ ,\pi_{\star}I_{l})$ in the epimorphism $k \to k_{\{\mathbf{0}\}}$ one obtains (as in the proof of 2.5) $0 \to k_{\{\mathbf{0}\}} \to \pi_{\star}I_{l}$, which has a retract (again, given by the residue map), and then $\pi_{\star}I_{l} = k_{\{\mathbf{0}\}} \oplus C$, with $C = \pi_{\star}I_{l}/k_{\{\mathbf{0}\}}$. The sheaf C is described as

$$C_0 = I_l / I_1 = I_{l-1}(1), \quad C_1 = I_l(1)$$

and the morphism $C_0 \xrightarrow{ix} C_1$ is injective, with cokernel $I_l(1)/x \cdot I_{l-1} = I_1(l) = k(l)$. Hence, we have an exact sequence

$$0 \to I_{l-1}(1) \to C \to k_{\{\mathbf{1}\}}(l) \to 0$$

which splits, because one has a section $k_{\{1\}}(l) \to C$, induced by the morphism $k(l) \to C_1$, $1 \mapsto (\mathrm{d} x)/x^l$. This ends the proof of (b) for n = 1, since $I_{l-1}(1) = (R/\mathfrak{m}_{l-1})(l)$. As before, the general case is obtained by additivity, as follows.

Now, $R = k[x_1, \ldots, x_n]$. Then

$$(R/\mathfrak{m}_l)^* = (k[x_1]/(x_1^l))^* \otimes_k \cdots \otimes_k (k[x_n]/(x_n^l))^*$$

hence,

$$\pi_{\star}(R/\mathfrak{m}_{l})^{*} = \bigotimes_{i=1}^{n} \phi_{i}^{-1}[k_{\{\mathbf{0}\}} \oplus k_{\{\mathbf{1}\}}(l) \oplus (k[x_{i}]/(x_{i}^{l-1})(l)], \quad \phi_{i} \colon \mathbb{A}_{F_{1}}^{n} \to \mathbb{A}_{F_{1}}^{1} \text{ the } i\text{-th projection},$$
$$= \bigoplus_{p \in \mathbb{A}_{F_{1}}^{n}} \left[\bigotimes_{i \in \widehat{p}} \left(k_{H_{i}} \oplus k_{U_{i}}(l) \right) \bigotimes_{i \in p} \left(k[x_{i}]/(x_{i}^{l-1}) \right)(l) \right].$$

Now,

$$\bigotimes_{i \in p} \left(k[x_i] / (x_i^{l-1}) \right)(l) = (R_{C_p} / \mathfrak{m}_{l-1})(l_p)$$

and

$$\bigotimes_{i\in\widehat{p}} (k_{H_i} \oplus k_{U_i}(l)) = \bigoplus_{q\le\widehat{p}} \left[\bigotimes_{i\in q} k_{H_i} \otimes_k \bigotimes_{i\in\widehat{p}-q} k_{U_i}(l) \right] = \bigoplus_{q\le\widehat{p}} k_{C_{\widehat{q}}} \otimes_k k_{U_{\widehat{p}-q}}(l_{\widehat{p}-q}) \\ = \bigoplus_{q\ge p} k_{C_q} \otimes_k k_{U_{q-p}}(l_{q-p})$$

This ends the proof of (b).

If k is a field and F is a sheaf of finite dimensional k-vector spaces on $\mathbb{A}_{F_1}^n$, Yanagawa computes (in [7, Thm. 3.3, Thm. 3.5]) the degree-wise components of $H^i_{\mathfrak{m}}(\pi^{\star}F)$ as cohomology groups with compact support of a certain sheaf $(\pi^{\star}F)^+$ on the geometric realization of $\mathbb{A}_{F_1}^n$. The next theorem generalizes Yanagawas's results for any ring k and any pointwise perfect complex F on $\mathbb{A}_{F_1}^n$, but computing $H^i_{\mathfrak{m}}(\pi^{\star}F)$ in terms of local cohomology groups of F (that is, we do not go through the geometric realization). See also [2].

Theorem 2.6. For any pointwise perfect $F \in D_{perf}(\mathbb{A}^n_{F_1})$, one has a canonical decomposition of \mathbb{Z}^n -graded k-modules:

$$H^{i}_{\mathfrak{m}}(\pi^{\star}F) = \bigoplus_{p \in \overline{\operatorname{supp}(F)}} H^{i-c_{p}}_{p}(U_{p},F) \otimes_{k} H^{c_{p}}_{\mathfrak{m}}(R_{C_{p}}), \quad c_{p} = \dim C_{p}$$

Taking $F = k_K$, with K a closed subset of $\mathbb{A}^n_{\mathbf{F}_1}$, one obtains Hochster's formula

$$H^{i}_{\mathfrak{m}}(R_{K}) = \bigoplus_{p \in K} H^{i-c_{p}-1}_{\mathrm{red}}(\mathrm{Link}_{K}(p), k) \otimes_{k} H^{c_{p}}_{\mathfrak{m}}(R_{C_{p}}).$$

Proof. By Theorem 2.1 one has

$$\mathbb{R}\operatorname{Hom}_{R}^{\bullet}(R/\mathfrak{m}^{l},\pi^{\star}F) = \mathbb{R}\operatorname{Hom}_{\mathbb{A}_{\mathbb{F}_{1}}^{n}}^{\bullet}(D(F),\pi_{\star}D(R/\mathfrak{m}^{l})).$$

Taking direct limit on l, one obtains

$$\mathbb{R}\Gamma_{\mathfrak{m}}(\pi^{\star}F) = \mathbb{R}\operatorname{Hom}_{\mathbb{A}_{F_{1}}^{n}}^{\bullet}(D(F), \pi_{\star}H_{\mathfrak{m}}^{n}(\Omega_{R/k}^{n}))[-n].$$
(2.6.1)

Indeed,

$$\mathbb{R}\Gamma_{\mathfrak{m}}(\pi^{\star}F) = \lim_{\substack{\to\\l}} \mathbb{R}\operatorname{Hom}_{R}^{\bullet}(R/\mathfrak{m}^{l},\pi^{\star}F) = \lim_{\substack{\to\\l}} \mathbb{R}\operatorname{Hom}_{\mathbb{A}_{F_{1}}^{n}}^{\bullet}(D(F),\pi_{\star}D(R/\mathfrak{m}^{l}))$$
$$= \lim_{\substack{\to\\l}} \mathbb{R}\operatorname{Hom}_{\mathbb{A}_{F_{1}}^{n}}^{\bullet}(\pi^{\star}D(F),D(R/\mathfrak{m}^{l})) \stackrel{\pi^{\star}D(F) \text{ is perfect}}{=} \mathbb{R}\operatorname{Hom}_{\mathbb{A}_{F_{1}}^{n}}^{\bullet}(\pi^{\star}D(F),\lim_{\substack{\to\\l}} D(R/\mathfrak{m}^{l}))$$
$$= \mathbb{R}\operatorname{Hom}_{R}^{\bullet}(\pi^{\star}D(F),\mathbb{R}\Gamma_{\mathfrak{m}}\Omega_{R/k}^{n}) = \mathbb{R}\operatorname{Hom}_{\mathbb{A}_{F_{1}}^{n}}^{\bullet}(D(F),\pi_{\star}\mathbb{R}\Gamma_{\mathfrak{m}}\Omega_{R/k}^{n})$$

and we conclude because $\mathbb{R}\Gamma_{\mathfrak{m}}\Omega_{R/k}^{n} = H_{\mathfrak{m}}^{n}(\Omega_{R/k}^{n})[-n].$

Now, combining (2.6.1) and Proposition 2.5,(a), we obtain

$$\mathbb{R}\Gamma_{\mathfrak{m}}(\pi^{\star}F) = \bigoplus_{p \in \mathbb{A}_{\mathbb{F}_{1}}^{n}} \mathbb{R}\operatorname{Hom}_{\mathbb{A}_{\mathbb{F}_{1}}^{n}}^{\bullet}(D(F), k_{C_{p}} \otimes_{k} H_{\mathfrak{m}}^{c_{p}}(R_{C_{p}}))[-n]$$

and one concludes because

$$\mathbb{R}\operatorname{Hom}_{\mathbb{A}_{F_1}^n}^{\bullet}(D(F), k_{C_p} \otimes_k H_{\mathfrak{m}}^{c_p}(R_{C_p})) \stackrel{(1.3.1)}{=} \mathbb{R}\Gamma_p(U_p, F)[n-c_p] \otimes_k H_{\mathfrak{m}}^{c_p}(R_{C_p}),$$

which is zero if $p \notin \overline{\operatorname{supp}(F)}$. Finally, for $F = k_K$, the formula follows from the equality $H^i_p(U_p, k_K) = H^{i-1}_{\operatorname{red}}(\operatorname{Link}_K(p), k)$. Indeed, let us denote $U^*_p = U_p - \{p\}$ and $\mathbb{R}\Gamma(U_p, \)$ (resp. $\mathbb{R}\Gamma(U^*_p, \)$) the right derived functor of the functor of sections in U_p (resp. in U^*_p); one has the exact triangle

$$\mathbb{R}\Gamma_p(U_p, k_K) \to \mathbb{R}\Gamma(U_p, k_K) \to \mathbb{R}\Gamma(U_p^*, k_K)$$

$$\downarrow \\ k \qquad \mathbb{R}\Gamma(U_p^* \cap K, k)$$

and then $\mathbb{R}\Gamma_p(U_p, k_K) = \mathbb{R}\Gamma_{\mathrm{red}}(U_p^* \cap K, k)[-1]$; one concludes by the homeomorphism $\mathrm{Link}_K(p) \simeq U_p^* \cap K$.

Remarks 2.7. (1) If k is a field, one can immediately compute the Hilbert series of $H^i_{\mathfrak{m}}(\pi^{\star}F)$ from Theorem 2.6, since $H^{c_p}_{\mathfrak{m}}(R_{C_p}) = \bigotimes_{i \in p} \frac{1}{x_i} k[x_i^{-1}]$. In particular, one obtains Hochster's formula for the Hilbert series of the Stanley–Reisner ring R_K ([6, Section II.4]).

(2) The R-module structure of the right hand term

p

$$\bigoplus_{p \in \operatorname{supp}(F)} H_p^{i-c_p}(U_p, F) \otimes_k H_{\mathfrak{m}}^{c_p}(R_{C_p})$$

can be tracked from the proof, as follows. First, let us consider the case n = 1, and the splitting $\pi_{\star}I = k_{\{0\}} \oplus H^1_{(x)}(k[x])$. The multiplication $I \xrightarrow{\cdot x} I$ induces a morphism of sheaves $\pi_{\star}I \xrightarrow{\pi_{\star}(\cdot x)} \pi_{\star}I$, which, via the splitting, becomes the morphism

$$k_{\{0\}} \oplus H^1_{(x)}(k[x]) \to k_{\{0\}} \oplus H^1_{(x)}(k[x])$$

defined by the matrix

$$\begin{pmatrix} 0 & \operatorname{Res}_0 \\ 0 & x \end{pmatrix}$$

where $x \colon H^1_{(x)}(k[x]) \to H^1_{(x)}(k[x])$ is just the multiplication by x and $\operatorname{Res}_0 \colon H^1_{(x)}(k[x]) \to k_{\{0\}}$ is the morphism corresponding to $\operatorname{Res}_0 \colon H^1_{(x)}(k[x]) \to k$, $P \mapsto \operatorname{Res}_0(P \,\mathrm{d} x)$, under the equality $\operatorname{Hom}_{\mathbb{A}^n_{\mathsf{F}_1}}(H^1_{(x)}(k[x]), k_{\{0\}}) = \operatorname{Hom}_k(H^1_{(x)}(k[x]), k).$

Then, the multiplication by x_j in the direct sum $\bigoplus_{p \in \text{supp}(F)} H_p^{i-c_p}(U_p, F) \otimes_k H_{\mathfrak{m}}^{c_p}(R_{C_p})$ is, on

each factor $H_p^{i-c_p}(U_p, F) \otimes_k H_{\mathfrak{m}}^{c_p}(R_{C_p})$, the morphism: (a) If $j \in p$, it is the morphism (let us denote $p' = p - \{j\}$)

$$H_{p}^{i-c_{p}}(U_{p},F) \otimes_{k} H_{\mathfrak{m}}^{c_{p}}(R_{C_{p}}) \longrightarrow H_{p}^{i-c_{p}}(U_{p},F) \otimes_{k} H_{\mathfrak{m}}^{c_{p}}(R_{C_{p}})$$
$$\oplus H_{p'}^{i-c_{p}+1}(U_{p'},F) \otimes_{k} H_{\mathfrak{m}}^{c_{p}-1}(R_{C_{p'}})$$

whose first component is just multiplication by x_j and the second one is the morphism

$$H_p^{i-c_p}(U_p,F) \otimes_k H_{\mathfrak{m}}^{c_p}(R_{C_p}) \xrightarrow{\delta_{pp'}^{i} \otimes \operatorname{Res}_{x_j=0}} H_{p'}^{i-c_p+1}(U_{p'},F) \otimes_k H_{\mathfrak{m}}^{c_p-1}(R_{C_{p'}})$$

where

$$\delta^{i}_{pp'} \colon H^{i-c_p}_p(U_p, F) \to H^{i-c_p+1}_{p'}(U_{p'}, F)$$

is the natural connecting and

$$\operatorname{Res}_{x_j=0} \colon H^{c_p}_{\mathfrak{m}}(R_{C_p}) \to H^{c_p-1}_{\mathfrak{m}}(R_{C_{p'}})$$

is the morphism induced by $\operatorname{Res}_0: H^1_{\mathfrak{m}}(k[x_j]) \to k$, taking into account the equality

$$H^{c_p}_{\mathfrak{m}}(R_{C_p}) = H^{c_p-1}_{\mathfrak{m}}(R_{C_{p'}}) \otimes_k H^1_{\mathfrak{m}}(k[x_j])$$

induced by the equality $R_{C_p} = R_{C_{p'}} \otimes_k k[x_j]$.

(b) The null morphism if $j \notin p$.

In [4, Thm. 1], Miyazaki computes the degree-wise components of $\operatorname{Ext}_{R}^{i}(R/\mathfrak{m}_{l}, R_{K})$ for any closed subset K of $\mathbb{A}_{F_{1}}^{n}$. The next theorem generalizes Miyazaki's result for any pointwise perfect complex F on $\mathbb{A}_{F_{1}}^{n}$. It is a consequence of Theorem 2.1 and Proposition 2.5, (b).

Theorem 2.8. Let us denote $\mathfrak{m}_l = (x_1^l, \ldots, x_n^l)$. For any pointwise perfect $F \in D_{perf}(\mathbb{A}_{F_1}^n)$, one has a canonical decomposition of \mathbb{Z}^n -graded k-modules:

$$\operatorname{Ext}_{R}^{i}(R/\mathfrak{m}_{l},\pi^{\star}F) = \bigoplus_{p \leq q \in \overline{\operatorname{supp}(F)}} H_{q}^{i-c_{q}}(U_{q},F) \otimes_{k} (R_{C_{p}}/\mathfrak{m}_{l-1})(l_{q}).$$

Taking $F = k_K$, with K a closed subset of $\mathbb{A}^n_{\mathbb{F}_1}$, one obtains

$$\operatorname{Ext}_{R}^{i}(R/\mathfrak{m}_{l}, R_{K}) = \bigoplus_{p \leq q \in K} H_{\operatorname{red}}^{i-c_{q}-1}(\operatorname{Link}_{K}(q), k) \otimes_{k} (R_{C_{p}}/\mathfrak{m}_{l-1})(l_{q}),$$

and taking the α -degree component in this equality (with $\alpha \in \mathbb{Z}^n$) one obtains Miyazaki's result ([4, Thm. 1]).

Proof. By Theorem 2.1

$$\mathbb{R}\operatorname{Hom}_{R}^{\bullet}(R/\mathfrak{m}_{l},\pi^{\star}F) = \mathbb{R}\operatorname{Hom}_{\mathbb{A}_{F_{1}}^{n}}^{\bullet}(D(F),\pi_{\star}D(R/\mathfrak{m}_{l})).$$

Now, $D(R/\mathfrak{m}_l) = \operatorname{Hom}_R(R/\mathfrak{m}_l, H^n_\mathfrak{m}(\Omega^n_{R/k}))[-n]$, so

$$\pi_{\bigstar} D(R/\mathfrak{m}_l) = \pi_{\bigstar} \operatorname{Hom}_R(R/\mathfrak{m}_l, H^n_\mathfrak{m}(\Omega^n_{R/k}))[-n].$$

By Proposition 2.5, (b),

$$\mathbb{R}\operatorname{Hom}_{R}^{\bullet}(R/\mathfrak{m}_{l},\pi^{\star}F) = \bigoplus_{q \ge p} \mathbb{R}\operatorname{Hom}_{\mathbb{A}_{\mathbb{F}_{1}}^{n}}^{\bullet}(D(F), k_{C_{q}} \otimes_{k} k_{U_{q-p}} \otimes_{k} (R_{C_{p}}/\mathfrak{m}_{l-1})(l_{q}))[-n]$$
$$= \bigoplus_{q \ge p} \mathbb{R}\operatorname{Hom}_{\mathbb{A}_{\mathbb{F}_{1}}^{n}}^{\bullet}(D(F), k_{C_{q}} \otimes_{k} k_{U_{q-p}}) \otimes_{k} (R_{C_{p}}/\mathfrak{m}_{l-1})(l_{q})[-n]$$

where the last equality follows because $R_{C_p}/\mathfrak{m}_{l-1}$ is a finite and free k-module. Then we are reduced to prove the following formula (for any $x \ge y \in \mathbb{A}^n_{\mathbb{F}_1}$):

$$\mathbb{R}\operatorname{Hom}_{\mathbb{A}_{\mathbb{F}_1}^n}^{\bullet}(D(F), k_{C_x} \otimes_k k_{U_y}) = \mathbb{R}\Gamma_x(U_x, F)[n - c_x] \quad \text{(which is zero if } x \notin \overline{\operatorname{supp}(F)}).$$

Let us denote $j: U_y \hookrightarrow \mathbb{A}^n_{\mathbb{F}_1}, C^y_x$ the closure of x in U_y (in other words, $C^y_x = C_x \cap U_y = j^{-1}(C_x)$). Notice that $U_y = \mathbb{A}^{n-c_y}_{\mathbb{F}_1}, q \mapsto q - y$, and $j^{-1}k_{\{1\}} = k_{\{1\}}$, so $j^{-1}D(F) = D(j^{-1}F)$. Then

$$\mathbb{R} \operatorname{Hom}_{\mathbb{A}_{\mathbb{F}_{1}}^{n}}^{\bullet}(D(F), k_{C_{x}} \otimes_{k} k_{U_{y}}) = \mathbb{R} \operatorname{Hom}_{\mathbb{A}_{\mathbb{F}_{1}}^{n}}^{\bullet}(D(F), j_{!}j^{-1}k_{C_{x}})$$

$$= \mathbb{R} \operatorname{Hom}_{\mathbb{A}_{\mathbb{F}_{1}}^{n-c_{y}}}^{\bullet}(j^{-1}D(F), k_{C_{x}}^{y}) \quad (\text{adjunction between } j^{-1} \text{ and } j_{!})$$

$$= \mathbb{R} \operatorname{Hom}_{\mathbb{A}_{\mathbb{F}_{1}}^{n-c_{y}}}^{\bullet}(D(j^{-1}F), k_{C_{x}}^{y})$$

$$\stackrel{(1.3.1)}{=} \mathbb{R} \Gamma_{x}(U_{x}, F)[n - c_{y} - \dim C_{x}^{y}]$$

and we conclude because dim $C_x^y = c_x - c_y$.

2.1. Comparison with other Hochster's type formulae. Let us compare our results with other Hochster's type formulae obtained more recently with a different approach (see [2] and specially [1]). Let us first consider Hochster's formula for local cohomology. Let K be a finite simplicial complex and R_K its associated Stanley-Reisner ring. The classical Hochster's formula computes $H^i_{\mathfrak{m}}(R_K)$, when k is a field, and Yanagawa generalizes this result for any squarefree module (over a field) and our Theorem 2.6 generalizes Yanagawa's result for any ring k and any complex of squarefree modules. A different approach to compute $H^i_{\mathfrak{m}}(R_K)$, which goes back to Yuzvinsky ([9]), is to see R_K as the global sections of a sheaf of k-algebras on the poset K^{op} : for each $p \in K$, let $R_p = R_{C_p}$ and, for any $p \leq q$, $R_q \to R_p$ the natural epimorphism. We have then a sheaf \mathcal{R}_K on K^{op} , defined by $(\mathcal{R}_K)_p = R_p$. The global sections of \mathcal{R}_K is the inverse limit of the $R_p's$, which amounts to be R_K :

$$\Gamma(K^{\mathrm{op}}, \mathcal{R}_K) = \lim_{\substack{\leftarrow \\ p \in K}} R_p = R_K.$$

From this point of view, Hochster's formula gives a description of the local cohomology of the limit $\lim_{\substack{\leftarrow\\p\in K}} R_p$ in terms of the local cohomology of the $R_p's$. This has been generalized by

replacing the simplicial complex K by an arbitrary poset P and considering a sheaf of k-algebras \mathcal{R} on P (as in [2]) or, more generally, a sheaf \mathcal{R} of A-modules on P (as in [1]). Under certain "degeneracy" conditions, one obtains a Hochster's type result: the local cohomology of the limit $\lim_{\leftarrow} R_p$ is computed in terms of the local cohomology of the $R_p's$. But Yanagawa's result (or our \leftarrow

Theorem 2.6) is not obtained from this perpective.

Regarding our Theorem 2.8, or Miyazaki's result, something similar happens. There are Hochster's type results for Ext's, obtained in [1], again under the perspective of comparing the Ext of a limit $\lim_{\leftarrow} R_p$ with the Ext of the $R_p's$. But these results do not yield Miyazaki's $p \in P$

result or our Theorem 2.8. For example, let \mathfrak{m} be a maximal ideal of a noetherian ring A, J an \mathfrak{m} -primary ideal (for example, $J = \mathfrak{m}^l$) and P_p an inverse system of finitely generated A-modules; then $\operatorname{Ext}_A^i(A/J, P_p)$ is a finitely generated module over the Artinian ring A/J, and then the degeneracy condition of [1, Thm. 7.1]

$\operatorname{Hom}_{A}(\operatorname{Ext}_{A}^{d_{p}}(A/J, P_{p}), \operatorname{Ext}_{A}^{d_{q}}(A/J, P_{q}) = 0$

is never satisfied if both $\operatorname{Ext}_{A}^{d_{p}}(A/J, P_{p})$ and $\operatorname{Ext}_{A}^{d_{q}}(A/J, P_{q})$ are non zero.

Statements and Declarations

Declarations of interest: none.

Data availability statement. Data sharing is not applicable to this article as no datasets were generated or analyzed during the current study.

References

- J. ÁLVAREZ MONTANER, A. F., BOIX AND S. ZARZUELA, On some local cohomology spectral sequences, Int. Math. Res. Not. IMRN(2020), no.19, 6197-6293.
- [2] M. BRUN, W. BRUNS AND T. RÖMER, Cohomology of partially ordered sets and local cohomology of section rings, Adv. Math. 208 (2007), no. 1, 210-235.

- [3] V. CARMONA SÁNCHEZ, C. MAESTRO PÉREZ, F. SANCHO DE SALAS AND J.F.. TORRES SANCHO, Homology and cohomology of finite spaces, J. Pure Appl. Algebra 224, no. 4, 106200, 38 pp. (2020).
- [4] M. MIYAZAKI, Characterizations of Buchsbaum complexes, Manuscripta Math. 63 (1989), no. 2, 245-254.
- [5] F. SANCHO DE SALAS AND A. TORRES SANCHO, A geometrization of Stanley-Reisner theory, arXiv:2303.06940 [math.AG].
- [6] R. STANLEY, Combinatorics and commutative algebra, Second edition. Progress in Mathematics, 41. Birkhäuser Boston, Inc., Boston, MA, 1996. x+164 pp.
- [7] K. YANAGAWA, Stanley-Reisner rings, sheaves, and Poincaré-Verdier duality, Math. Res. Lett. 10 (2003), no. 5-6, 635-650.
- [8] K. YANAGAWA, Sheaves on finite posets and modules over normal semigroup rings, J. Pure Appl. Algebra 161 (2001), no. 3, 341-366.
- [9] S. YUZVINSKY, Cohen-Macaulay rings of sections, Adv. in Math. 63 (1987), no.2, 172-195.

DEPARTAMENTO DE MATEMÁTICAS AND INSTITUTO UNIVERSITARIO DE FÍSICA FUNDAMENTAL Y MATEMÁTICAS (IUFFYM)

Universidad de Salamanca

Plaza de la Merced 1-4, 37008 Salamanca Spain

Email address: fsancho@usal.es Email address: atorressancho@usal.es