Noether's theorem and Ward-Takahashi identities from homotopy algebras

Keisuke Konosu¹ and Jojiro Totsuka-Yoshinaka²

¹Graduate School of Arts and Sciences, The University of Tokyo Komaba, Meguro-ku, Tokyo 153-8902, Japan konosu-keisuke@g.ecc.u-tokyo.ac.jp, ²Department of Physics, Kyoto University, Kitashirakawa, Kyoto 606-8502, Japan george.yoshinaka@gauge.scphys.kyoto-u.ac.jp,

Abstract

We derive the new identity in homotopy algebras which directly corresponds to the Schwinger-Dyson equations in quantum field theory. As an application, we derive the Ward-Takahashi identities. We demonstrate that the Ward-Takahashi identities are reproduced in several examples. In general, our formula contains divergence. We mediate this problem by introducing stubs known in the context of string field theory. With the regularization, we can calculate the anomaly such as axial U(1) anomaly in vector-like U(1) gauge theory.

Contents

1	Introduction A_{∞} algebras and quantum A_{∞} algebras Correlation functions from quantum A_{∞} algebras			1	
2				3	
3				10	
	3.1	Correl	ation functions for scalar field theories	10	
	3.2	Correl	ation functions and the algebraic Schwinger-Dyson equation	15	
	3.3	The al	gebraic derivation of the Schwinger-Dyson equations	17	
4	Noe	ether's	theorem and Ward-Takahashi identity	21	
	4.1	Infinit	esimal symmetry in homotopy algebras	21	
	4.2	4.2 Noether's theorem in homotopy algebra		22	
	4.3 Ward-Takahashi identity		25		
	4.4 Examples			27	
		4.4.1	The $U(1)$ symmetry in complex scalar field theories $\ldots \ldots \ldots \ldots$	27	
		4.4.2	The energy-momentum tensor in scalar field theories	33	
		4.4.3	The chiral $U(1)$ symmetry in vector-like gauge theories $\ldots \ldots \ldots$	36	
5	Reg	Regularization with stubs 4			
6	Conclusions and discussion			46	
\mathbf{A}	A Proof: $(2.14) \Leftrightarrow (2.46)$			48	
в	3 The proof for the Schwinger-Dyson equations for scalar field theory			49	

1 Introduction

Homotopy algebras such as A_{∞} algebras [1, 2, 3, 4, 5, 6] and L_{∞} algebras [7, 8] have contributed in the context of constructing the action of string field theory [7, 9, 10, 11, 12, 13, 14, 15, 16]. The homotopy structure ensures the gauge invariance of the action explicitly. Recently, homotopy algebras have been used for various purposes beyond constructing the action. For instance, we can reproduce the Feynman diagram expansion [17, 18, 19, 20, 21, 22, 23, 24, 25]. We can also relate covariant and light-cone string field theories by homotopy algebras [26]. Notably, they are applicable to not only string field theory but also ordinary quantum field theories [27,28, 29, 30, 31, 32, 33]. Particularly, recent proposals utilizing homotopy algebras to express correlation functions [34, 35, 36] are noteworthy.¹ One advantage of describing quantum field theory using homotopy algebras is that it allows descriptions independent of the details of the theory. For instance, string field theory and scalar field theory have different structures and the former is far more complex than the latter. Expressions based on homotopy algebras, however, formally represent both of the theories in a similar form. While solving string field theory poses significant challenges due to its complexity, various analytical techniques are available in quantum field theory. By translating the properties and analytical methods of quantum field theory into the language of homotopy algebras, we expect that we can directly apply or extend them for applications in string field theory.

Symmetry plays a crucial role in the analysis of quantum field theory. In particular, global symmetry is an important concept related to conserved charges and anomalies. In the context of homotopy algebras, there are several discussions regarding global symmetry [44, 19], but they remain at the classical level. In this paper, we provide an expression of the Ward-Takahashi identity in terms of homotopy algebras, based on the expression of correlation functions given in [34, 35, 36].² To derive this, we use a new identity that we refer to as the algebraic Schwinger-Dyson equation. Moreover, this equation improves the proof of the Schwinger-Dyson equation from the point of the formula for correlation functions in terms of homotopy algebras presented in [34, 35, 36], which enable us to extend to general theories easier. The Ward-Takahashi identity in terms of homotopy algebras we derived contains a term corresponding to an anomaly. This term is generally ill-defined and requires regularization. We applied regularization with stubs [7]. which is widely used in the context of string field theory. In the context of homotopy algebras, research on adding stubs has seen significant development in recent years [46, 47, 48, 49]. By adding stubs, we can perform the heat kernel regularization while preserving the structure of homotopy algebras, making it the optimal regularization for the discussions in this paper. The regularization method we introduce is related to the work [50], in which the anomaly and the regularization with stubs are discussed in the framework of the Batalin-Vilkovisky formalism.³

The rest of the paper is organized as follows. In section 2, we briefly review A_{∞} algebras and quantum A_{∞} algebras. In section 3, we review the formula for correlation functions in terms of homotopy algebras for the scalar field theories [34, 35]. Subsequently, we discuss correlation functions for the general theories and introduce the *algebraic Schwinger-Dyson equation*, which are important in our discussion. In section 4, we provide expressions for Noether's theorem and the Ward-Takahashi identity in terms of homotopy algebras. We then illustrate these expressions with three examples, the U(1) symmetry in complex scalar field theories, the translation

¹In the context of the Batalin-Vilkovisky formalism [37, 38, 39], see [40, 41]. See also [42, 43] for related works.

²Though the approach is different from ours, there is an unpublished work deeply related to our work by Hiroaki Matsunaga [45].

 $^{^{3}}$ The approach is a bit different since we are directly evaluating correlation functions.

symmetry in scalar field theories, and the chiral U(1) symmetry in vector-like gauge theories. In section 5, we explain the regularization with stubs and calculate the values of anomalies in the above examples. As is well known, non-zero anomalies can be calculated only in the third example.

Through the paper, we mostly follow the convention [51] when we describe quantum field theory.

2 A_{∞} algebras and quantum A_{∞} algebras

In this section, we explain A_{∞} algebras and their extensions, commonly referred to as quantum A_{∞} algebras. These algebras are defined within the context of graded vector space⁴ denoted as \mathcal{H} . In our setup, consider the graded vector space spanned by the basis $\{e_a(x), \tilde{e}^a(x)\}$. Here the index *a* represents discrete labels and *x* represents continuous label. In this paper, we associate the discrete label with the types of fields or the indices of Lorentz and gauge groups, and the continuous label is associated with the coordinates of a *d*-dimensional spacetime. We denote the subspace generated by the basis $\{e_a(x)\}$ as \mathcal{H}_1 , and the subspace generated by $\{\tilde{e}^a(x)\}$ as \mathcal{H}_2 . The general element of of \mathcal{H} can be written as

$$F = \sum_{a} \int d^{d}x \left[f^{a}(x) e_{a}(x) + \widetilde{f}_{a}(x) \widetilde{e}^{a}(x) \right], \qquad (2.1)$$

where coefficients f^a and \tilde{f}_a are functions.⁵ The identity map on \mathcal{H} is denoted by \mathbb{I} . The degree is determined by the Grassmann parity and the degree of $\tilde{e}^a(x)$ is defined to be opposite to that of $e_a(x)$. If the basis vector $e_a(x)$ has an even degree, $\tilde{e}^a(x)$ has an odd degree, and vice versa. We assume that the degree of coefficients and basis vectors do not depend on continuous labels x, and denote their degree as deg (f^a) and deg (e_a) . Furthermore, when the element F of \mathcal{H} is homogeneous with respect to the degree, we denote it as deg(F). For the sake of simplicity in notation, we write the sign factor $(-1)^{\text{deg}(f^a)}$ as $(-1)^{f^a}$.

We introduce the three kinds of operator acting on this \mathcal{H} , the degree-odd linear map $Q: \mathcal{H} \to \mathcal{H}$, the degree-odd multilinear map $m_n: \mathcal{H}^{\otimes n} \to \mathcal{H}$ for $n \geq 0$, and the degree-odd symplectic form $\omega: \mathcal{H} \otimes \mathcal{H} \to \mathbb{C}$. These operators are graded commutative with the coefficients. For instance, Q acts as

$$Q(f^{a}(x) e_{a}(x)) = (-1)^{f^{a}} f^{a}(x) Q e_{a}(x).$$
(2.2)

⁴Strictly speaking, this space is not the vector space, but the module when we consider Grassmann odd fields. In this paper, these differences do not contribute to our discussions.

⁵In this paper, we assume that spacetime is flat. On a curved spacetime, it is necessary to replace the integral measure with $\sqrt{g} d^d x$.

The operator Q is defined to satisfy the following conditions:

$$Q^2 = 0 \tag{2.3}$$

$$\omega(F_1, QF_2) = -(-1)^{F_1} \omega(QF_1, F_2).$$
(2.4)

The first condition means that the operator Q is nilpotent, and the second condition is called *cyclicity*. The symplectic form ω is defined for the $e_a(x)$ and $\tilde{e}^a(x)$ as follows:

$$\omega (e_a(x), \tilde{e}^b(y)) = (-1)^{e_a} \delta^b_a \delta(x-y),$$

$$\omega (\tilde{e}^a(x), e_b(y)) = (-1)^{\tilde{e}_a} \delta^a_b \delta(x-y),$$

$$\omega (e_a(x), e_b(y)) = \omega (\tilde{e}^a(x), \tilde{e}^b(y)) = 0.$$
(2.5)

When x = y, the above two equations are defined by taking a limit $x \to y$. As it can be seen from the definition, the symplectic form ω satisfies the graded antisymmetric property:

$$\omega(F_1, F_2) = -(-1)^{F_1 F_2} \omega(F_2, F_1).$$
(2.6)

It is convenient to define the bra notation of the symplectic form $\langle \omega |$ as

$$\langle \omega | F_1 \otimes F_2 = \omega(F_1, F_2).$$
(2.7)

In terms of homotopy algebras, the action is given by

$$S = -\frac{1}{2}\omega\left(\Phi, Q\Phi\right) - \sum_{n=0}^{\infty} \frac{1}{n+1}\omega\left(\Phi, m_n\left(\underbrace{\Phi, \dots, \Phi}_{n}\right)\right), \qquad (2.8)$$

where we used the operators defined above, and Φ represents the degree-even element of \mathcal{H}_1 :

$$\Phi = \sum_{a} \int \mathrm{d}^{d}x \, \phi^{a}(x) \, e_{a}(x) \,, \qquad (2.9)$$

where $\{\phi^a(x)\}\$ are fields. The condition that the element Φ has even degree is required so that the first term on the right-hand side of (2.8) does not vanish by the cyclicity of Q. Since the element Φ has even degree,

$$\deg(\phi^a) = \deg(e_a). \tag{2.10}$$

For the simple descriptions, it is convenient to define M_n by

$$M_1 = Q + m_1, \quad M_n = m_n,$$
 (2.11)

for $n \neq 1$. The action (2.8) is invariant under the gauge transformation of the element Φ given by

$$\delta_{\Lambda}\Phi = \sum_{k=1}^{\infty} \sum_{l=0}^{k-1} M_k(\underbrace{\Phi \otimes \ldots \otimes \Phi}_{l} \otimes \Lambda \otimes \underbrace{\Phi \otimes \ldots \otimes \Phi}_{k-l-1}), \qquad (2.12)$$

where Λ is the degree-odd gauge parameter, if the symplectic form ω and operators M_n satisfy the following two kinds of conditions, which are extensions of the nilpotency and the cyclicity of Q. The former one is called A_{∞} relations.⁶ A few of them are

$$0 = M_1(M_0), (2.13a)$$

$$0 = M_1(M_1(\Phi_1)) + M_2(M_0, \Phi_1) + (-1)^{\Phi_1} M_2(\Phi_1, M_0),$$
(2.13b)

$$0 = M_1(M_2(\Phi_1, \Phi_2)) + (-1)^{\Phi_1} M_2(M_1(\Phi_1), \Phi_2) + (-1)^{\Phi_1} M_2(\Phi_1, M_1(\Phi_2)) + M_2(M_0, \Phi_1, \Phi_2) + (-1)^{\Phi_1} M_2(\Phi_1, M_0, \Phi_2) + (-1)^{\Phi_1 + \Phi_2} M_2(\Phi_1, \Phi_2, M_0), \qquad (2.13c) 0 = M_1(M_3(\Phi_1, \Phi_2, \Phi_3)) + M_2(M_2(\Phi_1, \Phi_2), \Phi_3) + (-1)^{\Phi_1} M_2(\Phi_1, M_2(\Phi_2, \Phi_3)) + M_3(M_1(\Phi_1), \Phi_2, \Phi_3) + (-1)^{\Phi_1} M_3(\Phi_1, M_1(\Phi_2), \Phi_3) + (-1)^{\Phi_1 + \Phi_2} M_3(\Phi_1, \Phi_2, M_1(\Phi_3)) + M_4(M_0, \Phi_1, \Phi_2, \Phi_3) + (-1)^{\Phi_1} M_4(\Phi_1, M_0, \Phi_2\Phi_3)$$

$$+ (-1)^{\Phi_1 + \Phi_2} M_4(\Phi_1, \Phi_2, M_0, \Phi_3) + (-1)^{\Phi_1 + \Phi_2 \Phi_3} M_4(\Phi_1, \Phi_2, \Phi_3, M_0).$$
(2.13d)

Another one is cyclicity of M_n :

$$\omega(\Phi_1, M_n(\Phi_2, \dots, \Phi_{n+1})) = -(-1)^{\Phi_1} \omega(M_n(\Phi_1, \dots, \Phi_n), \Phi_{n+1}).$$
(2.14)

When an algebra satisfies A_{∞} relations, it is called an A_{∞} algebra. Additionally, if it possesses cyclicity, it is referred to as a cyclic A_{∞} algebra.

In order to manipulate A_{∞} algebras, it is convenient to introduce the *coalgebra representa*tion.⁷ In the context of coalgebra representation, we extend the operators on $\mathcal{H}^{\otimes n}$ to those on the tensor algebra:

$$T\mathcal{H} = \mathcal{H}^{\otimes 0} \oplus \mathcal{H} \oplus \mathcal{H}^{\otimes 2} \oplus \mathcal{H}^{\otimes 3} \oplus \cdots .$$
(2.15)

Here $\mathcal{H}^{\otimes 0}$ is isomorphic to \mathbb{C} , and we denote the single basis vector **1**, which satisfies

$$\begin{aligned}
\mathbf{1} \otimes \mathbf{1} &= \mathbf{1}, \\
\mathbf{1} \otimes F &= F \otimes \mathbf{1} = F,
\end{aligned}$$
(2.16)

for $F \in \mathcal{H}$. We denote the projection from $T\mathcal{H}$ to $\mathcal{H}^{\otimes n}$ by π_n . The tensor algebra equips a product $\nabla : T\mathcal{H} \otimes' T\mathcal{H} \to T\mathcal{H}$ defined by

$$\nabla (A \otimes' B) = A \otimes B, \qquad (2.17)$$

⁶More precisely, this relation is called *weak* A_{∞} *relations*, and, in particular, when $M_0 = 0$, it is called A_{∞} relations.

⁷The coalgebra representation of A_{∞} algebras is explained in detail in many papers. See, for example, appendix A of [52] and [20]. We mostly follow the conventions used in these papers and [35].

for $A, B \in T\mathcal{H}$, and a coproduct $\triangle : T\mathcal{H} \to T\mathcal{H} \otimes' T\mathcal{H}$ defined by

for $F_i \in \mathcal{H}$. Here \otimes' is the tensor product for the tensor algebra. The following relationship holds between the projection π_n , the product \bigtriangledown , and the coproduct \bigtriangleup :

$$\pi_n = \bigtriangledown \left(\pi_k \otimes' \pi_l \right) \bigtriangleup, \tag{2.19}$$

for k + l = n. The operator that plays a central role in the coalgebra representation is called the *coderivation*, and it is a linear map $\mathbf{c} : T\mathcal{H} \to T\mathcal{H}$ satisfying

$$\triangle \mathbf{c} = \left(\mathbf{I} \otimes' \mathbf{c} + \mathbf{c} \otimes' \mathbf{I} \right) \triangle, \qquad (2.20)$$

where \mathbf{I} is the identity map on the tensor algebra defined by

$$\mathbf{I} = \mathbf{1} + \mathbb{I} + \mathbb{I}^{\otimes 2} + \mathbb{I}^{\otimes 3} + \cdots .$$
(2.21)

An important property of the coderivation is that the graded commutator of two coderivations also becomes a coderivation. We can easily confirm that the graded commutator for two coderivation \mathbf{c}, \mathbf{c}'

$$[\mathbf{c}, \mathbf{c}'] = \mathbf{c} \, \mathbf{c}' - (-1)^{\mathbf{c}\mathbf{c}'} \, \mathbf{c}' \, \mathbf{c} \,, \qquad (2.22)$$

satisfies the condition (2.20). The degree-odd coderivation \mathbf{M}_n can be constructed from the multilinear map M_n as follows:

$$\mathbf{M}_{n} (F_{1} \otimes \cdots \otimes F_{m}) = 0, \quad \text{for} \quad m < n,$$

$$\mathbf{M}_{n} (F_{1} \otimes \cdots \otimes F_{n}) = M_{n} (F_{1}, \dots, F_{n}),$$

$$\mathbf{M}_{n} (F_{1} \otimes \cdots \otimes F_{m}) = \sum_{k=0}^{m-n} (-1)^{F_{1} + \dots + F_{k}} F_{1} \otimes \dots F_{k} \otimes M_{n} (F_{k+1}, \dots, F_{k+n})$$

$$\otimes F_{k+n+1} \otimes \cdots \otimes F_{m} \quad \text{for} \quad m > n,$$

$$(2.23)$$

where $F_1, \ldots, F_n \in \mathcal{H}$. We denote the sum of \mathbf{M}_n as \mathbf{M} :

$$\mathbf{M} = \sum_{n=1}^{\infty} \mathbf{M}_n \,. \tag{2.24}$$

The multilinear map M_n can be written as

$$M_n = \pi_1 \mathbf{M}_n = \pi_1 \mathbf{M} \pi_n \,. \tag{2.25}$$

It is clear from this construction that, in general, the coderivation \mathbf{c} is completely determined by π_1 -projection $\pi_1 \mathbf{c}$. Under the above notation, the A_{∞} relation can be written as the following simple equation:

$$\pi_1 \mathbf{M}^2 = 0. (2.26)$$

This equation implies

$$\mathbf{M}^2 = 0, \qquad (2.27)$$

i.e., **M** is nilpotent. This is because $\mathbf{M}^2 = \frac{1}{2} [\mathbf{M}, \mathbf{M}]$ is a coderivation and is completely determined by the π_1 -projection. Furthermore, the cyclicity (2.14) can be written as

$$\langle \omega | \pi_2 \mathbf{M}_n = 0. \tag{2.28}$$

In some cases, it is convenient to consider Q and m_1 separately instead of considering M_1 . We introduce degree-odd coderivations \mathbf{Q} and \mathbf{m}_n associated to Q and m_n , respectively. We also define

$$\mathbf{m} = \sum_{n=0}^{\infty} \mathbf{m}_n \,. \tag{2.29}$$

Then, the relation (2.3) is promoted to

$$\mathbf{Q}^2 = 0, \qquad (2.30)$$

and the A_{∞} relation (2.27) is rewritten as

$$(\mathbf{Q} + \mathbf{m})^2 = 0.$$
 (2.31)

We define the group-like element $\frac{1}{1-\Phi}$ on $T\mathcal{H}$ corresponding to the degree even element of Φ of \mathcal{H}_1 that constitutes the action as

$$\frac{1}{1-\Phi} = \mathbf{1} + \Phi + \Phi \otimes \Phi + \dots , \qquad (2.32)$$

which satisfies

$$\Delta \frac{1}{1 - \Phi} = \frac{1}{1 - \Phi} \otimes' \frac{1}{1 - \Phi}.$$
(2.33)

Moreover, as it is clear from the definition, this group element is invariant under the permutation of elements in \mathcal{H} . In other words, for the permutation map defined by $T^{(\sigma)}: T\mathcal{H} \to T\mathcal{H}$

$$T^{(\sigma)}(F_1 \otimes \cdots \otimes F_n) = (-1)^{\epsilon(\sigma)} F_{\sigma(1)} \otimes \cdots \otimes F_{\sigma(n)}, \qquad (2.34)$$

where $\sigma \in S_n$ and the sign factor $(-1)^{\epsilon(\sigma)}$ represents the appropriate sign that arises from the permutation, then

$$T^{(\sigma)} \frac{1}{1 - \Phi} = \frac{1}{1 - \Phi} \,. \tag{2.35}$$

The action (2.8) can be written as

$$S = -\sum_{n=0}^{\infty} \frac{1}{n+1} \langle \omega | \bigtriangledown (\pi_1 \otimes' \pi_1 \mathbf{M}_n) \bigtriangleup \frac{1}{1-\Phi}$$
(2.36)

with the group-like element. When there is no confusion, \bigtriangledown and \triangle are often omitted and the action is simply written as

$$S = -\sum_{n=0}^{\infty} \frac{1}{n+1} \langle \omega | (\mathbb{I} \otimes \pi_1 \mathbf{M}_n) \frac{1}{1-\Phi}.$$
(2.37)

We have presented all the ingredients to describe the quantum field theory at the tree level. The action with cyclic A_{∞} algebra is a solution of the classical master equation in the Batalin-Vilkovisky formalism. At the quantum level, the action must rather be a solution of the quantum master equation. To satisfy the quantum master equation, we need to extend the algebra to the quantum A_{∞} algebra, which is defined by the relation

$$\left(\mathbf{M} + i\hbar \mathbf{U}\right)^2 = 0, \qquad (2.38)$$

or

$$\left(\mathbf{Q} + \mathbf{m} + i\hbar \,\mathbf{U}\right)^2 = 0\,,\tag{2.39}$$

where \hbar is the parameter and physically the Dirac constant. Here, the multilinear map m_n , which is π_1 -projection of the coderivation \mathbf{m}_n , generally differs from the classical case. In this case, the multilinear map m_n can be written as

$$m_n = \sum_{g=0}^{\infty} m_{g,n} ,$$
 (2.40)

in the power series expansion of \hbar , and the zeroth order $m_{0,n}$ coincides with the classical case. Here and in what follows, we use the same notation to denote the operators before and after the \hbar correction. Next, let us explain the operator **U**. The operator **U** is the degree-odd second-order coderivation⁸ promoted by the map U from $\mathcal{H}^{\otimes 0}$ to $\mathcal{H}^{\otimes 2}$ given by

$$U = \sum_{a} \int d^{d}x \left(e_{a}(x) \otimes \tilde{e}^{a}(x) + \tilde{e}^{a}(x) \otimes e_{a}(x) \right).$$
(2.41)

The second-order coderivation \mathbf{U} is constructed by

$$\mathbf{U} \mathbf{1} = U = \sum_{a} \int d^{d}x \left(e_{a}(x) \otimes \widetilde{e}^{a}(x) + \widetilde{e}^{a}(x) \otimes e_{a}(x) \right)$$
$$\mathbf{U} \left(F_{1} \otimes \dots \otimes F_{n} \right) = \sum_{\substack{i,j,k \geq 0 \\ i+j+k=n}} \sum_{a} \int d^{d}x \, (-1)^{F_{1}+\dots+F_{i}+\widetilde{e}_{a}(F_{i+1}+\dots+F_{i+j})}$$
$$\left(F_{1} \otimes \dots \otimes F_{i} \otimes e_{a}(x) \otimes F_{i+1} \otimes \dots \otimes F_{i+j} \\ \otimes \widetilde{e}^{a}(x) \otimes F_{i+j+1} \otimes \dots \otimes F_{i+j+k} \right) + \left(e_{a} \leftrightarrow \widetilde{e}^{a} \right),$$
$$(2.42)$$

for $F_1, \ldots, F_n \in \mathcal{H}$, where $(e_a \leftrightarrow \tilde{e}^a)$ represents the term that the operators e_a and \tilde{e}^a are exchanged in the first term. Unlike usual coderivations (also called the first-order coderivation), **U** is not determined by π_1 -projection : $\pi_1 \mathbf{U} = 0$, but is determined by π_2 -projection : $\pi_2 \mathbf{U} = U$. If we use the coderivations $\mathbf{e}_a(x)$ and $\tilde{\mathbf{e}}^a(x)$ defined by

$$\mathbf{e}_{a}(x) \mathbf{1} = e_{a}(x)$$

$$\mathbf{e}_{a}(x) (F_{1} \otimes \cdots \otimes F_{n}) = \sum_{k=0}^{n} (-1)^{e_{a}(F_{1}+\cdots+F_{k})} F_{1} \otimes \cdots \otimes F_{k} \otimes e_{a}(x) \otimes F_{k+1} \otimes \cdots \otimes F_{n}$$

$$(2.43)$$

for $F_1, \ldots, F_n \in \mathcal{H}$, and similarly for $\tilde{\mathbf{e}}^a(x)$, then the second-order coderivation U can be rewritten as

$$\mathbf{U} = \sum_{a} \int \mathrm{d}^{d}x \, \mathbf{e}_{a}(x) \,\widetilde{\mathbf{e}}^{a}(x) \,. \tag{2.44}$$

An important property of U is that it is normalized by

$$\mathbb{I} = -(\mathbb{I} \otimes \langle \omega |)(U \otimes \mathbb{I})$$

= $\sum_{a} \int d^{d}x \left((-1)^{\tilde{e}^{a}} e_{a}(x) \otimes \langle \omega | (\tilde{e}_{a}(x) \otimes \mathbb{I}) + (-1)^{e_{a}} \tilde{e}^{a}(x) \otimes \langle \omega | (e_{a}(x) \otimes \mathbb{I}) \right),$ ^(2.45)

By using this property, the cyclicity (2.14) can also be expressed as

$$\sum_{a} \int d^{d}x \left[M_{n} \left(e_{a}(x) \otimes \mathbb{I}^{\otimes n-1} \right) \otimes \widetilde{e}^{a}(x) + (-1)^{e_{a}} e_{a}(x) \otimes M_{n} \left(\mathbb{I}^{\otimes n-1} \otimes \widetilde{e}^{a}(x) \right) \right] + \left(e_{a}(x) \leftrightarrow \widetilde{e}^{a}(x) \right) = 0,$$

$$(2.46)$$

⁸For the detailed definition of higher-order coderivation, see [8].

The equivalence of two expression (2.14) and (2.46) is proved in appendix A. The relation (2.38) contains two types of relations. One is called the *main identity*:

$$\pi_1 \left(\mathbf{M} + i\hbar \,\mathbf{U} \,\right)^2 = 0 \,. \tag{2.47}$$

This is an extension of the A_{∞} relations (2.26). In particular, the contribution from $i\hbar \pi_1 \mathbf{M} \mathbf{U}$ can often be canceled by the appropriate renormalization, in which case those satisfying A_{∞} relations (2.26) satisfy this main identity as well. Another one is

$$\sum_{a} \int d^{d}x \left[M_{n} \pi_{n} \mathbf{e}_{a}(x) \otimes \widetilde{e}^{a}(x) + (-1)^{e_{a}} e_{a}(x) \otimes M_{n} \pi_{n} \widetilde{\mathbf{e}}^{a}(x) \right]$$

$$+ (e_{a}(x) \leftrightarrow \widetilde{e}^{a}(x)) = 0, \qquad (2.48)$$

which is derived from $\pi_2(\mathbf{M} + i\hbar \mathbf{U})^2 = 0$. When the multilinear map M_n is commutative, as in the case of C_{∞} algebras or L_{∞} algebras, this relation is equivalent to the cyclicity (2.46). In the case of general A_{∞} algebras, this relation does not imply the cyclicity. However, in this paper, we assume the cyclicity of M_n .

3 Correlation functions from quantum A_{∞} algebras

In this section, we describe correlation functions in terms of quantum A_{∞} algebras. In §3.1, we first review the case for scalar field theories following [34, 35, 36].⁹. In §3.2, we show an important identity, the *algebraic Schwinger-Dyson equation*, which contains the Schwinger-Dyson equations. This identity plays a key role in proving the algebraic version of the Ward-Takahashi identities. In §3.3, we confirm that we can derive the Schwinger-Dyson equation from the algebraic Schwinger-Dyson equation.

3.1 Correlation functions for scalar field theories

In this subsection, we give a brief review on correlation functions for scalar field theories in terms of quantum A_{∞} algebras following [35, 36]. As in §2, we consider the graded vector space

$$\mathcal{H} = \mathcal{H}_1 \oplus \mathcal{H}_2 \,. \tag{3.1}$$

When we consider theories without gauge symmetry, we only need basis vectors corresponding to the spacetime ghost number 0 sector and the counterparts. In scalar field theory, we denote the degree-even basis vector of \mathcal{H}_1 by c(x) and the degree-odd basis vector of \mathcal{H}_2 by d(x).

⁹As mentioned in the introduction, there are several approaches to describe correlation functions. In this paper, we take the approach first described in [34] We mostly follow the convention in [35, 36].

In terms of homotopy algebra, we denote the action by

$$S = -\frac{1}{2}\omega\left(\Phi, Q\Phi\right) - \sum_{n=0}^{\infty} \frac{1}{n+1}\omega\left(\Phi, m_n\left(\Phi, \dots, \Phi\right)\right), \qquad (3.2)$$

where Φ is the degree-even element of \mathcal{H}_1 . The element Φ is expanded as

$$\Phi = \int \mathrm{d}^d x \,\varphi(x) \,c(x) \,, \tag{3.3}$$

where $\varphi(x)$ is the real scalar field. We define the symplectic form ω by

$$\begin{pmatrix} \omega(c(x_1), c(x_2)) & \omega(c(x_1), d(x_2)) \\ \omega(d(x_1), c(x_2)) & \omega(d(x_1), d(x_2)) \end{pmatrix} = \begin{pmatrix} 0 & \delta^d(x_1 - x_2) \\ -\delta^d(x_1 - x_2) & 0 \end{pmatrix}.$$
(3.4)

We define the degree-odd operator Q by

$$Q c(x) = (-\partial^2 + m^2) d(x), \qquad Q d(x) = 0,$$
 (3.5)

where m is the mass of the scalar field. From these definitions, the kinetic term is reproduced in terms of homotopy algebras:

$$-\frac{1}{2}\omega\left(\Phi,Q\Phi\right) = -\frac{1}{2}\int \mathrm{d}^{d}x\left[\partial_{\mu}\varphi(x)\partial^{\mu}\varphi(x) + m^{2}\varphi(x)^{2}\right].$$
(3.6)

Note that Q is niloptent:

$$Q^2 = 0. (3.7)$$

Operators m_n are degree-odd map from $\mathcal{H}^{\otimes n}$ to \mathcal{H} . For example, the cubic interaction in φ^3 theory is described by

$$m_2(c(x_1), c(x_2)) = -\frac{g}{2} \delta^d(x_1 - x_2) d(x_1), \qquad (3.8)$$

where g is the coupling constant. We define $m_2(c(x_1), d(x_2)), m_2(d(x_1), c(x_2))$ and $m_2(d(x_1), d(x_2))$ to be zero. In theories without gauge symmetry, operators m_n are defined to be nonzero only when m_n map from $\mathcal{H}_1^{\otimes n}$ to \mathcal{H}_2 .

Associated to the symplectic form ω , we can define the operator **U** by

$$\mathbf{U} = \int \mathrm{d}^d x \, \mathbf{c}(x) \, \mathbf{d}(x) \,, \tag{3.9}$$

where $\mathbf{c}(x)$ and $\mathbf{d}(x)$ are coderivations given by

$$\pi_{1} \mathbf{c}(x) \mathbf{1} = c(x), \quad \pi_{1} \mathbf{c}(x) \pi_{n} = 0, \pi_{1} \mathbf{d}(x) \mathbf{1} = d(x), \quad \pi_{1} \mathbf{d}(x) \pi_{n} = 0$$
(3.10)

for n > 0. Then, the main identity

$$\pi_1 \left(\mathbf{Q} + \sum_n \mathbf{m}_n + i\hbar \mathbf{U} \right)^2 = 0 \tag{3.11}$$

holds trivially, where \mathbf{Q} and \mathbf{m}_n are coderivations associates with Q and m_n , respectively. Since we can easily confirm that the operators \mathbf{Q} and \mathbf{m}_n meet the cyclicity (2.4) and (2.14), these operators satisfy quantum A_{∞} relations.

Let us consider the description of correlation functions. To extract the physical information from homotopy algebras, the projection operator P onto the subspace of \mathcal{H} plays the key role. As in [34], we take P to be

$$P = 0 \tag{3.12}$$

to calculate correlation functions. Intuitively, the operator $\mathbb{I} - P$ corresponds to the region where we carry out the path integral and the definition (3.12) means that we carry out the full path integration. In the coalgebra representation, the operator **P** associated with P is given by

$$\mathbf{P} = \pi_0 \,. \tag{3.13}$$

Associated to the projector P, we define the contracting homotopy h by

$$Qh + hQ = \mathbb{I} - P$$
, $hP = 0$, $Ph = 0$, $h^2 = 0$. (3.14)

In addition, we impose

$$\omega(F_1, h F_2) = (-1)^{F_1} \omega(h F_1, F_2).$$
(3.15)

In the current case, the second and third definitions in (3.14) are trivially satisfied due to (3.12). Then, we choose

$$h c(x) = 0, \qquad h d(x) = \int d^d y \,\Delta(x - y) \,c(y),$$
(3.16)

where $\Delta(x-y)$ is the Feynman propagator of scalar field theories given by

$$\Delta(x-y) = \int \frac{\mathrm{d}^d k}{(2\pi)^d} \, \frac{e^{ik\,(x-y)}}{k^2 + m^2 - i\epsilon} \,. \tag{3.17}$$

Associated with the contracting homotopy h, we define the operator \mathbf{h} in the coalgebra representation by¹⁰

$$\mathbf{h} = h \,\pi_1 + \sum_{n=2}^{\infty} (\mathbb{I}^{\otimes (n-1)} \otimes h) \,\pi_n \,. \tag{3.18}$$

¹⁰In general, the projector P is involved with the action of **h** on $T\mathcal{H}$. Since the projection operator is defined by (3.13), terms that contain P are trivially dropped.

The definitions (3.14) can be extended in the coalgebra representations by

$$\mathbf{Q}\mathbf{h} + \mathbf{h}\mathbf{Q} = \mathbf{I} - \mathbf{P}, \quad \mathbf{h}\mathbf{P} = 0, \quad \mathbf{P}\mathbf{h} = 0, \quad \mathbf{h}^2 = 0.$$
 (3.19)

Then, we can describe the formula for correlation functions in terms of quantum A_{∞} algebras by operators defined above. The formula is given by

$$\langle \Phi^{\otimes n} \rangle = \pi_n \mathbf{f} \mathbf{1} \,, \tag{3.20}$$

where

$$\Phi^{\otimes n} = \underbrace{\Phi \otimes \Phi \otimes \ldots \otimes \Phi}_{n} \tag{3.21}$$

and

$$\mathbf{f} = \frac{1}{\mathbf{I} + \mathbf{h} \,\mathbf{m} + i\hbar \,\mathbf{h} \,\mathbf{U}} \,. \tag{3.22}$$

The inverse of $\mathbf{I} + \mathbf{h} \mathbf{m} + i\hbar \mathbf{h} \mathbf{U}$ is defined by the formal power series:

$$\frac{1}{\mathbf{I} + \mathbf{h}\,\mathbf{m} + i\hbar\,\mathbf{h}\,\mathbf{U}} = \mathbf{I} + \sum_{n=1}^{\infty} (-1)^n \left(\,\mathbf{h}\,\mathbf{m} + i\hbar\,\mathbf{h}\,\mathbf{U}\,\right)^n.$$
(3.23)

Intuitively, operators π_n , **h**, **m**, and **U** correspond to the number of external lines, propagators, vertices, and loops, respectively in Feynman diagrams. In the formula (3.20), the usual correlation functions in quantum field theory appear as the coefficients of basis vectors:

$$\langle \Phi^{\otimes n} \rangle = \langle \underbrace{\Phi \otimes \Phi \otimes \ldots \otimes \Phi}_{n} \rangle$$

$$= \int d^{d}x_{1} d^{d}x_{2} \dots d^{d}x_{n} \langle \varphi(x_{1}) \varphi(x_{2}) \dots \varphi(x_{n}) \rangle c(x_{1}) \otimes c(x_{2}) \otimes \dots \otimes c(x_{n}) .$$

$$(3.24)$$

We define 11

$$\langle \omega_n(x_1,\ldots,x_n)| = (-1)^n \langle \omega| (d(x_1) \otimes \mathbb{I}) \cdots \langle \omega| (d(x_n) \otimes \mathbb{I}).$$
(3.25)

Then, we can extract ordinary correlation functions in quantum field theory by

$$\langle \varphi(x_1) \cdots \varphi(x_n) \rangle = \langle \omega_n(x_1, \dots, x_n) | \pi_n \mathbf{f} \mathbf{1}.$$
 (3.26)

This is verified by

$$\varphi(x_1) \dots \varphi(x_n) = \langle \omega_n(x_1, \dots, x_n) | \frac{1}{1 - \Phi}, \qquad (3.27)$$

and

$$\Phi^{\otimes n} = \pi_n \frac{1}{1 - \Phi} \,. \tag{3.28}$$

¹¹Notice that we use slightly different notations from the papers [34], [35] and [36]. These two notations are essentially the same.

Another important property of the formula (3.20) is that the quantity $\pi_n \mathbf{f} \mathbf{1}$ is symmetric under the permutations of tensor products. This is confirmed indirectly through (3.20).

As an example, we calculate the two-point function of the free scalar theory. From the formula (3.26), the two-point function for the free scalar theory is given by

$$\langle \varphi(x_1) \varphi(x_2) \rangle = \langle \omega_2(x_1, x_2) | \pi_2 \mathbf{f} \mathbf{1},$$
 (3.29)

where

$$\mathbf{f} = \frac{1}{\mathbf{I} + i\hbar\,\mathbf{h}\,\mathbf{U}}\,.\tag{3.30}$$

In fact, $\pi_2 \mathbf{f} \mathbf{1}$ for the free theory can be simplified as

$$\pi_2 \mathbf{f} \mathbf{1} = -i\hbar \pi_2 \mathbf{h} \mathbf{U} \mathbf{1}. \tag{3.31}$$

Since **U1** is given by

$$\mathbf{U}\,\mathbf{1} = \int d^d x \left(\, c(x) \otimes d(x) + d(x) \otimes c(x) \, \right), \tag{3.32}$$

and the contracting homotopy ${\bf h}$ acts on ${\cal H}^{\otimes 2}$ as

$$\mathbf{h}\,\pi_2 = \left(\,\mathbb{I}\otimes h\,\right)\pi_2\,,\tag{3.33}$$

we obtain

$$\mathbf{h} \mathbf{U} \mathbf{1} = \int \mathrm{d}^d x \left[c(x) \otimes h \, d(x) + d(x) \otimes h \, c(x) \right]. \tag{3.34}$$

Since the contacting homotopy h act on the basis vector c(x) as h c(x) = 0, we obtain

$$\mathbf{h} \mathbf{U} \mathbf{1} = \int \mathrm{d}^d x \, \left[\, c(x) \otimes h \, d(x) \, \right] \,. \tag{3.35}$$

Therefore, we obtain

$$\pi_2 \mathbf{f} \mathbf{1} = -i\hbar \pi_2 \mathbf{h} \mathbf{U} \mathbf{1}$$

= $-i\hbar \int d^d x \int d^d y \left[c(x) \otimes \Delta(x-y) c(y) \right],$ (3.36)

When we substitute the above results into (3.29), we reproduce the result of ordinary quantum field theory:

$$\langle \omega_2 | \pi_2 \mathbf{f} \mathbf{1} = \frac{\hbar}{i} \Delta (x_1 - x_2)$$

= $\langle \varphi(x_1) \varphi(x_2) \rangle .$ (3.37)

In general, the formula (3.20) is verified by showing that correlation functions from (3.20) satisfy the Schwinger-Dyson equations. In [34] and the following papers, it follows from the trivial identity

$$f^{-1} f 1 = 1. (3.38)$$

We present this proof in the Appendix B. In this paper, we prove this fact using the *algebraic* Schwinger-Dyson equations in the next subsection.

3.2 Correlation functions and the algebraic Schwinger-Dyson equation

Let us extend the above discussions of correlation functions for scalar field theory to general gauge-fixed theories. For a gauge-fixed theory, the kinetic term of the action (2.8) is nondegenerate, so it is possible to construct h such that it makes P in (3.14) equal to zero using the propagator. At this time, the vector space \mathcal{H} can be decomposed into two subspaces projected by hQ and Qh, with the former being \mathcal{H}_1 and the latter being \mathcal{H}_2 :

$$h Q e_a(x) = e_a(x), \quad Q h e_a(x) = 0,$$

$$h Q \tilde{e}^a(x) = 0, \quad Q h \tilde{e}^a(x) = \tilde{e}^a(x).$$
(3.39)

Notice that we need to add ghost fields in the gauge-fixed theory. In the homotopy algebraic point of view, we need to add an additional basis associated with the ghost. Then, the basis $\{e_a(x)\}$ contains both degree-even and degree-odd basis vectors in general. Since the element Φ in the action is in \mathcal{H}_1 , this implies

$$h\Phi = 0. (3.40)$$

This corresponds to choosing the gauge fixing condition.

As in the scalar field example above, consider the quantity

$$\mathbf{f} = \frac{1}{\mathbf{I} + \mathbf{h}\,\mathbf{m} + i\hbar\,\mathbf{h}\,\mathbf{U}}\,,\tag{3.41}$$

under P = 0. Here, the second-order coderivation **U** is defined in (2.44). As in the example above, we claim that correlation functions in terms of homotopy algebras for the gauge-fixed field theory are given by

$$\langle \phi^{a_1}(x_1)\phi^{a_2}(x_2)\dots\phi^{a_n}(x_n)\rangle = \langle \omega_n^{a_1,\dots,a_n}(x_1,\dots,x_n)| \ \pi_n \mathbf{f} \mathbf{1},$$
 (3.42)

where the operator $\langle \omega_n^{a_1,\dots,a_n}(x_1,\dots,x_n) |$ is defined by

$$\left\langle \omega_n^{a_1,\dots,a_n}(x_1,\dots,x_n)\right\rangle = (-1)^n \left\langle \omega \right| \left(\widetilde{e}^{a_1}(x_1) \otimes \pi_1\right) \otimes \dots \otimes \left\langle \omega \right| \left(\widetilde{e}^{a_n}(x_n) \otimes \pi_1\right).$$
(3.43)

In the same way as scalar field theory, the formula can also be written as

$$\left\langle \Phi^{\otimes n} \right\rangle = \pi_n \,\mathbf{f} \,\mathbf{1} \,. \tag{3.44}$$

The quantity f 1 satisfies the following important relation:

$$(\mathbf{M} + i\hbar \mathbf{U}) \mathbf{f} \mathbf{1} = (\mathbf{Q} + \mathbf{m} + i\hbar \mathbf{U}) \mathbf{f} \mathbf{1} = 0.$$
(3.45)

We call this identity the *algebraic Schwinger-Dyson equation* because we can derive the wellknown Schwinger-Dyson equations from this identity. The proof is given in the next subsection. In [34] and the following papers, the Schwinger-Dyson equations are obtained from (3.38). This derivation works well, but as it can be seen in the proof described in Appendix B, we need to act the concrete kinetic operators on (3.38) according to the model. In our case, we do not need such an operation and it is written by using only operators in homotopy algebras. Thus, it is more suitable to carry out the proof in general theories. Let us prove (3.45):

$$\begin{aligned} \mathbf{Q} \mathbf{f} \mathbf{1} &= \begin{bmatrix} \mathbf{Q}, \mathbf{f} \end{bmatrix} \mathbf{1} \\ &= \begin{bmatrix} \mathbf{Q}, \frac{1}{\mathbf{I} + \mathbf{h} (\mathbf{m} + i\hbar \mathbf{U})} \end{bmatrix} \mathbf{1} \\ &= -\frac{1}{\mathbf{I} + \mathbf{h} (\mathbf{m} + i\hbar \mathbf{U})} \begin{bmatrix} \mathbf{Q}, \mathbf{h} (\mathbf{m} + i\hbar \mathbf{U}) \end{bmatrix} \frac{1}{\mathbf{I} + \mathbf{h} (\mathbf{m} + i\hbar \mathbf{U})} \mathbf{1} \\ &= -\mathbf{f} (\mathbf{m} + i\hbar \mathbf{U}) \mathbf{f} \mathbf{1} + \mathbf{f} \mathbf{h} \begin{bmatrix} \mathbf{Q}, (\mathbf{m} + i\hbar \mathbf{U}) \end{bmatrix} \mathbf{f} \mathbf{1} \\ &= -\mathbf{f} (\mathbf{m} + i\hbar \mathbf{U}) \mathbf{f} \mathbf{1} - \mathbf{f} \mathbf{h} (\mathbf{m} + i\hbar \mathbf{U}) \end{bmatrix} \mathbf{f} \mathbf{1} \\ &= -\mathbf{f} \begin{bmatrix} \mathbf{I} + \mathbf{h} (\mathbf{m} + i\hbar \mathbf{U}) \end{bmatrix} (\mathbf{m} + i\hbar \mathbf{U}) \mathbf{f} \mathbf{1} \\ &= -\mathbf{f} \begin{bmatrix} \mathbf{I} + \mathbf{h} (\mathbf{m} + i\hbar \mathbf{U}) \end{bmatrix} (\mathbf{m} + i\hbar \mathbf{U}) \mathbf{f} \mathbf{1} \\ &= -(\mathbf{m} + i\hbar \mathbf{U}) \mathbf{f} \mathbf{1} . \end{aligned}$$
(3.46)

We used the quantum A_{∞} relation (2.39) at the fifth equality.

As it can be inferred from (3.44), the quantity **f 1** is an element of $T\mathcal{H}$ that has properties similar to the group-like element $\frac{1}{1-\Phi}$. In fact, it vanishes when **h** acts on it:

$$h f 1 = 0.$$
 (3.47)

In other words, each output is an element of \mathcal{H}_1 . Furthermore, it is invariant under the permutation of elements in \mathcal{H} :

$$T^{(\sigma)} \mathbf{f} \, \mathbf{1} = \mathbf{f} \, \mathbf{1} \,. \tag{3.48}$$

Previously, this property was not proven by the direct computation of $\mathbf{f} \mathbf{1}$, however, this can be verified at each order by using the cyclicity (2.46) of m_n and the formula

$$\sum_{a} \int \mathrm{d}^{d}x \, e_{a}(x) \otimes h \, \widetilde{e}^{a}(x) = \sum_{a} \int \mathrm{d}^{d}x \, (-1)^{e_{a}} h \, \widetilde{e}^{a}(x) \otimes e_{a}(x) \,, \tag{3.49}$$

which is derived as

$$\sum_{a} \int d^{d}x \, e_{a}(x) \otimes h \, \widetilde{e}^{a}(x) = \sum_{a,b} \int d^{d}x \, d^{d}y \, (-1)^{\widetilde{e}^{b}} \, e_{a}(x) \otimes e_{b}(y) \otimes \omega \left(\widetilde{e}^{b}(y), h \, \widetilde{e}^{a}(x) \right)$$
$$= -\sum_{a,b} \int d^{d}x \, d^{d}y \, e_{a}(x) \otimes e_{b}(y) \, \omega \left(\widetilde{e}^{a}(x), h \, \widetilde{e}^{b}(y) \right)$$
$$= \sum_{b} \int d^{d}y \, (-1)^{e_{b}} h \, \widetilde{e}^{b}(y) \otimes e_{b}(y) \,, \qquad (3.50)$$

where we used the normalization (2.45) and the property (3.15) of h. For instance, the \hbar^2 order of $\pi_2 \mathbf{f} \mathbf{1}$ is

$$\hbar^{2} \sum_{a,b} \int d^{d}x \, d^{d}y \left[(-1)^{e_{a}+e_{b}} e_{b}(y) \otimes hm_{3} \left(e_{a}(x), h \, \widetilde{e}^{a}(x), h \, \widetilde{e}^{b}(y) \right) + (-1)^{e_{a}+e_{b}+e_{a}e_{b}} e_{a}(x) \otimes hm_{3} \left(e_{b}(y), h \, \widetilde{e}^{a}(x), h \, \widetilde{e}^{b}(y) \right) + (-1)^{e_{a}+e_{b}} e_{a}(x) \otimes hm_{3} \left(h \, \widetilde{e}^{a}(x), e_{b}(y), h \, \widetilde{e}^{b}(y) \right) \right].$$
(3.51)

Due to the cyclicity of m_n and the property (3.15) of h, we get

$$\hbar^{2} \sum_{a,b} \int \mathrm{d}^{d}x \,\mathrm{d}^{d}y \left[(-1)^{e_{b}} hm_{3} \left(h \,\widetilde{e}^{a}(x), \, e_{b}(y), \, h \,\widetilde{e}^{b}(y) \right) \otimes e_{a}(x) \right. \\ \left. + (-1)^{e_{b}+e_{a}e_{b}} hm_{3} \left(e_{b}(y), \, h \,\widetilde{e}^{a}(x), h \widetilde{e}^{b}(y) \right) \otimes e_{a}(x) \right. \\ \left. + (-1)^{e_{a}} hm_{3} \left(e_{a}(x), \, h \,\widetilde{e}^{a}(x), \, h \widetilde{e}^{b}(y) \right) \otimes e_{b}(y) \right].$$

$$(3.52)$$

This is just a form where the output has been permuted. We can verify (3.48) in each order as well in principle. On the other hand, the action of the coproduct Δ is different:

$$\Delta \mathbf{f} \, \mathbf{1} \neq \mathbf{f} \, \mathbf{1} \otimes' \mathbf{f} \, \mathbf{1} \,. \tag{3.53}$$

This corresponds to the fact that the correlation function is generally not factorized.

3.3 The algebraic derivation of the Schwinger-Dyson equations

Let us show that the Schwinger-Dyson equations can be derived from the algebraic Schwinger-Dyson equation (3.45). In the path integral formalism, correlation functions are defined by

$$\langle \phi^{a_1}(x_1)\phi^{a_2}(x_2)\dots\phi^{a_{n-1}}(x_{n-1})\rangle = \frac{1}{Z}\int \mathcal{D}\phi\,\phi^{a_1}(x_1)\phi^{a_2}(x_2)\dots\phi^{a_{n-1}}(x_{n-1})e^{\frac{i}{\hbar}S}\,,$$
 (3.54)

where

$$Z = \int \mathcal{D}\phi \, e^{\frac{i}{\hbar}S} \,. \tag{3.55}$$

Since

$$\frac{1}{Z} \int \mathcal{D}\phi \, \frac{\delta}{\phi^a(x_n)} \Big[\phi^{a_1}(x_1) \phi^{a_2}(x_2) \dots \phi^{a_{n-1}}(x_{n-1}) e^{\frac{i}{\hbar}S} \Big] = 0 \,, \tag{3.56}$$

the Schwinger-Dyson equations are obtained:

$$\sum_{i=1}^{n-1} (-1)^{\phi^a \sum_{j=1}^i \phi^{a_{j-1}}} \langle \phi^{a_1}(x_1) \dots \phi^{a_{i-1}}(x_{i-1}) \delta_a^{a_i} \delta^d(x_i - x_n) \phi^{a_{i+1}}(x_{i+1}) \dots \phi^{a_n}(x_n) \rangle + \frac{i}{\hbar} (-1)^{\phi^a \sum_{j=1}^{n-1} \phi^{a_j}} \left\langle \phi^{a_1}(x_1) \dots \phi^{a_{n-1}}(x_{n-1}) \frac{\delta S}{\delta \phi^a(x_n)} \right\rangle = 0,$$
(3.57)

where we introduced auxiliary parameter $\phi^{a_0} = 0$.

To show the equivalence between (3.45) and (3.57), let us introduce the operator

$$\pi_n(\mathbb{I}^{\otimes (n-1)} \otimes P_{\tilde{e}^a}), \qquad (3.58)$$

where $P_{\tilde{e}^a}$ is the projection onto the subspace of \mathcal{H}_2 spanned by \tilde{e}^a for fixed label a. For example, in scalar field theory described in §3.1, P_d project onto the space spanned by d(x). Acting the above operator on the algebraic Schwinger-Dyson equation (3.45), we obtain

$$\pi_n(\mathbb{I}^{\otimes (n-1)} \otimes P_{\tilde{e}^a}) \operatorname{\mathbf{M}} \mathbf{f} \, \mathbf{1} + i\hbar \, \pi_n(\mathbb{I}^{\otimes (n-1)} \otimes P_{\tilde{e}^a}) \operatorname{\mathbf{U}} \pi_{n-2} \, \mathbf{f} \, \mathbf{1} = 0.$$
(3.59)

Since

$$\pi_n \mathbf{f} \, \mathbf{1} = \langle \Phi^{\otimes n} \rangle \,, \tag{3.60}$$

we can derive

$$\sum_{k=0}^{\infty} \langle \Phi^{\otimes (n-1)} \otimes P_{\tilde{e}^a} M_k(\Phi \otimes \ldots \otimes \Phi) \rangle + i\hbar \pi_n (\mathbb{I}^{\otimes (n-1)} \otimes P_{\tilde{e}^a}) \mathbf{U} \langle \Phi^{\otimes (n-2)} \rangle = 0.$$
(3.61)

Let us consider the first term on the left-hand side of (3.61). From the relation (2.45), we obtain

$$M(\Phi \otimes \ldots \otimes \Phi) = -\sum_{a} \int d^{d}x \, \tilde{e}^{a}(x) \frac{\delta S}{\delta \phi^{a}(x)} \,.$$
(3.62)

Then, the first term of (3.61) is written as

$$\sum_{k=0}^{\infty} \langle \Phi^{\otimes (n-1)} \otimes P_{\tilde{e}^a} M_k(\Phi \otimes \ldots \otimes \Phi) \rangle = - \left\langle \Phi^{\otimes (n-1)} \otimes \int \mathrm{d}^d y \, \tilde{e}^a(y) \, \frac{\delta S}{\delta \phi^a(y)} \right\rangle . \tag{3.63}$$

Notice that the superscript a is not summed over all the fields, but corresponds to the field labeled by a. Let us next consider the second term of (3.61). We obtain

$$i\hbar \pi_{n} (\mathbb{I}^{\otimes (n-1)} \otimes P_{\widetilde{e}}) \mathbf{U} \langle \Phi^{\otimes (n-2)} \rangle$$

$$= i\hbar \sum_{i=1}^{n-1} \int d^{d}y (\mathbb{I}^{\otimes (i-1)} \otimes e_{a}(y) \otimes \mathbb{I}^{\otimes (n-i-1)} \otimes \widetilde{e}^{a}(y)) \pi_{n-2} \mathbf{f} \mathbf{1}$$

$$= i\hbar \sum_{i=1}^{n-1} \int d^{d}y \langle \Phi^{\otimes (i-1)} \otimes e_{a}(y) \otimes \Phi^{\otimes (n-i-1)} \otimes \widetilde{e}^{a}(y) \rangle.$$
(3.64)

Let us comment on the last line. Since

$$\langle \Phi^{\otimes (n-2)} \rangle = (-1)^{\sigma} \int \mathrm{d}^{d} y_{1} \dots \mathrm{d}^{d} y_{n-2} \langle \phi^{a_{1}}(y_{1}) \dots \phi^{a_{n-2}}(y_{n-2}) \rangle (e_{a_{1}}(y_{1}) \otimes \dots \otimes e_{a_{n-2}}(y_{n-2})),$$
(3.65)

where $(-1)^{\sigma}$ is the appropriate sign factor, the term

$$\left(\mathbb{I}^{\otimes (i-1)} \otimes e_{a}(y) \otimes \mathbb{I}^{\otimes (n-i-1)} \otimes \widetilde{e}^{a}(y)\right) \pi_{n-2} \mathbf{f} \mathbf{1} = \left(\mathbb{I}^{\otimes (i-1)} \otimes e_{a}(y) \otimes \mathbb{I}^{\otimes (n-i-1)} \otimes \widetilde{e}^{a}(y)\right) \left\langle \Phi^{\otimes (n-2)} \right\rangle$$

$$(3.66)$$

should be understood as

$$\int \mathrm{d}^{d} y_{1} \dots \mathrm{d}^{d} y_{n-2} (-1)^{\sigma + (e_{a_{1}} + \dots e_{a_{i-1}}) + \tilde{e}^{a}(e_{a_{i}} + \dots e_{a_{n-2}})} \langle \phi^{a_{1}}(y_{1}) \dots \phi^{a_{n-2}}(y_{n-2}) \rangle \times (e_{a_{1}}(y_{1}) \otimes \dots \otimes e_{a_{i-1}}(y_{i-1}) \otimes e_{a}(y) \otimes e_{a_{i}}(y_{i}) \otimes \dots \otimes \tilde{e}^{a}(y)).$$

$$(3.67)$$

In the last line of (3.64), we carry out the formal calculation:

$$(\mathbb{I}^{\otimes (i-1)} \otimes e_a(y) \otimes \mathbb{I}^{\otimes (n-i-1)} \otimes \widetilde{e}^a(y)) \langle \Phi^{\otimes (n-2)} \rangle = \langle (\mathbb{I}^{\otimes (i-1)} \otimes e_a(y) \otimes \mathbb{I}^{\otimes (n-i-1)} \otimes \widetilde{e}^a(y)) \Phi^{\otimes (n-2)} \rangle = \langle (\Phi^{\otimes (i-1)} \otimes e_a(y) \otimes \Phi^{\otimes (n-i-1)} \otimes \widetilde{e}^a(y)) \rangle .$$

$$(3.68)$$

In fact, we find that the calculation in (3.67) coincides with that of (3.68).

Then, the equation (3.61) becomes

$$-\left\langle \Phi^{\otimes(n-1)} \otimes \int \mathrm{d}^d y \, \tilde{e}^a(y) \, \frac{\delta S}{\delta \phi^a} \right\rangle + i\hbar \, \sum_{i=1}^{n-1} \int \mathrm{d}^d y \, \langle \, \Phi^{\otimes(i-1)} \otimes e_a(y) \otimes \Phi^{\otimes(n-i-1)} \otimes \tilde{e}^a(y) \, \rangle = 0 \,.$$
(3.69)

To extract ordinary correlation functions, we use

$$\langle \omega_n^{a_1,\dots,a_n}(x_1,\dots,x_n)| = (-1)^n \langle \omega | \left(\tilde{e}^{a_1}(x_1) \otimes \pi_1 \right) \otimes \dots \otimes \langle \omega | \left(\tilde{e}^{a_n}(x_n) \otimes \pi_1 \right).$$
(3.70)

defined in (3.43). Let us act the operator

$$\langle \omega_n^{a_1,\dots,a_{n-1}}(x_1,\dots,x_{n-1})|\otimes \langle \omega| \left(e_a(x_n)\otimes \mathbb{I} \right)$$
(3.71)

from the left on the both sides in (3.69). After the action of the operator (3.71) on the first term of the left-hand side, we obtain

$$\langle \omega_n^{a_1,\dots,a_{n-1}}(x_1,\dots,x_{n-1})| \otimes \langle \omega | (e_a(x_n) \otimes \mathbb{I}) \left[-\left\langle \Phi^{\otimes (n-1)} \otimes \int \mathrm{d}^d y \, \widetilde{e}^a(y) \, \frac{\delta S}{\delta \phi^a(y)} \right\rangle \right]$$

$$= -\left\langle \phi^{a_1}(x_1)\dots\phi^{a_{n-1}}(x_{n-1}) \int \mathrm{d}^d y \, \langle \omega | \left(e_a(x_n) \otimes \widetilde{e}^a(y) \, \frac{\delta S}{\delta \phi^a(y)}\right) \right\rangle$$

$$= (-1)^{e^a+1} \left\langle \phi^{a_1}(x_1)\dots\phi^{a_{n-1}}(x_{n-1}) \frac{\delta S}{\delta \phi^a(x_n)} \right\rangle.$$

$$(3.72)$$

Let us next consider the second term of the left-hand side of (3.69) after the action of the

operator (3.71). We obtain

$$\begin{split} \left\langle \omega_{n}^{a_{1},\dots,a_{n-1}}(x_{1},\dots,x_{n-1})\right| \otimes \left\langle \omega\right| \left(e_{a}(x_{n})\otimes\mathbb{I}\right) \\ & \left[i\hbar\sum_{i=1}^{n-1}\int\mathrm{d}^{d}y\,\left\langle \Phi^{\otimes(i-1)}\otimes e_{a}(y)\otimes\Phi^{\otimes(n-i-1)}\otimes\widetilde{e}^{a}(y)\right\rangle\right] \\ &=i\hbar\sum_{i=1}^{n-1}\int\mathrm{d}^{d}y\,\left\langle \omega_{n}^{a_{1},\dots,a_{n-1}}(x_{1},\dots,x_{n-1})\right|\left\langle \Phi^{\otimes(i-1)}\otimes e_{a}(y)\otimes\Phi^{\otimes(n-i-1)}\otimes\left\langle \omega\right|\left(e_{a}(x_{n})\otimes\widetilde{e}^{a}(y)\right)\right\rangle \\ &=\left(-1\right)^{e^{a}}i\hbar\sum_{i=1}^{n-1}\left\langle \omega_{n}^{a_{1},\dots,a_{n-1}}(x_{1},\dots,x_{n-1})\right|\left\langle \Phi^{\otimes(i-1)}\otimes e_{a}(x_{n})\otimes\Phi^{\otimes(n-i-1)}\right\rangle \\ &=\left(-1\right)^{e^{a}}i\hbar\sum_{i=1}^{n-1}(-1)^{e^{a}\sum_{j=i+1}^{n-1}e^{a_{j}}} \\ &\times\left\langle \left[-\left\langle \omega\right|\left(\widetilde{e}^{a_{1}}(x_{1})\otimes\Phi\right)\right]\dots\left[-\left\langle \omega\right|\left(\widetilde{e}^{a_{i}}(x_{i})\otimes e^{a}(x_{n})\right)\dots\left[-\left\langle \omega\right|\left(\widetilde{e}^{a_{n-1}}(x_{n-1})\otimes\Phi\right)\right]\right\rangle \\ &\qquad (3.73) \end{split}$$

In the third line, no sign factor appears since $\langle \omega | (e_a(x_n) \otimes \mathbb{I}) \rangle$ is degree $e^a + 1$ and $e^a(x)$ has degree $e^a(x)$, that is, either $\langle \omega | (e_a(x_n) \text{ or } e^a(x) \rangle$ is degree even. In the fifth line, the sign factor arise due to the exchange of e^a and $\langle \omega | (\tilde{e}^{a_{j-1}}(x_{n-1}) \otimes \mathbb{I}) \rangle$ for j > i. Then, the normalization (2.5) yields

$$i\hbar \sum_{i=1}^{n-1} (-1)^{e^a \sum_{j=i+1}^{n-1} e^{a_j}} \langle \phi^{a_1}(x_1) \dots \delta_a^{a_i} \delta^d(x_i - x_n) \dots \phi^{a_{n-1}}(x_{n-1}) \rangle.$$
(3.74)

Therefore, the equation (3.69) is reduced to

$$(-1)^{e^{a}+1} \left\langle \phi^{a_{1}}(x_{1}) \dots \phi^{a_{n-1}}(x_{n-1}) \frac{\delta S}{\delta \phi^{a}(x_{n})} \right\rangle + i\hbar \sum_{i=1}^{n-1} (-1)^{e^{a} \sum_{j=i+1}^{n-1} e^{a_{j}}} \left\langle \phi^{a_{1}}(x_{1}) \dots \delta_{a}^{a_{i}} \delta^{d}(x_{i}-x) \dots \phi^{a_{n-1}}(x_{n-1}) \right\rangle = 0.$$
(3.75)

By multiplying $(-1)^{e^a \sum_{j=1}^{n-1} e^{a_j}}$ on the above equation, we obtain

$$-\frac{i}{\hbar}(-1)^{e^{a}\sum_{j=1}^{n-1}e^{a_{j}}}\left\langle\phi^{a_{1}}(x_{1})\dots\phi^{a_{n-1}}(x_{n-1})\frac{\delta S}{\delta\phi^{a}(x_{n})}\right\rangle -\sum_{i=1}^{n-1}(-1)^{e^{a}\sum_{j=1}^{i}e^{a_{j-1}}}\langle\phi^{a_{1}}(x_{1})\dots\delta_{a}^{a_{i}}\delta^{d}(x_{i}-x)\dots\phi^{a_{n-1}}(x_{n-1})\rangle = 0,$$
(3.76)

where we used the relation $(-1)^{e^a} = (-1)^{e^a e^{a_i}}$ in the second term of the left-hand side. Since the degree of fields and their associated basis vectors of \mathcal{H}_1 coincides, this is exactly the Schwinger-Dyson equations in quantum field theory.

4 Noether's theorem and Ward-Takahashi identity

In quantum field theory, one of the most important formulas relating to symmetries is the Ward-Takahashi identity, which is given by

$$\langle \phi^{a_1}(x_1)\dots\phi^{a_n}(x_n)\,\partial_\mu j^\mu(x)\,\rangle = -i\hbar\,\sum_{i=1}^n\,\delta^d(x-x_i)\,\langle\,\phi^{a_1}(x_1)\dots\delta\phi^{a_i}(x_i)\dots\phi^{a_n}(x_n)\,\rangle\,.$$
(4.1)

The above Ward-Takahashi identity holds only when no anomaly exists. In the remainder of this paper, we will investigate how this formula is described in terms of homotopy algebras.

4.1 Infinitesimal symmetry in homotopy algebras

In this subsection, we introduce a degree-even multilinear operator $A_n : \mathcal{H}^{\otimes n} \to \mathcal{H}$, which corresponds to an infinitesimal transformation:

$$\delta \Phi = \sum_{a} \int d^{d}x \, \delta \phi^{a}(x) \, e_{a}(x) = A_{0}(\mathbf{1}) + A_{1}(\Phi) + A_{2}(\Phi, \Phi) + \cdots \,. \tag{4.2}$$

We assume that the operator A_n is cyclic:

$$\omega\left(\Phi_{1}, A_{n}\left(\Phi_{2}, \dots, \Phi_{n+1}\right)\right) = -\omega\left(A_{n}\left(\Phi_{1}, \dots, \Phi_{n}\right), \Phi_{n+1}\right), \qquad (4.3)$$

We denote the coderivations promoted from A_n by \mathbf{A}_n . We also introduce the operator

$$\mathbf{A} = \sum_{n=0}^{\infty} \mathbf{A}_n \,. \tag{4.4}$$

Consequently, the transformation (4.2) can be compactly expressed by

$$\delta \Phi = \pi_1 \mathbf{A} \frac{1}{1 - \Phi} \,. \tag{4.5}$$

In the same way, the cyclicity (4.3) is written as

$$\langle \omega | \pi_2 \mathbf{A} = 0. \tag{4.6}$$

The variation of the action (2.8) due to this transformation is given by

$$\delta S = -\sum_{n=0}^{\infty} \frac{1}{n+1} \omega \left(\Phi, \pi_1[\mathbf{M}, \mathbf{A}] \pi_n \frac{1}{1-\Phi} \right).$$

$$(4.7)$$

From this expression, we find that it is sufficient for A to satisfy

$$[\mathbf{M}, \mathbf{A}] = 0. \tag{4.8}$$

for the infinitesimal transformation generated by \mathbf{A} to be the symmetry transformation. Here and in what follows, we only consider a symmetry transformation by \mathbf{A} which satisfies (4.8). Furthermore, we assume that if all of the inputs of A_n are elements of \mathcal{H}_1 , A_n maps to \mathcal{H}_1 . In other words, it satisfies

$$h A_n(\Phi_1 \otimes \Phi_2 \otimes \ldots \otimes \Phi_n) = 0, \qquad (4.9)$$

where Φ_1 , Φ_2 and Φ_n are elements of \mathcal{H}_1 . This assumption implies that this transformation is compatible with the gauge fixing. To define A_n cyclically, it is important to note that nontrivial values should generally be assigned even when only one of the inputs is an element of \mathcal{H}_2 .

4.2 Noether's theorem in homotopy algebra

Before discussing the homotopy description of the Ward-Takahashi identity, let us consider the homotopy description of Noether's theorem. As it is well known, when the action remains invariant under an infinitesimal transformation, a conserved current $j^{\mu}(x)$ associated with the transformation exists. The conserved current satisfies the conservation law:

$$\partial_{\mu} j^{\mu}(x) \approx 0, \qquad (4.10)$$

where we used \approx to represent that the equality holds under the equation of motion. In the ordinary procedure in quantum field theory, one way to obtain the conserved current is to replace the infinitesimal parameter ϵ to $\epsilon(x)$. In this process, the infinitesimal transformation can be modified as follows: For a symmetry transformation $\delta\phi(x)$, let us define $\delta_{\epsilon}\phi(x)$ by

$$\delta_{\epsilon}\phi(x) = \epsilon(x)\,\delta\phi(x)\,,\tag{4.11}$$

where $\epsilon(x)$ is arbitrary function. The conserved current is obtained from the variation of the action under this transformation:

$$\delta_{\epsilon} S = -\int \mathrm{d}^d x \,\epsilon(x) \,\partial_{\mu} \,j^{\mu}(x) \,. \tag{4.12}$$

Let us translate the above procedure in terms of homotopy algebras. As seen in the previous subsection, the symmetry transformation can be described by a degree-even cyclic operator A_n satisfying the condition (4.8). We define a degree-even cyclic operator A_n^{ϵ} with respect to the transformation $\delta_{\epsilon}\phi(x)$ by

$$\delta_{\epsilon}\Phi = \sum_{a} \int \mathrm{d}^{d}x \,\epsilon(x) \,\delta\phi^{a}(x) \,e_{a}(x) = A_{0}^{\epsilon}(\mathbf{1}) + A_{1}^{\epsilon}(\Phi) + A_{2}^{\epsilon}(\Phi,\Phi) + \cdots, \qquad (4.13)$$

The coderivation promoted by A_n^{ϵ} is denoted as \mathbf{A}_n^{ϵ} and we define \mathbf{A}^{ϵ} as

$$\mathbf{A}^{\epsilon} = \sum_{n=0}^{\infty} \mathbf{A}_{n}^{\epsilon} \,. \tag{4.14}$$

If we define the operator \mathcal{O}_{ϵ} as

$$\mathcal{O}_{\epsilon} e_a(x) = \epsilon(x) e_a(x), \quad \mathcal{O}_{\epsilon} \tilde{e}^a(x) = \epsilon(x) \tilde{e}^a(x), \qquad (4.15)$$

then \mathbf{A}_n^{ϵ} can be written as

$$A_n^{\epsilon}(\Phi_1,\ldots,\Phi_n) = \mathcal{O}_{\epsilon} A_n(\Phi_1,\ldots,\Phi_n), \qquad (4.16)$$

on $\mathcal{H}_1^{\otimes n}$. In addition, for this A_n^{ϵ} to be cyclic it should be defined as

$$A_n^{\epsilon}(\Phi_1,\ldots,\widetilde{\Phi},\ldots,\Phi_{n-1}) = A_n(\Phi_1,\ldots,\mathcal{O}_{\epsilon}\widetilde{\Phi},\ldots,\Phi_{n-1}), \qquad (4.17)$$

for $\Phi_1, \ldots, \Phi_{n-1} \in \mathcal{H}_1$ and $\widetilde{\Phi} \in \mathcal{H}_2$. The operator \mathbf{A}^{ϵ} no longer satisfies $[\mathbf{M}, \mathbf{A}^{\epsilon}] = 0$. The variation of the action for the transformation is written as

$$\delta_{\epsilon}S = -\sum_{n=0}^{\infty} \frac{1}{n+1} \omega \left(\Phi, \pi_{1} \left[\mathbf{M}, \mathbf{A}^{\epsilon} \right] \pi_{n} \frac{1}{1-\Phi} \right)$$

$$= -\sum_{n=0}^{\infty} \frac{1}{n+1} \langle \omega | \left(\mathbb{I} \otimes \pi_{1} \left[\mathbf{M}, \mathbf{A}^{\epsilon} \right] \pi_{n} \right) \frac{1}{1-\Phi}.$$
(4.18)

Since this is the counterpart of (4.12) in terms of homotopy algebras, we find

$$\int \mathrm{d}^d x \,\epsilon(x) \,\partial_\mu \, j^\mu(x) = \sum_{n=0}^\infty \, \frac{1}{n+1} \,\langle \omega | \, \left(\,\mathbb{I} \,\otimes\, \pi_1 \left[\,\mathbf{M}, \,\mathbf{A}^\epsilon \,\right] \pi_n \,\right) \frac{1}{1-\Phi} \,. \tag{4.19}$$

The operator \mathcal{O}_{ϵ} can also be written as

$$\mathcal{O}_{\epsilon} = \sum_{a} \int \mathrm{d}^{d} x \,\epsilon(x) \left[(-1)^{\widetilde{e}^{a}} e_{a}(x) \otimes \langle \omega | \, (\widetilde{e}^{a}(x) \otimes \mathbb{I}) + (-1)^{e_{a}} \widetilde{e}^{a}(x) \otimes \langle \omega | \, (e_{a}(x) \otimes \mathbb{I}) \right],$$

$$(4.20)$$

where we used the normalization of U described in (2.45). We can rewrite the right-hand side of the equation (4.19) utilizing (4.20). The cyclicity of operators **M** and **A** allows us to express (4.19) as

$$\int \mathrm{d}^d x \,\epsilon(x) \,\partial_\mu \, j^\mu(x) = \langle \omega | \left(\,\pi_1 \mathbf{A}^\epsilon \otimes \pi_1 \mathbf{M} \,\right) \frac{1}{1 - \Phi} \,. \tag{4.21}$$

Since the action on $\mathcal{H}_1^{\otimes n}$ of $\pi_1 \mathbf{A}^{\epsilon}$ can be written as

$$\pi_1 \mathbf{A}^{\epsilon} = \mathcal{O}_{\epsilon} \, \pi_1 \, \mathbf{A} = \sum_a \int \mathrm{d}^d x \, \epsilon(x) \, (-1)^{\widetilde{e}^a} \, e_a(x) \otimes \langle \omega | \left(\widetilde{e}^a(x) \otimes \pi_1 \mathbf{A} \right), \tag{4.22}$$

substituting this into (4.21) yields

$$\int \mathrm{d}^d x \,\epsilon(x) \,\partial_\mu \, j^\mu(x) = -\sum_a \int \mathrm{d}^d x \,\epsilon(x) \big(\left\langle \omega \right| \left(e_a(x) \otimes \pi_1 \mathbf{M} \right) \otimes \left\langle \omega \right| \left(\widetilde{e}^a(x) \otimes \pi_1 \mathbf{A} \right) \big) \frac{1}{1 - \Phi} \,.$$
(4.23)

Here, we omitted the product and the coproduct. To be more precise, we can write

$$-\sum_{a} \int \mathrm{d}^{d} x \,\epsilon(x) \,\bigtriangledown \left(\left\langle \omega \right| \left(e_{a}(x) \otimes \pi_{1} \mathbf{M} \right) \otimes' \left\langle \omega \right| \left(\widetilde{e}^{a}(x) \otimes \pi_{1} \mathbf{A} \right) \right) \bigtriangleup \frac{1}{1 - \Phi} \,. \tag{4.24}$$

From this expression, we can see that $\partial_{\mu} j^{\mu}(x)$ can be written in terms of homotopy algebras by

$$\partial_{\mu} j^{\mu}(x) = -\sum_{a} \left(\left| \left| \left| \left| e_{a}(x) \otimes \pi_{1} \mathbf{M} \right| \right| \otimes \left| \left| \left| \left| \widetilde{e}^{a}(x) \otimes \pi_{1} \mathbf{A} \right| \right| \right| \right| \right) \frac{1}{1 - \Phi}, \quad (4.25)$$

From the property of group-like element (2.33), it is understood that the right-hand side of (4.24) is proportional to the equation of motion

$$\pi_1 \mathbf{M} \frac{1}{1 - \Phi} = 0.$$
 (4.26)

Then, Noether's theorem is rederived in terms of homotopy algebras:

$$\partial_{\mu} j^{\mu}(x) = 0. \qquad (4.27)$$

Furthermore, integrating the right-hand side and using (2.45), we obtain

$$\langle \omega | (\pi_1 \mathbf{A} \otimes \pi_1 \mathbf{M}) \frac{1}{1 - \Phi}.$$
 (4.28)

This can also be written as

$$\sum_{n=0}^{\infty} \frac{1}{n+1} \left\langle \omega \right| \left(\mathbb{I} \otimes \pi_1 \left[\mathbf{M}, \mathbf{A} \right] \pi_n \right) \frac{1}{1-\Phi} \,. \tag{4.29}$$

Now, since **A** is a symmetry transformation, it commutes with **M**, so this is zero. Therefore, this is consistent with the fact that the right-hand side of (4.25) is a total derivative.

4.3 Ward-Takahashi identity

Now that we have determined the expression corresponding to the derivative of the conserved current $\partial_{\mu} j^{\mu}(x)$, we can proceed to consider the expectation value $\langle \partial_{\mu} j^{\mu}(x) \rangle$. Recall that the equation

$$\left\langle \frac{1}{1-\Phi} \right\rangle = \mathbf{f} \, \mathbf{1} \tag{4.30}$$

holds, it is understood that replacing the group-like element $\frac{1}{1-\Phi}$ with **f** 1 corresponds to taking the expectation value. Hence, the expectation value of $\partial_{\mu} j^{\mu}(x)$ is expressed as

$$\langle \partial_{\mu} j^{\mu}(x) \rangle = -\sum_{a} \left(\langle \omega | (e_{a}(x) \otimes \pi_{1} \mathbf{M}) \otimes \langle \omega | (\widetilde{e}^{a}(x) \otimes \pi_{1} \mathbf{A}) \right) \mathbf{f} \mathbf{1}.$$
(4.31)

This quantity should vanish in the absence of anomalies. When an anomaly exists, it should be non-zero. Let us examine this from the perspective of homotopy algebras. Since the operator \mathbf{M} is the coderivation, the right-hand side of (4.31) can be expressed as

$$\langle \partial_{\mu} j^{\mu}(x) \rangle = -\sum_{a} (-1)^{e_{a}} (\langle \omega | (e_{a}(x) \otimes \pi_{1}) \otimes \langle \omega | (\tilde{e}^{a}(x) \otimes \pi_{1}\mathbf{A})) \mathbf{M} \mathbf{f} \mathbf{1}$$

$$+ \sum_{a} (-1)^{e_{a}} (\langle \omega | (e_{a}(x) \otimes \pi_{1}) \otimes \langle \omega | (\tilde{e}^{a}(x) \otimes \pi_{1}\mathbf{A}\mathbf{M})) \mathbf{f} \mathbf{1}.$$

$$(4.32)$$

Notice that the second term vanishes due to the definition of the symplectic form. Here, using the algebraic Schwinger-Dyson equation (3.45), the first term can be expressed as

$$i\hbar \sum_{a,b} \int \mathrm{d}^d y \, (-1)^{e_a} \big(\langle \omega | \, (e_a(x) \otimes \pi_1) \otimes \langle \omega | \, (\widetilde{e}^a(x) \otimes \pi_1 \mathbf{A}) \, \big) \, \mathbf{e}_b(y) \, \widetilde{\mathbf{e}}_b(y) \, \mathbf{f} \, \mathbf{1} \,. \tag{4.33}$$

Since

$$\langle \omega | (e_a(x) \otimes e_b(y)) = 0, \qquad (4.34)$$

we obtain

$$i\hbar \sum_{a,b} \int \mathrm{d}^d y \, (-1)^{e_a + \widetilde{e}^b \, e_a} \left(\left| \left| \left(e_a(x) \otimes \widetilde{e}_b(y) \right) \otimes \left| \left(\widetilde{e}^a(x) \otimes \pi_1 \, \mathbf{A} \, \mathbf{e}_b(y) \right) \right| \right) \mathbf{f} \, \mathbf{1} \, . \tag{4.35}$$

By the normalization (2.5), we ultimately obtain

$$\langle \partial_{\mu} j^{\mu}(x) \rangle = -i\hbar \sum_{a} \langle \omega | (\pi_1 \mathbf{A} \mathbf{e}_a(x) \otimes \widetilde{e}^a(x)) \mathbf{f} \mathbf{1}.$$
 (4.36)

In general, the right-hand side of this equation includes divergences. For instance, if a symmetry transformation **A** is simply linear and scales by constant, this quantity is proportional to $\delta^d(0)$. Hence, some appropriate regularization is necessary. We will discuss regularization in detail later. From these facts, this quantity is expected to represent anomalies.¹² Indeed, as we will see in section 5, it can be confirmed to coincide with known anomalies in specific examples.

As an extension of (4.19), the expectation value between the fields and $\partial_{\mu} j^{\mu}(x)$ can be written as

$$\langle \phi^{a_1}(x_1) \dots \phi^{a_n}(x_n) \,\partial_\mu \, j^\mu(x) \rangle = -\sum_a \left(\left\langle \omega_n^{a_1,\dots,a_n}(x_1,\dots,x_n) \right| \pi_n \otimes \left\langle \omega \right| \left(e_a(x) \otimes \pi_1 \mathbf{M} \right) \otimes \left\langle \omega \right| \left(\tilde{e}^a(x) \otimes \pi_1 \mathbf{A} \right) \right) \mathbf{f} \, \mathbf{1} \,.$$
 (4.37)

From this expression, we can derive the Ward-Takahashi identity as follows. As before, the coderivation \mathbf{M} can be move to just in front of $\mathbf{f1}$ and using the algebraic Schwinger-Dyson equation (3.45), we obtain

$$i\hbar \sum_{a,b} \int \mathrm{d}^{d} y (-1)^{e_{a}} \left(\left\langle \omega_{n}^{a_{1},\dots,a_{n}}(x_{1},\dots,x_{n}) \right| \pi_{n} \right.$$

$$\otimes \left\langle \omega \right| \left(\left. e_{a}(x) \otimes \pi_{1} \right) \otimes \left\langle \omega \right| \left(\left. \widetilde{e}^{a}(x) \otimes \pi_{1} \mathbf{A} \right) \right) \mathbf{e}_{b}(y) \, \widetilde{\mathbf{e}}^{b}(y) \, \mathbf{f} \, \mathbf{1} \, .$$

$$(4.38)$$

Due to the normalization (2.5),

$$\langle \, \omega | \, \left(e_a(x) \otimes \pi_1 \right) \tag{4.39}$$

does not vanish only when the input is the element of \mathcal{H}_2 . Then, we obtain

$$\langle \phi^{a_1}(x_1) \dots \phi^{a_n}(x_n) \partial_\mu j^\mu(x) \rangle$$

$$= -i\hbar \sum_a \left(\langle \omega_n^{a_1,\dots,a_n}(x_1,\dots,x_n) | \pi_n \otimes \langle \omega | (\pi_1 \mathbf{A} \otimes \tilde{e}^a(x)) \right) \mathbf{e}_a(x) \mathbf{f} \mathbf{1}$$

$$= -i\hbar \sum_a (-1)^{e_a} \left(\langle \omega_n^{a_1,\dots,a_n}(x_1,\dots,x_n) | \pi_n \mathbf{e}_a(x) \otimes \langle \omega | (\pi_1 \mathbf{A} \otimes \tilde{e}^a(x)) \right) \mathbf{f} \mathbf{1}$$

$$- i\hbar \sum_a \left(\langle \omega_n^{a_1,\dots,a_n}(x_1,\dots,x_n) | \pi_n \otimes \langle \omega | (\pi_1 \mathbf{A} \mathbf{e}_a(x) \otimes \tilde{e}^a(x)) \right) \mathbf{f} \mathbf{1} .$$

$$(4.40)$$

The second term on the right-hand side corresponds to the anomaly. Let us rewrite the first term to make its meaning clear. According to the definition of $\langle \omega_n^{a_1,\ldots,a_n}(x_1,\ldots,x_n)|$, this term can be expressed as the sum of cases where $e_a(x)$ enters one of the *n* slots,

$$-i\hbar\sum_{a}\sum_{i=1}^{n}(-1)^{n+e_{a}+e_{a}(e_{a_{i+1}}+\dots+e_{a_{n}})}\Big(\langle\omega|\left(\widetilde{e}^{a_{1}}(x_{1})\otimes\pi_{1}\right)\otimes\dots\otimes\langle\omega|\left(\widetilde{e}^{a_{i}}(x_{i})\otimes e_{a}(x)\right)\otimes\dots\otimes\langle\omega|\left(\widetilde{e}^{a_{n}}\otimes\pi_{1}\right)\otimes\langle\omega|\left(\pi_{1}\mathbf{A}\otimes\widetilde{e}^{a}(x)\right)\Big)\mathbf{f}\mathbf{1}.$$

$$(4.41)$$

 $^{^{12}}$ In addition to the contribution of anomalies, these divergences are related to the renormalization of composite operators. We will discuss this point in §6.

With the careful treatment of sign factors, we obtain

$$-i\hbar \sum_{i=1}^{n} (-1)^{n} \,\delta^{d}(x-x_{i}) \Big(\left\langle \omega \right| \left(\widetilde{e}^{a_{1}}(x_{1}) \otimes \pi_{1} \right) \otimes \cdots \otimes \left\langle \omega \right| \left(\widetilde{e}^{a_{i}}(x_{i}) \otimes \pi_{1} \mathbf{A} \right) \otimes \cdots \\ \otimes \left\langle \omega \right| \left(\widetilde{e}^{a_{n}}(x_{n}) \otimes \pi_{1} \right) \Big) \,\mathbf{f} \,\mathbf{1} \,, \tag{4.42}$$

where we used the normalization (2.5). Again, from the definition of $\langle \omega_n^{a_1,\ldots,a_n}(x_1,\ldots,x_n)|$, it can be expressed as

$$-i\hbar \sum_{i=1}^{n} \delta^{d}(x-x_{i}) \left\langle \omega_{n}^{a_{1},\dots,a_{n}}(x_{1},\dots,x_{n}) \right| \left(\pi_{i-1} \otimes \pi_{1} \mathbf{A} \otimes \pi_{n-i} \right) \mathbf{f} \mathbf{1}.$$

$$(4.43)$$

Since the operator **A** generates the infinitesimal transformation (4.5) and the replacement of $\frac{1}{1-\Phi}$ by **f 1** corresponds to taking the expectation value, we find that this term is a contact term,

$$-i\hbar \sum_{a} (-1)^{e_a} \left(\left\langle \omega_n^{a_1,\dots,a_n}(x_1,\dots,x_n) \right| \pi_n \mathbf{e}_a(x) \otimes \left\langle \omega \right| (\pi_1 \mathbf{A} \otimes \widetilde{e}^{a}(x)) \right) \mathbf{f} \mathbf{1}$$

= $-i\hbar \sum_{i} \int \mathrm{d}^d x \, \epsilon(x) \, \delta^d(x-x_i) \left\langle \phi^{a_1}(x_1) \dots \delta \phi^{a_i}(x_i) \dots \phi^{a_n}(x_n) \right\rangle \,.$ (4.44)

In summary, the Ward-Takahashi identity can be viewed in terms of homotopy algebras as

$$-\sum_{a} \left(\left\langle \omega_{n}^{a_{1},\dots,a_{n}}(x_{1},\dots,x_{n}) \right| \pi_{n} \otimes \left\langle \omega \right| \left(e_{a}(x) \otimes \pi_{1} \mathbf{M} \right) \otimes \left\langle \omega \right| \left(\widetilde{e}^{a}(x) \otimes \pi_{1} \mathbf{A} \right) \right) \mathbf{f} \mathbf{1}$$

$$= -i\hbar \sum_{i=1}^{n} \delta^{d}(x-x_{i}) \left\langle \omega_{n}^{a_{1},\dots,a_{n}}(x_{1},\dots,x_{n}) \right| \left(\pi_{i-1} \otimes \pi_{1} \mathbf{A} \otimes \pi_{n-i} \right) \mathbf{f} \mathbf{1}$$

$$-i\hbar \sum_{a} \left(\left\langle \omega_{n}^{a_{1},\dots,a_{n}}(x_{1},\dots,x_{n}) \right| \pi_{n} \otimes \left\langle \omega \right| \left(\pi_{1} \mathbf{A} \mathbf{e}_{a}(x) \otimes \widetilde{e}^{a}(x) \right) \right) \mathbf{f} \mathbf{1}.$$

$$(4.45)$$

The left-hand side represents the expectation value of fields and the derivative of the current $\partial_{\mu} j^{\mu}(x)$, while the first term on the right-hand side corresponds to the contact term, and the second term represents the anomaly.

4.4 Examples

In this subsection, we give several examples of the main formula (4.45).

4.4.1 The U(1) symmetry in complex scalar field theories

Let us consider the U(1) symmetry in complex scalar field theory. We consider the action given by

$$S = -\int d^d x \left[\varphi^{\dagger}(x) (-\partial^2 + m^2) \varphi(x) + \sum_{n=1}^{\infty} \frac{1}{2n} g_n (\varphi^{\dagger}(x) \varphi(x))^n \right].$$
(4.46)

The action is invariant under the U(1) transformation given by

$$\begin{aligned} \varphi(x) &\to \varphi'(x) = e^{-i\alpha} \varphi(x), \\ \varphi^{\dagger}(x) &\to \varphi'^{\dagger}(x) = e^{i\alpha} \varphi^{\dagger}(x). \end{aligned} \tag{4.47}$$

The infinitesimal transformations $\varphi(x) \to \varphi(x) + \delta \varphi(x)$ and $\varphi^{\dagger}(x) \to \varphi^{\dagger}(x) + \delta \varphi^{\dagger}(x)$ are given by

$$\begin{aligned} \delta\varphi(x) &\to -i\,\alpha\,\varphi(x)\,,\\ \delta\varphi^{\dagger}(x) &\to i\,\alpha\,\varphi^{\dagger}(x)\,, \end{aligned} \tag{4.48}$$

In terms of quantum A_{∞} algebras, we consider an action of the form

$$S = -\frac{1}{2}\omega(\Phi, Q\Phi) - \sum_{n=1}^{\infty} \frac{1}{2n}\omega(\Phi, m_{2n-1}(\Phi, \dots, \Phi)), \qquad (4.49)$$

where

$$\Phi = \int d^d x \left[\varphi(x) c^{\dagger}(x) + \varphi^{\dagger}(x) c(x) \right], \qquad (4.50)$$

is the degree-even element of \mathcal{H}_1 and c(x) is the basis vector of \mathcal{H}_1 and $c^{\dagger}(x)$ is its hermitian conjugate. Associated with the basis vector c(x), we introduce a degree-odd basis vector d(x)of \mathcal{H}_2 and its hermitian conjugate $d^{\dagger}(x)$. Then, the non-vanishing symplectic form ω is defined by

$$\omega (c(x_1), d^{\dagger}(x_2)) = \delta^d (x_1 - x_2),
\omega (c^{\dagger}(x_1), d(x_2)) = \delta^d (x_1 - x_2),
\omega (d^{\dagger}(x_2), c(x_1)) = -\delta^d (x_1 - x_2),
\omega (d(x_2), c^{\dagger}(x_1)) = -\delta^d (x_1 - x_2).$$
(4.51)

We define the operator Q by

$$Q c(x) = (-\partial^2 + m^2) d(x), \quad Q d(x) = 0,$$

$$Q c^{\dagger}(x) = (-\partial^2 + m^2) d^{\dagger}(x), \quad Q d^{\dagger}(x) = 0,$$
(4.52)

to obtain

$$-\frac{1}{2}\omega\left(\Phi,Q\Phi\right) = -\int d^{d}x\,\varphi^{\dagger}(x)\left(-\partial^{2}+m^{2}\right)\varphi(x)\,.$$

$$(4.53)$$

To define the operators m_{2n-1} , which describe interaction terms, let $e_1(x_1), \ldots, e_{2n-1}(x_{2n-1})$ be the 2n-1 inputs of m_{2n-1} . If the operator m_{2n-1} contains n pieces of c and n-1 pieces of c^{\dagger} , we define

$$m_{2n-1}(e_1(x_1),\ldots,e_{2n-1}(x_{2n-1})) = \frac{n!(n-1)!}{2(2n-1)!} g_n \,\delta^d(x_1-x_2)\cdots\delta^d(x_1-x_{2n-1}) \,d(x_1)\,, \quad (4.54)$$

and if the operator m_{2n-1} contains n-1 pieces of c and n pieces of c^{\dagger} , we define

$$m_{2n-1}(e_1(x_1),\ldots,e_{2n-1}(x_{2n-1})) = \frac{n!(n-1)!}{2(2n-1)!} g_n \,\delta^d(x_1-x_2)\cdots\delta^d(x_1-x_{2n-1}) \,d^{\dagger}(x_1) \,. \tag{4.55}$$

Otherwise, we define

$$m_{2n-1}(e_1(x_1),\ldots,e_{2n-1}(x_{2n-1})) = 0.$$
 (4.56)

Associated with the operator Q, we introduce the contracting homotopy h by

$$h c(x) = 0, \qquad h d(x) = \int d^{d}y \,\Delta(x - y) \,c(y),$$

$$h c^{\dagger}(x) = 0, \qquad h d^{\dagger}(x) = \int d^{d}y \,\Delta(x - y) \,c^{\dagger}(y),$$
(4.57)

where $\Delta(x-y)$ is the scalar propagator given by

$$\Delta(x-y) = \int \frac{d^d k}{(2\pi)^d} \frac{e^{ik(x-y)}}{k^2 + m^2 - i\epsilon} \,. \tag{4.58}$$

The correlation functions of this theory in terms of homotopy algebras are given by the formula (3.42) with the operators defined above.

Next, let us consider the symmetry transformations in terms of homotopy algebras. We introduce the degree-even operator A by

$$A c(x) = i \alpha c(x), \qquad A d(x) = i \alpha d(x),$$

$$A c^{\dagger}(x) = -i \alpha c^{\dagger}(x), \qquad A d^{\dagger}(x) = -i \alpha d^{\dagger}(x).$$
(4.59)

This operator correctly reproduces the symmetry transformation (4.48).

$$A\Phi = \int d^{d}x \left[\varphi(x) A c^{\dagger}(x) + \varphi^{\dagger}(x) A c(x)\right]$$

=
$$\int d^{d}x \left\{\varphi(x) \left[-i\alpha c^{\dagger}(x)\right] + \varphi^{\dagger}(x) \left[i\alpha c(x)\right]\right\}$$

=
$$\int d^{d}x \left\{\left[-i\alpha \varphi(x)\right] c^{\dagger}(x) + \left[i\alpha \varphi^{\dagger}(x)\right] c(x)\right\}$$

=
$$\int d^{d}x \left[\delta\varphi(x) c^{\dagger}(x) + \delta\varphi^{\dagger}(x) c(x)\right]$$

=
$$\delta\Phi.$$

(4.60)

It can be easily verified that $[\mathbf{M}, \mathbf{A}] = 0$ holds.

The Ward-Takahashi identity in terms of homotopy algebras (4.45) are described by

$$-\sum_{a\in\{c,c^{\dagger}\}} \left(\left\langle \omega_{n}^{a_{1},\dots,a_{n}}(x_{1},\dots,x_{n})\right| \pi_{n} \otimes \left\langle \omega\right| \left(e_{a}(x) \otimes \pi_{1}\mathbf{M} \right) \otimes \left\langle \omega\right| \left(\tilde{e}^{a}(x) \otimes \pi_{1}\mathbf{A} \right) \right) \mathbf{f} \mathbf{1} \\ = -i\hbar \sum_{i=1}^{n} \delta^{d}(x-x_{i}) \left\langle \omega_{n}^{a_{1},\dots,a_{n}}(x_{1},\dots,x_{n})\right| \left(\pi_{i-1} \otimes \pi_{1}\mathbf{A} \otimes \pi_{n-i} \right) \mathbf{f} \mathbf{1} \\ -i\hbar \sum_{a\in\{c,c^{\dagger}\}} \left(\left\langle \omega_{n}^{a_{1},\dots,a_{n}}(x_{1},\dots,x_{n})\right| \pi_{n} \otimes \left\langle \omega\right| \left(\pi_{1}\mathbf{A}\mathbf{e}_{a}(x) \otimes \tilde{e}^{a}(x) \right) \right) \mathbf{f} \mathbf{1} ,$$

$$(4.61)$$

where

$$e_a(x) = \begin{cases} c(x) & (a=c) \\ c^{\dagger}(x) & (a=c^{\dagger}) \end{cases}$$

$$(4.62)$$

and

$$\widetilde{e}^{a}(x) = \begin{cases} d^{\dagger}(x) & (a=c) \\ d(x) & (a=c^{\dagger}) \end{cases}$$

$$(4.63)$$

Let us first consider the case for free theory. In this case, the identities (4.61) can be written as

$$-\sum_{a\in\{c,c^{\dagger}\}} \left(\left\langle \omega_{n}^{a_{1},\dots,a_{n}}(x_{1},\dots,x_{n})\right| \pi_{n} \otimes \left\langle \omega\right| \left(e_{a}(x)\otimes\pi_{1}\mathbf{Q}\right)\otimes \left\langle \omega\right| \left(\widetilde{e}^{a}(x)\otimes\pi_{1}\mathbf{A}\right) \right) \mathbf{f} \mathbf{1} \\ = -i\hbar\sum_{i=1}^{n} \delta^{d}(x-x_{i}) \left\langle \omega_{n}^{a_{1},\dots,a_{n}}(x_{1},\dots,x_{n})\right| \left(\pi_{i-1}\otimes\pi_{1}\mathbf{A}\otimes\pi_{n-i}\right) \mathbf{f} \mathbf{1} \\ -i\hbar\sum_{a\in\{c,c^{\dagger}\}} \left(\left\langle \omega_{n}^{a_{1},\dots,a_{n}}(x_{1},\dots,x_{n})\right| \pi_{n}\otimes\left\langle \omega\right| \left(\pi_{1}\mathbf{A}\mathbf{e}_{a}(x)\otimes\widetilde{e}^{a}(x)\right) \right) \mathbf{f} \mathbf{1}.$$

$$(4.64)$$

For simplicity, we take n = 2k, $a_1 = a_2 = \ldots = a_k = c^{\dagger}$ and $a_{k+1} = a_{k+2} = \ldots = a_{2k} = c$. Then, the operator $\langle \omega_n^{a_1,\ldots,a_n}(x_1,\ldots,x_n) |$ is

$$\langle \omega_{2k}^{a_1,\dots,a_{2k}}(x_1,\dots,x_{2k})| = \langle \omega | (d(x_1)\otimes\pi_1)\otimes\dots\otimes\langle \omega | (d(x_k)\otimes\pi_1) \\ \otimes \langle \omega | (d^{\dagger}(x_{k+1})\otimes\pi_1)\otimes\dots\otimes\langle \omega | (d^{\dagger}(x_{2k})\otimes\pi_1).$$

$$(4.65)$$

Since

$$\pi_n \mathbf{f} \mathbf{1} = \langle \Phi^{\otimes n} \rangle = \left\langle \left[\int d^d x \left(\varphi(x) c^{\dagger}(x) + \varphi^{\dagger}(x) c(x) \right) \right]^{\otimes n} \right\rangle, \qquad (4.66)$$

the left-hand side of (4.64) is given by

$$-\int d^{d}y \, d^{d}z \, \langle \varphi(x_{1}) \dots \varphi(x_{k}) \varphi^{\dagger}(x_{k+1}) \dots \varphi^{\dagger}(x_{2k}) \\ \left[\varphi(y) \varphi^{\dagger}(z) \, \langle \omega | \, (c(x) \otimes Q \, c^{\dagger}(y) \,) \otimes \langle \omega | \, (d^{\dagger}(x) \otimes A \, c(z)) \\ + \varphi^{\dagger}(y) \, \varphi(z) \, \langle \omega | \, (c^{\dagger}(x) \otimes Q \, c(y) \,) \otimes \langle \omega | \, (d(x) \otimes A \, c^{\dagger}(z) \,) \, \right] \right\rangle.$$

$$(4.67)$$

The quantity inside the square brackets can be calculated as follows. From the definition (4.52) and (4.59), we obtain

$$-\int d^{d}y \, d^{d}z \, \varphi(y) \, \varphi^{\dagger}(z) \, \langle \omega | \, (c(x) \otimes Q \, c^{\dagger}(y) \,) \otimes \langle \omega | \, (d^{\dagger}(x) \otimes A \, c(z)) \\ = -\int d^{d}y \, d^{d}z \, \varphi(y) \, \varphi^{\dagger}(z) \, \langle \omega | \, (c(x) \otimes (-\partial_{y}^{2} + m^{2}) \, d^{\dagger}(y) \,) \otimes \langle \omega | \, (d^{\dagger}(x) \otimes i\alpha \, c(z)) \\ = -i\alpha \, \int d^{d}y \, d^{d}z \, (-\partial_{y}^{2} + m^{2}) \, \varphi(y) \, \varphi^{\dagger}(z) \, \langle \omega | \, (c(x) \otimes d^{\dagger}(y) \,) \otimes \langle \omega | \, (d^{\dagger}(x) \otimes c(z) \,) \,,$$

$$(4.68)$$

and

$$-\int d^{d}y \, d^{d}z \, \varphi^{\dagger}(y) \, \varphi(z) \, \langle \omega | \, (c^{\dagger}(x) \otimes Q \, c(y) \,) \otimes \langle \omega | \, (d(x) \otimes A \, c^{\dagger}(z) \,)$$

$$= -\int d^{d}y \, d^{d}z \, \varphi^{\dagger}(y) \, \varphi(z) \, \langle \omega | \, (c^{\dagger}(x) \otimes (-\partial_{y}^{2} + m^{2}) \, d(y)) \otimes \langle \omega | \, (d(x) \otimes (-i\alpha) \, c^{\dagger}(z) \,) \quad (4.69)$$

$$= i\alpha \, \int d^{d}y \, d^{d}z \, (-\partial_{y}^{2} + m^{2}) \, \varphi^{\dagger}(y) \, \varphi(z) \, \langle \omega | \, (c^{\dagger}(x) \otimes d(y) \,) \otimes \langle \omega | \, (d(x) \otimes c^{\dagger}(z) \,) \,.$$

Then, the quantity inside the square brackets in (4.67) can be expressed as

$$i\alpha \left(\varphi(x)\,\partial^2 \varphi^{\dagger}(x) - \varphi^{\dagger}(x)\,\partial^2 \varphi(x)\,\right). \tag{4.70}$$

We find that this equals to $\partial_{\mu} j^{\mu}(x)$, where $j^{\mu}(x)$ denotes the Noether current associated with the symmetry (4.48). Hence, the left-hand side of (4.64) is equal to

$$\langle \varphi(x_1) \varphi(x_2) \dots \varphi(x_k) \varphi^{\dagger}(x_{k+1}) \varphi^{\dagger}(x_{k+2}) \dots \varphi^{\dagger}(x_{2k}) \partial_{\mu} j^{\mu}(x) \rangle.$$
 (4.71)

Next, let us consider the second term of the left-hand side of (4.64). The relation (4.60) yields

$$(\pi_{i-1} \otimes \pi_1 \mathbf{A} \otimes \pi_{2k-i}) \mathbf{f} \mathbf{1} = \langle \Phi^{\otimes i-1} \otimes \delta \Phi \otimes \Phi^{\otimes (2k-i)} \rangle, \qquad (4.72)$$

and we find that the first term of the right-hand side of (4.64) is given by

$$-i\hbar \sum_{i=1}^{2k} \delta^{d}(x-x_{i}) \left\langle \omega_{2k}^{a_{1},\dots,a_{2k}}(x_{1},\dots,x_{2k}) \right| (\pi_{i-1}\otimes\pi_{1}\mathbf{A}\otimes\pi_{2k-i}) \mathbf{f} \mathbf{1}$$

$$= -i\hbar \sum_{i=1}^{k} \delta^{d}(x-x_{i}) \left\langle \varphi(x_{1})\dots\varphi(x_{i-1})\delta\varphi(x_{i})\varphi(x_{i+1})\dots\varphi(x_{k})\varphi^{\dagger}(x_{k+1})\dots\varphi^{\dagger}(x_{2k})\right\rangle$$

$$-i\hbar \sum_{i=1}^{k} \delta^{d}(x-x_{i}) \left\langle \varphi(x_{1})\dots\varphi(x_{k})\varphi^{\dagger}(x_{k+1})\dots\varphi^{\dagger}(x_{k+i-1})\delta\varphi^{\dagger}(x_{k+i})\varphi^{\dagger}(x_{k+i+1})\dots\varphi^{\dagger}(x_{2k})\right\rangle.$$

$$(4.73)$$

The second term of the right-hand side of (4.64) is given by

$$-i\hbar \sum_{a\in\{c,c^{\dagger}\}} \left(\left\langle \omega_{2k}^{a_{1},\dots,a_{n}}(x_{1},\dots,x_{2k}) \right| \pi_{2k} \otimes \left\langle \omega \right| \left(\pi_{1} \operatorname{\mathbf{Ae}}_{a}(x) \otimes \widetilde{e}^{a}(x) \right) \right) \operatorname{\mathbf{f}} \operatorname{\mathbf{1}}$$

$$= -i\hbar \left\langle \varphi(x_{1})\dots\varphi(x_{k}) \varphi^{\dagger}(x_{k+1})\dots\varphi^{\dagger}(x_{2k}) \right[\left\langle \omega \right| \left(Ac(x) \otimes d^{\dagger}(x) \right) + \left\langle \omega \right| \left(Ac^{\dagger}(x) \otimes d(x) \right) \right] \right\rangle.$$

(4.74)

We obtain

$$\langle \omega | (Ac(x) \otimes d^{\dagger}(x)) + \langle \omega | (Ac^{\dagger}(x) \otimes d(x)) = i\alpha \, \delta^{d}(0) - i\alpha \, \delta^{d}(0) \,. \tag{4.75}$$

This expression is ill-defined and requires regularization, however, for the convenience, we ignore the divergence and then, this term vanishes. Details regarding the regularization are discussed in §5. Then, the identities (4.64) can be expressed as

$$\langle \varphi(x_1) \varphi(x_2) \dots \varphi(x_k) \varphi^{\dagger}(x_{k+1}) \varphi^{\dagger}(x_{k+2}) \dots \varphi^{\dagger}(x_{2k}) \partial_{\mu} j^{\mu}(x) \rangle$$

$$= -i\hbar \sum_{i=1}^{k} \delta^{d}(x-x_i) \langle \varphi(x_1) \dots \varphi(x_{i-1}) \delta\varphi(x_i) \varphi(x_{i+1}) \dots \varphi(x_k) \varphi^{\dagger}(x_{k+1}) \dots \varphi^{\dagger}(x_{2k}) \rangle$$

$$- i\hbar \sum_{i=1}^{k} \delta^{d}(x-x_i) \langle \varphi(x_1) \dots \varphi(x_k) \varphi^{\dagger}(x_{k+1}) \dots \varphi^{\dagger}(x_{k+i-1}) \delta\varphi^{\dagger}(x_{k+i}) \varphi^{\dagger}(x_{k+i+1}) \dots \varphi^{\dagger}(x_{2k}) \rangle .$$

$$(4.76)$$

This is exactly the Ward-Takahashi identity for the free complex scalar theory. Actually, for the action (4.46), the interaction term does not affect the identity (4.61). In the interaction theory, the term that contains

$$\sum_{n=1}^{\infty} \sum_{a \in \{c,c^{\dagger}\}} \langle \omega | (e_a(x) \otimes \pi_1 \mathbf{m}_{2n-1}) \otimes \langle \omega | (\tilde{e}^a(x) \otimes \pi_1 \mathbf{A})$$
(4.77)

is added to the identities for the free theory. We can verify that this term vanishes. It is sufficient to evaluate its action on

$$\Phi^{\otimes 2n} = \left[\int d^d x \left(\varphi(x) c^{\dagger}(x) + \varphi^{\dagger}(x) c(x) \right) \right]^{\otimes 2n}.$$
(4.78)

Terms in which the number of c's and c^{\dagger} 's is not equal vanish trivially due to the definition of m_{2n-1} . There are two types of non-trivial terms. One type is the term where \mathbf{m}_{2n-1} takes n pieces of c and n-1 pieces of c^{\dagger} as inputs, and \mathbf{A} takes a c as a input. There are $\binom{2n}{n}$ such terms. These terms are evaluated as

$$-i\alpha \frac{n! (n-1)!}{2(2n-1)!} g_n \left(\varphi(x)\varphi^{\dagger}(x)\right)^n \times \binom{2n}{n} = -\frac{i\alpha g_n}{2} (\varphi(x)\varphi^{\dagger}(x))^n \,. \tag{4.79}$$

Another type is the term where \mathbf{m}_{2n-1} takes n-1 pieces of c and n pieces of c^{\dagger} as input, and **A** takes a c^{\dagger} as input. There are also $\binom{2n}{n}$ such terms. These terms are evaluated as

$$i\alpha \frac{n! (n-1)!}{2(2n-1)!} g_n \left(\varphi(x)\varphi^{\dagger}(x)\right)^n \times \binom{2n}{n} = \frac{i\alpha g_n}{2} (\varphi(x)\varphi^{\dagger}(x))^n , \qquad (4.80)$$

and cancel out with the former. Hence, the interaction term does not affect the current and obtain the same form of the Ward-Takahashi identities as (4.76).

4.4.2 The energy-momentum tensor in scalar field theories

In this subsection, we consider the energy-momentum tensor in scalar field theories. We use the setup described in §3.1. We consider a scalar field theory whose action is given by

$$S = \int \mathrm{d}^d x \left(-\frac{1}{2} \partial^\mu \varphi(x) \,\partial_\mu \varphi(x) - \sum_{n=0}^\infty \frac{1}{n+1} g_n \,\varphi(x)^{n+1} \right),\tag{4.81}$$

This theory has the spacetime transformation symmetry:

$$\varphi(x) \to \varphi(x-a),$$
 (4.82)

where a is the constant spacetime transformation. The infinitesimal transformation $\varphi(x) \rightarrow \varphi(x) + \delta \phi(x)$ of (4.82) is given by

$$\delta\varphi(x) = -a^{\nu} \,\partial_{\nu} \,\varphi(x) \,. \tag{4.83}$$

From Noether's theorem, the conserved current associated with the transformation (4.82) is given by

$$j^{\mu}(x) = a_{\nu} T^{\mu\nu} , \qquad (4.84)$$

where $T^{\mu\nu}$ is the energy-momentum tensor. The energy-momentum tensor is given by

$$T^{\mu\nu} = -\frac{\partial \mathcal{L}(x)}{\partial (\partial_{\mu} \varphi_a(x))} + \eta^{\mu\nu} \mathcal{L}(x) , \qquad (4.85)$$

where $\mathcal{L}(x)$ is the lagrangian density and $\eta^{\mu\nu}$ is a metric defined by

$$\eta^{\mu\nu} = \text{diag}(-1, 1, 1, 1).$$
(4.86)

In terms of homotopy algebras, we can rewrite the above action to

$$S = -\frac{1}{2}\omega\left(\Phi, Q\Phi\right) - \sum_{n=0}^{\infty} \frac{1}{n+1}\omega\left(\Phi, m_n\left(\Phi, \dots, \Phi\right)\right), \qquad (4.87)$$

where

$$Q c(x) = (-\partial^2 + m^2) d(x), \qquad Q d(x) = 0,$$
 (4.88)

and

$$m_n(c(x_1),\ldots,c(x_n)) = g_{n+1}\,\delta^d(x_1-x_2)\ldots\delta^d(x_1-x_n)\,d(x_1)\,,\tag{4.89}$$

for n > 0 and

$$m_0 \mathbf{1} = \int \mathrm{d}^d x \ g_1 \, d(x) \,. \tag{4.90}$$

To consider the symmetry tansformation (4.83), we introduce degree-even operators

$$A c(x) = a_{\nu} \partial^{\nu} c(x) ,$$

$$A d(x) = a_{\nu} \partial^{\nu} d(x) .$$
(4.91)

Since

$$A\Phi = \int d^{d}x \left[\varphi(x) A c(x)\right]$$

=
$$\int d^{d}x a_{\nu}\varphi(x) \partial^{\nu}c(x)$$

=
$$\int d^{d}x \left[-a_{\nu} \partial^{\nu} \varphi(x)\right] c(x)$$

=
$$\int d^{d}x \,\delta\varphi(x) c(x)$$

=
$$\delta\Phi,$$

(4.92)

this correctly reproduces the symmetry transformation (4.83). It can be easily verified that $[\mathbf{M}, \mathbf{A}] = 0$ holds.

The Ward-Takahashi identities in terms of homotopy algebras (4.45) are described by

$$-\left(\left\langle \omega_{n}(x_{1},\ldots x_{n})| \pi_{n} \otimes \left\langle \omega\right| (c(x) \otimes \pi_{1}\mathbf{M}) \otimes \left\langle \omega\right| (d(x) \otimes \pi_{1}\mathbf{A})\right) \mathbf{f} \mathbf{1} \\ = -i\hbar \sum_{i=1}^{n} \delta^{d}(x-x_{i}) \left\langle \omega_{n}(x_{1},\ldots x_{n})\right| (\pi_{i-1} \otimes \pi_{1}\mathbf{A} \otimes \pi_{n-i}) \mathbf{f} \mathbf{1} \\ -i\hbar \left(\left\langle \omega_{n}(x_{1},\ldots x_{n})\right| \pi_{n} \otimes \left\langle \omega\right| (\pi_{1}\mathbf{A}\mathbf{c}(x) \otimes d(x))\right) \mathbf{f} \mathbf{1}.$$

$$(4.93)$$

Since

$$\pi_n \mathbf{f} \mathbf{1} = \langle \Phi^{\otimes n} \rangle = \left\langle \left[\int \mathrm{d}^d x \, \varphi(x) c(x) \right]^{\otimes n} \right\rangle \,, \tag{4.94}$$

the left-hand side of (4.93) is given by

$$-\left(\left\langle\omega_{n}(x_{1},\ldots x_{n})|\,\pi_{n}\,\otimes\left\langle\,\omega\right|\,\left(\,c(x)\otimes\pi_{1}\mathbf{M}\,\right)\otimes\left\langle\,\omega\right|\,\left(\,d(x)\otimes\pi_{1}\mathbf{A}\,\right)\,\right)\,\mathbf{f}\,\mathbf{1}\right.\\ =-\left\langle\left.\varphi(x_{1})\ldots\varphi(x_{n})\,\left\langle\,\omega\right|\,\left(\,c(x)\otimes\pi_{1}\,\mathbf{M}\,\frac{1}{1-\Phi}\,\right)\otimes\left\langle\,\omega\right|\,\left(\,d(x)\otimes A\,\Phi\,\right)\,\right\rangle\,.$$

$$(4.95)$$

Let us calculate

$$- \langle \omega | \left(c(x) \otimes \pi_{1} \mathbf{M} \frac{1}{1 - \Phi} \right) \otimes \langle \omega | (d(x) \otimes A \Phi)$$

=
$$- \langle \omega | (c(x) \otimes Q \Phi) \otimes \langle \omega | (d(x) \otimes A \Phi)$$

$$- \sum_{n=0}^{\infty} \langle \omega | (c(x) \otimes m_{n}(\Phi, \dots, \Phi)) \otimes \langle \omega | (d(x) \otimes A \Phi) .$$
(4.96)

Let us first consider the free part. For simplicity, we split Q into two parts:

$$Q = Q^{\rm kin} + Q^{\rm mass} \,, \tag{4.97}$$

$$Q^{\text{kin}} c(x) = -\partial^2 d(x), \quad Q^{\text{mass}} c(x) = m^2 d(x).$$
 (4.98)

The kinetic part can be calculated as follows,

$$- \langle \omega | (c(x) \otimes Q^{\operatorname{kin}} \Phi) \otimes \langle \omega | (d(x) \otimes A \Phi) \rangle$$

$$= - \int d^{d}y \, d^{d}z \, \langle \omega | (c(x) \otimes \varphi(y) (-\partial_{y}^{2} d(y))) \otimes \langle \omega | (d(x) \otimes \varphi(z) a_{\nu} \partial_{z}^{\nu} c(z)) \rangle$$

$$= -a_{\nu} \int d^{d}y \, d^{d}z \, \partial_{y}^{2} \varphi(y) \, \partial_{z}^{\nu} \varphi(z) \, \langle \omega | (c(x) \otimes d(y)) \otimes \langle \omega | (d(x) \otimes c(z)) \rangle$$

$$= a_{\nu} \, \partial^{2} \varphi(x) \, \partial^{\nu} \varphi(x)$$

$$= a_{\nu} \, \partial_{\mu} \left(\partial^{\mu} \varphi(x) \, \partial^{\nu} \varphi(x) - \frac{1}{2} \, \eta^{\mu\nu} \, \partial^{\rho} \varphi(x) \, \partial_{\rho} \varphi(x) \right)$$

$$= a_{\nu} \, \partial_{\mu} T_{\operatorname{kin}}^{\mu\nu}(x) \,, \qquad (4.99)$$

where $T_{\rm kin}^{\mu\nu}$ is the energy-momentum tensor which is the contribution from the kinetic term. Since the contribution of the operator $Q^{\rm mass}$ can be calculated in the same way as that of the operator m_1 , we omit the calculation.

Let us next consider the interaction part:

$$-\sum_{n=0}^{\infty} \langle \omega | (c(x) \otimes m_n(\Phi, \dots, \Phi)) \otimes \langle \omega | (d(x) \otimes A \Phi).$$
(4.100)

This can be calculated as follows:

$$- \langle \omega | (c(x) \otimes m_n(\Phi, \dots, \Phi)) \otimes \langle \omega | (d(x) \otimes A \Phi) \rangle$$

$$= - \int d^d y d^d z g_{n+1} \varphi(y)^n \varphi(z) \langle \omega | (c(x) \otimes d(y)) \otimes \langle \omega | (d(x) \otimes a_\nu \partial^\nu c(z)) \rangle$$

$$= -a_\nu g_{n+1} \varphi(x)^n \partial^\nu \varphi(x) \qquad (4.101)$$

$$= -a_\nu \frac{g_{n+1}}{n+1} \partial_\mu (g^{\mu\nu} \varphi(x)^{n+1})$$

$$= a_\nu \partial_\mu T_{n+1}^{\mu\nu},$$

where $T_{n+1}^{\mu\nu}$ is the energy-momentum tensor which is the contribution from the (n + 1)-point interactions. Combining all the contributions, the left-hand side of (4.93) can be expressed as

$$-\left(\left\langle\omega_n(x_1,\ldots x_n)\right|\pi_n\otimes\left\langle\omega\right|(c(x)\otimes\pi_1\mathbf{M})\otimes\left\langle\omega\right|(d(x)\otimes\pi_1\mathbf{A})\right)\mathbf{f}\mathbf{1} \\ = a_\nu\left\langle\varphi(x_1)\ldots\varphi(x_n)\,\partial_\mu\,T^{\mu\nu}(x)\right\rangle.$$
(4.102)

The right-hand side of (4.93) can be evaluated in the same as the example of a complex scalar field and the first term can be written as

$$-i\hbar \sum_{i=1}^{n} \delta^{d}(x-x_{i}) \langle \omega_{n}(x_{1},\ldots,x_{n}) | (\pi_{i-1}\otimes\pi_{1}\mathbf{A}\otimes\pi_{n-i}) \mathbf{f} \mathbf{1}$$

$$= -i\hbar \sum_{i=1}^{n} \delta^{d}(x-x_{i}) \langle \varphi(x_{1})\ldots\varphi(x_{i-1}) \delta\varphi(x_{i}) \varphi(x_{i+1})\ldots\varphi(x_{n}) \rangle .$$

$$(4.103)$$

The second term includes an ill-defined factor

$$\langle \omega | (A c(x) \otimes d(x)) = \langle \omega | (a_{\nu} \partial^{\nu} c(x) \otimes d(x)) = \lim_{y \to x} a_{\nu} \partial_{x}^{\nu} \delta^{d}(x - y).$$
(4.104)

Thus, it requires regularization, which will be described in $\S5$. Here, we ignore this term. We finally find that the equation (4.93) is equivalent to

$$a_{\nu} \langle \varphi(x_{1}) \dots \varphi(x_{n}) \partial_{\mu} T^{\mu\nu}(x) \rangle$$

= $-i\hbar \sum_{i=1}^{n} \delta^{d}(x - x_{i}) \langle \varphi(x_{1}) \dots \varphi(x_{i-1}) \delta\varphi(x_{i}) \varphi(x_{i+1}) \dots \varphi(x_{n}) \rangle.$ (4.105)

This is exactly the Ward-Takahashi identity for the energy-momentum tensor of scalar field theory.

4.4.3 The chiral U(1) symmetry in vector-like gauge theories

Finally, let us consider the chiral U(1) symmetry in a vector-like gauge theory. Consider the action of a massless Dirac fermion given by

where s and t are spinor indices¹³ running from 1 to 4, $\not \!\!\!\!D$ is defined by

$$\mathcal{D}(x) = \gamma^{\mu} (\partial_{\mu} - iA_{\mu}(x)), \qquad (4.107)$$

and γ^{μ} is a γ matrix satisfying

$$\{\gamma^{\mu}, \gamma^{\nu}\} = -2\,\eta^{\mu\nu}\,. \tag{4.108}$$

 $^{^{13}}$ We take Einstein summation convention. We also omit the spinor indices if there is no confusion.

Furthermore, we treat $A_{\mu}(x)$ as the background field. This theory has a chiral U(1) symmetry:

$$\Psi(x) \to e^{-i\alpha\gamma_5} \Psi(x) ,
\overline{\Psi}(x) \to \overline{\Psi}(x) e^{-i\alpha\gamma_5} ,$$
(4.109)

where γ_5 is defined by

$$\gamma_5 = i\gamma^0 \gamma^1 \gamma^2 \gamma^3 \,. \tag{4.110}$$

The infinitesimal transformations are given by

$$\delta \Psi(x) = -i\alpha \gamma_5 \Psi(x), \delta \bar{\Psi}(x) = -i\alpha \overline{\Psi}(x) \gamma_5.$$
(4.111)

In terms of homotopy algebras, the above action can be rewritten as

$$S = -\frac{1}{2}\omega(\Phi, Q\Phi), \qquad (4.112)$$

where

$$\Phi = \int d^4x \, \left[\Psi_s(x) \,\overline{\theta}_s(x) + \overline{\Psi}_s(x) \,\theta_s(x) \right], \qquad (4.113)$$

is the degree-even element of \mathcal{H}_1 and $\{\theta_s(x), \overline{\theta}_s(x)\}$ are the degree-odd basis vectors of \mathcal{H}_1 .¹⁴ We introduce the degree-even basis vectors $\{\lambda_s(x), \overline{\lambda}_s(x)\}$ of \mathcal{H}_2 satisfying the normalization:

$$\omega(\theta_s(x), \overline{\lambda}_t(y)) = -\delta_{st} \,\delta^4(x-y),
\omega(\overline{\theta}_s(x), \lambda_t(y)) = -\delta_{st} \,\delta^4(x-y),
\omega(\overline{\lambda}_s(x), \theta_t(y)) = \delta_{st} \,\delta^4(x-y),
\omega(\lambda_s(x), \overline{\theta}_t(y)) = \delta_{st} \,\delta^4(x-y).$$
(4.114)

We do not need to introduce a basis vector for the background field A_{μ} . We define Q by

where \overleftarrow{D} is defined by

$$\overleftarrow{\mathcal{D}}(x) = \gamma^{\mu} (-\overleftarrow{\partial}_{\mu} - iA_{\mu}(x)), \qquad (4.116)$$

¹⁴In [35, 36], the basis vector $\overline{\theta}_s(x)$ is the Dirac adjoint of $\theta_s(x)$ to be consistent with the reality condition. Our notation is slightly different from the previous papers and the "adjoint" of the basis vector should be minus the Dirac adjoint.

and $\overleftarrow{\partial}_{\mu}$ denotes the differential acting from the right. Then, we reproduce the action in terms of homotopy algebras:

$$-\frac{1}{2}\omega(\Phi, Q\Phi) = \int d^4x \,\overline{\Psi}(x) \, i \, \mathcal{D}(x) \,\Psi(x) \,. \tag{4.117}$$

Associated with the operator Q, we introduce the contracting homotopy h by

$$h \theta_s(x) = 0, \quad h \lambda_s(x) = i \left[\mathcal{D}(x) \int_0^\infty d\tau \, e^{-\tau \mathcal{D}^2(x)} \right]_{st} \theta_t(x),$$

$$h \overline{\theta}_s(x) = 0, \quad h \overline{\lambda}_s(x) = -i \overline{\theta}_t(x) \left[\overleftarrow{\mathcal{D}}(x) \int_0^\infty d\tau \, e^{-\tau \overleftarrow{\mathcal{D}}^2(x)} \right]_{ts}.$$
(4.118)

The degree-even operator A corresponding to the symmetry transformation (4.111) is defined as

$$A \theta_s(x) = -i\alpha (\gamma_5)_{st} \theta_t(x), \quad A \lambda_s(x) = i\alpha (\gamma_5)_{st} \lambda_t(x), A \overline{\theta}_s(x) = -i\alpha \overline{\theta}_t(x) (\gamma_5)_{ts}, \quad A \overline{\lambda}_s(x) = i\alpha \overline{\lambda}_t(x) (\gamma_5)_{ts}.$$

$$(4.119)$$

It reproduces the the symmetry transformation as

$$A \Phi = \int d^4x \left[\Psi_s(x) A \overline{\theta}_s(x) + \overline{\Psi}_s(x) A \theta_s(x) \right]$$

= $-i\alpha \int d^4x \left[\Psi_s(x) (\gamma_5)_{ts} \overline{\theta}_t(x) + \overline{\Psi}_s(x) (\gamma_5)_{st} \theta_t(x) \right]$
= $\int d^4x \left[\delta \Psi_t(x) \overline{\theta}_t(x) + \delta \overline{\Psi}_t(x) \theta_t(x) \right]$
= $\delta \Phi$. (4.120)

Moreover, we can confirm that the operator A describes the symmetry since

$$[Q, A] \theta_{s}(x) = -i\alpha (\gamma_{5})_{st} Q \theta_{t}(x) + i \mathcal{D}_{st}(x) A \lambda_{t}(x)$$

$$= -\alpha (\gamma_{5})_{st} \mathcal{D}_{tu}(x) \lambda_{u}(x) - \alpha \mathcal{D}_{st}(x) (\gamma_{5})_{tu} \lambda_{u}(x)$$

$$= -\alpha (\gamma_{5})_{st} \mathcal{D}_{tu}(x) \lambda_{u}(x) + \alpha (\gamma_{5})_{st} \mathcal{D}_{tu}(x) \lambda_{u}(x)$$

$$= 0,$$
(4.121)

where we used $\{\gamma_5, \gamma^{\mu}\} = 0$. In the same way, we obtain

$$[Q, A]\overline{\theta}_s(x) = 0. \qquad (4.122)$$

The Ward-Takahashi identity in terms of homotopy algebras (4.45) is described by

$$-\sum_{a\in\{\theta_{s},\overline{\theta}_{s}\}} \left(\left\langle \omega_{n}^{a_{1},\dots,a_{n}}(x_{1},\dots,x_{n})\right| \pi_{n} \otimes \left\langle \omega\right| \left(e_{a}(x)\otimes\pi_{1}\mathbf{Q}\right)\otimes \left\langle \omega\right| \left(\widetilde{e}^{a}(x)\otimes\pi_{1}\mathbf{A}\right) \right) \mathbf{f} \mathbf{1} \\ = -i\hbar\sum_{i=1}^{n} \delta^{4}(x-x_{i}) \left\langle \omega_{n}^{a_{1},\dots,a_{n}}(x_{1},\dots,x_{n})\right| \left(\pi_{i-1}\otimes\pi_{1}\mathbf{A}\otimes\pi_{n-i}\right) \mathbf{f} \mathbf{1} \\ -i\hbar\sum_{a\in\{\theta_{s},\overline{\theta}_{s}\}} \left(\left\langle \omega_{n}^{a_{1},\dots,a_{n}}(x_{1},\dots,x_{n})\right| \pi_{n}\otimes\left\langle \omega\right| \left(\pi_{1}\mathbf{A}\mathbf{e}_{a}(x)\otimes\widetilde{e}^{a}(x)\right) \right) \mathbf{f} \mathbf{1} ,$$

$$(4.123)$$

where

$$e_a(x) = \begin{cases} \theta_s(x) & (a = \theta_s) \\ \overline{\theta}_s(x) & (a = \overline{\theta}_s) \end{cases}$$
(4.124)

and

$$\widetilde{e}^{a}(x) = \begin{cases} \overline{\lambda}_{s}(x) & (a = \theta_{s}) \\ \lambda^{s}(x) & (a = \overline{\theta}_{s}) \end{cases}$$
(4.125)

For simplicity, we take n = 2k, $a_i = \overline{\theta}_{s_i}$ for $i = 1, \ldots, k$ and $a_i = \theta_{s_i}$ for $i = k+1, \ldots, 2k$. Then, the operator $\langle \omega_n^{a_1,\ldots,a_n}(x_1,\ldots,x_n) |$ is

$$\langle \omega_{2k}^{a_1,\dots,a_{2k}}(x_1,\dots,x_{2k}) | = \langle \omega | (\lambda_{s_1}(x_1) \otimes \pi_1) \otimes \dots \otimes \langle \omega | (\lambda_{s_k}(x_k) \otimes \pi_1) \\ \otimes \langle \omega | (\overline{\lambda}_{s_{k+1}}(x_{k+1}) \otimes \pi_1) \otimes \dots \otimes \langle \omega | (\overline{\lambda}_{s_{2k}}(x_{2k}) \otimes \pi_1) .$$

$$(4.126)$$

Since

$$\pi_n \mathbf{f} \mathbf{1} = \left\langle \Phi^{\otimes n} \right\rangle = \left\langle \left[\int \mathrm{d}^4 x \, \left(\Psi_s(x) \,\overline{\theta}_s(x) + \overline{\Psi}_s(x) \,\theta_s(x) \right) \right]^{\otimes n} \right\rangle, \qquad (4.127)$$

the left-hand side of (4.123) is given by

$$-\int d^{4}y \, d^{4}z \, \langle \Psi_{s_{1}}(x_{1}) \dots \Psi_{s_{k}}(x_{k}) \, \overline{\Psi}_{s_{k+1}}(x_{k+1}) \dots \overline{\Psi}_{s_{2k}}(x_{2k}) \\ \left[\Psi_{t}(y) \, \overline{\Psi}_{u}(z) \, \langle \omega | \left(\theta_{s}(x) \otimes Q \, \overline{\theta}_{t}(y)\right) \otimes \langle \omega | \left(\overline{\lambda}_{s}(x) \otimes A \, \theta_{u}(z)\right) \right. \\ \left. + \overline{\Psi}_{t}(y) \, \Psi_{u}(z) \, \langle \omega | \left(\overline{\theta}_{s}(x) \otimes Q \, \theta_{t}(y)\right) \otimes \langle \omega | \left(\lambda_{s}(x) \otimes A \, \overline{\theta}_{u}(z)\right) \right] \rangle \,.$$

$$(4.128)$$

Since

and

$$-\int d^{4}y \, d^{4}z \, \overline{\Psi}_{t}(y) \, \Psi_{u}(z) \, \langle \omega | \left(\overline{\theta}_{s}(x) \otimes Q \, \theta_{t}(y) \right) \otimes \langle \omega | \left(\lambda_{s}(x) \otimes A \, \overline{\theta}_{u}(z) \right)$$

$$= -\alpha \, \overline{\Psi}(x) \, \overleftarrow{D}(x) \, \gamma_{5} \, \Psi(x) \,,$$

$$(4.130)$$

we obtain

$$-\int d^{4}y \, d^{4}z \left[\Psi_{t}(y) \,\overline{\Psi}_{u}(z) \, \langle \omega | \left(\theta_{s}(x) \otimes Q \,\overline{\theta}_{t}(y) \right) \otimes \langle \omega | \left(\overline{\lambda}_{s}(x) \otimes A \,\theta_{u}(z) \right) \right. \\ \left. + \overline{\Psi}_{t}(y) \,\Psi_{u}(z) \, \langle \omega | \left(\overline{\theta}_{s}(x) \otimes Q \,\theta_{t}(y) \right) \otimes \langle \omega | \left(\lambda_{s}(x) \otimes A \,\overline{\theta}_{u}(z) \right) \right]$$

$$= \partial_{\mu} \left[\alpha \,\overline{\Psi}_{s}(x) \,\gamma^{\mu} \,\gamma_{5} \,\Psi(x) \right]$$

$$= \partial_{\mu} \, j_{5}^{\mu}(x) \,, \qquad (4.131)$$

where $j_5^{\mu}(x)$ denotes the Noether current associated with the symmetry (4.111). Hence, the left-hand side of (4.123) is equal to

$$\left\langle \Psi_{s_1}(x_1)\dots\Psi_{s_k}(x_k)\,\overline{\Psi}_{s_{k+1}}(x_{k+1})\dots\overline{\Psi}_{s_{2k}}(x_{2k})\,\partial_\mu\,j_5^\mu\,\right\rangle\,.\tag{4.132}$$

The first term on the right-hand side of (4.123) can be expressed similarly to the previous examples as

$$-i\hbar \sum_{i=1}^{n} \delta^{4}(x-x_{i}) \left\langle \omega_{n}^{a_{1},...,a_{n}}(x_{1},...x_{n}) \right| \left(\pi_{i-1} \otimes \pi_{1}\mathbf{A} \otimes \pi_{n-i} \right) \mathbf{f} \mathbf{1}$$

$$= -i\hbar \sum_{i=1}^{k} \delta^{4}(x-x_{i}) \left\langle \Psi_{s_{1}}(x_{1}) \dots \Psi_{s_{i-1}}(x_{s-1}) \delta \Psi_{s_{i}}(x_{i}) \Psi_{s_{i+1}}(x_{i+1}) \dots \Psi_{s_{k}}(x_{k}) \right.$$

$$\overline{\Psi}_{s_{k+1}}(x_{k+1}) \dots \overline{\Psi}_{s_{2k}}(x_{2k}) \left\langle \Psi_{s_{1}}(x_{1}) \dots \Psi_{s_{k}}(x_{k}) \right\rangle$$

$$-i\hbar \sum_{i=k+1}^{2^{k}} \delta^{4}(x-x_{i}) \left\langle \Psi_{s_{1}}(x_{1}) \dots \Psi_{s_{k}}(x_{k}) \right\rangle$$

$$\overline{\Psi}_{s_{k+1}}(x_{k+1}) \dots \overline{\Psi}_{s_{i-1}}(x_{i-1}) \delta \overline{\Psi}_{s_{i}}(x_{i}) \overline{\Psi}_{s_{i+1}}(x_{i+1}) \dots \overline{\Psi}_{s_{2k}}(x_{2k}) \left\langle ...\right\rangle.$$

$$(4.133)$$

The second term still contains an ill-defined factor

$$\langle \omega | \left(A \,\theta_s(x) \otimes \overline{\lambda}_s(x) \right) + \langle \omega | \left(A \,\overline{\theta}_s(x) \otimes \lambda_s(x) \right)$$

= 2 *i* \alpha \lim_{y \rightarrow x} \text{tr} \gamma_5 \delta^4(x-y) .
 (4.134)

As is well known, the chiral symmetry is anomalous, and this factor does not vanish after the regularization. We will explore how to regularize this term from the homotopy algebraic point of view in the next section.

5 Regularization with stubs

As we mentioned in the previous section, we need to regularize the term

$$-i\hbar \sum_{a} \langle \omega | (\pi_1 \mathbf{A} \mathbf{e}_a(x) \otimes \widetilde{e}^a(x)) \mathbf{f} \mathbf{1}, \qquad (5.1)$$

to mediate the divergence. To preserve the homotopy algebraic structure, we introduce the regularization with *stubs*. The concept of stubs [7] is widely used in the context of string field theory, and they can be used for the regularization. Stubs in quantum field theory are discussed in [50]. In this regularization, the operators \mathbf{M} and \mathbf{U} are modified by the heat kernel in the way to preserve the homotopy algebraic structure. By repeating the previous arguments using these modified operators, we obtain the correct anomalous term. Let us comment on technical

facts about the heat kernel. In the Lorentzian signature, the heat kernel does not regularize path integrals. The solution is that we formally calculate and when we carry out the integral involving the heat kernel, we introduce the Wick rotation.

Let us explain the detailed treatment of stubs in homotopy algebras. In [46, 47, 48, 49, 50], adding stubs in terms of homotopy algebras is described by using the homological perturbation lemma (HPL) [53]. Before considering stubs, we explain the statement of the HPL. Consider two degree-even maps $\pi : \mathcal{H} \to \mathcal{H}'$ and $\iota : \mathcal{H}' \to \mathcal{H}$, where \mathcal{H}' denotes image of π , and degree-odd map $\eta : \mathcal{H} \to \mathcal{H}$ satisfying

$$\pi \iota = \mathbb{I}, \tag{5.2}$$

$$Q\eta + \eta Q = \mathbb{I} - \iota \pi , \qquad (5.3)$$

$$\eta^2 = \eta \iota = \pi \eta = 0. \tag{5.4}$$

From the condition (5.2), it follows that $\iota \pi$ becomes projection:

$$(\iota \pi)(\iota \pi) = \iota (\pi \iota) \pi = \iota \pi.$$
(5.5)

These maps can be promoted to operators on the tensor algebra as follows:

$$\boldsymbol{\pi} = \sum_{n=1}^{\infty} \pi^{\otimes n} ,$$

$$\boldsymbol{\iota} = \sum_{n=1}^{\infty} \iota^{\otimes n} ,$$
(5.6)

$$\boldsymbol{\eta} = \sum_{n=1}^{\infty} \sum_{k=0}^{n-1} \mathbb{I}^{\otimes k} \otimes \boldsymbol{\eta} \otimes (\iota \pi)^{\otimes n-k-1}$$

These operators also satisfy

$$\boldsymbol{\pi}\,\boldsymbol{\iota} = \mathbf{I}\,,\tag{5.7}$$

$$\mathbf{Q}\,\boldsymbol{\eta} + \boldsymbol{\eta}\,\mathbf{Q} = \mathbf{I} - \boldsymbol{\iota}\,\boldsymbol{\pi}\,,\tag{5.8}$$

$$\boldsymbol{\eta}^2 = \boldsymbol{\eta}\,\boldsymbol{\iota} = \boldsymbol{\pi}\,\boldsymbol{\eta} = 0\,. \tag{5.9}$$

According to the HPL, if the operator **M** satisfies the quantum A_{∞} relations

$$(\mathbf{M} + i\hbar \mathbf{U})^2 = 0,$$
 (5.10)

the operators

$$\pi' = \pi \frac{1}{\mathbf{I} + (\mathbf{m} + i\hbar \mathbf{U}) \boldsymbol{\eta}},$$

$$\iota' = \frac{1}{\mathbf{I} + \boldsymbol{\eta} (\mathbf{m} + i\hbar \mathbf{U})} \iota,$$

$$\eta' = \frac{1}{\mathbf{I} + \boldsymbol{\eta} (\mathbf{m} + i\hbar \mathbf{U})} \boldsymbol{\eta},$$
(5.11)

correspond to the perturbed operators of π, ι and η , respectively, and satisfy

$$\pi' \iota' = \mathbf{I} \,, \tag{5.12}$$

$$\left(\mathbf{M} + i\hbar \mathbf{U}\right)\boldsymbol{\eta}' + \boldsymbol{\eta}'\left(\mathbf{M} + i\hbar \mathbf{U}\right) = \mathbf{I} - \boldsymbol{\iota}' \boldsymbol{\pi}', \qquad (5.13)$$

$$(\eta')^2 = \eta' \iota' = \pi' \eta' = 0.$$
 (5.14)

When we can define a coderivation \mathbf{M}' and a second-order coderivation \mathbf{U}' as

$$\begin{split} \mathbf{M}' &= \boldsymbol{\pi} \, \mathbf{M} \, \boldsymbol{\iota}' \,, \\ \mathbf{U}' &= \boldsymbol{\pi} \, \mathbf{U} \, \boldsymbol{\iota} \,, \end{split}$$
 (5.15)

these operators satisfy

$$(\mathbf{M}' + i\hbar \mathbf{U}')^2 = 0,$$
 (5.16)

that is, operator \mathbf{M}' satisfies the quantum A_{∞} relations on \mathcal{H}' . Furthermore, operators $\boldsymbol{\pi}'$ and $\boldsymbol{\iota}'$ meet the chain map conditions:

$$(\mathbf{M}' + i\hbar \mathbf{U}') \boldsymbol{\pi}' = \boldsymbol{\pi}' (\mathbf{M} + i\hbar \mathbf{U}),$$

$$\boldsymbol{\iota}' (\mathbf{M}' + i\hbar \mathbf{U}') = (\mathbf{M} + i\hbar \mathbf{U}) \boldsymbol{\iota}'.$$
 (5.17)

To apply the HPL to adding stub, we define two degree-even maps π, ι by

$$\pi e_a(x) = \iota e_a(x) = K_{\lambda/2} e_a(x), \quad \pi \tilde{e}^a(x) = \iota \tilde{e}^a(x) = K_{\lambda/2} \tilde{e}^a(x).$$
(5.18)

Here, $K_{\lambda/2}$ is a heat kernel labeled by the positive parameter λ . The degree-odd map η that satisfies (5.3) for these π and ι is defined by

$$\eta = (1 - K_{\lambda}) h. \tag{5.19}$$

Let us see the example of the stub in scalar field theory. In such a case, we define

$$K_{\lambda} = e^{-\lambda(-\partial^2 + m^2)}, \qquad (5.20)$$

by using the heat kernel of a scalar field. Then, the basis vectors of \mathcal{H} is projected as follows:

$$\pi c(x) = \iota c(x) = e^{-\frac{\lambda}{2}(-\partial^2 + m^2)} c(x), \quad \pi d(x) = \iota d(x) = e^{-\frac{\lambda}{2}(-\partial^2 + m^2)} d(x), \quad (5.21)$$

where we used the basis vectors defined in §3.1. We will see that this projection serves as the regularization by analyzing the contact terms in the Ward-Takahashi identities. In addition, note that the limit $\lambda \to 0$ corresponds to the original theory without stubs.

Unfortunately, these π , ι and η do not satisfy the conditions (5.2) and (5.4) except $\eta^2 = 0$. In this case, the operator **M**' does not become coderivations. According to [46], the solution is simple: we can simply pretend that these conditions hold. More precise treatment is done in Appendix A of [47]. Alternatively, we can ensure that the conditions are met by defining the product appropriately with respect to $K_{\lambda/2}$ and K_{λ} . If we define the products as

$$K_{\lambda/2} \cdot K_{\lambda/2} = K_{\lambda/2} \cdot K_{\lambda} = K_{\lambda} \cdot K_{\lambda/2} = K_{\lambda} \cdot K_{\lambda} = K_{\lambda}, \qquad (5.22)$$

then the operator $\pi \iota$ can be treated as a projection.

We define the symprectic form $\langle \omega' |$ on \mathcal{H}' by

$$\langle \omega' | = \langle \omega | (\iota \otimes \iota) . \tag{5.23}$$

By using the operators \mathbf{M}' and the symplectic form $\langle \omega' |$, the stubbed action S' on \mathcal{H}' is defined as

$$S' = -\sum_{n=0}^{\infty} \frac{1}{n+1} \left\langle \omega' \right| \left(\mathbb{I} \otimes \pi_1 \mathbf{M}'_n \right) \boldsymbol{\pi} \frac{1}{1-\Phi} \,.$$
(5.24)

This action is quasi-isomorphic to the original action and reproduces the original action with the limit $\lambda \to 0$. Starting from this stubbed action, we can construct the counterparts of the correlation function **f 1** in the stubbed theory **f' 1** as

$$\mathbf{f}' \mathbf{1} = \frac{1}{\mathbf{I} + \mathbf{h}' \left(\mathbf{m}' + i\hbar \mathbf{U}' \right)} \mathbf{1}, \qquad (5.25)$$

where \mathbf{m}' and \mathbf{h}' are defined by

$$\mathbf{m}' = \mathbf{M}' - \boldsymbol{\pi} \mathbf{Q} \boldsymbol{\iota} ,$$

$$\mathbf{h}' = \boldsymbol{\pi} \mathbf{h} \boldsymbol{\iota} .$$
(5.26)

Here, the operators \mathbf{h}' is defined to satisfy

$$\pi \mathbf{Q} \boldsymbol{\iota} \, \mathbf{h}' + \mathbf{h}' \, \pi \, \mathbf{Q} \, \boldsymbol{\iota} = \mathbf{I} \,. \tag{5.27}$$

As discussed in § 3, the algebraic Schwinger-Dyson equation holds for \mathbf{f}' :

$$\left(\mathbf{M}' + i\hbar\mathbf{U}'\right)\mathbf{f}'\mathbf{1} = 0.$$
(5.28)

Let us consider the transformation \mathbf{A}' defined by

$$\mathbf{A}' = \boldsymbol{\pi} \, \mathbf{A} \, \boldsymbol{\iota} \,, \tag{5.29}$$

as counterpart of the symmetry transformation **A**. This **A'** is not a symmetry transformation because it does not commute with **M'** for $\lambda > 0$. However, for $\lambda \to 0$, it commutes with **M'** and becomes a symmetry transformation. By repeating the arguments of § 4 using $\mathbf{f}' \mathbf{1}$ and \mathbf{A}' , and finally setting $\lambda \to 0$, we obtain the regularized anomalous term:

$$-i\hbar \lim_{\lambda \to 0} \sum_{a} \langle \omega' | (\pi_{1} \mathbf{A}' \boldsymbol{\pi} \mathbf{e}_{a}(x) \boldsymbol{\iota} \otimes \boldsymbol{\pi} \, \tilde{e}^{a}(x)) \mathbf{f}' \mathbf{1}$$

$$= -i\hbar \lim_{\lambda \to 0} \sum_{a} \langle \omega | (\iota \boldsymbol{\pi} \pi_{1} \mathbf{A} \boldsymbol{\iota} \boldsymbol{\pi} \mathbf{e}_{a}(x) \boldsymbol{\iota} \otimes \iota \boldsymbol{\pi} \, \tilde{e}^{a}(x)) \mathbf{f}' \mathbf{1}.$$
(5.30)

Since the projection $\iota \pi$ satisfies

$$\langle \omega | (\iota \pi \otimes \mathbb{I}) = \langle \omega | (\mathbb{I} \otimes \iota \pi) = \langle \omega | (\iota \pi \otimes \iota \pi), \iota \pi e_a(x) \otimes \tilde{e}^a(x) = e_a(x) \otimes \iota \pi \tilde{e}^a(x) = \iota \pi e_a(x) \otimes \iota \pi \tilde{e}^a(x),$$

$$(5.31)$$

the regularized anomalous term can be also expressed as

$$-i\hbar \lim_{\lambda \to 0} \sum_{a} \langle \omega | \left(\pi_1 \mathbf{A} \, \boldsymbol{\iota} \, \boldsymbol{\pi} \, \mathbf{e}_a(x) \boldsymbol{\iota} \otimes \widetilde{e}^{\,a}(x) \right) \mathbf{f}' \, \mathbf{1} \,.$$
 (5.32)

In particular, when the symmetry transformation \mathbf{A} is linear as in the examples in the previous section, it can be simplified as

$$-i\hbar \lim_{\lambda \to 0} \sum_{a} \langle \omega | \left(A K_{\lambda} e_{a}(x) \otimes \widetilde{e}^{a}(x) \right).$$
(5.33)

Due to the smearing effect by the heat kernel over the delta function, this term is regularized.

Let us compute the anomaly for the example in the previous section. We first consider the case of the U(1) symmetry in complex scalar theories. The symmetry transformation A is defined by (4.59), and the stub is chosen as

$$\pi c(x) = \iota c(x) = e^{-\frac{\lambda}{2}(-\partial^2 + m^2)} c(x), \quad \pi d(x) = \iota d(x) = e^{-\frac{\lambda}{2}(-\partial^2 + m^2)} d(x),$$

$$\pi c^{\dagger}(x) = \iota c^{\dagger}(x) = e^{-\frac{\lambda}{2}(-\partial^2 + m^2)} c^{\dagger}(x), \quad \pi d^{\dagger}(x) = \iota d^{\dagger}(x) = e^{-\frac{\lambda}{2}(-\partial^2 + m^2)} d^{\dagger}(x).$$
(5.34)

Then, the anomalous term can be evaluated as

$$-i\hbar \lim_{\lambda \to 0} \sum_{a \in \{c,c^{\dagger}\}} \langle \omega | \left(A K_{\lambda} e_{a}(x) \otimes \widetilde{e}^{a}(x) \right) \\ = \hbar \alpha \lim_{\lambda \to 0} \lim_{y \to x} \left[e^{-\lambda(-\partial_{x}^{2} + m^{2})} \delta^{d}(x-y) - e^{-\lambda(-\partial_{x}^{2} + m^{2})} \delta^{d}(x-y) \right] \\ = i\hbar \alpha \lim_{\lambda \to 0} \left[\int \frac{\mathrm{d}^{d}k}{(2\pi)^{d}} e^{-\lambda(k^{2} + m^{2})} - \int \frac{\mathrm{d}^{d}k}{(2\pi)^{d}} e^{-\lambda(k^{2} + m^{2})} \right] \\ = 0,$$

$$(5.35)$$

where we carried out the Wick rotation in the third line. Therefore, we confirm that there is no anomaly in this case. Let us next consider the case of the energy-momentum tensor in scalar theories. The symmetry transformation A is defined by (4.91) and the stub is the same as above. Then, the anomalous term can be evaluated as

$$-i\hbar \lim_{\lambda \to 0} \langle \omega | \left(A K_{\lambda} c(x) \otimes d(x) \right) \\ = -i\hbar a_{\mu} \lim_{\lambda \to 0} \lim_{y \to x} e^{-\lambda(-\partial_{x}^{2} + m^{2})} \partial_{x}^{\mu} \delta^{d}(x-y) \\ = i\hbar a_{\mu} \lim_{\lambda \to 0} \int \frac{\mathrm{d}^{d}k}{(2\pi)^{d}} k^{\mu} e^{-\lambda(k^{2} + m^{2})} \\ = 0.$$
(5.36)

Again, we carried out the Wick rotation in the third line. Therefore, we confirm that there is no anomaly in this case.

Finally, let us consider the chiral U(1) symmetry in vector-like gauge theories. The symmetry transformation is defined by (4.119), and the stub can be chosen as

The anomalous term can be calculated as

$$- i\hbar \lim_{\lambda \to 0} \left[\langle \omega | \left(A K_{\lambda} \theta_{s}(x) \otimes \overline{\lambda}_{s}(x) \right) + \langle \omega | \left(A K_{\lambda} \overline{\theta}_{s}(x) \otimes \lambda_{s}(x) \right) \right] \\ = \hbar \alpha \lim_{\lambda \to 0} \lim_{y \to x} \operatorname{tr} \left[\gamma_{5} e^{-\lambda \overline{p}^{2}(x)} \delta^{4}(x-y) + \delta^{4}(x-y) e^{-\lambda \overline{p}^{2}(x)} \gamma_{5} \right] \\ = \hbar \alpha \lim_{\lambda \to 0} \int \frac{\mathrm{d}^{4}k}{(2\pi)^{4}} \operatorname{tr} \left[\gamma_{5} e^{-ikx} e^{-\lambda \overline{p}^{2}(x)} e^{ikx} + \gamma_{5} e^{ikx} e^{-\lambda \overline{p}^{2}(x)} e^{-ikx} \right],$$
(5.38)

where we carried out the Wick rotation in the last line. As is well known, this quantity can be evaluated as follows. Since

$$e^{-ikx} D_{\mu} e^{ikx} = D_{\mu} + i k_{\mu}, \qquad (5.39)$$

we have

$$e^{-ikx} e^{-\lambda \mathcal{D}^2(x)} e^{ikx} = e^{-\lambda(k-iD)^2 + \frac{i\lambda}{2}\gamma^{\mu}\gamma^{\nu}F_{\mu\nu}}$$
(5.40)

where $F_{\mu\nu}$ denotes the field strength of background field A_{μ}

$$F_{\mu\nu} = i \left[D_{\mu} , D_{\nu} \right]. \tag{5.41}$$

Integration over k yields a factor λ^{-2} and the formulas

$$\operatorname{tr} \gamma_5 = \operatorname{tr} \gamma_5 \gamma^{\mu} \gamma^{\nu} = 0, \quad \operatorname{tr} \gamma_5 \gamma^{\mu} \gamma^{\nu} \gamma^{\rho} \gamma^{\sigma} = -4 \, i \, \epsilon^{\mu\nu\rho\sigma}, \qquad (5.42)$$

leads

$$-i\hbar \lim_{\lambda \to 0} \left[\left\langle \omega \right| \left(A K_{\lambda} \theta_{s}(x) \otimes \overline{\lambda}_{s}(x) \right) + \left\langle \omega \right| \left(A K_{\lambda} \overline{\theta}_{s}(x) \otimes \lambda_{s}(x) \right) \right] \\ = \hbar \alpha \lim_{\lambda \to 0} \int \frac{\mathrm{d}^{4}k}{(2\pi)^{4}} \operatorname{tr} \left[\gamma_{5} e^{-ikx} e^{-\lambda \overline{\mathcal{P}}^{2}(x)} e^{ikx} + \gamma_{5} e^{ikx} e^{-\lambda \overline{\mathcal{P}}^{2}(x)} e^{-ikx} \right] \\ = -\frac{i\hbar \alpha}{64 \pi^{2}} \operatorname{tr} [\gamma_{5} \gamma^{\mu} \gamma^{\nu} \gamma^{\rho} \gamma^{\sigma}] F_{\mu\nu} F_{\rho\sigma}$$

$$= -\frac{\hbar \alpha}{16 \pi^{2}} \epsilon^{\mu\nu\rho\sigma} F_{\mu\nu} F_{\rho\sigma} , \qquad (5.43)$$

where we carried out the Wick rotation in the third line. This quantity is a well-known anomaly.¹⁵

6 Conclusions and discussion

In this paper, we introduced the identity

$$\left(\mathbf{Q} + \mathbf{m} + i\hbar \mathbf{U}\right)\mathbf{f}\mathbf{1} = 0, \qquad (6.1)$$

which we call the *algebraic Schwinger-Dyson equation* and confirm that this is exactly the Schwinger-Dyson equation in terms of homotopy algebras. Then, we applied this identity to the formula for the expectation values of the derivative of the Noether current, and obtained the Ward-Takahashi identity in terms of homotopy algebras:

$$\langle \partial_{\mu} j^{\mu}(x) \rangle = -i\hbar \sum_{a} \langle \omega | (\pi_1 \mathbf{A} \mathbf{e}_a(x) \otimes \widetilde{e}^a(x)) \mathbf{f} \mathbf{1}.$$
 (6.2)

In our formula, the terms associated with

$$\sum_{a} \langle \omega | (\pi_1 \mathbf{A} \mathbf{e}_a(x) \otimes \widetilde{e}^a(x)) \mathbf{f} \mathbf{1}$$
(6.3)

contain divergence. As we discussed in §4.3 and §5, these terms are related to contact terms and anomalies, and we need appropriate regularizations. In our treatment, adding stubs makes the theory finite, and then we consider the original theory as the limit of the stub parameter $\lambda \to 0$. Moreover, these divergences are related to the renormalization of the composite operators. The process we make a replacement

$$\frac{1}{1-\Phi} \longrightarrow \mathbf{f} \, \mathbf{1} \tag{6.4}$$

in §4 corresponds to take the expectation values of fields. We formally dealt with this process in the general case. In the examples described in §4.4, the results are consistent with the

¹⁵The ordinary derivation in quantum field theory is written in many textbooks. See, for example, §77 of [51].

results by ordinary prescriptions. We expect that our treatment is consistent with the ordinary prescriptions in quantum field theory in general. The treatment of these divergences originates from how to define the composite operator in terms of homotopy algebras, and it is important to consider this problem deeply.

Let us also comment on the extension to the gauge theory. In our example, we treated the background gauge field but the extension of the dynamical gauge field is straightforward. In this paper, the gauge is fixed at the beginning since we are considering correlation functions. This prescription provides a simple algebraic description, but the gauge symmetry is implicit. It is interesting and important to consider the direct relation to [45, 50].

Let us mention future works. The motivation of this work is to apply this method to string field theory. Since the description of homotopy algebras is universal, we hope that this work contributes to finding and analyzing symmetries and anomalies in string field theory. Our work is based on the perturbative expansion, however, recently one of the authors are working on the nonperturbative description of correlation functions in homotopy algebras [54]. Combined with these works, we hope that our work contributes to describing the nonperturbative anomaly from the homotopy algebraic point of view.

We can also consider to extend this work to the higher form symmetry [55]. In the context of higher form symmetry, the understanding of the Ward-Takahashi identity plays an important role in describing the symmetry. Our formula enables us to deal with the Ward-Takahashi identity from the algebraic point of view. We expect that the algebraic approach for the Ward-Takahashi identity provides effective tools to analyze the problem.

Lastly, it is interesting to develop our method to define quantum field theory from a mathematical point of view. It is important to relate our method to the formalism developed by Costello and Gwilliam using factorization algebras [56, 57].

We hope that our research will contribute to revealing the quantum aspect of quantum field theory and string field theory.

Acknowledgments

We would like to thank Yuji Okawa for discussions and comments on the earlier version of the draft, and Hiroaki Matsunaga, Hayato Kanno, and Masashi Kawahira for discussions. J. T. Y. would like to thank Hiroshi Kunitomo and Taichi Tsukamoto for discussions. The work of J. T. Y. is supported in part by JSPS KAKENHI Grant Nos. JP23KJ1311.

A **Proof:** $(2.14) \Leftrightarrow (2.46)$

For notational simplicity, in this section index a and x are written collectively as A. First, we show $(2.14) \Rightarrow (2.46)$. Let us consider the quantity

$$\sum_{A} \left[M_n(e_A \otimes \mathbb{I}^{\otimes n-1}) \otimes \tilde{e}_A + M_n(\tilde{e}_A \otimes \mathbb{I}^{\otimes n-1}) \otimes e_A \right].$$
(A.1)

By using the normalization (2.45), this quantity can be rewritten as

$$\sum_{A} \left[M_{n}(e_{A} \otimes \mathbb{I}^{\otimes n-1}) \otimes \tilde{e}^{A} + M_{n}(\tilde{e}_{A} \otimes \mathbb{I}^{\otimes n-1}) \otimes e_{A} \right]$$

$$= \sum_{A,B} \left[(-1)^{e_{B}} \tilde{e}^{B} \otimes \omega \left(e_{B}, M_{n}(e_{A} \otimes \mathbb{I}^{\otimes n-1}) \right) \otimes \tilde{e}^{A} + (-1)^{\tilde{e}^{B}} e_{B} \otimes \omega \left(\tilde{e}^{B}, M_{n}(e_{A} \otimes \mathbb{I}^{\otimes n-1}) \right) \otimes \tilde{e}^{A} + (-1)^{e_{B}} \tilde{e}^{B} \otimes \omega \left(e_{B}, M_{n}(\tilde{e}^{A} \otimes \mathbb{I}^{\otimes n-1}) \right) \otimes e_{A} + (-1)^{\tilde{e}_{B}} e_{B} \otimes \omega \left(\tilde{e}^{B}, M_{n}(\tilde{e}^{A} \otimes \mathbb{I}^{\otimes n-1}) \right) \otimes e_{A} \right]$$

$$= \sum_{A,B} \left[(-1)^{e_{A}e_{B}} \tilde{e}_{B} \otimes \tilde{e}^{A} \otimes \omega \left(e_{B}, M_{n}(e_{A} \otimes \mathbb{I}^{\otimes n-1}) \right) + (-1)^{\tilde{e}_{A}e_{B}} \tilde{e}_{B} \otimes e_{A} \otimes \omega \left(\tilde{e}^{B}, M_{n}(\tilde{e}^{A} \otimes \mathbb{I}^{\otimes n-1}) \right) + (-1)^{\tilde{e}^{A}\tilde{e}_{B}} e_{B} \otimes e_{A} \otimes \omega \left(\tilde{e}_{B}, M_{n}(\tilde{e}^{A} \otimes \mathbb{I}^{\otimes n-1}) \right) \right]. \quad (A.2)$$

In the second equality, we used that $\omega(...)$ is an element of $\mathcal{H}^{\otimes 0}$ and can be commuted with any element of \mathcal{H} with appropriate sign. If we assume (2.14), ω 's inputs can be replaced cyclically with appropriate sign and we find that the first and third terms in the left-hand side of (A.2) are

$$\sum_{B} -(-1)^{\tilde{e}_{B}} \tilde{e}^{B} \otimes \left[(-1)^{e_{A}} \tilde{e}^{A} \otimes \omega \left(e_{A}, M_{n}(\mathbb{I}^{\otimes n-1} \otimes e_{B}) \right) + (-1)^{\tilde{e}^{A}} e_{A} \otimes \omega \left(\tilde{e}^{A}, M_{n}(\mathbb{I}^{\otimes n-1} \otimes e_{B}) \right) \right]$$
$$= \sum_{B} -(-1)^{\tilde{e}^{B}} \tilde{e}^{B} \otimes M_{n}(\mathbb{I}^{\otimes n-1} \otimes e_{B}), \qquad (A.3)$$

where we used (2.45) again. In the same way, the second and fourth terms in the left-hand side of (A.2) are

$$\sum_{B} -(-1)^{e_{B}} e_{B} \otimes M_{n}(\mathbb{I}^{\otimes n-1} \otimes \widetilde{e}^{B}) .$$
(A.4)

This completes the proof of $(2.14) \Rightarrow (2.46)$. On the other hand, the proof of $(2.46) \Rightarrow (2.14)$ can be obtained by following the above proof in reverse.

B The proof for the Schwinger-Dyson equations for scalar field theory

Following [34, 35, 36], let us show that the formula

$$\pi_n \mathbf{f} \, \mathbf{1} = \langle \, \Phi^{\otimes n} \, \rangle \tag{B.1}$$

satisfies the Schwinger-Dyson equations. In the path integral formalism, correlation functions are defined by

$$\langle \varphi(x_1)\varphi(x_2)\dots\varphi(x_n)\rangle = \frac{1}{Z}\int \mathcal{D}\varphi\,\varphi(x_1)\varphi(x_2)\dots\varphi(x_n)e^{\frac{i}{\hbar}S},$$
 (B.2)

where

$$Z = \int \mathcal{D}\varphi \, e^{\frac{i}{\hbar}S} \,. \tag{B.3}$$

Since

$$\frac{1}{Z} \int \mathcal{D}\varphi \, \frac{\delta}{\delta\varphi(x_n)} \Big[\varphi(x_1)\varphi(x_2)\dots\varphi(x_{n-1})e^{\frac{i}{\hbar}S}\Big] = 0\,, \tag{B.4}$$

the Schwinger-Dyson equations

$$\sum_{i=1}^{n-1} \langle \varphi(x_1) \dots \varphi(x_{i-1}) \delta^d(x_i - x_n) \varphi(x_{i+1}) \dots \varphi(x_{n-1}) \rangle + \frac{i}{\hbar} \langle \varphi(x_1) \varphi(x_2) \dots \varphi(x_{n-1}) \frac{\delta S}{\delta \varphi(x_n)} \rangle = 0$$
(B.5)

are derived.

Let us show (B.5) in terms of homotopy algebras. Since

$$\pi_n \mathbf{f}^{-1} \mathbf{f} \, \mathbf{1} = 0 \tag{B.6}$$

for n > 0, the relations

$$\pi_n \mathbf{f} \, \mathbf{1} + \pi_n \, \mathbf{h} \, \mathbf{m} \, \mathbf{f} \, \mathbf{1} + i\hbar \, \pi_n \, \mathbf{h} \, \mathbf{U} \, \mathbf{f} \, \mathbf{1} = 0 \tag{B.7}$$

hold. We act the operator

$$\langle \omega_n(x_1, \dots, x_n) | = (-1)^n \langle \omega | (d(x_1) \otimes \mathbb{I}) \cdots \langle \omega | (d(x_n) \otimes \mathbb{I})$$
(B.8)

from the above relations:

$$\langle \omega_n(x_1,\ldots,x_n) | \pi_n \mathbf{f} \mathbf{1} + \langle \omega_n(x_1,\ldots,x_n) | \pi_n \mathbf{h} \mathbf{m} \mathbf{f} \mathbf{1} + i\hbar \langle \omega_n(x_1,\ldots,x_n) | \pi_n \mathbf{h} \mathbf{U} \mathbf{f} \mathbf{1} = 0.$$
(B.9)

Since

$$\pi_n \mathbf{f} \, \mathbf{1} = \langle \, \Phi^{\otimes n} \, \rangle \,, \tag{B.10}$$

the first term of the left-hand side is

$$\langle \omega_n(x_1, \ldots, x_n) | \pi_n \mathbf{f} \mathbf{1} = \langle \varphi(x_1) \varphi(x_2) \ldots \varphi(x_n) \rangle.$$
 (B.11)

Let us next consider the second term of the left-hand side of (B.9).

$$\pi_n \mathbf{h} \mathbf{m} \mathbf{f} \mathbf{1} = \sum_{k=0}^{\infty} \mathbb{I}^{\otimes (n-1)} \otimes hm_k(\Phi^{\otimes k}).$$
(B.12)

Notice that

$$m_k(\Phi^{\otimes k}) = -\int \mathrm{d}^d x \, d(x) \, \frac{\delta S_{\rm int}}{\delta \varphi(x)} \,, \tag{B.13}$$

where S_{int} is the interaction part of the action. Then, we obtain

$$\pi_n \mathbf{h} \mathbf{m} \mathbf{f} \mathbf{1} = -\int \mathrm{d}^d x \, \mathrm{d}^d y \, \Phi^{\otimes (n-1)} \otimes \Delta(x-y) \, c(y) \frac{\delta S_{\mathrm{int}}}{\delta \varphi(x)} \,. \tag{B.14}$$

Acting the operator (B.8),

$$\langle \omega_n(x_1,\ldots,x_n) | \pi_n \mathbf{h} \mathbf{m} \mathbf{f} \mathbf{1} = -\int d^d x \left\langle \varphi(x_1) \ldots \varphi(x_{n-1}) \Delta(x-x_n) \frac{\delta S_{\text{int}}}{\delta \varphi(x)} \right\rangle.$$
 (B.15)

Let us next consider the third term of the left-hand side of (B.9):

$$i\hbar \pi_{n} \mathbf{h} \mathbf{U} \mathbf{f} \mathbf{1}$$

$$= i\hbar \sum_{i=1}^{n-1} \int \mathrm{d}^{d} x \langle \Phi^{\otimes (i-1)} \otimes c(x) \otimes \Phi^{\otimes (n-i-1)} \otimes h d(x) \rangle$$

$$= i\hbar \sum_{i=1}^{n-1} \int \mathrm{d}^{d} x \mathrm{d}^{d} y \langle \Phi^{\otimes (i-1)} \otimes c(x) \otimes \Phi^{\otimes (n-i-1)} \otimes \Delta(x-y) c(y) \rangle.$$
(B.16)

Then, we obtain

$$i\hbar \langle \omega_n(x_1, \ldots, x_n) | \pi_n \mathbf{h} \mathbf{U} \mathbf{f} \mathbf{1} = i\hbar \sum_{i=1}^{n-1} \int d^d x \, \Delta(x_i - x_n) \langle \varphi(x_1) \ldots \varphi(x_{i-1}) \varphi(x_{i+1}) \ldots \varphi(x_{n-1}) \rangle.$$
(B.17)

Therefore, the relations (B.9) become

$$\langle \varphi(x_1) \varphi(x_2) \dots \varphi(x_n) \rangle - \int d^d x \left\langle \varphi(x_1) \dots \varphi(x_{n-1}) \Delta(x - x_n) \frac{\delta S_{\text{int}}}{\delta \varphi(x)} \right\rangle + i\hbar \sum_{i=1}^{n-1} \int d^d x \, \Delta(x_i - x_n) \langle \varphi(x_1) \dots \varphi(x_{i-1}) \varphi(x_{i+1}) \dots \varphi(x_{n-1}) \rangle = 0.$$
 (B.18)

Acting the operator $(-\partial_{x_n}^2 + m^2)$, we obtain

$$-\left\langle \varphi(x_1) \dots \varphi(x_{n-1}) \frac{\delta S}{\delta \varphi(x_n)} \right\rangle + i\hbar \sum_{i=1}^{n-1} \delta(x_i - x_n) \left\langle \varphi(x_1) \dots \varphi(x_{i-1}) \varphi(x_{i+1}) \dots \varphi(x_{n-1}) \right\rangle = 0.$$
(B.19)

This is exactly the Schwinger-Dyson equations (B.5).

References

- J. D. Stasheff, "Homotopy associativity of *H*-spaces. I," Trans. of the Amer. Math. Soc. 108, 275 (1963).
- [2] J. D. Stasheff, "Homotopy associativity of H-spaces. II," Trans. of the Amer. Math. Soc. 108, 293 (1963).
- [3] E. Getzler and J. D. S. Jones, " A_{∞} -algebras and the cyclic bar complex," Illinois J. Math **34**, 256 (1990).
- [4] M. Markl, "A cohomology theory for A(m)-algebras and applications," J. Pure Appl. Algebra 83, 141 (1992).
- [5] M. Penkava and A. S. Schwarz, " A_{∞} algebras and the cohomology of moduli spaces," hep-th/9408064.
- M. R. Gaberdiel and B. Zwiebach, "Tensor constructions of open string theories. 1: Foundations," Nucl. Phys. B505, 569 (1997) [hep-th/9705038].
- B. Zwiebach, "Closed string field theory: Quantum action and the Batalin-Vilkovisky master equation," Nucl. Phys. B 390, 33-152 (1993) [arXiv:hep-th/9206084 [hep-th]].
- [8] M. Markl, "Loop homotopy algebras in closed string field theory," Commun. Math. Phys. 221, 367-384 (2001) [arXiv:hep-th/9711045 [hep-th]].
- C. Maccaferri and J. Vošmera, "The classical cosmological constant of open-closed string field theory," JHEP 10, 173 (2022) [arXiv:2208.00410 [hep-th]].
- [10] C. Maccaferri, A. Ruffino and J. Vošmera, "The nilpotent structure of open-closed string field theory," JHEP 08, 145 (2023) [arXiv:2305.02843 [hep-th]].
- [11] H. Kajiura and J. Stasheff, "Homotopy algebras inspired by classical open-closed string field theory," Commun. Math. Phys. 263, 553-581 (2006) [arXiv:math/0410291 [math.QA]].
- [12] H. Kajiura and J. Stasheff, "Open-closed homotopy algebra in mathematical physics," J. Math. Phys. 47, 023506 (2006) [arXiv:hep-th/0510118 [hep-th]].
- [13] T. Erler, S. Konopka and I. Sachs, "Resolving Witten's superstring field theory," JHEP 04, 150 (2014) [arXiv:1312.2948 [hep-th]].
- [14] T. Erler, Y. Okawa and T. Takezaki, "Complete Action for Open Superstring Field Theory with Cyclic A_{∞} Structure," JHEP **08**, 012 (2016) [arXiv:1602.02582 [hep-th]].

- [15] H. Kunitomo and T. Sugimoto, "Heterotic string field theory with cyclic L_{∞} structure," PTEP **2019**, no.6, 063B02 (2019) [arXiv:1902.02991 [hep-th]].
- [16] H. Kunitomo, "Open-closed homotopy algebra in superstring field theory," PTEP 2022, no.9, 093B07 (2022) [arXiv:2204.01249 [hep-th]].
- [17] H. Kajiura, "Noncommutative homotopy algebras associated with open strings," Rev. Math. Phys. 19, 1-99 (2007) [arXiv:math/0306332 [math.QA]].
- [18] A. Sen, "Wilsonian Effective Action of Superstring Theory," JHEP 01, 108 (2017) [arXiv:1609.00459 [hep-th]].
- [19] H. Erbin, C. Maccaferri, M. Schnabl and J. Vošmera, "Classical algebraic structures in string theory effective actions," JHEP 11, 123 (2020) [arXiv:2006.16270 [hep-th]].
- [20] D. Koyama, Y. Okawa and N. Suzuki, "Gauge-invariant operators of open bosonic string field theory in the low-energy limit," [arXiv:2006.16710 [hep-th]].
- [21] A. S. Arvanitakis, O. Hohm, C. Hull and V. Lekeu, "Homotopy Transfer and Effective Field Theory I: Tree-level," Fortsch. Phys. 70, no.2-3, 2200003 (2022) [arXiv:2007.07942 [hep-th]].
- [22] A. S. Arvanitakis, O. Hohm, C. Hull and V. Lekeu, "Homotopy Transfer and Effective Field Theory II: Strings and Double Field Theory," Fortsch. Phys. 70, no.2-3, 2200004 (2022) [arXiv:2106.08343 [hep-th]].
- [23] R. Bonezzi, C. Chiaffrino, F. Diaz-Jaramillo and O. Hohm, "Tree-level Scattering Amplitudes via Homotopy Transfer," [arXiv:2312.09306 [hep-th]].
- [24] S. Konopka, "The S-Matrix of superstring field theory," JHEP 11, 187 (2015) [arXiv:1507.08250 [hep-th]].
- [25] H. Kunitomo, "Tree-level S-matrix of superstring field theory with homotopy algebra structure," JHEP 03, 193 (2021) [arXiv:2011.11975 [hep-th]].
- [26] T. Erler and H. Matsunaga, "Mapping between Witten and lightcone string field theories," JHEP 11, 208 (2021) [arXiv:2012.09521 [hep-th]].
- [27] O. Hohm and B. Zwiebach, " L_{∞} Algebras and Field Theory," Fortsch. Phys. 65, no.3-4, 1700014 (2017) [arXiv:1701.08824 [hep-th]].
- [28] B. Jurčo, L. Raspollini, C. Sämann and M. Wolf, " L_{∞} -Algebras of Classical Field Theories and the Batalin-Vilkovisky Formalism," Fortsch. Phys. **67**, no.7, 1900025 (2019) [arXiv:1809.09899 [hep-th]].
- [29] A. Nützi and M. Reiterer, "Amplitudes in YM and GR as a Minimal Model and Recursive Characterization," Commun. Math. Phys. **392**, no.2, 427-482 (2022) [arXiv:1812.06454 [mathph]].
- [30] A. S. Arvanitakis, "The L_{∞} -algebra of the S-matrix," JHEP **07**, 115 (2019) [arXiv:1903.05643 [hep-th]].

- [31] T. Macrelli, C. Sämann and M. Wolf, "Scattering amplitude recursion relations in Batalin-Vilkovisky-quantizable theories," Phys. Rev. D 100, no.4, 045017 (2019) [arXiv:1903.05713 [hepth]].
- [32] B. Jurčo, T. Macrelli, C. Sämann and M. Wolf, "Loop Amplitudes and Quantum Homotopy Algebras," JHEP 07, 003 (2020) [arXiv:1912.06695 [hep-th]].
- [33] C. Saemann and E. Sfinarolakis, "Symmetry Factors of Feynman Diagrams and the Homological Perturbation Lemma," JHEP **12**, 088 (2020) [arXiv:2009.12616 [hep-th]].
- [34] Y. Okawa, "Correlation functions of scalar field theories from homotopy algebras," [arXiv:2203.05366 [hep-th]].
- [35] K. Konosu and Y. Okawa, "Correlation functions involving Dirac fields from homotopy algebras I: the free theory," [arXiv:2305.11634 [hep-th]].
- [36] K. Konosu, "Correlation functions involving Dirac fields from homotopy algebras II: the interacting theory," [arXiv:2305.13103 [hep-th]].
- [37] I. A. Batalin and G. A. Vilkovisky, "Gauge Algebra and Quantization," Phys. Lett. B 102, 27-31 (1981).
- [38] I. A. Batalin and G. A. Vilkovisky, "Quantization of Gauge Theories with Linearly Dependent Generators," Phys. Rev. D 28, 2567-2582 (1983) [erratum: Phys. Rev. D 30, 508 (1984)].
- [39] A. S. Schwarz, "Geometry of Batalin-Vilkovisky quantization," Commun. Math. Phys. 155, 249-260 (1993) [arXiv:hep-th/9205088 [hep-th]].
- [40] O. Gwilliam and T. Johnson-Freyd, "How to derive Feynman diagrams for finite-dimensional integrals directly from the BV formalism," [arXiv:1202.1554 [math-ph]].
- [41] C. Chiaffrino, O. Hohm and A. F. Pinto, "Homological quantum mechanics," JHEP 02, 137 (2024) [arXiv:2112.11495 [hep-th]].
- [42] T. Masuda and H. Matsunaga, "Perturbative path-integral of string fields and the A_{∞} structure of the BV master equation," PTEP **2022**, no.11, 113B04 (2022) [arXiv:2003.05021 [hep-th]].
- [43] M. Doubek, B. Jurčo and J. Pulmann, "Quantum L_{∞} Algebras and the Homological Perturbation Lemma," Commun. Math. Phys. **367**, no.1, 215-240 (2019) [arXiv:1712.02696 [math-ph]].
- [44] T. Erler, "Supersymmetry in Open Superstring Field Theory," JHEP 05, 113 (2017) [arXiv:1610.03251 [hep-th]].
- [45] H. Matsunaga, "Homotopy algebra & symmetry generators in QFT," talk in the workshop of Homotopy Algebra of Quantum Field Theory and Its Application, Yukawa Institute for Theoretical Physics, Kyoto University, Japan, March 2021.
- [46] M. Schnabl and G. Stettinger, "Open string field theory with stubs," JHEP 07, 032 (2023) [arXiv:2301.13182 [hep-th]].
- [47] H. Erbin and A. H. Fırat, "Open string stub as an auxiliary string field," [arXiv:2308.08587 [hep-th]].

- [48] M. Schnabl and G. Stettinger, "More on stubs in open string field theory," [arXiv:2402.00308 [hep-th]].
- [49] C. Maccaferri, R. Poletti, A. Ruffino and B. Valsesia, "Adding stubs to quantum string field theories," [arXiv:2403.10471 [hep-th]].
- [50] C. Chiaffrino and I. Sachs, "QFT with stubs," JHEP 06, 120 (2022) [arXiv:2108.04312 [hep-th]].
- [51] M. Srednicki, "Quantum Field Theory," Cambridge University Press (2007).
- [52] T. Erler, "Relating Berkovits and A_{∞} superstring field theories; small Hilbert space perspective," JHEP 10, 157 (2015) [arXiv:1505.02069 [hep-th]].
- [53] M. Crainic, "On the perturbation lemma, and deformations," [arXiv:math/0403266 [math.AT]].
- [54] K. Konosu and Y. Okawa, "Nonperturbative correlation functions from homotopy algebras," in preparation
- [55] D. Gaiotto, A. Kapustin, N. Seiberg and B. Willett, JHEP 02, 172 (2015) doi:10.1007/JHEP02(2015)172 [arXiv:1412.5148 [hep-th]].
- [56] K. Costello and O. Gwilliam, "Factorization Algebras in Quantum Field Theory: Volume 1," Cambridge University Press (2016).
- [57] K. Costello and O. Gwilliam, "Factorization Algebras in Quantum Field Theory: Volume 2," Cambridge University Press (2021).