Neutrino Mixing from a Fresh Perspective

Pralay Chakraborty, ^{1,*} Manash Dey, ^{1,†} Biswajit Karmakar, ^{2,‡} and Subhankar Roy^{1,§}

¹Physics Department, Gauhati University, India

²Institute of Physics, University of Silesia, Katowice, Poland

(Dated: May 20, 2024)

We propose a new mixing scheme for neutrinos which eventually can be linked to a neutrino mass matrix texture bearing a suitable correlation $(m_{22}+2\,m_{13}=0)$ among its elements. The texture predicts three mass eigenvalues in the light of normal ordering, in addition to the Majorana phases. The texture is realized from $A_4\times Z_{10}\times Z_2$ group in the framework of general type-I+II seesaw mechanism

I. INTRODUCTION

In the realm of particle physics, neutrinos are mysterious creatures that can transform from one type to another as they traverse space. This phenomenon is called neutrino oscillation and this idea is first introduced by Bruno Pontecorvo in the year 1957 [1]. The neutrino oscillation arises from the fact that the three neutrino flavour states ($\nu_{l=e,\mu,\tau}$) are admixtures of three mass eigenstates ($\nu_{i=1,2,3}$) with corresponding mass values ($m_{i=1,2,3}$). The Pontecorvo-Maki-Nakagawa-Sakata (PMNS) matrix [2] is the mathematical representation that connects the flavour states to the mass states, providing a framework for understanding how neutrinos transform between different flavours as they propagate through space. The neutrino oscillation demands the existence of a non-zero neutrino mass, contradicting the predictions of the Standard Model (SM) [3–5]. In this regard, to understand the origin of neutrino mass, it is necessary to explore theories that go beyond the SM. The neutrino mass matrix (M_{ν}) originates from the Yukawa Lagrangian, which is formulated within the framework of the seesaw mechanism [6, 7]. In the literature, the widely used seesaw mechanisms are the type-I seesaw [8–10] and type-II seesaw mechanism [11–14].

The neutrino mass matrix M_{ν} contains twelve parameters: three mass eigenvalues $(m_{i=1,2,3})$, three mixing angles $(\theta_{12}, \theta_{13}, \theta_{23})$, three CP phases (δ, α, β) , and three unphysical phases (ϕ_1, ϕ_2, ϕ_3) . In recent years, the experiments predict the mixing angles with high precision but the precise determination of Dirac CP phase δ and the octant of the atmospheric mixing angle θ_{23} yet to be settled [15]. In addition, the experiments are unable to provide exact values for individual mass eigenvalues; instead, they offer predictions for two mass-squared differences. In addition, the Majorana phases are physical parameters which are still untouched in the experiments. In order to get the information of the physical parameters, we need to diagonalize M_{ν} : diag $(m_1, m_2, m_3) = V^T.M_{\nu}.V$, where, V is the PMNS matrix. The Particle Data Group (PDG) [16] has adopted a parametrization scheme for U known as standard parametrization as shown in the following,

$$V = P_{\phi}.U.P_{M},\tag{1}$$

where, $P_{\phi} = \text{diag}(e^{i\phi_1}, e^{i\phi_2}, e^{i\phi_3})$ and $P_M = \text{diag}(e^{i\alpha}, e^{i\beta}, 1)$. The matrix U is represented as shown in the following,

$$U = \begin{bmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{bmatrix} \times \begin{bmatrix} c_{13} & 0 & s_{13} e^{-i\delta} \\ 0 & 1 & 0 \\ -s_{13} e^{i\delta} & 0 & c_{13} \end{bmatrix} \times \begin{bmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{bmatrix},$$
 (2)

where, $s_{ij} = \sin \theta_{ij}$ and $c_{ij} = \cos \theta_{ij}$. By using the standard parametrization, the Jarlskog invariant (J_{CP}) [17] can be expressed as shown below,

$$J_{CP} = s_{13}c_{13}^2 s_{12}c_{12}s_{23}c_{23}\sin\delta. (3)$$

In literature, several mixing patterns were proposed. Among them, bimaximal mixing (BM) [18], tri-bimaximal mixing (TBM) [19], hexagonal mixing (HM) [20] and golden ratio (GR) [21] are notable. All of these fixed or constant

^{*} pralay@gauhati.ac.in

[†] manashdey@gauhati.ac.in

[‡] biswajit.karmakar@us.edu.pl

[§] subhankar@gauhati.ac.in

mixing schemes predict vanishing value for the reator mixing angle θ_{13} . For instance, in tri-bimaximal mixing, specific values are assigned to the mixing parameters: $s_{12} = \frac{1}{\sqrt{3}}$, $s_{23} = \frac{1}{\sqrt{2}}$ and $s_{13} = 0$. This mixing scheme holds significance due to its inherent connection to the $\mu - \tau$ symmetric neutrino mass matrix, which is particularly fascinating from the model-building perspective [22, 23]. In this regard, the tri-bimaximal mixing has been investigated within the framework of various discrete flavour symmetries [24, 25], for a recent review on discrete flavour symmetries and their implication in neutrino mixing, see Ref. [26] and for specific examples, see Refs. [27, 28] and references there in. It is to be emphasized that the recent experiments have ruled out the possibility of $s_{13} = 0$. However, the fundamental philosophy behind the mixing pattern and its connection to specific textures of the neutrino mass matrix remains sacred. All mixing patterns such as bimaximal, tri-bimaximal, hexagonal, golden ratio are characterized by constant values of the neutrino mixing angles and can be explained within the frameworks of non-Abelian dscrete flavour symmetries. In that light, the aim of this paper is to propose a new mixing pattern in terms of a fixed or constant mixing scheme and deviating from $s_{13} = 0$ (hence accommodating Dirac CP phase δ) which is consistent with the experiments:

$$s_{12} = (2\sqrt{2})/5,$$
 $s_{13} = 1/(3\sqrt{5}),$ $s_{23} = 3/4,$ $\delta = 3\pi/4.$




FIG. 1. The comparison of the tri-bimaximal, proposed and best fit [15] values of the mixing angles and Dirac CP phase δ .

For better visualization, we have compared the experimental best fit values [15], our proposed values of the mixing angles and tri-bimaximal mixing angles as illustrated in Fig. 1. In order to derive the proposed mixing scheme from a neutrino mass matrix, we adhere to the following texture bearing a suitable correlation $m_{22} + 2m_{13} = 0$ as shown in the following,

$$M_{\nu} = \begin{bmatrix} A & B & F \\ B & -2F & G \\ F & G & J \end{bmatrix}, \tag{4}$$

where, A, B, F, G and J are complex parameters.

The plan for rest of the paper is outlined as follows: In Section (II), we show the formalism and numerical analysis to show the compatibility of the proposed mixing scheme and the mass matrix presented in Eq. (4). Predictions involving neutrino masses, associated Majorana phases, and effective mass parameters appearing in the neutrinoless double beta decay. Section (III) is devoted to the relation of the proposed mixing scheme with a specific model. As an example, here we have showed a model for the proposed mixing scheme based on A_4 discrete flavour symmetry and study of model parameters. Finally, we summarize and conclude in Section (IV).

II. FORMALISM AND NUMERICAL ANALYSIS

The structure of the PMNS matrix in the light of proposed mixing pattern can be written as,

$$U_{P} = \begin{pmatrix} \frac{2\sqrt{187}}{15\sqrt{5}} & \frac{4\sqrt{12}}{15\sqrt{5}} & -\frac{(1+i)}{3\sqrt{10}} \\ \frac{\sqrt{17}(1-i)}{20\sqrt{10}} - \frac{7}{5\sqrt{2}} & \frac{\sqrt{119}}{20} + \frac{(1-i)}{10\sqrt{5}} & \frac{\sqrt{11}}{2\sqrt{5}} \\ \frac{3}{5\sqrt{2}} + \frac{\sqrt{119}(1-i)}{60\sqrt{10}} & \frac{3\sqrt{17}}{20} + \frac{\sqrt{7}(1-i)}{30\sqrt{5}} & \frac{\sqrt{75}}{6\sqrt{5}} \end{pmatrix}.$$
 (5)




FIG. 2. Shows the correlation plots between Re[J] and Im[J] vs Re[G].




FIG. 3. Shows the correlation plots between (a) Δm_{21}^2 vs Δm_{31}^2 . (b) m_1 vs m_2 and m_3 . (c) m_1 vs Σm_i . (d) α vs β . (e) Lightest neutrino mass (m_1) vs $m_{\beta\beta}$.

We now carry out a model-independent analysis of the texture given in Eq. (4) which can be diagonalized by U_P . To get the information of the physical parameters, we diagonalize M_{ν} in the following way,

$$\operatorname{diag}(\tilde{m_1}, \tilde{m_2}, m_3) = U_P^T M_{\nu} U_P, \tag{6}$$

where, $\tilde{m_1} = m_1 e^{-2i\alpha}$ and $\tilde{m_2} = m_2 e^{-2i\beta}$, where, $m_{i=1,2,3}$ are the real positive mass eigen values and α, β are associated Majorana phases. By applying the diagonalizing conditions, we restrict the number of free parameters to three Re[G], Re[J] and Im[J] and express three mass eigenvalues and two Majorana phases in terms of the said parameters. We generate a set of random numbers for Re[G], Re[J], and Im[J] ensuring that $\Sigma m_i \leq 0.12, eV$ [29] and that the two mass squared differences align with experimental observations [15]. For graphical visualization, we obtain the correlation plots. The Fig. (2) shows how the texture parameters Re[G], Re[J], and Im[J] vary. From the Figs. 3(a)-3(d), we identify the regions to where the observational parameters are constrained, where, the correlation between the parameters in the $\Delta m_{21}^2 - \Delta m_{31}^2$, $m_1 - m_{2,3}$, $m_1 - \sum m_i$ and $\alpha - \beta$ are visualized. The maximum and minimum values of the physical parameters are listed in Table I. In addition, the Jarlskog invariant (J_{CP}) , which determines the size of the leptonic CP violation, is obtained from the proposed mixing scheme as 0.0238. The

(7)

Parameters	Minimum Value	Maximum Value
m_1/eV	0.018	0.021
m_2/eV	0.020	0.022
m_3/eV	0.052	0.055
$\sum m_i/\text{eV}$	0.091	0.098
$\alpha/^{\circ}$	-13.96	-3.18
β/°	-29.37	-0.75
$m_{\beta\beta}/\text{meV}$	17.27	20.22

TABLE I. Shows the maximum and minimum values of m_1 , m_2 , m_3 , Σm_i , α , β and $m_{\beta\beta}$.

parameter $m_{\beta\beta}$ which is the effective Majorana neutrino mass is an observational parameter and several experiments have provided the upper bounds of $m_{\beta\beta}$: SuperNEMO(Se⁸²) as 67 – 131 meV, GERDA(Ge⁷⁶) as 104 – 228 meV, EXO-200(Xe¹³⁶) as 111 – 477 meV, CUORE(Te¹³⁰) as 75 – 350 meV and KamLAND-Zen(Xe¹³⁶) as 61 – 165 meV [30–36]. In this regard, the variation of the parameter $m_{\beta\beta}$ with respect to the lightest neutrino mass is visualized from the proposed (see Fig. 3(e)). We emphasize that the texture only yields the normal hierarchy for neutrino masses when diagonalized with the proposed mixing matrix.

III. THE MODEL

To realize the texture shown in the earlier section, compatible with the proposed mixing scheme, we now consider a framework based on non-Abelian discrete flavour symmetry. As an example, we have considered A_4 discrete flavour symmetry, which is the smallest group having a three-dimensional irreducible representation where three SM lepton doublets can be accommodated and unified into the triplet representation. Here, the whole framework is embedded within $A_4 \times Z_{10} \times Z_2$ symmetry. The light neutrino mass is generated via most general type-I+II seesaw mechanism [37]. Hence, in this regard, we extend the field content of the SM by adding three right-handed neutrinos ($\nu_{e_R}, \nu_{\mu_R}, \nu_{\tau_R}$) and two scalar triplets ($\Delta_{1,2}$). To implement the flavour symmetry and appropriate texture of the mass matrix, we also include seven scalar singlets (Φ , ξ , η , χ , ψ , κ , ς). The transformation properties of the field content associated with $A_4 \times Z_{10} \times Z_2$ symmetry are summarised in Table II.

Fields	D_{l_L}	l_R	$ u_{l_R}$	Н	Δ_1	Δ_2	Φ	ξ	η	χ	ψ	κ	ς
$SU(2)_L$	2	1	1	2	3	3	1	1	1	1	1	1	1
A_4	3	(1,1'',1')	$(1'', 1, 1^{'})$	1	1	1	3	1	1"	1	3	3	1
Z_{10}	0	(0,0,0)	(0,4,8)	0	3	1	0	2	5	6	7	9	5
Z_2	1	(1, 1, 1)	(1, 1, 1)	-1	1	1	-1	1	1	1	1	1	1

TABLE II. The transformation properties of various fields under $A_4 \times Z_{10} \times Z_2$.

The $A_4 \times Z_{10} \times Z_2$ invariant Lagrangian can be constructed in the following manner,

$$\begin{split} -\mathcal{L}_{Y} &= \frac{y_{e}}{\Lambda} (\overline{D}_{l_{L}} \Phi)_{1} H \, e_{R_{1}} + \frac{y_{\mu}}{\Lambda} (\overline{D}_{l_{L}} \Phi)_{1'} \, H \mu_{R_{1''}} + \frac{y_{\tau}}{\Lambda} (\overline{D}_{l_{L}} \Phi)_{1''} H \tau_{R_{1'}} + \frac{y_{1}}{\Lambda} (\overline{D}_{l_{L}} \Phi)_{1'} \\ & \qquad \qquad \tilde{H} \nu_{e_{R_{1''}}} + \frac{y_{2}}{\Lambda^{2}} (\overline{D}_{l_{L}} \Phi)_{1} \, \tilde{H} \nu_{\mu_{R_{1}}} \chi + \frac{y_{3}}{\Lambda^{2}} (\overline{D}_{l_{L}} \Phi)_{1''} \, \tilde{H} \nu_{\tau_{R_{1'}}} \xi + \frac{y_{a}}{2} (\overline{\nu^{c}}_{\mu_{R}} \nu_{\mu_{R}})_{1} \, \xi_{1} \\ & \qquad \qquad + \frac{y_{b}}{2} \left[(\overline{\nu^{c}}_{e_{R}} \nu_{\tau_{R}}) + (\overline{\nu^{c}}_{\tau_{R}} \nu_{e_{R}})_{1}_{1} \, \xi_{1} + \frac{y_{c}}{2\Lambda} (\overline{\nu^{c}}_{e_{R}} \nu_{e_{R}})_{1} \, \eta_{1''} \, \varsigma + \frac{y_{t}}{\Lambda} (\overline{D}_{l_{L}} \, D_{l_{L}}^{c})_{3_{S}} \, \psi \Delta_{1} \\ & \qquad \qquad + \frac{y_{s}}{\Lambda} (\overline{D}_{l_{L}} \, D_{l_{L}}^{c})_{3_{S}} \, \kappa \, \Delta_{2} \, + h.c. \end{split}$$

The product rules under A_4 are given in appendix A. The details for the A_4 vacuum alignment have been discussed in Appendix B. The additional symmetry groups Z_{10} and Z_2 are considered to restrict the next to leading order corrections. The details of Z_{10} group can be found in Refs. [38–40]. Here Λ is the cut-off scale of the theory. The charged lepton mass matrix under the choice of vacuum alignment $\langle \Phi \rangle_{\circ} = v_{\Phi} (1, 0, 0)^T$ and $\langle H \rangle_{\circ} = v_H$ can be derived from Eq.(7) as shown below,

$$M_{l} = \begin{bmatrix} \frac{y_{e}}{\Lambda} v_{\Phi} v_{H} & 0 & 0\\ 0 & \frac{y_{\mu}}{\Lambda} v_{\Phi} v_{H} & 0\\ 0 & 0 & \frac{y_{\tau}}{\Lambda} v_{\Phi} v_{H} \end{bmatrix}, \tag{8}$$

We choose the vacuum alignments $\langle \eta \rangle_{\circ} = v_{\eta}$, $\langle \chi \rangle_{\circ} = v_{\chi}$, $\langle \xi \rangle_{\circ} = v_{\xi}$, $\langle \varsigma \rangle_{\circ} = v_{\zeta}$ to derive the Dirac neutrino mass matrix (M_D) and right handed neutrino mass matrix (M_R) as shown in the following,

$$M_D = \begin{bmatrix} 0 & \frac{y_2}{\Lambda^2} v_{\Phi} v_H v_{\chi} & 0\\ \frac{y_1}{\Lambda} v_{\Phi} v_H & 0 & 0\\ 0 & 0 & \frac{y_3}{\Lambda^2} v_{\Phi} v_H v_{\xi} \end{bmatrix}, \tag{9}$$

$$M_R = \begin{bmatrix} \frac{y_c}{2\Lambda} v_{\eta} v_{\varsigma} & 0 & \frac{y_b}{2} v_{\xi} \\ 0 & \frac{y_a}{2} v_{\xi} & 0 \\ \frac{y_b}{2} v_{\xi} & 0 & 0 \end{bmatrix}, \tag{10}$$

The two matrices $M_{T_2}^1$ and $M_{T_2}^2$ from two type-II seesaw mechanisms can be derived from Eq.(7) by considering the vacuum alignment $\langle \Delta_1 \rangle_{\circ} = v_{\Delta_1}$, $\langle \Delta_2 \rangle_{\circ} = v_{\Delta_2}$, $\langle \psi \rangle_{\circ} = v_{\psi}(0, 1, -1)^T$ and $\langle \kappa \rangle_{\circ} = v_{\kappa}(0, 1, -1)^T$ as shown in the following,

$$M_{T_2}^1 = \begin{bmatrix} 0 & \frac{1}{3\Lambda} y_t \, v_{\Delta_1} \, v_{\psi} & -\frac{1}{3\Lambda} y_t \, v_{\Delta_1} \, v_{\psi} \\ \frac{1}{3\Lambda} y_t \, v_{\Delta_1} \, v_{\psi} & \frac{2}{3\Lambda} y_t \, v_{\Delta_1} \, v_{\psi} & 0 \\ -\frac{1}{3\Lambda} y_t \, v_{\Delta_1} \, v_{\psi} & 0 & -\frac{2}{3\Lambda} y_t \, v_{\Delta_1} \, v_{\psi} \end{bmatrix}, \tag{11}$$

$$M_{T_2}^2 = \begin{bmatrix} \frac{2}{3} \frac{1}{\Lambda} y_s v_\kappa v_{\Delta_2} & -\frac{1}{3\Lambda} y_s v_\kappa v_{\Delta_2} & 0\\ -\frac{1}{3\Lambda} y_s v_\kappa v_{\Delta_2} & 0 & -\frac{1}{3\Lambda} y_s v_\kappa v_{\Delta_2}\\ 0 & -\frac{1}{3\Lambda} y_s v_\kappa v_{\Delta_2} & \frac{2}{3\Lambda} y_s v_\kappa v_{\Delta_2} \end{bmatrix}.$$
 (12)

The effective neutrino mass matrix M_{ν} is constructed after taking the contributions from type-I $(-M_D M_R^{-1} M_D^T)$ and two type-II seesaw mechanisms as shown below,

$$M_{\nu} = \begin{bmatrix} A & B & F \\ B & -2F & G \\ F & G & J \end{bmatrix}, \tag{13}$$

where, A, B, F, G, J are purely complex parameters. The said texture parameters can be expressed in terms of the model parameters as shown in the following,

$$A = \frac{2 y_s v_{\Delta_2} v_{\kappa}}{3 \Lambda} - \frac{2 y_2^2 v_H^2 v_{\chi}^2 v_{\phi}^2}{\Lambda^4 y_a v_{\xi}}, \tag{14}$$

$$B = \frac{y_t v_{\Delta_1} v_{\psi}}{3 \Lambda} - \frac{y_s v_{\Delta_2} v_{\kappa}}{3 \Lambda},$$

$$F = -\frac{y_t v_{\Delta_1} v_{\psi}}{3 \Lambda},$$
(15)

$$F = -\frac{y_t \, v_{\Delta_1} \, v_{\psi}}{3 \, \Lambda}, \tag{16}$$

$$G = -\frac{2y_1 y_3 v_H^2 v_\phi^2}{\Lambda^3 y_b} - \frac{y_s v_{\Delta_2} v_\kappa}{3\Lambda}, \tag{17}$$

$$J = \frac{2 y_3^2 y_c v_H^2 v_\zeta v_\eta v_\phi^2}{\Lambda^5 y_h^2} + \frac{2 y_s v_{\Delta_2} v_\kappa}{3 \Lambda} - \frac{2 y_t v_{\Delta_1} v_\psi}{3 \Lambda}.$$
 (18)

It is important to mention that it is very difficult to predict the Yukawa couplings individually. However, the present work predicts the combinations of the model parameters $Y_1 = \frac{y_t v_{\Delta_1} v_{\psi}}{\Lambda}$, $Y_2 = \frac{y_s v_{\Delta_2} v_{\kappa}}{\Lambda}$, $Y_3 = \frac{\Lambda^4 y_a v_{\xi}}{2y_2^2 v_H^2 v_{\chi}^2 v_{\phi}^2}$, $Y_4 = \frac{\Lambda^3 y_b}{2y_1 y_3 v_H^2 v_{\phi}^2}$ and $Y_5 = \frac{y_c \Lambda v_\zeta v_\eta}{y_1^2 v_H^2 v_\phi^2}$. These combinations can be expressed in terms of the texture parameters A, B, F, G and J as shown below,

$$Y_1 = -3F,$$
 (19)

$$Y_2 = -3(B+F), (20)$$

$$Y_3 = -\frac{1}{A + 2B + 2F},\tag{21}$$

$$Y_4 = \frac{2}{B + F - G'},\tag{22}$$

$$Y_5 = \frac{2(2B+F)}{(B+F-G)^2}. (23)$$




FIG. 4. Shows the correlation plots between (a) $|Y_1|$ and $|Y_2|$. (b) $|Y_4|$ vs $|Y_3|$ and $|Y_5|$. (c) Arg $[Y_1]$ vs Arg $[Y_{2,3,4,5}]$.

Model Parameter	Minimum Value	Maximum Value				
$ Y_1 /\mathrm{eV}$	0.465	0.054				
$ Y_2 /\mathrm{eV}$	0,032	0.039				
$ Y_3 /{\rm eV}^{-1}$	27.70	41.59				
$ Y_4 /{\rm eV}^{-1}$	50.97	64.82				
$ Y_5 /{\rm eV}^{-1}$	23.47	26.86				
$Arg[Y_1]/^{\circ}$	-177.99	13.85				
$Arg[Y_2]/^{\circ}$	-31.73	150.80				
$Arg[Y_3]/^{\circ}$	-170.04	26.22				
$Arg[Y_4]/^{\circ}$	-171.01	19.01				
$Arg[Y_5]/^{\circ}$	-66.59	151.65				

TABLE III. Allowed values of the model parameters compatible with 3σ range of neutrino oscillation data[15]

•

For graphical representation, we generate correlation plots to visualize how the absolute values and arguments of the combinations of the model parameters vary. It is evident from Figs. 3(a)-3(c) that the studied parameters are constrained to certain bounds. The maximum and minimum values of the model parameters are listed in Table III corresponding to the 3σ range of neutrino oscillation data [15].

Deriving the proposed model-independent neutrino mass matrix texture as shown in Eq. (4) from the symmetry framework has some inherent challenges. Moreover, it's crucial to ensure the independence of parameters A, B, F, G, and J from each other. This is achieved by including contributions from both type-I and type-II contributions, leading to an increase in the field content. This is a necessary step to achieve the high productivity of the framework. To obtain the texture in its exact form, we introduce the field ς , to cut short certain non-leading terms in the Yukawa Lagrangian. Similarly, we need to cut short specific elements within the right-handed Majorana neutrino mass matrix which requires two additional scalar fields ζ and χ . If Δ_1 and Δ_2 were chosen as A_4 triplets, the field content could have been reduced. However, in that case, certain components within Δ_i will not get vevs and consequently, there remains the finite possibility that these components will contribute towards the neutrino mass via radiative correction. To avoid this, we need to treat Δ_1 and Δ_2 as singlets under A_4 and introduce the A_4 scalar triplet fields ψ and κ to restore the invariance of the Lagrangian.

IV. SUMMARY AND DISCUSSION

In the present work, we have proposed a new mixing scheme aligning with the philosophy of historical fixed/constant mixing schemes such as bimaximal, tri-bimaximal, golden ratio, and hexagonal mixing schemes. Unlike its predecessors, in the proposed mixing scheme, the Dirac CP phase δ is fixed at $3\pi/4$. The other mixing angles are fixed at $s_{12} = (2\sqrt{2})/5$, $s_{13} = 1/(3\sqrt{5})$ and $s_{23} = 3/4$. The proposed mixing scheme is special in the sense that it can be linked to a neutrino mass matrix texture bearing a suitable correlation $(m_{22} + 2 m_{13} = 0)$ among its elements. The texture predicts three non-vanishing neutrino mass eigenvalues which follow normal ordering ensuring that $\Sigma m_i \leq 0.12$ eV. In addition, the texture constraints of the Majorana phase α with minimum and maximum values -13.96° and -3.18° respectively. On the other hand, the phase β is constrained with minimum and maximum values -29.37° and -0.75° respectively. The effective mass parameter $m_{\beta\beta}$ appearing in the neutrinoless double beta decay is also predicted in the range with the minimum value 17.28 MeV and maximum value 20.22 MeV respectively which falls within the sensitivity of future experiments such as LEGEND-1000 [41]. We demonstrate that the texture compatible with the proposed mixing scheme can be derived from the interplay of general type-I+II seesaw mechanisms in the framework of $A_4 \times Z_{10} \times Z_2$ group. As a future prospect, we aim to study the baryon asymmetry of the Universe and lepton flavour violating decays in the context of the proposed texture. The detailed analysis in this regard is left for a future work.

ACKNOWLEDGEMENT

The research work of PC is supported by Innovation in Science Pursuit for Inspired Research (INSPIRE), Department of Science and Technology, Government of India, New Delhi, vide grant No. IF190651. MD acknowledges the financial support from the Council of Scientific and Industrial Research (CSIR), Government of India, New Delhi, through a NET Junior Research Fellowship vide grant No. 09/0059(15346)/2022-EMR-I. The work of BK has been supported in part by the Polish National Science Center (NCN) under grant 2020/37/B/ST2/02371, the Freedom of Research, Mobility of Science, and the Research Excellence Initiative of the University of Silesia in Katowice.

Appendix A: Product Rules of A₄

The discrete group A_4 is a subgroup of SU(3) which has 12 elements. The group has four irreducible representations, out of which three are one-dimensional and one is three-dimensional. In T-basis, the product of two triplets gives,

$$3 \times 3 = 1 + 1' + 1'' + 3_S + 3_A,\tag{A1}$$

where,

$$1 = (a_1b_1 + a_2b_3 + a_3b_2) \tag{A2}$$

$$1' = (a_3b_3 + a_1b_2 + a_2b_1) \tag{A3}$$

$$1'' = (a_2b_2 + a_1b_3 + a_3b_1) (A4)$$

$$(3 \times 3)_S = \frac{1}{3} \begin{bmatrix} 2a_1b_1 - a_2b_3 - a_3b_2 \\ 2a_3b_3 - a_1b_2 - a_2b_1 \\ 2a_2b_2 - a_1b_3 - a_3b_1 \end{bmatrix}, \tag{A5}$$

$$(3 \times 3)_{S} = \frac{1}{3} \begin{bmatrix} 2a_{1}b_{1} - a_{2}b_{3} - a_{3}b_{2} \\ 2a_{3}b_{3} - a_{1}b_{2} - a_{2}b_{1} \\ 2a_{2}b_{2} - a_{1}b_{3} - a_{3}b_{1} \end{bmatrix},$$

$$(3 \times 3)_{A} = \frac{1}{2} \begin{bmatrix} a_{2}b_{3} - a_{3}b_{2} \\ a_{1}b_{2} - a_{2}b_{1} \\ a_{3}b_{1} - a_{1}b_{3} \end{bmatrix}.$$

$$(A4)$$

$$(A5)$$

The trivial singlet can be obtained from the following singlet product rules,

$$1 \times 1 = 1, \ 1' \times 1'' = 1, \ 1'' \times 1' = 1.$$
 (A7)

Appendix B: The Scalar Potential

The $A_4 \times Z_{10} \times Z_2$ invariant scalar potential can be constructed in the following manner,

$$V = V(\Phi) + V(H) + V(\Delta_1) + V(\Delta_2) + V(\eta) + V(\xi) + V(\chi) + V(\psi) + V(\kappa)$$

$$+V(\zeta) + V(\Phi H) + V(\Phi \Delta) + V(\Phi \Delta_2) + V(\Phi \eta) + V(\Phi \xi) + V(\Phi \chi) + V(\Phi \psi)$$

$$+V(\Phi \kappa) + V(\Phi \zeta) + V(H\Delta) + V(H\Delta_2) + V(H\eta) + V(H\xi) + V(H\chi) + V(H\psi) + V(H\kappa) + V(H\zeta) + V(\Delta_1 \Delta_2) + V(\Delta_1 \eta) + V(\Delta_1 \xi) + V(\Delta_1 \chi) + V(\Delta_1 \psi)$$

$$+V(\Delta_1 \kappa) + V(\Delta_1 \zeta) + V(\Delta_2 \eta) + V(\Delta_2 \xi) + V(\Delta_2 \chi) + V(\Delta_2 \psi) + V(\Delta_2 \kappa) + V(\Delta_2 \zeta)$$

$$+V(\eta \xi) + V(\eta \chi) + V(\eta \psi) + V(\eta \kappa) + V(\eta \zeta) + V(\xi \chi) + V(\xi \psi) + V(\xi \kappa) + V(\xi \zeta) + V(\chi \psi) + V(\chi \kappa) + V(\chi \zeta) + V(\psi \kappa) + V(\psi \zeta) + V(\kappa \zeta),$$
(B1)

where,

$$V(\Phi) = \mu_{\Phi}^{2}(\Phi^{\dagger}\Phi) + \lambda_{1}^{\Phi}(\Phi^{\dagger}\Phi)(\Phi^{\dagger}\Phi) + \lambda_{2}^{\Phi}(\Phi^{\dagger}\Phi)_{1'}(\Phi^{\dagger}\Phi)_{1''} + \lambda_{3}^{\Phi}(\Phi^{\dagger}\Phi)_{3_{s}}(\Phi^{\dagger}\Phi)_{3_{s}} + \lambda_{4}^{\Phi}(\Phi^{\dagger}\Phi)_{3_{s}}$$
(B2)
$$V(H) = \mu_{H}^{2}(H^{\dagger}H) + \lambda^{H}(H^{\dagger}H)(H^{\dagger}H),$$
(B3)
$$V(\Delta_{1}) = \mu_{\Delta_{1}}^{2}Tr(\Delta_{1}^{\dagger}\Delta_{1}) + \lambda_{1}^{\Delta}Tr(\Delta_{1}^{\dagger}\Delta_{1})Tr(\Delta_{1}^{\dagger}\Delta_{1}),$$
(B4)
$$V(\Delta_{2}) = \mu_{\Delta_{2}}^{2}Tr(\Delta_{2}^{\dagger}\Delta_{2}) + \lambda^{\Delta_{2}}Tr(\Delta_{2}^{\dagger}\Delta_{2})Tr(\Delta_{2}^{\dagger}\Delta_{2}),$$
(B5)
$$V(\eta) = \mu_{\eta}^{2}(\eta^{\dagger}\eta) + \lambda^{\eta}(\eta^{\dagger}\eta)(\eta^{\dagger}\eta),$$
(B6)
$$V(\xi) = \mu_{\xi}^{2}(\xi^{\dagger}\xi) + \lambda^{\xi}(\xi^{\dagger}\xi)(\xi^{\dagger}\xi),$$
(B7)
$$V(\chi) = \mu_{\chi}^{2}(\chi^{\dagger}\chi) + \lambda^{\chi}(\chi^{\dagger}\chi)(\chi^{\dagger}\chi),$$
(B8)
$$V(\zeta) = \mu_{\psi}^{2}(\chi^{\dagger}\psi) + \lambda_{1}^{\psi}(\psi^{\dagger}\psi)(\psi^{\dagger}\psi) + \lambda_{2}^{\psi}(\psi^{\dagger}\psi)_{1'}(\psi^{\dagger}\psi)_{1''} + \lambda_{3}^{\psi}(\psi^{\dagger}\psi)_{3_{s}}(\psi^{\dagger}\psi)_{3_{s}} + \lambda_{4}^{\psi}(\psi^{\dagger}\psi)_{3_{s}}$$
(B9)
$$V(\psi) = \mu_{\psi}^{2}(\psi^{\dagger}\psi) + \lambda_{1}^{\psi}(\psi^{\dagger}\psi)(\psi^{\dagger}\psi) + \lambda_{2}^{\psi}(\psi^{\dagger}\psi)_{1'}(\psi^{\dagger}\psi)_{1''} + \lambda_{3}^{\psi}(\psi^{\dagger}\psi)_{3_{s}}(\psi^{\dagger}\psi)_{3_{s}} + \lambda_{4}^{\psi}(\psi^{\dagger}\psi)_{3_{s}}$$
(B10)
$$V(\kappa) = \mu_{\kappa}^{2}(\kappa^{\dagger}\kappa) + \lambda_{1}^{\kappa}(\kappa^{\dagger}\kappa)(\kappa^{\dagger}\kappa) + \lambda_{2}^{\kappa}(\kappa^{\dagger}\kappa)_{1'}(\kappa^{\dagger}\kappa)_{1''} + \lambda_{3}^{\kappa}(\kappa^{\dagger}\kappa)_{3_{s}}(\kappa^{\dagger}\kappa)_{3_{s}} + \lambda_{4}^{\psi}(\kappa^{\dagger}\kappa)_{3_{s}}$$
(B11)
$$V(\Phi H) = \lambda_{1}^{\Phi}(\Phi^{\dagger}\Phi)(H^{\dagger}H),$$
(B12)
$$V(\Phi \Delta_{1}) = \lambda_{1}^{\Phi\Delta_{1}}(\Phi^{\dagger}\Phi)Tr(\Delta_{1}^{\dagger}\Delta_{1}),$$
(B13)
$$V(\Phi \Delta_{2}) = \lambda_{1}^{\Phi\Delta_{2}}(\Phi^{\dagger}\Phi)Tr(\Delta_{1}^{\dagger}\Delta_{2}),$$
(B16)
$$V(\Phi \psi) = \lambda_{2}^{\Phi}(\Phi^{\dagger}\Phi)(\psi^{\dagger}\psi),$$
(B15)
$$V(\Phi \psi) = \lambda_{2}^{\Phi}(\Phi^{\dagger}\Phi)(\psi^{\dagger}\psi),$$
(B16)

 $V(\Phi\zeta) = \lambda_1^{\Phi\zeta}(\Phi^{\dagger}\Phi)(\zeta^{\dagger}\zeta),$ (B18) $V(\Phi\psi) \; = \; \lambda_1^{\Phi\psi}(\Phi^{\dagger}\Phi)(\psi^{\dagger}\psi) + \lambda_2^{\Phi\psi}[(\Phi^{\dagger}\Phi)_{1'}(\psi^{\dagger}\psi)_{1''} + (\Phi^{\dagger}\Phi)_{1''}(\psi^{\dagger}\psi)_{1'}] + \lambda_3^{\Phi\psi}[(\Phi^{\dagger}\Phi)_{1''}(\psi^{\dagger}\psi)_{1''}] + \lambda_3^{\Phi\psi}$ $(\Phi^{\dagger}\Phi)_{3_{-}}(\psi^{\dagger}\psi)_{3_{-}} + \lambda_{4}^{\Phi\psi}[(\Phi^{\dagger}\Phi)_{3_{-}}(\psi^{\dagger}\psi)_{3_{-}} + (\Phi^{\dagger}\Phi)_{3_{-}}(\psi^{\dagger}\psi)_{3_{-}}] + \lambda_{5}^{\Phi\psi}$

$$(\Phi^{\dagger}\Phi)_{3_s}(\psi^{\dagger}\psi)_{3_s} + \lambda_4^{\dagger} [(\Phi^{\dagger}\Phi)_{3_s}(\psi^{\dagger}\psi)_{3_a} + (\Phi^{\dagger}\Phi)_{3_a}(\psi^{\dagger}\psi)_{3_s}] + \lambda_5^{\dagger}$$

$$(\Phi^{\dagger}\Phi)_{3_a}(\psi^{\dagger}\psi)_{3_a}, \tag{B19}$$

$$V(\Phi\kappa) = \lambda_1^{\Phi\kappa}(\Phi^{\dagger}\Phi)(\kappa^{\dagger}\kappa) + \lambda_2^{\Phi\kappa}[(\Phi^{\dagger}\Phi)_{1'}(\kappa^{\dagger}\kappa)_{1''} + (\Phi^{\dagger}\Phi)_{1''}(\kappa^{\dagger}\kappa)_{1'}] + \lambda_3^{\Phi\kappa}(\Phi^{\dagger}\Phi)_{3_s} (\kappa^{\dagger}\kappa)_{3_s} + \lambda_4^{\Phi\kappa}[(\Phi^{\dagger}\Phi)_{3_s}(\kappa^{\dagger}\kappa)_{3_a} + (\Phi^{\dagger}\Phi)_{3_a}(\kappa^{\dagger}\kappa)_{3_s}] + \lambda_5^{\Phi\kappa}(\Phi^{\dagger}\Phi)_{3_a}(\kappa^{\dagger}\kappa)_{3_a},$$
(B20)

$$V(H\Delta_1) = \lambda_1^{H\Delta_1}(H^{\dagger}H) Tr(\Delta_1^{\dagger}\Delta_1) + \frac{1}{\Lambda}(H^T i\sigma_2 \Delta_1^{\dagger} H \Phi \psi), \tag{B21}$$

$$V(H\Delta_2) = \lambda_1^{H\Delta_2}(H^{\dagger}H) Tr(\Delta_2^{\dagger}\Delta_2) + \frac{1}{\Lambda} (H^T i\sigma_2 \Delta_2^{\dagger} H \Phi \psi), \tag{B22}$$

(B23)

(B17)

(B57)

```
V(H\eta) = \lambda_1^{H\eta} (H^{\dagger}H)(\eta^{\dagger}\eta),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           (B24)
            V(H\xi) = \lambda_1^{H\xi}(H^{\dagger}H)(\xi^{\dagger}\xi),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          (B25)
           V(H\chi) = \lambda_1^{H\chi}(H^{\dagger}H)(\chi^{\dagger}\chi),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          (B26)
           V(H\psi) = \lambda_1^{H\psi}(H^{\dagger}H)(\psi^{\dagger}\psi),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          (B27)
            V(H\kappa) = \lambda_1^{H\kappa}(H^{\dagger}H)(\kappa^{\dagger}\kappa),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          (B28)
             V(H\zeta) = \lambda_1^{H\zeta}(H^{\dagger}H)(\zeta^{\dagger}\zeta),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          (B29)
V(\Delta_1 \Delta_2) = \lambda_1^{\Delta_1 \Delta_2} (\Delta_1^{\dagger} \Delta_1) Tr(\Delta_2^{\dagger} \Delta_2),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          (B30)
         V(\Delta_1 \eta) = \lambda_1^{\Delta_1 \eta} Tr(\Delta_1^{\dagger} \Delta_1)(\eta^{\dagger} \eta),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          (B31)
         V(\Delta_1 \xi) = \lambda_1^{\Delta_1 \xi} Tr(\Delta_1^{\dagger} \Delta_1)(\xi^{\dagger} \xi),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          (B32)
       V(\Delta_1 \chi) = \lambda_1^{\Delta_1 \chi} Tr(\Delta_1^{\dagger} \Delta_1)(\chi^{\dagger} \chi),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           (B33)
       V(\Delta_1 \psi) = \lambda_1^{\Delta_1 \psi} Tr(\Delta_1^{\dagger} \Delta_1)(\psi^{\dagger} \psi),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           (B34)
       V(\Delta_1 \kappa) = \lambda_1^{\Delta_1 \kappa} Tr(\Delta_1^{\dagger} \Delta_1)(\kappa^{\dagger} \kappa),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           (B35)
        V(\Delta_1 \zeta) = \lambda_1^{\Delta_1 \zeta} Tr(\Delta_1^{\dagger} \Delta_1)(\zeta^{\dagger} \zeta),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           (B36)
       V(\Delta_2 \eta) = \lambda_1^{\Delta_2 \eta} Tr(\Delta_2^{\dagger} \Delta_2)(\eta^{\dagger} \eta),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          (B37)
        V(\Delta_2 \xi) = \lambda_1^{\Delta_2 \xi} Tr(\Delta_2^{\dagger} \Delta_2)(\xi^{\dagger} \xi),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          (B38)
       V(\Delta_2 \chi) = \lambda_1^{\Delta_2 \chi} Tr(\Delta_2^{\dagger} \Delta_2)(\chi^{\dagger} \chi),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          (B39)
      V(\Delta_2 \psi) = \lambda_1^{\Delta_2 \psi} Tr(\Delta_2^{\dagger} \Delta_2)(\psi^{\dagger} \psi),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          (B40)
       V(\Delta_2 \kappa) = \lambda_1^{\Delta_2 \kappa} Tr(\Delta_2^{\dagger} \Delta_2)(\kappa^{\dagger} \kappa),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          (B41)
        V(\Delta_2 \zeta) = \lambda_1^{\Delta_2 \zeta} Tr(\Delta_2^{\dagger} \Delta_2)(\zeta^{\dagger} \zeta),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          (B42)
                  V(\eta \xi) = \lambda_1^{\eta \xi} (\eta^{\dagger} \eta) (\xi^{\dagger} \xi),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          (B43)
                V(\eta \chi) = \lambda_1^{\eta \chi} (\eta^{\dagger} \eta) (\chi^{\dagger} \chi),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          (B44)
                V(\eta\psi) = \lambda_1^{\eta\psi}(\eta^{\dagger}\eta)(\psi^{\dagger}\psi),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          (B45)
                  V(\eta \kappa) = \lambda_1^{\eta \kappa} (\eta^{\dagger} \eta) (\kappa^{\dagger} \kappa),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          (B46)
                  V(\eta\zeta) = \lambda_1^{\eta\zeta}(\eta^{\dagger}\eta)(\zeta^{\dagger}\zeta),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          (B47)
                  V(\xi \chi) = \lambda_1^{\xi \chi}(\xi^{\dagger} \xi)(\chi^{\dagger} \chi),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           (B48)
                 V(\xi\psi) = \lambda_1^{\xi\psi}(\xi^{\dagger}\xi)(\psi^{\dagger}\psi),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           (B49)
                  V(\xi \kappa) = \lambda_1^{\xi \kappa} (\xi^{\dagger} \xi) (\kappa^{\dagger} \kappa),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           (B50)
                  V(\xi\zeta) = \lambda_1^{\xi\zeta}(\xi^{\dagger}\xi)(\zeta^{\dagger}\zeta),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          (B51)
              V(\chi\psi) = \lambda_1^{\chi\psi}(\chi^{\dagger}\chi)(\psi^{\dagger}\psi),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          (B52)
                 V(\chi\kappa) = \lambda_1^{\chi\kappa}(\chi^{\dagger}\chi)(\kappa^{\dagger}\kappa),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          (B53)
                  V(\chi\zeta) = \lambda_1^{\chi\zeta}(\chi^{\dagger}\chi)(\zeta^{\dagger}\zeta),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          (B54)
               V(\psi\kappa) = \lambda_1^{\psi\kappa}(\psi^{\dagger}\psi)(\kappa^{\dagger}\kappa) + \lambda_2^{\psi\kappa}[(\psi^{\dagger}\psi)_{1'}(\kappa^{\dagger}\kappa)_{1''} + (\psi^{\dagger}\psi)_{1''}(\kappa^{\dagger}\kappa)_{1'}] + \lambda_3^{\psi\kappa}[(\psi^{\dagger}\psi)_{1''}(\kappa^{\dagger}\kappa)_{1''}] + \lambda_3^{\psi\kappa}[(\psi^{\dagger}\psi)_{1''}(\kappa^{\dagger}\psi)_{1''}] + \lambda_3^{\psi\kappa}[(\psi^
                                                                           (\psi^{\dagger}\psi)_{3_{\circ}}(\kappa^{\dagger}\kappa)_{3_{\circ}} + \lambda_{4}^{\psi\kappa}[(\psi^{\dagger}\psi)_{3_{\circ}}(\kappa^{\dagger}\kappa)_{3_{\circ}} + (\psi^{\dagger}\psi)_{3_{\circ}}(\kappa^{\dagger}\kappa)_{3_{\circ}}] + \lambda_{5}^{\psi\kappa}
                                                                           (\psi^{\dagger}\psi)_{3a}(\kappa^{\dagger}\kappa)_{3a}
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           (B55)
                V(\psi\zeta) = \lambda_1^{\psi\zeta}(\psi^{\dagger}\psi)(\zeta^{\dagger}\zeta),
                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          (B56)
```

The following equations hold good for the chosen vacuum expectation values (VEVs) $\langle \Phi \rangle_{\circ} = v_{\Phi} (1,0,0)^{T}$, $\langle H \rangle_{\circ} = v_{H}$, $\langle \eta \rangle_{\circ} = v_{\eta}$, $\langle \chi \rangle_{\circ} = v_{\chi}$, $\langle \xi \rangle_{\circ} = v_{\xi}$, $\langle \varsigma \rangle_{\circ} = v_{\varsigma}$, $\langle \Delta_{1} \rangle_{\circ} = v_{\Delta_{1}}$, $\langle \Delta_{2} \rangle_{\circ} = v_{\Delta_{2}}$, $\langle \psi \rangle_{\circ} = v_{\psi} (0,1,-1)^{T}$ and $\langle \kappa \rangle_{\circ} = v_{\kappa} (0,1,-1)^{T}$, based on the minimization conditions of the scalar potential:

 $V(\kappa\zeta) = \lambda_1^{\kappa\zeta}(\kappa^{\dagger}\kappa)(\zeta^{\dagger}\zeta),$

(B70) (B71)

$$\frac{dV}{d\Phi_1} = \frac{1}{9} v_{\Phi} \left(-\frac{24}{\Lambda} \mu_2 v_{\Delta}, v_H^2 v_\chi^2 \Lambda_1^{\Phi X} + 9 v_{\Delta_2}^2 \Lambda_1^{\Phi X_2} + 9 v_{\eta}^2 \Lambda_1^{\Phi H} + 9 v_{\kappa}^2 \Lambda_1^{\Phi H} + 4 v_{\kappa}^2 \lambda_3^{\Phi \kappa} \right. \\ + 9 \mu_{\Phi}^2 - 18 v_{\phi}^2 \Lambda_1^{\Phi W} + 4 v_{\phi}^2 \lambda_3^{\Phi W} + 2 v_{\phi}^2 \left(9 \Lambda_1^{\Phi} + 4 \lambda_3^{\Phi} \right) + 9 \Lambda_1^{\Phi V} v_{\phi}^2 + 9 \Lambda_1^{\phi V} v_{\phi}^2 \right. \\ + 9 v_{\Delta}^2 \Lambda_1^{\Phi \Delta_1} \right) = 0, \qquad (B58)$$

$$\frac{dV}{d\Phi_2} = \frac{1}{9} v_{\Phi} \left(\frac{12}{\Lambda} v_H^2 v_{\Delta_1} \left(\mu_2 v_{\kappa} - \mu_1 v_{\psi} \right) + 18 v_{\kappa}^2 \lambda_2^{\Phi \kappa} + 2 v_{\kappa}^2 \lambda_3^{\Phi \kappa} - 3 v_{\kappa}^2 \lambda_4^{\Phi \kappa} + 9 v_{\phi}^2 \lambda_2^{\Phi \psi} \right) \\ - 2 v_{\phi}^2 \lambda_3^{\Phi \Psi} + 3 v_{\phi}^2 \lambda_1^{\Phi \Psi} \right) = 0, \qquad (B59)$$

$$\frac{dV}{d\Phi_3} = \frac{1}{9} v_{\Phi} (v_{\kappa}^2 (9 \lambda_2^{\Phi \kappa} - 2 \lambda_3^{\Phi \kappa} - 3 \lambda_4^{\Phi \kappa}) + v_{\phi}^2 \left(\frac{12}{\Lambda^2} \mu_1 v_H^2 v_{\Delta_1} + 9 \lambda_2^{\Phi \psi} - 2 \lambda_3^{\Phi \psi} \right) \\ - 3 \lambda_4^{\Phi \psi} \right) = 0, \qquad (B60)$$

$$\frac{dV}{d\psi_1} = -\frac{4}{3\Lambda} \mu_1 v_H^2 v_{\Delta_1} v_{\phi}^2 + \frac{1}{9} v_{\psi} (v_{\kappa}^2 (9 \lambda_2^{\Phi \psi} + 4 \lambda_3^{\Phi \psi}) - 6 v_{\phi}^2 \lambda_4^{\Psi}) = 0, \qquad (B61)$$

$$\frac{dV}{d\psi_2} = \frac{1}{9} v_{\Psi} \left(-9 v_H^2 \Lambda_1^{H \Psi} - 9 \mu_{\phi}^2 - 9 \Lambda_1^{4 \psi} v_{\Delta_1}^2 - 9 v_{\Delta_2}^2 \lambda_1^{4 \psi} v_{\phi} - 9 \lambda_1^{4 \psi} v_{\phi}^2 - 9 \lambda_1$$

 $+\lambda_1^{\eta\chi}v_{\chi}^2 - 2\lambda_1^{\eta\psi}v_{\psi}^2 + \lambda_1^{\Phi\eta}v_{\Phi}^2) = 0,$

$$\frac{dV}{d\xi} = v_{\xi}(v_{H}^{2}\lambda_{1}^{H\xi} + \mu_{\xi}^{2} + \lambda_{1}^{\Delta_{1}\xi}v_{\Delta_{1}}^{2} + v_{\Delta_{2}}^{2}\lambda_{1}^{\Delta_{2}\xi} + \lambda_{1}^{\xi\zeta}v_{\zeta}^{2} + \lambda_{1}^{\eta\xi}v_{\eta}^{2} + \lambda_{1}^{\xi\kappa}v_{\kappa}^{2} + 2\lambda^{\xi}v_{\xi}^{2} + \lambda_{1}^{\xi\kappa}v_{\chi}^{2} - 2\lambda_{1}^{\xi\psi}v_{\psi}^{2} + \lambda_{1}^{\Phi\xi}v_{\Phi}^{2}) = 0,$$
(B72)

$$\frac{dV}{d\chi} = v_{\chi}(v_{H}^{2}\lambda_{1}^{H\chi} + \mu_{\chi}^{2} + \lambda_{1}^{\Delta_{1}\chi}v_{\Delta_{1}}^{2} + v_{\Delta_{2}}^{2}\lambda_{1}^{\Delta_{2}\chi} + \lambda_{1}^{\chi\zeta}v_{\zeta}^{2} + \lambda_{1}^{\eta\chi}v_{\eta}^{2} + \lambda_{1}^{\chi\kappa}v_{\kappa}^{2} + 2\lambda^{\chi}v_{\chi}^{2} + \lambda_{1}^{\xi\chi}v_{\zeta}^{2} + \lambda_{1}^{\eta\chi}v_{\eta}^{2} + \lambda_{1}^{\chi\kappa}v_{\kappa}^{2} + 2\lambda^{\chi}v_{\chi}^{2} + \lambda_{1}^{\chi\kappa}v_{\zeta}^{2} + \lambda_{1}^{\eta\chi}v_{\eta}^{2} + \lambda_{1}^{\chi\kappa}v_{\zeta}^{2} + \lambda_{$$

$$\frac{dV}{d\zeta} = v_{\zeta}(\mu_{\zeta}^{2} + v_{H}^{2}\lambda_{1}^{H\zeta} + \lambda_{1}^{\Delta_{1}\zeta}v_{\Delta_{1}}^{2} + v_{\Delta_{2}}^{2}\lambda_{1}^{\Delta_{2}\zeta} + 2\lambda^{\zeta}v_{\zeta}^{2} + \lambda_{1}^{\eta\zeta}v_{\eta}^{2} + \lambda_{1}^{\kappa\zeta}v_{\kappa}^{2} + \lambda_{1}^{\xi\zeta}v_{\xi}^{2} + \lambda_{1}^{\eta\zeta}v_{\chi}^{2} - 2\lambda_{1}^{\psi\zeta}v_{\psi}^{2}) = 0.$$
(B74)

- [1] B. Pontecorvo, "Mesonium and anti-mesonium," Sov. Phys. JETP 6 (1957) 429.
- [2] Z. Maki, M. Nakagawa, and S. Sakata, "Remarks on the unified model of elementary particles," Prog. Theor. Phys. 28 (1962) 870–880.
- [3] S. L. Glashow, "Partial Symmetries of Weak Interactions," Nucl. Phys. 22 (1961) 579-588.
- [4] S. Weinberg, "A Model of Leptons," Phys. Rev. Lett. 19 (1967) 1264–1266.
- [5] A. Salam, "Weak and Electromagnetic Interactions," Conf. Proc. C 680519 (1968) 367–377.
- [6] M. Yoshimura, "Unified Gauge Theories and the Baryon Number of the Universe," Phys. Rev. Lett. 41 (1978) 281–284.
 [Erratum: Phys.Rev.Lett. 42, 746 (1979)].
- [7] E. K. Akhmedov, G. C. Branco, and M. N. Rebelo, "Seesaw mechanism and structure of neutrino mass matrix," *Phys. Lett. B* 478 (2000) 215–223, arXiv:hep-ph/9911364.
- [8] P. Minkowski, " $\mu \to e \gamma$ at a Rate of One Out of 10^9 Muon Decays?," Phys. Lett. B 67 (1977) 421–428.
- [9] M. Gell-Mann, P. Ramond, and R. Slansky, "Complex Spinors and Unified Theories," Conf. Proc. C 790927 (1979) 315-321, arXiv:1306.4669 [hep-th].
- [10] R. N. Mohapatra and G. Senjanovic, "Neutrino Mass and Spontaneous Parity Nonconservation," Phys. Rev. Lett. 44 (1980) 912.
- [11] M. Magg and C. Wetterich, "Neutrino Mass Problem and Gauge Hierarchy," Phys. Lett. B 94 (1980) 61–64.
- [12] T. P. Cheng and L.-F. Li, "Neutrino Masses, Mixings and Oscillations in SU(2) x U(1) Models of Electroweak Interactions," Phys. Rev. D 22 (1980) 2860.
- [13] G. Lazarides, Q. Shafi, and C. Wetterich, "Proton Lifetime and Fermion Masses in an SO(10) Model," Nucl. Phys. B 181 (1981) 287–300.
- [14] R. N. Mohapatra and G. Senjanovic, "Neutrino Masses and Mixings in Gauge Models with Spontaneous Parity Violation," Phys. Rev. D 23 (1981) 165.
- [15] M. C. Gonzalez-Garcia, M. Maltoni, and T. Schwetz, "NuFIT: Three-Flavour Global Analyses of Neutrino Oscillation Experiments," *Universe* 7 (2021) no. 12, 459, arXiv:2111.03086 [hep-ph].
- [16] Particle Data Group Collaboration, P. A. Zyla et al., "Review of Particle Physics," PTEP 2020 (2020) no. 8, 083C01.
- [17] C. Jarlskog, "Commutator of the Quark Mass Matrices in the Standard Electroweak Model and a Measure of Maximal CP Nonconservation," Phys. Rev. Lett. 55 (1985) 1039.
- [18] V. D. Barger, S. Pakvasa, T. J. Weiler, and K. Whisnant, "Bimaximal mixing of three neutrinos," *Phys. Lett. B* 437 (1998) 107–116, arXiv:hep-ph/9806387.
- [19] P. F. Harrison, D. H. Perkins, and W. G. Scott, "Tri-bimaximal mixing and the neutrino oscillation data," *Phys. Lett. B* 530 (2002) 167, arXiv:hep-ph/0202074.
- [20] C. H. Albright, A. Dueck, and W. Rodejohann, "Possible Alternatives to Tri-bimaximal Mixing," Eur. Phys. J. C 70 (2010) 1099-1110, arXiv:1004.2798 [hep-ph].
- [21] G.-J. Ding, L. L. Everett, and A. J. Stuart, "Golden Ratio Neutrino Mixing and A₅ Flavor Symmetry," Nucl. Phys. B 857 (2012) 219–253, arXiv:1110.1688 [hep-ph].
- [22] Z.-z. Xing, "Flavor structures of charged fermions and massive neutrinos," Phys. Rept. 854 (2020) 1–147, arXiv:1909.09610 [hep-ph].
- [23] E. Ma, "Tribimaximal neutrino mixing from a supersymmetric model with A4 family symmetry," Phys. Rev. D 73 (2006) 057304, arXiv:hep-ph/0511133.
- [24] E. Ma, "A(4) symmetry and neutrinos with very different masses," Phys. Rev. D 70 (2004) 031901, arXiv:hep-ph/0404199.
- [25] G. Altarelli and F. Feruglio, "Tri-bimaximal neutrino mixing from discrete symmetry in extra dimensions," Nucl. Phys. B 720 (2005) 64–88, arXiv:hep-ph/0504165.
- [26] G. Chauhan, P. S. B. Dev, I. Dubovyk, B. Dziewit, W. Flieger, K. Grzanka, J. Gluza, B. Karmakar, and S. Zieba, "Phenomenology of Lepton Masses and Mixing with Discrete Flavor Symmetries," arXiv:2310.20681 [hep-ph].

- [27] B. Karmakar and A. Sil, "Nonzero θ_{13} and leptogenesis in a type-I seesaw model with A_4 symmetry," *Phys. Rev. D* **91** (2015) 013004, arXiv:1407.5826 [hep-ph].
- [28] D. Borah and B. Karmakar, "A₄ flavour model for Dirac neutrinos: Type I and inverse seesaw," Phys. Lett. B 780 (2018) 461–470, arXiv:1712.06407 [hep-ph].
- [29] Planck Collaboration, N. Aghanim et al., "Planck 2018 results. VI. Cosmological parameters," Astron. Astrophys. 641 (2020) A6, arXiv:1807.06209 [astro-ph.CO]. [Erratum: Astron.Astrophys. 652, C4 (2021)].
- [30] H. Ejiri, "Neutrino-Mass Sensitivity and Nuclear Matrix Element for Neutrinoless Double Beta Decay," Universe 6 (2020) no. 12, 225.
- [31] M. Agostini, G. Benato, J. A. Detwiler, J. Menéndez, and F. Vissani, "Toward the discovery of matter creation with neutrinoless $\beta\beta$ decay," Rev. Mod. Phys. **95** (2023) no. 2, 025002, arXiv:2202.01787 [hep-ex].
- [32] CUORE Collaboration, D. Q. Adams et al., "Improved Limit on Neutrinoless Double-Beta Decay in ¹³⁰Te with CUORE," Phys. Rev. Lett. 124 (2020) no. 12, 122501, arXiv:1912.10966 [nucl-ex].
- [33] **GERDA** Collaboration, M. Agostini *et al.*, "Probing Majorana neutrinos with double-β decay," *Science* **365** (2019) 1445, arXiv:1909.02726 [hep-ex].
- [34] KamLAND-Zen Collaboration, A. Gando et al., "Search for Majorana Neutrinos near the Inverted Mass Hierarchy Region with KamLAND-Zen," Phys. Rev. Lett. 117 (2016) no. 8, 082503, arXiv:1605.02889 [hep-ex]. [Addendum: Phys.Rev.Lett. 117, 109903 (2016)].
- [35] **SuperNEMO** Collaboration, R. Arnold *et al.*, "Measurement of the distribution of ^207Bi depositions on calibration sources for SuperNEMO," *JINST* **16** (2021) no. 07, T07012, arXiv:2103.14429 [physics.ins-det].
- [36] CUORE Collaboration, C. Alduino et al., "Double-beta decay of ¹³⁰Te to the first 0⁺ excited state of ¹³⁰Xe with CUORE-0," Eur. Phys. J. C 79 (2019) no. 9, 795, arXiv:1811.10363 [nucl-ex].
- [37] E. Ma, "Neutrino mass from triplet and doublet scalars at the TeV scale," Phys. Rev. D 66 (2002) 037301, arXiv:hep-ph/0204013.
- [38] A. E. Cárcamo Hernández and I. de Medeiros Varzielas, "Δ(27) framework for cobimaximal neutrino mixing models," Phys. Lett. B 806 (2020) 135491, arXiv:2003.01134 [hep-ph].
- [39] M. Dey and S. Roy, "Revisiting the Dirac Nature of Neutrinos," arXiv:2403.12461 [hep-ph].
- [40] M. Dey and S. Roy, "Unveiling Neutrino Mysteries with $\Delta(27)$ Symmetry," arXiv:2309.14769 [hep-ph].
- [41] **LEGEND** Collaboration, N. Abgrall *et al.*, "The Large Enriched Germanium Experiment for Neutrinoless $\beta\beta$ Decay: LEGEND-1000 Preconceptual Design Report," arXiv:2107.11462 [physics.ins-det].