Asymmetry models and separability for multi-way contingency tables with ordinal categories

Hisaya Okahara * Kouji Tahata

Department of Information Sciences, Tokyo University of Science

Abstract

In this paper, we propose a model that indicates the asymmetry structure for cell probabilities in multivariate contingency tables with the same ordered categories. The proposed model is the closest to the symmetry model in terms of the f-divergence under certain conditions and incorporates various asymmetry models as special cases, including existing models. We elucidate the relationship between the proposed model and conventional models from several aspects of divergence in f-divergence. Furthermore, we provide theorems showing that the symmetry model can be decomposed into two or more models, each imposing less restrictive parameter constraints than the symmetry condition. We also discuss the properties of goodness-of-fit statistics, particularly focusing on the likelihood ratio test statistics and Wald test statistics. Finally, we summarize the proposed model and discuss some problems and future work.

1 Introduction

Multivariate categorical data with ordinal variables is very common in various fields such as psychology, education, and social science, and is typically summarized into a multi-way contingency table when these variables are commensurable. A significant objective of statistical analysis of contingency tables is to discover the association among these variables. Descriptive models are useful for reducing complexity only if the parameters can be readily interpreted as representing structures of the data. When these variables are expected to be highly associated, it is worth exploring whether the association pattern exhibits a symmetry structure instead of independence. The emphasis on the symmetry structure is for evaluation purposes, such as in panel studies and in the study of social mobility. Particularly, in the analysis of occupational mobility tables, researchers have found that the cells on the main diagonal, where each variable takes the same value, often need to be treated separately from the other cells in the table, and that analyzing the table with the cells on the main diagonal deleted is often worthwhile (see [8, 9, 11]). The issues of symmetry have been addressed in many studies, for example, [1, 19, 20]. Although models having symmetry structures are straightforward to interpret, these models have quite strong constraints on the parameters. Therefore, we explore various models with asymmetry structures that have less restrictive parameters and more beneficial properties.

For two-way contingency tables, [9, 10] discussed the diagonals-parameter symmetry (DPS) model, and [1] focused on a special case of the DPS model, denoted as the linear DPS (LDPS) model. Furthermore, [22] introduced the extended LDPS (ELDPS) model. The LDPS and ELDPS models relax the strong constraints of the symmetry (S) model by introducing parameters that depend on the distance from the main diagonal and have theoretical justification. For ordinal matched pairs data, it is often reasonable to assume an underlying continuous distribution that is approximately a bivariate normal distribution with equal marginal standard deviations, and the LDPS and ELDPS models satisfy the

^{*}Email : hisaya.okahara@gmail.com. Postal address : Yamazaki, Noda, 2788510, Chiba, Japan.

relationships between the joint probabilities in the discrete case. More generally, the LDPS model can be extended to the ordinal quasi-symmetry (OQS) model by replacing the integer scores with fixed monotonic scores. The quasi-symmetry (QS) model further generalizes the OQS model by relaxing the ordinal constraints, allowing for more flexible parameter structures. Recent studies have further extended these models. [12, 13] generalized the QS and OQS models based on f-divergence, which was introduced by [4, 5] and encompasses a wide family of divergences, including Cressie-Read power divergence. Additionally, [20] introduced a model that bridges the gap between these models. These generalized models are particularly important because they can improve the goodness-of-fit of the models without requiring additional parameters.

For multi-way contingency tables, [21, 24] considered linear diagonals-parameter symmetry (LS^T) and extended LS^T (ELS^T) models as the extension of the LDPS and ELDPS models, and introduced generalized LS^T (GLS^T) model. Generally, the LS^T and ELS^T models may be appropriate for an ordinal table if it is reasonable to assume an underlying multivariate normal distribution with moment restrictions, while the GLS^T model does not impose any constraints. Despite the importance of these models, there has been limited research on generalizing models for multi-way contingency tables. To address this issue, we propose a model that enhances the goodness-of-fit without increasing the number of parameters. This parsimony in parameters allows for more interpretable and computationally efficient modeling of the stochastic structure. Specifically, we introduce an f-divergence based asymmetry model for multi-way contingency tables with ordinal categories, which is less restrictive than the S^T model. Furthermore, under certain conditions, the proposed model is the closest model to the S^T model, allowing for unequal marginal means, variances, and correlations while preserving key properties of the GLS^T model.

The rest of this paper is organized as follows. Section 2 introduces the notation used throughout the article and reviews the conventional models for contingency tables with the same ordinal variables. Section 3 broadens the scope to the generalized family of the GLS^T model based on the *f*-divergence, which is the closest model to the symmetry model under certain conditions. Section 4 discusses some special cases of the proposed model and its dependent structures, and investigates its properties from different perspectives for a multivariate contingency table. Section 5 provides partitions of the S^T model, which allows us to investigate the reasons for a poor fit of the S^T model. Section 6 shows the relationship between these models and corresponding goodness-of-fit statistics, and presents properties of several types of test statistics. Finally, Section 7 summarizes the paper and discusses future work.

2 Notation and background

In this section, we introduce the notation used throughout the paper and provide an overview of conventional models. Additional notation will be introduced in Section 6. For a comprehensive list of notation, see Table 2 in Appendix A.

2.1 Notation

Let $V = \{1, \ldots, T\}$ be an index set, and we use $(X_j, j \in V)$ to denote variables with $X_j \in \mathcal{I} = \{1, \ldots, r\}$. Let $\mathcal{I}_V = \mathcal{I} \times \cdots \times \mathcal{I} = \mathcal{I}^{|V|}$, where |V| represents the number of elements in the set V. Elements of \mathcal{I}_V are referred to as cells of the contingency table, and there are r^T cells in total. We generically denote a cell by i, with $i = (i_1, \ldots, i_T) \in \mathcal{I}_V$. The joint probability mass function of $X = (X_1, \ldots, X_T)$ is denoted by π , with

$$\pi_{\boldsymbol{i}} = \Pr(X_1 = i_1, \dots, X_T = i_T), \quad \boldsymbol{i} \in \mathcal{I}_V,$$

which satisfies $\sum_{i \in \mathcal{I}_V} \pi_i = 1$ and $\pi_i > 0$.

To explicitly describe the S^T model, we define the following sets.

$$D(\mathbf{i}) = D(i_1, \dots, i_T)$$

= { $(j_1, \dots, j_T) \in \mathcal{I}_V \mid (j_1, \dots, j_T)$ is any permutation of (i_1, \dots, i_T) }

The cell probabilities that satisfy the symmetry structure are defined as follows.

$$\pi_{\boldsymbol{i}}^{S} = \frac{1}{|D(\boldsymbol{i})|} \sum_{\boldsymbol{j} \in D(\boldsymbol{i})} \pi_{\boldsymbol{j}}, \quad \boldsymbol{i} \in \mathcal{I}_{V}.$$
(1)

2.2 Conventional models

Let $\mathbf{p} = (p_i)$ and $\mathbf{q} = (q_i)$ are two discrete finite multivariate probability distributions. The *f*-divergence between \mathbf{p} and \mathbf{q} is defined as

$$I^{C}(\mathbf{p}:\mathbf{q}) = \sum_{i \in \mathcal{I}_{V}} q_{i} f\left(\frac{p_{i}}{q_{i}}\right),$$

where f is a strictly convex function on $(0, \infty)$ with f(1) = 0, $f(0) = \lim_{x \to 0} f(x)$, $0 \cdot f(0/0) = 0$ and $0 \cdot f(a/0) = a \lim_{t \to \infty} [f(t)/t]$ (see [5, 6]).

For two-way contingency tables, let X_1 and X_2 denote the row and column variables, and take the values $i, j \in \mathcal{I}$, respectively. Assuming a set of the integer scores, the ELS² model is expressed as

$$\frac{\pi_{ij}}{\pi_{ji}} = \alpha^{j-i} \beta^{j^2 - i^2}, \quad i < j.$$

The ELS² model is reduced to the LS² model when $\beta = 1$. Herein, we consider random variables Z_1 and Z_2 having a joint bivariate normal distribution with means $E(Z_1) = \mu_1$ and $E(Z_2) = \mu_2$, variances $Var(Z_1) = \sigma_1^2$ and $Var(Z_2) = \sigma_2^2$, and correlation $Cor(Z_1, Z_2) = \rho$. Then the joint bivariate normal density function $f(z_1, z_2)$ satisfies

$$\frac{f(z_1, z_2)}{f(z_2, z_1)} = \exp\left[-\frac{z_1 - z_2}{2(1 - \rho^2)} \left\{ \left(\frac{1}{\sigma_1^2} - \frac{1}{\sigma_2^2}\right)(z_1 + z_2) - 2\left(\frac{\mu_1}{\sigma_1^2} - \frac{\mu_2}{\sigma_2^2}\right) - \frac{2\rho(\mu_1 - \mu_2)}{\sigma_1\sigma_2} \right\} \right].$$

[1] described the relationship between the LS² model and the joint bivariate normal distribution as follows: When $\sigma_1^2 = \sigma_2^2$, the ratio $f(z_1, z_2)/f(z_2, z_1)$ takes the form $\xi^{z_2-z_1}$ for some constant ξ . Hence, the LS² model may be appropriate for a square ordinal contingency table if it is reasonable to assume an underlying bivariate normal distribution with equal marginal variances. On the other hand, [22] elucidated that the ELS² model, rather than the LS² model, would be more appropriate. This is the case if the underlying bivariate normal distribution does not require equal marginal variances.

Assume that $\{u_i\}$ is a set of known scores $u_1 < u_2 < \cdots < u_r$ that can be assigned to the row and columns. [20] proposed the asymmetry $(AS_k[f])$ model based on the *f*-divergence that fills the gap between the QS[*f*] model [13] and the OQS[*f*] model [12] by considering a natural extension on the asymmetry. For a given $k \in \{1, \ldots, r-1\}$, the AS_k[*f*] model is defined as

$$\pi_{ij} = \pi_{ij}^S F^{-1} \left(\sum_{h=1}^k u_i^h \alpha_h + \gamma_{ij} \right), \quad i, j \in \mathcal{I},$$

where $\gamma_{ij} = \gamma_{ji}$, $\pi_{ij}^S = (\pi_{ij} + \pi_{ji})/2$, f is a twice-differentiable and strictly convex function, and F(t) = f'(t). The AS_k[f] model is reduced to the OQS[f] model when k = 1 and the QS[f] model when k = r - 1. For $k \in \{2, \ldots, r - 2\}$, the AS_k[f] model is more restrictive than the QS[f] model, but is less restrictive than the OQS[f] model. Thus, it provides intermediate asymmetry models to fill the gap between the QS[f] and OQS[f] models.

For multi-way r^T contingency tables, [24] considered the GLS^T models defined as

$$\pi_{\boldsymbol{i}} = \left(\prod_{s=1}^{T} \alpha_s^{i_s}\right) \left(\prod_{t=1}^{T} \beta_t^{i_t^2}\right) \left(\prod_{s=1}^{T-1} \prod_{t=s+1}^{T} \gamma_{st}^{i_s i_t}\right) \psi_{\boldsymbol{i}}, \quad \boldsymbol{i} \in \mathcal{I}_V,$$
(2)

where $\psi_i = \psi_j$ and $j \in D(i)$. To ensure identifiability, we may set constraints such as $\alpha_T = \beta_T = \gamma_{(T-1)T} = 1$ without loss of generality. The special cases of this model are the ELS^T model with $\{\gamma_{st} = 1\}$, the LS^T model with $\{\beta_s = 1\}$ and $\{\gamma_{st} = 1\}$, and the S^T model with $\{\alpha_s = 1\}$, $\{\beta_s = 1\}$ and $\{\gamma_{st} = 1\}$. For a multivariate table with the same classifications, [24] argued that the GLS^T model may be appropriate if it is reasonable to assume an underlying multivariate normal distribution without any moment restrictions. Moreover, [21, 24] provided the decomposition properties of the S^T model using these models.

In this paper, we focus on an asymmetry model based on f-divergence that incorporates moment constraints specific to multivariate contingency tables. Furthermore, we discuss the relationship between the proposed model and the GLS^T model and present the advantageous properties of the proposed model.

3 Modeling based on the f-divergence

We define the proposed model, denoted as f-divergence based GLS^T ($\text{GLS}^T[f]$) model, and provide its construction in the following theorem.

Definition 1. The $GLS^{T}[f]$ model is defined by

$$\pi_{i} = \pi_{i}^{S} F^{-1} \left(\sum_{s=1}^{T} \alpha_{s} i_{s} + \sum_{s=1}^{T} \beta_{s} i_{s}^{2} + \sum_{s=1}^{T-1} \sum_{t=s+1}^{T} \gamma_{st} i_{s} i_{t} + \psi_{i} \right), \quad i \in \mathcal{I}_{V},$$
(3)

where $\psi_i = \psi_j$ and $j \in D(i)$. For identifiability, we may set, e.g., $\alpha_T = \beta_T = \gamma_{(T-1)T} = 0$ without loss of generality.

Theorem 1. The GLS^T[f] model is the closest to the S^T model in terms of the f-divergence under the conditions where the sums $\{\sum_{j \in D(i)} \pi_j\}, \sum_{i \in \mathcal{I}_V} i_s^h \pi_i \ (s \in V)$ for fixed $h \in \{1, 2\}$, and $\sum_{i \in \mathcal{I}_V} i_s i_t \pi_i \ (s, t \in V, s < t)$ are given.

Proof. For $T \geq 3$, consider minimizing the f-divergence under conditions that

$$v_{i}^{S} = \sum_{j \in D(i)} \pi_{j}, \quad i \in \mathcal{I}_{V},$$
$$v_{s}^{h} = \sum_{i \in \mathcal{I}_{V}} i_{s}^{h} \pi_{i}, \quad s \in V, \ h \in \{1, 2\},$$

and

$$v_{st} = \sum_{i \in \mathcal{I}_V} i_s i_t \pi_i, \quad s, t \in V, \ s < t,$$

are given. Note that the superscript of v^S_i does not denote an exponent. The Lagrange function is written as

$$L(\{\pi_i\}) = I^C(\boldsymbol{\pi} : \boldsymbol{\pi}^S) + \sum_{\substack{h \in \{1,2\} \\ s \in V}} \sum_{s \in V} \lambda_s^{(h)} \left(\sum_{i \in \mathcal{I}_V} i_s^h \pi_i - v_s^h \right) + \sum_{\substack{s,t \in V \\ s < t}} \lambda_{st} \left(\sum_{i \in \mathcal{I}_V} i_s i_t \pi_i - v_{st} \right) + \sum_{i \in \mathcal{I}_V} \phi_i \left(\sum_{j \in D(i)} \pi_j - v_i^S \right).$$

Equating the partial derivative of $L(\{\pi_i\})$ to 0 with respect to π_i gives

$$f'\left(\frac{\pi_{\boldsymbol{i}}}{\pi_{\boldsymbol{i}}^{S}}\right) + \sum_{h \in \{1,2\}} \sum_{s \in V} \lambda_{s}^{(h)} i_{s}^{h} + \sum_{\substack{s,t \in V\\s < t}} \lambda_{st} i_{s} i_{t} + \sum_{\boldsymbol{j} \in D(\boldsymbol{i})} \phi_{\boldsymbol{j}} = 0$$

Let $f', -\lambda_s^{(1)}, -\lambda_s^{(2)}, -\lambda_{st}$ and $-\sum_{j \in D(i)} \phi_j$ denote $F, \alpha_s, \beta_s, \gamma_{st}$ and ψ_i , respectively. We have

$$F\left(\frac{\pi_{\boldsymbol{i}}}{\pi_{\boldsymbol{i}}^{S}}\right) = \sum_{s \in V} \alpha_{s} i_{s} + \sum_{s \in V} \beta_{s} i_{s}^{2} + \sum_{\substack{s,t \in V\\s < t}} \gamma_{st} i_{s} i_{t} + \psi_{\boldsymbol{i}},$$

where $\psi_i = \psi_j$ and $j \in D(i)$. Since f is a strictly convex function, it follows that F'(x) = f''(x) > 0 for all x. Hence, F is strictly monotone, which ensures that F^{-1} exists. From the above equation, we obtain

$$\pi_{\boldsymbol{i}} = \pi_{\boldsymbol{i}}^{S} F^{-1} \left(\sum_{s \in V} \alpha_{s} i_{s} + \sum_{s \in V} \beta_{s} i_{s}^{2} + \sum_{\substack{s,t \in V\\s < t}} \gamma_{st} i_{s} i_{t} 1 + \psi_{\boldsymbol{i}} \right), \quad \boldsymbol{i} \in \mathcal{I}_{V},$$

where $\psi_i = \psi_j$ and $j \in D(i)$.

Similarly, we can define two new models as special cases of the $\text{GLS}^T[f]$ model: the *f*-divergence based ELS^T ($\text{ELS}^T[f]$) model, obtained by setting { $\gamma_{st} = 0$ }, and the *f*-divergence based LS^T ($\text{LS}^T[f]$) model, obtained by setting { $\beta_s = 0$ } and { $\gamma_{st} = 0$ }. Furthermore, we establish the corresponding corollaries for these models, which can be derived using the same approach as the $\text{GLS}^T[f]$ model, as follows.

Corollary 1. The ELS^T[f] model is the closest to the S^T model in terms of the f-divergence under the conditions where the sums $\{\sum_{j \in D(i)} \pi_j\}$ and $\sum_{i \in \mathcal{I}_V} i_s^h \pi_i$ $(s \in V)$ for fixed $(h \in \{1, 2\})$ are given.

Corollary 2. The $\mathrm{LS}^{T}[f]$ model is the closest to the S^{T} model in terms of the *f*-divergence under the conditions where the sums $\{\sum_{j \in D(i)} \pi_{j}\}$ and $\sum_{i \in \mathcal{I}_{V}} i_{s} \pi_{i}$ $(s \in V)$ are given.

More generally, the $\mathrm{LS}^{T}[f]$ model can be extended to the *f*-divergence based OQS (OQS[*f*]) model by replacing the integer scores with fixed monotonic scores, as discussed in [15]. This extension allows for more flexible modeling of ordinal data while maintaining the desirable properties of the $\mathrm{LS}^{T}[f]$ model.

4 Properties of the $GLS^{T}[f]$ model

4.1 Variants of the proposed model

This section shows that the $\text{GLS}^T[f]$ model is reduced to some existing models by choosing a divergence in *f*-divergence. From (3) and the constraints $\pi_i^S = \pi_j^S$ and $\psi_i = \psi_j$ for any $j \in D(i)$,

$$\sum_{\boldsymbol{j}\in D(\boldsymbol{i})} \pi_{\boldsymbol{j}} = \pi_{\boldsymbol{i}}^{S} \sum_{\boldsymbol{j}\in D(\boldsymbol{i})} F^{-1} \left(\sum_{s=1}^{T} \alpha_{s} j_{s} + \sum_{s=1}^{T} \beta_{s} j_{s}^{2} + \sum_{s=1}^{T-1} \sum_{t=s+1}^{T} \gamma_{st} j_{s} j_{t} + \psi_{\boldsymbol{j}} \right),$$

$$\Leftrightarrow \quad |D(\boldsymbol{i})| = \sum_{\boldsymbol{j}\in D(\boldsymbol{i})} F^{-1} \left(\sum_{s=1}^{T} \alpha_{s} j_{s} + \sum_{s=1}^{T} \beta_{s} j_{s}^{2} + \sum_{s=1}^{T-1} \sum_{t=s+1}^{T} \gamma_{st} j_{s} j_{t} + \psi_{\boldsymbol{j}} \right).$$
(4)

We can explicitly represent the stochastic structures using (4) as follows.

(i) If $f(x) = x \log x$ (x > 0), then the *f*-divergence reduces to the Kullback-Leibler divergence [14], $F^{-1}(x) = e^{x-1}$, and the $\text{GLS}^T[f]$ model (3) becomes

$$\pi_{\boldsymbol{i}} = \pi_{\boldsymbol{i}}^{S} \exp\left(\sum_{s=1}^{T} \alpha_{s} i_{s} + \sum_{s=1}^{T} \beta_{s} i_{s}^{2} + \sum_{s=1}^{T-1} \sum_{t=s+1}^{T} \gamma_{st} i_{s} i_{t} + \psi_{\boldsymbol{i}} - 1\right), \quad \boldsymbol{i} \in \mathcal{I}_{V},$$

which, with the help of (4), is transformed to

$$\pi_{i} = \pi_{i}^{S} \frac{|D(i)|a_{i}}{\sum_{j \in D(i)} a_{j}}, \quad i \in \mathcal{I}_{V},$$

$$(5)$$

with

$$a_{i} = \exp\left(\sum_{s=1}^{T} \alpha_{s} i_{s} + \sum_{s=1}^{T} \beta_{s} i_{s}^{2} + \sum_{s=1}^{T-1} \sum_{t=s+1}^{T} \gamma_{st} i_{s} i_{t}\right).$$

It can be easily shown that (5) is equivalent to the GLS^T model proposed by [24], which exhibits a departure from the S^T model in the form of a ratio. The following theorem, proved in Appendix B, gives the fact.

Theorem 2. When we set $f(x) = x \log x$ (x > 0), the $\text{GLS}^T[f]$ model is equivalent to GLS^T model.

Additionally, the following corollaries can be proposed as well.

Corollary 3. When we set $f(x) = x \log x$ (x > 0), the ELS^T[f] model is equivalent to ELS^T model.

Corollary 4. When we set $f(x) = x \log x$ (x > 0), the $LS^T[f]$ model is equivalent to LS^T model.

From equation (5), the following equation can be derived.

$$\frac{\pi_i}{\pi_j} = \frac{a_i}{a_j},\tag{6}$$

where $j \in D(i)$. From equation (6), it can be observed that the ratio of cell probabilities at symmetric positions is expressed in terms of the parameters $\{\alpha_s\}, \{\beta_s\}, \text{ and } \{\gamma_{st}\}$. Furthermore, by taking the logarithm of both sides, we can confirm that the ratio of cell probabilities is represented as a linear relationship in terms of the logarithmic function.

(ii) If $f(x) = (1-x)^2$, then the *f*-divergence reduces to the Pearsonian distance [16], $F^{-1}(x) = x/2+1$, and the GLS^T[*f*] model (3) becomes

$$\pi_{i} = \pi_{i}^{S} \left(\frac{1}{2} \sum_{s=1}^{T} \alpha_{s} i_{s} + \frac{1}{2} \sum_{s=1}^{T} \beta_{s} i_{s}^{2} + \frac{1}{2} \sum_{s=1}^{T-1} \sum_{t=s+1}^{T} \gamma_{st} i_{s} i_{t} + \frac{1}{2} \psi_{i} + 1 \right), \quad i \in \mathcal{I}_{V}.$$

Considering (4), it can be verified that the model reduces to

$$\pi_{i} = \pi_{i}^{S} \left(a_{i} - \frac{1}{|D(i)|} \sum_{j \in D(i)} a_{j} \right), \quad i \in \mathcal{I}_{V},$$

$$(7)$$

with

$$a_{i} = \frac{1}{2} \sum_{s=1}^{T} \alpha_{s} i_{s} + \frac{1}{2} \sum_{s=1}^{T} \beta_{s} i_{s}^{2} + \frac{1}{2} \sum_{s=1}^{T-1} \sum_{t=s+1}^{T} \gamma_{st} i_{s} i_{t}.$$

This model represents a departure from the symmetry model in the form of a difference. Under certain constraints, this model is the closest to the S^T model when the divergence is measured by the Pearsonian distance. Thus, we shall refer to this model as the Pearsonian GLS^T (PGLS^T) model. Then, the following equation can be derived.

$$\frac{\pi_{\boldsymbol{i}}}{\pi_{\boldsymbol{i}}^{S}} - \frac{\pi_{\boldsymbol{j}}}{\pi_{\boldsymbol{i}}^{S}} = a_{\boldsymbol{i}} - a_{\boldsymbol{j}} \iff \pi_{\boldsymbol{i}}^{c} - \pi_{\boldsymbol{j}}^{c} = \frac{a_{\boldsymbol{i}} - a_{\boldsymbol{j}}}{|D(\boldsymbol{i})|},$$
(8)

where $j \in D(i)$ and

$$\pi_{\boldsymbol{i}}^c = \frac{\pi_{\boldsymbol{i}}}{\sum_{\boldsymbol{j} \in D(\boldsymbol{i})} \pi_{\boldsymbol{j}}}$$

Equation (8) shows that the PGLS^T model represents the structure of the difference between two conditional probabilities, which is linearly expressed in terms of the parameters $\{\alpha_s\}, \{\beta_s\}$, and $\{\gamma_{st}\}$.

(iii) If

$$f_{\lambda}(x) = \frac{x(x^{\lambda} - 1)}{\lambda(\lambda + 1)}, \quad F_{\lambda}^{-1}(x) = \left(\lambda x + \frac{1}{\lambda + 1}\right)^{\frac{1}{\lambda}}, \quad x > 0$$

where λ is a real-valued parameter, then *f*-divergence reduces to the Cressie-Read power divergence [18], and model (1) becomes

$$\pi_{\boldsymbol{i}} = \pi_{\boldsymbol{i}}^{S} \left(\lambda \left(\sum_{s=1}^{T} \alpha_{s} i_{s} + \sum_{s=1}^{T} \beta_{s} i_{s}^{2} + \sum_{s=1}^{T-1} \sum_{t=s+1}^{T} \gamma_{st} i_{s} i_{t} + \psi_{\boldsymbol{i}} \right) + \frac{1}{\lambda + 1} \right)^{\frac{1}{\lambda}}, \quad \boldsymbol{i} \in \mathcal{I}_{V}.$$
(9)

When $\lambda = 0$ or $\lambda = -1$, $f_0(x) = \lim_{\lambda \to 0} [f_\lambda(x)]$ and $f_{-1}(x) = \lim_{\lambda \to -1} [f_\lambda(x)]$. Under certain constraints, model (9) is the closest to the S^T model when the divergence is measured by the Cressie-Read power divergence. This model also reduces to model (5) when $\lambda = 0$ and model (7) when $\lambda = 1$. Additionally, when $\lambda = -1/2$, we consider a model based on the Hellinger distance, which is referred to as the Hellinger GLS^T (HGLS^T) model.

In a similar manner to the above, the following relationship can be derived.

$$\left(\frac{\pi_{\boldsymbol{i}}}{\pi_{\boldsymbol{i}}^{S}}\right)^{\lambda} - \left(\frac{\pi_{\boldsymbol{j}}}{\pi_{\boldsymbol{i}}^{S}}\right)^{\lambda} = \lambda \left(a_{\boldsymbol{i}}^{\lambda} - a_{\boldsymbol{j}}^{\lambda}\right) \iff (\pi_{\boldsymbol{i}}^{c})^{\lambda} - (\pi_{\boldsymbol{j}}^{c})^{\lambda} = \frac{\lambda \left(a_{\boldsymbol{i}}^{\lambda} - a_{\boldsymbol{j}}^{\lambda}\right)}{|D(\boldsymbol{i})|^{\lambda}},\tag{10}$$

where $j \in D(i)$ and

$$a_{i} = \left(\sum_{s=1}^{T} \alpha_{s} i_{s} + \sum_{s=1}^{T} \beta_{s} i_{s}^{2} + \sum_{s=1}^{T-1} \sum_{t=s+1}^{T} \gamma_{st} i_{s} i_{t}\right).$$

It can be verified that equation (10) reduces to equation (8) when $\lambda = 1$. This equation represents the difference of conditional probabilities raised to the power of λ as a linear relationship. Furthermore, it suggests that by adjusting the parameter λ , the model may exhibit robustness to the underlying assumed probability distribution.

It is also possible to consider other models by using different types of divergence in the f-divergence framework, where the divergence is minimized under certain conditions. Notably, the GLS^T model (5), which represents the log-linear form, and the PGLS^T model (7), which shows the difference between two conditional probabilities in symmetric cells, have stochastic structures that are straightforward to interpret.

4.2 Interpretation of the proposed model

In this section, we provide an interpretation of the parameters $\boldsymbol{\alpha} = (\alpha_1, \ldots, \alpha_T)$, $\boldsymbol{\beta} = (\beta_1, \ldots, \beta_T)$, and $\boldsymbol{\gamma} = (\gamma_{12}, \ldots, \gamma_{(T-1)T})$ of the proposed model, as well as how the choice of the divergence captures stochastic structures. To facilitate understanding for the reader, we consider the GLS^T model expressed as

$$\log \pi_{i} = \psi + \sum_{s=1}^{T} \alpha_{s} i_{s} + \sum_{s=1}^{T} \beta_{s} i_{s}^{2} + \sum_{s=1}^{T-1} \sum_{t=s+1}^{T} \gamma_{st} i_{s} i_{t} + \psi_{i}, \quad i \in \mathcal{I}_{V},$$
(11)

where $\psi_{i} = \psi_{j}$ and $j \in D(i)$. We may set, e.g., $\sum \alpha_{s} = \sum \beta_{s} = \sum \sum \gamma_{st} = 0$ without loss of generality. In the following, we provide an interpretation of the GLS^T model (11) from the log-linear model perspective. For simplicity, we present a saturated model for a three-way contingency table as follows.

$$\log \pi_{(i_1, i_2, i_3)} = \mu + \sum_{u=1}^{3} \mu_u(i_u) + \sum_{s=1}^{2} \sum_{t=s+1}^{3} \mu_{st}(i_s, i_t) + \mu_{123}(i_1, i_2, i_3), \quad (i_1, i_2, i_3) \in \mathcal{I}_{\{1, 2, 3\}}$$
(12)

where, for example,

$$\sum_{i_u=1}^r \mu_u(i_u) = 0, \quad \sum_{i_u=1}^r \mu_{123}(i_1, i_2, i_3) = 0, \quad 1 \le u \le 3,$$
$$\sum_{i_s=1}^r \mu_{st}(i_s, i_t) = \sum_{i_t=1}^r \mu_{st}(i_s, i_t) = 0, \quad 1 \le s < t \le 3.$$

The $\{\mu_s(i_s)\}\$ are effect terms that represent the main effects of the levels for each variable. The $\{\mu_{st}(i_s, i_t)\}\$ and $\{\mu_{123}(i_1, i_2, i_3)\}\$ are interaction terms that reflect deviations from independence (see [2]). For a multi-way contingency table, we shall consider higher-order interaction terms.

While log-linear models are commonly used to describe associations among variables in terms of interaction, the GLS^T model focuses on characterizing associations conditioned on the symmetric structure. For the GLS^T model (11), the first term ψ is an intercept corresponding to μ in (12). The second term, $\sum_s \alpha_s i_s$, expresses the linear effects of the levels for each variable, while the third term, $\sum_s \beta_s i_s^2$, captures the non-linear effects. In other words, these two terms can be regarded as explanatory variables that approximate the main effects as the response variables in a regression-like manner. The fourth terms, $\sum \sum_{s < t} \gamma_{st} i_s i_t$, represent the first-order interactions between each variables and correspond to the third terms in (12). The last term, ψ_i , represents the $(T-1)^{\text{th}}$ order interaction among all variables under the assumption of a symmetric structure. Note that the GLS^T model is given within the log-linear framework conditioned on the symmetric structure. While the log-linear models can specify how associations among categorical variables, the GLS^T model focuses on associations among symmetric cells. Notably, when all elements in γ equal 0, we interpret this as all first-order interactions being the same, not absent; the association terms in the log-linear model reflect deviations from independence.

Analogously, we can also represent associations using different divergences. In general, we give the $\operatorname{GLS}^{T}[f]$ model with conditional probabilities π_{i}^{c} as

$$F(|D(\mathbf{i})| \ \pi_{\mathbf{i}}^{c}) = \sum_{s=1}^{T} \alpha_{s} i_{s} + \sum_{s=1}^{T} \beta_{s} i_{s}^{2} + \sum_{s=1}^{T-1} \sum_{t=s+1}^{T} \gamma_{st} i_{s} i_{t} + \psi_{\mathbf{i}}, \quad \mathbf{i} \in \mathcal{I}_{V},$$

where $\psi_i = \psi_j$ and $j \in D(i)$. The proposed model enables a more detailed description of each effect and interaction. For instance, we can evaluate the linear effects by contrasting α and the first-order interactions with γ . Furthermore, if a function F is useful in the sense that a linear model with explanatory parameters is plausible for that link, it is noteworthy that the proposed model characterizes

the stochastic structure with respect to that link function. Thus, akin to the concept of link functions in generalized linear models (GLMs), the function F allows us to construct various paradigms that extend beyond the log-linear model framework.

5 Partition of symmetry

Since the S^T model rarely fits real data due to its strong restrictions, it is often necessary to consider more relaxed models that impose fewer constraints on the cell probabilities. In this section, we are interested in sets of models that, when combined, are equivalent to the S^T model. In other words, we seek to identify groups of models such that the S^T model holds if and only if all the models in a given group hold simultaneously. This approach allows us to decompose the S^T model into a collection of less restrictive models, which can be used to analyze the structure of the data more effectively. By examining the goodness-of-fit of these component models, as discussed in the next section (Section 6), we can gain insights into the specific aspects of the data that contribute to the poor fit of the S^T model. This information can be valuable in identifying the sources of the lack of fit and can guide the development of more appropriate models that better capture the underlying structure of the data.

In the following subsections, we will introduce several groups of models that, when combined, are equivalent to the S^T model. By studying these model groups and their relationships to the S^T model, we aim to provide a more comprehensive understanding of the structure of multivariate categorical data and to facilitate the selection of appropriate models for a given dataset.

Firstly, consider the marginal mean equality (ME^T) model defined as

$$\mu_1 = \mu_u, \quad u \in V,$$

where $\mu_u = E(X_u)$.

Secondly, consider the marginal variance equality (VE^T) model defined as

$$\sigma_1^2 = \sigma_u^2, \quad u \in V,$$

where $\sigma_u^2 = \operatorname{Var}(X_u)$.

Finally, consider the marginal correlation equality (CE^T) model defined as

$$\rho_{12} = \rho_{st}, \quad s, t \in V, \ s < t,$$

where

$$\rho_{st} = \frac{\mathrm{E}(X_s X_t) - \mu_s \mu_t}{\sigma_s \sigma_t}.$$

In addition, we can consider models that incorporate the conditions of several models. We consider the marginal mean and variance equality (MV^T) model defined as

$$\mu_1 = \mu_u$$
 and $\sigma_1^2 = \sigma_u^2$, $u \in V$,

and the marginal mean, variance, and correlation equality (MVC^T) model defined as

$$\mu_1 = \mu_u, \ \ \sigma_1^2 = \sigma_u^2 \ \ \text{and} \ \ \rho_{12} = \rho_{st}, \quad u, s, t \in V, \ s < t.$$

The CE^T and MVC^T models are defined under the condition of $T \ge 3$. The following theorem and corollaries detail how the S^T model can be decomposed into various combinations of models.

Theorem 3. The S^T model holds if and only if all the $GLS^T[f]$, ME^T , VE^T , and CE^T models hold.

Theorem 3 provides a comprehensive decomposition of the S^T model into four models. This result is particularly useful for identifying the sources of lack of fit in the S^T model. The proof of this theorem is provided in Appendix B. In addition to Theorem 3, we present four corollaries that offer alternative decompositions of the S^T model. These corollaries can be useful in different contexts, depending on the research question and the nature of the data. **Corollary 5.** The S^T model holds if and only if both the $GLS^T[f]$ and MVC^T models hold.

Corollary 6. The S^T model holds if and only if all the $ELS^T[f]$, ME^T and VE^T models hold.

Corollary 7. The S^T model holds if and only if both the $ELS^T[f]$ and MV^T models hold.

Corollary 8. The S^T model holds if and only if both the $LS^T[f]$ and ME^T models hold.

The proofs for these corollaries are omitted as they follow similar lines of reasoning as the proof of Theorem 3.

Remark 1. The theorem and corollaries presented above provide a powerful tool for investigating the reasons behind a poor fit of the S^T model to a given dataset. By decomposing the S^T model into less restrictive models, such as the $GLS^T[f]$, ME^T , VE^T , and CE^T models, we can identify which aspects of the data are not adequately captured by the S^T model. This approach allows us to pinpoint the specific properties of the data that contribute to the lack of fit, such as unequal marginal means, variances, or correlations.

6 Property of goodness-of-fit statistics

Let n_i denote the observed frequency in the *i*th cell, for $i \in \mathcal{I}_V$, in a r^T contingency table. Assume that the cell counts follow a multinomial distribution. Let m_i and \hat{m}_i denote the expected frequency in the *i*th cell and its maximum likelihood estimate under a given model, respectively. The maximum likelihood estimates (MLEs) of the expected frequencies can be obtained using the Newton-Raphson method to solve the log-likelihood equations. For instance, we can utilize the package Rsolnp in R to calculate the MLEs under any model.

To assess the goodness-of-fit of a model, one can use various test statistics such as the likelihood ratio Chi-square statistic. For a given model M, the likelihood ratio Chi-square statistic is defined as

$$G^{2}(\mathbf{M}) = 2 \sum_{\boldsymbol{i} \in \mathcal{I}_{V}} n_{\boldsymbol{i}} \log\left(\frac{n_{\boldsymbol{i}}}{\hat{m}_{\boldsymbol{i}}}\right),$$

with the corresponding degrees of freedom (df). Table 1 presents the df for various models in the context of an r^T contingency table, including the S^T , $GLS^T[f]$, $ELS^T[f]$, $LS^T[f]$, MVC^T , MV^T , ME^T , VE^T , and CE^T models.

Table 1: Degrees of freedom for various models in an r^T contingency table

Models	df
\mathbf{S}^T	$r^T - L^{(r,T)}$
$\mathrm{GLS}^T[f]$	$r^{T} - L^{(r,T)} - (T^{2} + 3T - 6)/2$
$\mathrm{ELS}^T[f]$	$r^T - L^{(r,T)} - 2(T-1)$
$\mathrm{LS}^T[f]$	$r^T - L^{(r,T)} - (T-1)$
MVC^T	$(T^2 + 3T - 6)/2$
MV^T	2(T-1)
ME^T	T-1
VE^T	T-1
CE^T	$(T^2 - T - 2)/2$
Note: $L^{(r,T)} = \binom{r+T-1}{T} = \frac{(r+T-1)!}{(r-1)!T!}$	

[7, 23] discussed the properties of test statistics for several models. In general, suppose that model M_3 holds if and only if both models M_1 and M_2 hold, and the following asymptotic equivalence holds.

$$G^{2}(M_{3}) = G^{2}(M_{1}) + G^{2}(M_{2}) + o_{p}(1),$$

where the df for M_3 equals the sum of the df for M_1 and M_2 . When both M_1 and M_2 are accepted with high probability, M_3 would also be accepted (see [3]).

To facilitate the discussion of the properties of goodness-of-fit statistics, we introduce some additional notation. Let \mathcal{P}_V be a subset of \mathcal{I}_V , defined as

$$\mathcal{P}_V = \{(\kappa_1, \ldots, \kappa_T) \mid \kappa_1 \leq \cdots \leq \kappa_T\} \subset \mathcal{I}_V.$$

For a set $V = \{1, \ldots, T\}$, we define a set $\mathcal{C}_V^{(2)}$ related to the parameters γ as

$$\mathcal{C}_{V}^{(2)} = \{ (\lambda_{1}, \lambda_{2}) \mid \lambda_{1} < \lambda_{2}, \ \lambda_{1} \leq T - 2 \} \subset V \times V.$$

We also introduce the concept of *lexicographical ordering* (\preceq) defined as

$$\forall \boldsymbol{v}, \boldsymbol{v}' \in \mathbb{N}^n, \ \boldsymbol{v} \leq \boldsymbol{v}' \ \Leftrightarrow \ \exists i \in \{1, \dots, n\}, \ (\forall j < i, v_j = v'_j) \land (v_i \leq v'_i).$$

Using the order \leq , we can define totally ordered sets (\mathcal{I}_V, \leq) , (\mathcal{P}_V, \leq) , and $(\mathcal{C}_V^{(2)}, \leq)$. For an arbitrary totally ordered set \mathcal{A} , we define a sequence of its elements $(\boldsymbol{a}_n)_{n\in\mathbb{N}}$ satisfying $\boldsymbol{a}_1 \leq \cdots \leq \boldsymbol{a}_{|\mathcal{A}|}$. This allows us to express each totally ordered set $\{\boldsymbol{\iota}_1, \ldots, \boldsymbol{\iota}_{|\mathcal{I}_V|}\}, \{\boldsymbol{\kappa}_1, \ldots, \boldsymbol{\kappa}_{|\mathcal{P}_V|}\}$ and $\{\boldsymbol{\lambda}_1, \ldots, \boldsymbol{\lambda}_{|\mathcal{C}_V^{(2)}|}\}$

as a permutation of the elements of \mathcal{I}_V , \mathcal{P}_V , and $\mathcal{C}_V^{(2)}$, respectively. These notations and concepts will be used in the following discussion of the properties of goodness-of-fit statistics for given models.

The $GLS^{T}[f]$ model may be expressed as

$$F\left(\frac{\pi_{i}}{\pi_{i}^{S}}\right) = \sum_{s=1}^{T-1} (i_{s} - i_{s+1})\alpha_{s} + \sum_{s=1}^{T-1} (i_{s}^{2} - i_{s+1}^{2})\beta_{s} + \sum_{s=1}^{T-2} \left(\sum_{t=s+1}^{T-1} (i_{s}i_{t} - i_{s}i_{t+1})\gamma_{st} + (i_{s}i_{T} - i_{s+1}i_{s+2})\gamma_{sT}\right) + \psi_{i}$$
(13)

where $\psi_i = \psi_j$ and $j \in D(i)$.

It is important to note that although the parameters $\{\alpha_s\}, \{\beta_s\}$ and $\{\gamma_{st}\}$ in (13) share the same notation as those in (3), they are not necessarily the same and are used here without ambiguity. Then, we discuss separating the S^T model under the condition of $T \geq 3$. Let

$$\boldsymbol{\pi} = \left(\pi_{\iota_{1}}, \dots, \pi_{\iota_{|\mathcal{I}_{V}|}}\right)^{\top},$$
$$F\left(\frac{\boldsymbol{\pi}}{\boldsymbol{\pi}^{S}}\right) = \left(F\left(\frac{\pi_{\iota_{1}}}{\pi_{\iota_{1}}^{S}}\right), \dots, F\left(\frac{\pi_{\iota_{|\mathcal{I}_{V}|}}}{\pi_{\iota_{|\mathcal{I}_{V}|}}^{S}}\right)\right)^{\top},$$
$$\boldsymbol{\theta} = (\boldsymbol{\alpha}, \boldsymbol{\beta}, \boldsymbol{\gamma}, \boldsymbol{\psi})^{\top},$$

where

$$\begin{aligned} \boldsymbol{\alpha} &= (\alpha_1, \dots, \alpha_{T-1}), \\ \boldsymbol{\beta} &= (\beta_1, \dots, \beta_{T-1}), \\ \boldsymbol{\gamma} &= \left(\gamma_{\boldsymbol{\lambda}_1}, \dots, \gamma_{\boldsymbol{\lambda}_{|\mathcal{C}_V^{(2)}|}}\right), \\ \boldsymbol{\psi} &= \left(\psi_{\boldsymbol{\kappa}_1}, \dots, \psi_{\boldsymbol{\kappa}_{|\mathcal{P}_V|}}\right). \end{aligned}$$

Then the $\operatorname{GLS}^T[f]$ model is expressed as

$$F\left(rac{oldsymbol{\pi}}{oldsymbol{\pi}^S}
ight) = oldsymbol{X}oldsymbol{ heta} = (oldsymbol{x}_1,\ldots,oldsymbol{x}_{d_2},oldsymbol{X}_T)oldsymbol{ heta}$$

where X_T is the $r^T \times L^{(r,T)}$ matrix of 1 or 0 elements determined from the structure of the $\text{GLS}^T[f]$ model, and

$$\begin{aligned} \boldsymbol{x}_1 &= \boldsymbol{J}_r \otimes \boldsymbol{1}_{r^{T-1}} - \boldsymbol{1}_r \otimes \boldsymbol{J}_r \otimes \boldsymbol{1}_{r^{T-2}}, \\ &\vdots \\ \boldsymbol{x}_{T-1} &= \boldsymbol{1}_{r^{T-2}} \otimes \boldsymbol{J}_r \otimes \boldsymbol{1}_r - \boldsymbol{1}_{r^{T-1}} \otimes \boldsymbol{J}_r, \\ \boldsymbol{x}_T &= \boldsymbol{J}_r^2 \otimes \boldsymbol{1}_{r^{T-1}} - \boldsymbol{1}_r \otimes \boldsymbol{J}_r^2 \otimes \boldsymbol{1}_{r^{T-2}}, \\ &\vdots \\ \boldsymbol{x}_{2T-2} &= \boldsymbol{1}_{r^{T-2}} \otimes \boldsymbol{J}_r^2 \otimes \boldsymbol{1}_r - \boldsymbol{1}_{r^{T-1}} \otimes \boldsymbol{J}_r^2, \\ \boldsymbol{x}_{2T-1} &= (\boldsymbol{J}_r \otimes \boldsymbol{1}_{r^{T-1}}) \odot (\boldsymbol{1}_r \otimes \boldsymbol{J}_r \otimes \boldsymbol{1}_{r^{T-2}}) - (\boldsymbol{1}_r \otimes \boldsymbol{J}_r \otimes \boldsymbol{1}_{r^{T-2}}) \odot (\boldsymbol{1}_{r^2} \otimes \boldsymbol{J}_r \otimes \boldsymbol{1}_{r^{T-3}}), \\ &\vdots \\ \boldsymbol{x}_{d_2} &= (\boldsymbol{1}_{r^{T-3}} \otimes \boldsymbol{J}_r \otimes \boldsymbol{1}_{r^2}) \odot (\boldsymbol{1}_{r^{T-2}} \otimes \boldsymbol{J}_r \otimes \boldsymbol{1}_r) - (\boldsymbol{1}_{r^{T-2}} \otimes \boldsymbol{J}_r \otimes \boldsymbol{1}_r) \odot (\boldsymbol{1}_{r^{T-1}} \otimes \boldsymbol{J}_r), \end{aligned}$$

where $\mathbf{1}_s$ is the $s \times 1$ vector of one element, $\mathbf{J}_r^l = (1^l, \dots, r^l)^\top$, " \otimes " denotes the Kronecker product, and " \odot " denotes Hadamard product. Note that $\mathbf{X}_T \mathbf{1}_{d_2} = \mathbf{1}_{r^T}$ holds, and the $r^T \times L$ matrix \mathbf{X} has full column rank where $L = d_2 + L^{(r,T)}$ and $d_2 = (T^2 + 3T - 6)/2$.

We also denote the linear space spanned by the columns of the matrix \boldsymbol{X} by $S(\boldsymbol{X})$ with dimension L. $S(\boldsymbol{X})$ is subspace of \mathbb{R}^{r^T} . Let \boldsymbol{U} be an $r^T \times d_1$, where $d_1 = r^T - L$, full column rank matrix such that the linear space is spanned by the column of \boldsymbol{U} . Namely, $S(\boldsymbol{U})$ is the orthogonal complement of the space $S(\boldsymbol{X})$. Thus, $\boldsymbol{U}^{\top}\boldsymbol{X} = \boldsymbol{O}_{d_1,L}$ where $\boldsymbol{O}_{d_1,L}$ is the $d_1 \times L$ zero matrix.

Let $h_1(\pi)$ and $h_2(\pi)$ be a vector of functions defined by $h_1(\pi) = U^{\top} F(\pi/\pi^S)$ and $h_2(\pi) = M\pi$ with the $d_2 \times r^T$ matrix $M = (x_1, \dots, x_{d_2})^{\top}$, respectively. Note that M^{\top} belongs to the space S(X), namely, $S(M^{\top}) \subset S(X)$. From Corollary 8, the S^T model is equivalent to the hypothesis $h_3(\pi) = (h_1^{\top}(\pi), h_2^{\top}(\pi))^{\top} = \mathbf{0}_{d_3}$, where $\mathbf{0}_s$ is the $s \times 1$ zero vector and $d_3 = d_1 + d_2$, because the GLS^T[f] model is equivalent to the hypothesis $h_1(\pi) = \mathbf{0}_{d_1}$ and the MVC^T model is equivalent to the hypothesis $h_2(\pi) = \mathbf{0}_{d_2}$.

Let $H_s(\pi)$ for s = 1, 2, 3 denote the $d_s \times r^T$ matrix of partial derivatives of $h_s(\pi)$ with respect to π , i.e., $H_s(\pi) = \partial h_s(\pi) / \partial \pi^\top$. Also, let $\Sigma(\pi) = \text{diag}(\pi) - \pi \pi^\top$, where $\text{diag}(\pi)$ denotes a diagonal matrix with the *i*th component of π as the *i*th diagonal component, and p denotes π with π_{λ_i} replaced by p_{λ_i} , where $p_{\lambda_i} = n_{\lambda_i}/n$ with $n = \sum_{\lambda} n_{\lambda}$. Since $\sqrt{n}(p - \pi)$ has an asymptotically multivariate normal distribution with mean $\mathbf{0}_{r^T}$ and covariance matrix $\Sigma(\pi)$, using the delta method, $\sqrt{n}(h_3(p) - h_3(\pi))$ has asymptotically a multivariate normal distribution with mean $\mathbf{0}_{r^T}$ and covariance matrix

$$oldsymbol{H}_3(oldsymbol{\pi}) oldsymbol{\Sigma}(oldsymbol{\pi}) oldsymbol{H}_3^ op (oldsymbol{\pi}) oldsymbol{\Sigma}(oldsymbol{\pi}) oldsymbol{H}_3^ op (oldsymbol{\pi}) oldsymbol{\Sigma}(oldsymbol{\pi}) oldsymbol{H}_1^ op (oldsymbol{\pi}) oldsymbol{\Sigma}(oldsymbol{\pi}) oldsymbol{H}_1^ op (oldsymbol{\pi}) oldsymbol{\Sigma}(oldsymbol{\pi}) oldsymbol{H}_2^ op (oldsymbol{\pi}) oldsymbol{\pi}) oldsymbol{H}_2^ op (oldsymbol{\pi}) oldsymbol{H}_2^ op (oldsymbol{\pi}) oldsymbol{\pi}) oldsymbol{\pi}_2^ op (oldsymbol{\pi}) oldsymbol{\pi}_2^ op (oldsymbol{\pi}) oldsymbol{\pi}) oldsymbol{\pi}_2^ op (oldsymbol{\pi}) oldsymbol{\pi}) oldsymbol{\pi} oldsymbol{\pi}) oldsymbol{\pi}_2^ op (oldsymbol{\pi}) oldsymbol{\pi}) oldsymbol{\pi} oldsymbol{\pi} oldsymbol{\pi}) oldsymbol{\pi} oldsymbol{\pi} oldsymbol{\pi} oldsymbol{\pi} oldsymbol{\pi} oldsymbol{\pi}) oldsymbol{\pi} oldsymbol{\pi} oldsymbol{\pi} oldsymbol{\pi} oldsymbol{\pi}) oldsymbol{\pi} oldsymbol{\pi} oldsymbol{\pi} oldsymbol{$$

Then,

$$oldsymbol{H}_1(oldsymbol{\pi}) = oldsymbol{U}^ op rac{\partial}{\partial oldsymbol{\pi}^ op} F\left(rac{oldsymbol{\pi}}{oldsymbol{\pi}^S}
ight),$$

and

$$H_2(\pi) = M$$

Thus, under the hypothesis $h_3(\pi) = \mathbf{0}_{d_3}$, since $H_1(\pi)\Sigma(\pi)H_2^{\top}(\pi) = O_{d_1,d_2}$ (*), we obtain that

$$W_3 = W_1 + W_2$$

where

$$W_s = n \boldsymbol{h}_s^{\top}(\boldsymbol{p}) \left(\boldsymbol{H}_s(\boldsymbol{p}) \boldsymbol{\Sigma}(\boldsymbol{p}) \boldsymbol{H}_s^{\top}(\boldsymbol{p}) \right)^{-1} \boldsymbol{h}_s(\boldsymbol{p}).$$

We can show (i) W_1 is Wald statistic for the $\text{GLS}^T[f]$ model, (ii) W_2 is that for the MVC^T model, and (iii) W_3 is that for the S^T model.

Thus, we can lead to the following theorem when $T \geq 3$.

Theorem 4. When the S^T model holds, the following equivalence holds:

$$W(\mathbf{S}^T) = W(\mathbf{GLS}^T[f]) + W(\mathbf{MVC}^T),$$

where W(M) denotes the Wald statistic for model M.

As the Wald statistic W_s asymptotically follows a chi-squared distribution with d_s df for s = 1, 2, 3, and given the asymptotic equivalence of the Wald statistic and the likelihood ratio statistic (see [17]), we obtain the following theorem.

Theorem 5. Under the S^T model, the following asymptotic equivalence holds:

$$G^2(\mathbf{S}^T) \simeq G^2(\mathbf{GLS}^T[f]) + G^2(\mathbf{MVC}^T).$$

Theorem 5 extends the decomposition result of Theorem 4 to the likelihood ratio statistics, which are more commonly used in practice.

7 Concluding remarks

In this paper, we introduced the $\operatorname{GLS}^{T}[f]$ model, a novel and flexible approach to modeling multivariate categorical data that incorporates the *f*-divergence to capture complex dependence structures. The key advantage of the $\operatorname{GLS}^{T}[f]$ model is that it is the closest model to the S^{T} model in terms of the *f*-divergence, under certain conditions on the marginal probabilities and symmetric cell probabilities. By applying a function f(x) that satisfies the conditions of the *f*-divergence, the $\operatorname{GLS}^{T}[f]$ model provides a general framework for constructing new models tailored to the specific characteristics of the data and the research objectives.

One of the most significant features of the $\operatorname{GLS}^T[f]$ model is its interpretability. As demonstrated in Section 4.2, the GLS^T model, which is a special case of the $\operatorname{GLS}^T[f]$ model when $f(x) = x \log x$ (x > 0), offers valuable insights into the dependence structures present in the data through its parameters. This interpretability is crucial for understanding the complex relationships among variables in multivariate categorical data and making substantive conclusions based on the model results.

Moreover, the $\operatorname{GLS}^T[f]$ model provides a unified approach to modeling multivariate categorical data, with the GLS^T model as a special case when the Kullback-Leibler divergence is chosen. By choosing appropriate divergence functions, researchers can construct models that are best suited for their specific data and research questions. This flexibility enables a more comprehensive and adaptive analysis of multivariate categorical data, potentially uncovering new insights and patterns that may not be captured by existing models.

The paper also presents several theoretical results, including the decomposition of the S^T model into the $GLS^T[f]$ model and other models, as well as the asymptotic equivalence of the likelihood ratio statistics for these models. These results provide a deeper understanding of the relationships among various models and their goodness-of-fit statistics, facilitating model selection and assessment.

In conclusion, the $\operatorname{GLS}^T[f]$ model introduced in this paper offers a powerful, flexible, and interpretable approach to modeling multivariate categorical data. Its ability to capture complex dependence structures and provide a general framework for constructing new models makes it a valuable tool for researchers across various fields. As future work, we plan to further investigate the properties of the $\operatorname{GLS}^T[f]$ model and extend the model to handle sparse contingency tables. We leave detailed investigations of these for future work.

Appendix

Appendix A Notation

Table 2 provides a summary of the notation used throughout the paper.

Table 2: Notation reference		
Symbol	Definition	
V	The indices set, usually $\{1, \ldots, T\}$	
\mathcal{I}	Levels of any variable, by default $\{1, \ldots, r\}$	
i	(i_1,\ldots,i_T) with $i_j \in \mathcal{I}$, generic notation to denote a cell	
\mathcal{I}_V	$\prod_{i \in V} \mathcal{I}$, the collection of all cells	
X	(X_1, \ldots, X_T) , collection of T variables with $X_j \in \mathcal{I}$	
π	Joint p.m.f of X , $\pi_i = \pi_{(i_1,\ldots,i_T)} = \Pr(X_1 = i_1,\ldots,X_T = i_T)$ for $i \in \mathcal{I}_V$	
\mathcal{P}_V	The subset of \mathcal{I}_V that imposes restrictions on the order of its elements.	
$\mathcal{C}_V^{(2)}$	The subset of $V \times V$, the collection of all index of γ .	
\preceq	The ordinal relation defined on finite vector space \mathbb{N}^n .	
ι	$\{\iota_1, \ldots, \iota_{ \mathcal{I}_V }\}$ is elements of \mathcal{I}_V ordered in ascending order.	
κ	$\{\kappa_1, \ldots, \kappa_{ \mathcal{P}_V }\}$ is elements of \mathcal{P}_V ordered in ascending order.	
λ	$\{\boldsymbol{\lambda}_1, \dots, \boldsymbol{\lambda}_{ \mathcal{C}_{v_v}^{(2)} }\}$ is elements of $\mathcal{C}_V^{(2)}$ ordered in ascending order.	

Appendix B Proofs

Proof of Theorem 2.

If $f(u) = u \log u(u > 0)$, then $\operatorname{GLS}^{T}[f]$ model (5) is represented to

$$\pi_{oldsymbol{i}} = \pi_{oldsymbol{i}}^S rac{|D(oldsymbol{i})|a_{oldsymbol{i}}}{\sum_{oldsymbol{j} \in D(oldsymbol{i})}}, \quad oldsymbol{i} \in \mathcal{I}_V,$$

with

$$a_{i} = \exp\left(\sum_{s=1}^{T} \alpha_{s} i_{s} + \sum_{s=1}^{T} \beta_{s} i_{s}^{2} + \sum_{s=1}^{T-1} \sum_{t=s+1}^{T} \gamma_{st} i_{s} i_{t}\right).$$

Now, this model can be written as follows.

$$\frac{\pi_{\boldsymbol{i}}}{\sum\limits_{\boldsymbol{j}\in D(\boldsymbol{i})}\pi_{\boldsymbol{j}}} = \frac{a_{\boldsymbol{i}}}{\sum\limits_{\boldsymbol{j}\in D(\boldsymbol{i})}a_{\boldsymbol{j}}}, \quad \boldsymbol{i}\in\mathcal{I}_{V}$$

This is an equivalent expression for the GLS^T model (2).

Proof of Theorem 3.

If the S^T model holds, then the $\text{GLS}^T[f]$, ME^T , VE^T and CE^T models hold. Assuming that these models hold, then we show that the S^T model holds. Let $\{\hat{\pi}_i\}$ denote the cell probabilities satisfying the structure of $\text{GLS}^T[f]$, ME^T , VE^T and CE^T simultaneously. Since $\{\hat{\pi}_i\}$ satisfy the $\text{GLS}^T[f]$ model,

$$F\left(\frac{\hat{\pi}_{\boldsymbol{i}}}{\hat{\pi}_{\boldsymbol{i}}^{S}}\right) = \sum_{s=1}^{T} \alpha_{s} i_{s} + \sum_{s=1}^{T} \beta_{s} i_{s}^{2} + \sum_{s< t} \gamma_{st} i_{s} i_{t} + \psi_{\boldsymbol{i}}, \quad \boldsymbol{i} \in \mathcal{I}_{V},$$

where $\psi_{i} = \psi_{j}, j \in D(i)$ and (1). Then for $i \in \mathcal{I}_{V}; j \in D(i)$,

$$F\left(\frac{\hat{\pi}_{i}}{\hat{\pi}_{i}^{S}}\right) - F\left(\frac{\hat{\pi}_{j}}{\hat{\pi}_{j}^{S}}\right) = \sum_{s=1}^{T} \alpha_{s}(i_{s} - j_{s}) + \sum_{s=1}^{T} \beta_{s}(i_{s}^{2} - j_{s}^{2}) + \sum_{s(B.1)$$

The sum of right side of equation (B.1) multiplied by $\hat{\pi}_i$ gives

$$\sum_{s=1}^{T} \sum_{\boldsymbol{i} \in \mathcal{I}_V} \alpha_s (i_s - j_s) \hat{\pi}_{\boldsymbol{i}} = 0,$$

because $\hat{\pi}_i$ satisfy the structure of ME^T,

$$\sum_{s=1}^{T} \sum_{i \in \mathcal{I}_V} \beta_s (i_s^2 - j_s^2) \hat{\pi}_i = 0,$$

because $\hat{\pi}_i$ satisfy the structure of ME^T and VE^T, and

$$\sum_{s < t} \sum_{i \in \mathcal{I}_V} \gamma_s (i_s i_t - j_s j_t) \hat{\pi}_i = 0,$$

because $\hat{\pi}_i$ satisfy the structure of ME^T, VE^T and CE^T. Therefore,

$$\sum_{\boldsymbol{i}\in\mathcal{I}_V}\hat{\pi}_{\boldsymbol{i}}\left(F\left(\frac{\hat{\pi}_{\boldsymbol{i}}}{\hat{\pi}_{\boldsymbol{i}}^S}\right)-F\left(\frac{\hat{\pi}_{\boldsymbol{j}}}{\hat{\pi}_{\boldsymbol{j}}^S}\right)\right)=0.$$

Here, we define the following set

$$D_n(\mathbf{i}) = D_n(i_1, \dots, i_T)$$

= { $\mathbf{j} = (j_1, \dots, j_T) \in \mathcal{I}_V \mid \mathbf{j}$ is any permutation of the *n* elements of $\mathbf{i} \in \mathcal{I}_V$ },

for $n \in V$, and function

$$G(\boldsymbol{i}, \boldsymbol{j}) = F\left(\frac{\hat{\pi}_{\boldsymbol{i}}}{\hat{\pi}_{\boldsymbol{i}}^S}\right) - F\left(\frac{\hat{\pi}_{\boldsymbol{j}}}{\hat{\pi}_{\boldsymbol{j}}^S}\right).$$

The function F being monotonically increasing, G(i, j) = 0 is satisfied only when $\hat{\pi}_i = \hat{\pi}_j$. Then, we will prove that the following proposition holds

$$\forall n \in V, \ \forall \boldsymbol{j} \in D_n(\boldsymbol{i}), \ \sum_{\boldsymbol{i} \in \mathcal{I}_V} \hat{\pi}_{\boldsymbol{i}} G(\boldsymbol{i}, \boldsymbol{j}) = 0 \ \Rightarrow \ \hat{\pi}_{\boldsymbol{i}} = \hat{\pi}_{\boldsymbol{j}}.$$
(B.2)

Firstly, we consider $j \in D_2(i)$, without loss of generality, permuting i_1 and i_2 in i. Since the following equivalence holds

$$\sum_{\boldsymbol{i}\in\mathcal{I}_{V}}\hat{\pi}_{\boldsymbol{i}}G(\boldsymbol{i},\boldsymbol{j})=0\Leftrightarrow\sum_{\substack{\boldsymbol{i}\in\mathcal{I}_{V}\\i_{1}< i_{2}}}(\hat{\pi}_{\boldsymbol{i}}-\hat{\pi}_{\boldsymbol{j}})G(\boldsymbol{i},\boldsymbol{j})=0,$$

we can easily obtain $\hat{\pi}_i = \hat{\pi}_j$. Thus, when n = 2, the proposition (B.2) holds. Then, assuming that proposition (B.2) holds for the case of n = k - 1, we will consider the case of n = k. We consider $\mathbf{j} \in D_k(\mathbf{i})$, without loss of generality, that is permutation of (i_1, \ldots, i_k) in \mathbf{i} . If $j_k = i_k$, we can consider $\mathbf{j} \in D_{k-1}(\mathbf{i})$, and, from assumption, (B.2) holds. If $j_k \neq i_k$, we can derive the following proposition holds from assumption, with $\mathbf{k} \in D_{k-1}(\mathbf{i})$ that is permutation of i_k and j_k in \mathbf{j}

$$\sum_{i \in \mathcal{I}_{V}} \hat{\pi}_{i} G(i, j) = 0 \Leftrightarrow \sum_{i \in \mathcal{I}_{V}} \hat{\pi}_{i} G(i, k) + \sum_{i \in \mathcal{I}_{V}} \hat{\pi}_{i} G(k, j) = 0$$
$$\Rightarrow \hat{\pi}_{i} = \hat{\pi}_{k} \wedge \sum_{k \in \mathcal{I}_{V}} \hat{\pi}_{k} G(k, j) = 0$$
$$\Rightarrow \hat{\pi}_{i} = \hat{\pi}_{k} = \hat{\pi}_{j}.$$

Therefore, the proposition (B.2) holds, namely, $\hat{\pi}_i = \hat{\pi}_j$ for any permutation of i, and $\{\hat{\pi}_i\}$ satisfy the structure of symmetry.

Proof of Equation (*).

Let us denote the matrix F and their elements F_{ij} as

$$\boldsymbol{F} = \frac{\partial}{\partial \boldsymbol{\pi}^{\top}} F\left(\frac{\boldsymbol{\pi}}{\boldsymbol{\pi}^{S}}\right),$$

and

$$F_{ij} = \begin{cases} \frac{1}{\pi_{\boldsymbol{\iota}_i}^S} \left(1 - \frac{\pi_{\boldsymbol{\iota}_i}}{|D(\boldsymbol{\iota}_i)|\pi_{\boldsymbol{\iota}_i}^S} \right) f''\left(\frac{\pi_{\boldsymbol{\iota}_i}}{\pi_{\boldsymbol{\iota}_i}^S}\right) & j = i, \\ -\frac{\pi_{\boldsymbol{\iota}_i}}{|D(\boldsymbol{\iota}_i)|(\pi_{\boldsymbol{\iota}_i}^S)^2} f''\left(\frac{\pi_{\boldsymbol{\iota}_i}}{\pi_{\boldsymbol{\iota}_i}^S}\right) & j \neq i \land \boldsymbol{\iota}_j \in D(\boldsymbol{\iota}_i), \\ 0 & otherwise. \end{cases}$$

Thus, we obtain

$$\boldsymbol{H}_1(\boldsymbol{\pi})\boldsymbol{\pi} = \boldsymbol{0}_{d_1}$$

and, under the hypothesis $h_3(\pi) = \mathbf{0}_{d_3}$, i.e., under the S^T model, we see

$$\boldsymbol{H}_{1}(\boldsymbol{\pi}) \operatorname{diag}(\boldsymbol{\pi}) = c \boldsymbol{U}^{\top} (\boldsymbol{I} - \boldsymbol{J}),$$

where c = f''(1) and the elements of J, J_{ij} , as

$$J_{ij} = \begin{cases} \frac{1}{|D(\boldsymbol{\iota}_j)|} & \boldsymbol{\iota}_j \in D(\boldsymbol{\iota}_i), \\ 0 & otherwise. \end{cases}$$

It is easily verified that J belong to the space S(X), that is, $S(J) \subset S(X)$. Therefore, noting that $U^{\top}J = O_{d_1,d_2}$ and $MU = O_{d_2,d_1}$, we obtained that under the S^T model

$$oldsymbol{H}_1(oldsymbol{\pi}) oldsymbol{\Sigma}(oldsymbol{\pi}) oldsymbol{H}_2^+(oldsymbol{\pi}) = oldsymbol{O}_{d_1,d_2}.$$

Acknowledgement

This work was supported by JSPS KAKENHI Grant Number JP20K03756. The authors also thank Dr. Kengo Fujisawa for his valuable advice and feedback throughout this research.

References

- Alan Agresti. A simple diagonals-parameter symmetry and quasi-symmetry model. Statistics & probability letters, 1(6):313–316, 1983.
- [2] Alan Agresti. Categorical Data Analysis. John Wiley & Sons, 2012.
- [3] John Aitchison. Large-sample restricted parametric tests. Journal of the Royal Statistical Society. Series B (Methodological), 24(1):234–250, 1962.
- [4] S. M. Ali and S. D. Silvey. A general class of coefficients of divergence of one distribution from another. Journal of the Royal Statistical Society. Series B (Methodological), 28(1):131–142, 1966.

- [5] I Csiszár. Eine informationstheoretische ungleichung und ihre anwendung auf beweis der ergodizitaet von markoffschen ketten. Magyer Tud. Akad. Mat. Kutato Int. Koezl., 8:85–108, 1964.
- [6] I. Csiszár and P. C. Shields. Information theory and statistics: A tutorial. Foundations and Trends in Communications and Information Theory, 1(4):417–528, 2004.
- [7] J. N. Darroch and S. D. Silvey. On testing more than one hypothesis. The Annals of Mathematical Statistics, 34(2):555–567, 1963.
- [8] Leo A. Goodman. How to ransack social mobility tables and other kinds of cross-classification tables. American Journal of Sociology, 75(1):1–40, 1969.
- [9] Leo A. Goodman. Some multiplicative models for the analysis of cross classified data. In Proceedings of the Sixth Berkeley Symposium on Mathematical Statistics and Probability, Volume 1: Theory of Statistics, volume 6, pages 649–697. University of California Press, 1972.
- [10] Leo A. Goodman. Multiplicative models for square contingency tables with ordered categories. *Biometrika*, 66(3):413–418, 1979.
- [11] Shelby J. Haberman. Log-linear models for frequency tables with ordered classifications. *Biometrics*, 30(4):589–600, 1974.
- [12] Maria Kateri and Alan Agresti. A class of ordinal quasi-symmetry models for square contingency tables. *Statistics & Probability Letters*, 77(6):598–603, 2007.
- [13] Maria Kateri and Takis Papaioannou. Asymmetry models for contingency tables. Journal of the American Statistical Association, 92(439):1124–1131, 1997.
- [14] S. Kullback and R. A. Leibler. On information and sufficiency. The Annals of Mathematical Statistics, 22(1):79–86, 1951.
- [15] Hisaya Okahara and Kouji Tahata. A generalized ordinal quasi-symmetry model and its separability for analyzing multi-way tables, 2024. arXiv:2405.04193.
- [16] Karl Pearson. X. On the criterion that a given system of deviations from the probable in the case of a correlated system of variables is such that it can be reasonably supposed to have arisen from random sampling. The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science, 50(302):157–175, 1900.
- [17] C.R Rao. Large Sample Theory and Methods. In *Linear Statistical Inference and Its Applications*, chapter 6, pages 382–443. John Wiley & Sons, Ltd, 1973.
- [18] Timothy R. C. Read and Noel A. C. Cressie. Goodness-of-Fit Statistics for Discrete Multivariate Data. Springer Science & Business Media, 2012.
- [19] K. Tahata, H. Yamamoto, and S. Tomizawa. Linear ordinal quasi-symmetry model and decomposition of symmetry for multi-way tables. *Mathematical Methods of Statistics*, 20(2):158–164, 2011.
- [20] Kouji Tahata. Separation of symmetry for square tables with ordinal categorical data. Japanese Journal of Statistics and Data Science, 3(2):469–484, 2020.
- [21] Kouji Tahata, Hideharu Yamamoto, and Sadao Tomizawa. Orthogonality of decompositions of symmetry into extended symmetry and marginal equimoment for multi-way tables with ordered categories. Austrian Journal of Statistics, 37(2):185–194, 2008.
- [22] Sadao Tomizawa. An extended linear diagonals-parameter symmetry model for square contingency tables with ordered categories. *Metron*, 49:401–409, 1991.

- [23] Sadao Tomizawa and Kouji Tahata. The analysis of symmetry and asymmetry : Orthogonality of decomposition of symmetry into quasi-symmetry and marginal symmetry for multi-way tables. Journal de la Société française de statistique & Revue de statistique appliquée, 148(3):3–36, 2007.
- [24] Hideharu Yamamoto, Toshiya Iwashita, and Sadao Tomizawa. Decomposition of symmetry into ordinal quasi-symmetry and marginal equimoment for multi-way tables. Austrian Journal of Statistics, 36(4):291–306, 2007.