Three sided com plex adaptative system s

R D 'hulst¹ and G J.R odgers

Departm ent of M athem atics and Statistics, B runel U niversity, Uxbridge, Middlesex, UB8 3PH, UK.

Abstract

W e introduce two three sided adaptative system s as toy m odels to m in ic the exchange of com m odities between buyers and sellers. These m odels are simple extensions of the minority game, exhibiting similar behaviour as well as some new features. The main di erence between our two models is that in the second, one choice appears as a comprom ise between the two other sides. Both models are investigated num erically and com pared with the originalm inority gam e.

Keywords: m inority gam e, econom y, optim ization, probability PACS: 89.90.+ n, 02.50 Le, 64.60 Cn, 87.10.+ e

1 Introduction

In this paper, we consider extensions of the bar-attendance problem model, the agents have to choose beintroduced by Arthur [1] and sim pli-tween three identical sides at each

and Zhang [2]. In its simplest form, ing with each other, agents on one the minority gam e minics the inter-side buying from the second side to naldynamic of the exchange of one sell to the third. This model mincommodity. Agents are allowed to ics the cyclic trading of goods. If we buy or sell this commodity at each group any two sides together and contime step. No attempt is made to sider the trading between this im agmodel any external factors that in- inary group and the third side, the troduce symmetric and asymmetric game with an uneven distribution of

three sided gam es as extensions of the m inority gam e.

In the symmetric three sided ed into a m inority gam e by Challet tim e step. These three sides are traduence the market. Here, we in-model reduces to a kind of minority

¹ corresponding author: Rene DH ulst@ bruneLac.uk

gam e is very strong.

model, the agents can buy or sell a m time steps constitutes the history commodity at each time step, but they can also be inactive, that is, they are allowed to miss a turn. In contrast to the symmetric model, the three choices are not equivalent, as being inactive appears as a com prom ise between buying and selling. This model can be thought of as an open minority game in the sense that the agents buying and selling are playing a m inority gam e with a variable number of agents at each turn.

2, the minority game In Sec. is brie y recalled and the two new three sided models are described in In Sec. 3, the symmetric detail. three sided model is num erically investigated, while in Sec. 4, the asym metric three sided model is investigated. Sec. 5 presents a com parison between the minority game and the two three sided models, as well as our conclusions.

2 The models

In the minority game, an odd number N of agents have to choose

the agents. Hence, the connection be- between two sides, 1 or 2, at each time tween this model and the minority step. An agent wins if he chooses the minority side. The record of which In the asymmetric three sided side was the winning side for the last of the system . The agents analyze the history of the system in order to make their next decision.

> In the symmetric three sided model, a number of agents N have to choose between three sides, 1, 2 or 3, at each time step. N is not a multiple of 3. The agents on side 1 buy from side 2 to sell to side 3, the agents on side 2 buy from side 3 to sell to side 1 and the agents on side 3 buy from side 1 to sell to side 2. This cyclic trading pattern is shown in Fig. 1. It is assumed that the protor loss at a side is relected in the di erence between the number of agents they are selling to and the number of agents they are buying from . For instance, Nз N_2 is a measure of the prot of side 1. Agents choosing the side with the highest pro twin and are rewarded with a point. A gents choosing the side with the lowest pro t bse and consequently bse a point. A gents choosing the side with the interm ediate pro t neither lose nor gain a point. Agents strive to maxim ize their total num ber of points.




Figure 1: Schem atic representation of the symmetric three sided model. The arrows indicate the direction of the exchange. N_i is the number of agents choosing side i.

a point. The agents choosing the number of points. larger group am ong buyers and sellers

In the asymmetric three sided lose and they lose a point. The points model, a number of agents N also of the inactive agents don't change. have to choose between three sides, If there is the same number of buy-1, 2 or 3, at each time step. 1 corre- ers and sellers, the points of all the sponds to selling, 2 to doing nothing agents remain unchanged. However, and 3 to buying. The agents buy- the inactive agents are recorded as ing or selling are said to be active, winners in the history of the system, while the agents doing nothing are on the grounds that they achieved the said to be inactive. The agents choos- same result as the buyers and sellers, ing the smaller group among buyers but without taking any risk. Again, and sellers w in and are rewarded with agents strive to maxim ize their total

In each model, the record of which agent makes a decision for the next side was the winning side for the last time step according to the history of m time steps constitutes the history the system. To be able to play, an of the system. For a given m, there agent must have a strategy that alare 3^m di erent histories. The 9 dif-lows him to make a decision for any ferent histories for m = 2 are listed of the 3^m di erent histories. The secin the rst column of table 1. Every ond and third columns of table 1 list

H istory		0	
II ISCOLÀ			
(1,1)	3	2	
(1,2)	1	3	
(1,3)	2	2	
(2,1)	2	1	
(2 , 2)	3	3	
(2 , 3)	1	1	
(3,1)	3	1	
(3 , 2)	2	2	
(3,3)	3	1	

two possible sets of decisions, and ⁰, that we will call strategies.

Each agent has at his disposal a xed set of s strategies chosen at random, multiple choices of the same strategy being allowed. At any one m om ent in time, the agent only uses one of these strategies to make a decision. To allow an agent to decide which strategy to use, every strategy is awarded points, which are called virtual points. The virtual points of and 0 component by component, we a strategy are the points the agent see that for some histories they can egy predicting the third side does not ences between the strategies can give

Table 1: The rst column lists all the possible histories of the system for the last 2 time steps (m = 2). A strategy is a set of decisions for all the di erent possible histories. Two example strategies and are shown in the second and third colum ns.

gain or lose any points. In the asymmetric model, in the case of an equal num ber of buyers and sellers, the virtualpoints of all strategies rem ain unchanged. An agent always plays with the strategy with the highest number of virtual points. W hen more than one strategy has the highest number of virtual points, one of them is chosen at random .

If we compare two strategies thinks he could have earned had he make the same prediction and for othplayed with this strategy. Hence, ers they can make di erent predicthe virtual points are rewarded us- tions. In the example in table 1, ing the same scheme as the points the decisions dier when the history given to the agents, the prediction of is (1,1), (1,2), (2,1), (3,1) and (3,3). a strategy being compared to the ac- To consider this feature, we have to tual decisions. A strategy predicting distinguish between the symmetric the winning side is awarded a virtual model and the asymmetric one. For point, a strategy predicting the losing the form er, the three sides are equivside loses a virtual point and a strat- alent and only the number of di er-

two strategies in the strategy space. The symmetric distance between two For the latter, there is a qualitative strategies corresponds to the probadi erence between the three sides. bility that these two strategies will pear in the de nition of the di erence that all the histories are equally likely between strategies.

sided model. As the three sides are the minority game [3]. Two strategies equivalent, a geom etrical representa- at $d_s = 0$ are correlated, two stratetion should put them at the same dis- gies at $d_s = 2=3$ are uncorrelated and tance from one another. A conve-two strategies at $d_s = 1$ are anticornient measure of the di erences be- related. tween two strategies and 0 is

$$d_{s} = \frac{1}{3^{m}} \frac{\hat{X}^{m}}{\sum_{i=1}^{m}} (1 - \sum_{i=1}^{m} \frac{1}{2})$$
 (1)

where (0) = 1, and (x) = 0 otherwise. d_s is de ned as the distance between strategies in the symmetric model. This de nition takes into account the geom etrical structure of the is a measure of the di erence between strategy space, including the equiva- the two strategies. da is de ned as lence between the three sides. In the the distance between strategies in the example of table 1, $d_s = 5=9$. By def- asymmetric model. This de nition inition, the symmetric distance is a takes into account the fact that buynumber ranging from 0 to 1.

distance d_s is de ned as a sum of 3th trary way. In the example of table 1, these terms is equal to 0 with prob- metric distance is a number ranging ability 1/3 or equal to 1 with proba-from 0 to 1. bility 2/3. The average distance between two strategies is $\overline{d}_s = 2=3$, metric distance d_a is de ned as a sum while the variance of the symmetric of 3^m term swe label $d_a^{(i)}$'s. When the

a measure of the di erence between distance distribution is $\frac{2}{s} = 2 = 3^{m+2}$. This qualitative di erence should ap- give di erent predictions, assuming to occur. The symmetric distance Consider rst the symmetric three corresponds to the distance de ned in

> In the asymmetric three sided minority game, selling is just the opposite decision to buying while doing nothing is a comprom ise. Consequently, the norm alized asymmetric distance between strategies and 0

$$d_{a} = \frac{1}{3^{m}} \frac{\dot{X}^{m}}{\sum_{i=1}^{m} \frac{j_{i}}{2}} \frac{j_{i}}{2}$$
(2)

ing is more di erent from selling than As Eq. (1) shows, the symmetric it is from being inactive in an arbiterms, which we label $d_s^{(i)}$'s. Each of $d_a = 4=9$. By de nition, the asym-

As shown by Eq. (2), the asym –

component of a strategy is equal to 2, 3this component can never give a d_a⁽ⁱ⁾ equal to 1. In other words, the inactive side has no side at distance 1 from itself. Considering all the possibilities, the probability to $nd a d^{(i)}$ of 0 is 1/3, of 0.5 is 4/9 and of 1 is 2/9. The average asymmetric distance between strategies is $\overline{d}_a = 4=9$, while the variance of the asymmetric distance distribution is equal to $a^2 = 11=3^{m+4}$. The interpretation of this asymmetric distance is ambiguous. In fact, the opposite to selling is buying, but the opposite to being inactive is being inactive. Hence, d_a is not a measure of the probability that two strategies would give opposite decisions. Two correlated strategies are at $d_s = 0$ from each other, two uncorrelated strategies are at $d_a = 1=2$ from each other, but two anticorrelated strategies can be at $d_a = 0$ or $d_a = 1$ from each other.

Numerical results for the symmetric m odel

In this section, we report on numerical investigations of the properties of the symmetric three sided model, interpreting the results using the symmetric distance.

Fig. 2 presents a typical result for the time evolution of the attendance at one side. The simulation is for N = 101 agents with s = 2 strategies each and a memory of m = 3. The result for the attendance at one side is very similar to the results of the m inority gam e, the m ean attendance being shifted to N = 3 instead of N = 2. G iven an agent choosing one side, the average distance between the strategy used by this agent and the strategies used by the other agents is d = 2=3. That is, around 2=3 of the agents should choose one of the two other sides. Hence, the average attendance at one side is N = 3.

The variance of the attendance at ing to 2N = 9, the random value, as m the memory m is presented at Fig. shape are obtained for the variance strategies. The result is again very ance of the number of losers. A lso, very high variance for m < 3, a m in- of agents on the more crowded side in un at m = 3 and the variance go- exhibit the same behaviour. Each

one side as a function of the size of goes to in nity. Curves of the same 3 for N = 101 agents with s = 2 of the number of winners or the varisimilar to the minority game, with a the maximum prot or the number




Figure 2: Numerical simulation of the attendance at one side for the symm etric three sided m odel. The choice for the parameters is N = 101, s = 2and m = 3.

of the strategies. Consequently, the the same as the kinetics of a random fact, the variance of the attendance e ect being obtained [4]. scales like $1 = \frac{2}{s}$. On the contrary, for

of these curves has a minimum for m > 3, the space of the strategies is m = 3. For m < 3, the number of very large, so that most of the stratestrategies used at each time step is gies used are uncorrelated. As a rea representative sample of the space sult, the kinetics of the system are variance of the attendance is directly walk. Between these two behaviours, related to the variance of the dis- form around 3, the agents organize tance distribution, $\frac{2}{s} = 2=3^{m+2}$. In them selves better, a crow d-anticrow d

Even for small values of m, the we suppose that 1= $^2_{\rm s}$ gives an estispace of strategies is very large, of size mate of the number of uncorrelated 3^{3^m} . But as in the m inority gam e, not strategies, the m ethod of Johnson et all the strategies are uncorrelated. If al. [4] can be used to nd an ana-




Figure 3: Variance of the number of agents at one side as a function of the size of the memory m () for N = 101 and s = 2 (symmetric model). The analytical results using the calculation of Johnson et al. [4] is also shown (2).

lytical expression for the variance of an independent agent. The analytthe attendance at one side. We fol- ical result obtained by this method lowed the original calculation in [4], is compared in Fig. 3 to the result with a size $a = 3^{m+2}=2$ for the space of the num erical simulations. The of strategies and a variance of 2/9 for curves agree qualitatively.

given to the agents and their strate- complex behaviour. The points given virtual points are steadily decreasing lution of the virtual points can be ap-

Fig. 4 (a) presents a typical re- with time. In contrast, the points sult for the average number of points given to the agents display a more gies. The parameters of the simula- to the agents increase very slowly for tion are N = 101, s = 2 and m = 3. m < 3 and then oscillate around 0 for Note that there are two di erent or di = m > 3. There seem to be no special nate scales. As Fig. 4 (a) shows, the behaviour for m = 3. The time evo-




Figure 4: (a) A verage points earned by the agents and their strategies as a function of the time for N = 101, s = 2 and m = 3. The ordinate on the left refers to the points of the agents (bold line) while the ordinate on the right refers to the points of their strategies (simple line). (b) Prot trate of the strategies as a function of m for N = 101 and s = 2.

proximated by a linear relation with a critical value m_c for the size of the a protrate. We de ne average number of points earned by a strategies is crowded and its geometpresents as a function of the mem - sults. As this structure is encoded in m < 3, the strategies are slowly los-driven by its distance distribution . tual points oscillate around 0. Hence, the strategies used are uncorrelated. the agents seem to be able to choose The kinetics of the system reduce to their strategy e ciently, in the sense agents choosing one of the three sides that the strategies they choose win at random. Hence, there is a transimore often than the average strat- tion from a system driven by its disegy. This behaviour is to be con-tance distribution to a random systrasted with the minority game where tem. the agents are not able to choose a strategy e ciently.

As a summary, the symmetric model is a direct extension of the minority game to three sides. The results found are very similar, with a glassy phase transition [5] when the increased. We num erically identied ric three sided game. In the gures, 1

as the memory. For $m < m_{c}$, the space of strategy at each time step. Fig. 4 (b) rical structure is apparent in the reory m for N = 101 and s = 2. For the distance de nition, the system is ing points, the worst results being ob- For m > m_c, the number of stratetained form = 3. Form > 3, the vir- gies used is not relevant as most of


4 Numerical results for the asym m etric m odel

We investigated num erically the size of the memory of the agents is di erent properties of the asymmetdenotes buying, 2, doing nothing and ing (the lower curve for sm all m val-3, selling.

ferent sides as a function of the size more and more to the agents guessof the memory m is plotted at Fig. 5 for N = 101 agents, playing with s = 2 strategies each. The number m, the number of winners is signifof agents in the winning side is also presented. For small m values, most guess value. Fig. 5 is interesting be-(the two superim posed upper curves). sides is clearly apparent. Just a few of them are doing noth-

As the size of the memory ues). The attendance of the three dif- is increased, the system corresponds ing at random between the three possibilities. Also, for small values of icantly more than 1/3, the random of the agents are buying or selling cause the di erence between the three

Figure 5 R. D'hulst and G. J. Rodgers



F igure 5: Num erical result for the average attendances at the three sides as functions of m for N = 101 and s = 2. The two upper curves (4 and) are for the selling and buying options. The lower curve () is for the inactive agents. The average number of winners is also presented (2) as a function ofm for the same choice of parameters.

tendances at the three sides and the around 2. The variance of the num variance of the number of winners ber of winners has a shape very sim are presented as functions of m for ilar to the one found in the minority N = 101 and s = 2. For m < 6, game. For sm all value of m, the varithe variance of the number of inac- ance diverges like a power law ofm; at tive agents is signicantly higher than m '7, it seems to reach a minimum 2N = 9, the value for agents guessing at and for higher values of m, it goes random. The variances of the num - asymptotically to a value near N = 9. ber of buyers and sellers has a min- However, the existence of a minimum in un at m = 2. The variances of at m = 7 could not be con m = 4 unthe three sides increase to 2N = 9 as m equivocally by the num erical simulaincreases. Hence, there seems to be tions.

In Fig. 6, the variance of the at- an organization of the agents for m




Figure 6 R. D'hulst and G. J. Rodgers

Figure 6: Variances of attendances at the three sides as functions of m for N = 101 and s = 2.4 and, selling and buying options, inactive agents. 2 is the variance of the number of winners as a function of m for the same choice of param eters.

m values, the behaviour of the sys- minority game, these two ratios are tem is directly related to the proper- 4/9 and 1/3 respectively. Hence, the ties of the distance distribution. The transition from a system driven by proportion of people buying or sell- the distance distribution to a system ing is of the same order as the av- of agent quessing at random is seen erage distance, 4/9, while the vari- directly in the attendance of the difance of the num ber of winners scales ferent sides. as $1 = \frac{2}{2}$, the inverse of the variance of the asym m etric distance distribution. These properties were also present in the m inority gam e. In this asym m etric model, the variance of the attendance at one side does not represent the wasted number of points. The wasted number of points is de ned to to win than the inactive side. In fact, mum points that can be earned by the system at each time step and the average points actually earned by the system at each time step. This is why we also have to consider the properties of the number of winners in addition to the properties of the attendances.

For higher values of m, the strateqy space is so large that most of the used strategies are uncorrelated. The system is similar to a system with agents choosing at random from the three sides. In them inority gam e, the relative attendance predicted by the distance distribution is the same as the one predicted by random guesses,

Fig. 5 and 6 show that for small that is 1/2. In the present three sided

Fig. 7 presents the average success rate of one side, that is, the probability that at any one moment in time, one side will win. As expected, the sides corresponding to buying and selling are symmetric and more likely be the di erence between the maxi- there are (N + 1)(N + 2)=2 di erent con gurations for the attendances of the 3 sides. Among these, only (N + 2)=2 m ake the inactive agents winners if N is even, (N + 1) = 2 if N is odd. Hence, if all the situations were equally likely to occur, the inactive agents would win at most about every N + 1 tim e steps. This is the order of the asymptotical value for the success rate of this side. For low values ofm, the success rate of the inactive side is higher than the asymptotical value, in plying that the agents playing are organizing them selves rather well. The transition between organized and non-organized agents is for m = 2 in F ig. 7.




Figure 7: Success rates of the three di erent sides as functions of m for N = 101 and s = 2. The two upper curves (4 and) are for selling and buying while the lower curve () is for the inactive agents.

Fig. 8 con m s the organization are able to choose strategies which of the agents. The pro t rates of the are m ore successful than the average agents and their strategies are shown ones. On the contrary, for m > 5, as functions of m. We de nea pro t they are doing worse than guessing rate as the average number of points at random. The curve of the pro t earned at each time step. For val- rate of the strategies suggests that the ues of m less than m = 5, the agents transition takes place for m = 2.

As a summary, in the asymmetric three sided m inority game, agents and the symmetric three sided m odel, playing with a small memory win a glassy phase transition [5] is found more points on average than agents at a particular value of the memory playing with a bigger memory in a m_c . For $m < m_c$, the geometrical pure population, that is, a population properties of the space of strategies of all the agents with the same mem - is apparent, especially the asymmetric strategies of the space of strategies




Figure 8: A verage pro t rates of the agents and their strategies as functions of m for N = 101 and s = 2.2 is for the prot trate of the agents while is for the pro t rate of their strategies.

try between the three sides. Most of on their pro t. Hence, the agents are tem is driven by the distance distri- are unlikely to win. bution. In contrast to the minority game, this property is seen directly in the number of agents on each side. 5For $m > m_{c}$, the strategies used are uncorrelated and the system is similar to a system of agents guessing at ran- models as extensions of the minority dom. Considering the adaptation of game. In the symmetric three sided the agents, they are unable to realize model, agents are given three equivthat the wiser choice is to decide to alent choices while in the asymmetbe inactive. In fact, more than half ric three sided model, agents have the of the active agents will bee. The opportunity to miss a turn and not agents are fooled because they base play. We have investigated these two their con dence in virtual points, not new models num erically and com -

the agents are playing and the sys- always tempted to play even if they

Conclusions

We introduced two three sided

nority gam e.

tance between the strategies of the of the geometrical structure of the agents. These distances incorporate space of strategies. in their de nitions the geometrical structure of the space of strategies. In vestigate both models analytically. the symmetric model, the geometrical The symmetric model, in particustructure of the space of strategies is lar, should be amenable to analytivery sim ilar to the one in them inority cal treatment, perhaps following the game. The distance gives a measure methods introduced in [6] for the two of the correlation between two strate- sided m inority gam e. gies. Conversely, the distance in the asymmetricmodel has no obvious interpretation.

A transition between a system driven by the distance distribution and a system of agents quessing at random was identi ed num erically in both models. However, in contrast to the m inority gam e, the agents m ake their highest pro t for m small and not at the transition value of m . In the distance driven phase, the agents organize them selves, as in the minority game. In contrast to the minority gam e, how ever, the average pro t rate of the agents is higher than the average pro t rate of the strategies, indicating that the agents are choosing their strategies e ciently. In the symmetric model, the transition is apparent in the variance of the num ber of agents choosing one side while in the asymmetric model the transi-

pared the results with the original mi- tion is seen in the num ber of agents it-

self. This latter property of the asym -In both models, we de ned a dis-metric model is a direct consequence

In the future, we intend to in-

References

- [1] W. B. Arthur, Inductive Reasoning and Bounded Rationality, Am. Econ. Assoc. Papers and Proc. 84 (1994) 406.
- [2] D. Challet and Y.-C. Zhang, Physica A 246 (1997) 407; 256 (1998) 514.
- [3] R. D'hulst and G. J. Rodgers, submitted to Physica A (adaporg/9902001).
- [4] N.F. Johnson, M. Hart and P. M.Hui, cond-m at/9811227.
- [5] R. Savit, R. M anuca and R. Riolo, adap-org/9712006.
- [6] D. Challet and M. Marsili, condm at/9904071.