

Cohomology of coherent sheaves on a proper scheme

GEORGE R. KEMPF
THE JOHNS HOPKINS UNIVERSITY

Let $f : X \rightarrow S$ be a proper morphism where $S = \text{Spec } A$ where A is a noetherian ring. Let $\mathcal{F}$ be a coherent sheaf on X . Then a theorem of Grothendieck says that the cohomology groups $H^i(X, \mathcal{F})$ are A -modules of finite type. We set out to give an algorithm to compute these A -modules when A is sufficiently algorithmic. We found general result about a $\Gamma(X, -)$ -acyclic resolution of $\mathcal{F}$.

We will first define a coherent sheaf $\mathcal{G}$ on X to be primitive if support $(\mathcal{G}) = \bar{x}$ where x is a point of X and the natural mapping $\mathcal{G} \rightarrow i_{x*}(\tilde{\mathcal{G}}_x)$ is injective where $i_x : \text{Spec } \mathcal{O}_{X,x} \rightarrow X$ is the natural morphism. The associated points of $\mathcal{G}$ is denoted by $\text{Ass}(\mathcal{G})$, which is a finite set.

The theorem we prove is the following.

Theorem 1. *There exists an inclusion $\mathcal{G} \xrightarrow{\alpha} \bigoplus_{x \in \text{Ass}(\mathcal{G})} \mathcal{H}_x$ where a) each $\mathcal{H}_x$ is a primitive coherent sheaf with support $\bar{x}$, b) each $\mathcal{H}_x$ is $\Gamma(X, -)$ -acyclic and its group of sections $\Gamma(X, \mathcal{H}_x)$ is an A -module of finite type, and c) α is an isomorphism on an open dense subset of the support of $\mathcal{G}$.*

This theorem gives a method to compute the cohomology of $\mathcal{F}$. One applies the theorem inductively to construct a $\Gamma(X, -)$ -acyclic coherent resolution $0 \rightarrow \mathcal{F} \rightarrow \mathcal{F}^*$ and simply computes $H^i(X, \mathcal{F})$ as the i -homology group of the complex $\Gamma(X, \mathcal{F}^*)$.

Thus Theorem 1 implies

Corollary (Grothendieck). *The cohomology groups $H^i(X, \mathcal{F})$ are A -modules of finite type.*

§1. A lemma.

We will use

Lemma 2. *If $\mathcal{F}$ is a coherent sheaf on X , there exists an embedding $\mathcal{F} \xrightarrow{\beta} \bigoplus_{x \in \text{Ass}(\mathcal{F})} \mathcal{G}_x$ where $\mathcal{G}_x$ is a primitive coherent sheaf with support $\bar{x}$ and β is an isomorphism in an open dense subset of support $(\mathcal{F})$.*

Proof. We have an injection $\mathcal{F} \hookrightarrow \bigoplus_{x \in \text{Ass}(\mathcal{F})} i_{x*}(\tilde{\mathcal{F}}_x)$ because the kernel would have no associated points. Similarly, we have the inclusion $\mathcal{F}_{x1} \xrightarrow{\varprojlim_n} F_x/m_x^n F$ where m_x is the maximal ideal of $\mathcal{O}_{x,x}$. As F is coherent it follows that $\mathcal{F} \hookrightarrow \bigoplus_{x \in \text{Ass}(\mathcal{F})} i_{x*}(\widetilde{F_x/m_x^n F_x})$ where the n_x are sufficiently big. This is our embedding if $\mathcal{G}_x$ is the image of $\mathcal{F}$ in $i_{x*}(\widetilde{F_x/m_x^n F_x})$. QED

By the lemma to prove the theorem we may assume that $\mathcal{G}$ is primitive with support $\bar{x}$ for some x in X .

§2. The proof.

To prove the theorem we will use basically Grothendieck's idea of using Chow coverings. Let Z be a closed subscheme of X such that the support $(Z) = \bar{x}$ and $\mathcal{G}$ admits the structure of a $\mathcal{O}_Z$ -module. We will modify Chow lemma to the non-reduce case.

Let $Z = \cup U_i$ be a finite affine covering of Z by dense open subsets. Let $\bar{U}_i \supset U_i$ be a projective imbedding for each i . Take W to be the schematic closure of $\cap U_i$ embedded in $X \times_S \prod \bar{U}_i$. Then by the usual reasoning the projection $W \rightarrow \prod \bar{U}_i$ is set-theoretically injective proper morphism. Thus W is a projective scheme over S .

Let $\mathcal{M}$ be a very ample sheaf on $\prod \bar{U}_i = K$. Then $\mathcal{N} = \pi_K^* \mathcal{M}$ is very ample on W and relatively ample for the projective morphism $\pi_Z^* : W \rightarrow Z$, which is an isomorphism between open dense subsets.

Now let $\mathcal{L}_x$ be the maximal primitive quotient of $\pi_Z^* \mathcal{G}$ with support W . By projective theory the $\mathcal{K}_x \equiv \mathcal{L}_x \otimes \mathcal{M}^{\otimes n}$ is $\Gamma(W, -)$ -acyclic and π_{Z*} -acyclic for $n \gg 0$. Thus $\pi_{Z*}(\mathcal{K}_*) = \mathcal{H}_x$ is a $\Gamma(X, -)$ -acyclic coherent primitive sheaf, by the degenerate Leray spectral sequence. It has support $\bar{x}$ and its sections are isomorphic to $\Gamma(W, \mathcal{K}_x)$, which is an A -module of finite type.

To get the inclusion $\mathcal{F} \rightarrow \mathcal{H}_x$ just multiply the natural map $\mathcal{F} \rightarrow \pi_{Z*} \mathcal{L}_x$ by a section of $\mathcal{M}^{\otimes n}$ which is general. This proves the theorem.

References

1. A. Grothendieck, *Éléments de Géométrie Algébrique III* Publications Mathématiques No. 111.
2. G. Kempf, Some elementary proofs of basis theorems in the cohomology of quasi-coherent sheaves, *Rocky Mountain Journal of Math.*, vol. 10, 1980, pp. 637-645.

AMS (MOS) SUBJECT CLASSIFICATION NUMBER 14F05.