Com plex dynam ics in a sim ple model of pulsations for Super-A sym ptotic G iant B ranch Stars

Andreea Munteanu^a, Enrique Garc a-Berro^{a,b}, Jordi Jose^{c,b}, & Emilia Petrisor^d

^a Departam ent de F sica Aplicada, Universitat Politecnica de Catalunya, Jordi Girona Salgado s/n, Modul B {5, Campus Nord, 08034 Barcelona, Spain

^b Institut d'Estudis Espacials de Catalunya, Edi ci Nexus, Gran Capita 2-4, 08034 Barœlona, Spain

^c Departament de F sica i Enginyeria Nuclear, Universitat Politecnica de Catalunya, Av. V ctor Balaguer, s/n, 08800, Vilanova i la Geltru (Barcelona), Spain

^d Departam entul de Matematica, Universitatea Politehnica Timisoara, Pta Regina Maria 1, 1900 Timisoara, Romania (April 10, 2024)

W hen interm ediate m ass stars reach their last stages of evolution they show pronounced oscillations. This phenom enon happens when these stars reach the so-called A sym ptotic G iant B ranch (AGB), which is a region of the Hertzsprung-Russell diagram located at about the same region of e ective tem peratures but at larger lum inosities than those of regular giant stars. The period of these oscillations depends on the mass of the star. There is growing evidence that these oscillations are highly correlated with m ass loss and that, as the m ass loss increases, the pulsations becom e m ore chaotic. In this paper we study a simple oscillator which accounts for the observed properties of this kind of stars. This oscillator was rst proposed and studied in [1] and we extend their study to the region of m ore m assive and lum inous stars | the region of Super-AGB stars. The oscillator consists of a periodic nonlinear perturbation of a linear Hamiltonian system. The formalism of dynam ical system s theory has been used to explore the associated Poincare m ap for the range of param eters typical of those stars. We have studied and characterized the dynam ical behaviour of the oscillator as the parameters of the model are varied, leading us to explore a sequence of local and global bifurcations. Am ong these, a tripling bifurcation is remarkable, which allows us to show that the Poincare map is a nontwist area preserving map. M eandering curves, hierarchical-islands traps and sticky orbits also show up. W e discuss the implications of the stickiness phenomenon in the evolution and stability of the Super-AGB stars.

PACS numbers: 97.10.Sj 05.45Pq, 95.10Fh, 82.40Bj 05.45Ac

I. IN TRODUCTION

The study of pulsating stars has attracted much attention from astronom ers. Pulsational instabilities are found in many phases of stellar evolution, and also for a wide range of stellar m asses (see [2] for an excellent review). M oreover, pulsational instabilities provide a unique opportunity to learn about the physics of stars and to derive useful constraints on the stellar physical mechanism s that would not be accessible otherwise. Within the theory of stellar pulsations, there are three basic characteristics of the motions that the associated model may include or not, namely the oscillations can be linear or nonlinear, adiabatic or not, and radial or nonradial [3]. A ctual pulsations of real stars must certainly involve som e degree of nonlinearity [4{6]. In fact, the irregular behaviour observed in many variable stars is, de nitely, the result of those nonlinear e ects [7]. Moreover, the fact that the observed pulsation amplitudes of variable stars of a given type do not show huge variations from star to star suggests the existence of a (nonlinear) lim it-cycle type of behaviour [8]. However the full set of nonlinear equations is so complicated that there are no realistic stellar models for which analytic solutions exist and thus the

investigations of nonlinear pulsations rely on numerical analysis. A coordingly, most recent theoretical studies of stellar pulsations proceed either through pure numerical hydrodynam ical codes $[9\{13\} \text{ or, conversely, they adopt a}$ set of sim plifying assumptions in order to be able to deal with a extraordinarily complex problem. Most of these simple models of stellar pulsation are based on a onezone type of model which may be visualized as a single, relatively thin, spherical mass shell on top of a rigid core. These models have helped considerably in clarifying som e of the complicated physics involved in stellar pulsations and the role played by di erent physical mechanism s.

AGB stars are de ned as stars that develop electron degenerate cores m ade of m atter which has experienced com plete hydrogen and helium burning, but not carbon burning. M ore lum inous stars with highly evolved cores are called Super-AGB stars and their core is m ade of a m ixture of oxygen-neon [14,15]. It is a well known result that the observational counterparts of Super-AGB stars | the so-called Long Period Variables (LPVs) | are radial pulsators. M oreover, long term photom etry of these stars has shown that their light curves (the variation of the lum inosity with tim e) usually have some irregularities [16,17] which lead to a high degree of in predictability. D ue to the lack of appropriate tools for an-

alyzing these irregular uctuations of the lum inosity or the stellar radius, the scienti c community did not pay much attention, until recently, to this category of variable stars. The developm ent of new nonlinear tim e-series analysis tools during the last decade has changed this situation. In particular, it has been found that these tools have rich applications in a broad range of astrophysical situations, which include time analysis of gam ma-ray bursts [18], of gravitationally lensed quasars [19] or X rays within galaxy clusters [20], and of variable white dwarfs [21]. These tools have also been used to analyze theoretical models of stellar pulsations. In particular, it has been proven that num erical hydrodynam ical m odels display cascades of period doublings [22{24] as well as tangent bifurcations [25,26]. To be more speci c, it has been con med [27] that the irregular pulsations of W V irm odels are indeed chaotic and, furtherm ore, it has been shown that the physical system generating the time series is equivalent to a system of three ordinary di erential equations. A similar approach has been used also for the study of the pulsations of other types of stars like, for instance two RV Tau stars, R Scuti [28] and AC Her [29], and has provided signi cant results concerning the underlying dynamics. This result is not a trivial one since it is not obvious at all that stellar pulsations can be fully described by such a simple system of ordinary di erential equations. This stem s from the fact that the classical method to physically describe stellar pulsations is based on the use of a hydrodynam ical code where the partial di erential equations of uid dynam ics are replaced by a discrete approximation consisting of N mass shells. Therefore, a set of 3N coupled nonlinear ordinary di erential equations must be solved, where N is typically of the order of 60. Thus, simple models do not only help in clarifying the basic behaviour of stellar pulsations but, actually, they may be able to reproduce with a reasonable accuracy the oscillations of real stars.

Several such simple models have been proposed and studied [30{37] including the one in [1]. This model was intended for study the linear, adiabatic and radial pulsations of AGB stars. The purpose of this paper is to analyze in depth the linear oscillator proposed by these authors, and to extend their study to more massive and lum inous stars. The paper is organized as follow s. In section II, we sum m arize the basic assumptions of the m odel and we mathematically describe the oscillator, whereas in section III we characterize its dynam ics and describe a sequence of bifurcations which occur as the param eters of the model are varied. In section IV we compare our results with those for the perturbed oscillator, in order to get a better physical insight. In section V we discuss our results and, nally, in section VI we draw our conclusions.

II. DESCRIPTION OF THE MODEL

In the search for a description which embodies the essentials of stellar oscillations, we follow the simple model of the driven keplerian oscillator derived in [1], and successfully applied to AGB stars. The model assumes that the compact stellar interior is decoupled from the extended outer layers. The driving originates in the stellar interior and consists of a pulsation generated by pressure waves. Moreover, we consider only the case of sinusoidal driving, where the outer layers are driven by the interior pressure waves that pass through a transition zone characterized by a certain coe cient of transmission. The motion is calculated at successive states of hydrostatic equilibrium . No back reaction of the outer layers on the inner ones is considered.

We denom inate the driving oscillator \the interior" and the driven oscillator \the mantle". These are separated by a transition zone through which the pressure waves from the interior propagate until they hit the mantle and dissipate. The driving oscillator is represented by variations of the interior radius, R c, around an equilibrium position, R_0 , according to: $R_c = R_0 + R_0 \sin !_c$ where and ! c are the fractional amplitude and the frequency of the driving, respectively. The mantle is represented by a single spherical shell of mass m at instantaneous radius R_m . In absence of any driving force, the equation of m otion is given by:

$$\frac{d^{2}R_{m}}{d^{2}} = \frac{4 R_{m}^{2}}{m}P - \frac{GM}{R_{m}^{2}}; \qquad (2.1)$$

is the time, P is the pressure inside $R_{\,\rm m}$, and where M is the mass of the rigid core. As in [1], we assume that the gas follows a polytropic equation of state with = 5=3. If the region where the pulsation occurs is assumed to be nearly isothermal (i.e., Rm R,with R the equilibrium radius of the star), the equation of motion near the mantle reduces to

$$\frac{d^2 r}{dt^2} = \frac{1}{r^2} \cdot 1 \cdot \frac{1}{r} + Q !^{4=3} \sin [! (t + r_0 + r_0 \sin ! t)] \quad (2.2)$$

where we have introduced the following set of nondim ensional variables:

r
$$R_m = R$$

r_0 $R_0 = R$
t $!_m$
! $!_c = !_m = r_0^{3=2}$

:

In this equations

(



FIG.1. Poincare m ap for Eq.(2.3) with ! = 20.1 (typical of AGB stars) and (a) = 0.1, (b) = 0.4, and for !' 3 (typical of Super-AGB stars) and (c) = 0.1, (d) = 0.4. In all cases, = 1. Note the di erence in the scales of the axes.

$$!_m \qquad \frac{GM}{R^3}$$
 ¹⁼²

is the characteristic dynam ical frequency of the system and Q the transm ission coe cient of the transition zone through which the pressure waves from the interior propagate. In the lim it r 1 (sm all oscillations) and absorbing som e term s into t as phase shifts, one gets:

$$\frac{d^2x}{dt^2} = x + \sin[! (t + 1)^{2-3} \sin[! t)] \quad (2.3)$$

where = Q!⁴⁼³ is the total driving am plitude, and x r 1. The parameter ! is a measure of the core/envelope ratio, and provides inform ation on the location of the source of the driving in the stellar interior. Note that for = 0, Eq.(2.3) transforms into the classical equation of the linear oscillator, x = x. All the interesting features of the motion are generated by the perturbation (the second term of the right-hand side of Eq.(2.3) with \neq 0) and its interaction with the unperturbed motion. The system we are dealing with is a periodic{tim e dependent H am iltonian system.

H (x;y;t) =
$$\frac{x^2 + y^2}{2}$$

 $\frac{1}{1} \cos[! (t x + !)^{2=3} \sin ! t)]$ (2.4)

The associated Poincare map is an area preserving map. Hence, its dynamics does not exhibit attractors or repellers.

III.CHARACTERIZATION OF THE OSCILLATOR

The study performed in [1] explored a limited set of values of the parameter space: = 0.5; 0.75; 1.0, and= 0:1; 0:2; 0:4 for values of ! around ' 20, which is characteristic of regular AGB stars (M 8M). Their main results and conclusions are that small values of ! produce more strongly chaotic pulsations and that for large values of stable orbits in some de nite regions of the phase space are obtained. Our aim is to extend the previous study to Super-AGB stars, which are characterized by a di erent value of ! . For the physical conditions found in Super-AGB stars, 8M М 11 M and R 450R, it turns out that ! > 3. W e will also explore a wider range of the parameter space than that investigated in [1]. Contrary to [1], who adopted a fourthorder predictor-corrector scheme for the num erical integration of Eq.(2.3), we use a fourth order Runge-K utta integrator with step-size control and dense output as described in [38]. We have tested several other integrators, speci cally designed for sti problem s [39], and we have obtained the same results for a given set of initial conditions. Thus we conclude that our num erical integrator is appropriate for problem under study.



FIG.2. The e ect of increasing on the Poincare map: (a) = 0.2; (b) = 0.3 and (c) = 0.4. = 0.5 in all cases.

A.AGB stars vs. Super-AGB stars: the role of !

In order to com pare the behaviour for two di erent values of! (and, thus, the di erences between pulsations of AGB and Super-AGB stars), Figure 1 show sthe Poincare m aps obtained with ! ' 20, typical of regular AGB stars (panels a and b), and those obtained for ! ' 3 typical of Super-AGB stars (panels c and d), for the same initial conditions. As ! increases the behaviour becomes m ore

irregular, and the islands and the structure of panels (a) and (b) quickly disappear, leading to a more chaotic behaviour. Consequently, we expect that the pulsations of Super-AGB stars will exhibit a more chaotic behaviour than those of AGB stars.

The phase portrait of our model is a typical example of an area-preserving map. For very small values of the perturbation parameter, it exhibits invariant circles on which them otion is quasi{periodic. Three periodic island chains surround the xed point of the map. For large perturbations, because of the breaking {up of the invariant circles and splitting of the separatrices of hyperbolic periodic points, chaotic zones are present, as well as islands in the stochastic sea. Until recently, it was thought that the existence of islands was not very important in determ ining the origin and character of chaos in H am iltonian systems. However, it has been recently shown that the boundaries of such islands are very interesting [40 [42]. In particular, crossing an island boundary means a transition from a regular periodic or quasiperiodic behaviour to an irregular one that lies in the stochastic sea. One of the particularities of this singular zone is illustrated in Figure 1a. O ne can see a central island em bedded in the dom ain of chaotic motion with a boundary separating the area of chaos from the region of regular motion. By changing a controlparam eter of the system, several bifurcations occur which in uence the topology of the boundary zone, leading to the appearance and disappearance of sm aller islands. Thism ay result in self-sim ilar hierarchical structures of islands which is crucial for understanding chaotic transport in the stochastic sea and the general dynam ics, since they represent high-order resonances.

Figure 1b illustrates another essential phenom enon typical of som e H am iltonian system s. It is linked to the existence of a com plex phase space topology in the neighborhood of som e islands: a trajectory can spend an inde nitely long time in the boundary layer of the island (ie., a dynam icaltrap). The main uncertainty in characterizing this phenom enon concerns the level of stickiness (that is, its characteristic trapping time) which depends on the parameters of the system in a still unknown manner. In general, the hierarchical structure of the islands in the boundary zone can explain the origin of the stickiness of the trajectory within this region, which is the reason why this behaviour is called hierarchical { islands trap [42]. Much attention was generally paid to understanding the structure of these islands because di erent islands correspond to di erent physical processes responsible for their origin. The boundary layer of an island m ay include higher-order resonant islands. A nother im portant reason is that the topology of the islands could be an indicator for the vicinity of bifurcations. However, no general description for the birth and collapse of hierarchical islands exists yet.



FIG.3. Poincare m aps for Eq.(2.3), = 0.3 and di erent values of . The successive birth of two dimerized island chains. The hom oclinic and heteroclinic connections appear in red color.

B. The role of the fractional driving am plitude,

In order to characterize in depth the behaviour of our system, we have perform ed a thorough param etric study by varying and , while keeping ! constant at the value typical of Super-AGB stars. Reasonable values of the fractional amplitude, , range from 0.1 to 0.4, whereas the total driving am plitude, , varies between 0.1 and 1.0. Since is the ratio between the amplitude of the internal driving and the radius of the star, the upper lim it considered in this study is 40%, which is physically sound. In general, a star is characterized by its com pactness, !, and the interior-mantle coupling strength $Q = = (!^{4=3})$, which should stay within the limits of 0 (0% transm ission) and 1 (100% transm ission) in order to maintain its physical meaning. Therefore some speci c combinations of (;) that lay outside this range for Q must be regarded with caution, since they have no physical meaning. According to all these considerations, we have investigated the dynam ics of the system associated to Eq.(2.3) rewritten below in the form of a perturbed oscillator,

$$\underline{x} = y$$
 (3.1)
 $\underline{y} = x + \sin[! (t + b \sin ! t)]$

where b = ! ²⁼³, m aintaining a constant value of ! ' 3, as discussed above.

In a rst step, we have studied the qualitative changes in the dynamics (i.e., bifurcations) as the parameters and are varied. Poincare maps corresponding to the case 2 (0:1 0:4) and = 0:5, for several initial conditions, are shown in Figure 2. The maps are characterized by the same geometric structure: a region, centered around (x;y) = (0;0), of closed orbits, surrounded by a region of chaotic orbits. As it can be seen in Figure 2, as increases, the central region of regular behaviour expands to the detrin ent of the stochastic sea.

We have also analyzed the behaviour of the system for negative values of . A quick look at the driving allows one to see that negative values of correspond to a mere change of phase in the driving and, therefore, correspond also to physically meaningful cases. How ever, and for the sake of clarity and conciseness, in this paper we will concentrate our e orts on the positive dom ain of this parameter. Nevertheless, we would like to remark at this point that the phase portrait of the Poincare map changes drastically depending on whether is positive, negative or alm ost zero.



FIG.4. Creation by saddle{ center bifurcation of a pair of three periodic orbits for = 0.3 and increasing values of . (a): = 0.0875 | the phase portrait at the threshold of the bifurcation; (b): the invariant circle with 6 cusps for the case (a); (c) = 0.095 | two intertwined orbits of period three have been created, inside each new born loop there is a period three elliptic point; (d): a sequence of heteroclinic and hom oclinic connections between the new born hyperbolic points; (e): = 0.11725 | the threshold of global bifurcation; (f): the new topology of the separatrices.

C. The role of the total driving am plitude,

Next we have focused on values of 1 in order to observe in detail the departure of our equation from the harm onic oscillator as this parameter increases. We restricted the study of the Poincare map to a rectangle limited by initial conditions close to reasonable values of the radius and velocity of the mantle. The restriction is also due to the approximations involved in deriving Eq.(2.3). For = 0 we get an integrable Ham iltonian system, whose integrals of motion are the tori given by the condition $x^2 + y^2 = C$; t = [0;2 = !). The Poincare map has the elliptic xed point (0,0), which corresponds

to a periodic orbit of period P = 2 = ! of the perturbed H am iltonian system .

We have chosen the range 2 (0;0:12) and = 0:3and we have obtained a cascade of local and global bifurcations. Note that the considered pairs of parameters (;) are located above the curve (), which is the curve of triplication of the elliptic xed point. A coording to [43,44] the corresponding Poincare map is in these cases a nontwistm ap. The sequence of detected local and global bifurcations is typical for such a map [45[48]. Figure 3 illustrates the birth in stages of two dimerized island chains containing periodic points of period three. This is a typical scenario of creation of new orbits in nontwist m aps. The upper-left panel exhibits the phase portrait of the near{to{integrable map. The elliptic xed point is surrounded by invariant circles. At = 0:03 starts the birth of the rst dimerized island chain, namely a regular invariant circle bifurcated to an invariant circle with three cusps. The newly born dimerized island chain is clearly seen in the bottom {left panel of the Figure 3, where a new circle with three cusps, closer to the the xed point can be seen. Both dimerized island chains are illustrated in the bottom {right panel. Between them the invariant circles are meanders. Note that each cusp is a point of saddle{center creation [49]. This scenario is a good example to illustrate the theoretical results concerning the creation of a twistless circle after triplication [43,44].



FIG.5. Poincare sections for = 0.3 and di erent values of : (a) = 0.2; (b) = 0.3, and (c) = 0.4. In red and green colors are represented evolutions of initial conditions close to the rst and respectively second separatrix.

A round the last born three{periodic dimerized island chain, a sequence of local and global bifurcations occurs.

This is illustrated in Figure 4, where it can be seen that two independent orbits of the sam e period three are created by cusp bifurcation. They evolve in such a way that, nally, interact with the orbits which belong to the rst dim erized island chain. As increases the new ly born elliptic points approach the hyperbolic points of the dim erized chain. W hen reaches a value of 0.11725 a globalbifurcation occurs: the new ly created orbits interfere, and the hyperbolic points of the dimerized island chain becom e hyperbolic points with hom oclinic eight-like orbits encircling the new created elliptic points. We have also noticed that as increases, the process of creation of three periodic orbits repeats outside the region of the Poincare map we focus on, and the new orbits form chains of vortices, not dimerized island chains. This behaviour is due to the oscillating character of the nontwist property of the perturbation.

In a second step we have paid attention to larger values of . In particular we have studied the range 2 (0.2;0.5). The increase in strength of the external perturbation destroys the separatrices (Figure 5, top) by

clothing every one of them in a stochastic layer (Figure 5, middle). As the thickness of the layers increases with the perturbation, depending on the positions of the separatrices in the phase space, they can merge form ing the stochastic sea (Figure 5, bottom).

There are m any islands, which the chaotic trajectory cannot penetrate. W ithin an island there are quasiperiodic motions (invariant tori) and regions of trapped chaos. The stronger the chaos is, the smaller the islands and the larger the fraction of phase space occupied by the stochastic sea. The coexistence of regions of regular dynamics (closed orbits) and regions of chaos in the phase space is a wonderful example of the property which di erentiates chaotic systems from ordinary random processes, where no stability islands are present. This property m akes possible the analysis of the onset of chaos and the appearance of m inim al regions of chaos. To sum m arize, the system undergoes the following bifurcations, as it results from Figures 3, 4 and 5:

- (a) For 2 [0;0:03), the phase portrait of the Poincare map is similar to that of the harmonic oscillator, with the elliptic xed point (x;y) = (0;0) surrounded by almost circular closed orbits.
- (b) For 2 [0:03;0:04), in addition to the central elliptic point, a dimerized island chain is born, namely, three periodic elliptic and hyperbolic points, each elliptic point being surrounded by a hom oclinic orbit to the corresponding hyperbolic point, and distinct hyperbolic points being connected by heteroclinic orbits.
- (c) For 2 [0:04;0:07), a second dim erized island chain is created, which is a typical con guration after the triplication of an elliptic xed point.
- (d) For 2 [0:07;0:2), a sequence of local and global bifurcations occurs which interphere with the rst

created dimerized island chain: a regular meander bifurcates to a meander with 6 cusps. A new dimerized island chain is created by saddle{center bifurcation. If we denote the cusps as 1;2;:::;6 in clockwise order, then the elliptic and hyperbolic points born from the odd cusps, form independent orbits of those born from the even cusps. This particular dimerized island chain approaches the original one, and as a result the hyperbolic points of the two chains are connected giving rise to a pattern of a complicated topological type (Figure 4f).

- (e) For 2 [02;03), we notice a more de nite chaotic behaviour created by the instability of the remainings of the rst heteroclinic orbit engul ng the periodic points of case (c).
- (f) For 2 [0:3;0:5], we notice the extension of the chaotic orbits in the central region, together with



FIG.6. (a): A m eander near an invariant circle with 35 cusps for = 0.4 and = 0.3; (b): two orbits, one lling the m eander and the second, a chaotic orbit, con ned by the m eander.

A nother in portant feature of the phase portrait characteristic of nontwist maps is the existence of meanders (i.e., invariant circles which fold exactly as a meander). Meanders are created between two successively born dimerized island chains, and between two chains of vortices. In twist standard{like maps they become usual circles after the reconnection of the two chains. Note that in our case the reconnection of the rst two created dimerized island chains, does not occur, and as increases the meanders break {up leaving instead a C antor set. The top panel of Figure 6 illustrates a meander near a pair of dimerized island chains containing periodic orbits of period 35. Meanders appear to be robust invariant circles, even when the nearby orbits are chaotic (Figure 6, bottom panel). This behaviour was observed in nontwist standard-like maps [47], but until now there is no explanation for this robustness.

IV.COM PARISON W ITH THE PERTURBED OSC ILLATOR

In order to get a better insight of the underlying physics of the oscillator studied so far, in this section we are going to compare it with the motion of a perturbed oscillator, which has been already studied extensively. This is in portant since, as it will be shown below, the form alappeareance of Eq.(2.3) does not dier very much from that of a perturbed oscillator. In order to make this clear consider the motion of a perturbed linear oscillator in the form of:

$$x + !_{0}^{2}x = sin[!x !t]$$
 (4.1)

The left-hand side of Eq.(4.1) describes very simple dynam ics of linear oscillations of frequency $!_0$. To ease the comparison with the perturbed oscillator, Eq.(2.3) can be rewritten as:

$$x + !_{0}^{2}x = [!x ! (t)t];$$
 (4.2)

where

$$!(t) = ! \frac{b\sin !t}{t} + 1$$
 (4.3)

Both equations share common features. It is important to point out here that for our system $!_0 = 1$, as it results from Eq.(2.3). The Ham iltonian associated with the system given by Eq.(4.1) is:

$$H = \frac{(x^{2} + !_{0}^{2}x^{2})}{2} - \frac{1}{!}\cos[!x !(t)t] \qquad (4.4)$$

C onsequently, for weak perturbations the results obtained so far should be similar to the case of a perturbed oscillator. And indeed this is the case, the only dimension resides in the period 3 hom oclinic orbit, that is in point d) of the summary given in section III.C. The creation of this period-3 hom oclinic structure was illustrated in Figure 2 as increases from = 0.2 to = 0.3. This suggests that the ultimate reason of this dimension might be the presence of the function ! (t) from Eq.(4.3) as a weak detuning of resonance.



FIG.7. Variations of radius (top) and velocity (bottom) as a function of time for !'3, = 0.3, = 0.5 and the initial condition (x₀;y₀) = (0.0;0.02). The escape velocity of the model star (v_{esc} ' 86 km/s) is represented as a red line.

e rewrite Eq.(4.2) with
$$\frac{6}{9}$$
 0 in the following way:
 $x + !_0^2 x = sin ! (t x + bsin ! t)]$ (4.5)

For = 0, Eq.(4.5) transform s into:

W

$$x + !_0^2 x = sin[!(t x)]$$
 (4.6)

All these ordinary di erential equations, together with the wellknown equation for the perturbed oscillator, have the same mathematical structure. It is an example of weak chaos, where the perturbation itself creates the separatrix network at a certain $_0$ and then destroys it as increases beyond 0 by producing channels of chaotic dynam ics. For an unperturbed equation, the unperturbed Ham iltonian H₀ intrinsically has separatrix structures and the perturbation clothes them in thin stochastic layers, an example of strong chaos. In the phase space there appear invariant curves (deform ed tori) em bracing the center which do not allow diusion in the radial direction. Inside these cells of the web, motion occurs along closed-orbits, around the elliptic points from the centers of the cells. With the increase of and creation of the stochastic layers, particles can wander along the channels of the new ly born web, a phenom enon that represents an universal instability and gives birth to chaotic uctuations. The heteroclinic structures form ed as the perturbation increases through this bifurcation are what di erentiates our equation from the typical equations of Ham iltonian chaos and in particular from those of the perturbed oscillator.

V.ASTROPHYSICAL INTERPRETATION OF THE RESULTS

In order to provide the reader with a feeling of the physical ranges for the stellar uctuations in radius and

velocity, in Figure 7 we show a particular time series of the model presented above. In this gure the variations of radius and velocity as functions of time are represented both in physical units (solar radii, km/s and years, respectively) and in adimensionalized units | as we have been doing so far.

G enerally speaking, the prim ary inform ation that one can derive from a generic computed or observed tim e series is the spectral distribution of energies (or am plitudes) of the light curve. We found that for some initial conditions, the resultant time series do not show a single frequency or a sm all set of frequencies but, in addition, linear combinations of the prim ary frequencies m ay also appear in the spectra. This behaviour has been found in real stars quite frequently, and in particular in LPVs, C epheids and RR Lyrae stars [50].

In any calculation of a time-varying phenom enon, like the one we are describing here, there is always the im portant practical question of when the sim ulations should be term inated. For full nonlinear hydrodynam ic sim ulations, even in the case in which a stable lim it cycle seem s to have set in, one often is worried about the fact that therm alchanges, which take place on a longer tim escales, could still be occurring. In a recent paper [52], the authors have carried the calculations of M ira variable pulsationsmuch further than in previous investigations and, to their surprise, they have found that the actual behaviour was signi cantly di erent from what it was previously thought. To be precise, they obtained a new modi ed \true" lim it cycle which is quite di erent from the earlier, \false" lim it cycle. In agreem ent with these results, the num erical integration of our system presents a sim ilar behavior, despite the very crude approach adopted here which results in an extrem e sim plicity of the model. This is shown in Figure 7 for a particular orbit: during

the rst 1400 yr the star oscillates quite regularly, and with small am plitudes. A fler this period of time, the quiet phase suddenly stops and pulsations become more violent and chaotic. Moreover, at late times the velocity of the outer layers exceeds the escape velocity and, hence, mass loss is very likely to occur, in accordance with the observations of LPV s, which are the observational counterparts of Super-AGB stars.



FIG.8. Poincare map for the orbit of Figure 7. Panel (a) includes the points R () and V () for the time interval (1000-1400) years, whereas panel (b) shows the points R () and V () for the time interval (1000-2000) years.

This behaviour | a quiet phase, followed by an extrem ely violent phase | is a typical case of a dynam ical trap discussed in section III. The Poincare m ap associated with the radius and velocity variations in Figure 7 is shown in Figure 8. None of these gures include the oscillations previous to 1000 years because the behaviour is sim ilar to the one of the tim e interval between 1000 and 1400 years. This example corroborates the suggestion, rst proposed by [53] and later found in the very detailed num erical calculations of [52] that the long-term e ects in M ira variables have an important role in our understanding of the mechanism which drives mass-loss. This result also argues in favour of the capability of simple m odels to capture the underlying dynam ics of the physical system. To put it in a di erent way, our m odel, despite its sim plicity, reproduces reasonably well the results obtained with full, sophisticated and nonlinear hydrodynam icm odels. It is nevertheless important to realize here that our m odel does not incorporate the secular e ects induced by the therm al changes and, hence, it is quite likely that this kind of behaviour is intrinsically associated with the physical characteristics of the oscillations of real stars.

In order to get a more precise physical insight a tim efrequency analysis would be very valuable. Consequently we devote the remaining of this section to such a purpose. M ost often, astronom ers have used wavelet transform s in their investigations of the tim e-frequency characteristics of variable light curves. However, [54] com pared the results obtained with di erent tim e-frequency m ethods using both real light-curves and synthetic signals. From these tests they stly concluded that the G abor transform [55] provides much more informative results on the high frequency part of the data than the wavelet transform, but, on the contrary, they also found that the tim efrequency analysis using the Choi-W illiam s distribution [56] is de nitely superior to both methods (at least on the data they used). Further investigations carried out by the same authors [51] led them to conclude that in general one cannot claim a priori that any of these m ethods is better than the others. In fact, this depends largely on the nature of the signal and on which kind of features one tries to enhance. Therefore, it is always advantageous to use simultanously several of them . How ever, as the Choi-W illiams distribution is generally accepted to be the most useful am ong all the methods, we will center our analysis using this tool, as it was done in [51].

The time-frequency analysis of a model computed with ! ' 3, = 0.3, = 0.5 and the initial condition $(x_0; y_0) = (0.0; 0.02)$, is shown in the central panel of Figure 9. In order to be used as a visual help this time series has been reproduced in the bottom panel of this gure. We have split the time series into two pieces, in order to better illustrate the stickiness of the oscillator, and we have computed the Fourier transforms of both pieces separately. These Fourier transforms are shown in the top panels of Figure 9, the left one corresponds to < 1400 yr, whereas the right one corresponds to

1400 yr. Note the di erence in the scales of the power, which is much larger for the right panel. As it can be seen in these panels the power is suddenly shifted from relatively large frequencies to much sm aller frequencies.



FIG.9. Choi-W illiam s distribution of the velocity variations (central panel), Fourier analysis for < 1440 yr (top left panel) and for 1440 yr (top right panel), and the corresponding time series (bottom panel).

This behavior is even more clearly shown in the Choi-W illiams distribution. Note as well the contribution of the frequency f ' 2.1 yr ¹ at small times, just before the beginning of the burst (at 1430 yr). Indeed, this small contribution could be at the origin of the transfer of power to f ' 1.2 yr ¹, which occurs inmediately after ' 1430 yr and which ultimately leads to the burst at ' 1440 yr. A lso remarkable is the small time elapsed since the beginning of the transfer of power. which is only

since the beginning of the transfer of power, which is only about 10 yr.

VI.CONCLUSIONS

W e have presented and analyzed the complex dynam ics of a forced oscillator which is interesting not only from the m athem atical point of view, but also because it describes with a reasonable degree of accuracy the m ain characteristics of some stellar oscillations. This m odel has been previously used to study the irregular pulsations of A sym ptotic G iant B ranch stars [1], and we have used it to study the pulsations of m ore m assive and lum inous stars, the so-called Super-A sym ptotic G iant B ranch stars. In doing this we have extended the previous studies to a range of the parameters specic for this stellar evolutionary phase. We have found, in agreement with the observations of Long Period Variables which are the observational counterparts of Super-A sym ptotic G iant B ranch stars, that the oscillator shows a chaotic behavior. It is in portant to realize as well that this kind of stars shows a more pronounced chaotic behavior than regular A sym ptotic G iant B ranch stars of sm aller m ass and lum inosity. We have also characterized in depth the full sequence of bifurcations as the physical parameters of the model are varied. We have found a rich set of local and global bifurcations which were not described in [1]. Among these, perhaps the most important one is a tripling bifurcation, but m eandering curves, hierar-

chical islands traps and sticky orbits also appear. Correspondingly, the resulting time series also show a rich behaviour. In particular, we have found that although there are light curves which show a rather regular behaviour for certain values of the param eters of the physical system and given initial conditions, there are as well som e other light curves which show clear beatings or linear combinations of two main frequencies up to term s of $2f_0 + 7f_1$, being f_0 the fundamental frequency and f_1 that of the st overtone, and even more complex orbits. For the parameters and initial conditions leading to more irregular behaviour, we noticed the existence of both clear chaotic pulsations and sudden changes from a lim it-cycle to chaotic pulsations, the latter being associated with the stickiness phenom enon characteristic of som e H am iltonian system s. For these orbits the velocity of the very outer layers clearly exceeds the escape velocity. Hence, for these chaotic pulsations mass-loss is very likely to occur, in good agreem ent with the observations, which correlate the degree of irregularity with the massloss rate. Regarding the stickiness of som e orbits which we have found for a set of parameters, perhaps the most in portant result is that the long-term e ects found in real stars are reproduced by our model, despite of its sim plicity, even though the driven oscillator studied here does not incorporate the e ects of secular changes. Hence it is quite likely that this kind of behaviour which has been already found in full hydrodynam ical simulations [52], is intrinsically associated with the physical characteristics of the oscillations of real stars and not to the long-term therm alchanges.

ACKNOW LEDGMENTS

This work has been supported by the DGES grant PB98{1183{C03{02, by the MCYT grant AYA2000{ 1785, and by the CIRIT grant 1995SGR-0602. We also would like to acknow ledge m any helpful discussions with G M.Zaslavsky.

- [L] V. Icke, A. Franck, H. Heske, Astron. & Astrophys., 258, 341 (1992)
- [2] A.Gautschy, H.Saio, Annu. Rev. Astron. & Astrophys., 34, 551 (1996)
- [3] JP. Cox, Theory of Stellar Pulsation, Princeton Univ. Press: Princeton (1980)
- [4] J.R. Buchler, A strophys. & Space Sci., 210, 9 (1993)
- [5] A.G autschy, H.Saio, Annu.Rev.Astron. & Astrophys., 33, 113 (1995)
- [6] J.R. Buchler, Z. Kollath, R. Cadmus, in Proceed-

ings of the 6^{th} Experim ental Chaos Conference, astroph/0106329 (2001)

- [7] K. Pollard, C. A loock, D R. A lves, D P. Bennett, K H. Cook, S L. Marshall, D. Minniti, R A. A llsman, T S. Axelrod, K C. Freeman, B A. Peterson, A W. Rodgers, K. Griest, J A. Guem, M L. Lehner, M. L., A. Becker, M R. Pratt, P J. Quinn, W. Sutherland, D L. Welch, in Variable Stars and the A strophysical Returns of Microlensing Surveys, Eds.: R. Ferlet, JP. Maillard and B. Raban, Editions Frontiers: Paris, p. 219 (1997)
- [8] J. Perdang, in The Numerical Modelling of Nonlinear Stellar Pulsation, Ed.: J.R. Buchler, K luwer: Dordrecht, p. 333 (1990)
- [9] S.Hofner, U.G. Jorgensen, R. Loidl, B. Aringer, Astron.
 & Astrophys., 340, 497 (1998)
- [10] S.Hofner, Astron. & Astrophys., 346, L9 (1999)
- [11] JM.W inters, T.LeBertre, K.S.Jeong, Ch.Helling, E. Sedlm ayr, Astron. & Astrophys., 361, 641 (2000)
- [12] Y J W . Sim is, V. Icke, C. Dom inik, Astron. & Astrophys., 371, 205 (2001)
- [13] G.Bono, M.Marconi, R.F. Stellingwerf, Astron. & Astrophys., 360, 245 (2000)
- [14] C. R itossa, E. G arc a-Berro, I. Iben, A strophys. J., 460, 489 (1996)
- [15] E.Garca-Berro, C.Ritossa, I.Iben, Astrophys. J., 485, 765 (1997)
- [16] D.Barthes, Astron. & Astrophys., 333, 647 (1998)
- [17] D. Barthes, X. Luri, R. A Ivarez, M. O. M ennessier, Astron. & Astrophys. Suppl. Ser., 140, 55 (1999)
- [18] J.P.Norris, R.J.Nem iro, J.D. Scargle, C.Kouveliotou, G.J.Fishman, C.A.Meegan, W.S.Paciesas, J.T.Bonnel, Astrophys. J., 424, 540 (1994)
- [19] J. H jorth, A strophys. J., 424, 106 (1994)
- [20] E. Slezak, F. Durret, D. Gerbal, Astron. J., 108, 1996 (1994)
- [21] M.J.Goupil, M.Auvergne, A.Baglin, Astron. & Astrophys., 250, 89 (1991)
- [22] J.R. Buchler, G.Kovacs, Astrophys. J., 320, L57 (1987)
- [23] G.Kovacs, J.R. Buchler, Astrophys. J., 334, 971 (1988)
- [24] T. Aikawa, Astrophys. & Space Sci., 164, 295 (1990)
- [25] J.R. Buchler, M. J. Goupil, G. Kovacs, Phys. Lett. A, 126,177 (1987)
- [26] T. Aikawa, Astrophys. & Space Sci., 139, 281 (1987)
- [27] T. Serre, Z.Kollath, J.R. Buchler, Astron. & Astrophys., 311, 833 (1996)
- [28] J.R. Buchler, T. Serre, Z.Kollath, Phys. Rev. Lett., 74, 842 (1995)
- [29] Z.Kollath, J.R. Buchler, T. Serre, J.M. attei, A stron. & A strophys., 329, 147 (1998)
- [30] N. Baker, in Stellar Evolution, Eds.: R F. Stein and A G W. Cameron, Plenum Press: New York, p. 333 (1966)
- [31] N. Baker, D. Moore, E A. Spiegel, Astron. J., 71, 845 (1966)
- [32] T J.Rudd, R M.Rosenberg, Astron. & Astrophys. 6, 193 (1970)
- [33] R.F. Stellingwerf, Astron. & Astrophys., 21, 91 (1972)
- [34] J.R. Buchler, O. Regev, Astrophys. J., 263, 312 (1982)
- [35] Y. Tanaka, M. Takeuti, A strophys. & Space Sci., 148, 229 (1988)
- [36] M. Saitou, M. Takeuti, Y. Tanaka, Publ. A stron. Soc.

Pac., 41, 297 (1989)

- [37] W. Unno, D.-R. X iong, A strophys. & Space Sci., 210, 7 (1993)
- [38] E. Hairer, S.P. Norsett, G. Wanner, Solving Ordinary Di erential Equations, Nonsti Problem s, Springer-Verlag: Berlin (1993)
- [39] L.Sham pine, M.Gordon, Computer Solution of Ordinary Dierential Equations, Freeman: San Francisco (1975)
- [40] G M .Zaslavsky, Phys. Today, 52, 39 (1999)
- [41] G M. Zaslavsky, M. Edelman, B A. Niyazov, Chaos, 7, 159 (1997)
- [42] G M .Zaslavsky, Physics of chaos in H am iltonian system s, Im perial College Press: London (1998)
- [43] H R. Dullin, JD. M eiss, D. Sterling, Nonlinearity, 13, 203 (2000)
- [44] V J.A mold, V V .K ozlov, A J.N eishtadt, D ynam icalSystem s III, Springer-Verlag: Berlin (1988)
- [45] D. del-Castillo-Negrete, JM. Greene, P.J. Morrison, Physica D, 91, 1 (1996)
- [46] D. del-Castillo-Negrete, JM. Greene, P.J. Morrison, Physica D, 100, 311 (1997)
- [47] C.Simo, Regular and Chaotic Dynamics, 3, 180 (1998)
- [48] E. Petrisor, submitted to Nonlinearity (2002)
- [49] J. K. Hale, H. Kocak, Dynamics and bifurcations, Springer-Verlag: Berlin (1991)
- [50] Z.Kollath, J.R. Buchler, in Stellar Pulsation { nonlinear studies, Eds: M. Takeuti & D. Sasselov, K luwer: Dordrecht, p. 29 (2001)
- [51] J.R. Buchler, Z.Kollath, in Stellar Pulsation { nonlinear studies, Eds: M. Takeuti & D. Sasselov, K luwer: Dordrecht, p. 185 (2001)
- [52] A. Ya'ari, Y. Tuchm an, Astrophys. J., 456, 350 (1996)
- [53] J.R. Buchler, W. R. Yueh, J.Perdang, Astrophys. J., 214, 510 (1977)
- [54] Z.Kollath, J.R. Buchler, in Nonlinear Signal and Im age A nalysis, Eds: J.R. Buchler and H.K andrup, New York A cadem y of Sciences Press: New York, 808, p.116 (1997)
- [55] D.Gabor, J.Inst.Elec.Eng., 93, 429 (1946)
- [56] H J.Choi, W J.W illiam s, EEE Trans.on A coust. Speech and Signal Processing, 37, 862 (1989)

