

Long-term variability survey of the old open cluster NGC 6791.

B. J. Mochejska¹, K. Z. Stanek

Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138

bmochejs@cfa.harvard.edu, kstanek@cfa.harvard.edu

J. Kaluzny

Copernicus Astronomical Center, Bartycka 18, 00-716 Warszawa

jka@camk.edu.pl

ABSTRACT

We present the results of a long-term variability survey of the old open cluster NGC 6791. The *BVI* observations, collected over a time span of 6 years, were analyzed using the ISIS image subtraction package. The main target of our observations were two cataclysmic variables B7 and B8. We have identified possible cycle lengths of about 25 and 18 days for B7 and B8, respectively. We tentatively classify B7 as a VY Scl type nova-like variable or a Z Cam type dwarf nova. B8 is most likely an SS Cygni type dwarf nova. We have also extracted the light curves of 42 other previously reported variable stars and discovered seven new ones. The new variables show long-period or non-periodic variability. The long baseline of our observations has also allowed us to derive more precise periods for the variables, especially for the short period eclipsing binaries.

Subject headings: open clusters and associations: individual: NGC 6791 – cataclysmic variables – binaries: eclipsing – stars: variables: other – color-magnitude diagrams

1. INTRODUCTION

The open cluster NGC 6791 is unique in many ways. At an age of about 8 Gyr it is believed to be the oldest open cluster in the Galaxy (Chaboyer et al. 1999, Kaluzny & Rucinski 1995). It is also probably the most metal-rich, with $[\text{Fe}/\text{H}]$ estimates ranging from 0.1 to 0.5 dex (Friel et al. 2002, Chaboyer et al. 1999, Peterson & Green 1998, Kaluzny & Rucinski 1995). NGC 6791 possesses two of the three cataclysmic variables (CVs) found in open clusters (Kaluzny et al. 1997), with the third residing in M67 (Gilliland et al. 1991)².

Variability in NGC 6791 was first studied by Kaluzny & Rucinski (1993), who discovered 17 variable stars, among them one CV candidate. Rucinski, Kaluzny & Hilditch (1996) reported ad-

ditional 5 variables, including another potential CV. The CV candidates were confirmed spectroscopically by Kaluzny et al. (1997). Mochejska et al. (2002; hereafter M02) discovered additional 47 variables, bringing the total to 69.

In this paper we present a long-term variability study of the open cluster NGC 6791. Over a time span of nearly 6 years we have collected 465 observations in *V*, 229 in *I* and 72 in *B*. Our main motivation was to study the long-term behavior of the two cataclysmic variables. This dataset also offered us the unique possibility of investigating long period and non-periodic variables in the cluster.

The paper is organized as follows: Section 2 describes the observations, §3 summarizes the data reduction and variable selection procedures and §4 contains the variable star catalog. Concluding remarks are found in §5.

¹Hubble Fellow

²Another unconfirmed as of yet candidate was reported by Mochejska & Kaluzny (1999) in NGC 7789.

2. OBSERVATIONS

The data analyzed in this paper were obtained between September 1996 and May 2002 at three telescopes, using four CCD cameras. The observations were collected during:

1. 28 nights between 8 Sep and 23 Oct 1996 on the 1.3 m McGraw-Hill Telescope at the Michigan-Dartmouth-MIT (MDM) Observatory, equipped with the front-illuminated, Loral 2048² CCD “Wilbur” (Metzger, Tonry & Luppino 1993). This dataset will hereafter be referred to as the *MDM* dataset.
2. 33 nights between 26 Oct 1996 and 9 Oct 1997 on the 1.2 m telescope at the F. L. Whipple Observatory (FLWO), equipped with the thinned, back-illuminated, AR coated Loral 2048² CCD “AndyCam” (Szentgyorgyi et al. 2002), hereafter *AndyCam* dataset.
3. 47 nights between 19 Sep 1998 and 22 May 2002 on the 1.2 m telescope at FLWO, equipped with the “4Shooter” CCD mosaic with four thinned, back-illuminated, AR coated Loral 2048² CCDs (Szentgyorgyi et al. 2002), hereafter *4Shooter* dataset. Only data from chip 3, centered on the cluster, were analyzed here.
4. 15 nights between Sep 30 and Nov 7, 1999 and between Oct 14 and 21, 2001 on the the Kitt Peak National Observatory³ (KPNO) 2.1 m telescope, equipped with the Tektronix 2048² CCD “T2KA”, hereafter *KPNO* dataset.

Typical exposure times for the 1 m/2 m telescopes, respectively, were 450/600 s in *B*, 450/300 s in *V* and 300/300 s in *I*. Some short exposures were also collected to investigate the variability of bright stars, which are normally saturated on the longer exposures. Typical 1 m/2 m telescope short exposure times were 45/30 s in *V* and 30/30 s in *I*. Short *B* exposures were only taken at KPNO

³Kitt Peak National Observatory is a division of NOAO, which are operated by the Association of Universities for Research in Astronomy, Inc. under cooperative agreement with the National Science Foundation.




Fig. 1.— Digital Sky Survey image of NGC 6791 showing the field of view of the 4Shooter chip3 (solid line), MDM (dotted), AndyCam (short-dashed) and KPNO (long-dashed) CCDs. North is up and east is to the left

and their exposure times varied between 25 and 180 s.

Over 6 years of monitoring of this cluster we have collected 391 long and 74 short exposures in *V*, 140/89 in *I* and 65/7 in *B*. The number of *BVI* images in each dataset is listed in Table 1.

3. DATA REDUCTION

3.1. Photometry

The preliminary processing of the CCD frames was performed with the standard routines in the IRAF ccdproc package.⁴ The KPNO data were corrected for CCD non-linearity at this stage, as described by Mochejska et al. (2001).

Photometry was extracted using the ISIS image subtraction package (Alard & Lupton 1998, Alard 2000). A brief outline of the applied reduction procedure is presented here. For a more detailed description the reader is referred to M02.

⁴IRAF is distributed by the National Optical Astronomy Observatories, which are operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the NSF.

The ISIS reduction procedure consists of the following steps: (1) transformation of all frames to a common (x, y) coordinate grid; (2) construction of a reference image from several best exposures; (3) subtraction of each frame from the reference image; (4) selection of stars to be photometered and (5) extraction of profile photometry from the subtracted images.

In further discussion we will refer to images called “template” and “reference”. By “template” we mean a single best quality exposure in a filter and dataset combination. A reference frame is a high S/N image constructed from the template and several other high quality exposures processed to match the template point-spread function (PSF) and background level. A template is chosen and a reference image constructed for each of the 19 filter and dataset combinations.

All computations were performed with the frames internally subdivided into four sections (`sub_x=sub_y=2`). Differential brightness variations of the background were fit with a first degree polynomial (`deg_bg=1`). A convolution kernel varying quadratically with position was used (`deg_spatial=2`). The psf width (`psf_width`) was set to 33 pixels for all chips, except for the KPNO data, where it was set to 15 pixels. We used a photometric radius (`radphot`) of 5 pixels for 4Shooter and AndyCam, 4 pixels for MDM and 3 pixels for KPNO data.

3.2. Calibration

The transformations of instrumental magnitudes to the standard system were derived from observations of 67 stars in 21 Landolt (1992) standard fields, collected with the 4Shooter on 4 November 1999. The following transformations were adopted:

$$\begin{aligned} v &= V + 2.1326 + 0.0439(B - V) + 0.1480(X - 1.25) \\ b - v &= 0.1275 + 0.9151(B - V) + 0.1074(X - 1.25) \\ v &= V + 2.1318 + 0.0385(V - I) + 0.1478(X - 1.25) \\ v - i &= -0.7390 + 1.0240(V - I) + 0.0880(X - 1.25) \end{aligned}$$

Figure 2 shows the V , $B - V$ and $V - I$ transformation residuals of the standard stars as a function of color. The calibrated 4Shooter reference image photometry is shown in $V/V-I$ and $V/B-V$ color-magnitude diagrams (CMDs) in Figure 3.




Fig. 2.— V , $B - V$ and $V - I$ residuals of Landolt (1992) standard stars as a function of color for the adopted transformation equations.

The zero points of the light curves of variable stars from the other datasets were adjusted to the 4Shooter data by adding an offset. It was computed as the mean offset for stars within a radius of 300 pixels around the variable, with the rejection of > 3 sigma outliers. The color term was not taken into account.

Additional offsets, if necessary, were applied between overlapping datasets for all variables, ie. MDM-FLWO and 4Shooter-KPNO, and between all datasets in case of periodic variables. The offsets are probably due in large part to the difference in color terms between the chips. Such offsets were added in 14% of the cases and typically were ≤ 0.06 mag. For some variables the offsets were large (up to 0.5 mag) because of blending with nearby stars in the datasets with inferior seeing.

3.3. Astrometry

The transformation from rectangular to equatorial coordinates was derived using 997 transformation stars from the USNO A-2 catalog (Monet




Fig. 4.— The distribution of preliminary cluster membership probabilities for 32 out of 51 variable stars, kindly supplied by Dr. Kyle Cudworth.

et al. 1998) for the 4Shooter *V*-band template. The average difference between the catalog and the computed coordinates for the transformation stars was $0''.14$ in right ascension and $0''.12$ in declination.

3.4. Variability Search

We used the standard ISIS procedure to search for variable stars. For every pixel it computes a median of absolute deviations on all subtracted images and performs a simple rejection of cosmic rays and defects. The result is stored as an image, where the variables are then identified as bright peaks. We used this method to search for new variables in all filter and dataset combinations.

We also extracted the light curves for all stars detected by DAOPHOT (Stetson 1987) on the template frames and searched them for variability using the index *J* (Stetson 1996), as described in more detail by M02.

To search for periodicity we used the analysis of variance statistic (Schwarzenberg-Czerny 1996).

4. VARIABLE STAR CATALOG

We have confirmed the variability of 37 out of 44 previously known variables within our field of view. Of the 7 variables not recovered, three are detached eclipsing binaries with no eclipses observed by us. Many of the variables (V29-V67) were discovered in higher S/N 900 s *R*-band observations (M02), hence it was anticipated that some fainter and/or lower amplitude objects might not be recovered. We have also discovered seven new variables.

The properties of the 51 variable stars are discussed in subsections 4.1 - 4.5. Tables 2-4 list the following parameters for the variables: the identification number, right ascension, declination, period and its error estimate (for eclipsing and periodic variables), *V*, *I* and *B*-band magnitude (maximum for non-periodic and eclipsing variables, flux weighted mean for periodic), variability amplitudes in each band (semiamplitudes for periodic variables) and preliminary cluster membership probabilities, kindly provided to us by Dr. Kyle Cudworth. Table 2 contains an additional column with comments on the variability type.

The variables are plotted on the CMD in Figure 3. Non-periodic variables are indicated by triangles, eclipsing binaries by squares and other periodic variables by circles. Filled symbols indicate membership probability above 75%, open symbols – below 75%, and skeletal – no membership probability estimate. The light curves, mostly in *V*, for a selection of variables are shown in Figures 7-3.4. The *BVI* light curves and finder charts for all variables and machine-readable versions of tables 2-4 are available for download via ftp from [cfa-ftp.harvard.edu](ftp://cfa-ftp.harvard.edu/pub/bmochejs/NGC6791) in the `/pub/bmochejs/NGC6791` directory.

The distribution of the cluster membership probabilities for the variables is shown in Figure 4. Proper motion data, available for 32 out of 51 variables, are in most cases conclusive in resolving the question of their cluster membership. Of those variables, 21 are likely cluster members and 8 are not, with probabilities in excess of 75% and below 25%, respectively. There are only three variables for which the membership is very uncertain ($25\% < P < 75\%$).




Fig. 3.— The $V/V-I$ and $V/B-V$ color-magnitude diagrams. Non-periodic variables are indicated by triangles, eclipsing binaries by squares and periodic variables by circles. Filled symbols indicate membership probability above 75%, open symbols – below 75%, and skeletal – no membership probability estimate.

4.1. Cataclysmic variables

Our main interest was to study the long-term behavior of the two cataclysmic variables B7 (a.k.a. V15) and B8. These are unique objects, as only three such stars have been found and confirmed in all open clusters. B7 is a cluster member, as shown by proper motion data, while for B8 no estimate is available. Both were spectroscopically confirmed to be CVs by Kaluzny et al. (1997).

During our monitoring the brighter cataclysmic variable B7 underwent a large drop in brightness, by 3 mag in V and 2 mag in I , and three outbursts of about 0.5-1 mag. Smaller amplitude variations, as well as long-term trends are also present in the light curve.

The large decline in the MDM data and intermediate brightness at other times would suggest that B7 may be an VY Scl type nova-like (NL) variable, which vary little about their mean magnitude, but occasionally fall in brightness by > 1 mag (Warner 1995). Another possibility is that it is a Z Cam type dwarf nova (DN), which exhibit periods of standstill at irregular intervals. One no-

table difference between these two types of CVs is that Z Cam DNe are brighter by 0.5-1 mag in outburst than in the standstill phase, while the apparent outbursts in VY Scl type NL variables reach at most the high state brightness. We have observed the variable at minimum light only once, when it fell from the mean brightness by 3 magnitudes, so we are unable to classify B7 based on this criterion. The 4Shooter and KPNO data phase rather well with a period of ~ 25.4 days, which is compatible with the typical cycle length of 10-30 days for a Z Cam type DN (Sterken & Jaschek 1996).

The other cataclysmic variable, B8, displayed several large outbursts about 3 magnitudes in amplitude, as well as a few smaller ones. The MDM and AndyCam data display incoherent variability, possibly due to flickering.

The variability of B8 is reminiscent of an SS Cygni type dwarf nova (Sterken & Jaschek 1996). The 4Shooter and KPNO light curve phases reasonably with a period of ~ 17.73 days, which if real, is shorter than the typical range of 30-100 days for these variables.




Fig. 5.— V-band light curves for 10 selected long period and non-periodic variable stars. Each dataset is plotted with a different shade of grey. The top window illustrates the distribution in time of the 15 sub-windows plotted for the variables.




Fig. 6.— Phased V -band light curves for the cataclysmic variables B7 and B8 (4Shooter and KPNO data only).

4.2. Newly Discovered Variables

We have identified a total of seven new variables in the field of NGC 6791.

V68, located 4 magnitudes above and 0.3 mag to the blue of the turnoff, is a periodic variable, with a period of ~ 15 days. It may be an RV Tau variable with twice the period.

V69 and V70 are very red, bright variables. V69 seems to be periodic, with a period of about 43 or 86 days. The period determinations are very uncertain and different for the V and B bands due to the star's long period and sparse sampling – the star was unsaturated only on some of the short exposures. V70 is an irregular variable with an amplitude of ~ 0.4 mag in V . The preliminary proper motion data are inconclusive as to the cluster membership of these variables, yielding probabilities of 47% for V69 and 62% for V70.

V71 is also irregular, with a V -band amplitude of ~ 0.3 mag. On the CMD it is located near the base of RGB. Proper motion data gives a membership probability of 55% for this variable.

V72 and V73 are located about 2 mag above the cluster turnoff. They exhibit irregular variability with V -band amplitudes of about 0.04 and 0.08 mag, respectively. When the observations between HJD 2452099 and 2452204 are excluded, the V -band light curve for V73 can be phased with a

period of ~ 34 days (Figure 7). Its membership probability is low (20%).

V74 exhibits irregular brightness variations with an amplitude of 0.04 mag in V . On the CMD it is located just blueward of the red clump. It is very likely a cluster member.

4.3. Previously known long period and non-periodic variables

During previous investigations the cluster was found to possess six other long period and non-periodic variables: V13, V19, V62 and V65-V67. V13 and V19 show irregular variability with sudden rises in brightness by 0.1-0.2 mag in V . V13 is a cluster member. We confirm the variability of V62, but due to its faintness and poor quality light curve, we are unable to fathom its nature. V65 seems to be periodic, with a period of ~ 11.3 days, when the MDM dataset and May 2002 4Shooter run are excluded (Figure 7). It does not belong to the cluster. Of the two variables on the RGB, V66 and V67 (confirmed member), the first is more or less periodic, with a period of ~ 49 or ~ 99 days (Figure 7), while the second seems to be irregular.

Of the five variables classified in previous investigations as detached eclipsing binaries with unknown periods, V10, V18, V20, V21 and V60, we have detected eclipses only in V18 and V20, but were unable to determine convincing periods. V10




Fig. 7.— Light curves for six selected eclipsing binaries (top two rows) and for three variables classified as long-period (bottom row). Each dataset is plotted with a different shade of grey. The light curves are in the V-band, with the exception of V68, which is in the I-band.

seems to have decreased its brightness by 0.1 mag between the 1999 and 2001 seasons. V60 also displays a long-term trend, especially visible in the 4Shooter data. Based on proper motions, V18 and V60 are most likely cluster members.

4.4. Previously known eclipsing binaries

Thanks to the long time span of our observations, we have derived more precise periods for the eclipsing binaries B4, V1, V3-V7, V9, V11, V12, V16 and V38. We have confirmed the variability of V2 and V14 with the previously reported periods, but were unable to improve their determinations. Variables V3-V5, V9, V16, V38, V12 are likely cluster members, while B4, V6 and V14

are non-members.

Figure 8 shows a plot of apparent distance modulus $(m - M)_V$ as a function of period for W UMa type contact binaries V1-V7. The absolute V magnitude was derived from the calibration of Rucinski (2000), based on B-V and V-I colors (left and right panel, respectively). A reddening of $E(B-V)=0.1$ (Chaboyer et al. 1999) was adopted. Filled circles indicate membership probability above 75%, open symbols – below 75%, and skeletal – no membership probability estimate. Of the three variables, which are high probability cluster members based on proper motions, V3 and V4 are within 0.7 mag of the cluster distance modulus $(m - M)_V = 13.42$ (dotted line; Chaboyer et




Fig. 8.— Apparent distance modulus as a function of period for the contact binaries V1-V7. The absolute V magnitude was derived from the calibration of Rucinski (2000), based on B-V and V-I colors (left and right panel, respectively). Filled circles indicate membership probability above 75%, open symbols – below 75%, and skeletal – no membership probability estimate.

al. 1999) and V5 is about a magnitude brighter than predicted from the calibration. V6, which is a non-member, based on its proper motion, is located ~ 2.5 magnitudes in front of the cluster. Of the three variables with no membership probability estimate, V2 and V7 are probably located at the distance of the cluster, while V1 seems closer by ~ 2 magnitudes.

V29, which was classified in M02 as an eclipsing binary with a period of ~ 0.4 days, shows a long-term rise in brightness with an amplitude of ~ 0.5 mag during our observations (the large amplitude in Table 3 is mainly due to scatter in the AndyCam data). The eclipses were not recovered.

For V31 and V32 we did not detect any coherent variability at the periods determined in M02. These periods are also not confirmed by the additional extensive R -band photometry for this cluster, collected by the authors (Mochejska et al. 2003, in preparation), where the variables only show weak long-term trends. V31 is a probable cluster member and V32 is most likely not.

V33, previously classified as an EB with a period of ~ 2.37 days, exhibits long term brightness variations. In the 4Shooter data we have observed a recovery by 0.15 mag to its maximum observed brightness. It also displays quasi-periodic variations with roughly the period reported in M02. This variable is most likely not a member of the cluster.

The light curve for V37 is very noisy and we were not able to recover the previously determined period. In the M02 data the variable underwent a 0.1 mag rise in brightness. It appears to have experienced a rapid increase in brightness by 0.6 mag at the beginning of the AndyCam run.

4.5. Previously Known Other Periodic Variables

We have confirmed the previously determined periods of V41, V48, V54, V56 and V58. We have confirmed the variability of V17, V45, V46 and V52 with previously derived periods, but were unable to improve their determinations. V45 seems

to exhibit long term brightness variations, besides short term variability. We were unable to confirm the periods of V42, V51 and V53, determined by M02. V42 seems to have steadily decreased its brightness by ~ 0.6 in V . Variables V17, V41, V42, V48, V53, V56, V58 are high probability cluster members, and V45 is most likely a non-member.

Based on its period and location on the color-magnitude diagrams, V17 might be a member of the newly proposed class of variable stars termed “red stragglers” (Albrow et al. 2001) or “sub-subgiant stars” (Mathieu et al. 2003). To date, six such stars have been found in 47 Tuc (Albrow et al. 2001) and two in M67 (Mathieu et al. 2003). Thus far, the origin and evolutionary status of these stars remains unknown.

5. CONCLUSIONS

In this paper we have presented *BVI* photometry for 51 variable stars in the field of the open cluster NGC 6791. Over 6 years of monitoring of this cluster we have collected 391 long and 74 short exposures in V , 140/89 in I and 65/7 in B . The light curves of the variables were extracted using the ISIS image subtraction package (Alard & Lupton 1998).

The main target of our observations were the two cataclysmic variables B7 and B8. Their light curves show outbursts characteristic of CVs. We have identified possible cycle lengths of 25 and 18 days for B7 and B8, respectively. We tentatively classify B7 as a VY Scl type nova-like variable or a Z Cam type dwarf nova. More observations of the variable at minimum brightness would help elucidate its nature. B8 is most likely an SS Cygni type dwarf nova.

We have also discovered seven new variables: two periodic and five long-period or non-periodic ones. In addition, we have extracted the light curves of 42 other previously reported variable stars. For 31 of them we confirm the variability period, if applicable, and type, given by M02 and for 17 we have derived new, more precise periods.

We would like to thank Gáspár Bakos, David Bersier, Martin Krockenberger, Lucas Macri, Dimitar Sasselov and Andy Szentgyorgyi for their help in obtaining the data and Alex Schwarzenberg-

Czerny for software to compute errors in the period determinations. We also thank Dr. Kyle Cudworth for kindly supplying us with his preliminary cluster membership data for the variables. We would also like to thank the referee, Ronald Gilliland, for a prompt and useful report, which significantly improved the paper. BJM was supported by the Polish KBN grant 5P03D004.21 and the Foundation for Polish Science stipend for young scientists. Support for BJM was provided by NASA through Hubble Fellowship grant HST-HF-01155.01-A from the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Incorporated, under NASA contract NAS5-26555.

REFERENCES

- Alard, C., Lupton, R. 1998, *ApJ*, 503, 325
- Alard, C. 2000, *A&AS*, 144, 363
- Albrow, M. D., Gilliland, R. L., Brown, T. M., Edmonds, P. D., Guhathakurta, P., & Sarajedini, A. 2001, *ApJ*, 559, 1060
- Chaboyer, B., Green, E. M., Liebert, J. 1999, *AJ*, 117, 1360
- Friel, E. D., Janes, K. A., Tavaréz, M., Scott, J., Katsanis, R., Lotz, J., Hong, L., & Miller, N. 2002, *AJ*, 124, 2693
- Gilliland, R. L. et al. 1991, *AJ*, 101, 541
- Kaluzny, J., Ruciński, S. M. 1995, *A&AS*, 114, 1 (KR95)
- Kaluzny, J., Ruciński, S. M. 1993, *MNRAS*, 265, 34
- Kaluzny, J., Stanek, K. Z., Garnavich, P. M. & Challis, P. 1997, *ApJ*, 491, 153
- Landolt, A. U. 1992, *AJ*, 104, 340
- Mathieu, R. D., van den Berg, M., Torres, G., Latham, D., Verbunt, F., & Stassun, K. 2003, *AJ*, 125, 246
- Mochejska, B. J. & Kaluzny, J. 1999, *Acta Astronomica*, 49, 351
- Mochejska, B. J., Kaluzny, J., Stanek, K. Z., Sasselov, D. D., & Szentgyorgyi, A. H. 2001, *AJ*, 121, 2032
- Mochejska, B. J., Stanek, K. Z., Sasselov, D. D., & Szentgyorgyi, A. H. 2002, *AJ*, 123, 3460 (M02)

- Monet, D., et al. 1998, USNO-A2.0, (U.S. Naval Observatory, Washington DC).
- Peterson, R. C. & Green, E. M. 1998, ApJ, 502, L39
- Ruciński, S. M., Kaluzny, J., Hilditch, R. W. 1996, MNRAS, 282, 705 (RKH)
- Schwarzenberg-Czerny, A. 1996, ApJ, 460, L107
- Sterken, C. & Jaschek, C. 1996, Light Curves of Variable Stars, A Pictorial Atlas, ISBN 0521390168, Cambridge University Press, 1996
- Stetson, P. B. 1996, PASP, 108, 851
- Stetson, P. B. 1987, PASP, 99, 191
- Szentgyorgyi, A. H. et al. 2002, in preparation
- Warner, B. 1995, Cataclysmic Variable Stars (Cambridge: Cambridge University Press)

TABLE 1
LOG OF OBSERVATIONS FOR NGC 6791

Telescope	CCD	" / pix	B	B _{sh}	V	V _{sh}	I	I _{sh}
MDM 1.3m	Wilbur	0.32	-	-	79	29	50	39
KPNO 2.1m	T2KA	0.30	6	7	28	16	16	11
FLWO 1.2m	AndyCam	0.32	40	-	139	-	40	6
FLWO 1.2m	4Shooter	0.33	19	-	145	29	34	33
Total			65	7	391	74	140	89

TABLE 2
LONG PERIOD OR NON-PERIODIC VARIABLES IN NGC 6791

ID	$\alpha_{2000}[^h]$	$\delta_{2000}[^\circ]$	B_{max}	V_{max}	I_{max}	A_B	A_V	A_I	MP [%]	Comments
B7	19.352056	37.799038	18.384	17.486	17.536	0.505	3.623	2.511	98	CV
B8	19.343262	37.747878	19.095	18.430	18.331	2.729	4.545	2.841	...	CV
V10	19.353280	37.799508	20.469	19.578	18.328	0.786	0.402	0.273	...	DEB*
V13	19.347147	37.728627	15.617	13.986	10.704	0.051	0.171	0.330	98	Irr
V18	19.347051	37.769253	18.612	17.658	16.711	0.111	0.524	0.446	90	DEB
V19	19.347874	37.764148	16.715	15.099	13.936	0.155	0.249	0.200	...	Irr
V20	19.348416	37.759655	18.295	17.295	16.332	0.251	0.377	0.234	...	DEB
V21	19.349261	37.760280	18.404	17.465	16.567	0.052	0.139	0.072	...	DEB*
V60	19.350193	37.762528	19.576	18.614	17.556	0.269	0.270	0.138	91	DEB*
V62	19.350848	37.731072	20.234	19.180	18.118	0.161	0.318	0.119	...	
V65	19.347909	37.791823	17.256	16.225	15.120	0.033	0.056	0.039	0	P $\sim$ 11.3 ^d ?
V66	19.352340	37.748702	17.174	15.886	14.529	0.184	0.308	0.190	...	P $\sim$ 49 ^d , 99 ^d ?
V67	19.351021	37.801037	18.170	16.851	15.542	0.113	0.180	0.103	92	Irr
V70	19.342278	37.739207	16.330	14.777	10.581	0.295	0.397	0.277	62	Irr
V71	19.352913	37.723582	18.433	17.252	16.033	0.154	0.307	0.134	55	Irr
V72	19.351705	37.694531	15.925	15.111	14.230	0.029	0.045	0.025	...	Irr
V73	19.348047	37.781267	15.728	14.833	13.871	0.036	0.080	0.051	20	P $\sim$ 34.5 ^d ?
V74	19.352001	37.743069	15.921	14.654	13.353	0.025	0.037	0.060	91	Irr

NOTE.—CV – cataclysmic variable; DEB – detached eclipsing binary; Irr – irregular; P – period.
* No eclipses observed.

TABLE 3
ECLIPSING BINARIES IN NGC 6791

ID	$\alpha_{2000}[^h]$	$\delta_{2000}[^\circ]$	P [d]	σ_P [d]	B_{max}	V_{max}	I_{max}	A_B	A_V	A_I	MP [%]
V1	19.346558	37.742248	0.2676758	1e-07	17.026	16.175	15.193	0.451	0.423	0.328	...
V2	19.354874	37.766763	0.2732310*	8e-07	20.672	19.572	18.468	0.376	0.432	0.286	...
V6	19.350752	37.813589	0.2790259	2e-07	16.178	15.398	14.602	0.106	0.117	0.106	0
V5	19.346260	37.813300	0.3126604	6e-07	18.062	17.167	16.250	0.107	0.073	0.113	98
V3	19.354377	37.769393	0.3175443	9e-07	19.471	18.524	17.414	0.255	0.188	0.132	78
V4	19.348396	37.806629	0.3255937	6e-07	18.689	17.727	16.725	0.136	0.139	0.128	98
V29	19.354796	37.751465	0.436533*	2e-06	20.971	19.757	18.340	0.483	1.024	0.329	...
V7	19.340273	37.821882	0.487558	7e-06	18.634	17.674	16.726	0.212	0.371	0.212	...
B4	19.353583	37.764288	0.796993	5e-06	17.805	17.812	17.999	0.075	0.089	0.142	4
V11	19.342581	37.804616	0.883053	2e-06	20.386	19.349	18.295	0.502	0.803	0.428	...
V12	19.345259	37.849029	1.52321	2e-05	18.528	17.491	16.460	0.419	0.324	0.290	96
V31	19.350686	37.785924	1.53617*	3e-05	18.112	17.078	16.079	0.054	0.067	0.032	97
V32	19.341003	37.787297	2.09576*	7e-05	20.250	19.289	18.194	0.289	0.358	0.245	2
V33	19.344395	37.731804	2.36626*	4e-05	17.322	16.233	14.918	0.108	0.162	0.057	0
V9	19.346635	37.777072	3.18670	4e-05	18.391	17.177	15.841	0.293	0.440	0.249	82
V37	19.355069	37.851982	3.2199*	1e-04	20.905	19.324	16.899	0.761	0.830	0.244	...
V38	19.351025	37.768327	3.90904	8e-05	19.732	18.713	17.663	0.293	0.327	0.168	92
V16	19.352108	37.802682	4.4232	1e-04	18.707	17.745	16.772	0.186	0.173	0.095	96
V14	19.347685	37.756901	11.296*	2e-03	19.540	18.557	17.526	0.209	0.172	0.105	0

NOTE.—* Periods taken from M02.

TABLE 4
OTHER PERIODIC VARIABLES IN NGC 6791

ID	$\alpha_{2000}[^h]$	$\delta_{2000}[^\circ]$	P [d]	σ_P [d]	$\langle B \rangle$	$\langle V \rangle$	$\langle I \rangle$	A_B	A_V	A_I	MP [%]
V41	19.347492	37.806873	0.481631	3e-06	20.285	19.244	17.708	0.059	0.087	0.052	77
V42	19.350057	37.714872	0.506767*	8e-06	20.679	19.631	18.336	0.018	0.061	0.042	92
V45	19.346127	37.701679	4.578*	1e-03	17.964	17.062	16.027	0.007	0.006	0.009	0
V46	19.355274	37.798951	5.148*	1e-03	19.777	18.624	17.273	0.073	0.027	0.020	...
V48	19.352078	37.718526	5.8278	6e-04	18.432	17.539	16.568	0.044	0.021	0.018	96
V17	19.344137	37.817930	6.1752*	9e-04	19.172	18.013	16.720	0.018	0.023	0.019	88
V51	19.353380	37.748572	6.644*	3e-03	21.072	20.003	18.708	0.051	0.017	0.009	...
V52	19.355798	37.772039	7.055*	6e-03	18.329	17.497	16.591	0.012	0.008	0.004	...
V53	19.350232	37.743193	7.470*	3e-03	19.657	18.781	17.822	0.019	0.009	0.006	86
V54	19.355196	37.726818	8.333	1e-03	17.461	16.485	15.441	0.022	0.017	0.009	...
V56	19.345907	37.763587	11.863	3e-03	17.989	17.042	16.070	0.011	0.009	0.005	98
V58	19.354038	37.801260	13.007	2e-03	18.362	17.520	16.599	0.009	0.016	0.009	87
V68	19.341891	37.848689	15.26	1e-02	14.786	14.211	13.834	0.033	0.003	0.031	...
V69	19.354467	37.779617	43.26	5e-02	15.357	13.844	10.650	0.011	0.055	0.023	47

NOTE.—* Periods taken from M02.