

The Evolution of a Structured Relativistic Jet and GRB Afterglow Light-Curves

Pawan Kumar

Astronomy Department, University of Texas, Austin, TX 78731

Jonathan Granot

Institute for Advanced Study, Princeton, NJ 08540

pk@astro.as.utexas.edu, granot@ias.edu

ABSTRACT

We carry out a numerical hydrodynamical modeling for the evolution of a relativistic collimated outflow, as it interacts with the surrounding medium, and calculate the light-curve resulting from synchrotron emission of the shocked fluid. The hydrodynamic equations are reduced to 1-D by assuming axial symmetry and integrating over the radial profile of the outflow, thus considerably reducing the computation time. We present results for a number of different initial jet structures, including several different power-laws and a Gaussian profile for the dependence of the energy per unit solid angle, ϵ , and the Lorentz factor, Γ , on the angle from the jet symmetry axis. Our choice of parameters for the various calculations is motivated by the current knowledge of relativistic outflows from gamma-ray bursts and the observed afterglow light-curves. Comparison of the light curves for different jet profiles with GRB afterglow observations provides constraints on the jet structure. One of the main results we find is that the transverse fluid velocity in the comoving frame (v_t) and the speed of sideways expansion, for smooth jet profiles, is typically much smaller than the speed of sound (c_s) throughout much of the evolution of the jet; v_t approaches c_s when ϵ along the jet axis becomes of order a few (ϵ for large angular gradient of ϵ , $\psi \ll c_s$ while ψ is still large). This result suggests that the dynamics of relativistic structured jets may be reasonably described by a simple analytic model where ϵ is independent of time, as long as ϵ along the jet-axis is larger than a few.

Subject headings: gamma-rays: bursts { gamma-rays: theory

1. Introduction

The great advance in our understanding of Gamma-Ray Bursts (GRBs) in the last few years has largely resulted from the observation and modeling of afterglow radiation (emission observed for days to months after the end of a GRB, in the X-ray, optical and radio bands). The basic procedure for obtaining information about the explosion, such as the energy release, opening angle of the emergent jet, the density of the medium in the immediate vicinity of the GRB etc., is by comparing the observed afterglow light-curve with the theoretically calculated flux (Wijers & Galama 1999; Granot, Piran & Sari 1999b; Chevalier & Li 2000; Panaitescu & Kumar 2001a,b, 2002). Most works on GRB jets assume a homogeneous (or 'top hat') jet, where all the hydrodynamic quantities of the jet, such as its Lorentz factor and energy density, are the same within some finite, well defined, opening angle around the jet axis, and drop to zero at larger angles.

A comparison of theoretically calculated light-curves, under several simplifying assumptions described below (and assuming a 'top hat' jet), with observed light-curves in X-ray, optical, and radio bands for 8 GRBs, has led to a number of interesting results (Panaitescu & Kumar 2001b). Perhaps the most remarkable discovery is that the kinetic energy in the relativistic outflow is nearly the same, within a factor of 5, for the set of eight GRBs. A similar result has been obtained by Piran et al. (2001) through a method which requires fewer assumptions. Frail et al. (2001) have also found that the energy radiated in GRBs does not vary much from one burst to another. The opening angle for GRB jets is found to be in the range of 2-20 degrees, and the density of the external medium in the vicinity of GRBs is estimated to be between 10^{-3} and 30 cm^{-3} . Moreover, there is no firm evidence for the density to vary as inverse squared distance in all but one case (Price et al. 2002; Panaitescu & Kumar, 2002), which is surprising in light of the currently popular model for GRBs (the collapsar model).

The possibility that GRB jets can display an angular structure, i.e. that the Lorentz factor, Γ , and energy per unit solid angle, E , in the GRB outflow can vary smoothly as power laws in the angle θ from the jet axis, was proposed by Meszaros, Rees & Wijers (1998). Recently, in view of the evidence described above for a roughly constant energy in the gamma-ray emission and in the kinetic energy of the afterglow shock, it has been suggested that GRB jets might have a universal structure, and the differences in the observed properties of GRBs and their afterglows arise due to different viewing angles, θ_{obs} , w.r.t the jet axis (Lipunov, Postnov & Prokhorov 2001; Rossi, Lazzati & Rees 2002; Zhang & Meszaros 2002). In this interpretation, the jet break in the light curve occurs when the Lorentz factor along the line of sight, $\Gamma(\theta_{\text{obs}})$, drops to Γ_{obs}^{-1} , so that the jet break time, t_j , is determined by the viewing angle, θ_{obs} , rather than by the opening angle of the 'top hat' jet, as in the

conventional interpretation.

The calculation of light-curves from a shock-heated, collimated, relativistic outflow has been carried out by a few research groups (Rhoads 1999; Panaitescu & Meszaros 1999; Kumar & Panaitescu 2000; Moderski, Sikora & Bulik 2000; Granot et al. 2002). However, most of the works to date have been based on a simplified model for the jet dynamics and on a number of ad-hoc assumptions. All the above works assume a 'top hat' jet, and furthermore, most of them model the dynamics of the jet at times much greater than the deceleration time as uniform expansion at the sound speed or the speed of light (in the local rest frame of the shocked fluid) — the results are nearly the same for both of these cases. Similar simplifications were made in the recent work on a universal structured jet (Rossi, Lazzati & Rees 2002). An exception to this, is the work of Granot et al. (2001), where the dynamics of an initial 'top hat' jet were calculated using a hydrodynamic simulation, and the resulting light curves were calculated numerically. However, such hydrodynamic simulations are very time consuming, and difficult to apply to a structured jet, so that there is currently no rigorous treatment of the hydrodynamic evolution of a structured jet. In this paper we develop such a rigorous treatment for the dynamics of structured jets, which at the same time is not very time consuming and may become practical to include in fits to afterglow observations.

Another simplification made in previous works (including all the works mentioned above), and in the lack of a better alternative, is also made in this work, is that the strength of the magnetic field and the energy in the electrons are determined by assuming that the energy densities of the magnetic field and of the electrons are constant fractions of the internal energy density of the shocked fluid. It is unclear how some of the simplifying assumptions in the afterglow light-curve modeling affect the overall burst parameters and properties we have inferred as described above.

Some progress has been made recently toward understanding the generation of magnetic fields in relativistic collisionless shocks: the numerical simulations of Medvedev (2002) show that magnetic fields generated behind collisionless relativistic shocks via the Weibel instability (Medvedev & Loeb 1999) do not decay to very low values within a short distance behind the shock, as was previously thought (Guzinov 1999, 2001), but rather approach a finite value in the bulk of the shocked fluid behind the shock, which might be compatible with the values inferred from afterglow observations. Moreover, the modeling of GRB afterglow light-curves indicates that the energy fraction in electrons is close to equipartition (Panaitescu & Kumar, 2001b), hence the parametrization of electron energy does not appear to be a serious drawback for current models. Thus, at present, one of the biggest uncertainties in the afterglow modeling is the assumption of a uniform jet and the simplified jet dynamics. The purpose of this paper is to remedy this situation and develop a much more realistic

model for GRB jets. Fitting afterglow observations with light-curves that are calculated using a realistic jet model & dynamics may both constrain the structure of GRB jets (the initial distribution of the Lorentz factor and energy per unit solid angle as a function of the angle from the jet axis), and provide more accurate estimates for the physical parameters, which include the external density profile and the parameters describing the micro-physics of relativistic collisionless shocks.

In the next section (x2) we discuss the evolution of structured jets. In x2.1 we describe our hydrodynamical scheme where we begin from the full hydrodynamic equations, assume axial symmetry and integrate over the radial structure, thus reducing the problem to a set of one dimensional partial differential equations that are solved numerically. The initial and boundary conditions are outlined in x2.2, while results for some physically interesting cases are shown in x2.3. In x3 we describe the light-curve calculation and compare the results of hydro simulations with a simplified model. The main results are summarized in x4.

2. Jet modeling

We begin with a brief description of the uniform jet model, and then we describe in some detail the evolution of a more realistic, structured, jet and the afterglow light-curves resulting from emission by the shock heated medium swept up by the jet.

Most calculations of GRB light-curves have assumed that the properties of the relativistic outflow do not vary across the jet, and that the jet dynamics is described by a uniform lateral expansion in the comoving frame, at close to the speed of sound, c_s , which for a hot relativistic plasma is ≈ 0.3 times the speed of light, c . These assumptions drastically simplify the calculation of the evolution of the jet opening angle, θ_j , with time: the increase in the lateral size of the jet in comoving time t_{co} is $c_s t_{co}$, and so the change to its angular size is $d\theta_j = c_s t_{co} / r = (c_s/c) r = (r/c) dt_{co}$, or,

$$\frac{d\theta_j}{dr} = \frac{c_s}{c r} :$$

This equation, together with the energy conservation equation, describe the dynamics of a uniform relativistic disk or a jet. The implication of this equation is that the jet opening angle θ_j starts to increase when Γ drops below Γ_j^{-1} , and from that time onward the jet opening angle is roughly Γ_j^{-1} . A detailed discussion of the uniform jet dynamics and lightcurve calculation can be found in a number of papers (e.g. Rhoads 1999; Panaitescu & Meszaros 1999; Sari

et al. 1999; Kumar & Panaitescu 2000). Such a uniform jet with sharp well defined edges shall be referred to as a 'top hat' jet.

However, 2D hydrodynamical simulations of the evolution of a jet that is initially uniform within some finite opening angle (Granot et al. 2001) have shown that the lateral expansion of the jet is smaller than that predicted by the simple models described above. This suggests that the assumption of lateral expansion at close to the sound speed in the comoving frame, that is made in most simple jet models, is not valid. Nevertheless, the light curves calculated from these simulations show a sharp jet break in the light curves, similar to that seen in most afterglow observations, around the time $t \sim t_j^{-1}$.

2.1. Dynamics of Structured Relativistic jets

Clearly, it is unrealistic to assume that the outflow from GRB explosions will be uniform within some finite opening angle, outside of which the Lorentz factor, Γ , and energy per unit solid angle, \mathcal{E} , decrease very sharply (i.e. a 'top hat' jet). A more realistic situation is that the Lorentz factor (LF), the energy density etc. are smooth functions of the angle, θ , from the jet axis, and possibly also of the distance, r , from the central source. The causality consideration suggests that the outflow is unlikely to be uniform over large angles, and moreover it provides a limit on how rapidly initial inhomogeneities can be smoothed out. Let the LF of the shell after elapsed time t since the explosion, measured in the lab frame, be $\Gamma(t)$. The comoving time corresponding to this is $t' = t/\Gamma$, and the distance traversed by sound waves during this interval is $c_s t' = ct/\Gamma^2$. Therefore, the angular size of a causally connected region is $\Delta\theta = 1/\Gamma^2$, and inhomogeneities on an angular scale of $\theta_{inh} > 1/\Gamma$, if present initially, will persist; the inhomogeneities can be smoothed out only when the LF has dropped below Γ_{inh}^{-1} . As an example, the large angular scale inhomogeneities for a jet of opening angle 5° start to decrease only when the bulk LF has dropped below $\Gamma \sim 10$, or roughly one day after the explosion (as seen by the observer). It should also be noted that if one were to start with a uniform jet, or a top-hat profile for the LF or \mathcal{E} , the large gradient at the edge will decrease with time and the jet will develop angular structure (e.g. Granot et al. 2001).

The remainder of this section is devoted to the solution of the relativistic hydrodynamic equations to describe the evolution of jets from GRBs. The starting point is the relativistic fluid equations (e.g. Landau & Lifshitz, 1959):

$$T_{;\mu}{}^{\mu} = 0 \quad ; \quad T^{\mu\nu}{}_{;\nu} = w u^\mu + p g^{\mu\nu}{}_{;\nu} \quad ; \quad (1)$$

where $T^{\mu\nu}$ is the energy-momentum tensor for an ideal fluid, u^μ is the 4-velocity of the fluid, $g^{\mu\nu}$ is the metric tensor, p is the pressure and $w = \rho + p$ is the proper enthalpy density,

where ρ and e are the proper rest mass density and internal energy density, respectively, and $c = 1$ in our units. We use a spherical coordinate system and assume the flow possesses axial symmetry about the z -axis, i.e. $u_\phi = \partial_\phi = 0$. Under these assumptions the t ; r and θ components of equation (1) are

$$\frac{\partial}{\partial t} (w^2 - p) + \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 w^2 v_r) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \theta} (\sin \theta w^2 v_\theta) = 0 ; \quad (2)$$

$$\frac{\partial}{\partial t} (w^2 v_r) + \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 w^2 v_r^2) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \theta} (\sin \theta w^2 v_r v_\theta) + \frac{\partial p}{\partial r} \frac{w^2 v^2}{r} = 0 ; \quad (3)$$

$$\frac{\partial}{\partial t} (w^2 v_\theta) + \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 w^2 v_r v_\theta) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \theta} (\sin \theta w^2 v_\theta^2) + \frac{1}{r} \frac{\partial p}{\partial \theta} + \frac{w^2 v_r v_\theta}{r} = 0 ; \quad (4)$$

where v_r and v_θ are the r and θ components of the fluid velocity, and $\gamma = (1 - v^2)^{-1/2}$ is the Lorentz factor of the fluid. Assuming that pair production has a negligible effect on the rest mass density, baryon number conservation implies

$$(\rho u)_r = \frac{\partial}{\partial t} (\rho) + \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \rho v_r) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \theta} (\sin \theta \rho v_\theta) = 0 ; \quad (5)$$

We assume an equation of state

$$p = (\gamma - 1)e = \frac{1}{\gamma} (w^2 - p) \quad \text{with} \quad \gamma = \frac{4 + 1}{3} ; \quad (6)$$

Equations (2)-(6) can be solved together to determine the structure and evolution of the outflow from GRBs. The computation time, for a 1 GHz clock speed computer, and for a modest resolution in r & θ coordinate of 100×1000 (in order to keep the error small in finite difference schemes) and 5000 time steps, is expected to take of order several hours to complete one run for one set of initial conditions; for comparison the 2-D relativistic jet hydrodynamics calculation of Miller and Hughes, reported in Granot et al. (2001), took several hours to days of computation time, for low to medium resolution runs, to follow the evolution for ~ 10 observer days, while an even longer computational time was required for the higher resolution runs. The successful modeling of light-curves of a single GRB to determine various parameters requires several thousand runs, and thus the computation time to model one GRB, using a 2-D code, is currently estimated to range between months to years. Using many processors simultaneously can help reduce the actual overall time required, but at any rate, this requires a great computational effort.

The computation time can be drastically decreased by reducing the problem to a 1-D system, by integrating out the radial dependence for all of the relevant variables, over the width of the outflow plus the swept-up material. The physical motivation for this is that jets in GRBs are in fact thin shells.¹ This procedure reduces the computing time drastically without introducing a significant loss of information as far as the emergent synchrotron emission is concerned; we find that the lightcurves from a relativistic spherical shock which has radial structure described by the self-similar Blandford-McKee (1976) solution is almost the same as in a model where the radial dependences have been integrated out and the shell thickness is taken to be zero (see Figure 5 of Granot, Piran & Sari 1999a).

The shock front is a two dimensional surface described by $r = R(\tau; t)$. The shocked fluid is concentrated in a thin shell of thickness $R - R = 4^{-2} R$ for a relativistic flow², and therefore it makes sense to integrate all the dependent variables, such as p, w, \dots , etc., over the width of the shell in the radial direction. We define quantities averaged over r , at a fixed and lab frame time t , as follows:

$$\bar{p} = \int_0^{Z_R} dr r^2 p ; \quad \bar{w}^2 = \int_0^{Z_R} dr r^2 p^2 ; \quad \bar{u} = \int_0^{Z_R} dr r^2 p u ; \quad (7)$$

$$\bar{u}_r = \int_0^{Z_R} dr r^2 p u_r ; \quad \bar{u}_r \bar{u} = \int_0^{Z_R} dr r^2 p u_r u ; \quad \bar{u}^2 = \int_0^{Z_R} dr r^2 p u^2 ; \quad (8)$$

$$\bar{M}_s \frac{dM_s}{d} = \int_0^{Z_R} dr r^2 ; \quad \bar{M}_0 \frac{dM_0}{d} = \int_0^{Z_R} dr r^2 \bar{M}_0 ; \quad (9)$$

where \bar{M}_s and \bar{M}_0 are the rest mass per unit solid angle of swept-up material and of the original ejecta, respectively, and \bar{M}_1 & \bar{M}_2 are dimensionless correlation coefficients, of order unity magnitude, which are taken to be independent of time. Integration of equation (2) times r^2 , over the radial interval corresponding to the width of the shell, when the shell is located at $R(t)$, yields

$$\frac{\partial}{\partial t} \bar{h}^{-2} + \frac{1}{\sin} \frac{\partial}{\partial} \frac{\sin u}{R} = \bar{M}_{ext}(R) R^2 \frac{\partial R}{\partial t} ; \quad (10)$$

¹At a distance r from the center of the explosion, the laboratory frame radial thickness of the ejecta plus the swept-up shock heated material moving with LF is $r = 4^{-2} R$, whereas its transverse dimension is r_j . Therefore the geometric shape of the system is that of a thin disk as long as $r_j \ll 4^{-2} R$.

²In fact, even as the flow becomes non-relativistic, we still expect $R = R_0 \tau^{0.1}$, as in the Sedov-Taylor self-similar solution, so that the thin shell approximation should still be reasonable.

where

$$\frac{\partial \bar{u}_r}{\partial t} + \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\frac{\sin \theta}{R} \bar{u}_r \right) - \frac{\bar{u}_r^2}{R} = 0 ; \quad (11)$$

An integration of equations (3) and (4) over the width of the shell gives

$$\frac{\partial}{\partial t} \bar{u}_r + \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\frac{\sin \theta}{R} \bar{u}_r \right) - \frac{\bar{u}_r^2}{R} = 0 ; \quad (12)$$

$$\frac{\partial}{\partial t} \bar{u} + \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\frac{\sin \theta}{R} \bar{u}^2 \right) + \frac{\bar{u}_r \bar{u}}{R} + \frac{1}{R} \frac{\partial}{\partial \theta} = 0 ; \quad (13)$$

and the closure relation, given below, is obtained by integrating the equation $\bar{u}^2 = 1 + u_r^2 + u^2$, times the pressure p , over the width of the shell

$$\bar{u}^2 - \bar{u}_r^2 = 1 ; \quad (14)$$

where $\bar{u}_r = \beta_3 \bar{u}$ is a constant factor; $\beta_3 = 1$ corresponds to the assumption that $\bar{u}_r = \bar{u}$.

In deriving these equations it was assumed that the ejecta moves with the shocked ISM, i.e. that the radial and the transverse components of the ejecta velocity are same as those of the swept up ISM. Under this assumption the mass continuity equation for the ejecta and the shocked ISM are respectively

$$\frac{\partial \bar{u}_0}{\partial t} + \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\frac{\sin \theta}{R} \bar{u}_0 \right) = 0 ; \quad (15)$$

$$\frac{\partial \bar{u}_s}{\partial t} + \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\frac{\sin \theta}{R} \bar{u}_s \right) = \frac{\partial R}{\partial t} ; \quad (16)$$

The velocity of the shock front is given by the shock jump conditions (Blandford & McKee 1976):

$$v_{sh} = \frac{1}{\beta_{ps} + 1} \frac{\beta_{ps} (\beta_{ps} + 1) + 2}{[\beta_{ps} + 1]^2} \quad (17)$$

where β_{ps} is the post shock LF. The shock jump conditions imply that in the rest frame of the fluid before the shock (which in our case is the lab frame), the direction of the velocity of the fluid just behind the shock is always perpendicular to the shock front. In order to

propagate the shock front in time, we assume that the average velocity of the shocked fluid is a good approximation for its value just behind the shock, and obtain

$$\frac{\partial R}{\partial t} = \frac{v_{sh}}{1 + (u = u_r)^2} : \tag{18}$$

Equations (10)–(18) are solved numerically with appropriate initial conditions (discussed below), to determine the evolution of the jet.

2.2. Initial & boundary conditions

The initial conditions are chosen at a lab frame time t_0 , after the internal shocks have ended, and before there is a significant deceleration due to the sweeping up of the external medium. We implement a number of different initial conditions, one of which is that the initial energy (including the rest mass energy) per unit solid angle, ϵ , and the initial Lorentz factor (minus 1) are power-law functions of r , outside of a core of opening angle θ_c . The initial energy per unit solid angle, $\epsilon(r; t_0)$, and the LF for the power-law model are

$$\epsilon(r; t_0) = \epsilon_0 r^a ; \quad \Gamma(r; t_0) = 1 + (\epsilon_0 - 1) r^b ; \tag{19}$$

where ϵ_0 and Γ_0 are the initial energy per unit solid angle and Lorentz factor at the jet axis, and

$$\Gamma = \frac{\Gamma_0}{1 + \frac{r^2}{c^2}} ; \tag{20}$$

Another initial condition we explore is a Gaussian profile for which $\epsilon(r; t_0)$ and $\Gamma(r; t_0)$ are proportional to $\exp(-r^2/2\sigma^2)$. The Gaussian jet profile was mentioned in Zhang & Meszaros (2001), however they did not calculate the jet dynamics or lightcurve for this case.

Equation (19), or its counterpart for the Gaussian case, are applied only as long as $\Gamma(r; t_0) > 1.1$. At larger angles, we assume a uniform outflow, the parameters of which are set by the continuity condition.

We assume that the velocity is initially purely in the radial direction, i.e.,

$$v_\theta(r; t_0) = 0 ; \quad v_r(r; t_0) = \frac{R}{1 + \frac{r^2}{c^2}} \Gamma(r; t_0) ; \tag{21}$$

while the initial radius is given by $R(r; t_0) = t_0 v_r(r; t_0)$.

The angular derivative of all dependent variables except \bar{u} is zero at the pole and the equator, whereas $\partial \bar{u} / \partial \theta$ at the pole is determined by the assumption of axisymmetry, and at the equator by the reflection symmetry; \bar{u} vanishes at the pole and at the equator.

2.3. Numerical Results

Equations (10)–(18) are solved using the two step Lax-Wendro scheme, to determine the evolution of the jet, for several different choices of initial conditions. The number of grid points in the angular directions is taken to be about 2000, and the time step is chosen to satisfy the Courant condition. The numerical solution respects global energy conservation to within 0.1%. We have looked into the dependence of the solution on the value of the dimensionless correlation parameters α_1 and α_2 , and find consistent solutions for α_1 between 0.8 and 1, and α_2 approximately between 0.9 and 1.2. Outside of this range of parameters the code is unstable and the solution unphysical, and the energy conservation is not satisfied. Inside this range the solution is not sensitive to the exact value of α_1 and α_2 .

The evolution of $\bar{u}(\theta)$, $\bar{v}(\theta)$, and $\bar{e}(\theta)$ are shown in figures 1-4 for (a;b) = (0,2), (2,0), (2,2), and for a Gaussian jet. Note that the transverse velocity in the comoving frame of the shocked fluid, $v^0 = \bar{v}$, is much less than the sound speed, $3^{-1/2}$, throughout much of the time, and approaches the sound speed only when the jet Lorentz factor on the axis has fallen to a value of order a few. Clearly, this result depends on the gradient of the initial LF or \bar{e} at the initial time, and therefore the transverse velocity is found to be largest for the Gaussian case, which has the highest gradient of all the models we have considered.

The small value for v^0 can be understood from equation (13). Ignoring the second order term in u , and noticing that the source term for u is the gradient of \bar{e} , we find that $\bar{v} \sim (\bar{e})^{1/2}$, where \bar{e} is the angular scale for the variation of \bar{e} or the energy density. Thus, we get an appreciable transverse velocity in the comoving frame only when $\bar{e} \sim 1$.

Another way of deriving this result is from the shock jump conditions, that imply that the velocity, v_{ps} , of the fluid just behind the shock in the rest frame of the fluid before the shock (the lab frame in our case) is perpendicular to the shock front. This implies that the angle between \hat{v}_{ps} and \hat{r} (i.e. $\hat{v}_{ps} \cdot \hat{r} = \cos \theta$) satisfies

$$\tan \theta = \frac{v}{v_r} = \frac{1}{R} \frac{\partial R}{\partial \theta} : \tag{22}$$

For a relativistic flow $v < 1$ so that $v \approx v_r \approx 1$ and $\tan \theta \approx v$. Thus we have $v \approx \frac{1}{R} \frac{\partial R}{\partial \theta}$, and since $R \approx (1 - \beta^2)^{-1/2} t$, this implies $v \approx \frac{3}{4} \frac{\partial \ln R}{\partial \theta}$, or $v \approx \frac{3}{4} \frac{1}{R} \frac{\partial R}{\partial \theta}$ and $u = v \approx \frac{3}{4} \frac{1}{R} \frac{\partial R}{\partial \theta}$, where θ is the angle over which \bar{e} varies appreciably. Therefore, this reproduces the result of the previous paragraph, as it is easy to show that the angular scale for the variation of \bar{e} and \bar{e} are similar. From the definition of \bar{e} (equation 7), we have $\bar{e} \approx \frac{pR^2}{R} \approx pR^3 \approx 4^{-2}$ and since the shock jump conditions imply $3p = e = 4^{-2} \bar{e}_{ext}(R)$, this gives $\bar{e}_{ext}(R) R^3 \approx 3$, i.e. \bar{e} is a power law in $R \approx (1 - \beta^2)^{-1/2} t$, and therefore varies over the same angular scales as \bar{e} .

The transfer of energy from small to large angles over the course of the evolution of the jet, from highly relativistic to mildly relativistic regime, is also found to be small (Fig. 3).

The Gaussian initial jet profile is a smooth and more realistic version of a 'top hat' jet, where the hydrodynamic quantities are roughly constant within some typical opening angle, and sharply drop outside of this angle (though they drop smoothly, and not in a step function as in the 'top hat' jet). We therefore expect the results for an initial Gaussian profile to be similar to those of an initial 'top hat' jet profile. The hydrodynamic evolution of the latter has been investigated using a 2D hydrodynamic simulation (Granot et al. 2001) and was found to be quite similar to our results, namely the lateral spreading of the jet was much smaller than the prediction of simple 'top hat' jet models, and there was very little lateral transfer of energy. The fact that our hydrodynamic results for an initial Gaussian profile are similar to the results of 2D hydrodynamic simulations of an initial 'top hat' jet profile, is very reassuring and gives us some confidence in our numerical scheme.

3. Light-curves

Once the jet dynamics and the pressure and density of the shocked fluid are known, the synchrotron plus inverse-Compton emissions are calculated from the fractional energies contained in the magnetic field and relativistic electrons which are parametrized by dimensionless numbers B and ϵ_e respectively. The electrons are assumed to be accelerated to a power law distribution of energies, $dN = d\epsilon_e / \epsilon_e^p$, promptly behind the shock, and then cool due to radiative losses and adiabatic cooling. The local emissivity, j^0_ν (the energy emitted per unit volume per unit time, frequency & solid angle in the comoving frame), is approximated by a broken power law, with breaks at the cooling frequency, ν_c^0 , the synchrotron frequency, ν_m^0 , and the self absorption frequency, ν_{sa}^0 .

Due to the curvature of the jet surface and its motion, photons arriving at some observed time t_{obs} were emitted at different space-time points $(r;t)$. The calculation of afterglow multi-wavelength light-curves takes into account appropriate integration over equal arrival time surface as given by the following equation for the flux density

$$F_\nu(t_{obs}; \hat{n}) = \frac{(1+z)^2}{d_L^2} \frac{d^4x j^0_\nu(r;t)}{[1 - v \cdot \hat{n}]^2} \quad (23)$$

where prime denotes quantities in the comoving frame of the fluid, \hat{n} is the direction to the observer (in the lab frame), ν is related to the observed frequency by the Doppler shift relation i.e. $\nu = (1+z) \nu' / (1 - v \cdot \hat{n})$, z and d_L are the redshift and luminosity distance of the burst, and d^4x is the Lorentz invariant 4-volume element. The observer time t_{obs}

is related to the lab-frame time and location of the source r by: $t_{\text{obs}} = t_r + r \hat{n}$, with $r \hat{n} = r(\cos \theta \cos \theta_{\text{obs}} + \sin \theta \sin \theta_{\text{obs}} \cos \phi)$.

The lightcurves for the four jet profiles described in 2.3 are shown in Figure 5, and the micro-physics parameters for the shocked gas can be found in the figure caption. The average thermal Lorentz factor of shock heated protons is given by $\bar{\gamma}_{\text{th}}(\theta; t) = 1 + \frac{1}{2} \left[\frac{v_s(\theta; t)}{c} \right]^2 = \left(\frac{v_s(\theta; t)}{c} \right)^2$, which turns out to be in very good agreement with the value one obtains from shock jump conditions.

The light curves for the Gaussian profile are rather similar to those for a 'top hat' jet, although the jet break for 'on axis' observers ($\theta_{\text{obs}} = 0$) is somewhat less sharp for the Gaussian profile, compared to a 'top hat' jet. The jet break is however still sufficiently sharp to be consistent with afterglow observations, in most cases.

For $a = 0$ and $b = 2$, the energy per unit solid angle (ϵ) is independent of θ , so that after the deceleration time, t_{dec} , the light curves for viewing angles, $\theta_{\text{obs}} < \theta_{\text{dec}}$, become the same as for a spherical flow with energy $4\pi \epsilon$; note that for this jet model $t_{\text{dec}} / t_{\text{obs}} = 8^{1/3} / 16^{1/3}$ and $t_{\text{dec}} / t_{\text{obs}} = 3^{1/6}$. These light curves do not show a jet break, and are therefore not compatible with afterglow lightcurves that have a break.

For $a = 2$ and $b = 0$ (i.e. $\epsilon / \theta^2 = \text{const}$, at initial time), there is a clear jet break for $\theta_{\text{obs}} \lesssim 2 \theta_c$. The reason for this is that in this case much of the observed flux comes from small angles around the line of sight and the sharp break results when $\theta < \theta_{\text{obs}}$. For $\theta_{\text{obs}} \gtrsim 7 \theta_c$ there is a flattening of the light curve just before the jet break, due to the contribution from the inner parts of the jet, which is not seen in the observational data. This feature provides some constraint on this jet model and is discussed in Granot & Kumar (2002).

For $a = b = 2$ (i.e. ϵ / θ^2 initially) the temporal decay slope before the jet break is steeper for small θ_{obs} and more moderate for large θ_{obs} , and the magnitude of the increase in the slope after the steepening of the lightcurve is larger for larger θ_{obs} . If such a correlation is found in the data, it would provide support for this jet profile. The jet break is smoother for small θ_{obs} , and sharper for large θ_{obs} , making it difficult to explain the sharp jet breaks and small inferred θ_{obs} (or opening angles for 'top hat' jets) that have been observed in quite a few afterglows.

It can be seen that for $a = b = 2$ and a small θ_0 ($\lesssim 200$ or less), the deceleration time, t_{dec} , is quite large for large viewing angles, so that the light curve has a rising part at $t < t_{\text{dec}}$ (Fig. 5). The fact that this is not seen in afterglow observations provides a lower limit on θ_0 or an upper limit on b , and thus can be used to constrain the structure of the jet. A more detailed analysis of the constraints that can be put on the jet profile from comparison

to afterglow observations is discussed in an accompanying paper (Granot & Kumar 2002).

The light-curves obtained for the hydro simulation of jets can be reproduced, quantitatively, by one or both of two simple and extreme models, that are described below. For more details on these models and their results we refer the reader to Granot & Kumar (2002). The two different simple models are referred to as model 1 and model 2.

In model 1, the energy per unit solid angle is assumed to retain its initial distribution, $(\epsilon; t) = (\epsilon_0; t)$. This represents the limiting case where there is very little lateral transport of energy. Model 2 attempts to make the opposite assumption, that is, it assumes the maximal averaging of ϵ over the angle θ that is consistent with causality. The latter is achieved by averaging ϵ over its initial distribution, over the range in θ out to which a sound wave could have propagated from the initial time t_0 . These two extreme assumptions are designed to bracket the expected range of possible behaviors for lateral energy transport. They are therefore expected to roughly cover the range of observed flux which a more rigorous treatment of the jet dynamics should give. In this sense they serve to quantify the uncertainties in the jet dynamics and light curves. For both models, the Lorentz factor is determined by energy conservation i.e.

$$\epsilon_0(\theta) (\epsilon; t) + \epsilon_s(\theta; t) (\gamma^2 - 1) = \epsilon(\theta; t): \tag{24}$$

4. Conclusion

We have carried out hydrodynamical simulations of a relativistic, collimated, axisymmetric outflow propagating into an external medium. For simplicity, we used a uniform density medium for the calculations presented in this work. However, the numerical scheme developed in this paper is good for any axially symmetric external density distribution (including power laws with the distance from the source, etc.). We have reduced the problem to a 1-D system of partial differential equations by integrating over the radial thickness of the outflow, at a fixed lab-frame time, thereby greatly reducing the computation time. The hydrodynamical results were used to calculate the synchrotron emission and lightcurves for a variety of observer angles w.r.t. the symmetry axis.

The model for GRB jets that is described in this paper is both rigorous and requires a very reasonable computational time, thus making it a useful tool for the study of GRB afterglows. In particular, it can be used in fits to afterglow observations, which can help constrain the jet profile, the external density profile, and the micro-physics parameters of

collisionless relativistic shocks.

We find that for jets with smoothly varying energy per unit solid angle and Lorentz factor, Γ , the maximum transverse fluid velocity in the comoving frame of the shocked fluid is typically substantially less than the speed of sound: the peak velocity is of order $v_{\perp} = (c/\Gamma)$, where θ is the angular scale for the variation of the energy per unit solid angle, or of Γ . Thus large transverse velocities, approaching the sound speed, are realized only when the energy density varies rapidly with θ or decreases to $(\Gamma)^{-1}$. For the jet profiles examined in this paper, the largest lateral velocity occurred for the Gaussian profile, for which the initial gradients were largest. In fact, in this case a shock wave in the lateral direction develops due to the steep initial angular profile. This did not occur for the other jet profile which had a more moderate angular dependence.

The energy per unit solid angle in the jet, E , is found to change very slowly with time for all four of the jet models we have analyzed, and to a reasonable approximation can be considered essentially constant in time until $t = 4R/c$, along the jet axis, has dropped to $4R/c$, which corresponds to about a week since the explosion in the observer frame (this is also the time when the transverse velocity is getting to be of order $0.2c$).

Therefore, a simple model where the energy per unit solid angle is taken to be time independent, and each element of the jet behaves as if it were part of a spherical flow with the same Γ , can serve as a useful approximation for the jet dynamics, as long as the jet is sufficiently relativistic. As can be seen from Figure 6, this simple model indeed seems to reproduce the light-curves obtained using the hydrodynamical simulations quite well. However, the simple model must necessarily breakdown when the transverse velocity becomes of order c , and the energy density is no longer constant. Unfortunately, this is also the regime when our hydrodynamical calculation becomes unstable.

We have calculated the observed light-curves in the R-band for several different jet angular profiles and viewing angles. We find that the light curves for a Gaussian jet profile are similar to those for a 'top hat' jet (as expected), and compatible with most observed jet breaks. The light curves for a jet with a constant energy per unit solid angle, E , but $\Gamma(t_0)$ decreasing with t , are similar to those for a spherical explosion, and thus not applicable to cases where we see a jet break in the light curve. For Jet profiles where initially $E \propto t^{-2}$ and Γ is either constant or $\propto t^{-2}$ do produce jet breaks in the light curves, and have the advantage that these models can reproduce the 'observed' narrow range for the total energy in GRB relativistic outflows (Panaitescu & Kumar, 2002; Piran et al. 2001). A companion paper (Granot & Kumar, 2002) discusses some preliminary constraints on jet profiles from qualitative comparison with observations.

JG thanks the support of the Institute for Advanced Study, funds for natural sciences.

REFERENCES

- Blandford, R D ., and McKee, C F . 1976, *Phys. Fluids*, 19, 1130
- Bloom, J.S., et al. 1999, *Nature*, 401, 453
- Bloom, J.S., et al. 2002, *astro-ph/0203391*
- Chevalier, R A ., & Li, Z.-Y . 2000, *ApJ*, 536, 195
- Fishman, G J., and Meegan, C A ., 1995, *Ann. Rev. A & A*, 33, 415
- Frail, D . et al., 2001, *ApJ*, 562, L55
- Galam, T J., et al. 2000, *ApJ*, 536, 185
- Goodman, J. 1997, *New Astronomy*, 2, 449
- Granot, J., & Kumar, P. 2002, submitted to *ApJ*.
- Granot, J., Piran, T ., & Sari, R . 1999a, *ApJ*, 513, 679
- Granot, J., Piran, T ., & Sari, R . 1999b, *ApJ*, 527, 236
- Granot, J., et al. 2001, in "GRBs in the Afterglow Era", eds. E. Costa, F. Frontera & J. Hjorth, Springer-Verlag Berlin, 312
- Granot, J., Panaitescu, A ., Kumar, P ., & Woosley, S. E . 2002, *ApJ*, 570, L61
- Gruzinov, A . 1999, *ApJ*, 525, L29
- Gruzinov, A . 2001, *ApJ*, 563, L15
- Kulkarni, S R . et al., 1999, *Nature* 398, 389
- Konigl, A ., & Granot, J., 2002, *astro-ph/0112087*
- Kumar, P ., & Panaitescu, A . 2000, *ApJ* 541, L9
- Landau, L D ., & Lifshitz, E M ., 1959, *Course of Theoretical Physics*, vol. 6, Fluid Mechanics
- Lipunov, V M ., Postnov, K A ., & Prokhorov, M E . 2001, *Astron. Rep.*, 45, 236
- Medvedev, M V ., 2002, in "Gamma-ray Bursts: the Brightest Explosions in the Universe"
- Medvedev, M V ., & Loeb, A . 1999, *ApJ*, 526, 697
- Meszáros, P ., Rees, M J., & Wijers, R A M J. 1998, *ApJ*, 499, 301
- Moderski, R ., Sikora, M ., & Bulik T . 2000, *ApJ*, 529, 151
- Norris, J., Marani, G . & Bonnell, J. 2000, *ApJ*, 534, 248

- Panaiteescu, A., & Meszaros, P., 1998, *ApJ*, 503, 314
- Panaiteescu, A., & Meszaros, P., 1999, *ApJ*, 526, 707
- Panaiteescu, A., & Kumar, P., 2001a, *ApJ* 554, 667
- Panaiteescu, A., & Kumar, P., 2001b, *ApJ* 560, L49
- Panaiteescu, A., & Kumar, P., 2002, *ApJ*, 571, 779
- Panaiteescu, A. & Meszaros, P. 1999, *ApJ*, 526, 707
- Piran, T., 1999, *Physics Reports*, 314, 575
- Piran, T., 2000, *Physics Reports*, 333, 529
- Piran, T., Kumar, P., Panaiteescu, A. & Piro, L. 2001, *ApJ*, 560, L167
- Priest, E. R., et al., 2002, *astro-ph/0203467*
- Reichart, D. E. 1999, *ApJ*, 521, L111
- Rhoads, J., 1999, *ApJ* 525, 737
- Rossi, E., Lazzati, D., & Rees, M. J. 2002, *MNRAS*, 332, 945
- Sari, R., Piran, T., & Halpern, J. 1999, *ApJ*, 519, L17
- van Paradijs, J., Kouveliotou, C., and Wijers, R., 2000, *Ann. Rev. A & A* 38, 379
- Wijers, R. A. M. J., & Galama, T. J. 1999, *ApJ*, 523, 177
- Zhang, B., & Meszaros, P. 2002, *ApJ*, 571, 876




Fig. 1. The different panels show the evolution of the Lorentz factor, $\bar{\Gamma}(\theta;t)$, the energy per unit solid angle, $\epsilon(\theta;t)$, and the transverse velocity in the comoving frame, $v_\theta/c = \bar{u}$, for a jet with an initially Gaussian profile, i.e. $\bar{\Gamma}$ & ϵ proportional to $\exp(-\theta^2/2\theta_c^2)$ at the initial time. The parameters are: $\theta_c = 0.035$ radian, $\epsilon(\theta=0;t_0) = 10^{53} = 4$ erg/sr, $\bar{\Gamma}(\theta=0;t_0) = 200$, and the density of the external medium is 10 particles per CC. At angles larger than about 65° when $\bar{\Gamma}$ drops below 1.1, the LF and ϵ are taken to be independent of θ . Note that the energy per unit solid angle does not change much with time except at large angles where it was small initially, and we see it increase with time. The sideways expansion of the jet can be seen in the bottom panel, which shows the edge of the "relativistic jet", i.e. the vertical line where the transverse velocity drops to zero, which is moving to larger angles with time; the jet edge, and its motion, can also be seen in the top left panel as a sharp drop in $\bar{\Gamma}$. The sharp jump in the lateral velocity and energy density may be understood as a formation of a shock wave in the lateral direction. The numerical scheme we use becomes unstable when $\bar{\Gamma}$ drops below 4 at $\theta = 0$ or $v=c$ becomes larger than about 0.4; the LF near the edge of the jet is close to 1.01 when the code becomes unstable. It should be noted that the transverse velocity depends on the gradient of the energy density in the transverse direction and on the value of LF locally and not along the jet axis. And so our result for the transverse velocity (that it remains below the sound speed throughout much of the jet evolution) is not compromised by the numerical instability. The lab-frame time increases monotonically from the curve with smallest $v=c$ to the curve with highest $v=c$ (the lower panel), and the time increases from the highest to the lowest curves in the top two panels.




Fig. 2. The different panels show the evolution of the Lorentz factor $\Gamma(\theta; t)$, the energy per unit solid angle, $\epsilon(\theta; t)$, and the transverse velocity in the comoving frame, $v_\theta/c = \bar{u}$, for a jet with initial $\Gamma = 1 + 199 = (1 + \beta^2 = \frac{2}{c})$ and $\epsilon = \epsilon_0 (1 + \beta^2 = \frac{2}{c})^{-1}$ with $\epsilon_0 = 10^{53} = 4$ erg/sr, and $\theta_c = 0.02$ radian; this initial profile corresponds to $a = 2$ & $b = 2$ in the notation of equation (19). The density of the external medium is 10 particles per CC. Note that the memory of the initial core angle, θ_c , is not erased with time, and in fact it remains unchanged until quite late times (see top left panel). Moreover, the energy per unit solid angle does not change much with time either, except at large angles where it was initially small, and we see it increase slightly with time. The transverse velocity v_θ^0 (see lower panel) remains quite small throughout much of the evolution of jet and even at late stages is about a factor of 3 smaller than the sound speed; the oscillations seen at early time in v_θ are almost certainly unphysical and numerical in origin.




Fig. 3. Same as figure 2, expect that $a = 2$ & $b = 0$ i.e. initially $\epsilon = \epsilon_0 (1 + \theta^2 = \frac{2}{c})^{-1}$ and $\Gamma(\theta; t_0) = 200$, with $\theta_c = 0.02$ radian. See caption for fig. 2 for other details. The minimum in v_θ at early times (the two lower curves on the bottom panel) is because v_θ is changing sign from negative values at small angles to positive values at larger angles and we have plotted v_θ/c . The reason for the negative velocity near the jet axis at early times is that in this model the deceleration time is largest at the pole ($R_{dec}; t_{dec} / \theta^{2=3} = (1 + \theta^2 = \frac{2}{c})^{-1=3}$); thus the pressure integrated over the shell thickness is close-to-zero at the pole and non-zero at large θ at times much less than the deceleration time scale at the pole. In this case a poleward transverse flow ensues. At $\theta > \theta_{dec}(t)$ the θ^{-2} profile dominates and induces a flow toward larger θ . At later time, much greater the deceleration time at the pole, the pressure integrated over the shell thickness, is indeed largest at the pole, as the referee mentioned, and the transverse velocity becomes positive everywhere. For $(a,b) = (2,1)$ model the transverse velocity is everywhere positive at all times as expected from this argument and we have varied this numerically.




Fig. 4. | Same as Figure 3, expect that $a = 0$ & $b = 2$, i.e. initially $(\dot{\theta}; \dot{t}) = 10^{53} = 4$ erg/sr, and $\bar{\Gamma} = 1 + 199 = (1 + \beta^2 = \frac{2}{c})$, with $\theta_c = 0.02$ rad. See caption of Fig. 2 for other details. Note that the jet becomes increasingly spherically symmetric with time; Γ is independent of θ up to an angle where shell deceleration has occurred, as in fact we expect when $\dot{\theta}$ is independent of angle.




Fig. 5. | Observed R-band light-curves for different viewing angles, θ_{obs} , w.r.t. the jet axis are shown for the four jet profiles presented in Figs. 1-4; the basic jet model parameters are shown in the top right corner of each panel (see equation 19 for the definition of a & b). The remaining model parameters are the same for all panels: $z = 1$, the energy fraction in electrons and magnetic field are respectively $f_e = 0.5$ & $B = 10^4$, the power-law index for electron distribution $p = 2.5$, and a constant external density $n = 10 \text{ cm}^{-3}$. The lightcurves were calculated using the hydrodynamic simulation results shown in figures 1-4 (see appropriate figure captions for the details of the jet models) and equation (23). For $a = b = 2$ the temporal decay slope before the jet break is more moderate for large θ_{obs} , and if γ_0 is not very large (200 in this figure) then the deceleration time for large θ_{obs} is rather large. For $a = 2$ and $b = 0$ the jet break is rather sharp (as seen in observations), but at $\theta_{\text{obs}} \approx 11.5^\circ$ & a few γ_0 there is a flattening of the light curve just before the jet break (which is not seen in observations).




Fig. 6. | A comparison of lightcurves obtained with hydro simulation of jet (solid line) and a simple jet evolution model (dotted curve) where the energy per unit solid angle is assumed to be time independent. The initial jet model is a power-law profile with $a = 2$, $b = 2$ and $\theta_c = 1.1^\circ$ (see equation 19 for the definition of a , b & θ_c), and the observer location w.r.t. the jet axis, θ_{obs} , is given in each panel. All the other parameters for the calculation are same as in figure 5 (see the caption for details). For clarity the observed flux for the simple model has been multiplied by a factor 2.