A post-W M A P perspective on in ation

Arthur Lue, Glenn D. Starkm an, $^{\mathtt{Y}}$ and Tanm ay $\mathtt{Vachaspat}^{\mathtt{Z}}$

Center for Education and Research in Cosmology and Astrophysics
Department of Physics
Case Western Reserve University
Cleveland, OH 44106{7079

Abstract

Recent results from the Wilkinson Microwave Anisotropy Probe have been called a corroboration, or even a con mation, of in ation. Yet, the results include features that require, at least, a signicant distortion of what is usually meant by in ation. At the same time, critics have leveled the charge that in ation is an arbitrarily pliable theory and is therefore beyond proof or disproof. This startling dissonance in attitudes toward in ation seems to have grown out of the lack of a clear fram ework with which to evaluate the in ationary paradigm. In this rhetorical pamphlet we reexam ine the in ationary paradigm, attempt to articulate explicitly how the paradigm and its descendant models are falsi able, and make a sober assessment of the successes and failures of in ation.

E-m ail: lue@ bifur.cw ru edu

^yE-m ail: glenn .starkm an@ cw ru .edu

^zE-m ail: tanm ay@ m onopole.cw ru .edu

I. IN TRODUCTION

The dawn of the 21st century has indeed yielded the promised golden age of modern cosmology. The wealth of observational data from both satellites and ground-based surveys provide an increasingly rened set of tools for probing and criticizing the increasingly coherent theoretical fram ework of the standard cosmological model: a hot big bang evolution of a universe led with cold dark matter, with an early period of in ation that provides atness and homogeneity in the observable Universe and which, at the same time, provides the source of primordial density uctuations from which all observed structure evolved.

The recent results from the rst year of data from NASA's Wilkinson Microwave Anisotropy Probe (WMAP) are a remarkable accomplishment, a tour deforce of fantastic and careful analysis. The NASA press conference announcing the results of WMAP claimed that the data provides a conmation, or at least a corroboration, of the inationary paradigm. This last phrase, the inationary paradigm, has given rise to considerable angst amongst cosmologists. The mantra that ination is not a theory, rather it is a paradigm, has been used by enthusiasts and detractors alike. Proponents claim that ination is a simple but powerful environment where one can study a large variety of models and answer a host of questions. Critics respond by questioning whether ination is really science under those circum stances, and assert that ination, as a paradigm rather than a theory, can be engineered to provide whatever result is necessary. Indeed, the claim that WMAP corroborates in ation merely commed the worst fears of ination detractors: how can one comma paradigm that can never be disproven?

The cosm ology community must surely demand that the pillars of its standard theoretical framework have me foundations in scientic principles: providing explanations of known information, one ering new predictions, and subjecting itself to falsic cation. Is in ation good science? Perhaps { what is clear is that the criticisms of in ation as a scientic paradigm are not entirely unjusticed. When wish to lay out a set of sober thoughts regarding in ation, both prohand con. Little in this discussion will be new. Consider this a rhetorical pamphlet rather than a paper, one where we attempt to collect and organize ideas that many have expressed, to give voice to the frustrations that many physicists and cosmologists have concerning the status of in ation as sound science, and to provide another perspective with which to continue the productive debate on the subject.¹

With apologies to our colleagues, given the nature of this document and the familiarity of the community with the subject matter, we have included no references.

A.An Allegory

Is the in ationary paradigm good science? By this we mean is it falsi able? A re there any principles or predictions that are inviolable? These are the stringent questions that must be asked of any scientic paradigm. But, it is worth contemplating an analogy before denouncing in ation.

Particle physics lays claim to a remarkable theoretical foundation, its Standard Model. This model, approximately thirty years old, has been tested to an exquisite degree and, by the standards of cosmology, holds up incredibly well. But, just as one can ask whether the in ationary paradigm is good science, one can as easily ask the same of the Standard Model. More accurately, we should ask whether the gauge principle is good science. Here, we view the gauge principle as the governing concept that all fundamental interactions are mediated by vector bosons that are universally coupled to fermionic matter, representing a perfectly respected gauge symmetry. This gauge principle arising out of quantum electrodynamics is the foundation for the Standard Model.

But is the gauge principle falsi able? What m predictions does it make? Just as for in ation, there are many models that are consistent with the paradigm, many gauge groups that may be considered, many variations on the theme. In the real world, one must take the rather cumbersome SU(3) SU(2) U(1) gauge group to explain all the data. Indeed, if the data were to be dierent, one would modify the gauge group or add more particles to explain every anomalous feature.

One can take this analogy even further. Taken in its simplest form, the gauge principle has de nite predictions. It must have massless gauge bosons for every gauge symmetry present. And while this prediction works extraordinarily well for electromagnetism, it doesn't work for the nuclear forces. The weak gauge bosons are not massless. One cannot even directly observe the gluons. Not all gauge symmetries are explicitly, or even approximately, respected.

Direct predictions of the sim plest manifestation of the gauge principle are categorically refuted by observation. A whole new system needs to be manufactured. Outrageous modications are made to the gauge paradigm such as the addition of a fundamental scalar Higgs boson with non-universal couplings and spontaneous symmetry breaking; and a non-perturbative realization of the gauge principle must be introduced for the color force in order to bring the gauge principle into line with observations. Should one then argue that the gauge principle is garbage? That it isn't science because one can modify it ad in nitum in order to the color away with the data? And yet the gauge paradigm is considered wildly successful. Why? Is this situation different from in ation?

B.Lessons

Particle physicists would be reluctant to characterize to the gauge principle as lacking the heft of real science, or being totally devoid of inviolable predictions, and therefore not falsi able. The answers to the provocative questions raised above are that, indeed, the gauge principle does have a set of inviolable principles: an exact (but possibly hidden) gauge sym metry, gauge bosons mediating the associated interactions, and universal couplings of those gauge bosons to matter. Each of these predictions is indeed con med by observation. All variants of the Standard Model, however baroque, must respect these principles.

In order to put in ation on the same footing as the gauge principle, we need to enum erate a similar set of inviolable principles. Put another way, we need to identify what makes in ation so appealing that it may suer many alterations. What are its inviolable predictions? What are its core principles? The frustration with in ation stems from the apparent scarcity of inviolable principles, thanks to the ingenuity of creative in ationary theorists, and the apparent scarcity of independent experiments with which to test the self-consistency of in ation in the conceivable future.

II. THE INFLATIONARY PARADIGM

W hat are the principles underlying an in ationary theory?

We start by de ning the classical in ationary paradigm as: accelerated expansion of an initially marginal super-horizon (or, super-Planck, if at t = 0) volume, proceeding for many doubling times (order 100), and ending everywhere (or at least over an exponentially larger super-horizon volume) with them alization and baryogenesis (su cient for successful nucleosynthesis). Implicit in this paradigm is some driving mechanism for the accelerated expansion and the appropriate initial conditions that would lead to it. In all realizations of which we are aware, the driving mechanism is some eld, usually referred to as the \in aton." Suitable initial conditions for the in aton are assumed, usually on the basis that all possible initial conditions are statistically realized. General relativity (GR) is taken to be the dynamics of spacetime.

This classical paradigm, which arises out of classical eld theory, must be promoted to a quantum paradigm. So long as we are interested in spacetime curvature scales much less than the Planck scale, we continue to treat gravity as classical; however, the in aton eld must be treated quantum eld theoretically? Here there are two levels of complexity which we denote the semiclassical in ationary paradigm and the quantum in ationary paradigm.

² A lso implicit has been a particular description of the vacuum state of the theory (the Bunch-D avies vacuum), extending possibly to trans-P lanckian energy scales (and hence sub-P lanckian length scales), although some researchers have begun to explore the robustness of this fram ework.

In the sem iclassical in ationary paradigm one is in the perturbative regime of the quantum theory and quantum uctuations can self-consistently be regarded as occurring against a background of the classical evolution of the in aton eld and the metric, at least over a range of length scales extending up beyond our current Hubble volume. This paradigm is the one appropriate to new in ationary and natural in ation models. Moreover, it is this paradigm that is in play whenever predictions of in ation are compared to observational data.

In the quantum in ationary paradigm, for at least some portion of the in ationary epoch, one is in the regime where backreaction of quantum—uctuations on the spacetime need to be taken into account. To do this properly, one would need to extend GR to include quantum elects. This paradigm is the one appropriate to eternal, stochastic or chaotic in ation. The progenitors of in ation have argued that the quantum paradigm is the most satisfying realization of the in ationary paradigm, especially to alleviate the tuning of initial conditions necessary to start in ation. In this scenario the Universe is bubbling with regions that are in ating. In ation never ends everywhere; nevertheless, there are pockets that stop in ating and subsequently thermalize. According to this quantum scenario, we live in one of the thermalized regions. In ation is therefore anthropic. Conditional probabilities for predictions are found with the condition that the thermalized region be inhabitable. The Universe is not hom ogeneous on the largest scales, but regions large enough to accommodate our visible Universe can be smooth enough.

The apparent simplicity of these paradigms makes in ation so attractive. Unfortunately, it also means that there are only a few generic features to characterize in ationary models of the Universe observationally or experimentally. Nevertheless, even these few ingredients do seem to have certain consequences:

A . H om ogeneous, Isotropic Entropy-Filled Universe.

That the accelerated expansion of the Universe ends everywhere is implicit in the sem iclassical paradigm. That it does so in the quantum paradigm is no less true, but much more subtle, incorporating generically the simultaneous truths that at any given place it eventually ends, but that it never ends everywhere, and the volume of space in which it has ended is vastly smaller than the volume in which it has not. Indeed, in this picture it is often justimed only anthropically why we do not inhabit a still-in ating region. Either way, we apparently must live in a region where the energy stored in the eld driving in ation, the in aton, was converted into other more prosaic forms of energy. The vast amount of in ationary expansion is followed in all generic models by a rapid injection of entropy and its thermalization (through either reheating or preheating). This is taken to be governed

by a Lagrangian density which is independent of space-time location. It is discult to put any measure on the predicted esciency of this process, but the reheat temperature must be high enough to allow nucleosynthesis.

In the sem iclassical paradigm, the vast in ationary expansion provides (almost) homogeneous initial conditions for entropy injection and them alization over some large length scale. This scale may however be limited (as in 4 theory) where, despite weak coupling (1) the sem iclassical approximation (= 1) fails on su ciently large scales. Thus, homogeneity su cient to accommodate the sem iclassical assumption over a moderate range of scales is a consequence of weak coupling.

In the quantum paradigm homogeneity seems to be an assumption that can be made self-consistently rather than a prediction. In this scenario, quantum uctuations can be large, though in ation might be quenched wherever this happens. Models exist in which the uctuations remain tamed.

B. Super-H orizon F luctuations

The inevitable quantum uctuations in the in aton eldwillbe stretched beyond the cosmic horizon and imprint them selves in the resulting energy density after reheating. Only after in ation stops and conventional big-bang evolution occurs will scales that left the horizon during in ation reenter the cosmic horizon. These uctuations thus appear super-horizon in scale. Unfortunately, there is no minimum predicted amplitude of scalar uctuations; their spectrum is model-dependent.

The same type of uctuations would be produced for any light (compared to in ationary Hubble scale), non-conformally-coupled eld, e.g., gravity waves. As with in aton uctuations, these eld uctuations will be super-horizon. However, unlike the in aton, these elds are not expected to carry the bulk of the Universe's energy density, and sifting for these particular signature elds may be challenging. The amplitude for super-horizon tensor (gravity-wave) uctuations is constrained from below by the requirement that the post-in ationary reheat temperature be larger than that necessary for nucleosynthesis. In principle this constraint oversa strictly falsi able prediction of the in ationary paradigm, though in practice the minimum amplitude is inaccessible for the foreseeable future.

C.Other Model-Independent Predictions

Of course, there are other predictions, such as the existence of in aton particles that should appear at the in ationary mass scale. These particles, however, may be extremely weakly coupled to conventional matter and may be dicult to observe, even if one had

access to such energies. Nevertheless, the in aton eld cannot be completely decoupled from standard model physics. A signi cant amount of reheating to conventional particles requires some amount of coupling. This coupling may in principle be exploited, putting in ation strictly within the regime of particle physics, and providing another avenue for the falsi cation of in ation. Unfortunately, unless the in ation energy scale is very low compared to the Planck scale (e.g. near energies of 1 TeV), this also remains an inaccessible possibility for the foreseeable future.

III.MODEL-DEPENDENT PREDICTIONS

Unfortunately, other predictions depend on the particular in ation model employed. As indicated earlier, the ingenuity of theorists has shown that the idea that the Universe can be hom ogenized with an early stage of accelerated expansion may be incorporated (with varying degrees of ease) in an overwhelm ingly diverse set of models. However, we may take the predictions made by the simplest models as a guide for what is more or less natural in an in ationary model.

We can imagine a scenario where hypothetical observers know very little about observational cosm ology except that the Universe is very old and led with matter. However, they have a great deal of understanding about the rest of physics, and in particular, have been led to believe that gravitation is intimately connected with the dynamics of spacetime and that GR should govern the evolution of the Universe. In so doing, they would have realized, as have we, that the age of the Universe, as determined from the ages of planetary and meteoroidalmaterial, is much greater than the only natural time scale of GR { the Planck time, and that the curvature scale of the Universe is much greater than the only natural length scale in GR { the Planck length. They might also have wondered where all the entropy in the Universe came from, and why, in particular, the total energy of everything they could see was much greater than the only naturalmass scale in GR { the Planck mass.

Faced with these problems { the age problem, the atness problem and the entropy problem { they might well have developed the beautiful paradigm of in ation: the idea that there was in the early history of the Universe an epoch of accelerated expansion driven by the energy density and negative pressure of the instantaneous vacuum state, which serves to atten the Universe, vastly increase the characteristic dynamical time scale of cosmology, and Ils the Universe with a relatively hom ogeneous and abundant \soup" of particles. In the absence of any substantial data, physicists in this world would turn to the most basic models of in ation to ascertain possible new predictions about cosmology.

The simplest versions of in ation involve a single scalar eld, m in imally coupled to gravity, with a potential polynomial in the eld, e.g., $V = {}^4$ where (x) is the in aton

eld, and is small. The in aton begins trapped in some state far away from the true vacuum, (t=0;x)=0 M_P , where M_P is the Planck mass. If 0 M_P $^{1=6}$, we are in the semiclassical paradigm. The in aton eld rolls slowly down the potential as the Universe engages in accelerated expansion. Eventually, the eld exits the slow-roll regime and coherently oscillates around the vacuum. This oscillation induces preheating and reheating to standard model particles, and the Universe subsequently evolves via a standard hot-big-bang model. If 0 M_P $^{1=6}$, we are in the quantum paradigm. The Universe begins in a stochastically in ating state but eventually transitions into a regime where (t) M_P $^{1=6}$ in some region; semiclassical behavior subsequently dominates. Evolution in this region proceeds as in the semiclassical paradigm.

A . Flat U niverse

In this simplest model, in ationary expansion attens the Universe beyond the ability of any likely experiment to discern a non-zero value for j 1j. Thus, the hypothetical cosmologists would conclude that j 1jshould be so small as to not be easily measurable.

One can see how this prediction can be easily avoided by looking beyond the simplest models. The original terrestrial (\old") in ationary models, in which in ation ended via a rst-order phase transition generically predicted that if we live in a single bubble of the true vacuum then the space-like hypersurfaces of constant curvature should be hyperbolic (<1). (When cosmological data suggested that indeed '0:3, this fact was used to argue that '0:1 1 was generic.) However, rst-order in ation (unless dressed up with double in ation, topologically-non-trivialmanifolds, or other complexications) fails to solve the suite of in ation-motivating cosmological problems. Moreover, if even in the simplest models, in ation can accommodate observably non-at universes by allowing in ation to turn o at exactly the correct number of e-foldings. Of course, this just-so possibility is often viewed as unpalatable and unnatural.

$$B. = < 1$$

In the sem iclassical regime, for an in ationary eld with a self-interaction potential V (), the amplitude of scalar uctuations (as opposed to vector or tensor modes) is

$$- \frac{V^{3=2}}{V^{9}M_{P_1}^{2}} : (3.1)$$

Speci cally, to nd the amplitude of uctuations on a particular scale, we evaluate the right-hand side of Eq. (3.1) at the value which held when that particular scale crossed out of the apparent horizon. It might seem that this easily could be much less than unity. However,

during slow-roll, there is a relationship between and the number of e-foldings until the end of in ation, N ,

$$N = \frac{2}{M_{Pl}^2}$$
: (3.2)

For the model V () = 4 , Eq. (3.1) m ay be recast as

$$-\frac{1}{k}$$
 1=2N $_{k}^{3=2}$; (3.3)

where = kj is the scalar uctuation amplitude of a given comoving wavenumber, k, where N_k is the number of e-foldings between when that scale left the inationary horizon and the end of ination. Those scales where = kj > 0 (1) actually probe the stochastic regime of the quantum in ationary paradigm, implying Eq. (3.3) is no longer valid.

We observe density uctuations in the Universe over a given range of com oving scales k whose N_k 100. Equation (3.3) then implies that even though may be small enough for weak-coupling to be self-consistent, density uctuations need not be small. For $=_k j = 1$, must be further ne-tuned; the smaller the observed uctuations, the more ne-tuned must be. A lternatively, one may venture into so-called natural in ation models, which exploit almost-symmetries (such as pseudo-goldstone modes or at directions in dynamically broken supersymmetry) to explain unexpectedly small density perturbations.

C.Adiabatic uctuations

Because the energy in the eld driving in ation is eventually converted into the thermal soup of radiation and matter—lling the Universe, the in—aton would be converted into uctuations in the cosm ic energy density, and thence, through the dynam ical response of the local geometry, into—uctuations in the metric, as well as the large-scale statistical distribution of matter in the Universe. Thus, the—uctuations would generically be adiabatic. In more complicated models of in—ation, however, the—uctuations can have a non-adiabatic component.

D.Gaussian uctuations

In the simplest in ationary models, the uctuations arise from the excitation of independent in aton modes. Therefore the statistics of each mode would be that of a Gaussian random eld. In more complicated in ationary models, it is seen that there can be small departures from Gaussianity.

E. (Very Nearly) Equal Power on All Scales

Equation (3.1) shows that = is a function only of V and V⁰. Since to realize a large number of e-folds of expansion V () must be very at, therefore the amplitude of uctuations generated on all scales should be nearly equal. The hypothetical cosmologists would therefore conclude that the spectrum of uctuations should be scale-free or very nearly so. In particular, unless the scale corresponding to the onset of in ation, or some other transitory event, just happens to have been stretched to a physically observable scale { less than the current horizon size yet larger than the scale on which non-linear dynam ics confuses the traces of the primordial uctuations { there should be no observable features in the primordial power spectrum that they would deduce when they some day make measurements of structure beyond their planetary system.

However, the detailed structure of the power spectrum depends on the exact form of the in aton potential, the potential can be tuned in such a way as to provide whatever power spectrum is necessary, within some broad constraints that slow-roll in ation require. It is no wonder why many cosmologists invariably point to this feature of in ation and regard it as dangerously epicyclic. That one can tune the spectrum with an arbitrarily pliable in aton potential to tmost any given spectrum is disturbingly unsatisfying.

IV.OBSERVATIONS AND EVALUATION OF THE PARADIGM

In our hypothetical scenario, eventually observational cosm ology as we know it would be revealed. We here sum marize current observations, and evaluate the in ationary paradigm in light of each piece of evidence.

1. A hom ogeneous, full U niverse. Measurements of the CMB probe primarily our past light cone, and mostly the surface of last scattering. A lthough O ccam's razor suggests that it is highly unlikely that we just happen to live at the center (within parts per billion by volume) of a spherically symmetric inhom ogeneous universe, direct observational probes of the interior of the light cone are harder to come by. However, observations of distant galaxies establish that element abundances are uniform across the Universe, suggesting that there were no large uctuations in the energy density or baryon number at the time of primordial nucleosynthesis. Having had ample time to investigate the details of the onset and dynamics of in ation, we may well be reluctant to claim that the homogeneity and isotropy of the Universe are really great successes of in ation since the onset of in ation in any particular patch of space requires that that patch be relatively homogeneous on super-horizon scales to begin with (although once it is, in ation can vastly in prove the homogeneity). Moreover, other theories

(such as variable speed of light and various braneworld scenarios) may also explain the homogeneity and isotropy, so these features are not terribly good discrim inators between theories. Consistent with the classical paradigm, the visible Universe has a very large entropy, S $^{\prime}$ 10^{87} .

2. Super-horizon uctuations. The observation of acoustic peaks in the angular power spectrum of the CMB and in particular, as discussed by the WMAP team, the anti-correlation between the temperature anisotropy and the E-mode polarization at 1 2 angular scales establishes that super-horizon scalar uctuations exist. This observation is a true cause of celebration for the in ationary paradigm. While other theories may also predict such uctuations, they really are a generic feature of all in ationary models. Tensor uctuations have not yet been observed. This sets a mildly interesting limit on the in ationary energy scale, but of course far above the minimum energy scale required by nucleosynthesis.

The rudim ents of the in ationary paradigm seem to hold up to scrutiny. However, as we commented, these are extremely limited, and lack a great deal of discriminatory power. What of predictions of the simplest models? How surprised would our hypothetical cosmologists be?

- 1. A at Universe. The discovery and clear denition of the rst peak in the angular power spectrum of the cosmic microwave background (CMB) established denitively that the Universe is at or nearly so, '1, in particular that 603 as had previously been widely considered. Analysis of the WMAP observations show that = 1:02 0:02. In the simplest models of in ation, which are the only ones we are considering here, is predicted to be unity to very high precision. This is in well with observation.
- 2. An extremely homogeneous Universe. The original discovery of the 2:7 K CMB radiation by Penzias and Wilson in 1965, was soon followed by electrons to measure any anisotropy in that background. However, it was not until 1991 that the instructions of the anisotropy was made by the Cosmic Background Explorer satellite. The long delay was due to the very small amplitude of the anisotropy, only parts per 105. Since then many experiments have measured this anisotropy and its properties. The netuning needed to achieve the observed = would concern our hypothetical cosmologists. Because we developed in ation with the foreknowledge that = 1, we have been more prepared to accept a priorith is netuning problem. As limits on = improved through the 1980's, the netuning grew ever more severe, but it did

so adiabatically, forestalling any increasing sense of concern.

- 3. A diabatic uctuations. All known observations are consistent with all uctuations being entirely adiabatic in nature. As reported by WMAP the to their data is not improved by adding any amount of isocurvature uctuations. This is good support for acausal generation of perturbations, and to invery well with the simplest models of in ation. So the consistency of adiabaticity is important, but the limits on non-adiabaticity remain weak. Also a number of in ationary models have been constructed that generate non-adiabatic uctuations.
- 4. Gaussian uctuations. No deviations from Gaussianity have been observed in the uctuation spectrum. The absence of any detected non-Gaussianity of the uctuations would likely be viewed as a relief, but hardly a coup, since very nearly Gaussian distributions are rather generic due to the central limit theorem; moreover, unless one knows what non-Gaussianity to look for, noting it is really like noting a needle in a very large haystack. It is again to be noted that there exist several in ationary models that predict non-Gaussian uctuations.
- 5. Lack of equal power on all scales. On scales characterized by 's from ten to several hundred, the angular power spectrum, as determined by many CMB experiments, and particularly by WMAP is nearly scale free. However, COBE-DMR found and WMAP has con imed that on angular scales greater than about 60, the two point angular correlation function of the CMB temperature uctuations nearly vanishes. The WMAP team has argued that the best tistandard CDM model is ruled out at the 99.85% con dence level based on comparisons of the observed C() with a Monte Carbo of 10⁵ realizations of the model. Mild adjustments of the model only improve that to a 99.7% exclusion. The absence of these correlations on large angular scales is a serious problem for in ation. This is not because there exist no in ationary models which accommodate it. Features in the in aton potential, two stage in ation, just-so in ation in a compact manifold, braneworld models, etc. all may hold promise of accommodating this data. However, unless such modications of eradditional testable predictions, they are, indeed, dangerously epicyclic.

V.CONCLUDING REMARKS

Post-W MAP statements have been made claiming that the predictions of in ation have been con med, and that in ation is a successful paradigm. However, careful consideration of the meaning of the term \in ationary paradigm "suggests that such statements are, at best, imprecise. Generic predictions of the in ationary paradigms depend on certain

assumptions that are rarely made explicit. Granting these assumptions, the essential predictions of homogeneity and isotropy, and the existence of super-horizon uctuations are indeed con med; however, only the latter is a post-in ation discovery.

Further implementation of the in ationary paradigm requires adopting a particular model. The simplest one—eld in ation models have met with limited success when confronted by new data. The Universe appears spatially—at, and—uctuations are adiabatic and Gaussian. However, the—uctuation amplitude is unnaturally small and are decidedly not scale-free on the largest angular scales. While the former requires only a—ne-tuning of Lagrangian parameters, the absence of large-scale power seems to demand models that are carefully designed. But, unless these new models yield testable predictions, this tack merely perpetuates the habits that in ation's critics abhor. Do we continue to accept in ation merely because there is no better alternative?

Science is not a democratic pursuit. It only takes one contradictory fact to consign a theory to the dustbin of history, or at least to take it o its pedestal and send it back to the workshop. On the other hand, when one poses a given paradigm, it always make sense to begin with the simplest incarnation of that paradigm. The degree to which a model must be engineered to reproduce the needed data should then be factored into a reassessment of the worth of the original idea. If a theory is repeatedly faced with contradictory facts which force a reengineering, at what point does it stop being good science? If this is to be the dawn of a new era of precision cosmology, it must involve not only precise determinations of an ever increasing number of new parameters, but also precision tests of the self-consistency of our theories which permit their dispassionate evaluation.

ACKNOW LEDGM ENTS

We are grateful to Serge W in itzki for clarifying remarks on the quantum in ationary scenario. This work is sponsored by DOE Grant DEFG 0295ER 40898 and CWRUO ce of the Provost.