# New SX Phoenicis Stars in the G lobular C luster M 53

Young-Beom Jeon<sup>1</sup>

Korea Astronom y Observatory, Daejeon 305-348, KOREA

Em ail: ybjeon@ boao.re.kr

M yung G yoon Læ

A stronom y Program, School of Earth and Environm ental Sciences, Seoul National

University, Seoul 151-742, KOREA<sup>2</sup> and

Department of Terrestrial Magnetism, Carnegie Institution of Washington, 5241 Broad

Branch Road, N.W., Washington, D.C.

Em ail: m glee@ astrog.snu.ac.kr

Seung-Lee K im

Korea Astronom y Observatory, Daejeon 305-348, KOREA

Em ail: slkim@kao.re.kr

and

HoLæ

Department of Earth Science Education, Korea National University of Education,

Choongbuk 363-791, Korea

Em ail: leehol190 boao.re.kr

Received \_\_\_\_\_; accepted \_\_\_\_\_

<sup>1</sup>A lso A stronom y P rogram, School of E arth and E nvironm ental Sciences, Seoul N ational U niversity, Seoul 151-742, KOREA

<sup>2</sup>Perm anent address

#### ABSTRACT

Through time-series CCD photom etry of them etal-poor globular cluster M 53, we have discovered eight new SX Phoenicis type stars (labeled from SXP1 to SXP8). All the new SX Phoenicis stars are located in the blue straggler star region of a color-m agnitude diagram of M 53. One of these stars (SXP2) is found to have very closely separated pulsation frequencies:  $f_1=f_2 = 0.9595$  where  $f_1$  and  $f_2$  are primary and secondary frequencies. This may be due to excitation of non-radial modes. Six of these SX Phoenicis stars are considered to be pulsating in the fundamental mode. They show a tight linear correlation for the fundamental mode for the period range of 1.36 < LogP [d] < 1.15: < V > = 3.010(0.262)LogP + 15.310(0.048) with an rm s scatter of 0.038, corresponding to  $< M_V > = 3.010\text{LogP}$  1.070 for an adopted distance modulus of (m M)<sub>V</sub> = 16:38 (Harris 1996).

Subject headings: G lobular clusters: individual (M 53 (NGC 5024)) | stars: blue stragglers | stars: oscillations | stars: variable stars

### 1. Introduction

SX Phoenicis stars are Population II pulsating variable stars with shorter periods (< 0<sup>d</sup>:1) and larger amplitudes of variability, compared with Scuti stars which are short period pulsating variable stars belonging to Population I. Interestingly most of known SX Phoenicis stars in globular clusters are located in the blue straggler star (BSS) region in the color-m agnitude diagram (CMD). It is long since the presence of the BSS in the globular cluster M 3 became known by Sandage (1953). Now there are many BSS discovered in

the globular clusters and open clusters in our Galaxy and in the elds of nearby galaxies. However, the origin of the BSS is still controversial. The BSSs are hotter and brighter than the main sequence turn-o stars so that it is needed som emechanisms to make the stars hotter and brighter (more massive) to explain the origin of the BSS. There have been suggested two classes of mechanisms to explain the origin of the BSS. One class involves a single star where mixing in the atm osphere can increase the life time of the main sequence. The other class is based on merger process where two low-mass stars are merged to form a more massive unevolved star via mass transfer or direct collisions between stars. These days the latter is preferred (Bailyn 1995). Recently a signi cant fraction of the BSS in globular clusters are known to be SX Phoenicis stars. The presence of these pulsating variable stars (as well as eclipsing variables) among the BSS provides an excellent opportunity to understand the origin of the BSS, because we can investigate physical properties and processes of the stars in more detail compared with the case of non-variable stars. Therefore the study of SX Phoenicis stars can provide us with in portant clues on the origin of the BSS.

It is only after 1980's that SX Phoenicis stars were discovered in the globular clusters. Since the rst discovery of SX Phoenicis stars in the globular cluster ! Centauri (N iss 1981), the discovery rate of these stars in the globular clusters has increased rapidly in the last decade, using the interm ediate-to-large telescopes. Recently, m any SX Phoenicis stars are being discovered in several globular clusters and nearby galaxies. Som e exam ples are ! Centauri (K aluzny et al. 1997), M 5 (K aluzny et al. 1999), M 15 (Jeon et al. 2001a), M 22 (K aluzny & Thom pson 2001), 47 Tuc (G illiland et al. 1998; B runtt et al. 2001), and M 55 (Pych et al. 2001). Rodr guez & Lopez-G onzalez (2000) listed a total of 122 SX Phoenicis stars belonging to 18 globular clusters and 27 stars in two external galaxies, covering inform ation published until January, 2000. However, the characteristics of the SX Phoenicis stars have not yet been fully explained by the present stellar evolution theories, requiring further investigation in observation and theory.

In addition, SX Phoenicis stars are known to show period-lum inosity (P-L) or period-m etallicity-lum inosity (P-Fe/H)L) relation, which can be a very useful distance indicator for globular clusters and nearby galaxies (M cN am ara 1997; N em ec et al. 1994). However, the number of known SX Phoenicis star in a given globular cluster is sm all and m ost studies use an inhom ogeneous sam ple based on di erent sources. Therefore there is seen a large variation am ong these relations. For exam ple, M cN am ara (1997) derived a slope of {3.7, while Pych et al. (2001) obtained a much atter slope of {2.9.

In 1999 we began time series CCD photom etry to search for variable BSSs in globular clusters using the 1.8m telescope at the Bohyunsan Optical A stronom y Observatory (BOAO) in Korea. Generally we can detect SX Phoenicis type stars brighter than V = 19.5 m ag for 200 seconds exposures, depending on the seeing condition. For larger am plitude variable stars such as W UM a type eclipsing binaries, the detecting m agnitude reaches 20.2 m ag for the same exposures. The rst result of our survey was discoveries of an SX Phoenicis star (Jeon et al. 2001a) and two W UM a type variable stars (Jeon et al. 2001b) in the globular cluster M 15.

In this paper we present the result of our survey for short period variable stars in the m etal-poor globular cluster M 53. M 53 ( $RA = 13^{h} 12^{m} 55^{s}3$ ,  $DEC = +18 10^{0} 09^{m}$ , J2000.0; Harris (1996)) has a very low m etallicity [Fe/H] = 1:99, a low interstellar reddening E(B = V) = 0.02, and an apparent V distance m odulus (m = M)<sub>V</sub> = 16:38 (Harris 1996). M any blue straggler stars are known to exist in M 53 (Rey et al. 1998). Some of these blue stragglers are expected to be SX Phoenicis stars, but no SX Phoenicis stars have been reported yet (R odr guez & Lopez-G onzalez 2000; C lem ent 2001). Here we report a discovery of eight SX Phoenicis stars in this cluster.

This paper is composed as follows. Observations and data reduction are described in

{ 4 {

Section 2. Section 3 presents the light curves of the new SX Phoenicis stars in M 53 and Section 4 discusses the characteristics of these stars, including the pulsation m odes and the period-lum inosity relation. Finally prim ary results are summarized in Section 5.

### 2. Observations and D ata R eduction

# 2.1. Observations

We obtained time series CCD in ages of M 53 for eleven nights from M arch 12th, 1999 to April 1st, 2001. A total of 45 and 381 frames were obtained for B and V bands, respectively. Because the observations were performed under various seeing condition, we adjusted exposure times depending on the seeing. The observation log is listed in Table 1.

### ED ITOR: PLACE TABLE 1 HERE.

The CCD in ages were obtained with a thinned SITe 2k CCD can era attached to the BOAO 1.8m telescope. The eld of view of a CCD in age is 11% 11% (0%3438 pixel<sup>1</sup>) at the f/8 Cassegrain focus of the telescope. The readout noise, gain and readout time of the CCD are 7.0 e , 1.8e = ADU and 100 seconds, respectively.

A greyscale m ap of a V-band CCD in age is shown in Figure 1. It shows only south-east region (7% 6%) of the cluster, out of the total observed eld of 11% 11%. Eight new SX Phoenicis stars are represented by circles and labeled from SXP1 to SXP8.

ED ITOR: PLACE FIGURE 1 HERE.

### 2.2. Data Reduction

Using the IRAF/CCDRED package, we processed the CCD in ages to correct overscan regions, trim unreliable subsections, subtract bias frames and correct at eld in ages. Instrumental magnitudes were obtained using the point spread function (PSF) tting photometry routine in the IRAF/DAOPHOT package (Massey & Davis 1992). The instrumental magnitudes of the stars in M 53 were transformed to the standard system using the BVI standard stars (Landolt 1992) observed on the photometric night of March 30, 2000. In Figure 2 we plotted CMDs for a total of about 18,000 stars on the (V, B V) plane. The left panel in Figure 2 shows the CMD for a central region at r< 1%, and the right panel shows the CMD for an outer region at r 1%.

### ED ITOR: PLACE FIGURE 2 HERE.

On the right panel, the main sequence (MS), the red giant branch (RGB) and the horizontal branch (HB) are clearly seen. In addition, there are about 100 stars at the brighter and bluer region above the MS turno (represented by a box), which are blue stragglers. Large scatters shown in the left panel CMD for r< 1% are due to crowding, so that we could not nd any variable stars at r < 1%.

W e applied the ensemble normalization technique (G illiland & Brown 1988; Jeon et al. 2001a) to normalize instrumental magnitudes between time-series CCD frames. We used about a hundred normalizing stars ranging from  $13^{n}$ . 7 to  $17^{n}$ .8 for the V-band and from  $13^{n}$ .7 to  $18^{n}$ .5 for the B-band, except for variable stars and central stars at r < 190 where the crowding is severe. For B-band data, we use them only for obtaining mean magnitudes, because the data quality was not good enough to apply frequency analysis. The normalization equation is

$$B \text{ or } V = m + c_1 + c_2 (B \quad V) + c_3 P_x + c_4 P_v$$
(1)

#### {6{

where B, V, and m are the standard and instrum entalm agnitudes of the normalizing stars, respectively.  $c_1$  is the zero point and  $c_2$  is the color one cient.  $c_3$  and  $c_4$  are used to correct position-dependent terms such as atmospheric di erential extinction and variable PSF.

### 3. Light Curves and Frequency Analysis

A fler photom etric reduction of the tim e-series frames, we inspected lum inosity variations for about 18,000 stars to search for variable stars. >From the variable star search we discovered eight new SX Phoenicis stars and recovered about tenth of previously known RR Lyrae stars. Here we report only the results on the SX Phoenicis stars. Detailed results on the RR Lyrae stars will be presented in a separate paper (Jeon et al. 2003).

The m ean m agnitudes and color indices of the SX Phoenicis stars are listed in Table 2. The R A .(J2000.0) and D ecl.(J2000.0) coordinates of the stars in Table 2 were obtained from the astrom etry using the G uide Star C atalogue (Version 1.1). Interestingly all the new SX Phoenicis stars are located in the south-east direction from the center of M 53. W e checked any possible system atic errors preventing us from noting variable stars in other region of the eld, but noting none. The coordinates, m ean m agnitudes and color indices of the eight new SX Phoenicis stars are listed in Table 2.

ED ITOR: PLACE TABLE 2 HERE.

Figure 3 displays V -band light curves of eight new SX Phoenicis stars. Because SXP3 is located near the cluster center and SXP4 close to the CCD edge, some data of SXP3 were lost by poor seeing and some data of SXP4 by daily variation of observing eld center. The light curves in Figure 3 show typical characteristics of SX Phoenicis stars, i.e., short periods and low am plitudes.

#### ED ITOR: PLACE FIGURE 3 HERE.

We have performed multiple-frequency analyses to derive the pulsating frequencies of the eight SX Phoenicis stars, using the discrete Fourier transform and linear least-square tting methods (K in & Lee 1996). Figure 4 displays the power spectra the eight SX Phoenicis stars we derived. Sm all inner panels in Figure 4 represent the pre-whitening processes. We searched for one to three pulsating frequencies for each star at the frequency region from 5 to 70 cycles day<sup>1</sup>. Low frequencies detected for the SXP1, SXP3, SXP4 and SXP8 might have resulted from variable seeing condition and/or drift during long observing runs from 1999 to 2001. Synthetic light curves obtained from these analyses are superim posed on the data in Figure 3, tting the data well.

### ED ITOR: PLACE FIGURE 4 HERE.

The results of the multiple-frequency analyses for the eight SX Phoenicis stars are sum marized in Table 3. In Table 3, we accepted as intrinsic pulsating frequencies with their amplitude signal-to-noise ratios larger than 4 (B reger et al. 1993) except for a frequency  $f_3$  of SX P2. A lthough the amplitude signal-to-noise ratio of SX P2, 3.8, is slightly sm aller than 4, its  $f_3$  seem s to be a harm onic frequency of  $f_1$ . The periods of prim ary modes for these SX Phoenicis stars range from 0.0385 days to 0.0701 days, and the sem i-amplitudes of the variability range from 0.030 m ag to 0.118 m ag.

# ED ITOR: PLACE TABLE 3 HERE.

3.1. The Characteristics of the individual SX Phoenicis star

SX P1: SX P1 has the longest prim ary period of 0.0701 days am ong the eight new SX

#### { 8 {

Phoenicis type stars of M 53. This star has two frequencies,  $f_1 = 142600$  cycles day <sup>1</sup> and  $f_2 = 27.5256$  cycles day <sup>1</sup>. This starm ight be a mono-periodic pulsator, because the second frequency  $f_2$  seems to be caused by 1 cycle day <sup>1</sup> alias e ect of a harmonic frequency of  $f_1$ ,  $2f_1 = 28.5200$  cycles day <sup>1</sup>. In F igure 3, the light curves of SXP1 shows an asymmetric sinusoidal feature, which is the characteristic of harmonic frequencies. The maximum amplitude of SXP1 is estimated to be V = 0.292 mag.

SX P 2: Light curves of this star in Figure 3 show an amplitudes change from day to day, in plying the excitation of multiple pulsating frequencies. We derived two closely separated frequencies of  $f_1 = 22.0450$  cycles day <sup>1</sup> and  $f_2 = 22.9750$  cycles day <sup>1</sup>, with a ratio of  $f_1=f_2 = 0.9595$ . These frequencies can be explained by excitation of non-radial modes. Pulsations with non-radial modes were found in several recent observational results of the SX Phoenicis stars: SX Phoenicis itself (G arrido & Rodr guez 1996), BL Cam (Zhou et al. 1999), V3 in 47 Tuc (G illiland et al. 1998), SX P1 in M 15 (Jeon et al. 2001a), and ten SX Phoenicis stars in M 55 (Pych et al. 2001). A first these two frequencies are pre-whitened, the third frequency is detected at  $f_3 = 43.0190$  cycles day <sup>1</sup>, which m ight have resulted from 1 cycle day <sup>1</sup> alias e ect of  $2f_1 = 44.0900$  cycles day <sup>1</sup>. Even if the amplitude signal-to-noise ratio for the third frequency is only 3.8, the light curves of SX P2 show asymmetric sinusoidal feature, supporting the existence of harm onic frequencies. The m axim um amplitude of SX P2 is estim ated to be V = 0.372 m ag, which is the largest value am ong the eight SX Phoenicis stars in M 53.

SX P 3, SX P 4, SX P 5, SX P 6, SX P 7: We could detect only a prim ary frequency for these ve stars. Prim ary frequencies and the maximum amplitudes of SX P 3, SX P 4, SX P 5, SX P 6 and SX P 7 are  $f_1 = 20.8796$ , 20.0114, 22.9880, 22.6000 and 24.1024 cycles day <sup>1</sup> and V = 0.204, 0.138, 0.082, 0.200 and 0.194 m ag, respectively. In Figure 4 SX P 4 and SX P 5 show some hints for the existence of harm onic frequencies above 40 cycles day <sup>1</sup>, but they are not conclusive.

SX P 8: SX P 8 has the shortest prim ary period of 0.0385 days and the smallest sem i-am plitude of 0.030 m ag am ong the eight SX Phoenicis stars of M 53. SX P 8 shows a typical feature of low am plitude Scuti stars (LAD S), i.e., the feature of com plicated oscillation pattern and several frequencies com pared to one or two stable frequencies of high am plitude Scuti stars (HAD S). We could detect three frequencies with am plitude signal-to-noise ratios over 4.0, although the data are not excellent. The prim ary, secondary and third frequencies are  $f_1 = 25.9800$ ,  $f_2 = 38.1942$  and  $f_3 = 8.2322$  cycles day <sup>1</sup>, respectively.

# 4. Discussion

# 4.1. Characteristics of the New SX Phoenicis Stars

In Figure 5, we show the position of the eight new SX Phoenicis stars in the color-magnitude diagram of M 53. All the SX Phoenicis stars discovered in M 53 are found to be located in the blue straggler region, brighter and bluer than the main sequence turno point. It is interesting that all of them are located in the red side in the BSS region of M 53. This indicates that the pulsational instability strip m ay cover only a part of the BSS region (a hotter region in this case) and that only those (or some of them) in this region can start pulsation. It is not yet known why some BSSs pulsate and some do not in the sam e pulational instability region. It needs further studies using several globular clusters with SX Phoenicis stars.

ED ITOR: PLACE FIGURE 5 HERE.

In Figure 6 we have compared the V -amplitudes and periods of the SX Phoenicis stars in M 53 with those of SX Phoenicis stars and Scuti stars in other globular clusters and eld SX Phoenicis stars. The sources of the data in Figure 6 are Rodr guez et al. (2000) for eld SX Phoenicis stars and Scuti stars, and Rodr guez & Lopez-G onzalez (2000) for SX Phoenicis stars in G alactic globular clusters. Figure 6 shows that the V -amplitudes and periods of the eight SX Phoenicis stars (star symbols) are consistent with those for other SX Phoenicis stars in globular clusters. The V -amplitudes of the SX Phoenicis stars are much larger than those of Scuti stars with the same period, and the periods of the SX Phoenicis stars the shortest end.

#### ED ITOR: PLACE FIGURE 6 HERE.

### 4.2. M ode Identi cation and Period-Lum inosity Relation

The period-lum inosity (P-L) relation of SX Phoenicis stars in the globular clusters can be very useful for estimating the distances to the clusters and nearby galaxies. However, it is not easy to derive the P-L relation from the observations, because there are often seen a mixture of di erent pulsation modes in the SX Phoenicis stars. Di erent pulsation modes of the SX Phoenicis stars follow di erent P-L relation (o sets in the zero points).

It is well known that observational identi cation of the pulsation modes of the SX Phoenicis star is di cult in general. M dN am ara (1997, 2000, 2001) suggested that the light am plitude and the degree of asymmetry of the light curves might be useful parameters in deciding the pulsating modes of the high am plitude stars. M dN am ara (2000) showed that most of the known SX Phoenicis stars at the fundamental mode show large am plitude (for example, larger than 0.15 m ag in the case of SX Phoenicis stars in ! C entauri) and asymmetric light curve. However, not all of the SX Phoenicis stars with large am plitude are at the fundam entalm ode, as shown by M dN am ara (2000). On the other hand, Pych et al. (2001) pointed out \Am plitudes generally yield no de nitive clues for the identi cation of m odes, except that large am plitudes are m ore likely to occur in radial pulsations". A nother possible way of identifying the pulsating m odes is to use the period-lum inosity relations and the period ratios of the SX Phoenicis stars in globular clusters, being applied in this study.

In Figure 7 we display the period and m ean V m agnitude relation for the SX Phoenicis stars in M 53 in comparison with other globular clusters and theoretical models. Several interesting features are seen in Figure 7. Figure 7 (a) illustrates that six SX Phoenicis stars in M 53 show a very good correlation between the period and m ean V m agnitude, while the other two stars (SX P2 and SX P8) are in the higher mode than the others. SX P1, the brightest in the sam ple, has a period m uch longer than the rest of the SX Phoenicis stars. This star is considered to be a member of M 53 for three reasons: rst,  $\ddagger$  is seen inside the blue straggler region in the color-m agnitude diagram; secondly,  $\ddagger$  is located closer to the center of M 53 in the sky than SX P4 which is clearly a member of M 53; and thirdly,  $\ddagger$  has period and V m agnitude consistent with those of other SX Phoenicis stars in M 53. W e derive a period – lum inosity relation using these six stars with 1:36 < Loop [d] < 1:15:

$$\langle V \rangle = 3.010 (0.262) LogP + 15.310 (0.048)$$
 (2)

with an rm s scatter of 0.038. This result corresponds to  $\langle M_V \rangle = 3.010 \text{LogP} = 1.070$  for an adopted distance m odulus of (m M)<sub>V</sub> = 16.38 (Harris 1996).

In Figure 7 (a) we compare the SX Phoenicis stars in M 53 and NGC 5466. The results for NGC 5466 were also obtained using the same procedures as for M 53 from the similar data so that they can be compared directly with those of M 53 (Jeon et al. 2003). We adopted distance m odulus of  $(m V)_0 = 16:15$  and zero foreground reddening for NGC 5466 (Harris 1996). It is shown clearly that the P-L relation of the SX Phoenicis stars in NGC 5466 agrees very well with that of M 53. The P-L relation for NGC 5466 has a slope very sim ilar to that for M 53 and a zero point slightly brighter than that for M 53. M ost of the SX Phoenicis stars in NGC 5466 are considered to be at the fundam ental pulsating m ode (see Jeon et al. (2003) for details). Therefore it is concluded that all SX Phoenicis stars (except for SXP2 and SXP8) in M 53 are probably at the fundam ental pulsating m ode. Then SXP2 and SXP8 are probably at the rst overtone m ode and at the second overtone m ode, respectively. Interestingly the am plitudes of SX Phoenicis stars in NGC 5466 are m uch larger than those of M 53 stars. This will be discussed in detail in Jeon et al. (2003).

In Figure 7 (b) we compare the SX Phoenicis stars in M 53 and M 55 where as many as 24 SX Phoenicis stars were discovered recently by Pych et al. (2001). We adopted distance m odulus of (m V)<sub>V</sub> = 13.86 (Pych et al. 2001) and foreground reddening of E (B V) = 0.07 for M 55 (Harris 1996). Note that Pych et al. (2001) derived, using the H IPPARCOS data for SX Phoenicis star itself, a distance m odulus, (m V)<sub>V</sub> = 13.86 0.25, which is very similar to the value given by Harris (1996), (m V)<sub>V</sub> = 13.87. Pych et al. (2001) used SX Phoenicis stars in M 55 to estimate the slope of the P-L relation and used H IPPARCOS data for SX Phoenicis star itself to calibrate the zero point, deriving a P-L relation for the fundam entalm ode:  $\langle M_V \rangle = 2.88LogP = 0.77$ . Figure 7 (b) shows that the P-L relation for M 53 agrees approximately with that for the fundam entalm ode of SX Phoenicis stars in M 55 and that the scatter of the P-L relation for M 53 is much smaller than that for M 55.

In Figure 7 (c) we compare the SX Phoenicis stars in M 53 and ! Centauriwhere as many as 34 SX Phoenicis stars were discovered by Kaluzny et al. (1996, 1997). Using the data for ! Centauri, M dN am ara (2000) derived steep P-L relations for the fundam entalm ode and rst overtone m ode,  $\langle V \rangle = 4:66LogP + 11:21$  and  $\langle V \rangle = 4:26LogP + 11:38$ , respectively. W e adopted distance m odulus of (m V)<sub>V</sub> = 14:05 0:11 and foreground reddening of E (B V) = 0:13 for ! Centauri given recently by Thom pson et al. (2001) who used a detached eclipsing binary in ! Centauri. These values are very similar to those in Harris (1996), (m V)<sub>V</sub> = 13:97 and E (B V) = 0:12.

Figure 7 (c) shows that the P-L relation for M 53 agrees approximately with that for the SX Phoenicis stars in ! Centauri which were considered to be at the rst overtone mode by M dN am ara (2000) (called A-sequence hereafter). Note that the steep slope for the rst overtone mode in ! Centauri is based on the data for the period range of log P < 1:3, where we stars at the fundamental mode in M 53 show a similar behavior. The SX Phoenicis stars at the fundamental mode in ! Centauri (called B-sequence hereafter) are 0.4 0.5 m ag fainter than those in M 53.

This result is very intriguing and the reasons for this discrepancy are not clear. ! Centauri has been known to be a very unique globular cluster among the G alactic globular clusters in several aspects: it is the most massive globular cluster in the M iky W ay; it has multiple stellar populations with a wide spread in metallicity from  $[Fe/H] = \{2.0 \text{ to } \{0.6 \text{ dex} (the metallicity of the main population is <math>[Fe/H] = 1.6 \text{ dex}$ ); and it has a very elongated structure. These features lead to a suggestion that it may be a remnant core of dwarf spheroidal galaxy which form ed via merging of several components (G nedin et al. (2002); Ferraro et al. (2002) and references therein). This peculiarity of ! C entaurim ay be related with the fact that it is the only cluster where more rst overtone mode SX Phoenicis stars were found than the fundam entalm ode stars (or in comparable number). Note that much fewer rst overtone mode stars were found than the fundam entalm ode stars (or in comparable number).

We checked several possibilities to understand this puzzling result: (1) If the A-sequence and the B-sequence are, respectively, indeed rst overtone mode and the fundamental mode as considered by M cN am ara (2000) (and if the P-L relation of SX Phoenicis stars is universal), then the distance modulus of either ! Centauri or M 53 must be wrong by as much as 0.4 0.5 m ag. This is very unlikely, considering the good agreement among the di erent estimates of the distance to these objects; (2) The A-sequence may be fundamental mode rather than instructione mode. Then it is not possible to explain the existence of the B-sequence without assuming that the B-sequence is another fundamental mode behind the A-sequence. This is also unlikely for the same reason as (1); (3) The A-sequence and the B-sequence may not be two independent modes. Instead both may belong to the same e fundamental mode with a large dispersion. This is contrary to the result given by M dN am ara (2000). However, it happens to be consistent with the theoretical P-L relation for the fundamental mode given by Templeton et al. (2002), as shown in Figure 7(d). Therefore this problem remains to be explained by further studies.

In Figure 7 (d) we can pare the P-L relation for M 53 with empirical relations given by M cN am ara (1997, 2002) and with theoretical relations given by Santolam azza et al. (2001) and Templeton et al. (2002). The slope of the P-L relation for the fundam ental m ode of M 53 is very similar to those in theoretical ones ({3.05 (Santolam azza et al. 2001) and {3.14 (Templeton et al. 2002) ). The six SX Phoenicis stars in M 53 are located at the right side of the blue edge of the fundam ental m ode relation given by Santolam azza et al. (2001), consistent with the conclusion that these stars are indeed the fundam ental m odes. On the other hand, the zero point of the M 53 relation is about 0.4 m ag brighter than that of the theoretical relation given by Templeton et al. (2002). The reason for this discrepancy is not known at this moment. The empirical P-L relations given by M cN am ara (1997, 2002) (< M<sub>V</sub> > = 3:725 log P 1:933 < M<sub>V</sub> > = 4:14 log P 2:46) are steeper than those for M 53 and the theoretical relations given by Santolam azza et al. (2001) and Templeton et al. (2002).

ED ITOR: PLACE FIGURE 7 HERE.

### 5. Sum m ary

We present time-series BV CCD photom etry of the metal-poor globular cluster M 53. we have discovered eight new SX Phoenicis type stars (labeled from SXP1 to SXP8). Physical parameters of these stars are summarized in Tables 2 and 3. All the new SX Phoenicis stars are located in the blue straggler star region of a color-magnitude diagram of M 53. One of these stars (SXP2) is found to have very closely separated pulsation frequencies:  $f_1=f_2 = 0.9595$  where  $f_1$  and  $f_2$  are primary and secondary frequencies. This may be due to excitation of non-radial modes.

Considering the position in the period-V m agnitude diagram, six of these SX Phoenicis stars are identified to be pulsating at the fundam ental mode, and SXP2 and SXP3 are probably at the st overtone and second overtone modes. We derive a period - lum inosity relation for the fundam ental mode for the period range of  $1.36 < \log P$  [d] < 1.15: < V > = 3.010(0.262)LogP + 15.310(0.048) with an rm s scatter of 0.038, corresponding to  $< M_V > = 3.010LogP$  1.070 for an adopted distance modulus of (m M)<sub>V</sub> = 16.38 (Harris 1996).

A uthors are very much grateful to Dr. D. H. M dN am ara, the referee, who provided very useful comments to improve this paper. M G L thanks the director and stall of the Department of Terrestrial M agnetism, the Carnegie Institution of W ashington, for their kind hospitality. M G L. was supported in part by the K orean Research Foundation G rant (KRF-2000-DP0450).

#### REFERENCES

- Bailyn, C.D. 1995, ARA & A, 33, 133
- Breger, M., Stich, J., & Garrido, R., et al. 1993, A & A, 271, 482
- Bruntt, H., Frandsen, S., Gilliland, R.L., Christensen-Dalsgaard, J., Petersen, J.O., Guhathakurta, P., Edmonds, P.D., & Bono, G. 2001, A & A, 371, 614
- Clement, C.M., Muzzin, A., Dufton, Q., Ponnanpalam, T., Wang, J., Burford, J., Richardson, A., Rosebery, T., Rowe, J., & Sawyer Hogg, H. 2001, AJ, 122, 2587
- Ferraro, F.R.; Bellazzini, M.; Pancino, E. 2002, ApJ, 573, 195
- Garrido, R., & Rodr guez, E. 1996, MNRAS, 281, 696
- Gilliland, R.L., Bono, G., Edmonds, P.D., Caputo, F., Cassisi, S., Petro, L.D., Saha, A., & Shara, M.M. 1998, ApJ, 507, 818
- Gilliland, R.L., & Brown, T.M. 1988, PASP, 100, 754
- G nedin, O leg Y .; Zhao, HongSheng; Pringle, J.E.; Fall, S.M ichael; Livio, M ario; M eylan, G eorges 2002, ApJ, 568, L23
- Harris, H.C. 1996, AJ, 112, 1487 (http://physun.physics.m.cm.aster.ca/G.lobular.html; revised in June 22, 1999)

Jeon, Y.-B., Kim, S.-L., Lee, H., & Lee, M.G. 2001a, AJ, 121, 2769

- Jeon, Y.-B., Kim, Lee, H., Kim, S.-L., & Lee, M.G. 2001b, IBVS, no. 5189
- Jeon, Y.-B., Lee, H., Kim, S.-L., & Lee, M.G. 2003, in preparation.
- Kaluzny, J., Kubiak, M., Szymanski, M., Udalski, A., Krzeminski, W., & Mateo, M. 1996, A & A S, 120, 139

- Kaluzny, J., Kubiak, M., Szymanski, M., Udalski, A., Krzeminski, W., Mateo, M., & Stanek, K. 1997, A & A S, 122, 471
- Kaluzny, J., Thompson, I., Krzeminski, W., & Pych, W. 1999, A&A, 350, 469
- Kaluzny, J., & Thompson, IB. 2001, A&A, 373, 899
- Kim, S.-L., & Lee, S.-W .1996, A&A, 310, 831
- Landolt, A.J. 1992, AJ, 104, 340
- Massey, P., & Davis, LE. 1992, A User's Guide to Stellar CCD photom etry with IRAF
- M dN am ara, D H . 1997, PASP, 109, 1221
- M dN am ara, D H . 1997, PASP, 112, 1096
- M dN am ara, D H . 1997, PA SP, 113, 335
- M dN am ara, D H. 2002, A SP Conf. Series 259, 116
- Nem ec, JM ., Nem ec, A F N ., & Lutz, T E . 1994, A J, 108, 222
- N iss, B. 1981, A & A, 98, 415
- Pych, W., Kaluzny, J., Krzeminski, W., Schwarzenberg-Czerny, A., & Thompson, IB. 2001, A&A, 367, 148
- Rey, S-C., Byun, Y-J., Chun, M-S. 1998, AJ, 116, 1775
- Rodr guez, E., & Lopez-Gonzalez, M.J. 2000, A&A, 359, 597
- Rodr guez, E., Lopez-Gonzalez, M. J., & Lopez de Coca, P. 2000, A & A S, 144, 469

Sandage, A. 1953, AJ, 58, 61

Santolam azza, P., Marconi, M., Bono, G., Caputo, F., Cassisi, S., & Gilliland, R.L.2001, ApJ, 554, 1124

Thompson, I.B., Kaluzny, J., Pych, W., Burley, G., Krzeminski, W., Paczynski, B., Persson, S.E., Preston, G.W. 2001, AJ, 121, 3089

Tem pleton, M., Basu, S., & Dem arque, P. 2002, ApJ, 576, 963

Zhou, A-Y., Rodr guez, E., Jiang, S-Y., Rolland, A., & Costa, V. 1999, MNRAS, 308, 631

This manuscript was prepared with the AAS  ${\rm I\!AT}_E X$  macros v5.0.



Fig. 1. A greyscale map of a V-band CCD in age of the globular cluster M 53. This in age shows a 7% 6% eld in the south-east region of the cluster, out of the observed eld of 11%. Eight new SX Phoenicis stars are labeled from SXP1 to SXP8.





Fig. 2. Color-m agnitude diagram s of M 53. The left panel is for a central region at r < 1%, and the right panel is for an outer region at r 1%. The box represents a blue straggler region.

# { 22 {



H.J.D.(+2450000)





Fig. 3. Observed light curves (dots) for eight new SX Phoenicis stars. Synthetic light curves (solid lines) obtained from the multiple-frequency analysis (see Table 3) are superim – posed on the data.



Fig. 4. Power spectra of eight new SX Phoenicis stars. W indow spectra are shown in the smallbox within each panel.



Fig. 5. Positions of eight new SX Phoenicis stars in the color-m agnitude diagram of M 53. Note that all they are located in the blue straggler region.



Fig. 6. | V am plitude versus period diagram. Star sym bols denote eight new SX Phoenicis stars in M 53, triangles represent eld SX Phoenicis stars, open circles indicate SX Phoenicis stars in other globular clusters, and dots denote Scuti stars.



Fig. 7. Mean magnitude <V> versus period diagram. Filled circles represent the SX Phoenicis stars in M 53. The solid line represent a linear t to the data for the fundam ental mode of M 53. (a) C om parison with the SX Phoenicis stars in NGC 5466 (open circles). The dash line represents a linear t to the data of the fundam entalmode of NGC 5466 (Jeon et al. 2003). (b) C om parison with the SX Phoenicis stars in M 55 (open squares). The dash line represents a linear t to the data for the fundam entalmode of M 55 (Pych et al 2001). (c) C om parison with the SX Phoenicis stars in ! C entauri (open triangles). The two dashed lines represents linear ts to the data for the A-sequence (left) and the B-sequence (right) of ! C entauri. (d) C om parison with the oretical P-L relations (short dashed lines) given by M cN am ara (1997, 2002) and with theoretical P-L relations. The long dashed line represents a theoretical relation for the fundam entalmode given by Tem pleton et al. (2002). The dot-

dashed lines, respectively, represent the blue edge of the fundam entalm ode, the rst overtone m ode and the second overtone m ode (from the bottom to top) given by Santolam azza et al. (2001).

| D ate     | Start H JD            | D uration | N <sub>obs</sub> | Seeing     | Exposure T in e         | Rem arks       |
|-----------|-----------------------|-----------|------------------|------------|-------------------------|----------------|
| (UT)      | (2,450,000+)          | (hours)   |                  | (arcsec)   | (seconds)               |                |
|           |                       |           |                  |            |                         |                |
| 1999 312  | 1250 <b>.</b> 173(V)  | 4.7       | 30 <b>(</b> V )  | 12 2.4(V)  | 300 <b>(</b> V )        |                |
| 2000 3 30 | 1634.007(V)           | 5.1       | 38 <b>(</b> V )  | 12 23(V)   | 180,240(V)              | standard stars |
|           | 1634.022(B)           |           | 12(B)            | 1.3 2.0(B) | 300,450(B)              |                |
| 2000 3 31 | 1635.018(V)           | 7.6       | 46 <b>(</b> V )  | 2.1 3.4(V) | 210,180(V)              |                |
|           | 1635.029(B)           |           | 14(B)            | 2.3 3.1(B) | 300 <b>(</b> B <b>)</b> |                |
| 2000 4 1  | 1636 <b>.</b> 153(V)  | 23        | 16(V)            | 22 2.4 (V) | 420,210(V)              | thin cloud     |
|           | 1636 <b>.</b> 165(B)  |           | 1(B)             | 2.6(B)     | 300 <b>(</b> B <b>)</b> |                |
| 2000 4 4  | 1639 <b>.</b> 196(V)  | 1.0       | 6 <b>(</b> V )   | 22 2.4 (V) | 210 (V)                 |                |
|           | 1639,201(B)           |           | 3(B)             | 2.3 3.0(B) | 300 <b>(</b> B <b>)</b> |                |
| 2000 4 5  | 1640.010(V)           | 5.2       | 38 <b>(</b> V )  | 2.8 3.6(V) | 210,300(V)              |                |
|           | 1640.025(B)           |           | 11(B)            | 3.0 3.7(B) | 300 <b>(</b> B )        |                |
| 2000 4 6  | 1641 <b>.</b> 103(V)  | 4.3       | 34 <b>(</b> V )  | 1.6 2.1(V) | 210,180(V)              | thin cloud     |
|           | 1641.133(B)           |           | 4 <b>(</b> B )   | 1.8 1.9(B) | 300 <b>(</b> B )        |                |
| 2001 326  | 1995.085(V)           | 6.4       | 42 <b>(</b> V )  | 3.1 4.0(V) | 400 (V )                | bad seeing     |
| 2001 3 29 | 1998.058(V)           | 5.9       | 34 <b>(</b> V )  | 1.9 2.8(V) | 350 500 (V)             |                |
| 2001 3 31 | 2000 <b>.</b> 000 (V) | 7.5       | 42 (V )          | 1.8 2.7(V) | 500 (V)                 |                |
| 2001 4 1  | 2000 <b>.</b> 988 (V) | 8.1       | 53 (V )          | 12 23(V)   | 300 600 (V)             |                |

Table 1. Observation Log.

| Name | RA (J2000.0)         | Dec(J2000.0) | < V >  | <b> <v></v></b> |
|------|----------------------|--------------|--------|-----------------|
|      |                      |              |        |                 |
| SXP1 | 13 13 03.38          | 18 09 25 3   | 18.814 | 0.284           |
| SXP2 | 13 12 49.71          | 18 07 26.0   | 18.915 | 0.310           |
| SXP3 | 13 13 01 <b>.</b> 95 | 18 10 13.4   | 19248  | 0,261           |
| SXP4 | 13 13 19.09          | 18 06 40.7   | 19.171 | 0.337           |
| SXP5 | 13 13 08 21          | 18 07 38.7   | 19.441 | 0.252           |
| SXP6 | 13 13 09.44          | 18 09 40.1   | 19.366 | 0.301           |
| SXP7 | 13 13 05.01          | 18 08 36.1   | 19.522 | 0.223           |
| SXP8 | 13 12 52.77          | 18 04 58.4   | 18.959 | 0264            |

Table 2. Observational Param eters of the Eight SX Phoenicis Stars.

=

=

| Name | Value | Frequency <sup>a</sup> | <sup>,</sup> ⊳Amp. <sup>ь</sup> | Phase <sup>b</sup> | S/N <sup>c</sup> | M ode        |                      | Remai    | ks              |
|------|-------|------------------------|---------------------------------|--------------------|------------------|--------------|----------------------|----------|-----------------|
|      |       |                        |                                 |                    |                  |              |                      |          |                 |
| SXP1 | $f_1$ | 14,2600                | 0":103                          | 0.6873             | 9.0              | F            |                      |          |                 |
|      | $f_2$ | 27.5256                | 0ª:043                          | {1.1868            | 5.1              | $2f_1$       | $f_2$                | $2f_1 =$ | 0 <b>:</b> 9944 |
| SXP2 | $f_1$ | 22.0450                | 0 <sup>m</sup> :118             | {0.0736            | 14.4             | 1H           | the highest amp.     |          |                 |
|      | $f_2$ | 22.9750                | 0 <sup>m</sup> :047             | 1.5520             | 72               | Non-radial   | $f_1 = f_2 = 0.9595$ |          |                 |
|      | $f_3$ | 43.0190                | 0ª:021                          | {1.3179            | 3.8              | $2f_1$       | $f_3$                | $2f_1 =$ | 1 <b>:</b> 0630 |
| SXP3 | $f_1$ | 20.8796                | 0 <sup>m</sup> :102             | 4.2981             | 7.0              | F            |                      |          |                 |
| SXP4 | $f_1$ | 20.0114                | 0 <sup>m</sup> :069             | 3.1873             | 7.9              | F            |                      |          |                 |
| SXP5 | $f_1$ | 22.9880                | 0ª:041                          | 0.3222             | 6.0              | F            |                      |          |                 |
| SXP6 | $f_1$ | 22.6000                | 0 <sup>m</sup> :100             | 2.6747             | 12.1             | F            |                      |          |                 |
| SXP7 | $f_1$ | 24.1024                | 0 <sup>m</sup> :097             | 2.0239             | 14.4             | F            |                      |          |                 |
| SXP8 | $f_1$ | 25.9800                | 0°° <b>.</b> 030                | 3.1414             | 82               | 2Н           |                      |          |                 |
|      | $f_2$ | 38.1912                | 0 <sup>m</sup> :015             | {1.5238            | 4.4              | N on-radial? |                      |          |                 |
|      | $f_3$ | 8 2322                 | 0 <sup>m</sup> :014             | {0.7535            | 4.3              | N on-radial? |                      |          |                 |

Table 3. Pulsating Properties of the Eight SX Phoenicis Stars.

<sup>a</sup>In cycles per day.

 $^{b}V = C \text{ onst} + _{j}A_{j} \cos f2 f_{j}(t t_{0}) + _{j}g; t_{0} = H JD 2,450,000.00.$ 

 $^{\rm c}{\rm Am}$  plitude signal-to-noise ratio introduced by B reger et al. (1993).