

A LOW-MASS CENTRAL BLACK HOLE IN THE BULGELESS SEYFERT 1 GALAXY NGC 4395

ALEXEI V. FILIPPENKO¹ AND LUIS C. HO²
To appear in The Astrophysical Journal (Letters).

ABSTRACT

NGC 4395 is one of the least luminous and nearest known type 1 Seyfert galaxies, and it also lacks a bulge. We present a *Hubble Space Telescope (HST)* I-band image of its nuclear region, and Keck high-resolution (~ 8 km s⁻¹) echelle spectra containing the Ca II near-infrared triplet. In addition to the unresolved point source, there is a nuclear star cluster of size $r \sim 3.9$ pc; the upper limit on its velocity dispersion is only 30 km s⁻¹. We thus derive an upper limit of $6.2 \times 10^6 M_{\odot}$ for the mass of the compact nucleus. Based on the amount of spatially resolved light in the *HST* image, a sizable fraction of this is likely to reside in stars. Hence, this estimate sets a stringent upper limit on the mass of the central black hole. We argue, from other lines of evidence, that the true mass of the black hole is likely to be $10^4 - 10^5 M_{\odot}$. Although the black hole is much less massive than those thought to exist in classical active galactic nuclei, its accretion rate of $L_{\text{bol}} = L_{\text{Edd}} \sim 2 \times 10^{-2}$ to 2×10^{-3} is consistent with the mass-luminosity relation obeyed by classical AGNs. This may explain why NGC 4395 has a high-excitation (Seyfert) emission-line spectrum; active galaxies having low-ionization spectra seem to accrete at significantly lower rates. NGC 4395, a pure disk galaxy, demonstrates that supermassive black holes are not associated exclusively with bulges.

Subject headings: galaxies: individual (NGC 4395) — galaxies: kinematics and dynamics — galaxies: nuclei — galaxies: Seyfert

1. INTRODUCTION

The nucleus of NGC 4395, an Sd III–IV galaxy (Sandage & Tammann 1981) at a distance of only 4.2 Mpc (D. Minniti et al., in preparation)³, exhibits strong, narrow emission lines with relative intensities similar to those of type 2 Seyfert galaxies. Spectra having high signal-to-noise ratios, however, reveal the presence of faint, broad components to the permitted hydrogen and helium lines; NGC 4395 therefore has a Seyfert 1 nucleus, one of the nearest and least luminous known (Filippenko & Sargent 1989). With $M_B = -10.8$ mag, it provides an exceptional opportunity to study the active galactic nucleus (AGN) phenomenon on luminosity scales comparable to those of the brightest stars. Moreover, unlike the case in other low-luminosity AGNs (e.g., M81; Filippenko & Sargent 1988; Ho, Filippenko, & Sargent 1996), starlight contamination of the AGN in NGC 4395 is almost negligible because the bulge of this dwarf galaxy ($M_B = -17.5$ mag) is essentially absent. The extreme late type of the host galaxy is highly unusual for AGNs, which tend to be found in bulge-dominated systems (e.g., Ho, Filippenko, & Sargent 1997). The nucleus of NGC 4395 is detectable as a variable X-ray source (Lira et al. 1999; Moran et al. 1999; Iwasawa et al. 2000; Shih, Iwasawa, & Fabian 2003; Moran et al. 2003) that remains unresolved at *Chandra* resolution (Ho et al. 2001). It further exhibits a highly compact radio core (Moran et al. 1999; Ho & Ulvestad 2001) with a brightness temperature in excess of 2×10^6 K (Wrobel, Fasnacht, & Ho 2001).

Here we present a careful search for stellar absorption lines in NGC 4395. The measured stellar velocity dispersion provides an upper limit to the mass of the compact nucleus, and hence on the mass of the central black hole that presumably powers the AGN. This measurement has special significance in view of the fact that NGC 4395 is a *bulgeless* galaxy. Recent studies point

to the prevalence of central, supermassive ($10^6 - 10^9 M_{\odot}$) black holes and their close connection with galaxy bulges (Magorrian et al. 1998; Gebhardt et al. 2000; Ferrarese & Merritt 2000). By contrast, supermassive black holes seem to have little relation to galaxy disks (Kormendy et al. 2003). Our mass limit on NGC 4395 provides another critical data point for the currently poorly constrained low-mass end of the mass spectrum of central black holes.

2. OBSERVATIONS

High-resolution ($R = 38,000$) spectra of the nucleus of NGC 4395 were obtained on 1994 April 14 UT using the HIRES echelle spectrometer (Vogt 1992; Vogt et al. 1994) at the Nasmyth focus of the Keck I 10-m telescope; details will be presented in a subsequent paper (A. V. Filippenko & L. C. Ho, in preparation). Our setting spanned the wavelength range 6338–8774 Å (16 orders), and the slit width was $1.15''$, yielding a spectral resolution of ~ 8 km s⁻¹ full width at half-maximum intensity (FWHM). Exposure times were 30, 60, and 60 minutes. To enable identification and removal of telluric lines, we also observed the sdF star HD 84937 (Oke & Gunn 1983), whose spectrum has relatively few features. Data reduction followed standard procedures similar to those of Ho & Filippenko (1995). The sky-subtracted, wavelength-calibrated, one-dimensional spectra correspond to an effective aperture of $1.15'' \times 2.9''$.

3. STELLAR VELOCITY DISPERSION

Our primary aim for the spectra was to detect, or set limits on, the Ca II near-infrared triplet (8498, 8542, 8662). Figure 1 shows the two relevant orders of the Keck echelle data for NGC 4395; the signal-to-noise ratio per resolution element in the continuum is ~ 35 . Also shown are spectra of the

¹ Department of Astronomy, University of California, Berkeley, CA 94720-3411; alex@astro.berkeley.edu.

² The Observatories of the Carnegie Institution of Washington, 813 Santa Barbara St., Pasadena, CA 91101; lho@ociw.edu.

³ We scale all distance-dependent quantities to this value, revised from the often-quoted distance of 2.6 Mpc from Rowan-Robinson (1985).




FIG. 1.— One order, and part of the next order, from the high-resolution spectrum (FWHM = 8 km s^{-1}) of NGC 4395 obtained with the Keck I 10-m telescope. Also shown is the sdF comparison star HD 84937. The gap between 8570 \AA and 8650 \AA is due to incomplete spectral coverage of the detector.

comparison star HD 84937 taken with the same instrumental setup. The Ca II lines are detected in NGC 4395, at a systemic velocity of 320 km s^{-1} , adjacent to the hydrogen Paschen emission lines. These higher-order members of the Paschen series are seldom seen in AGN spectra; they are detected in NGC 4395 because its recombination lines have exceptionally high equivalent widths. Their presence, however, presents a challenge for measuring the Ca II lines in NGC 4395, since both components have comparable strengths. The differences between the laboratory wavelengths of the Ca II lines and the adjacent Paschen lines are 4.5 , 3.3 , and 2.8 \AA for Ca II 8498 , 8542 , and 8662 (respectively), and the FWHM of the Paschen lines is 1.7 \AA . Judging by the relative strengths and the profiles of the three Ca II lines, it appears that the 8542 and 8662 lines are substantially affected by Paschen emission. By simultaneously fitting two Gaussians, one to the absorption and the other to the emission component, we measure equivalent widths (EWs) of 0.42 , 0.44 , and 0.37 \AA for Ca II 8498 , 8542 , and 8662 (respectively). In the absence of emission-line contamination, the relative strengths of the three lines in old stellar populations are approximately $0.4:1:0.9$ (Terlevich, Díaz, & Terlevich 1990).

Given these special circumstances, conventional methods of deriving stellar velocity dispersions (e.g., Tonry & Davis 1981) are of questionable utility in this instance. Instead, we adopt a more straightforward strategy, namely by directly comparing the spectrum of NGC 4395 to diluted and velocity-broadened spectra of template stars. This method is similar to that discussed by Barth, Ho, & Sargent (2002), but here we limit the fitting only to the Ca II 8498 line, since it is least contaminated by emission (due to the relatively large wavelength separation and the weakness of the adjacent Paschen line; see Fig. 1). Figure 2 compares the spectrum of NGC 4395 to those of four K-giant stars⁴ (luminosity classes I–III), whose Ca II 8498 line has been diluted to match the strength in the galaxy. Unlike weaker metal lines, note that the intrinsic widths of the Ca II

triplets are *not* negligible at echelle resolution. They range from $20\text{--}25 \text{ km s}^{-1}$ for K-type giants and bright giants to 30 km s^{-1} for red supergiants (Fig. 2; Ho et al. 2003, in preparation). The dotted curve shows that the diluted spectrum of HD 214868 (K2 III) convolved with a Gaussian with $\sigma = 25 \text{ km s}^{-1}$ provides a reasonable match to NGC 4395. On the other hand, we note that HD 206778 (K2 Ib), whose Ca II 8498 line has an intrinsic $\sigma = 30 \text{ km s}^{-1}$, also gives a decent fit, with *no* additional broadening. Thus, we conservatively adopt $\sigma < 30 \pm 5 \text{ km s}^{-1}$ as the line-of-sight velocity dispersion of the nucleus, where the error bar reflects our estimate of the probable systematic uncertainty.

4. SIZE OF THE NUCLEAR STAR CLUSTER

In order to determine the physical size of the region from which the stellar Ca II lines originate, we have used GALFIT (Peng et al. 2002) to decompose an archival *Hubble Space Telescope* (*HST*) WFPC2 F814W (*I*-band) image (GO-6464; obtained on 1998 January 6 UT) of the nuclear region of NGC 4395 into several components, as shown in Figure 3. We find (1) an unresolved point source with $I = 17.42 \pm 0.1 \text{ mag}$, (2) a fairly round (axis ratio 0.83 ± 0.01 , position angle 99°), compact source with a Sersic profile (index = 2.63 ± 0.05), and (3) an extended component that can be fit with an exponential profile.

We identify the compact source, whose half-light radius is 0.19 (3.9 pc), with a nuclear star cluster that produces the observed Ca II lines; based on the *HST* photometry, it contributes 58% of the light in the effective spectral aperture. The extended component accounts for only 4% of the flux in this aperture; accordingly, we neglect it here. Thus, we assign a radius of 3.9 pc to the stellar component detected in the Keck spectrum of the nucleus of NGC 4395.

The measured EW of the Ca II 8498 line, 0.42 \AA , is only about 25% of that found in typical old stellar populations

⁴ We did not observe any velocity template stars at Keck in this setting. These stars come from observations made with the Shane 3-m telescope at Lick Observatory (Ho et al. 2003, in preparation); they have $R < 50,000$.




FIG. 2.— Region around Ca II 8498 absorption and Paschen 16 emission for the nucleus of NGC 4395 and four standard stars. The continuum has been normalized to unity. From top to bottom, the stars are HD 213179 (K2 II), HD 214868 (K2 III), HD 6953 (K7 III), and HD 206778 (K2 Ib). The dotted curve superposed on the spectrum of NGC 4395 is the spectrum of HD 214868 broadened by $v = 25 \text{ km s}^{-1}$.

(Terlevich et al. 1990). This may suggest that the nonstellar component contributes $\sim 75\%$ of the *I*-band light, inconsistent with the smaller (38%) contribution that we find for the point source in the *HST* image. However, NGC 4395 has a very low metallicity compared with typical spiral galaxies (e.g., Garnett 2002), and it is well known that the Ca II EW is small in stars having low metal abundance ($\sim 0.4\text{--}1.0 \text{ \AA}$ for Ca II 8498; Díaz, Terlevich, & Terlevich 1989). Roy et al. (1996) derive a value of $\log(O/H) = -3.7$ for an H II region within $\sim 0.1 \text{ kpc}$ of the nucleus, and Kraemer et al. (1999) suggest $\log(O/H) = -3.5$ for clouds in the narrow-line region. Since we do not know the precise EW of the undiluted Ca II 8498 line in the nuclear star cluster of NGC 4395, the amount of Ca II dilution produced by the nonstellar continuum is best estimated from the *HST I*-band image.

Note that the low metallicity inferred for the central star cluster in NGC 4395 is not inconsistent with the high metallicities found in the broad-line regions (BLRs) of high-redshift quasars (e.g., Pentericci et al. 2002, and references therein). The stellar cluster in NGC 4395 is much larger (relative to the probable black-hole mass) than the volume occupied by the BLR in luminous quasars; moreover, there is a well-known quasar metallicity-luminosity correlation, in light of which we expect low metal abundance even in the BLR of NGC 4395.

5. THE CENTRAL MASS OF NGC 4395

The observed radial velocity dispersion of the stellar absorption lines can be used to set an upper limit to the virial mass of the central star cluster (or star cluster plus black hole). For a bound system with negligible rotational support, $M \leq 2.5 h^2 v_r^2 / r_h = G$ (Spitzer 1969), where $h^2 v_r^2$ is the mean-square velocity, r_h is the half-mass radius, and G is the gravitational constant. Under the assumption of an isotropic velocity distribution, $h^2 v_r^2 = 3 \sigma^2$, and so $M \leq 7.5 \sigma^2 r_h = G$. As discussed above, the most secure constraint on σ is that of the Ca II 8498 line:

$\sigma < 30 \text{ km s}^{-1}$. The best measurement on the size of the cluster comes from the *HST* image, which gives $r_h = 3.9 \text{ pc}$ (Fig. 3). If we assume that light traces mass, then $r_h \approx r_{\text{light}}$, and $M < 6.2 \times 10^6 M_\odot$.




FIG. 3.— Decomposition of the *HST*/WFPC2 F814W image of the nuclear region of NGC 4395. Data points are shown with error bars. Undulations in the unresolved point-source profile are an artifact of diffraction. The scale length of the exponential disk may be erroneous by a factor of ~ 2 ; there was not enough detector area to properly estimate the background sky.

The derived upper limit to the mass of the nucleus of NGC 4395 is a factor of $\sim 10\text{--}100$ larger than those of typical Galactic globular clusters (Mandushev, Spassova, & Staneva 1991), and is several times more massive than the largest young “super star clusters” (SSCs) found in starburst systems (Ho & Filippenko 1996; Larsen et al. 2001; Smith & Gallagher 2001). However, if the actual mass is substantially less than the upper limit, and if only part of the measured mass comes from stars (the rest being the black hole), then the true mass may be comparable to that of the most massive globular clusters and SSCs. Indeed, adopting an apparent *I*-band magnitude of 16.8 for the stellar component of the nucleus (based on the photometric decomposition of the *HST* image), $M_I = -11.3 \text{ mag}$, within the realm of SSCs (e.g., Barth et al. 1995; Whitmore & Schweizer 1995; Maoz et al. 2001) and quite typical of nuclear star clusters in other late-type spiral galaxies (Böker et al. 2002). In the absence of reliable information on the age of the stellar population in the nucleus of NGC 4395, it is difficult to judge whether the inferred limit on the mass-to-light ratio [$M/L_I < 2(M/L_I)_{\text{SSC}}$] is abnormally high or not. The data, therefore, do not require the presence of excess dark mass. The nucleus of NGC 4395 may consist of an essentially normal cluster of stars, with a small black hole at its center accreting matter and producing the observed nonstellar activity. Indeed, the AGN bolometric luminosity of $\sim 3 \times 10^{40} \text{ erg s}^{-1}$ (Moran et al. 1999) can be produced by a $\sim 250 M_\odot$ black hole accreting at the Eddington limit.

It is instructive to examine other indirect methods to estimate black hole masses in AGNs. From photoionization modeling of the narrow-line region and BLR in NGC 4395, Kraemer et al. (1999) find that the broad H β line originates from a region with a radius of $r_{\text{BLR}} = 3 \times 10^{-4} \text{ pc}$. Assuming that the line-emitting gas is gravitationally bound, that the orbits are randomly oriented, and that the observed width of the H β line traces the velocity dispersion of the gas, the virial mass follows from $M_{\text{BH}} \approx 2 r_{\text{BLR}} \sigma^2 / G$. Choosing $\sigma = (\sqrt{3}/2) \text{ FWHM}$ (Netzer 1990) and $\text{FWHM} = 1500 \text{ km s}^{-1}$ for H β (Kraemer et al. 1999),

$M_{\text{BH}} \approx 1.2 \times 10^5 M_{\odot}$.

The size of the BLR can also be obtained through the luminosity-size relation recently established for type 1 AGNs studied with reverberation mapping. Equation (6) of Kaspi et al. (2000) relates r_{BLR} to the monochromatic luminosity of the featureless continuum at 5100 \AA , $L(5100 \text{ \AA})$. From the spectral information given in Filippenko & Sargent (1989) and Filippenko, Ho, & Sargent (1993) [$F(4400 \text{ \AA}) = 0.42 \text{ mJy}$], we find $L(5100 \text{ \AA}) = 6.6 \times 10^{39} \text{ erg s}^{-1}$ and $r_{\text{BLR}} = 3.3 \times 10^{-5} \text{ pc}$. Although one might legitimately question whether the luminosity-size relation can be extrapolated to the ultra-low luminosity regime of NGC 4395, it is interesting to note that this value of the BLR radius is a factor of 10 smaller than that obtained through photoionization modeling, remarkably consistent with a similar trend found in more luminous Seyfert 1 nuclei (Peterson 1993). With the smaller value of r_{BLR} , the virial mass is consequently an order of magnitude lower, $M_{\text{BH}} \approx 1.3 \times 10^4 M_{\odot}$.

Finally, the black hole mass in NGC 4395 can be estimated from its X-ray variability characteristics. Shih et al. (2003) applied the formalism of Hayashida et al. (1998) to *ASCA* observations of NGC 4395 and find $M_{\text{BH}} \approx 10^4 - 10^5 M_{\odot}$. Although the systematic uncertainties of this method are not well understood, it is encouraging that the X-ray-based mass, within its relatively large allowable range, turns out to be consistent with the virial mass (calculated using r_{BLR} from the luminosity-size relation).

To summarize, indirect estimates suggest that the putative black hole in NGC 4395 has a mass $M_{\text{BH}} \approx 10^4 - 10^5 M_{\odot}$. If so, the AGN is radiating at 2×10^{-2} to 2×10^{-3} of the Eddington limit, consistent with the extrapolation of the mass-luminosity relation for classical type 1 Seyfert nuclei and QSOs (Kaspi et al. 2000, eq. 11). Thus, the nucleus of NGC 4395 appears to be a low-mass analog of a QSO or type 1 Seyfert, with a “normal” accretion rate relative to the Eddington limit. By contrast, LINERs have healthy-sized black holes ($M_{\text{BH}} > 10^6 M_{\odot}$) but very low accretion rates ($L_{\text{bol}} = L_{\text{Edd}} < 10^{-3}$; Ho 1999; Ho et al. 2000). This supports the hypothesis that the accretion rate is one of the main factors in dictating the level of excitation exhibited by the ultraviolet/optical spectrum: NGC 4395 has a high-excitation spectrum, as in classical Seyferts and QSOs, while the spectra of LINERs are dominated by low-excitation emission lines.

6. IMPLICATIONS FOR BLACK HOLE DEMOGRAPHY

The recent discovery of a tight correlation between black hole mass and the stellar velocity dispersion of the bulge (Gebhardt et al. 2000; Ferrarese & Merritt 2000), the $M_{\text{BH}} - \sigma$ relation, has inspired many efforts to couple the formation and growth of supermassive black holes to the formation of their host galaxies (e.g., Kauffmann & Haehnelt 2000; Burkert & Silk 2001). An important clue is that the connection seems to involve primarily the bulge component, not the disk (Kormendy et al. 2003). The most stringent limit to date comes from *HST* stellar-dynamical observations of the nucleus of the Scd galaxy M33, also a bulgeless system: Gebhardt et al. (2001) find that any central black hole would have to be less massive than $1500 M_{\odot}$. The M_{BH} limit for M33 falls significantly below the mass predicted from the $M_{\text{BH}} - \sigma$ relation.

Do supermassive black holes *ever* form in pure disk systems? Yes, as illustrated by NGC 4395. Our assessment in § 5 suggests that the central black hole in NGC 4395 is likely to have a mass between 10^4 and $10^5 M_{\odot}$, with a firm upper limit of $6.2 \times 10^6 M_{\odot}$. These mass estimates are not inconsistent with the $M_{\text{BH}} - \sigma$ relation of Tremaine et al. (2002). For $\sigma = 30 \text{ km s}^{-1}$, it predicts $M_{\text{BH}} = 6.6 \times 10^4 M_{\odot}$. Although current ground-based surveys indicate that unambiguous type 1 Seyfert galaxies like NGC 4395 are quite rare, Ho et al. (1997) show that more subtle signatures of nonstellar activity (based on the spectrum of the narrow emission lines) can be found in 10% of late-type spirals (Sc and later).

The W. M. Keck Observatory, made possible by the generous financial support of the W. M. Keck Foundation, is operated as a scientific partnership between the California Institute of Technology, the University of California, and NASA. We are grateful to the Keck staff for their assistance with the observations. We thank Aaron Barth, Ryan Chornock, and Gibor Basri for useful comments, and especially Chien Peng for producing Figure 3 and for doing the relevant analysis. The excellent suggestions of an anonymous referee helped improve the paper. Support for this work was provided by the NSF through grants AST-9417213 and AST-9987438, by NASA through grant NAG 5-3556, and again by NASA through grant AR-7527 from the Space Telescope Science Institute, which is operated by AURA, Inc., under NASA contract NAS5-26555. The research of L. C. H. is supported by the Carnegie Institution of Washington.

REFERENCES

- Barth, A. J., Ho, L. C., Filippenko, A. V., & Sargent, W. L. W. 1995, *AJ*, 110, 1009
 Barth, A. J., Ho, L. C., & Sargent, W. L. W. 2002, *AJ*, 124, 2607
 Böker, T., van der Marel, R. P., Laine, S., Rix, H.-W., Sarzi, M., Ho, L. C., & Shields, J. C. 2002, *AJ*, 123, 1389
 Burkert, A., & Silk, J. 2001, *ApJ*, 554, L151
 Díaz, A. I., Terlevich, E., & Terlevich, R. 1989, *MNRAS*, 239, 325
 Ferrarese, L., & Merritt, D. 2000, *ApJ*, 539, L9
 Filippenko, A. V., Ho, L. C., & Sargent, W. L. W. 1993, *ApJ*, 410, L75
 Filippenko, A. V., & Sargent, W. L. W. 1988, *ApJ*, 324, 134
 ——. 1989, *ApJ*, 342, L11
 Garnett, D. R. 2002, *ApJ*, 581, 1019
 Gebhardt, K., et al. 2000, *ApJ*, 539, L13
 ——. 2001, *AJ*, 122, 2469
 Hayashida, K., Miyamoto, S., Kitamoto, S., Negoro, H., & Inoue, H. 1998, *ApJ*, 500, 642
 Ho, L. C. 1999, *ApJ*, 516, 672
 Ho, L. C., & Filippenko, A. V. 1995, *ApJ*, 444, 165 [Erratum: 464, 818 (1996)]
 ——. 1996, *ApJ*, 472, 600
 Ho, L. C., Filippenko, A. V., & Sargent, W. L. W. 1996, *ApJ*, 462, 183
 ——. 1997, *ApJ*, 487, 568
 Ho, L. C., Rudnick, G., Rix, H.-W., Shields, J. C., McIntosh, D. H., Filippenko, A. V., Sargent, W. L. W., & Eracleous, M. 2000, *ApJ*, 541, 120
 Ho, L. C., & Ulvestad, J. S. 2001, *ApJS*, 133, 77
 Ho, L. C., et al. 2001, *ApJ*, 549, L51
 Iwasawa, K., Fabian, A. C., Almaini, O., Lira, P., Lawrence, A., Hayashida, K., & Inoue, H. 2000, *MNRAS*, 318, 879
 Kauffmann, G., & Haehnelt, M. 2000, *MNRAS*, 311, 576
 Kormendy, J., et al. 2003, *ApJ*, submitted
 Kraemer, S. B., Ho, L. C., Crenshaw, D. M., Filippenko, A. V., & Shields, J. C. 1999, *ApJ*, 520, 564
 Larsen, S. S., Brodie, J. P., Elmegreen, B. G., Efremov, Y. N., Hodge, P. W., & Richter, T. 2001, *ApJ*, 556, 801
 Lira, P., Lawrence, A., O’Brien, P., Johnson, R. A., Terlevich, R., & Bannister, N. 1999, *MNRAS*, 304, 109
 Magorrian, J., et al. 1998, *AJ*, 115, 2285
 Mandushev, G., Spassova, N., & Staneva, A. 1991, *A&A*, 252, 94
 Maoz, D., Barth, A. J., Ho, L. C., Sternberg, A., & Filippenko, A. V. 2001, *AJ*, 121, 3048
 Moran, E. C., Eracleous, M., Leighly, K. M., Chartas, G., Filippenko, A. V., Ho, L. C., & Blanco, P. R. 2003, *ApJ*, submitted

- Moran, E. C., Filippenko, A. V., Ho, L. C., Shields, J. C., Belloni, T., Comastri, A., Snowden, S. L., & Sramek, R. A. 1999, *PASP*, 111, 801
- Netzer, H. 1990, in *Active Galactic Nuclei*, ed. R. D. Blandford, H. Netzer, & L. Woltjer (Berlin: Springer), 57
- Oke, J. B., & Gunn, J. E. 1983, *ApJ*, 266, 713
- Peng, C. Y., Ho, L. C., Impey, C. D., & Rix, H.-W. 2002, *AJ*, 124, 266
- Pentericci, L., et al. 2002, *AJ*, 123, 2151
- Peterson, B. M. 1993, *PASP*, 105, 247
- Rowan-Robinson, M. 1985, *The Cosmological Distance Ladder* (New York: Freeman)
- Roy, J.-R., Belly, J., Dutil, Y., & Martin, P. 1996, *ApJ*, 460, 284
- Sandage, A. R., & Tammann, G. A. 1981, *A Revised Shapley-Ames Catalog of Bright Galaxies* (Washington, DC: Carnegie Inst. of Washington)
- Shih, D. C., Iwasawa, K., & Fabian, A. C. 2003, *MNRAS*, in press
- Spitzer, L., Jr. 1969, *ApJ*, 158, L139
- Smith, L. J., & Gallagher, J. S., III. 2001, *MNRAS*, 326, 1027
- Terlevich, E., Díaz, A. I., & Terlevich, R. 1990, *MNRAS*, 242, 271
- Tonry, J., & Davis, M. 1981, *ApJ*, 246, 666
- Tremaine, S., et al. 2002, *ApJ*, 574, 740
- Vogt, S. S. 1992, in *High Resolution Spectroscopy with the VLT*, ed. M.-H. Ulrich (Garching: ESO), 223
- Vogt, S. S., et al. 1994, *Proc. SPIE*, 2198, 362
- Whitmore, B. C., & Schweizer, F. 1995, *AJ*, 109, 960
- Wrobel, J. M., Fasnacht, C. D., & Ho, L. C. 2001, *ApJ*, 553, L23